

**BIOLOGY DATA BOOK**

## FOREWORD

The Biology Data Book is the third of the Biological Handbooks to be issued under the general direction of the Committee on Biological Handbooks of the Federation of American Societies for Experimental Biology, Washington, D. C. This volume continues a series of handbooks prepared under the auspices of the National Academy of Sciences-National Research Council, the first of which was published in 1952.

Participation in this undertaking was fulfilled under National Institutes of Health Grant No. GM 06533, National Science Foundation Grant No. GN 255, and Air Force Contract No. AF 33(657)-10802.

Dr. J. W. Heim, Technical Director of the Biophysics Laboratory, Aerospace Medical Research Laboratories, Aerospace Medical Division, Wright-Patterson Air Force Base, Ohio, was the technical monitor for the Air Force. The Air Force participation in this undertaking is in support of Project No. 7164, "Biomedical Criteria for Aerospace Flight," Task No. 716406, "Systemization of Biological Knowledge."

The Committee on Biological Handbooks acknowledges with thanks the contribution of 470 botanists, zoologists, and basic medical scientists who have contributed so generously with their time and advice. The Committee thanks the National Institutes of Health, the National Science Foundation, and the Aerospace Medical Research Laboratories for the generous support and cooperation that have made possible the production of this book.

This technical report has been reviewed and is approved.

J. W. HEIM, PhD  
Technical Director  
Biophysics Laboratory

## ABSTRACT

The Biology Data Book has been compiled to present numerical data of biology and medicine in a convenient and accessible form for reference, and to standardize accepted constants as a basis for correlation, establish common standards for statistical studies, and provide normal values for research. The biology data are organized in the form of tables, diagrams, charts, and graphs, arranged under the following headings: Genetics and Cytology, Reproduction, Development and Growth, Morphology, Nutrition and Digestion, Metabolism, Respiration and Circulation, Blood, Biological Regulators and Toxins, Biophysical and Biochemical Characteristics, Environment and Survival, Parasitism, and Materials and Methods. Seven appendices provide information concerning estimated number of species, taxonomic classification for living plants and animals, geologic distribution, atomic weights, as well as logarithms and antilogarithms. A detailed index completes the book. The contents have been authenticated by 470 authorities in the fields of biology and medicine. The review process of the tables was designed to eliminate, insofar as possible, errors of transcription and material of questionable validity.

## COMMITTEE ON BIOLOGICAL HANDBOOKS

RAYMUND L. ZWEMER, *Chairman*

PHILIP L. ALTMAN, *Executive Secretary*

SCOTT ADAMS  
National Library of Medicine  
Bethesda, Maryland  
CARL R. BREWER†  
National Institutes of Health  
Bethesda, Maryland  
GEORGE B. BROWN  
Sloan-Kettering Institute  
for Cancer Research  
Rye, New York

T. C. BYERLY  
U.S. Department of Agriculture  
Washington, D.C.  
F. S. CHEEVER\*  
University of Pittsburgh  
School of Medicine  
Pittsburgh, Pennsylvania  
J. W. HEIM  
Aerospace Medical Research  
Laboratories,  
Wright-Patterson Air Force Base,  
Ohio

MILTON O. LEE  
Federation of American Societies  
for Experimental Biology  
Washington, D.C.  
H. S. MAYERSON\*  
Tulane University School of Medicine  
New Orleans, Louisiana  
ILEEN E. STEWART†  
National Science Foundation  
Washington, D.C.

## BIOLOGY DATA BOOK ADVISORY COMMITTEE

RAYMUND L. ZWEMER, *Chairman*

TUNIS BAKER  
Hope College  
Holland, Michigan  
C. S. CHADWICK  
Emory and Henry College  
Emory, Virginia

F. M. CLARK  
University of Illinois  
Urbana, Illinois

HERMAN C. KRANZER  
Temple University  
Philadelphia, Pennsylvania  
THURLO B. THOMAS  
Carleton College  
Northfield, Minnesota

## HANDBOOK STAFF

PHILIP L. ALTMAN, *Director*

DOROTHY S. DITTMER, *Editor*

BETTY R. CONNERS  
MARY J. GRANT

SAKI HIMEL  
PHYLLIS JAY

KATHRYN F. NAYLOR  
M. LOUISE STANTON

† liaison  
\* *ex officio*



# CONTENTS

CONTRIBUTORS AND REVIEWERS . . . . .	xii
ABBREVIATIONS AND SYMBOLS . . . . .	xviii
INTRODUCTION . . . . .	xix

## I. GENETICS AND CYTOLOGY

1. Chromosome Numbers: Animals . . . . .	1
Part I. Vertebrates . . . . .	1
Part II. Invertebrates . . . . .	3
2. Chromosome Numbers: Plants . . . . .	6
Part I. Nonvascular . . . . .	6
Part II. Vascular . . . . .	8
3. Sex Linkage: Man . . . . .	11
4. Linkage Groups: Vertebrates . . . . .	13
Part I. Guinea Pig . . . . .	13
Part II. Mouse . . . . .	13
Part III. Rabbit . . . . .	17
Part IV. Rat . . . . .	18
Part V. Chicken . . . . .	19
5. Linkage Groups: Invertebrates . . . . .	20
Part I. Fruit Fly . . . . .	20
Part II. Parasitic Wasp . . . . .	29
Part III. Silkworm . . . . .	31
6. Linkage Groups: Plants . . . . .	33
Part I. <i>Neurospora crassa</i> . . . . .	33
Part II. <i>Chlamydomonas reinhardi</i> . . . . .	37
Part III. Corn . . . . .	39
Part IV. Tomato . . . . .	41
7. Genetic Code . . . . .	43
8. Cell Types: Seed Plants . . . . .	44
9. Tissue Growth Characteristics: Mammals . . . . .	46
10. Cell Division Frequency: Microorganisms . . . . .	51
Part I. Protozoa . . . . .	51
Part II. Viruses and Bacteria . . . . .	51
11. Organic Compounds Affecting Cell Division . . . . .	53

## II. REPRODUCTION

12. Propagation: Mammals . . . . .	57
13. Propagation: Birds . . . . .	59
Part I. Nest Building, Incubation, and Parental Care of Young . . . . .	59
Part II. Clutch Size . . . . .	60
Part III. Hatching Success: Precocial Species . . . . .	60
Part IV. Hatching and Fledging Success: Altricial Species . . . . .	61
14. Propagation: Reptiles . . . . .	62
15. Propagation: Amphibians . . . . .	63
16. Propagation: Fishes . . . . .	64
17. Propagation: Aquatic Invertebrates . . . . .	66
18. Propagation and Metamorphosis: Insects . . . . .	67
19. Propagation and Development: Invertebrates . . . . .	69
Part I. Metazoa . . . . .	69
Part II. Protozoa . . . . .	71
20. Breeding Systems: Angiosperms . . . . .	72
21. Propagation Methods: Cultivated Plants . . . . .	73
22. Seed Germination: Herbaceous Plants . . . . .	75
23. Seed Germination: Forest Trees, North American . . . . .	76

## III. DEVELOPMENT AND GROWTH

24. Early Prenatal Development: Man [Drawings] . . . . .	79
25. Germ Layers and Derivatives: Eutherian Mammals [Diagram] . . . . .	80
26. Time Variations in Developmental Stages: Mammals and Birds . . . . .	82
27. Characterization of Developmental Stages . . . . .	82
Part I. Man . . . . .	82
Part II. Rat . . . . .	84
Part III. Swine . . . . .	86
Part IV. Chick . . . . .	88
Part V. Frog . . . . .	89
Part VI. Salmonid Fishes . . . . .	91

# Contents

28. Growth: Mammals . . . . .	93
Part I. Body Weight and Height: Man . . . . .	93
Part II. Body Weight: Rodents . . . . .	94
Part III. Body Weight: Mammals Other than Man and Rodents . . . . .	97
29. Growth: Vertebrates Other than Mammals . . . . .	101
Part I. Body Weight: Birds . . . . .	101
Part II. Body Length: Reptiles and Amphibians . . . . .	102
Part III. Body Length and Weight: Fishes . . . . .	104
30. Life Spans: Animals . . . . .	106
Part I. Vertebrates . . . . .	106
Part II. Invertebrates . . . . .	109
31. Development and Life Spans: Forest Trees, North American . . . . .	110
32. Life Spans: Seeds . . . . .	111
Part I. In Air-dry Storage . . . . .	111
Part II. Undisturbed in Soil . . . . .	112
Part III. At Various Temperatures . . . . .	113
33. Life Spans: Pollen . . . . .	114
34. Growth Rates: Plant Tissues . . . . .	115

## IV. MORPHOLOGY

35. Body Composition with Increasing Weight and Age: Man [Graphs] . . . . .	119
36. Body Surface Area: Mammals . . . . .	120
Part I. Surface Area for Known Weight and Height: Man . . . . .	120
Part II. Constants for Use in Surface Area Formula: Mammals . . . . .	120
37. Brain: Man [Drawings] . . . . .	122
Part I. Regions and Functions . . . . .	123
Part II. Cortical Cerebral Regions and Functions [Drawings] . . . . .	124
Part III. Nuclei of Metathalamus and Dorsal Thalamus . . . . .	125
Part IV. Tracts . . . . .	127
38. Autonomic Nervous System: Man [Drawing] . . . . .	130
Part I. Sympathetic Connections . . . . .	131
Part II. Parasympathetic Connections . . . . .	134
Part III. Ganglia . . . . .	136
Part IV. Plexuses . . . . .	138
39. Digestive Enzymes: Vertebrates . . . . .	139
40. Comparative Anatomy of the Circulatory System: Vertebrates . . . . .	142
Part I. Heart . . . . .	142
Part II. Blood Vessels . . . . .	144
Part III. Lymphatics . . . . .	148
41. Comparative Anatomy of the Endocrine System: Vertebrates . . . . .	152
42. Comparative Anatomy of the Skeletal System: Mammals . . . . .	158
Part I. Axial Skeleton . . . . .	158
Part II. Appendicular Skeleton . . . . .	160

## V. NUTRITION AND DIGESTION

43. Nutrients: Chemical Elements . . . . .	165
44. Nutrients: Lipids . . . . .	167
45. Nutrients: Proteins, Peptides, and Amino Acids . . . . .	168
46. Nutrients: Purines and Pyrimidines . . . . .	171
47. Nutrients: Vitamins and Related Compounds . . . . .	172
48. Nutrients: Miscellaneous Growth Factors . . . . .	175
49. Nutrients: Carbon, Nitrogen, and Sulfur . . . . .	177
Part I. Carbon Sources . . . . .	177
Part II. Nitrogen Sources . . . . .	178
Part III. Sulfur Sources . . . . .	180
50. Pathways of Protein Digestion: Man and Laboratory Mammals [Diagram] . . . . .	183
51. Pathways of Carbohydrate Digestion: Man and Laboratory Mammals [Diagram] . . . . .	184
52. Pathways of Lipid Digestion: Man and Laboratory Mammals [Diagram] . . . . .	185
53. Excretion Products: Man . . . . .	186
Part I. Urine . . . . .	186
Part II. Feces . . . . .	190

## VI. METABOLISM

54. Pathways of Mineral Metabolism: Laboratory Mammals [Diagram] . . . . .	192
55. Pathways of Lipid Metabolism: Mammals [Diagram] . . . . .	197
56. Pathways of Carbohydrate Metabolism [Diagram] . . . . .	198
57. Pathways of Amino Acid Metabolism . . . . .	199
58. Pathways of Nucleoprotein Catabolism [Diagram] . . . . .	201
59. Pathways of Purine and Pyrimidine Catabolism [Diagram] . . . . .	202
60. Metabolic Interrelationships: Carbohydrate, Fat, and Protein [Diagram] . . . . .	203

# Contents

61. Krebs Cycle [Diagram] . . . . .	204
62. Cytochrome System [Diagram] . . . . .	205
63. Properties of Cytochromes: Animals and Higher Plants . . . . .	206
64. Pathways of Biosynthesis: Purines [Diagram] . . . . .	208
65. Pathways of Biosynthesis: Pyrimidines [Diagram] . . . . .	209
66. Pathways of Biosynthesis: Chlorophyll [Diagram] . . . . .	210
67. Pathways of Photosynthesis: Carbon Dioxide Reduction Cycle [Diagram] . . . . .	211
68. Pathways of Sucrose Synthesis: Intermediates [Diagram] . . . . .	212
69. Photosynthesis: Apparent Rates . . . . .	212
Part I. Maximum Rates: Natural Conditions, Various Locales . . . . .	212
Part II. Maximum Rates: Near-optimum Conditions . . . . .	214
Part III. Average Rates . . . . .	215
70. Carbon Production and Photosynthetic Efficiency . . . . .	216
Part I. Estimated Annual Carbon Production . . . . .	216
Part II. Energy Utilization in Photosynthesis . . . . .	216
71. Nitrogen Fixation . . . . .	216
Part I. Rhizobia-inoculated Legumes . . . . .	216
Part II. Characteristics of Nitrogen-fixing Organisms . . . . .	217
72. Nitrogen Cycle in Nature [Diagram] . . . . .	218

## VII. RESPIRATION AND CIRCULATION

73. Characteristics of Respiratory Media . . . . .	219
74. Lung Ventilation: Vertebrates . . . . .	220
75. Oxygen Consumption . . . . .	221
Part I. Mammals . . . . .	221
Part II. Vertebrates Other than Mammals . . . . .	221
Part III. Invertebrates Other than Protozoa . . . . .	223
Part IV. Protozoa . . . . .	224
76. Respiration Rates . . . . .	225
Part I. Bacteria . . . . .	225
Part II. Myxophyta and Fungi . . . . .	225
Part III. Lichens, Algae, and Bryophytes . . . . .	227
Part IV. Tracheophyta . . . . .	228
77. Heart Rates . . . . .	234
Part I. Man . . . . .	234
Part II. Vertebrates Other than Man . . . . .	234
Part III. Invertebrates . . . . .	237
78. Arterial Blood Pressure . . . . .	238
Part I. Man . . . . .	238
Part II. Animals Other than Man . . . . .	239
79. Vascular and Capillary Pressures . . . . .	241
Part I. Vascular Pressures: Man . . . . .	241
Part II. Relationship of Peripheral Arterial to Central Arterial Pressure: Man . . . . .	242
Part III. Venous Blood Pressure: Man . . . . .	242
Part IV. Capillary Blood Pressure: Vertebrates . . . . .	242

## VIII. BLOOD

80. Blood Group Systems: Man . . . . .	245
Part I. Phenotypes and Genotypes of the A-B-O System . . . . .	245
Part II. Partial List of Allelic Genes of the M-N System . . . . .	246
Part III. Phenotypes and Genotypes of the Rh-Hr System . . . . .	247
Part IV. Partial List of Allelic Genes of the Rh-Hr System . . . . .	248
81. Heredity of Blood Groups and Types: Man . . . . .	249
Part I. A-B-O Exclusion . . . . .	249
Part II. M-N Exclusion . . . . .	249
Part III. Rh-Hr Exclusion . . . . .	249
82. Distribution of Blood Groups and Types in Various Populations: Man . . . . .	250
Part I. A-B-O Groups . . . . .	250
Part II. M-N Types . . . . .	251
Part III. Rh-Hr Types . . . . .	252
83. Blood Coagulation Theories [Diagrams] . . . . .	253
Part I. According to F. C. Monkhouse and W. W. Coon (1963) . . . . .	253
Part II. According to P. A. Owren (1963) . . . . .	254
Part III. According to A. J. Quick (1963) . . . . .	256
Part IV. According to W. H. Seegers (1963) . . . . .	257
Part V. According to L. M. Tocantins (1960) . . . . .	258
84. Acid-Base Balance . . . . .	259
Part I. Acid-Base Values: Man . . . . .	259
Part II. Acid-Base Values: Vertebrates . . . . .	259
Part III. Normal Ionic Patterns, Arterial Blood: Man [Graphs] . . . . .	262
Part IV. Classification of Acid-Base Disturbances: Man . . . . .	262

# Contents

85. Blood Volumes . . . . .	263
Part I. Vertebrates . . . . .	263
Part II. Insects . . . . .	266
86. Erythrocyte and Platelet Values . . . . .	267
Part I. Erythrocyte and Hemoglobin Values: Vertebrates . . . . .	267
Part II. Blood Platelet Count: Mammals . . . . .	271
87. Leukocyte Counts . . . . .	272
Part I. Man . . . . .	272
Part II. Vertebrates Other than Man . . . . .	273
88. Bone Marrow Differential Cell Counts . . . . .	275
Part I. Rib: Dog . . . . .	275
Part II. Sternum: Man . . . . .	275
Normal Blood and Marrow Cells: Man [Color Plate] . . . . .	<i>facing page</i> 276

## IX. BIOLOGICAL REGULATORS AND TOXINS

89. Enzymes . . . . .	277
Part I. Catalytic Action . . . . .	277
Part II. Physical and Kinetic Properties . . . . .	282
Part III. Chemical Composition . . . . .	288
90. Hormones: Vertebrates . . . . .	290
91. Endocrine Organs and Hormones: Invertebrates . . . . .	304
92. Relative Activity of Growth Regulators: Plants . . . . .	307
Part I. Cell Elongation of Oat Coleoptiles . . . . .	307
Part II. Stem Curvature of Slit Pea and Leaf Expansion of Bean . . . . .	308
93. Antimetabolites . . . . .	309
94. Antibiotics . . . . .	312
Part I. Physical and Chemical Characteristics . . . . .	312
Part II. Biological Activity . . . . .	319
95. Anticoagulants . . . . .	325
96. Animal Toxins . . . . .	328
Part I. Reptiles . . . . .	328
Part II. Toads . . . . .	334
Part III. Marine Organisms . . . . .	336
97. Plant Toxins . . . . .	344

## X. BIOPHYSICAL AND BIOCHEMICAL CHARACTERISTICS

98. Carbohydrates: Physical and Chemical Characteristics . . . . .	351
Part I. Natural Monosaccharides: Aldoses and Ketoses . . . . .	351
Part II. Natural Monosaccharides: Amino Sugars . . . . .	355
Part III. Natural Alditols and Inositols (with Inososes and Inosamines) . . . . .	356
Part IV. Natural Aldonic, Uronic, and Aldaric Acids . . . . .	358
Part V. Natural Carbohydrate Phosphate Esters . . . . .	360
Part VI. Natural Oligosaccharides . . . . .	364
99. Glycosides: Characteristics, Occurrence, and Uses . . . . .	368
100. Fatty Acids: Physical and Chemical Characteristics . . . . .	370
101. Fats and Oils: Physical and Chemical Characteristics . . . . .	380
102. Waxes: Physical and Chemical Characteristics . . . . .	382
103. Phosphatides and Cerebrosides: Physical and Chemical Characteristics . . . . .	383
104. Sterols: Physical and Chemical Characteristics . . . . .	385
105. Proteins: Physical and Chemical Characteristics . . . . .	388
106. Amino Acids: Physical and Chemical Characteristics . . . . .	392
107. Vitamins and Provitamins: Physical and Chemical Characteristics . . . . .	394
108. Various Cells and Cell Parts: Chemical Composition . . . . .	398
109. Animal Tissues and Organs: Water Content . . . . .	401
110. Cell Sap: Chemical Composition . . . . .	404
111. Plant Tissues and Organs: Mineral Composition . . . . .	405
Part I. Major Elements . . . . .	405
Part II. Minor Elements . . . . .	411

## XI. ENVIRONMENT AND SURVIVAL

112. Hibernation: Mammals and Birds . . . . .	417
113. Diapause: Insects and Mites . . . . .	419
114. Dispersion of Small Organisms . . . . .	420
Part I. Invertebrates . . . . .	420
Part II. Viruses, Bacteria, and Fungi . . . . .	426
Part III. Pollen and Seeds . . . . .	428
115. Effect of Temperature on Inactivation and Survival: Viruses . . . . .	431
Part I. Animal Viruses . . . . .	431
Part II. Plant Viruses . . . . .	432



# Contents

116. Effect of Temperature on Growth and Survival: Rickettsia and Bacteria . . . . .	438
Part I. Optimum Temperature for Growth. . . . .	438
Part II. Thermal Death Time . . . . .	439
117. Effect of Temperature on Growth and Survival: Fungi . . . . .	440
118. Temperature Tolerances: Algae . . . . .	441
119. Soil pH: Spermatophytes. . . . .	442
120. Shade Tolerance: Vascular Plants . . . . .	443
121. Effect of Light on Development: Angiosperms . . . . .	443
Part I. Various Wavelengths. . . . .	443
Part II. Various Exposures . . . . .	444
122. Photoperiod, with Temperature Interactions, for Flowering: Angiosperms . . . . .	446
123. Factors Affecting Protoplasmic Streaming: Plants . . . . .	448
Part I. Temperature. . . . .	448
Part II. Sudden Changes of Temperature. . . . .	448
Part III. Light Intensity: Avena Coleoptile . . . . .	449
Part IV. Various Wavelengths: Avena Coleoptile . . . . .	450
Part V. Oxygen . . . . .	450
124. Factors Affecting Transpiration Rates: Angiosperms . . . . .	451
Part I. Various Conditions . . . . .	451
Part II. Variation in Soil Conditions: Corn . . . . .	452
Part III. Diurnal Variation: Corn . . . . .	452
Part IV. Annual Variation . . . . .	453
125. Factors Affecting Osmotic Potential: Vascular Plants . . . . .	453
Part I. Species Variation: Leaves. . . . .	453
Part II. Physical and Environmental Variation. . . . .	454
Part III. Variation in Depth of Rooting . . . . .	456
Part IV. Variation in Habitat . . . . .	456
Part V. Variation in Ecologic Groups. . . . .	456
126. Maximum Permissible Occupational Exposure to Radiation: Man. . . . .	457
Part I. Dose Equivalent to Body Organs . . . . .	457
Part II. Type of Radiation . . . . .	457
Part III. Internal Concentration of Radionuclides . . . . .	458
127. Late Effects of Irradiation: Mammals. . . . .	468

## XII. PARASITISM

128. Arthropod Parasites: Mammals and Birds. . . . .	477
129. Arthropod Pests: Plants and Plant Products . . . . .	481
130. Helminth and Protozoan Parasites: Mammals and Birds . . . . .	486
Part I. Man. . . . .	486
Part II. Vertebrates Other than Man . . . . .	490
131. Nematode Parasites: Plants . . . . .	494
132. Viral Diseases: Animals . . . . .	498
133. Viral Diseases: Plants. . . . .	500
134. Rickettsial Parasites: Mammals and Birds . . . . .	503
135. Bacterial Parasites: Mammals and Birds . . . . .	504
136. Bacterial Parasites: Plants . . . . .	506
137. Fungal Parasites: Plants . . . . .	508
Part I. Field, Fruit, and Vegetable Crops . . . . .	508
Part II. Forest Trees . . . . .	511
138. Mistletoe Parasites: Forest Trees. . . . .	514
139. Fungal Parasites: Man. . . . .	516
Part I. Superficial Mycoses . . . . .	516
Part II. Deep Mycoses. . . . .	518

## XIII. MATERIALS AND METHODS

140. Culture Media: Protozoa . . . . .	523
Part I. Parasitic Amoebae . . . . .	523
Part II. Trichomonadidae. . . . .	524
Part III. Trypanosomatidae . . . . .	526
Part IV. Phytomastigina . . . . .	528
141. Culture Media: Animal Tissues. . . . .	529
Part I. Balanced Salt Solutions . . . . .	529
Part II. Tissue Culture Media. . . . .	530
142. Culture Media: Plants . . . . .	534
Part I. Bacteria . . . . .	534
Part II. Fungi . . . . .	536
Part III. Algae . . . . .	536
Part IV. Higher Plants . . . . .	537
143. Culture Media: Plant Tissues . . . . .	538
Part I. Balanced Salt Solutions . . . . .	538
Part II. Tissue Culture Media. . . . .	538

# Contents

144. Natural Sea Water . . . . .	539
Part I. General Characteristics, Salinity, and Constituents . . . . .	539
Part II. Surface Temperature of the Oceans . . . . .	540
Part III. Relation of Chlorinity and Salinity to Density . . . . .	540
Part IV. Oxygen Saturation from Normal Dry Atmosphere . . . . .	540
Part V. Pressure-Depth Gradient . . . . .	541
145. Artificial Sea Water . . . . .	541
146. Normal Solutions . . . . .	543
147. Buffer Solutions: pH Ranges . . . . .	543
148. Weak Acids and Bases: pK Values . . . . .	544
149. Acid-Base Indicators: pH Ranges . . . . .	545
150. Oxidation-Reduction Indicators . . . . .	545
151. Radionuclides Used in Biological Research . . . . .	546
152. Anesthetics . . . . .	547
153. Fixatives and Clearing Agents . . . . .	549
Part I. Fixatives . . . . .	549
Part II. Clearing Agents . . . . .	551
154. Staining Methods . . . . .	551
Part I. Living Materials . . . . .	551
Part II. Fixed Materials . . . . .	553
155. Histochemical Tests . . . . .	557

## APPENDICES

Appendix I. Estimated Number of Species: Animal and Plant Kingdoms . . . . .	561
Appendix II. Taxonomic Classification: Living Animals . . . . .	562
Appendix III. Taxonomic Classification: Living Plants . . . . .	566
Part I. Nonvascular Plants . . . . .	566
Part II. Vascular Plants . . . . .	567
Appendix IV. Geologic Distribution: Animals and Plants . . . . .	570
Appendix V. Formulas, Factors, and Constants . . . . .	571
Part I. Conversion Formulas . . . . .	571
Part II. Conversion Factors . . . . .	572
Part III. Numerical Constants and Binomial Coefficients . . . . .	578
Part IV. Physical Constants . . . . .	579
Appendix VI. Atomic Weights . . . . .	579
Appendix VII. Logarithms and Antilogarithms . . . . .	580
Part I. Four-Place Logarithms . . . . .	580
Part II. Four-Place Antilogarithms . . . . .	582

INDEX . . . . .	585
-----------------	-----

## CONTRIBUTORS AND REVIEWERS

- ABBOTT, R. TUCKER  
Academy of Natural Sciences  
Philadelphia, Pennsylvania
- ACHOR, LEONARD B.  
Sandoz Pharmaceuticals  
Hanover, New Jersey
- AHMADJIAN, V.  
Clark University  
Worcester, Massachusetts
- ALDRICH, FREDERICK A.  
Memorial University  
St. Johns, Newfoundland, Canada
- ALLEN, FRED H., JR.  
Blood Grouping Laboratory  
Boston, Massachusetts
- ALLEN, MARY BELLE  
Kaiser Foundation Research  
Institute  
Richmond, California
- ALLEN, WILLIAM W.  
University of California  
Berkeley, California
- ALLFREY, VINCENT G.  
Rockefeller Institute  
New York, New York
- ALTLAND, PAUL D.  
National Institutes of Health  
Bethesda, Maryland
- AMBROSE, CHARLES TESCH  
Harvard University  
Cambridge, Massachusetts
- ANDERSEN, AXEL L.  
USDA, Crops Research Division  
East Lansing, Michigan
- ANDERSON, DONALD B.  
University of North Carolina  
Chapel Hill, North Carolina
- ANDERSON, LEWIS E.  
Duke University  
Durham, North Carolina
- ANDREW, WARREN  
Indiana University  
Indianapolis, Indiana
- ANDREWARTHA, H. G.  
University of Adelaide  
Adelaide, South Australia
- APPLEMAN, MILO D.  
University of Southern California  
Los Angeles, California
- AREY, LESLIE B.  
Northwestern University  
Chicago, Illinois
- ARIMOTO, KUNITARO  
National Institute of Nutrition  
Tokyo, Japan
- ARMER, SISTER JOSEPH MARIE  
Incarnate Word College  
San Antonio, Texas
- ARMSTRONG, J. M.  
University of Adelaide  
Adelaide, South Australia
- ASDELL, S. A.  
Cornell University  
Ithaca, New York
- BAILEY, LOWELL F.  
University of Arkansas  
Fayetteville, Arkansas
- BALLARD, W. W.  
Dartmouth College  
Hanover, New Hampshire
- BANKS, HARLAN P.  
Cornell University  
Ithaca, New York
- BARRATT, R. W.  
Dartmouth College  
Hanover, New Hampshire
- BARRETT, HAROLD W.  
Jacksonville University  
Jacksonville, Florida
- BARRINGTON, E. J. W.  
University of Nottingham  
Nottingham, England
- BARTELMEZ, GEORGE W.  
224 Agnes Avenue  
Missoula, Montana
- BARTGIS, I. LOUISE  
National Institutes of Health  
Bethesda, Maryland
- BARTON, LELA V.  
Boyce Thompson Institute for  
Plant Research  
Yonkers, New York
- BASS, DAVID E.  
U. S. Quartermaster Research &  
Engineering Command  
Natick, Massachusetts
- BASSHAM, JAMES A.  
University of California  
Berkeley, California
- BATEMAN, ANGUS J.  
Christie Hospital & Holt Radium  
Institute  
Withington, Manchester, England
- BATES, ROGER G.  
National Bureau of Standards  
Washington, D. C.
- BAWDEN, F. C.  
Rothamsted Experiment Station  
Harpenden, Hertfordshire,  
England
- BAXTER, DOW V.  
University of Michigan  
Ann Arbor, Michigan
- BENNETT, C. W.  
USDA, Field Crops Research  
Branch  
Salinas, California
- BENNETT, L. R.  
University of California  
Los Angeles, California
- BENNETT, W. F.  
Agricultural & Mechanical College  
of Texas  
College Station, Texas
- BENSON, ANDREW A.  
Pennsylvania State University  
University Park, Pennsylvania
- BERN, HOWARD A.  
University of California  
Berkeley, California
- BILLINGS, MARTA S.  
University of California  
Los Angeles, California
- BING, ARTHUR  
Cornell University  
Farmingdale, New York
- BIRD, ORSON D.  
Parke, Davis & Co.  
Ann Arbor, Michigan
- BISHOP, DAVID W.  
Carnegie Institution of Washington  
Baltimore, Maryland
- BISHOPP, FRED C.  
3823 East River Drive  
Fort Myers, Florida
- BLAIR, ALBERT P.  
University of Tulsa  
Tulsa, Oklahoma
- BLANDAU, RICHARD J.  
University of Washington  
Seattle, Washington
- BOHNING, RICHARD H.  
Ohio State University  
Columbus, Ohio
- BONNER, JAMES F.  
California Institute of Technology  
Pasadena, California
- BONNYCASTLE, DESMOND D.  
Seton Hall College of Medicine &  
Dentistry  
Jersey City, New Jersey
- BOWMAN, H. H. M.  
Toledo Hospital  
Toledo, Ohio
- BRASE, KARL D.  
Cornell University  
Geneva, New York
- BRAUNWALD, EUGENE  
National Institutes of Health  
Bethesda, Maryland
- BRECHER, GEORGE  
National Institutes of Health  
Bethesda, Maryland
- BRIDGMAN, CHARLES F.  
University of California  
Los Angeles, California
- BRIGGS, GEORGE M.  
University of California  
Berkeley, California
- BROUN, G. O.  
St. Louis University  
St. Louis, Missouri
- BROWN, ELLEN  
University of California  
San Francisco, California
- BROWN, GEORGE B.  
Sloan-Kettering Institute for  
Cancer Research  
Rye, New York
- BROWN, JAMES W.  
Chemical Corps Biological  
Laboratories  
Fort Detrick, Frederick, Maryland
- BROWN, RELIS B.  
Florida State University  
Tallahassee, Florida
- BUCK, JOHN B.  
National Institutes of Health  
Bethesda, Maryland
- BURKHOLDER, W. H.  
Greycourt Apartments  
Ithaca, New York
- BURNS, GEORGE W.  
Ohio Wesleyan University  
Delaware, Ohio



# Contrails

- BUTLER, L.  
University of Toronto  
Toronto, Ontario, Canada
- CAGLE, FRED R.  
Tulane University  
New Orleans, Louisiana
- CALDER, D. M.  
University College of Wales  
Plas Gogerddan, near Aberystwyth,  
Wales
- CALESNICK, BENJAMIN  
Hahnemann Medical College  
Philadelphia, Pennsylvania
- CALHOUN, JOHN B.  
National Institutes of Health  
Bethesda, Maryland
- CALVIN, MELVIN  
University of California  
Berkeley, California
- CAMPBELL, BERRY  
Los Angeles County Hospital  
Los Angeles, California
- CAMPBELL, JACK J. R.  
University of British Columbia  
Vancouver, British Columbia,  
Canada
- CANTINO, EDWARD C.  
Michigan State University  
East Lansing, Michigan
- CAPLIN, SAMUEL M.  
Los Angeles State College  
Los Angeles, California
- CARLANDER, KENNETH D.  
Iowa State University  
Ames, Iowa
- CARLETON, RALPH K.  
Curry College  
Milton, Massachusetts
- CARRIKER, MELBOURNE R.  
U. S. Fish & Wildlife Service  
Oxford, Maryland
- CARSCALLEN, LEONA J.  
College of Medical Evangelists  
Loma Linda, California
- \*CASTLE, W. E.
- CAVE, MARION S.  
University of California  
Berkeley, California
- CHASTAIN, SARAH  
University of California  
Los Angeles, California
- CHEN, K. K.  
Indiana University  
Indianapolis, Indiana
- CHRISTENSEN, P. AGERHOLM  
South African Institute for Medical  
Research  
Johannesburg, Union of South Africa
- CHRISTIE, JESSE R.  
Route 1  
Newport, Nova Scotia, Canada
- CLAPP, GRACE L.  
1245 Palisade Avenue  
Windsor, Connecticut
- CLARK, F. M.  
University of Illinois  
Urbana, Illinois
- CLARKE, NORMAN E.  
Providence Hospital  
Detroit, Michigan
- COLE, LaMONT C.  
Cornell University  
Ithaca, New York
- CONKLIN, RUTH E.  
Vassar College  
Poughkeepsie, New York
- COON, WILLIAM W.  
University of Michigan  
Ann Arbor, Michigan
- COOPER, J. P.  
University College of Wales  
Plas Gogerddan, near Aberystwyth,  
Wales
- COPENHAVER, WILFRED M.  
Columbia University  
New York, New York
- CORLEY, RALPH C.  
Purdue University  
Lafayette, Indiana
- CORNMAN, IVOR  
Lerner Marine Laboratory  
Miami, Florida
- COWAN, IAN McTAGGART  
University of British Columbia  
Vancouver, British Columbia,  
Canada
- CRONKITE, EUGENE P.  
Brookhaven National Laboratory  
Upton, Long Island, New York
- CROWN, R. M.  
Louisiana State University  
Baton Rouge, Louisiana
- CUNNINGHAM, CHARLES H.  
Michigan State University  
East Lansing, Michigan
- CUTKOMP, LAURENCE K.  
University of Minnesota  
St. Paul, Minnesota
- D'AMATO, FRANCESCO  
Istituto di Genetica della  
Università  
Pisa, Italy
- DAMON, ALBERT  
Harvard University  
Boston, Massachusetts
- DARBY, RICHARD T.  
U. S. Quartermaster Research &  
Engineering Command  
Natick, Massachusetts
- DARLINGTON, C. D.  
University of Oxford  
Oxford, England
- DAVIS, DAVID E.  
Pennsylvania State University  
University Park, Pennsylvania
- DAWE, ALBERT R.  
Office of Naval Research  
Chicago, Illinois
- DeGARIS, CHARLES F.  
University of Oklahoma  
Oklahoma City, Oklahoma
- DeMARSH, Q. B.  
University of Washington  
Seattle, Washington
- DeRITTER, E.  
Hoffman-La Roche Inc.  
Nutley, New Jersey
- DEUTSCH, MARSHALL E.  
Becton, Dickinson & Co.  
Englewood Cliffs, New Jersey
- DIAMOND, LOUIS S.  
National Institutes of Health  
Bethesda, Maryland
- DIANZANI, MARIO U.  
University of Cagliari  
Sardinia, Italy
- DICKEY, ROBERT S.  
Cornell University  
Ithaca, New York
- DIGGS, L. W.  
University of Tennessee  
Memphis, Tennessee
- do AMARAL, AFRANIO  
Instituto Butantan  
São Paulo, Brazil
- DOWNES, R. J.  
USDA Agricultural Research  
Center  
Beltsville, Maryland
- DOZIER, BYRD K.  
University of Tennessee  
Memphis, Tennessee
- DuBOIS, R. CALLERY  
505 W. Chestnut Hill Avenue  
Philadelphia, Pennsylvania
- \*DUCA, CHARLES J.
- DUGGAN, T. L.  
Loyola University  
New Orleans, Louisiana
- DUNLAP, J. S.  
Washington State University  
Pullman, Washington
- DUPRÉ, MARGARET V.  
State University College  
Buffalo, New York
- EAMES, A. J.  
Cornell University  
Ithaca, New York
- EATON, ORSON N.  
4320 Clagett Road  
Hyattsville, Maryland
- EBERSOLD, W. T.  
University of California  
Los Angeles, California
- EDGAR, S. A.  
Auburn University  
Auburn, Alabama
- ELISBERG, EDWARD I.  
104 S. Michigan Avenue  
Chicago, Illinois
- ELWYN, DAVID H.  
Michael Reese Hospital & Medical  
Center  
Chicago, Illinois
- ERDMAN, LEWIS W.  
USDA, Soil & Water Conservation  
Research Division  
Beltsville, Maryland
- EVANS, ROBERT JOHN  
Michigan State University  
East Lansing, Michigan
- FAUST, ERNEST CARROLL  
Tulane University  
New Orleans, Louisiana
- FERGUSON, JOHN H.  
University of North Carolina  
Chapel Hill, North Carolina
- FITCH, HENRY S.  
University of Kansas  
Lawrence, Kansas

\* Deceased



# Contrails

- FITCH, JOHN E.  
State Department of Fish & Game  
Terminal Island, California
- FLEMISTER, LAUNCE J.  
Swarthmore College  
Swarthmore, Pennsylvania
- FLOCK, EUNICE V.  
Mayo Foundation  
Rochester, Minnesota
- FOGG, G. E.  
Westfield College  
London, England
- FORSTER, ROBERT E.  
University of Pennsylvania  
Philadelphia, Pennsylvania
- FORWARD, DOROTHY F.  
University of Toronto  
Toronto, Ontario, Canada
- FOSTER, ADRIANCE S.  
University of California  
Berkeley, California
- FREED, S. CHARLES  
Mt. Zion Hospital  
San Francisco, California
- FREIS, EDWARD D.  
Veterans Administration Hospital  
Washington, D. C.
- FRIEDMAN, LORRAINE  
Tulane University  
New Orleans, Louisiana
- FROBISHER, MARTIN  
P.O. Box 267  
Harwich, Massachusetts
- FULTON, ROBERT W.  
University of Wisconsin  
Madison, Wisconsin
- FURMAN, DEANE P.  
University of California  
Berkeley, California
- GARB, SOLOMON  
University of Missouri  
Columbia, Missouri
- GEORG, LUCILLE K.  
U. S. Public Health Service  
Atlanta, Georgia
- GEYER, ROBERT P.  
Harvard University  
Cambridge, Massachusetts
- GIDDENS, JOEL  
University of Georgia  
Athens, Georgia
- GLASER, KURT  
University of Maryland  
Baltimore, Maryland
- GLUCKSMANN, A.  
Strangeways Research Laboratory  
Cambridge, England
- GORBMAN, AUBREY  
Columbia University  
New York, New York
- GORDON, HAROLD THOMAS  
University of California  
Berkeley, California
- GORDON, MORRIS A.  
State Department of Health  
Albany, New York
- GOTS, JOSEPH S.  
Long Island Biological Association  
Cold Spring Harbor, New York
- GRAHAM, JOHN B.  
University of North Carolina  
Chapel Hill, North Carolina
- GRANICK, S.  
Rockefeller Institute  
New York, New York
- GRAY, PETER  
University of Pittsburgh  
Pittsburgh, Pennsylvania
- GRAYDON, JOHN J.  
Commonwealth Serum Laboratories  
Parkville, Victoria, Australia
- GREEN, MARGARET C.  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- GREULACH, VICTOR A.  
University of North Carolina  
Chapel Hill, North Carolina
- GRIFFITH, JOHN QUINTIN, Jr.  
Griffith Foundation for Medical  
Research  
Philadelphia, Pennsylvania
- GRODZINSKI, Z.  
Jagellonian University  
Kraków, Poland
- GROSSMAN, MORTON I.  
Veterans Administration Center  
Los Angeles, California
- GUEST, GEORGE M.  
University of Cincinnati  
Cincinnati, Ohio
- HAEUSSLER, G. J.  
USDA, Bureau of Entomology &  
Plant Quarantine  
Washington, D. C.
- HAGEN, CHARLES W., JR.  
Indiana University  
Bloomington, Indiana
- HALDE, CARLYN  
University of California  
San Francisco, California
- HALL, FRANK G.  
Duke University  
Durham, North Carolina
- HALSTEAD, BRUCE W.  
World Life Research Institute  
Reche Canyon, Colton, California
- HAMERSLAG, FRANK E.  
Wyeth Laboratories, Inc.  
Philadelphia, Pennsylvania
- HAMILTON, HOWARD L.  
Iowa State University  
Ames, Iowa
- HAMRE, CHRISTOPHER J.  
University of North Dakota  
Grand Forks, North Dakota
- HANSARD, SAM L.  
Louisiana State University  
Baton Rouge, Louisiana
- HARDY, ROSS  
Long Beach State College  
Long Beach, California
- HARRAR, E. S.  
Duke University  
Durham, North Carolina
- HARRELL, GEORGE T.  
University of Florida  
Gainesville, Florida
- HART, J. SANFORD  
National Research Council  
Ottawa, Canada
- HARTMAN, OLGA  
University of Southern California  
Los Angeles, California
- HARTROFT, W. STANLEY  
Washington University  
St. Louis, Missouri
- HARWOOD, H. J.  
Durkee Famous Foods  
Chicago, Illinois
- HASKINS, R. H.  
National Research Council  
Saskatoon, Saskatchewan, Canada
- HASTINGS, A. BAIRD  
Harvard University  
Cambridge, Massachusetts
- HAUROWITZ, FELIX  
Indiana University  
Bloomington, Indiana
- HEISLER, CHARLES R.  
Oregon State College  
Corvallis, Oregon
- HEMINGWAY, ALLAN  
University of California  
Los Angeles, California
- HENDERSON, LAVANIEL L., SR.  
Texas Southern University  
Houston, Texas
- HENSCHEL, AUSTIN  
U. S. Quartermaster Research &  
Engineering Command  
Natick, Massachusetts
- HERNANDEZ, THOMAS  
Louisiana State University  
New Orleans, Louisiana
- HERRMANN, ROY G.  
Eli Lilly & Co.  
Indianapolis, Indiana
- HERTIG, ARTHUR T.  
Harvard University  
Boston, Massachusetts
- HESE, CLARON O.  
University of California  
Davis, California
- HEWITT, HAROLD B.  
Westminster School of Medicine  
Horseferry Road, London, England
- HILL, BERTON F.  
National Academy of Sciences  
Washington, D. C.
- HIMWICH, WILLIAMINA A.  
State Research Hospital  
Galesburg, Illinois
- HOCK, RAYMOND J.  
University of California  
Pig Pine, California
- HOLLANDER, FRANKLIN  
Mount Sinai Hospital  
New York, New York
- HOLMES, FRANCIS O.  
Rockefeller Institute  
New York, New York
- HOUSE, HOWARD L.  
Canadian Department of Agriculture  
Belleville, Ontario, Canada
- HOWELL, ROBERT W.  
USDA, Regional Soybean Laboratory  
Urbana, Illinois
- HUTT, F. B.  
Cornell University  
Ithaca, New York
- IDLER, D. R.  
Fisheries Research Board  
Halifax, Nova Scotia, Canada

# Contrails

- IRVIN, J. LOGAN  
University of North Carolina  
Chapel Hill, North Carolina
- JAQUES, LOUIS B.  
University of Saskatchewan  
Saskatoon, Saskatchewan, Canada
- JENNISON, MARSHALL W.  
Syracuse University  
Syracuse, New York
- JOHNSON, B. CONNOR  
University of Illinois  
Urbana, Illinois
- JOHNSON, ELTON L.  
University of Minnesota  
St. Paul, Minnesota
- JOHNSON, RICHARD P.  
Louisa, Virginia
- JONES, GALEN E.  
University of California  
La Jolla, California
- JONES, JACK COLVARD  
University of Maryland  
College Park, Maryland
- JONES, RUTH McCLUNG  
Winthrop College  
Rock Hill, South Carolina
- JUSTICE, O. L.  
USDA, Agricultural Research Center  
Beltsville, Maryland
- KAHN, BERND  
National Academy of Sciences  
Washington, D. C.
- KALISZEWSKI, BARBARA FREEMAN  
Boston University  
Boston, Massachusetts
- KASSANIS, B.  
Rothamsted Experiment Station  
Harpenden, Hertfordshire,  
England
- KATZ, MAX  
University of Washington  
Seattle, Washington
- KEMP, NORMAN E.  
University of Michigan  
Ann Arbor, Michigan
- KENDEIGH, S. CHARLES  
University of Illinois  
Urbana, Illinois
- KIESSELBACH, T. A.  
University of Nebraska  
Lincoln, Nebraska
- KIKKAWA, H.  
Osaka University  
Kitaku, Osaka, Japan
- KIRKHAM, WILLIAM R.  
Oklahoma A. & M. College  
Stillwater, Oklahoma
- KISCH, Bruno  
71 Maple Street  
Brooklyn, New York
- KLEIN, RICHARD M.  
New York Botanical Garden  
Bronx Park, New York, New York
- KLEINER, ISRAEL S.  
New York Medical College  
New York, New York
- KNIPLING, E. F.  
USDA, Entomology Research  
Division  
Beltsville, Maryland
- KNOBLOCH, IRVING W.  
Michigan State University  
East Lansing, Michigan
- KOLLROS, JERRY J.  
University of Iowa  
Iowa City, Iowa
- KOSER, STEWART A.  
University of Chicago  
Chicago, Illinois
- KRAMER, PAUL J.  
Duke University  
Durham, North Carolina
- KRATZER, F. H.  
University of California  
Davis, California
- KRAUSS, BEATRICE  
Pineapple Research Institute  
Honolulu, Hawaii
- KROGMAN, W. M.  
Philadelphia Center for Research  
in Child Growth  
Philadelphia, Pennsylvania
- KRUTA, VLADISLAV  
Masaryk University  
Komenského nám. 2, BRNO,  
Czechoslovakia
- KUCK, KATHRYN D.  
c/o Emory University Medical  
School  
Atlanta, Georgia
- \*KUNTZ, ALBERT
- LANSFORD, EDWIN M., JR.  
University of Texas  
Austin, Texas
- LARSON, EDWARD  
University of Miami  
Coral Gables, Florida
- LATIMER, HOMER B.  
University of Kansas  
Lawrence, Kansas
- LATYSZEWSKI, M.  
Institute of Animal Genetics  
Edinburgh, Scotland
- LEE, JOHN J.  
Haskins Laboratories  
New York, New York
- LEES, A. D.  
Agricultural Research Council  
Cambridge, England
- LEVINE, E. E.  
Harvard University  
Boston, Massachusetts
- LEVINE, NORMAN D.  
University of Illinois  
Urbana, Illinois
- LEVINE, PHILIP  
Ortho Research Foundation  
Raritan, New Jersey
- LEVINE, VICTOR E.  
Creighton University  
Omaha, Nebraska
- LEVITT, J.  
University of Missouri  
Columbia, Missouri
- LIGHT, AMOS E.  
Wellcome Research Laboratories  
Tuckahoe, New York
- LIMARZI, LOUIS R.  
University of Illinois  
Chicago, Illinois
- LINDQUIST, A. W.  
USDA, Entomology Research  
Division  
Beltsville, Maryland
- LINDSAY, HUGH A.  
West Virginia University  
Morgantown, West Virginia
- LINK, ROGER P.  
University of Illinois  
Urbana, Illinois
- LITTLE, ELBERT L., JR.  
USDA, U.S. Forest Service  
Washington, D. C.
- LOCHHEAD, JOHN H.  
University of Vermont  
Burlington, Vermont
- LOEFER, JOHN B.  
Office of Naval Research  
Pasadena, California
- LOGAN, J. E.  
Department of National Health &  
Welfare  
Ottawa, Ontario, Canada
- LOMBARD, ELNA A.  
Medical College of Georgia  
Augusta, Georgia
- LOOSANOFF, VICTOR L.  
U.S. Fish & Wildlife Service  
Milford, Connecticut
- LOVE, R. M.  
Torry Research Station  
Aberdeen, Scotland
- LYMAN, CHARLES P.  
Harvard University  
Cambridge, Massachusetts
- McCHESNEY, EVAN W.  
Sterling-Winthrop Research  
Institute  
Rensselaer, New York
- McCUTCHEON, F. HAROLD  
University of Pennsylvania  
Philadelphia, Pennsylvania
- McILRATH, WAYNE J.  
University of Chicago  
Chicago, Illinois
- McKUSICK, VICTOR A.  
Johns Hopkins Hospital  
Baltimore, Maryland
- McLOUD, E. S.  
S. C. Johnson & Son, Inc.  
Racine, Wisconsin
- McMEEKIN, T. L.  
USDA Eastern Utilization Division  
Philadelphia, Pennsylvania
- MACOUN, HORACE W.  
University of California  
Los Angeles, California
- MAHER, GEORGE G.  
Clinton Corn Processing Co.  
Clinton, Iowa
- MAHLSTEDE, JOHN P.  
Iowa State University  
Ames, Iowa
- MAKINO, SAJIRO  
University of Hokkaido  
Sapporo, Japan
- MANDELS, GABRIEL R.  
U.S. Quartermaster Research &  
Engineering Command  
Natick, Massachusetts

\* Deceased

# Contrails

- MANVILLE, RICHARD H.  
U.S. Dept. Interior, Wildlife  
Research  
Washington, D. C.
- MARAMOROSCH, KARL  
Boyce Thompson Institute for Plant  
Research  
Yonkers, New York
- MARKLEY, KLARE S.  
Correio de Copacabana  
Rio de Janeiro, Brazil
- MASTER, ARTHUR M.  
125 East 72nd Street  
New York, New York
- MAYERSON, H. S.  
Tulane University  
New Orleans, Louisiana
- MEISTER, ALTON  
Tufts University  
Boston, Massachusetts
- MENDLOWITZ, MILTON  
2 East 95th Street  
New York, New York
- MEYER, MARION P.  
University of Wisconsin  
Madison, Wisconsin
- MIGDALSKI, EDWARD C.  
Yale University  
New Haven, Connecticut
- MILLS, CLARENCE A.  
Cincinnati General Hospital  
Cincinnati, Ohio
- MINTON, SHERMAN A., JR.  
Indiana University  
Indianapolis, Indiana
- MITCHELL, G. A. G.  
University of Manchester  
Manchester, England
- MONIE, I. W.  
University of California  
San Francisco, California
- MONKHOUSE, FRANK C.  
University of Toronto  
Toronto, Ontario, Canada
- MOOG, FLORENCE  
Washington University  
St. Louis, Missouri
- MORGAN, F. G.  
Commonwealth Serum Laboratories  
Parkville, Victoria, Australia
- MORGAN, KARL Z.  
Oak Ridge National Laboratory  
Oak Ridge, Tennessee
- MORRISON, PETER R.  
University of Wisconsin  
Madison, Wisconsin
- MORTON, C. V.  
Smithsonian Institution  
Washington, D. C.
- MORTON, JULIA F.  
University of Miami  
Coral Gables, Florida
- MORTON, R. K.  
University of Adelaide  
Adelaide, S. A., Australia
- MOSBY, HENRY S.  
Virginia Polytechnic Institute  
Blacksburg, Virginia
- MOSES, HAROLD E.  
Purdue University  
Lafayette, Indiana
- MOYER, ELIZABETH K.  
Boston University  
Boston, Massachusetts
- MUIR, ROBERT M.  
University of Iowa  
Iowa City, Iowa
- MUSACCHIA, X. J.  
St. Louis University  
St. Louis, Missouri
- MYERS, JACK  
University of Texas  
Austin, Texas
- NARDONE, ROLAND M.  
Catholic University of America  
Washington, D. C.
- NICE, MARGARET MORSE  
5725 Harper Avenue  
Chicago, Illinois
- NIEMER, WILLIAM T.  
Creighton University  
Omaha, Nebraska
- NIRENBERG, MARSHALL W.  
National Institutes of Health  
Bethesda, Maryland
- NOLTMANN, ERNST A.  
University of California  
Riverside, California
- NOVITSKI, E.  
University of Oregon  
Eugene, Oregon
- O'BRIEN, JOHN S.  
University of Southern California  
Los Angeles 33, California
- \*O'CONNOR, R. J.
- OLIVE, LINDSAY S.  
Columbia University  
New York, New York
- OLSON, F. C. W.  
Radio Corporation of America  
Princeton, New Jersey
- OLSON, RODNEY A.  
National Institutes of Health  
Bethesda, Maryland
- OMAN, PAUL W.  
A.P.O. 143, Box ND  
San Francisco, California
- OSER, BERNARD L.  
Food & Drug Research Labora-  
tories, Inc.  
Maspeth, New York
- OSGOOD, EDWIN E.  
University of Oregon  
Portland, Oregon
- OWREN, PAUL A.  
Rikshospitalet  
Oslo, Norway
- PAGNUCCO, RINALDO G.  
Ohio State University  
Columbus, Ohio
- PARKER, RAYMOND C.  
University of Toronto  
Toronto, Ontario, Canada
- PATTEN, BRADLEY M.  
University of Michigan  
Ann Arbor, Michigan
- PAVCEK, PAUL L.  
Rhineland, Wisconsin
- PELLETIER, RÉAL L.  
McGill University  
Montreal, Quebec, Canada
- PERLMAN, D.  
35 University Place  
Princeton, New Jersey
- PETT, L. BRADLEY  
Department of National Health &  
Welfare  
Ottawa, Ontario, Canada
- PFEIFFER, NORMA E.  
14 Odell Avenue  
Yonkers, New York
- PHILIP, CORNELIUS B.  
U.S. Public Health Service  
Hamilton, Montana
- PISEK, A.  
Botanisches Institut der  
Universität  
Innsbruck, Austria
- PORTER, B. A.  
USDA, Entomology Research  
Division  
Beltsville, Maryland
- PORTER, JOHN N.  
American Cyanamid Co.  
Pearl River, New York
- POTTS, CARL G.  
USDA, Agricultural Research  
Center  
Beltsville, Maryland
- PRITHAM, GORDON H.  
Pennsylvania State University  
University Park, Pennsylvania
- PROVASOLI, LUIGI  
Haskins Laboratories  
New York, New York
- PURVIS, E. R.  
147 North Sixth Avenue  
Highland Park, New Jersey
- QUICK, ARMAND J.  
Marquette University  
Milwaukee, Wisconsin
- RAAF, JOHN  
833 Southwest 11th Avenue  
Portland, Oregon
- REDFIELD, ALFRED C.  
Woods Hole Oceanographic  
Institute  
Woods Hole, Massachusetts
- REHDER, HARALD A.  
Smithsonian Institution  
Washington, D. C.
- REICH, HANS  
Landon Foundation Research  
Institute of Chemotherapy  
Colorado Springs, Colorado
- REKERS, PAUL E.  
1400 North Vermont Avenue  
Los Angeles, California
- REYER, RANDALL W.  
West Virginia University  
Morgantown, West Virginia
- REYNOLDS, MONICA  
University of Pennsylvania  
Kennett Square, Pennsylvania
- RHOADES, M. M.  
Indiana University  
Bloomington, Indiana

\* Deceased



# Contributors

- RICHARDS, OSCAR W.  
American Optical Co.  
Southbridge, Massachusetts
- RICHERT, DAN A.  
State University of New York  
Syracuse, New York
- RICK, CHARLES M.  
University of California  
Davis, California
- RIGDON, R. H.  
University of Texas  
Galveston, Texas
- RITCHER, PAUL O.  
Oregon State College  
Corvallis, Oregon
- ROBB, JANE SANDS  
Route 1, Box 149  
Biloxi, Mississippi
- ROBBINS, W. REI  
Rutgers University  
New Brunswick, New Jersey
- ROBERTS, R. H.  
University of Wisconsin  
Madison, Wisconsin
- ROBINSON, R. A.  
National Bureau of Standards  
Washington, D. C.
- ROCKSTEIN, MORRIS  
University of Miami  
Miami, Florida
- RODBARD, SIMON  
Chronic Disease Research Institute  
Buffalo, New York
- ROE, EUGENE I.  
USDA, U.S. Forest Service  
St. Paul, Minnesota
- ROE, JOSEPH H.  
George Washington University  
Washington, D. C.
- ROGERS, WILLIAM M.  
Columbia University  
New York, New York
- ROLLIN, S. F.  
USDA, Agricultural Research Center  
Beltsville, Maryland
- ROOT, RAYMOND W.  
City University of New York  
New York, New York
- ROSSETTI, VICTORIA  
Instituto Biológico  
São Paulo, Brazil
- RUBIN, SAUL H.  
Hoffman-La Roche, Inc.  
Nutley, New Jersey
- RUDOLF, PAUL O.  
University of Minnesota  
St. Paul, Minnesota
- RUSOFF, LOUIS LEON  
Louisiana State University  
Baton Rouge, Louisiana
- RUSSELL, FINDLAY E.  
Loma Linda University  
Los Angeles, California
- RUSSELL, JANE A.  
Emory University  
Atlanta, Georgia
- SAGER, RUTH  
Columbia University  
New York, New York
- SALLACH, H. J.  
University of Wisconsin  
Madison, Wisconsin
- SALTMAN, PAUL  
University of Southern California  
Los Angeles, California
- SAMUELS, GEORGE  
Agricultural Experiment Station  
Rio Piedras, Puerto Rico
- SASSER, J. N.  
North Carolina State College  
Raleigh, North Carolina
- SAUBERLICH, HOWERDE E.  
Fitzsimons General Hospital  
Denver, Colorado
- SAWIN, PAUL B.  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- SAX, KARL  
Harvard University  
Cambridge, Massachusetts
- SCHAEFER, ARNOLD EDWARD  
National Institutes of Health  
Bethesda, Maryland
- SCHÖTTLER, WERNER H. A.  
Sydney Ross Co.  
Rio de Janeiro, Brazil
- SCHUBERT, LEO  
Council of Chief State School  
Officers  
Washington, D. C.
- SCOTT, J. P.  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- SCOTT, ROLAND B.  
Freedmen's Hospital  
Washington, D. C.
- SEEGERS, WALTER H.  
Wayne State University  
Detroit, Michigan
- SELIGER, VÁCLAV  
Salmovska 5  
Prague, Czechoslovakia
- SELLMER, GEORGE P.  
Upsala College  
East Orange, New Jersey
- SENDROY, JULIUS, JR.  
National Naval Medical Center  
Bethesda, Maryland
- SHANNON, F. A.  
Box 276  
Wickenburg, Arizona
- SHAW, CHARLES E.  
Zoological Society of San Diego  
San Diego, California
- SHELTON, MAURICE  
Agricultural & Mechanical College  
of Texas  
McGregor, Texas
- SHIVE, WILLIAM  
University of Texas  
Austin, Texas
- SHORE, MARY S.  
University of Maryland  
College Park, Maryland
- SHUSTER, CARL N., JR.  
University of Delaware  
Newark, Delaware
- SIEGEL, JACK M.  
Pabst Laboratories  
Milwaukee, Wisconsin
- SILBERSCHMIDT, KARL M.  
Instituto Biológico  
São Paulo, Brazil
- \*SILVERMAN, MILTON  
SINGER, RICHARD B.  
New England Mutual Life Insurance  
Co.  
Boston, Massachusetts
- SIRI, WILLIAM E.  
University of California  
Berkeley, California
- SKUTCH, ALEXANDER F.  
Finca "Los Cusingos"  
San Isidro del General, Costa Rica
- SLATE, GEORGE L.  
Cornell University  
Geneva, New York
- SLOTTA, KARL H.  
University of Miami  
Miami, Florida
- SMITH, CLEMENT A.  
Boston Lying-In Hospital  
Boston, Massachusetts
- SNELL, GEORGE D.  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- SOMERS, G. FRED  
University of Delaware  
Newark, Delaware
- SOROKIN, CONSTANTINE  
University of Maryland  
College Park, Maryland
- STANLEY, W. W.  
University of Tennessee  
Knoxville, Tennessee
- STARR, RICHARD C.  
Indiana University  
Bloomington, Indiana
- \*STEINBAUER, GEORGE P.  
STEVENS, RUSSELL B.  
George Washington University  
Washington, D. C.
- STEVENSON, JAMES A. F.  
University of Western Ontario  
London, Ontario, Canada
- STRICKLAND, W. N.  
Dartmouth College  
Hanover, New Hampshire
- STROUD, ROBERT  
U.S. Public Health Service  
Cincinnati, Ohio
- STRUCKMEYER, BURDEAN E.  
University of Wisconsin  
Madison, Wisconsin
- SUTIN, JEROME  
Yale University  
New Haven, Connecticut
- \*SVERDRUP, H. U.  
SWETT, WALTER W.  
USDA, Dairy Cattle Research  
Branch  
Beltsville, Maryland
- TAMURA, T.  
Fisheries Research Board of  
Canada  
Halifax, Nova Scotia, Canada

\* Deceased

- TANNER, VASCO M.  
Brigham Young University  
Provo, Utah
- TEMPLETON, GEORGE S.  
17118 Merrill Avenue  
Fontana, California
- TERRY, LUTHER L.  
U.S. Public Health Service  
Washington, D. C.
- THIMANN, KENNETH V.  
Harvard University  
Cambridge, Massachusetts
- THOMAS, THURLO B.  
Carleton College  
Northfield, Minnesota
- THOMPSON, RANDALL L.  
National Institutes of Health  
Bethesda, Maryland
- THOMSON, JOHN F.  
Argonne National Laboratory  
Argonne, Illinois
- TIETZE, CHRISTOPHER  
National Committee on Maternal  
Health  
New York, New York
- TOBIE, ELEANOR J.  
National Institutes of Health  
Bethesda, Maryland
- \*TOCANTINS, LEANDRO M.
- TUPPER, RONALD  
Medical College of Saint  
Bartholomew's Hospital  
London, England
- TURNER, ROBERT A.  
Duke Laboratories, Inc.  
South Norwalk, Connecticut
- TURRELL, FRANKLIN M.  
University of California  
Riverside, California
- VAN BRUGGEN, JOHN T.  
University of Oregon  
Portland, Oregon
- VANDEBELT, J. M.  
Parke, Davis & Co.  
Ann Arbor, Michigan
- VAN LIERE, EDWARD J.  
West Virginia University  
Morgantown, West Virginia
- VAN PILSUM, JOHN F.  
University of Minnesota  
Minneapolis, Minnesota
- VAN WAGENEN, GERTRUDE  
Yale University  
New Haven, Connecticut
- VAN WAGTENDONK, W. J.  
9720 Southwest 114th Street  
Miami, Florida
- von BONIN, GERHARDT  
Mount Zion Hospital  
San Francisco, California
- von BRAND, THEODOR  
National Institutes of Health  
Bethesda, Maryland
- WAINIO, WALTER W.  
Rutgers University  
New Brunswick, New Jersey
- WALKER, HENRY  
University of Alabama  
University, Alabama
- WALKER, RICHARD B.  
University of Washington  
Seattle, Washington
- WALKER, SHEPPARD M.  
University of Louisville  
Louisville, Kentucky
- WARD, WILFRED H.  
USDA, Western Utilization Division  
Albany, California
- WARREN, KATHERINE BREHME  
National Institutes of Health  
Bethesda, Maryland
- WARTH, ALBIN H.  
29 York Court  
Baltimore, Maryland
- WATTS, R. W. E.  
Medical College of Saint  
Bartholomew's Hospital  
London, England
- WAY, KATHARINE  
National Academy of Sciences  
Washington, D. C.
- WAYMOUTH, CHARITY  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- WEAGLEY, JOHN L.  
Upper Ironia Road  
Mendham, New Jersey
- WEBB, RAYMON E.  
USDA, Crops Research Division  
Beltsville, Maryland
- WEDGWOOD, RALPH J.  
Western Reserve University  
Cleveland, Ohio
- WEINTRAUB, ROBERT L.  
Chemical Corps Biological  
Laboratories  
Fort Detrick, Frederick,  
Maryland
- WELT, ISAAC D.  
Institute for Advancement of  
Medical Communication  
Washington, D. C.
- WETMORE, RALPH H.  
Harvard University  
Cambridge, Massachusetts
- WHERRY, EDGAR T.  
University of Pennsylvania  
Philadelphia, Pennsylvania
- WHITE, FRED N.  
University of Texas  
Dallas, Texas
- WHITE, PHILIP R.  
Roscoe B. Jackson Memorial  
Laboratory  
Bar Harbor, Maine
- WHITING, P. W.  
University of Pennsylvania  
Philadelphia, Pennsylvania
- WIENER, ALEXANDER S.  
64 Rutland Road  
Brooklyn, New York
- WILKES, A.  
Department of Agriculture  
Belleville, Ontario, Canada
- \*WILLIAMS, BERT C.
- WILLS, E. D.  
Medical College of Saint  
Bartholomew's Hospital  
London, England
- WINDLE, WILLIAM F.  
National Institutes of Health  
Bethesda, Maryland
- WINTROBE, M. M.  
Salt Lake County General Hospital  
Salt Lake City, Utah
- WITSCHI, EMIL  
University of Basel  
Basel, Switzerland
- WOLF, FREDERICK T.  
Vanderbilt University  
Nashville, Tennessee
- WOLFENBARGER, D. O.  
University of Florida  
Homestead, Florida
- WOLFROM, MELVILLE L.  
Ohio State University  
Columbus, Ohio
- WOODBURY, ROBERT A.  
University of Tennessee  
Memphis, Tennessee
- WOOLLEY, D. W.  
Rockefeller Institute  
New York, New York
- WRIGHT, IRVING S.  
Cornell University  
New York, New York
- WRIGHT, SEWALL  
University of Wisconsin  
Madison, Wisconsin
- WYMAN, DONALD  
Harvard University  
Boston, Massachusetts
- YOCUM, L. EDWIN  
1322 Weber Drive  
Clearwater, Florida
- YOUNG, I. MAUREEN  
St. Thomas's Hospital  
London, England
- ZAUMEYER, WILLIAM J.  
USDA, Crops Research Division  
Beltsville, Maryland
- ZBARSKY, S. H.  
University of British Columbia  
Vancouver, British Columbia,  
Canada
- ZIPKIN, ISADORE  
National Institutes of Health  
Bethesda, Maryland
- ZOBELL, CLAUDE E.  
University of California  
La Jolla, California
- ZUCKER, LOIS M.  
Laboratory of Comparative  
Pathology  
Stow, Massachusetts

\* Deceased

# Contrails

## ABBREVIATIONS AND SYMBOLS

### Measurements

ht	= height	wt	= weight
mi	= mile	lb	= pound
ft	= foot	g	= gram
in.	= inch	kg	= kilogram
m	= meter	mg	= milligram
km	= kilometer	$\mu$ g	= microgram
dm	= decimeter	$\mu\mu$ g	= micromicrogram
cm	= centimeter	mEq	= milliequivalent
mm	= millimeter	gr	= grain
$\mu$	= micron	M	= mole
m $\mu$	= millimicron	mM	= millimole
Å	= Ångström unit	$\mu$ M	= micromole
yr	= year	L	= liter
mo	= month	ml	= milliliter
wk	= week	$\mu$ l	= microliter
da	= day	I.U.	= international unit
hr	= hour	ppm	= parts per million
min	= minute	vol %	= volume percent
sec	= second	$^{\circ}$ C	= degrees centigrade
cgs	= centimeter-gram-second	$^{\circ}$ F	= degrees Fahrenheit
rpm	= revolutions per minute	cal	= calorie
ft-c	= foot-candle	kcal	= kilocalorie
atm	= atmosphere	BTU	= British thermal unit
sq	= square	>	= greater than
cu	= cubic	<	= less than

### Biological and Chemical Specifications

♂	= male	<i>d</i>	= <i>dextro</i> (rotatory)
♀	= female	<i>l</i>	= <i>levo</i> (rotatory)
sp.	= species (singular)	<i>D</i>	= <i>dextro</i> (in configurational sense only)
spp.	= species (plural)	<i>L</i>	= <i>levo</i> (in configurational sense only)
po	= oral	<i>m</i>	= meta
rec	= rectal	<i>o</i>	= ortho
sc	= subcutaneous	<i>p</i>	= para
im	= intramuscular	<i>M</i>	= molar
ip	= intraperitoneal	<i>N</i>	= normal, or <i>nitro</i>
iv	= intravenous	<i>O</i>	= <i>oxy</i>
RBC	= red blood cell (erythrocyte)	<i>S</i>	= <i>sulf</i> or <i>sulfo</i>
WBC	= white blood cell (leukocyte)	STP	= standard temperature and pressure
CNS	= central nervous system		
CSF	= cerebrospinal fluid		



# Contrails

## INTRODUCTION

The *Biology Data Book* is a volume of broad scope and limited coverage designed to serve as a basic reference in the field of biology. It is a radical revision of the *Handbook of Biological Data* published in 1956 by the W. B. Saunders Company.

Much has been learned over the past eight years from users of the old handbook. In order to incorporate their suggestions for improvement, i.e., larger type, literature citations, and a detailed index, it became obvious that the number of tables to be included in the *Biology Data Book* would have to be restricted to a more discriminating selection. The Committee on Biological Handbooks assigned the task of choosing the basic tables for the new general reference book to a specially appointed *Biology Data Book* Advisory Committee. Copies of the old handbook and of the specialized volumes in the Biological Handbooks series were sent to the members of the Advisory Committee, who used these books for two years in daily work situations. On the basis of frequency of referral, the Advisory Committee selected 143 tables for extensive revision and updating, and recommended the inclusion of 12 additional tables containing data of fundamental importance and current relevance.

The space limitations affecting subject coverage also imposed restrictions on the number of species to be included in the *Biology Data Book*. The Advisory Committee approved a list of approximately 400 species, which included the more common animals and plants, certain physiologically unique forms, and the size extremes within taxonomic groups. Frequently data were not accessible for a plant or animal appearing on the list, but were available for a related form. In such cases, the information for the related organism was used in the tabulation. In the tables on toxins and parasitism, the inclusion of data was dependent on whether the victim or host, rather than the offending organism, appeared on the approved list.

The *Biology Data Book* has been organized in the form of quantitative and descriptive tables, charts, and diagrams, and arranged in 13 sections for the convenience of the user. Contents of the volume have been authenticated by 470 leading investigators in the fields of botany, zoology, and medicine. The review process to which the data have been subjected was designed to eliminate, insofar as possible, material of questionable validity and errors of transcription.

An explanatory headnote, serving as an introduction to the subject matter, may precede a table. More frequently, tables are prefaced by a short headnote containing such important information as units of measurement, abbreviations, definitions, and estimate of the range of variation. To interpret the data, reading of the related headnote is essential.

The main conventions used throughout the data book have been adapted from the *Style Manual for Biological Journals*, published in 1960 for the Conference of Biological Editors by the American Institute of Biological Sciences. The terminology has been checked against *Webster's Third New International Dictionary*, published in 1961 by G. & C. Merriam Company.

Appended to the tables are the names of the contributors, and a list of the literature citations arranged in


alphabetical sequence. The reference abbreviations conform to the *1961 Chemical Abstracts List of Periodicals* published by the American Chemical Society.

It is suggested that the table of contents be used in conjunction with the index: the table of contents to determine the scope of the data for a particular topic, and the index to locate data for a specific subject or organism. To facilitate identification, the index includes the taxonomic orders for animals, and the family for plants.

.....

Values are generally presented as a mean and the lower and upper limit of the range of individual values about the mean. This range may be estimated in several ways, the method depending on the information available. Letter designations (a, b, c, d) identify types of ranges in descending order of accuracy.

(a) When the group of values is relatively large, a 95% range is derived by curve fitting. A recognized type of normal frequency curve is fitted to a group of measured values, and the extreme 2.5% of the area under the curve at each end is excluded (*see illustration*).



(b) When the group of values is too small for curve fitting, as is usually the case, a 95% range is estimated by a simple statistical calculation. Assuming a normal symmetrical distribution, the standard deviation is multiplied by a factor of 2, then subtracted from and added to the mean to give the lower and upper range limits.

(c) A less dependable, but commonly applied, procedure takes as range limits the lowest value and the highest value of the reported sample group of measurements. It underestimates the 95% range for small samples and overestimates for larger sample sizes, but may be used in preference to the preceding method where there is marked asymmetry in the position of the mean within the sample range.

(d) Another estimate of the lower and upper limits of the range of variation is based on the judgment of an individual experienced in measuring the quantity in question. The trustworthiness of such limits should not be underestimated.

# *Contrails*



# I. GENETICS AND CYTOLOGY

## 1. CHROMOSOME NUMBERS: ANIMALS

For information on additional species, consult references 2 and 26, Part I.

### Part I. VERTEBRATES

**Diploid** (column C): s = spermatogonium; o = oogonium; m = somatic cell. **Haploid** (column D): ♂(I) = primary spermatocyte; ♂(II) = secondary spermatocyte.

	Species	Common Name	Chromosome Number		Sex Type	Reference
			Diploid	Haploid		
	(A)	(B)	(C)	(D)	(E)	(F)
Mammalia						
1	<i>Homo sapiens</i>	Man	46s, o, m	23♂(I, II)	X-Y♂	60
2	<i>Bos taurus</i>	Cattle	60m	.....	X-Y♂	22,52
3	<i>Camelus bactrianus</i>	Bactrian camel	.....	35♂(I)	.....	39
4	<i>Canis familiaris</i>	Dog	78m	.....	X-Y♂	3
5	<i>Capra hircus</i>	Goat	60s	30♂(I, II)	X-Y♂	22,23
6	<i>Cavia porcellus</i>	Guinea pig	64m	.....	X-Y♂	3
7	<i>Dasyypus novemcinctus</i>	Nine-banded armadillo	60m	.....	X-O♂?	42,43
8	<i>Didelphis marsupialis virginiana</i>	Virginia opossum	22s, m	11♂(I, II)	X-Y♂	41
9	<i>Equus caballus</i>	Horse	64m	.....	X-Y♂	52
10	<i>Erinaceus europaeus</i>	European hedgehog	48s	24♂(I)	X-Y♂	4,5
11	<i>Felis catus</i>	Cat	38m	.....	X-Y♂	3
12	<i>Macaca mulatta</i>	Rhesus monkey	42	.....	X-Y♂	7
13	<i>Mesocricetus auratus</i>	Golden hamster	44m	.....	.....	3
14	<i>Mus musculus</i>	House mouse	40s	20♂(I, II)	X-Y♂	20
15	<i>Mustela vison</i>	Mink	30	.....	.....	55
16	<i>Myotis myotis</i>	Common brown bat	44s	22♂(I)	X-Y♂	5
17	<i>Ondatra zibethica</i>	Muskrat	54s	.....	.....	27
18	<i>Ornithorhynchus anatinus</i>	Platypus	70±10s	.....	.....	33
19	<i>Oryctolagus cuniculus</i>	European rabbit	44s	22♂(I)	X-Y♂	23,33
20	<i>Ovis aries</i>	Sheep	54m	.....	X-Y♂	34
21	<i>Phocaena dalli</i>	Dall's porpoise	44s	22♂(I, II)	X-Y♂	25
22	<i>Rattus norvegicus</i>	Norway rat	42m	.....	X-Y♂	28
23	<i>Sciurus carolinensis</i>	Gray squirrel	48s	.....	X-Y♂	8
24	<i>Sorex araneus</i>	European shrew	23s	11♂(I); 11,12♂(II)	X <sub>1</sub> X <sub>2</sub> -Y♂	4,5
25	<i>Sus scrofa</i>	Swine	40s	20♂(I, II)	X-Y♂	21,23
Aves						
26	<i>Anas platyrhynchos</i>	Mallard duck	80s	40♂(I)	.....	69
27	<i>Anser albifrons</i>	White-fronted goose	82s	41♂(I)	.....	69
28	<i>Columba livia</i>	Street pigeon	80s, 79o	40♂(I)	X-X♂, X-O♀	70
29	<i>Cygnus cygnus</i>	Whooper swan	80s	.....	.....	69
30	<i>Gallus domesticus</i>	Chicken	78s, 77o	39♂(I)	X-X♂, X-O♀	66,67
31	<i>Larus crassirostris</i>	Black-tailed gull	64s	.....	.....	61
32	<i>Meleagris gallopavo</i>	Turkey	82s, 81o	41♂(I)	X-X♂, X-O♀	68
33	<i>Passer domesticus</i>	House sparrow	54-60s, m	23♂(I)?	X-X♂, X-O♀	46
34	<i>Phasianus colchicus</i>	Ring-necked pheasant	82s, 81o	41♂(I)	X-X♂, O♀	66
35	<i>Pica pica</i>	Black-billed magpie	82s, 81o	.....	X-O♀	56
36	<i>Turdus merula</i>	Blackbird	60-85s, o, m	.....	X-X♂, X-O♀	62
Reptilia						
37	<i>Alligator mississippiensis</i>	American alligator	32s	16♂(I)	.....	48
38	<i>Ancistrodon acutus</i>	Mexican copperhead	36s	18♂(I)	X-X♂	37
39	<i>Anguis fragilis</i>	Slowworm	44s, o	22♂(I)	X-Y♀?	30
40	<i>Anotis carolinensis</i>	American "chameleon"	36s	18♂(I, II)	X-X♂	32
41	<i>Caretta caretta</i>	Loggerhead turtle	58s, 57o	.....	X-X♂, X-O♀	38
42	<i>Chrysemys marginata</i>	Painted turtle	.....	17♂(I)	X-O♂	11
43	<i>Emys orbicularis</i>	European pond turtle	50s	25♂(I)	X-X♂	32
44	<i>Eumeces elegans</i>	Elegant skink	26s	.....	.....	29,35
45	<i>Heloderma suspectum</i>	Gila monster	38s	19♂(I, II)	X-X♂	31,32
46	<i>Naja naja</i>	Indian cobra	38s	19♂(I, II)	X-X♂	37
47	<i>Natrix tigrina</i>	Japanese water snake	40s	20♂(I, II)	X-X♂	36

continued

# Contrails

## 1. CHROMOSOME NUMBERS: ANIMALS

### Part I. VERTEBRATES

	Species	Common Name	Chromosome Number		Sex Type	Reference
			Diploid	Haploid		
	(A)	(B)	(C)	(D)	(E)	(F)
Reptilia						
48	<i>Sceloporus spinosus</i>	Spiny fence lizard	22s	11♂(I)	XX-O♂?	40
49	<i>Sphenodon punctatus</i>	Tuatara	36s	18♂(I, II)	.....	12
50	<i>Sternotherus odoratus</i>	Musk turtle	50s	25♂(I, II)	.....	47
51	<i>Thamnophis butleri</i>	Butler's garter snake	37s	18♂(I)	XX-Y♂?	59
Amphibia						
52	<i>Ambystoma tigrinum</i>	Tiger salamander	28s	14♂(I)	X-O♂	6
53	<i>Amphiuma means</i>	Two-toed amphiuma	.....	12♂(I, II)	.....	14
54	<i>Bufo americanus</i>	American toad	.....	11♂(I, II)	X-Y♂?	65
55	<i>Bufo arenarum</i>	Sand toad	22s	11♂(I, II)	.....	50,51
56	<i>Cryptobranchus alleganiensis</i>	Hellbender	62s	31♂(I, II)	.....	16
57	<i>Hyla arborea</i>	Tree frog	24s	12♂(I, II)	.....	10,63
58	<i>Necturus maculosus</i>	Mud puppy	.....	12♂(I)	.....	13
59	<i>Rana catesbeiana</i>	American bullfrog	26s	13♂(I, II)	.....	24
60	<i>R. pipiens</i>	Leopard frog	26s	13♂(I)	.....	45
61	<i>Triturus cristatus</i>	Crested newt	24s	12♂(I)	.....	63
62	<i>T. viridescens</i>	Common newt	22m	.....	.....	9
63	<i>Xenopus laevis</i>	Clawed frog	36s	.....	.....	63,64
Pisces						
64	<i>Anguilla anguilla</i>	European freshwater eel	36s, o?	.....	.....	49
65	<i>Carassius auratus</i>	Goldfish	94s	47♂(I, II)	.....	15,19
66	<i>Coregonus albula</i>	European lake whitefish	80m	.....	.....	57
67	<i>Cyprinus carpio</i>	Carp	104s	52♂(I, II)	.....	15,18
68	<i>Esox lucius</i>	Northern pike	18m	.....	.....	58
69	<i>Fundulus heteroclitus</i>	Mummichog	45m	.....	.....	44
70	<i>Lepidosiren paradoxa</i>	South American lungfish	38m	19♂(I)	.....	1
71	<i>Osmerus eperlanus</i>	European smelt	58m	.....	.....	57
72	<i>Perca fluviatilis</i>	European perch	28m	.....	.....	58
73	<i>Protopterus annectans</i>	West African lungfish	34s	17♂(I)	.....	63
74	<i>Salmo salar</i>	Atlantic salmon	60m	30♂(I)	.....	57
75	<i>S. trutta</i>	Brown trout	80m	40♂(I)	.....	57
76	<i>Salvelinus fontinalis</i>	Eastern brook trout	84m	.....	.....	57
Chondrichthyes						
77	<i>Raja meerdervoortii</i>	Skate	104s	52♂(I, II)	.....	17
78	<i>Squalus suckleyi</i>	Pacific spiny dogfish	62s	31♂(I, II)	.....	17
Agnatha						
79	<i>Myxine glutinosa</i>	Atlantic hagfish	ca.52s, m	26♂(I, II)	.....	53,54

Contributor: Makino, Sajiro

References: [1] Agar, W. E. 1912. Quart. J. Microscop. Sci. 58:285. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Awa, A., M. Sasaki, and S. Takayama, 1959. Japan. J. Zool. 12:257. [4] Bovey, R. 1949. Experientia 5:72. [5] Bovey, R. 1949. Rev. Suisse Zool. 56:371. [6] Carrick, R. 1934. Trans. Roy. Soc. Edinburgh 58:63. [7] Chu, E. H. Y., and N. H. Giles. 1957. Am. Naturalist 91:273. [8] Cross, J. C. 1931. J. Morphol. 52:373. [9] Fankhauser, G. 1941. Ibid. 68:161. [10] Galgano, M. 1933. Arch. Ital. Anat. Embriol. 32:171. [11] Jordan, D. S., S. Tanaka, and J. O. Snyder. 1914. Science 39:178. [12] Keenan, R. D. 1932. J. Anat. 67:1. [13] King, H. D. 1912. Anat. Record 6:405. [14] McGregor, J. H. 1899. J. Morphol. 15(Suppl.):57. [15] Makino, S. 1934. Nippon Idengaku Zasshi 9:100. [16] Makino, S. 1935. J. Morphol. 58:573. [17] Makino, S. 1937. Cytologia (Tokyo), Fujii Jubilai Vol. (2):867. [18] Makino, S. 1939. Cytologia (Tokyo) 9:430. [19] Makino, S. 1941. Ibid. 12:96. [20] Makino, S. 1941. J. Fac. Sci. Hokkaido Imp. Univ., VI, 7:305. [21] Makino, S. 1943-44. Cytologia (Tokyo) 13:170. [22] Makino, S. 1943-44. Ibid. 13:247. [23] Makino, S.

continued

## 1. CHROMOSOME NUMBERS: ANIMALS

### Part I. VERTEBRATES

1944. Dobutsugaku Zasshi 56:8. [24] Makino, S. 1947. Kromosomo (Tokyo) 3-4:137. [25] Makino, S. 1948. Chromosoma 3:220. [26] Makino, S. 1951. An atlas of the chromosome numbers in animals. Ed. 2. Iowa State College Press, Ames. [27] Makino, S. 1953. Science 118:630. [28] Makino, S., and T. C. Hsu. 1954. Cytologia (Tokyo) 19:23. [29] Makino, S., and E. Momma. 1949. Ibid. 15:153. [30] Margot, A. 1946. Rev. Suisse Zool. 53:555. [31] Matthey, R. 1931. Bull. Soc. Vaudoise Sci. Nat. 57:269. [32] Matthey, R. 1931. Rev. Suisse Zool. 38:117. [33] Matthey, R. 1949. Les chromosomes des vertébrés. Rouge, Lausanne. [34] Melander, Y. 1959. Hereditas 45:649. [35] Momma, E. 1948. Dobutsugaku Zasshi 58. [36] Nakamura, K. 1928. Mem. Coll. Sci. Kyoto Imp. Univ., B, 4:1. [37] Nakamura, K. 1935. Ibid., B, 10:341. [38] Nakamura, K. 1949. Kromosomo (Tokyo) 5. [39] Novikov, I. I. 1940. Tr. Inst. Genet. Akad. Nauk SSSR 13:285. [40] Painter, T. S. 1921. J. Exptl. Zool. 34:281. [41] Painter, T. S. 1924. Ibid. 39:197. [42] Painter, T. S. 1925. Am. Naturalist 59:385. [43] Painter, T. S. 1925. Science 61:423. [44] Pinney, E. 1918. J. Morphol. 31:225. [45] Porter, K. R. 1941. Biol. Bull. 80:238. [46] Riley, G. M. 1938. Cytologia (Tokyo) 9:165. [47] Risley, P. L. 1936. Ibid. 7:232. [48] Risley, P. L. 1942. Anat. Record 84:513. [49] Rodolico, A. 1933. Pubbl. Staz. Zool. Napoli 13(2):180. [50] Saez, F. A., P. Rojas, and E. de Robertis. 1936. Inst. Museo Univ. Nacl. La Plata 2. [51] Saez, F. A., P. Rojas, and E. de Robertis. 1936. Z. Zellforsch. Mikroskop. Anat. 24:727. [52] Sasaki, M. S., and S. Makino. 1962. J. Heredity 53:157. [53] Schreiner, A., and K. E. Schreiner. 1904. Anat. Anz. 24:561. [54] Schreiner, A., and K. E. Schreiner. 1904. Arch. Biol. (Liege) 21:183. [55] Shioda, G., and M. S. Sasaki. 1962. Dobutsugaku Zasshi 71:98. [56] Suzuki, K. 1949. Nippon Idengaku Zasshi 24:90. [57] Svårdson, G. 1945. Medd. Statens Und. Foersoeeksanstalt Soetvatt.-Fisk. 23. [58] Svårdson, G., and T. Wickbom. 1939. Hereditas 25:472. [59] Thatcher, L. E. 1922. Science 56:372. [60] Tjio, J. H., and A. Levan. 1956. Hereditas 42:1. [61] Udagawa, T. 1954. Annotationes Zool. Japon. 27:91. [62] Unger, H. 1936. Z. Zellforsch. Mikroskop. Anat. 25:476. [63] Wickbom, T. 1945. Hereditas 31:241. [64] Wickbom, T. 1949. Ibid. 35:33. [65] Witschi, E. 1933. Cytologia (Tokyo) 4:174. [66] Yamashina, Y. 1943. J. Fac. Sci. Hokkaido Imp. Univ., VI, 8:307. [67] Yamashina, Y. 1944. Cytologia (Tokyo) 13:270. [68] Yamashina, Y. 1946. Seibutsu 1. [69] Yamashina, Y. 1951. Iden No Sogo Kenkyu 2. [70] Yamashina, Y., and S. Makino. 1946. Seibutsu 1.

### Part II. INVERTEBRATES

**Diploid** (column D): s = spermatogonium; o = oogonium; m = somatic cell. **Haploid** (column E):  $\sigma(I)$  = primary spermatocyte;  $\sigma(II)$  = secondary spermatocyte;  $\varphi(I)$  = primary oocyte;  $\varphi(II)$  = secondary oocyte.

	Class	Species	Common Name	Chromosome Number		Reference
				Diploid	Haploid	
	(A)	(B)	(C)	(D)	(E)	(F)
Chordata						
1	Cephalochordata <sup>1</sup>	<i>Branchiostoma lanceolatum</i>	Amphioxus	24o	12 $\varphi(I,II)$	10
2	Ascidacea	<i>Ciona intestinalis</i>	Sea squirt	18m	.....	5
Echinodermata						
3	Asteroidea	<i>Asterias forbesi</i>	Starfish	36m	18 $\sigma(I,II)$	23
4	Echinoidea	<i>Arbacia punctulata</i>	Sea urchin	ca. 40m	.....	68
5		<i>Echinavachmius parma</i>	Sand dollar	52m	.....	36
6	Holothuroidea	<i>Stichopus regalis</i>	Sea cucumber	28-36s	16-18 $\sigma(I)$	19
Arthropoda						
7	Arachnida	<i>Ixodes ricinus</i>	Sheep tick	28s	.....	46

<sup>1/1</sup> Subphylum.

continued

# 1. CHROMOSOME NUMBERS: ANIMALS

## Part II. INVERTEBRATES

	Class	Species	Common Name	Chromosome Number		Reference
				Diploid	Haploid	
(A)	(B)	(C)	(D)	(E)	(F)	
Arthropoda						
8	Arachnida	<i>Tegenaria domestica</i>	House spider	43s	23♂(I)	53
9	Merostomata	<i>Tachypleus tridentatus</i>	King crab	26s	13♂(I,II)	48
10	Crustacea	<i>Artemia salina</i>	Brine shrimp	42m	21♀(I,II)	1
11		<i>Astacus fluviatilis</i>	Crayfish	.....	ca. 58♂(I)	52
12		<i>Cyclops viridis</i>	Cyclops	12o	6♀(I)	63
13		<i>Daphnia magna</i>	Water flea	20s,m	10♀(I)	40
14		<i>Homarus sp.</i>	Lobster	.....	18♂(I)	31
15		<i>Lepas anatifera</i>	Goose barnacle	26s,o	13♂(I), 13♀(I)	75
16		<i>Potamon dehaami</i>	River crab	82s	41♂(I,II)	77
17	Insecta	<i>Aedes albopictus</i>	Mosquito	6s,o	.....	59
18		<i>Apis mellifera</i>	Honeybee	16o	16♂(I,II)	13,14
19		<i>Bombyx mori</i>	Silkworm	56s	28♂(I,II)	47
20		<i>Calliphora erythrocephala</i>	Bluebottle fly	12s,o	6♂(I,II)	24
21		<i>Cimex lectularius</i>	Bedbug	30-34s,33-41o	18-21♂(I)	11
22		<i>Ctenocephalides canis</i>	Dog flea	14♂m	.....	25,26
23		<i>Drosophila melanogaster</i>	Fruit fly	8s,o	4♀(I)	20
24		<i>Dytiscus marginalis</i>	Diving beetle	38s	19♂(I,II)	55
25		<i>Ephestia kuehniella</i>	Mediterranean flour moth	.....	30♂(I,II); 30♀(I); 29,30♀(II)	71
26		<i>Formica sanguinea</i>	Red ant	ca. 48m	24♀(I,II)	57
27		<i>Habrobracon juglandis</i>	Parasitic wasp	20o	10♀(I)	62
28		<i>Leptinotarsa signaticolis</i>	Potato beetle	34s	17♂(I); 16, 17♂(II)	74
29		<i>Locusta migratoria</i>	Migratory locust	23s	12♂(I); 11, 12♂(II)	76
30		<i>Magicicada septendecim</i>	Periodical cicada	19s,20o	10♂(I); 9,10♂(II)	58
31		<i>Mantis religiosa</i>	Praying mantis	27s	13♂(I); 13,14♂(II)	27
32		<i>Melanoplus differentialis</i>	Differential grasshopper	24♀m	12♀(I,II)	60
33		<i>Musca domestica</i>	Housefly	12s,o	6♂(I,II)	65
34		<i>Pediculus capitis</i>	Head louse	12m	.....	16
35		<i>Periplaneta americana</i>	American cockroach	33s, 34♀m	17♂(I); 16, 17♂(II)	66
36		<i>Pieris brassicae</i>	European cabbageworm	30s,o	15♂(I,II), 15♀(I)	15
37	<i>Popillia japonica</i>	Japanese beetle	18s	9♂(I,II)	78	
38	<i>Samia cynthia</i>	Cynthia moth	26s,o, ♀m	13♂(I,II), 13♀(I,II)	12	
39	<i>Tenebrio molitor</i>	Yellow mealworm	20s,o, ♀m	10♂(I,II)	64	
40	<i>Thermobia domestica</i>	Firebrat	34s, 36o	18♂(I); 16, 18♂(II)	49	
41	Onychophora	<i>Peripatus sp.</i>	Peripatus	28s,o,m	14♂(I,II)	39
Annelida						
42	Hirudinea	<i>Herpobdella bistrata</i>	Leech	18s	9♂(II)	73
43	Oligochaeta	<i>Enchytraeus humiculator</i>	White worm	32s,o	16♀(I)	69
44		<i>Lumbricus terrestris</i>	Earthworm	36m	.....	41
45	Polychaeta	<i>Nereis limbata</i>	Clam worm	20-30m	14♀(I)	3,4
Mollusca						
46	Cephalopoda	<i>Sepia officinalis</i>	Cuttlefish	.....	6♀(I)	35
47	Bivalvia	<i>Macra sp.</i>	Bar clam	24m	12♀(I,II)	29
48		<i>Unio sp.</i>	Pearl mussel	.....	16♀(I,II)	33
49	Gastropoda	<i>Aplysia limacina</i>	Sea hare	24m	.....	6,7
50		<i>Doris bifida</i>	Sea lemon	32m	16♀(I,II)	61
51		<i>Helix pomatia</i>	Land snail	54s	27♂(I), 27♀(I)	50
52		<i>Lymnaea japonica</i>	Freshwater snail	36s	18♂(I)	22
Aschelminthes						
53	Nematoda	<i>Ascaris lumbricoides</i>	Large roundworm	43s; 48o; 43, 48m	24♂(I); 19, 24♂(II); 24♀(I,II)	72
54		<i>Rhabditis sp.</i>	Free-living roundworm	13s; 14o; 13, 14m	7♂(I); 6, 7♂(II); 7♀(I,II)	30
55	Rotifera	<i>Asplanchna intermedia</i>	Rotifer	24s, ♂m	12♂(I)	67
Platyhelminthes						
56	Cestoda	<i>Taenia pisiformis</i>	Dog tapeworm	.....	13-15♀(I)	70
57	Trematoda	<i>Fasciola hepatica</i>	Liver fluke	12o,m	6♀(I,II)	56

continued



## 1. CHROMOSOME NUMBERS: ANIMALS

### Part II. INVERTEBRATES

Class	Species	Common Name	Chromosome Number		Reference	
			Diploid	Haploid		
(A)	(B)	(C)	(D)	(E)	(F)	
Platyhelminthes						
58	Trematoda	<i>Schistosoma haematobium</i>	Human blood fluke	14s	8♂(I); 6, 8♂(II); 8♀(II)	34
59	Turbellaria	<i>Planaria torva</i>	Flatworm	16m	8♀(I,II)	37,38
Cnidaria						
60	Scyphozoa	<i>Aurelia flavedula</i>	Scyphomedusa	18-20m	9-10♀(I)	21
61	Hydrozoa	<i>Gonionemus murbachii</i>	Hydromedusa	24-25s; ca. 24o,m	ca. 12♂(I,II)	2
62		<i>Hydra vulgaris attenuata</i>	Freshwater hydra	32s,o,m	16♂(I)	45
63		<i>Obelia geniculata</i>	Marine hydra	34o	17♀(I)	18
Porifera						
64	Desmospongiae	<i>Spongilla lacustra</i>	Freshwater sponge	10-12m	.....	42,43
65	Calcarea	<i>Scypha ciliatum</i>	Marine sponge	26m	13♀(I)	17
Protozoa						
66	Ciliata	<i>Didinium nasutum</i>	Carnivorous ciliate	16	8	51
67		<i>Stentor coeruleus</i>	Heterotrichous ciliate	28	14	44
68	Rhizopoda	<i>Amoeba proteus</i>	Free-living amoeba		ca. 50 <sup>a</sup>	8
69		<i>Entamoeba histolytica</i>	Parasitic amoeba		6 <sup>a</sup>	28
70	Mastigophora	<i>Euglena gracilis</i>	Green flagellate		ca. 45 <sup>a</sup>	32
71		<i>Trypanosoma equiperdum</i>	Trypanosome		3 <sup>a</sup>	54
72		<i>Volvox globator</i>	Pale-green flagellate	.....	5	9

/a/ Uncertain whether diploid or haploid.

Contributor: Makino, Sajiro

References: [1] Artom, C. 1928. Compt. Rend. Soc. Biol. 99(Suppl.):29. [2] Bigelow, H. B. 1907. Bull. Museum Comp. Zool. Harvard Univ. 48:287. [3] Bonnevie, K. 1907. Biol. Bull. 13:57. [4] Bonnevie, K. 1908. Arch. Zellforsch. 2:201. [5] Boveri, T. 1890. Jena. Z. Naturw. 24. [6] Carazzi, D. 1905. Arch. Ital. Anat. Embriol. 4:231. [7] Carazzi, D. 1905. Ibid. 4:459. [8] Carter, L. 1919. Proc. Roy. Phys. Soc. Edinburgh 20:193. [9] Cave, M. S., and M. A. Pocock. 1951. Am. J. Botany 38:800. [10] Cerfontaine, P. 1906-07. Arch. Biol. (Liege) 22:229. [11] Darlington, C. D. 1939. J. Genet. 39:101. [12] Dederer, P. H. 1928. J. Morphol. 45:599. [13] Doncaster, L. 1906. Anat. Anz. 29:490. [14] Doncaster, L. 1907. Ibid. 31:168. [15] Doncaster, L. 1912. Proc. Cambridge Phil. Soc. 16:491. [16] Doncaster, L., and H. G. Cannon. 1920. Quart. J. Microscop. Sci. 64:303. [17] Dubosq, O., and O. Tuzet. 1937. Arch. Zool. Exptl. Gen. 79(2):157. [18] Faulkner, G. H. 1929. Quart. J. Microscop. Sci. 73:225. [19] Field, G. W. 1895. J. Morphol. 11:235. [20] Guyénot, E., and A. Naville. 1929. Cellule Rec. Cytol. Histol. 39:25. [21] Hargitt, G. T. 1910. J. Morphol. 21:593. [22] Inaba, A. 1950. Nippon Idengaku Zasshi 25:222. [23] Jordan, H. E. 1908. Papers Tortugas Lab. Carnegie Inst. Wash. 1:1. [24] Keuneke, W. 1924. Z. Zellforsch. Mikroskop. Anat. 1:357. [25] Kichijo, H. 1941. Botany Zool. (Tokyo) 9. [26] Kichijo, H. 1941. Nippon Idengaku Zasshi 17:122. [27] King, R. L. 1931. J. Morphol. 52:525. [28] Kofoid, C. A., and O. Swezy. 1925. Univ. Calif. (Berkeley) Publ. Zool. 26:331. [29] Kostanecki, K. 1911. Arch. Mikroskop. Anat. Entwicklungsmech. 78(2):1. [30] Kröning, F. 1923. Arch. Zellforsch. 17:63. [31] Labbé, A. 1904. Compt. Rend. 138:96. [32] Leedale, G. F. 1958. Nature 181:502. [33] Lilje, F. R. 1901. J. Morphol. 17:227. [34] Lindner, E. 1914. Arch. Zellforsch. 12:516. [35] Loyez, M. 1906. Arch. Anat. Microscop. Morphol. Exptl. 8:69. [36] Matsui, K. 1924. J. Coll. Agr. Imp. Univ. Tokyo 7:211. [37] Mattiesen, E. 1903. Zool. Anz. 27:81. [38] Mattiesen, E. 1904. Z. Wiss. Zool. 77:274. [39] Montgomery, T. H., Jr. 1900. Zool. Jahrb. 14:277. [40] Mortimer, C. H. 1935. Naturwissenschaften 23:476. [41] Muldal, S. 1949. John Innes Hort. Inst. Ann. Rept. 39:21. [42] Müller, K. 1911. Arch. Entwicklungsmech. Organ. 32:397. [43] Müller, K. 1911. Ibid. 32:557. [44] Mulsow, W. 1913. Arch. Protistenk. 28:363. [45] Niiyama, H. 1944. Cytologia (Tokyo) 13:204.

continued

## 1. CHROMOSOME NUMBERS: ANIMALS

### Part II. INVERTEBRATES

- [46] Nordenskiöld, E. 1920. Parasitology 12:159. [47] Oguma, K. 1919. Dobutsugaku Zasshi 31. [48] Okada, A. 1938. J. Sci. Hiroshima Univ., B(1), 6:37. [49] Perrot, J. L. 1933. Z. Zellforsch. Mikroskop. Anat. 18:573. [50] Perrot, J. L., and M. Perrot. 1938. Compt. Rend. 207:1005. [51] Prandtl, H. 1906. Arch. Protistenk. 7:229. [52] Prowazek, S. 1902. Z. Wiss. Zool. 71:445. [53] Revell, S. H. 1947. Heredity 1:337. [54] Roskin, G., and S. Schischliawewa. 1928. Arch. Protistenk. 60:460. [55] Schäfer, F. 1907. Zool. Jahrb. 23:535. [56] Schellenberg, A. 1911. Arch. Zellforsch. 6:443. [57] Schleip, W. 1908. Zool. Jahrb. 26:651. [58] Shaffer, E. L. 1920. Biol. Bull. 38:83. [59] Sinoto, Y., and K. Suzuki. 1943. Igaku To Seibutsugaku 3:175. [60] Slifer, E. H., and R. L. King. 1934. J. Morphol. 56:593. [61] Smallwood, W. M. 1905. Morphol. Jahrb. 33:87. [62] Speicher, K. G., and B. R. Speicher. 1938. Biol. Bull. 74:247. [63] Stella, E. 1931. Intern. Ges. Hydrobiol. Hydrog. 26:112. [64] Stevens, N. M. 1906. Carnegie Inst. Wash. Publ. 36(2). [65] Stevens, N. M. 1908. J. Exptl. Zool. 5:453. [66] Suomalainen, E. 1946. Ann. Acad. Sci. Fennicae, A(4), 10. [67] Tauson, A. 1927. Z. Zellforsch. Mikroskop. Anat. 4:652. [68] Tennent, D. H. 1912. J. Exptl. Zool. 12:391. [69] Vejdovsky, F. 1907. Sitzber. Kgl. Boehm. Ges. Wiss. Math. Naturw. Kl. (1). [70] Von Janicki, C. 1907. Z. Wiss. Zool. 87:685. [71] Wagner, H. O. 1930. Z. Zellforsch. Mikroskop. Anat. 12:749. [72] Walton, A. C. 1924. Ibid. 1:167. [73] Wendrowsky, V. 1928. Ibid. 8:153. [74] Wieman, H. L. 1910. J. Morphol. 21:135. [75] Witschi, E. 1935. Biol. Bull. 68:263. [76] Wu, J. S. 1938. Cytologia (Tokyo) 9:334. [77] Yanagita, T. 1944. Sci. Rept. Tokyo Bunrika Daigaku, B, 6. [78] Yosida, T. 1949. Trans. Sapporo Nat. Hist. Soc. 18:43.

## 2. CHROMOSOME NUMBERS: PLANTS

For information on additional species of plants, consult reference 2, Part I.

### Part I. NONVASCULAR

Many of the chromosome numbers are of doubtful accuracy since the small size of the chromosomes makes it difficult to determine exact counts.

Class and Species	Haploid Number	Reference	Class and Species	Haploid Number	Reference
(A)	(B)	(C)	(A)	(B)	(C)
<b>Myxophyta</b>					
Myxomyceteae					
1	<i>Comatricha nigra</i>	ca. 30	15	<i>Venturia inaequalis</i>	4-6
2	<i>Physarum polycephalum</i>	ca. 90	16		7
Acrasieae					
3	<i>Dictyostelium discoideum</i>	7	17	<i>Agaricus campestris</i>	4
Fungi					
Phycomycetes					
4	<i>Phycomyces nitens</i>	2	18		9
5		ca. 12	19		12
6	<i>Saprolegnia ferax</i>	7 or more	20	<i>Lycoperdon piriforme</i>	2
Ascomycetes					
7	<i>Aspergillus niger</i>	2	21		6
8	<i>Neurospora crassa</i>	7	22	<i>Panus torulosus</i>	6
9		9	23	<i>Puccinia graminis</i>	2
10	<i>Penicillium</i> sp.	2	24		6
11	<i>Peziza vesiculosa</i>	8	25	<i>Ustilago hordei</i>	2
12	<i>Saccharomyces cerevisiae</i>	2	Lichenes		
13		4	Ascolichenes		
14	<i>Schizosaccharomyces octosporus</i>	4	26	<i>Cladonia cristatella</i>	4
			27	<i>Dermatocarpon fluviatile</i>	6-8
			28	<i>Lecanora dispersa</i>	3
			29	<i>Lecidea crustulata</i>	2
			Algae		
			Chrysochyta		
			30	<i>Vaucheria sessilis</i>	7-10

/1/ Division.

continued

## 2. CHROMOSOME NUMBERS: PLANTS

### Part I. NONVASCULAR

	Class and Species	Haploid Number	Reference		Class and Species	Haploid Number	Reference
	(A)	(B)	(C)		(A)	(B)	(C)
	Algae				Bryophyta		
	Chlorophyta <sup>1</sup>				Musci		
31	<i>Acetabularia wettsteinii</i>	ca. 10	57	47	<i>Hylocomium splendens</i>	10	68
32	<i>Chlamydomonas moewusii</i>	8	8	48		11	69
33		36±2	55	49		12	79
34	<i>Cladophora glomerata</i>	32	36	50	<i>Hypnum cupressiforme</i>	10	68
35		48	56,59-61	51		16	3
36	<i>Oedogonium</i> spp.	9,13,17-19	67	52	<i>Mnium undulatum</i>	6	24,27
37	<i>Spirogyra majuscula</i>	34-36	16	53		7	42
38	<i>Ulothrix zonata</i>	4	17,58	54	<i>Polytrichum juniperinum</i>	7	3,34,80
39		10	52,53	55	<i>Sphagnum girgensohnii</i>	23	26
	Phaeophyta <sup>1</sup>				Hepaticae		
40	<i>Ectocarpus siliculosus</i>	8	62	56	<i>Marchantia polymorpha</i>	9-11	20,21
41		8,9	46	57	<i>Riccia fluitans</i>	8	7,66
42	<i>Fucus vesiculosus</i>	10	14	58		16	38
43		32	78		Anthocerotae		
44	<i>Laminaria digitata</i>	27-31	45	59	<i>Anthoceros laevis</i>	4	11,33
	Rhodophyta <sup>1</sup>			60		5	47
45	<i>Polysiphonia nigrescens</i>	30	4	61		6	48,65
46	<i>Porphyrta umbilicalis</i>	5	32	62		8	37

<sup>1</sup>/ Division.

**Contributors:** (a) Olive, Lindsay S., (b) Cave, Marion S., (c) Anderson, Lewis E., (d) Ahmadjian, V.

**References:** [1] Ahmadjian, V. Unpublished. Clark Univ., Worcester, Mass., 1963. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Anderson, L. E., and H. Crum. 1958. Bull. Natl. Museum Can. Contrib. Botany 160:1. [4] Austin, A. P. 1956. Nature 178:370. [5] Backus, E. J., and G. W. Keitt. 1940. Bull. Torrey Bot. Club 67:765. [6] Badian, J. 1937. Bull. Intern. Acad. Polon. Sci., B, 5:61. [7] Berrie, G. K. 1958. Ph.D. Thesis. London Univ., England. [8] Buffaloe, N. D. 1958. Bull. Torrey Bot. Club 85:157. [9] Burgeff, H. 1915. Flora (Jena) 108:353. [10] Colson, B. 1935. Ann. Botany (London) 49:1. [11] Davis, B. M. 1899. Bot. Gaz. 28:89. [12] Day, P. R., D. M. Boone, and G. W. Keitt. 1956. Am. J. Botany 43:835. [13] Elisei, F. G. 1939. Ist. Botan. Univ. Crittogam. Pavia Atti, Ser. 4, 11:13. [14] Farmer, J. B., and J. L. Williams. 1896. Ann. Botany (London) 10:479. [15] Ganeson, A. T. 1959. Compt. Rend. Trav. Lab. Carlsberg 31:149. [16] Geitler, L. 1935. Ber. Deut. Botan. Ges. 53:270. [17] Gross, I. 1931. Arch. Protistenk. 73:206. [18] Guilliermond, A. 1904. Rev. Gen. Botan. 16:129. [19] Hanatschek, H. 1932. Arch. Protistenk. 78:497. [20] Haupt, G. 1932. Z. Induktive Abstammungs- Vererbungslehre 62:367. [21] Haupt, G. 1933. Ibid. 63:390. [22] Heim, P. 1952. Rev. Mycol. 17:3. [23] Heim, P. 1954. Ibid. 19:201. [24] Heitz, E. 1942. Arch. Julius Klaus-Stift. Vererbungsforsch., Sozialanthropol. Rassenhyg. 17:444. [25] Höhnk, W. 1935. Naturw. Ver. Bremen 29:308. [26] Holmen, K. 1955. Bot. Tidskr. 52:37. [27] Holmen, K. 1958. Ibid. 54:23. [28] Hughes, D. T. 1961. Nature 190:285. [29] Hüttig, W. 1931. Z. Botan. 24:529. [30] Julian, J. B. 1958. Can. J. Botany 36:607. [31] Kharbush, S. S. 1927. Ann. Sci. Nat. Botan. Biol. Vegetale 10:285. [32] Krishnamurthy, V. 1959. Ann. Botany (London), N.S. 23:147. [33] Lander, C. A. 1935. Am. J. Botany 22:42. [34] Lewis, K. R. 1957. Trans. Brit. Bryol. Soc. 3:279. [35] Lindegren, C. C., and S. Rumann. 1938. J. Genet. 36:395. [36] List, H. 1930. Arch. Protistenk. 72:453. [37] Lorbeer, G. 1924. Ber. Deut. Botan. Ges. 42:231. [38] Lorbeer, G. 1934. Jahrb. Wiss. Botan. 80:567. [39] McClintock, B. 1945. Am. J. Botany 32:671. [40] McGinnis, R. C. 1956. J. Heredity 47:255. [41] Maire, R. 1902. Thesis. Paris. [42] Mazzeo, M. 1941. Nuovo Giorn. Botan. Ital. 48:613. [43] Moreau, F. 1913. Thesis. Paris. [44] Moreau, F., and F. Moreau. 1931. Rev. Gen. Botan. 43:465. [45] Naylor, M. 1956. Ann. Botany (London), N.S. 20:431. [46] Papenfuss, G. F. 1935. Bot. Gaz. 96:421.

*continued*



## 2. CHROMOSOME NUMBERS: PLANTS

### Part I. NONVASCULAR

- [47] Proskauer, J. 1958. In L. Rabenhorst, ed. Kryptogamen-Flora. Akademische Verlagsgesellschaft, Leipzig. v. 6, pp. 1303-1319. [48] Rink, W. 1935. Flora (Jena) 130:87. [49] Ross, I. K. 1961. Am. J. Botany 48:244. [50] Sappin-Trouffy, P. 1896. Botaniste 5:59. [51] Sarazin, A. 1938. Compt. Rend. 206:275. [52] Sarma, Y. S. R. K. 1957. Nature 180:46. [53] Sarma, Y. S. R. K. 1958. Brit. Phycol. Bull. 6:22. [54] Sass, J. E. 1928. Papers Mich. Acad. Sci. 9:287. [55] Schaechter, M., and E. D. DeLamater. 1955. Am. J. Botany 42:417. [56] Schussnig, B. 1928. Oesterr. Botan. Z. 77:62. [57] Schussnig, B. 1929. Ber. Deut. Botan. Ges. 47:266. [58] Schussnig, B. 1930. Z. Zellforsch. Mikroskop. Anat. 10:642. [59] Schussnig, B. 1944. Ber. Deut. Botan. Ges. 62:5. [60] Schussnig, B. 1951. Svensk Botan. Tidskr. 45:597. [61] Schussnig, B. 1954. Arch. Protistenk. 100:287. [62] Schussnig, B., and E. Kothbauer. 1934. Oesterr. Botan. Z. 83:81. [63] Singleton, J. R. 1953. Am. J. Botany 40:124. [64] Stevens, R. B. 1941. Ibid. 28:59. [65] Tatuno, S. 1941. J. Sci. Hiroshima Univ., B(2), 4:73. [66] Tatuno, S. 1957. Ibid., B(2), 8:81. [67] Tschermak, E. 1944. Chromosoma 2:493. [68] Vaarama, A. 1950. Botan. Notiser, p. 239. [69] Vaarama, A. 1953. Bryologist 56:169. [70] Von Stosch, H. A. 1937. Ber. Deut. Botan. Ges. 55:362. [71] Wakayama, K. 1931. Cytologia (Tokyo) 2:291. [72] Wang, D. T. 1932. Compt. Rend. 195:1041. [73] Wang, D. T. 1934. Botaniste 26:539. [74] Whelden, R. M. 1940. Mycologia 32:630. [75] Widra, A., and E. D. DeLamater. 1955. Am. J. Botany 42:423. [76] Wilson, C. M. 1952. Proc. Natl. Acad. Sci. U.S. 38:659. [77] Wilson, C. M., and I. K. Ross. 1957. Am. J. Botany 44:345. [78] Yamanouchi, S. 1909. Botan. Gaz. 47:173. [79] Yano, K. 1957. Mem. Takada Branch Niigata Univ. 1:85. [80] Yano, K. 1957. Ibid. 1:129.

### Part II. VASCULAR

For additional information on gymnosperms and angiosperms, consult reference 18.

Species (Common Name)	Diploid Number	Refer- ence	Species (Common Name)	Diploid Number	Refer- ence
(A)	(B)	(C)	(A)	(B)	(C)
Pteridophyta			17	<i>Tsuga canadensis</i> (eastern hemlock)	24 67
1	<i>Adiantum pedatum</i> (American maidenhair)	58 48	Angiospermae (Monocotyledoneae)		
2	<i>Equisetum arvense</i> (field horsetail)	216 8	18	<i>Allium cepa</i> (garden onion)	16,32 11
3	<i>Lycopodium clavatum</i> (club moss)	68 8	19	<i>Asparagus officinalis</i> (garden asparagus)	20 55
4	<i>Polypodium virginianum</i> (rock polypody)	74,148 8,48	20	<i>Avena sativa</i> (common oat)	42 22
5	<i>Selaginella selaginoides</i> (spike moss)	18 8	21	<i>Elodea canadensis</i> (Canada waterweed)	48 65
Gymnospermae			22	<i>Gladiolus</i> spp. <sup>1</sup> (gladiolus)	30,45,60,75 2
6	<i>Abies concolor</i> (white fir)	24 67	23	<i>Hordeum vulgare</i> (barley)	14 42
7	<i>Cupressus sempervirens</i> (Italian cypress)	22 51	24	<i>Iris versicolor</i> (blue-flag iris)	72,84,105 68
8	<i>Ginkgo biloba</i> (ginkgo)	24 45	25	<i>Lilium</i> spp. (lily)	24 29,64
9	<i>Juniperus virginiana</i> (eastern red cedar)	22,33 74	26	<i>Oryza sativa</i> (rice)	24 1
10	<i>Larix occidentalis</i> (western larch)	24 67	27	<i>Phleum pratense</i> (timothy)	42 54
11	<i>Picea glauca</i> (white spruce)	24 67	28	<i>Phoenix dactylifera</i> (date palm)	36 4
12	<i>Pinus palustris</i> (longleaf pine)	24 49	29	<i>Poa pratensis</i> (Kentucky bluegrass)	36-123 56
13	<i>Sequoia gigantea</i> (giant sequoia)	22 39	30	<i>Tridactylis virginiana</i> (Virginia spiderwort)	38-96 40
14	<i>Taxodium distichum</i> (bald cypress)	22 73	31	<i>Triticum aestivum</i> (wheat)	24 14
15	<i>Taxus baccata</i> (English yew)	24 12	32	<i>Yucca</i> spp. (yucca)	42 61
16	<i>Thuja occidentalis</i> (northern white cedar)	22 67	33	<i>Zea mays</i> (corn)	60 72,82
			34		20 60
			Angiospermae (Dicotyledoneae)		
			35	<i>Acer saccharinum</i> (silver maple)	52 77

/1/ Cultivated.

continued



## 2. CHROMOSOME NUMBERS: PLANTS

### Part II. VASCULAR

Species (Common Name)			Diploid Number	Refer- ence	Species (Common Name)			Diploid Number	Refer- ence
(A)			(B)	(C)	(A)			(B)	(C)
Angiospermae (Dicotyledoneae)									
36	<i>Alnus rubra</i> (red alder)	28	83	61	<i>Juglans nigra</i> (black walnut)	32	85		
37	<i>Antirrhinum majus</i> (snap- dragon)	16	59	62	<i>Lactuca sativa</i> (lettuce)	18	79		
38	<i>Beta vulgaris</i> (common beet)	18	47,84	63	<i>Lycopersicon esculentum</i> (tomato)	24	3		
39	<i>Betula lenta</i> (sweet birch)	28	86	64	<i>Magnolia</i> spp. (magnolia)	38,76,114	34		
40	<i>Capsicum frutescens</i> (bush red pepper)	24	69	65	<i>Malus pumila</i> (common apple)	34,51	16		
41	<i>Carya tomentosa</i> (mockernut hickory)	64	85	66	<i>Medicago sativa</i> (alfalfa)	16	6		
42	<i>Catalpa speciosa</i> (northern catalpa)	40	70	67		32,64	81		
43	<i>Chrysanthemum maximum</i> (Pyrenees chrysanthemum)	85,90,126,148, 154,160,171	21	68	<i>Nicotiana tabacum</i> (common tobacco)	48	28		
44	<i>Cinchona ledgeriana</i> (ledger- bark cinchona)	34	19	69	<i>Oenothera biennis</i> (common evening primrose)	14	16		
45	<i>Citrus limon</i> (lemon)	18,36	44	70	<i>Pastinaca sativa</i> (parsnip)	22	57		
46	<i>C. sinensis</i> (sweet orange)	45	44	71	<i>Persea americana</i> (American avocado)	24	7		
47	<i>Cornus florida</i> (flowering dogwood)	22	20	72	<i>Phaseolus vulgaris</i> (kidney bean)	22	78		
48	<i>Cucumis sativus</i> (cucumber)	14	30	73	<i>Phlox</i> spp. (phlox)	14,21,28	24,25,53		
49	<i>Cucurbita pepo</i> (pumpkin)	40	10,23	74	<i>Pisum sativum</i> (garden pea)	14	63		
50	<i>Daucus carota</i> (carrot)	18	26,31	75	<i>Populus tremuloides</i> (quaking aspens)	38	71		
51	<i>Digitalis purpurea</i> (common foxglove)	56	9	76	<i>Prunus amygdalus</i> (almond)	16	15		
52	<i>Fagopyrum esculentum</i> (buckwheat)	16	36	77	<i>P. domestica</i> (garden plum)	48	15		
53	<i>Fagus sylvatica</i> (European beech)	24	38	78	<i>P. persica</i> (peach)	16	15		
54	<i>Fragaria virginiana</i> (Virgin- ia strawberry)	56	33	79	<i>Pyrus communis</i> (pear)	34,51	17		
55	<i>Fraxinus americana</i> (white ash)	46,92,138	87	80	<i>Quercus alba</i> (white oak)	24	75		
56	<i>Glycine soja</i> (soybean)	40	62	81	<i>Raphanus sativus</i> (garden radish)	18	41		
57	<i>Gossypium hirsutum</i> (upland cotton)	52	89	82	<i>Rheum officinale</i> (medicinal rhubarb)	22	37		
58	<i>Helianthus annuus</i> (common sunflower)	34	27	83	<i>Rhododendron</i> spp. (rhodo- dendron)	26,39,52,78, 104,156	35		
59	<i>Ilex aquifolium</i> (English holly)	40	50	84	<i>Ribes</i> spp. (currant)	16	13,52,90		
60	<i>Ipomoea batatas</i> (sweet po- tato)	90	80	85	<i>Rosa</i> spp. <sup>1</sup> (rose)	14,21,28	88		
				86	<i>Salix alba</i> (white willow)	76	5		
				87	<i>Solanum tuberosum</i> (potato)	48	76		
				88	<i>Trifolium pratense</i> (red clover)	14	46		
				89	<i>Ulmus americana</i> (American elm)	28	43		
				90		56	66		
				91	<i>Vicia faba</i> (broad bean)	12,14	32		
				92	<i>Vitis vinifera</i> (European grape)	38,57,76	58		

/1/ Cultivated.

Contributors: (a) Darlington, C. D., (b) Morton, C. V., (c) Sax, Karl

References: [1] Avdulov, N. P. 1931. Bull. Appl. Botany Genet. Plant Breeding (USSR), C, 43. [2] Bamford, R. 1935. J. Agr. Res. 51:945. [3] Barton, D. W. 1950. Am. J. Botany 37:639. [4] Beal, J. M. 1937. Botan. Gaz. 99:400. [5] Blackburn, K. B., and J. W. H. Harrison. 1924. Ann. Botany (London) 38:361. [6] Bolton, J. L., and J. E. R. Greenshields. 1950. Science 112:275. [7] Bowden, W. M. 1940. Chronica Botan. (Leiden) 6:123. [8] Britton, D. M. 1953. Am. J. Botany 40:575. [9] Buxton, B. H., and W. C. F. Newton. 1928. J. Genet. 19:269. [10] Castetter, E. F. 1930. Am. J. Botany 17:41. [11] D'Amato, F. 1948. Caryologia 1:48. [12] Dark, S. O. S. 1932. Ann. Botany (London) 46:965. [13] Darlington, C. D. 1929. Genetica (Haag) 11:267. [14] Darlington, C. D. 1929. J. Genet. 21:207. [15] Darlington, C. D. 1930. Ibid. 22:65. [16] Darlington, C. D. 1931. Ibid. 24:405. [17] Darlington, C. D., and A. A. Moffett. 1930. Ibid. 22:129. [18] Darlington, C. D., and A. P. Wylie. 1955.

continued

## 2. CHROMOSOME NUMBERS: PLANTS

### Part II. VASCULAR

- Chromosome atlas of flowering plants. Allen and Unwin, London. [19] Dawson, R. F. 1948. *Lloydia* 11:81.
- [20] Dermen, H. 1932. *J. Arnold Arboretum Harvard Univ.* 13:410. [21] Dowrick, G. J. 1952. *Heredity* 6:365.
- [22] Emme, H. 1930. *Zuechter* 2:65. [23] Erwin, A. T., and E. S. Haber. 1930. *Iowa Agr. Expt. Sta. Bull.* 263:343. [24] Flory, W. S. 1934. *Cytologia (Tokyo)* 6:1. [25] Flory, W. S. 1937. *Ibid.*, Fujii Jubilaei Vol. (1):171. [26] Gardé, A., and N. Gardé. 1951. *Genet. Iberica* 3:23. [27] Geisler, F. 1931. *Butler Univ. Botan. Studies* 2:53. [28] Goodspeed, T. H. 1945. *Univ. Calif. (Berkeley) Publ. Botany* 18:335. [29] Goodspeed, T. H., F. M. Uber, and P. Avery. 1935. *Ibid.* 18:33. [30] Heimlich, L. F. 1927. *Proc. Natl. Acad. Sci. U. S.* 13:113. [31] Heiser, C. B., and T. W. Whitaker. 1948. *Am. J. Botany* 35:179. [32] Hirayoshi, I., and M. Matsumura. 1952. *Japan. J. Breeding* 1:219. [33] Ichijima, K. 1926. *Genetics* 11:590. [34] Janaki-Ammal, E. K. Unpublished. *India Botanical Survey, Calcutta*, 1953. [35] Janaki-Ammal, E. K., I. C. Enoch, and M. Bridgewater. 1950. *Rhododendron Camellia Yearbook (London)* 5:78. [36] Jaretzky, R. 1927. *Ber. Deut. Botan. Ges.* 45:48. [37] Jaretzky, R. 1928. *Jahrb. Wiss. Botan.* 69:357. [38] Jaretzky, R. 1930. *Planta* 10:120. [39] Jensen, H., and A. Levan. 1941. *Hereditas* 27:220. [40] Juhl, H. 1952. *Flora (Jena)* 139:462. [41] Karpechenko, G. D. 1924. *Bull. Appl. Botany Plant Breeding (Leningrad)* 13:4. [42] Kihara, H. 1924. *Mem. Coll. Sci. Kyoto Imp. Univ.* 1:1. [43] Krause, O. 1930. *Ber. Deut. Botan. Ges.* 48:9. [44] Krug, C. A. 1943. *Botan. Gaz.* 104:602. [45] Lee, C. L. 1954. *Am. J. Botany* 41:545. [46] Levan, A. 1942. *Hereditas* 28:245. [47] Levan, A. 1942. *Ibid.* 28:345. [48] Manton, I. 1950. *Problems of cytology and evolution in the Pteridophyta*. Cambridge Univ. Press, London. [49] Mathews, A. C. 1932. *J. Elisha Mitchell Sci. Soc.* 48:101. [50] Maude, P. F. 1940. *New Phytologist* 39:17. [51] Mehra, P. N., and T. N. Khoshoo. 1948. *Proc. 34th Indian Sci. Congr.* 1947, (3):167. [52] Meurman, O. 1928. *Hereditas* 11:289. [53] Meyer, J. R. 1944. *Genetics* 29:199. [54] Myers, W. M. 1944. *J. Agr. Res.* 68:21. [55] Nagao, S. 1938. *Comment. Papers Agron. Akemine (Japan.)* [56] Nissen, Ø. 1950. *Agron. J.* 42:136. [57] Ogawa, K. 1929. *Mem. Coll. Sci. Kyoto Imp. Univ.* 4:309. [58] Olmo, H. P. 1937. *Cytologia (Tokyo)*, Fujii Jubilaei Vol. (1):606. [59] Propach, H. 1935. *Planta* 23:349. [60] Rhoades, M. M. 1950. *J. Heredity* 41:58. [61] Sachs, L. 1953. *J. Agr. Sci.* 43:204. [62] Sakai, B. 1951. *Kromosomo (Tokyo)* 11:425. [63] Sansome, E. R. 1933. *Cytologia (Tokyo)* 5:15. [64] Sansome, E. R., and L. La Cour. 1934. *Lily Yearbook*, p. 40. [65] Santos, J. K. 1924. *Botan. Gaz.* 77:353. [66] Sax, K. 1933. *J. Arnold Arboretum Harvard Univ.* 14:82. [67] Sax, K., and H. J. Sax. 1933. *Ibid.* 14:356. [68] Simonet, M. 1934. *Ann. Sci. Nat. Botan. Biol. Vegetale, Ser. 10*, 16:229. [69] Sinha, N. P. 1950. *Indian J. Genet. Plant Breeding* 10:36. [70] Smith, E. C. 1941. *J. Arnold Arboretum Harvard Univ.* 22:219. [71] Smith, E. C. 1943. *Ibid.* 24:275. [72] Snoad, B. 1952. *Rept. John Innes Hort. Inst.* 42:47. [73] Stebbins, G. L. 1948. *Science* 108:5. [74] Stiff, M. L. 1951. *Virginia J. Sci., N.S.* 2:317. [75] Sugiura, T. 1931. *Botan. Mag. (Tokyo)* 45:353. [76] Swaminathan, M. S. 1954. *Genetics* 39:59. [77] Taylor, W. R. 1920. *Contrib. Botan. Lab. Univ. Penna.* 5:111. [78] Thomas, P. T. Unpublished, 1955. [79] Thompson, R. C., T. W. Whitaker, and W. F. Kosar. 1941. *J. Agr. Res.* 63:91. [80] Ting, Y. C., and A. E. Kehr. 1953. *J. Heredity* 44:207. [81] Tomé, G. A. 1947. *Rev. Fac. Agron. Vet. Univ. Buenos Aires* 11:299. [82] Watkins, G. M. 1936. *Am. J. Botany* 23:328. [83] Wetzell, R. 1929. *Dissertation. Marburg Univ., Germany.* [84] Winge, Ø. 1917. *Compt. Rend. Trav. Lab. Carlsberg* 13:131. [85] Woodworth, R. H. 1930. *Am. J. Botany* 17:863. [86] Woodworth, R. H. 1931. *J. Arnold Arboretum Harvard Univ.* 12:206. [87] Wright, J. W. 1944. *J. Forestry* 42:489. [88] Wylie, A. P. 1954. *Am. Rose Ann.* 39:36. [89] Zaitzew, G. S. 1927. *Bull. Appl. Botany Genet. Plant Breeding (USSR)* 18:1. [90] Zielinski, Q. B. 1953. *Botan. Gaz.* 114:265.

# Contrails

## 3. SEX LINKAGE: MAN

For additional information, consult reference 23.

	Mutation	Phenotypic Expression			Reference
		Hemizygote XY	Heterozygote XX	Homozygote XX	
	(A)	(B)	(C)	(D)	(E)
1	Agammaglobulinemia	Gamma globulin absent	Absent	Unknown	18
2	Albinism, ocular	Melanin absent from eye	Mosaic pigmentary pattern of fundus oculi	Unknown	39
3	Aldrich syndrome	Eczema, thrombocytopenia	Absent	Unknown	1
4	Amelogenesis imperfecta	Defective enamel of teeth	Defective enamel in some families	Unknown	15,44
5	Anemia, hypochromic (Rundles, Falls)	Anemia	Splenomegaly, minor red cell change	Unknown	30
6	Angiokeratoma, diffuse	Dermal, vascular, neural, and renal lesions	Mild involvement	Unknown	43
7	Atrophy, peroneal	Atrophy of calf muscles	Occasional manifestation	Unknown	2
8	Blood group Xg <sup>a</sup>	Erythrocytes agglutinate with antiserum	Erythrocytes agglutinate with antiserum	Erythrocytes agglutinate with antiserum	24
9	Choroideremia	Night blindness, constricted visual fields, blindness	Depigmented retina	Unknown	21
10	Color blindness, partial Deutan series	Red blindness	Defect in only a few	Red blindness	10
11	Protan series	Green blindness	Mild defect revealed by special tests	Green blindness	10
12	Deaf-mutism	Profound deafness at birth	Absent	Profound deafness at birth	26,32
13	Diabetes insipidus, nephrogenic	High urinary output unaffected by pitressin	Slight increase in urinary output	Unknown	40
14	Dysplasia Anhidrotic ectodermal	Widespread ectodermal defects	Patchy defect	Unknown	9
15	Spondylo-epiphyseal	Dwarfism, with changes especially of spine and hips	Absent	Unknown	17
16	Dystrophy Macular	Loss of central vision	Absent	Unknown	34
17	Muscular (Duchenne)	Progressive atrophy of muscles	Mild serum enzymatic changes	Unknown	7,36
18	Glucose-6-PO <sub>4</sub> dehydrogenase deficiency	Hemolytic anemia with drugs	Absent	Hemolytic anemia with drugs	6
19	Hemeralopia	Night blindness with myopia	Absent	Unknown	34
20	Hemophilia Classical <sup>1</sup>	Severe bleeder	Absent	Severe bleeder	5,16,25,33
21	Mild <sup>2</sup>	Mild bleeder	Slight occasional manifestation	Unknown	14
22	Hurler syndrome	Dwarfism, mucopolysaccharide deposits, mental deficiency	Absent	Unknown	22
23	Hydrocephalus, congenital	Stenosis of aqueduct of Sylvius	Absent	Unknown	11
24	Hypoparathyroidism	Low serum Ca <sup>++</sup> , tetany	Absent	Unknown	27
25	Hypophosphatemia	Low serum inorganic phosphorus	Low serum inorganic phosphorus	Unknown	42
26	Ichthyosis simplex	Scaly skin	Absent	Scaly skin	9
27	Idiocy	Idiocy with microcephaly	Absent	Unknown	3
28	Keratosis follicularis (Laméris)	Multiple horny skin growths	Absent	Unknown	9
29	Lowe's syndrome	Glaucoma, mental retardation, renal tubule defect	Absent	Unknown	37
30	Megalocornea	Large cornea	Occasional manifestation	Unknown	13
31	Microphthalmia	Abnormally small eyes and blindness	Absent	Absent	28
32	Nystagmus	Severe involuntary movement of eyeball	Slight involuntary movement of eyeball	Unknown	29,34
33	Ophthalmoplegia	Paralysis of eye muscles, myopia; knee jerks absent	Knee jerks absent	Unknown	31

/1/ Gene symbol = *h*. /2/ Gene symbol = *h<sup>m</sup>*.

*continued*



3. SEX LINKAGE: MAN

Mutation	Phenotypic Expression			Reference
	Hemizygote $\bar{X}Y$	Heterozygote $\bar{X}X$	Homozygote $\bar{X}\bar{X}$	
(A)	(B)	(C)	(D)	(E)
34 Paraplegia, spastic	Spastic paralysis of legs	Absent	Unknown	19
35 Plasma thromboplastin component deficiency	Severe bleeder	Slight manifestation	Unknown	4,20
36 Retinal detachment, congenital	Retinal detachment and blindness	Absent	Blindness <sup>a</sup>	8,35,41
37 Retinitis pigmentosa	Choroidoretinal degeneration	Tapetal reflex	Unknown	12
38 Sclerosis, diffuse cerebral (Pelizaeus-Merzbacher)	Severe central nervous system involvement	Absent	Unknown	38

<sup>a</sup>/ Questionable.

Contributors: (a) Graham, John B., (b) McKusick, Victor A.

References: [1] Aldrich, R. A., A. G. Steinberg, and D. C. Campbell. 1954. *Pediatrics* 13:133. [2] Allan, W. 1939. *Arch. Internal Med.* 63:1123. [3] Allan, W., et al. 1944. *Am. J. Mental Deficiency* 48:325. [4] Barrow, E. M., et al. 1960. *J. Lab. Clin. Med.* 55:936. [5] Bell, J., and M. B. S. Haldane. 1937. *Proc. Roy. Soc. (London)*, B, 123:119. [6] Childs, B., et al. 1958. *Bull. Johns Hopkins Hosp.* 102:21. [7] Chung, C. S., et al. 1960. *Am. J. Human Genet.* 12:52. [8] Clark, E. 1898. *Trans. Ophthalmol. Soc. U. K.* 18:136. [9] Cockayne, E. A. 1933. *Inherited abnormalities of the skin and its appendages.* Oxford Univ. Press, London. [10] Crone, R. A. 1959. *Am. J. Ophthalmol.* 48:231. [11] Edwards, J. H. 1961. *Arch. Disease Childhood* 36:486. [12] Falls, H. F., and C. W. Cotterman. 1948. *Arch. Ophthalmol. (Chicago)* 40:685. [13] Gates, R. R. 1946. *Human genetics.* Macmillan, New York. [14] Graham, J. B., W. W. McLendon, and K. M. Brinkhous. 1953. *Am. J. Med. Sci.* 225:46. [15] Haldane, J. B. S. 1937. *J. Heredity* 28:58. [16] Israëls, M. C. G., et al. 1951. *Lancet* 1:1375. [17] Jacobsen, A. W. 1939. *J. Am. Med. Assoc.* 113:121. [18] Janeway, C. A., and D. Gitlin. 1957. *Advan. Pediat.* 9:65. [19] Johnston, A. W., and V. A. McKusick. 1962. *Am. J. Human Genet.* 14:83. [20] Lewis, J. H., and J. H. Ferguson. 1953. *Proc. Soc. Exptl. Biol. Med.* 82:445. [21] McCulloch, C., and R. J. P. McCulloch. 1948. *Trans. Am. Acad. Ophthalmol. Otolaryngol.* 52:160. [22] McKusick, V. A. 1960. *Heritable disorders of connective tissue.* C. V. Mosby, St. Louis. [23] McKusick, V. A. 1962. *Quart. Rev. Biol.* 37(2):69. [24] Mann, J. D., et al. 1962. *Lancet* 1:8. [25] Murakami, U., et al. 1951. *Nagoya J. Med. Sci.* 14:58. [26] Parker, N. 1958. *Am. J. Human Genet.* 10:196. [27] Peden, V. H. 1960. *Ibid.* 12:323. [28] Roberts, J. A. F. 1937. *Brit. Med. J.* 2:1213. [29] Rucker, C. W. 1949. *Am. J. Human Genet.* 1:52. [30] Rundles, R. W., and H. F. Falls. 1946. *Am. J. Med. Sci.* 211:641. [31] Salleras, A., and J. C. Ortiz de Zárate. 1950. *Brit. J. Ophthalmol.* 34:662. [32] Satalff, J., et al. 1955. *Am. J. Human Genet.* 7:201. [33] Snyder, L. H. 1946. *Principles of heredity.* D. C. Heath, Boston. [34] Sorsby, A. 1951. *Genetics in ophthalmology.* Butterworth, London. [35] Sorsby, A., et al. 1951. *Brit. J. Ophthalmol.* 35:1. [36] Stephens, F. E., and F. H. Tyler. 1951. *Am. J. Human Genet.* 3:111. [37] Streiff, E. B., W. Straub, and L. Tolay. 1958. *Ophthalmologica* 135:632. [38] Tyler, H. R. 1958. *Arch. Neurol. Psychiat.* 80:162. [39] Waardenburg, P. J., and J. van den Bosch. 1958. *Ann. Human Genet.* 21:101. [40] Williams, R. H., and C. Henry. 1947. *Ann. Internal Med.* 27:840. [41] Wilson, W. M. G. 1949. *Can. Med. Assoc. J.* 60:580. [42] Winters, R. W., et al. 1958. *Medicine* 37:97. [43] Wise, D. 1962. *Quart. J. Med., N.S.* 31:177. [44] Witkop, C. J. 1957. *Acta Genet. Statist. Med.* 7:236.

## 4. LINKAGE GROUPS: VERTEBRATES

The size or length of a linkage map reflects the extent of genetics investigation rather than the number of genes possessed by the animal. Capital letters (in columns giving Gene Symbol, Linkage, and Mutation) indicate dominant genes.

### Part I. GUINEA PIG

*Cavia porcellus* has 32 pair of chromosomes ( $\neq 1$  pair), including an XY pair in males. Linkage groups have been found for 2 pair.

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group I						
1	<i>R</i>	<i>R--Px</i>	43.8 $\pm$ 1.6	Rough fur	Rough fur, at least on hind toes	1-4
2	<i>Px</i>	<i>Px--R</i>	43.8 $\pm$ 1.6	Pollex	Tendency to atavistic return of thumb, little toe, and, on rare occasions, big toe	
Linkage Group II						
3	<i>si</i>	<i>si--m</i>	21.7 $\pm$ 5.2	silvered (stationary from birth)	Silver-coated fur; incomplete recessive	5
4	<i>m</i>	<i>m--si</i>	21.7 $\pm$ 5.2	modifier	Modifies rough fur effect; homozygote high-grade roughness	

Contributor: Wright, Sewall

References: [1] Castle, W. E., and A. Forbes. 1906. Carnegie Inst. Wash. Publ. 49:3. [2] Wright, S. 1928. Genetics 13:508. [3] Wright, S. 1941. Ibid. 26:650. [4] Wright, S. 1949. J. Exptl. Zool. 112:303. [5] Wright, S. 1959. Genetics 44:387.

### Part II. MOUSE

*Mus musculus* has 20 pair of chromosomes; linkage groups have been found for 19 pair.

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group I						
1	<i>fr</i>	<i>fr--sh-1</i>	16	frizzy	Fine thin hair, curled vibrissae	31
2	<i>ol</i> <sup>1</sup>	<i>ol--c</i>	17	oligodactyly	Reduced number of digits	48
3	<i>H-1</i>	<i>H-1--c</i>	7	Histocompatibility-1	Susceptibility to tissue transplants	91,93
4	<i>Hb</i> <sup>1</sup>	<i>Hb--c</i>	5 <sup>2</sup> , 2 <sup>3</sup>	Hemoglobin pattern	Electrophoretic pattern of hemoglobin	76
5	<i>sh-1</i>	<i>sh-1--c</i>	4 <sup>2</sup> , 3 <sup>3</sup>	shaker-1	Circling, head shaking, deafness	31,45,46
6	<i>c</i>	<i>c--p</i>	16 <sup>2</sup> , 12 <sup>3</sup>	albino	Absence of pigment in hair and eyes	31,45,46, 76
7	<i>hf</i>	<i>c--hf</i>	3	hepatic fusion	Fusion of left median and left lateral lobes of liver	4
8	<i>tp</i>	<i>tp--p</i>	5	taupe	Reduced pigment in coat	79
9	<i>H-4</i> <sup>1</sup>	<i>H-4--p</i>	0	Histocompatibility-4	Susceptibility to tissue transplants	93
10	<i>p</i>	<i>hf--p</i>	13	pink-eyed dilution	Pink eyes, reduced black or brown pigment	4
11	<i>qv</i>	<i>p--qv</i>	12	quivering	Locomotor instability, pronounced trembling in adults, priapism in old males	103
12	<i>da</i>	<i>p--da</i>	17	dark	Darkens back of agouti or yellow mice	28
13	<i>pu</i>	<i>p--pu</i>	22	pudgy	Tail short or absent, torso shortened	81
Linkage Group II						
14	<i>lu</i>	<i>lu--dse</i>	17	luxoid	Tibial hemimelia and preaxial polydactyly	44
15	<i>d</i>	<i>d--se</i>	0.1	dilute	Clumped pigment granules in hair	39
16	<i>se</i> <sup>1</sup>	<i>dse--du</i>	20	short ear	Reduced cartilaginous skeleton	44,90

/1/ Listed order not established. /2/ For heterozygous females. /3/ For heterozygous males.

continued

## 4. LINKAGE GROUPS: VERTEBRATES

### Part II. MOUSE

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group II						
17	<i>sv</i> <sup>1</sup>	<i>se--sv</i>	1	Snell's waltzer (recessive)	Circling, head shaking	43
18	<i>tk</i> <sup>1</sup>	<i>dse--tk</i>	11	tail kinks	Kinky tail, abnormal cervical and upper thoracic vertebrae	29
19	<i>du</i>	<i>du--dse</i>	20	ducky	Waddling gait	44,90
Linkage Group III						
20	<i>pn</i>	<i>pn--s</i>	30	pugnose	Frontal and nasal bones short and wide	53
21	<i>s</i> <sup>4</sup>	<i>s--hr</i>	8	piebald	Unpigmented areas of fur	87
22	<i>ag</i> <sup>1</sup>	<i>ag--hr</i>	0	agitans	Impaired locomotion, tremor, death at 3-4 wk	49
23	<i>hr</i>	<i>hr--W</i>	42	hairless	Hair shed beginning at 10-14 da	38
24	<i>wl</i> <sup>1,6</sup>	<i>hr--wl</i>	4	wabblers-lethal	Impaired locomotion, death at 3-4 wk	59
25	<i>pi</i>	<i>hr--pi</i>	36	pirouette	Circling, head shaking, deafness	18
26	<i>W</i>	<i>pi--W</i>	7	Dominant spotting	White spotting and dilution of coat color, macrocytic anemia, sterility	17
27	<i>Ph</i> <sup>1</sup>	<i>W--Ph</i>	0.1	Patch	White spotting	47
28	<i>le</i> <sup>1</sup>	<i>W--le</i>	12	light ears	Dilution of coat color	58
29	<i>lx</i>	<i>W--lx</i>	18	luxate	Tibial hemimelia, preaxial polydactyly	7
30	<i>rl</i>	<i>lx--rl</i>	16	reeler	Impaired locomotion, death at 3-4 wk	24
Linkage Group IV						
31	<i>r</i>	<i>r--si</i>	15	rodless retina	Absence of rods	52
32	<i>si</i>	<i>si--pg</i>	close	silvered	Absence or reduction of pigment in coat	30
33	<i>pg</i> <sup>1</sup>	<i>pg--si</i>	close	pigmy	Small size	30
34	<i>av</i> <sup>1</sup>	<i>si--av</i>	33	Ames' waltzer (recessive)	Circling, head shaking	83
Linkage Group V						
35	<i>Ra</i>	<i>Ra--a</i>	22	Ragged	Thin coat	11,60,72
36	<i>Op</i>	<i>Op--a</i>	27	Opossum	Very thin coat; probably an allele of <i>Ra</i>	40
37	<i>H-3</i> <sup>1</sup>	<i>H-3--a</i>	10	Histocompatibility-3	Susceptibility to tissue transplants	91
38	<i>kr</i>	<i>kr--a</i>	1	Kreisler (recessive)	Circling, head shaking, deafness	48,58
39	<i>bp</i>	<i>bp--a</i>	0.3	brachypodism	Short feet	77
40	<i>a</i>	<i>a--un</i>	5	non-agouti	Removes yellow band from hairs	5,35
41	<i>un</i>	<i>un--we</i>	7 <sup>a</sup> , 5 <sup>a</sup>	undulated	Wavy tail and abnormal vertebral column	26,35
42	<i>we</i>	<i>we--pa</i>	4 <sup>a</sup> , 2 <sup>a</sup>	wellhaartig	Wavy coat and vibrissae	26,35,58
43	<i>mg</i> <sup>1</sup>	<i>a--mg</i>	13 <sup>a</sup> , 10 <sup>a</sup>	mahogany	Dark coat, especially ears and tail	60
44	<i>pa</i>	<i>pa--ro</i>	1	pallid	Pink eyes, reduction of pigment in coat, frequent absence of otoliths	26
45	<i>ro</i>	<i>a--fi</i>	36 <sup>a</sup> , 27 <sup>a</sup>	rough	Air spaces in hair abnormal, waved vibrissae	9,97
46	<i>dm</i> <sup>1</sup>	<i>a--dm</i>	13	diminutive	Small size, malformed vertebrae and ribs	95
47	<i>fi</i>	<i>pa--fi</i>	19	fidget	Circling, head shaking, occasional polydactyly	6
48	<i>Sd</i>	<i>fi--Sd</i>	22	Danforth's short tail	Short tail, urogenital abnormalities	97
Linkage Group VI						
49	<i>N</i>	<i>N--Ca</i>	1 <sup>a</sup> , 3 <sup>a</sup>	Naked	Hair breaks off near skin level	12,66,69
50	<i>Ca</i>	<i>Ca--bt</i>	4 <sup>a</sup> , 11 <sup>a</sup>	Caracul	Wavy coat and vibrissae	66,69
51	<i>hl</i>	<i>Ca--hl</i>	2 <sup>a</sup> , 6 <sup>a</sup>	hair-loss	Loses hair, usually naked by 2-3 mo	50
52	<i>Ht</i> <sup>1</sup>	<i>Ca--Ht</i>	2 <sup>a</sup> , 3 <sup>a</sup>	High tail	Tail emerges high, short and thick at base, not kinked	80,81
53	<i>bt</i>	<i>hl--bt</i>	9	belted	White belt	50
Linkage Group VII						
54	<i>Re</i>	<i>Re--Al</i>	7	Rex	Wavy coat and vibrissae	14,58
55	<i>Al</i>	<i>Al--sh-2</i>	21	Alopecia	Hair thin and patchy beginning at 1 or 2 mo	58
56	<i>ti</i> <sup>1,6</sup>	<i>Re--ti</i>	20 <sup>a</sup> , 21 <sup>a</sup>	tipsy	Muscular incoordination, rabbit-like gait	84
57	<i>Tr</i> <sup>7</sup>	<i>Re--Tr</i>	23	Trembler	Convulsions in young, head trembling in adults	32

/1/ Listed order not established. /2/ For heterozygous females. /3/ For heterozygous males. /4/ *s--W* recombination, 47% [33]. /5/ *wl--W* recombination, 43% [59]. /6/ *ti--ut* recombination, 9% [84]. /7/ *Tr--sh-2* recombination, 3% [32].

continued

# 4. LINKAGE GROUPS: VERTEBRATES

## Part II. MOUSE

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group VII						
58	<i>sh-2<sup>a</sup></i>	<i>Re--sh-2<sup>1</sup></i>	28 <sup>a</sup> , 19 <sup>a</sup>	shaker-2	Circling, head shaking, deafness	10,23,70
59	<i>vt<sup>1,9</sup></i>	<i>Re--vt</i>	27 <sup>a</sup> , 18 <sup>a</sup>	vestigial	Tail short or absent	67
60	<i>wa-2</i>	<i>vi--wa-2</i>	23	waved-2	Wavy coat and vibrissae	68
61	<i>Tm<sup>1</sup></i>	.....	.....	Pulmonary tumors	Susceptibility to spontaneous and induced pulmonary tumors	96
Linkage Group VIII						
62	<i>m</i>	<i>m--Pt</i>	3	misty	Dilute coat color, tail and belly spots	55
63	<i>Pt</i>	<i>Pt--b</i>	5	Pintail	Short tail	51,55
64	<i>b</i>	<i>m--b</i>	5	brown	Brown instead of black pigment	86,100
65	<i>an<sup>1</sup></i>	<i>b--an</i>	5	anemia	Macrocytic anemia throughout life	48
66	<i>vc</i>	<i>b--vc</i>	7	vacillans	Muscular incoordination	85,86
67	<i>wi<sup>1</sup></i>	<i>b--wi</i>	6	whirler	Circling, head shaking	55
68	<i>wd</i>	<i>b--wd</i>	31	waddler	Swaying of hindquarters during locomotion	102
Linkage Group IX						
69	<i>T</i>	<i>T--Fu</i>	4	Brachyury	Short tail	2,20-22
70	<i>Fu</i>	<i>Fu--tf</i>	1	Fused	Tail and vertebral abnormalities	19
71	<i>tf<sup>1</sup></i>	<i>T--tf</i>	8	tufted	Successive waves of hair loss and regrowth from anterior to posterior	65
72	<i>H-2</i>	<i>Fu--H-2</i>	4	Histocompatibility-2	Susceptibility to tissue transplantation	1,2,89
Linkage Group X						
73	<i>v</i>	<i>v--ji</i>	18	waltzer	Circling, head shaking, deafness	88
74	<i>ji</i>	<i>ji--v</i>	18	jittery	Muscular incoordination, death at 3-4 wk	88
Linkage Group XI						
75	<i>tc<sup>1</sup></i>	<i>tc--mi</i>	8	truncate	Short tail, often with intermediate vertebrae of tail or sacrum missing	57
76	<i>mi</i>	<i>mi--px</i>	3	microphthalmia	Reduced pigment, failure of bone resorption	8
77	<i>px</i>	<i>px--wa-1</i>	1	postaxial hemimelia	Postaxial side of limbs defective	8
78	<i>wa-1</i>	<i>wa-1--Lc</i>	8	waved-1	Waved hair and vibrissae	75
79	<i>Lc</i>	<i>mi--wa-1</i>	11	Lurcher	Swaying of hindquarters and falling to one side	3
80	<i>ob<sup>1</sup></i>	<i>mi--ob</i>	29	obese	Obesity with hyperglycemia	16
Linkage Group XII						
81	<i>ru</i>	<i>ru--je</i>	49	ruby eye	Reduced pigmentation of eyes and hair	15,27,37,98
82	<i>je</i>	<i>je--ru</i>	49	jerker	Circling, head shaking, deafness	15,27,39,98
Linkage Group XIII						
83	<i>Lp</i>	<i>Lp--ln</i>	38 <sup>a</sup> , 35 <sup>a</sup>	Loop tail	Looped tail, abnormal behavior	94
84	<i>py</i>	<i>py--ln</i>	38 <sup>a</sup> , 24 <sup>a</sup>	polydactyly	Preaxial polydactyly	34,71
85	<i>dr<sup>1</sup></i>	<i>dr--Dh</i>	22	dreher	Circling, head shaking	63
86	<i>Dh</i>	<i>Dh--ln</i>	2	Dominant hemimelia	Preaxial hemimelia, absence of spleen	63
87	<i>ln</i>	<i>ln--Sp</i>	5	leaden	Clumped pigment granules in hair	71,94
88	<i>th<sup>1</sup></i>	<i>th--ln</i>	5	tilted head	Head tilted to right or left side	61
89	<i>Sp</i>	<i>Sp--fz</i>	40 <sup>a</sup> , 33 <sup>a</sup>	Spotch	White spotting on belly, feet, and tail	71,94
90	<i>fz</i>	<i>ln--fz</i>	43 <sup>a</sup> , 36 <sup>a</sup>	fuzzy	Thin wavy hair and vibrissae	71,94
Linkage Group XIV						
91	<i>cr</i>	<i>cr--ch</i>	15	crinkled	Absence of guard hairs and zigzags	54,74
92	<i>ch</i>	<i>ch--f</i>	18	congenital hydrocephalus	Severe reduction in cartilaginous skeleton	54,74
93	<i>f</i>	<i>f--ch</i>	18	flexed tail	Anemia at birth, flexed tail, belly spot	54,74

/1/ Listed order not established. /a/ For heterozygous females. /a/ For heterozygous males. /a/ *sh-2--wa-2* recombination, 24% for heterozygous females and 30% for heterozygous males [10,14,36,92,101]. /a/ *sh-2--vt* recombination, 2% [68].

continued



## 4. LINKAGE GROUPS: VERTEBRATES

### Part II. MOUSE

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group XV						
94	<i>Tw</i>	<i>Tw--ax</i>	0	Twirler	Circling, head shaking	62
95	<i>ax</i>	<i>ax--Tw</i>	0	ataxia	Muscular incoordination, death at 3-4 wk	62
Linkage Group XVI						
96	<i>Va</i>	<i>Va--de</i>	28	Varitint-waddler	Dilute and spotted coat, circling, head shaking, deafness	13
97	<i>de</i>	<i>de--Va</i>	28	droopy ear	Ears set low on head, pinnae project laterally	13
Linkage Group XVII						
98	<i>sa</i>	<i>sa--bg</i>	9	satin	Silky hair texture with high sheen	82
99	<i>bg</i>	<i>bg--sa</i>	9	beige	Diluted coat color	82
Linkage Group XVIII						
100	<i>Hk</i>	<i>Hk--Os</i>	17	Hook	Short tail, anus displaced toward tail	42
101	<i>Os</i>	<i>Os--tg</i>	0	Oligosyndactylism	Digits reduced in number and fused	41
102	<i>tg</i> <sup>1</sup>	<i>tg--Os</i>	0	tottering	Wobbly gait, occasional convulsions	41
Linkage Group XX (Sex Chromosome)						
103	<i>Bn</i>	<i>Bn--Ta</i>	12	Bent	Short crooked tail	73
104	<i>Gy</i> <sup>1</sup>	<i>Gy--Ta</i>	close	Gyro	Circling; abnormal development of long bones and ribs in males	64
105	<i>Ta</i>	<i>Ta--Mo</i>	4	Tabby	Dark transverse stripes	25,26
106	<i>Blo</i>	<i>Ta--Blo</i>	3	Blotchy	Irregular patches of light fur, males viable	78
107	<i>Mo</i>	<i>Mo--Ta</i>	4	Mottled	Patches of light hair, males die in utero	25,26
108	<i>To</i> <sup>1</sup>	<i>Bn--To</i>	22	Tortoise	Like <i>Mo</i> ; possibly an allele	56
109	<i>jp</i>	<i>Ta--jp</i>	20	jimpy	Muscular incoordination, death at 3-4 wk	73
110	<i>sf</i> <sup>1</sup>	<i>Ta--sf</i>	44	scurfy	Scaliness, tight skin, death at 3-4 wk	99

<sup>1</sup>/ Listed order not established.

*Contributors:* (a) Green, Margaret C., (b) Snell, George D.

*References:* [1] Allen, S. L. 1955. *Cancer Res.* 15:315. [2] Allen, S. L. 1955. *Genetics* 40:627. [3] Bunker, H., and G. D. Snell. 1948. *J. Heredity* 39:28. [4] Bunker, L. E., Jr. 1959. *Ibid.* 50:40. [5] Carter, T. C. 1947. *Heredity* 1:367. [6] Carter, T. C. 1951. *J. Genet.* 50:264. [7] Carter, T. C. 1951. *Ibid.* 50:300. [8] Carter, T. C. Unpublished. Univ. Genetics Dept., Medical Research Council, Edinburgh, 1958. [9] Carter, T. C., and H. Grüneberg. 1950. *Heredity* 4:373. [10] Carter, T. C., and R. J. S. Phillips. 1953. *Z. Induktive Abstammungs-Vererbungslehre* 85:564. [11] Carter, T. C., and R. J. S. Phillips. 1954. *J. Heredity* 45:151. [12] Cooper, C. B. 1939. *Ibid.* 30:212. [13] Curry, G. A. 1959. *J. Embryol. Exptl. Morphol.* 7:39. [14] Dickie, M. M. 1955. *J. Heredity* 46:31. [15] Dickie, M. M. Unpublished. Roscoe B. Jackson Memorial Laboratory, Bar Harbor, Maine, 1958. [16] Dickie, M. M., and P. W. Lane. 1957. *Mouse News Letter* 17:52. [17] Dickie, M. M., and G. W. Woolley. 1946. *J. Heredity* 37:335. [18] Dickie, M. M., and G. W. Woolley. 1948. *Ibid.* 39:288. [19] Dunn, L. C. 1958. *Mouse News Letter* 18:24. [20] Dunn, L. C., and E. Caspari. 1945. *Genetics* 30:543. [21] Dunn, L. C., and S. Gluecksohn-Waelsch. 1953. *Ibid.* 38:512. [22] Dunn, L. C., and S. Gluecksohn-Waelsch. 1954. *J. Genet.* 52:383. [23] Falconer, D. S. 1947. *Heredity* 1:133. [24] Falconer, D. S. 1952. *Ibid.* 6:255. [25] Falconer, D. S. 1953. *Z. Induktive Abstammungs-Vererbungslehre* 85:210. [26] Falconer, D. S. 1954. *Ibid.* 86:263. [27] Falconer, D. S. 1956. *Mouse News Letter* 15:24. [28] Falconer, D. S. 1957. *Ibid.* 17:40. [29] Falconer, D. S. 1961. *Ibid.* 25:30. [30] Falconer, D. S., and J. W. B. King. 1953. *Ibid.* 9(Suppl.):7. [31] Falconer, D. S., and G. D. Snell. 1952. *J. Heredity* 43:53. [32] Falconer, D. S., and W. R. Sobey. 1953. *Ibid.* 49:159. [33] Fisher, R. A. 1946. *Am. Naturalist* 80:568. [34] Fisher, R. A. 1953. *Heredity* 7:91. [35] Fisher, R. A., and

*continued*



## 4. LINKAGE GROUPS: VERTEBRATES

### Part II. MOUSE

W. Landauer. 1953. *Am. Naturalist* 87:116. [36] Fisher, R. A., M. F. Lyon, and A. R. G. Owen. 1947. *Heredity* 1:355. [37] Fisher, R. A., and G. D. Snell. 1948. *Ibid.* 2:271. [38] Gates, W. H., and T. Pullig. 1945. *Genetics* 30:4. [39] Goodwins, I. R., and M. A. C. Vincent. 1955. *Heredity* 9:413. [40] Green, E. L., and S. J. Mann. 1961. *J. Heredity* 52:223. [41] Green, M. C. 1960. *Mouse News Letter* 22:34. [42] Green, M. C. 1960. *Ibid.* 23:34. [43] Green, M. C. 1961. *Ibid.* 25:38. [44] Green, M. C. 1961. *J. Heredity* 52:73. [45] Grüneberg, H. 1935. *J. Genet.* 31:157. [46] Grüneberg, H. 1936. *Ibid.* 33:255. [47] Grüneberg, H., and G. M. Truslove. 1960. *Genet. Res.* 1:69. [48] Hertwig, P. 1942. *Z. Induktive Abstammungs- Vererbungslehre* 80:220. [49] Hoecker, G., et al. 1954. *J. Heredity* 45:10. [50] Hollander, W. F. 1959. *Mouse News Letter* 20:34. [51] Hollander, W. F., and L. C. Strong. 1951. *J. Heredity* 42:179. [52] Keeler, C. E. 1930. *Howe Lab. Ophthalmol. Bull.* 3. [53] Kidwell, J. F. 1961. *Mouse News Letter* 24:39. [54] King, J. W. B. 1956. *Nature* 178:1126. [55] Lane, P. W. 1960. *Mouse News Letter* 23:35. [56] Lane, P. W. 1960. *Ibid.* 23:36. [57] Lane, P. W. 1961. *Ibid.* 25:38. [58] Lane, P. W. Unpublished. Roscoe B. Jackson Memorial Laboratory, Bar Harbor, Maine, 1960. [59] Lane, P. W., and M. M. Dickie. 1961. *J. Heredity* 52:159. [60] Lane, P. W., and M. C. Green. 1960. *Ibid.* 51:228. [61] Larsen, M. M. 1961. *Mouse News Letter* 24:60. [62] Lyon, M. F. 1958. *J. Embryol. Exptl. Morphol.* 6:105. [63] Lyon, M. F. 1961. *Genet. Res.* 2:92. [64] Lyon, M. F. 1961. *Mouse News Letter* 24:34. [65] Lyon, M. F., and R. J. S. Phillips. 1959. *Heredity* 13:23. [66] Mallyon, S. A. 1951. *Nature* 168:118. [67] Michie, D. 1955. *J. Genet.* 53:270. [68] Michie, D. 1955. *Ibid.* 53:280. [69] Murray, J. M., and G. D. Snell. 1945. *J. Heredity* 36:266. [70] Nasrat, G. E. 1956. *Proc. Zool. Soc. (Bengal)* 9:85. [71] Parsons, P. A. 1958. *Heredity* 12:77. [72] Parsons, P. A. 1958. *Ibid.* 12:357. [73] Phillips, R. J. S. 1954. *Z. Induktive Abstammungs- Vererbungslehre* 86:322. [74] Phillips, R. J. S. 1956. *J. Heredity* 47:302. [75] Phillips, R. J. S. 1960. *J. Genet.* 57:35. [76] Popp, R. A., and W. St. Amand. 1960. *J. Heredity* 51:141. [77] Runner, M. N. 1959. *Ibid.* 50:81. [78] Russell, L. B. 1960. *Mouse News Letter* 23:58. [79] Russell, L. B. 1961. *Ibid.* 25:64. [80] St. Amand, W., and M. B. Cupp. 1957. *Ibid.* 16:37. [81] St. Amand, W., and M. B. Cupp. 1957. *Ibid.* 17:88. [82] St. Amand, W., and M. B. Cupp. 1958. *Ibid.* 19:38. [83] Schaible, R. H. 1961. *Ibid.* 24:38. [84] Searle, A. G. 1961. *Genet. Res.* 2:122. [85] Sirlin, J. L. 1956. *J. Genet.* 54:42. [86] Sirlin, J. L. 1957. *Heredity* 11:259. [87] Snell, G. D. 1931. *Genetics* 16:42. [88] Snell, G. D. 1945. *J. Heredity* 36:279. [89] Snell, G. D. 1952. *Heredity* 6:247. [90] Snell, G. D. 1955. *J. Heredity* 46:27. [91] Snell, G. D. 1958. *J. Natl. Cancer Inst.* 21:843. [92] Snell, G. D., and L. W. Law. 1939. *J. Heredity* 30:447. [93] Snell, G. D., and L. C. Stevens. 1961. *Immunology* 4:366. [94] Snell, G. D., et al. 1954. *Heredity* 8:271. [95] Stevens, L. C., and J. A. Mackensen. 1961. *Mouse News Letter* 24:41. [96] Tatchell, J. A. H. 1961. *Nature* 190:837. [97] Wallace, M. E. 1957. *Heredity* 11:223. [98] Wallace, M. E. 1958. *Ibid.* 12:453. [99] Welshons, W. J., and L. B. Russell. 1959. *Proc. Natl. Acad. Sci. U. S.* 45:560. [100] Woolley, G. W. 1945. *J. Heredity* 36:269. [101] Wright, M. E. 1947. *Heredity* 1:349. [102] Yoon, C. H. 1961. *J. Heredity* 52:279. [103] Yoon, C. H., and E. P. Les. 1957. *Ibid.* 48:176.

### Part III. RABBIT

*Oryctolagus cuniculus* has 22 pair of chromosomes; linkage groups have been found for 6 pair. For additional information, consult references 7-9, 12.

Gene Symbol	Locus	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	
Linkage Group I					
1	<i>c</i>	0	albinism	Coat color alleles vary from chinchilla to complete albinism	1,2,5
2	<i>y</i>	14,4	yellow fat	Yellow fat	3,11
3	<i>b</i>	42,8	brown	Brown coat	3,11

*continued*

# Contrails

## 4. LINKAGE GROUPS: VERTEBRATES

### Part III. RABBIT

Gene Symbol	Locus	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)	(E)
Linkage Group II				
4 <i>du</i>	0	dutch pattern	White belt on colored background	3,11
5 <i>En</i>	1.2	English	Colored spots on white background	1,4,6
6 <i>l</i>	14.3	angora hair	Increase in hair fiber length	1,4
Linkage Group III				
7 <i>r<sub>1</sub></i>	0	<i>rex<sub>1</sub></i>	Short, plushlike coat	3,11
8 <i>r<sub>2</sub></i>	17.2	<i>rex<sub>2</sub></i>	Short, plushlike coat	2,4
Linkage Group IV				
9 <i>a</i>	0	non-agouti	Black coat	3,11
10 <i>dw</i>	14.7	dwarf	Small size, lethal shortly after birth	3,11
11 <i>w</i>	29.9	wide-banded agouti	Wide banding of agouti hairs	5,10
Linkage Group V				
12 <i>br</i>	0	brachydactyly	Abnormality of toes	11
13 <i>f</i>	28.3	furless	Fur restricted to extremities	5
14 <i>an</i>	36.8	erythrocyte agglutination	Erythrocytes agglutinate	14
Linkage Group VI				
15 <i>E</i>	0	Extension	Extension of dark pigment	11
16 <i>At</i>	26.2	Production of atropinesterase	Production of atropinesterase	13

Contributors: (a) Sawin, Paul B., (b) Novitski, E.

References: [1] Castle, W. E. 1926. Carnegie Inst. Wash. Publ. 337:3. [2] Castle, W. E. 1936. Proc. Natl. Acad. Sci. U. S. 22:222. [3] Castle, W. E. 1940. Mammalian genetics. Harvard Univ. Press, Cambridge. [4] Castle, W. E., and N. Nachtsheim. 1933. Proc. Natl. Acad. Sci. U. S. 19:1006. [5] Castle, W. E., and P. B. Sawin. 1941. Ibid. 27:519. [6] Pease, M. S. 1928. Verhandl. Ver. Intern. Kongr. Vererbungswiss. 2:1153. [7] Rifaat, O. M. 1954. Heredity 8:107. [8] Robinson, R. 1956. J. Genet. 54:358. [9] Robinson, R. 1958. Bibliog. Genet. (Haag) 17:229. [10] Sawin, P. B. 1934. J. Heredity 25:477. [11] Sawin, P. B. 1944. Proc. Natl. Acad. Sci. U. S. 30:220. [12] Sawin, P. B. 1955. Advan. Genet. 7:183. [13] Sawin, P. B., and D. Glick. 1943. Proc. Natl. Acad. Sci. U. S. 29:55. [14] Sawin, P. B., M. A. Griffin, and C. A. Stuart. 1944. Ibid. 30:217.

### Part IV. RAT

*Rattus norvegicus* has 21 pair of chromosomes; linkage groups have been found for 5 pair. Seven genes have been found to be independent of linkage groups I-V, and of each other, and are provisionally regarded as markers of 7 additional chromosome pair: jaundice (*j*), curly coat (*Cu<sub>2</sub>*), cataract (*Ca*), blue dilution of coat (*d*), hooded coat pattern (*h*), cowlick (*cu*), and shaker (*sr*). [3, 5, 6, 9, 10]

Gene Symbol	Locus	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)	(E)
Linkage Group I				
1 <i>p</i>	0	pink <i>e; e</i>	Coat yellow, eyes pink	1, 2, 7, 14
2 <i>r</i>	20.5	red-eyed yellow	Coat yellow, eyes red	
3 <i>c</i>	21	albinism	Absence of pigment from coat and eyes	
4 <i>l</i>	24.3	lethal	Skeletal abnormalities	
5 <i>w</i>	66.3	waltzing	Runs in circles	
Linkage Group II				
6 <i>Sh</i>	0	Shaggy	Hair and vibrissae curved	2, 7-9, 13
7 <i>Cu</i>	4	Curly	Hairs of coat and vibrissae curved	

continued

## 4. LINKAGE GROUPS: VERTEBRATES

### Part IV. RAT

Gene Symbol	Locus	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	
Linkage Group II					
8	<i>an</i>	14.3	anemia	Lack of erythrocytes; young anemic	2, 7-9, 13
9	<i>in</i>	28	incisorless	Incisors lacking	
10	<i>s</i>	47	silvered	Coat silvered	
11	<i>b</i>	52	brown	Black pigment of coat and eyes replaced by brown	
Linkage Group III					
12	<i>n</i>	0	naked	Naked except for short fuzzy coat	2, 4, 12
13	<i>hr</i>	34.7	hairless	Hair lost at approximately 4 wk	
14	<i>wo</i>	75	wobbly	Ataxic locomotion	
Linkage Group IV					
15	<i>k</i>	0	kinky	Hairs of coat and vibrissae kinky	2, 8
16	<i>st</i>	34.1	stub	Short stubby tail	
Linkage Group V					
17	<i>A</i>	0	Agouti	Fur color agouti, wild type	2, 11
18	<i>f</i>	44.6	fawn	Coat tawny blue to fawn	

Contributors: (a) Castle, W. E., (b) Novitski, E.

References: [1] Castle, W. E. 1916. Carnegie Inst. Wash. Publ. 241:175. [2] Castle, W. E. 1947. Proc. Natl. Acad. Sci. U. S. 33:109. [3] Castle, W. E. 1951. Genetics 36:254. [4] Castle, W. E. 1955. J. Heredity 46:84. [5] Castle, W. E., E. R. Dempster, and H. C. Shurrager. 1955. Ibid. 46:9. [6] Castle, W. E., and H. D. King. 1940. Proc. Natl. Acad. Sci. U. S. 26:578. [7] Castle, W. E., and H. D. King. 1941. Ibid. 27:394. [8] Castle, W. E., and H. D. King. 1944. Ibid. 30:79. [9] Castle, W. E., and H. D. King. 1947. J. Heredity 38:341. [10] Castle, W. E., and H. D. King. 1948. Proc. Natl. Acad. Sci. U. S. 34:135. [11] Castle, W. E., and H. D. King. 1949. Ibid. 35:545. [12] Castle, W. E., H. D. King, and A. L. Daniels. 1941. Ibid. 27:250. [13] King, H. D., and W. E. Castle. 1935. Ibid. 21:390. [14] King, H. D., and W. E. Castle. 1937. Ibid. 23:56.

### Part V. CHICKEN

*Gallus domesticus* has 39 pair of chromosomes; linkage groups have been found for 6 pair (groups IV and V may eventually be joined).

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)	(E)	(F)
Linkage Group I (Sex Chromosome)					
1	<i>ko</i>	<i>ko--B</i>	13	head streak in down	1, 2
2	<i>B</i>	<i>B--ld</i>	10	Barring	
3	<i>Sd</i>	<i>B--Sd</i>	<1	Dilution	
4	<i>ld</i>	<i>ld--br</i>	27	Inhibitor	
5	<i>br</i>	<i>br--Li</i>	10	brown eyes	
6	<i>Li</i>	<i>Li--S</i>	16	Light down	
7	<i>S</i>	<i>S--al</i>	1.2	Silver	
8	<i>al</i>	<i>al--K</i>	1.6	albinism	
9	<i>K</i>	<i>K--dw</i>	6.6	Slow feathering	
10	<i>dw</i>	<i>dw--S</i>	7	dwarf	
11	<i>px</i>	<i>al--px</i>	11	paroxysm	
12	<i>n</i>	<i>px--n</i>	6	naked	
13	<i>sh</i>	<i>n--sh</i>	14	shaker	
14	<i>xl</i> <sup>1</sup>			lethal	
15	<i>j</i> <sup>1</sup>			jittery	

/1/ Listed order not established.

continued

## 4. LINKAGE GROUPS: VERTEBRATES

### Part V. CHICKEN

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Linkage Group II						
16	<i>Cp</i>	<i>Cp--R</i>	0.4	Creoper	Achondroplasia	1
17	<i>R</i>	<i>R--U</i>	30	Rose comb	Rose comb	
18	<i>U</i>	<i>U--R</i>	30	Uropygial	Bifurcation of uropygial papilla	
Linkage Group III						
19	<i>fr</i>	<i>fr--Cr</i>	46	fray	Defective wing and tail feathers	1,3
20	<i>Cr</i>	<i>Cr--I</i>	12.5	Crest	Topknot and cerebral hernia	
21	<i>I</i>	<i>I--F</i>	17	Dominant white	White plumage	
22	<i>F</i>	<i>F--I</i>	17	Frizzling	Recurved feathers	
Linkage Group IV						
23	<i>O</i>	<i>O--P</i>	5	Blue egg	Eggshell blue	1
24	<i>P</i>	<i>P--ma</i>	33	Pea comb	Pea comb	
25	<i>ma</i>	<i>ma--Na</i>	46	marbling	Pattern in down of chick	
26	<i>Na</i>	<i>Na--ma</i>	46	Naked neck	Pterylae reduced	
Linkage Group V						
27	<i>Na</i>	<i>Na--h</i>	43	Naked neck	Pterylae reduced	3
28	<i>h</i>	<i>h--Fl</i>	11	silkie	Barbules lack hooklets	
29	<i>Fl</i>	<i>Fl--h</i>	11	Flightless	Remiges break off	
Linkage Group VI						
30	<i>D</i>	<i>D--M</i>	26	Duplex comb	Bifurcation of comb	1,3
31	<i>M</i>	<i>M--Po</i>	33	Multiple spurs	Multiple spurs	
32	<i>Po</i>	<i>Po--M</i>	33	Polydactyly	Supernumerary digits	

Contributor: Hutt, F. B.

References: [1] Hutt, F. B. 1949. Genetics of the fowl. McGraw-Hill, New York. [2] Hutt, F. B. 1960. Heredity 15:97. [3] Warren, D. C. 1949. Genetics 34:333.

## 5. LINKAGE GROUPS: INVERTEBRATES

The size or length of a linkage map reflects the extent of genetics investigation rather than the number of genes possessed by the insect. Capital letters (in columns giving Gene Symbol, Linkage, and Mutation) indicate dominant genes.

### Part I. FRUIT FLY

*Drosophila melanogaster* has 4 pair of chromosomes; linkage groups have been found for all 4 pair. For information on other species of *Drosophila*, consult the following references: *D. affinis* [30, 39], *D. ananassae* [17, 25-27, 30], *D. hydei* [32-34], *D. montium* [28, 29], *D. persimilis* [10, 18, 35], *D. prosaltans* [31], *D. pseudoobscura* [24, 30, 39, 42], *D. similans* [30, 38, 41], *D. subobscura* [2, 6-9, 13-16, 21-23, 36, 37], *D. virilis* [3-5, 19, 30], *D. willistoni* [12, 20]. **Gene Symbol** (column A): *l* = lethal; *l* = the number one; *l* = the letter 1. **Locus** (column B): (Dp) = duplication; (Df) = deficiency. A number of loci have recently proved to be pseudoallelic (show crossing over with low frequency within subdivisions of an individual locus); such loci are indicated in column C as (pseudo).

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
X Chromosome [11, 43]				
1	<i>l(1)Jl</i>	0	lethal (1) Jacobs-Muller	Almost completely lethal; survivors scute, sterile
2	<i>l(1)55a</i>	0-	lethal (1) 55a	Lethal, heterozygote hyperviable
3	<i>su-w<sup>a</sup></i>	0-	suppressor of apricot	<i>w<sup>a</sup></i> eye color made to resemble <i>w<sup>co</sup></i>
4	<i>y</i>	0	yellow	Body yellow; bristles and hairs yellow or brown in different alleles
5	<i>brc</i>	0	brachymacrochaete	Macrochaetes reduced

continued



# Contrails

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
X Chromosome [11,43]				
6	<i>ac</i>	0+	achaete	Postdorsocentrals missing; intraocular and eye hairs fewer
7	<i>Hw</i>	0+ (Dp)	Hairy-wing	Extra bristles along wing veins, on head and thorax
8	<i>sc</i>	0+	scute	Scutellar bristles missing, others missing or reduced
9	<i>svr</i>	0+	silver	Body silvery, bristles dark
10	<i>su-s</i>	0+	suppressor of sable	Suppresses <i>s</i> and <i>v</i>
11	<i>dor</i>	0+	deep orange	Eyes orange; female sterile
12	<i>l(1)7e</i>	0+	lethal (1) 7e	Dies in larval stage
13	<i>saw</i>	0+	sawtooth	Wing hairs serrated
14	<i>su-b</i>	0.1	suppressor of black	Suppresses <i>b</i>
15	<i>om</i>	0.1±	ommatidia	Eyes slightly rough
16	<i>M(1)Bld</i>	0.1+ (Df)	Minute (1) Blond	Extreme minute (small bristles, low viability, homozygous lethal)
17	<i>l(1)7</i>	0.3	lethal (1) 7	Dies as late larva; tumors present
18	<i>fla</i>	0.3±	flat eye	Eyes small, flat
19	<i>sta</i>	0.3±	stubarista	Antennae and aristae short, bristles reduced, eye rotated
20	<i>tw</i>	0.4±	twisted	Abdomen twisted counterclockwise
21	<i>mwi</i>	0.4±	misheld wings	Wings divergent, upheld; eyes oval
22	<i>uq</i>	0.5±	unequal wings	Wings short, often unequal
23	<i>kz</i>	0.7	kurz	Bristles short, fine; postscutellars often absent
24	<i>rey</i>	0.7±	rough eye	Eyes small, rough
25	<i>pn</i>	0.8	prune	Eyes brownish, darkening with age, often mottled
26	<i>mk</i>	0.8±	murky	Body and eyes dark; female sterile
27	<i>gt</i>	0.9	giant	Giant larva, pupa, adult; variable
28	<i>rsc</i>	0.9±	reduplicated sex combs	Sex combs on all six legs of male
29	<i>fc</i>	0.9±	faulty chaete	Bristles short, thin; some absent
30	<i>ovi</i>	0.9±	ovioculus	Eyes small, egg-shaped; male sterile
31	<i>z</i>	1.0	zeste	Eyes yellow in female at 25°C; temperature-sensitive; interacts with <i>w</i> alleles
32	<i>fb</i>	1.0±	fine bristle	Bristles short, fine
33	<i>l(1)ml</i>	1.0	lethal (1) melanomalike	Dies as late larva, melanotic inclusions
34	<i>bsc</i>	1.1	bent scutellars	Scutellars and other bristles often bent
35	<i>mis</i>	1.3±	misproportioned	Abdomen abnormal in shape and size
36	<i>w</i>	1.5	white (pseudo)	White eyes, ocelli, testes, malpighian tubes
37	<i>rst<sup>2</sup></i>	1.7	roughest <sup>2</sup>	Eyes rough, body dwarfed, some bristles reduced
38	<i>To</i>	2.3-	Tousled	Thoracic bristles disarranged, duplicated
39	<i>Co</i>	3.0± (Dp)	Confluens	Wing veins thick, with deltas
40	<i>nd</i>	3.0±	notchoid (pseudo with <i>spl-fa-N</i> )	Eyes small, wings notched
41	<i>spl</i>	3.0±	split (pseudo with <i>spl-fa-N</i> )	Eyes rough, small; bristles often split or missing
42	<i>fa</i>	3.0±	facet (pseudo with <i>spl-fa-N</i> )	Eyes rough, wings nicked
43	<i>N</i>	3.0± (often Df)	Notch (pseudo with <i>spl-fa-N</i> )	Wings notched; male lethal
44	<i>Ax</i>	3.0±	Abruptex	Wings short, veins incomplete; thorax with mid-furrow
45	<i>rud</i>	3.3±	ruddle	Eyes reddish brown
46	<i>slc</i>	3.6±	slim chaete	Bristles fine, short
47	<i>Sc</i>	4.0±	Scotched eye	Ommatidia disarranged; male lethal
48	<i>dm</i>	4.6	diminutive	Body and bristles small; female sterile
49	<i>M(1)3E</i>	5.0±	Minute (1) 3E	Slight minute (body small, bristles fine, homozygous lethal)
50	<i>su<sup>X</sup>-dx</i>	5.0±	suppressor of deltex	<i>dx</i> made nearly +; male fertile
51	<i>ec</i>	5.5	echinus	Eyes rough, large; facets large
52	<i>mj</i>	5.5±	macrofine	Body small, macrochaetes fine
53	<i>te</i>	5.6±	tenuchaete	Bristles fine, short; eyes dark
54	<i>Oc</i>	5.7±	Ocellarless	One or both ocellar bristles missing
55	<i>mo</i>	6.7±	microoculus	Eyes small, wings narrow
56	<i>amb</i>	6.8±	amber	Body pale yellow, bristles reduced; male sterile
57	<i>bi</i>	6.9	bifid	Wing veins fused into bifid stalk
58	<i>M(1)4BC</i>	7.0±	Minute (1) at 4BC	Strong minute (body small, bristles fine, homozygous lethal)
59	<i>peb</i>	7.3±	pebbled	Eyes slightly roughened
60	<i>lac</i>	7.3±	lacquered	Body color light, glistening; eyes small
61	<i>rb</i>	7.5	ruby	Eyes clear ruby, darkening to garnet

continued

# Contrails

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
X Chromosome [11,43]				
62	<i>dow</i>	8.0	downy	Bristles fuzzy; male sterile
63	<i>rg</i>	11.0	rugose	Eyes rough; wings thin, frayed
64	<i>bo</i>	12.5	bordeaux	Eyes dark wine
65	<i>omm</i>	12.8	ommatoreductum	Peripheral ommatidia absent, giving rough eye; head, thorax abnormal
66	<i>cx</i>	13.6	curlex	Wings bent upward
67	<i>cv</i>	13.7	crossveinless	Crossveins absent or nearly so
68	<i>mur</i>	14.3	murrey	Eyes reddish purple; body size, bristles reduced
69	<i>rmp</i>	14.4±	rumpled	Wings unexpanded, bristles disarranged
70	<i>rux</i>	15.0	roughex	Eyes small, rough
71	<i>Ext</i>	15.2±	Extras	Wing veins thickened, extra veins present
72	<i>vs</i>	16.3	vesiculated	Wings warped, divergent, blistered
73	<i>dx</i>	17.0	deltex	Wings thickened and with deltas
74	<i>ov</i>	17.5	oval	Eyes oval and rough
75	<i>tmc</i>	17.5±	tonomacrochaetes	Macrochaetes thin; abdomen pale
76	<i>shf</i> <sup>2</sup>	17.9	shifted <sup>2</sup>	Wing veins shifted closer together
77	<i>cm</i>	18.9	carmine	Eyes dark ruby
78	<i>scp</i>	19.3	scooped	Wings upturned, warped
79	<i>bis</i>	19.8± (Df)	bistre	Eyes and ocelli dark brown; male sterile
80	<i>ct</i>	20.0	cut	Wings cut to points, scalloped
81	<i>sn</i>	21.0	singed (pseudo)	Bristles and hairs curled; female sterile
82	<i>l(1)mys</i>	21.7	lethal(1)myospheroid	Dies as embryo with spheroid muscles
83	<i>ha</i>	22.7±	hair bristles	Bristles fine, short; fly small
84	<i>oc</i>	23.1	ocelliless	Ocelli absent; female sterile
85	<i>pam</i>	23.1	platinum	Male body and bristles almost colorless, bristle bases dark; sterile
86	<i>gg</i> <sup>2</sup>	23.1±	goggle <sup>2</sup>	Eyes bulging, head bristles fewer
87	<i>ptg</i>	23.2	pentagon	Thoracic trident and scutellar spot dark
88	<i>ccw</i>	23.4±	concave wing	Wings reduced, concave
89	<i>ch-b</i>	23.8	chilblained-b	Tarsi conglutinated
90	<i>thd</i>	25.0±	tiny-bristloid	Bristles medium-fine; fly small; viability good
91	<i>Lg</i>	27.0±	Large	Body large; late-hatching
92	<i>dd</i> <sup>2</sup>	27.2	displaced <sup>2</sup>	Antennae sunken; eyes and head deformed
93	<i>t</i>	27.5	tan	Body yellowish, antennae light yellow
94	<i>amx</i>	27.7-	almondex	Eyes narrow, rough; female sterile
95	<i>lz</i>	27.7	lozenge (pseudo)	Eyes narrow, facets abnormal; female usually sterile
96	<i>tar</i>	27.7±	tarry	Femur and tibia blackened
97	<i>dvr</i>	28.1	divers	Wings short, dark; with <i>y</i> , wings curled
98	<i>sma</i>	29.9±	smaller	Body size reduced
99	<i>su-Cbx</i>	30.0±	suppressor of Contra-bithorax	Almost completely suppresses <i>Cbx</i> effect in male
100	<i>tpw</i>	30.8±	tapered wing	Wings short, pointed at L3 vein tip
101	<i>flp</i>	31.0±	flap wings	Wings concave; eyes bulging, rough
102	<i>ny</i>	32.0±	notchy	Wing tips nicked
103	<i>en-we</i>	32.0±	enhancer of white-eosin	With <i>w<sup>e</sup></i> alleles, gives nearly white eyes; suppresses <i>f</i>
104	<i>sto</i>	32.5	stocky	Fly short; eyes large, pear-shaped
105	<i>clm</i>	32.6±	clumpy marginals	Marginal wing hairs clumped
106	<i>ras</i>	32.8	raspberry	Eyes dark ruby
107	<i>ww</i>	32.9±	wider-wing	Wings short, broad
108	<i>v</i>	33.0	vermilion (pseudo)	Eyes bright vermilion, ocelli colorless
109	<i>osh</i>	33.0±	outshifted	Wings short, divergent; body light tan
110	<i>dwx</i>	33.2	dwarfex	Body small, wings coarse
111	<i>sbr</i>	33.4	small bristle	Bristles small, some missing
112	<i>csk</i>	33.4±	costakink	Wings reduced, costal vein kinked
113	<i>bla</i>	33.6±	bladder-wing	Wings deformed, with bladders; male sterile
114	<i>m</i>	36.1+	miniature	Wings small, dark
115	<i>dy</i>	36.2-	dusky	Wings small, dark
116	<i>ty-l</i>	36.4	tiny-like	Bristles short, fine
117	<i>trb</i>	37.0±	thread bristle	Bristles short, fine
118	<i>fw</i>	38.3	furrowed	Eyes furrowed, scutellum short, bristles gnarled
119	<i>alo</i>	38.3±	alopecia	Microchaetes nearly absent
120	<i>ups</i>	40.8	upright scutellars	Posterior scutellars vertical
121	<i>som</i>	40.8	sombre	Body dark, eyes dull

continued

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
X Chromosome [11, 43]				
122	<i>up</i>	41.0±	upheld	Wings held upright
123	<i>pun</i>	41.1±	purty	Fly small, late-hatching
124	<i>taw</i>	41.1±	tawny	Head and thorax dark, abdomen light
125	<i>wy</i>	41.9	wavy	Wings waved, curled upward
126	<i>kk</i>	42.0±	kinky	Bristles bent or forked
127	<i>s</i>	43.0	sable	Body dark
128	<i>cop</i>	43.3±	copper	Eyes brownish red
129	<i>ten</i>	43.9	tenuischaete	Bristles short, thin; body small
130	<i>g</i>	44.4	garnet (pseudo)	Eyes garnet pink
131	<i>ty</i>	44.5	tiny	Bristles, body small; female sterile
132	<i>na</i>	45.2	narrow abdomen	Abdomen cylindrical; female sterile
133	<i>shp</i>	47.5±	shrimp	Overall size reduction
134	<i>thb</i>	47.6±	thin bristle	Bristles thin
135	<i>pl</i>	47.9	pleated	Wings pleated longitudinally
136	<i>rim</i>	48.1±	rimy	Eyes brownish with white hairs; wings pleated
137	<i>sge</i>	48.4±	shifted genitals	Genitalia rotated
138	<i>thm</i>	48.9	thin-macros	Macrochaetes thin
139	<i>vb</i>	49.3	vibrissae	Vibrissae in tuft
140	<i>mgt</i>	49.6±	midget	Body small; late-hatching
141	<i>thv</i>	49.7±	thick vein	Wing veins thick; eyes small, dark
142	<i>sla</i>	50.0±	slimma	Body narrow
143	<i>sd</i>	51.5	scalloped	Wing margins excised
144	<i>exi</i>	51.5±	exiguous	Body small, dark
145	<i>tc</i>	51.6±	tinychaete	Bristles fine
146	<i>Bg</i>	51.6	Bag	Wings short, blunt, inflated
147	<i>sml</i>	51.9±	small thorax	Head and thorax small
148	<i>dru</i>	52.3±	droopy wing	Fly small, wings drooped; male sterile
149	<i>ber</i>	52.2±	berrytail	Abdomen narrow, with berrylike posterior protrusion bearing abnormal genitalia
150	<i>msc</i>	52.6±	melanoscutellum	Scutellum dark; eyes and wings abnormal in shape
151	<i>Shw</i>	53.3	Shaker-downheld	Legs, abdomen shake under ether; wings droop
152	<i>sl</i>	53.5	small-wing	Wings short, oblong; eyes large
153	<i>mc</i>	54.0	microchaete	Hairs irregular, bristles small
154	<i>un</i>	54.4	uneven	Eyes rough, small
155	<i>r9</i>	54.5	rudimentary <sup>9</sup>	Wings truncated; female sterile
156	<i>acc</i>	54.5±	acclinal wing	Wings upheld, sloping
157	<i>if3</i>	55.0±	inflated <sup>3</sup>	Wings inflated, veins thickened
158	<i>M(1)0</i>	56.6	Minute (1) 0	Minute (bristles fine, viability low, homozygous lethal)
159	<i>f</i>	56.7	forked (pseudo)	Bristles short, gnarled
160	<i>B</i>	57.0 (Dp)	Bar	Eyes narrow bar in homozygote, kidney-shaped in heterozygote
161	<i>der</i>	57.2±	deranged	Thoracic bristles disarranged; wings upheld
162	<i>Sh</i>	58.0	Shaker	Legs, abdomen shake under light ether
163	<i>siw</i>	58.5±	side wing	Wings held parallel to sides of abdomen
164	<i>od</i>	59.2	outstretched	Wings divergent
165	<i>sv</i>	59.2	small-eye	Eyes small, rounded
166	<i>Bx</i>	59.4	Beadex	Wings excised
167	<i>rwg</i>	59.5±	reduced wings	Wings short, upheld; wing hairs disarranged
168	<i>fu</i>	59.5	fused	Wing veins fused; ocelli, ocellar bristles reduced or absent
169	<i>hdp</i>	59.6±	heldup	Wings upheld
170	<i>bk</i>	59.8±	buckled	Wings misshapen, divergent
171	<i>crk</i>	60.1±	crooked setae	Bristles disarranged, abnormal
172	<i>smd</i>	60.1±	smalloid	Body size reduced
173	<i>ton</i>	60.1±	tonochaete	Bristles short, fine
174	<i>meg</i>	61.9±	megaoculus	Eyes rough; eyes, wings abnormally shaped
175	<i>M(1)36f</i>	62.0±	Minute (1) 36f	Slight minute (bristles fine, homozygous lethal)
176	<i>car</i>	62.5	carnation	Eyes dark ruby
177	<i>M(1)n</i>	62.7	Minute (1) n	Minute type (fine bristles, low viability, homozygous lethal)
178	<i>fo</i>	63.0±	folded	Wings unexpanded
179	<i>kno</i>	63.9±	knobbyhead	Head small, abnormal; male infertile
180	<i>sw</i>	64.0	short-wing	Wings trimmed, warped; eyes reduced, rough
181	<i>su-f</i>	64.0±	suppressor of forked	Certain <i>f</i> alleles made nearly +

continued



## 5. LINKAGE GROUPS: INVERTEBRATES

### Part 1. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
X Chromosome [11,43]				
182	<i>mel</i>	64.1±	melanized	Body slightly dark, eyes dull red
183	<i>wa-l</i>	64.4±	warty-like	Ommatidia disarranged
184	<i>ot</i>	65.1±	outheld	Wings held out; male inviable, sterile
185	<i>bb<sup>5</sup></i> (called <i>bb</i> )	66.0	hobbed	Bristles small, sclerites irregular
Chromosome II [1]				
186	<i>net</i>	0-	net	Extreme plexus venation
187	<i>al</i>	0	aristaless	Aristae reduced, scutellars divergent
188	<i>l(2)gl</i>	0±	lethal (2) giant larva	Larval lethal
189	<i>ocr</i>	0	ochracea	Eye color light, darkening with age
190	<i>ex</i>	0.1	expanded	Wings broad, spread; eyes rough
191	<i>ds</i>	0.3	dachsous	Wings shorter, crossveins closer
192	<i>S</i>	1.3	Star (pseudo)	Eyes small, rough; homozygous lethal
193	<i>Su-S</i>	1.3±	Suppressor of Star	Suppresses <i>S</i> ; <i>Su-S/S</i> is +
194	<i>ast</i>	1.3±	asteroid	Eyes small, rough
195	<i>shr</i>	2.3±	shrunken	Body small, wizened
196	<i>shv</i>	3.8±	short vein	Constant terminal gaps in veins L2 and L4
197	<i>ho</i>	4.0	heldout	Wings extended
198	<i>fes</i>	5.0±	female-sterile	Eggs do not develop
199	<i>E-S</i>	6.0±	Enhancer of Star	Increases expression of <i>S</i>
200	<i>Cy</i>	7.0	Curly	Wings curled upward; homozygous lethal
201	<i>l(2)ay</i>	8.3	lethal (2) ay	Lethal
202	<i>Dt</i>	10.0±	Detached	Vein L2 does not reach margin
203	<i>ang</i>	10.5	angle wing	Wings held up from dorsal surface
204	<i>ed</i>	11.0	echinoid	Eyes large, rough
205	<i>M(2)C</i>	11.0-12.0 (Df)	Minute (2) Curry	Fairly strong minute
206	<i>ft</i>	12.0	fat	Body short, fat; scutellar bristles far apart
207	<i>G</i>	12.0	Gull	Wings large, spread; homozygous lethal
208	<i>M(2)z</i>	12.9±	Minute (2) z	Medium minute
209	<i>M(2)B</i>	13.0 (Df)	Minute (2) Bridges	Medium minute
210	<i>dp</i>	13.0	dumpy	Wings truncated; vortices on thorax
211	<i>dw-24F</i>	13.0±	dwarf in 24F	Eyes dull, body dwarfed
212	<i>M(2)S1</i>	15.0	Minute (2) Schultz' 1	Strong minute
213	<i>l(2)cg</i>	15.0±	lethal (2) comb-gap	Lethal from <i>cg</i> stock
214	<i>Sk</i>	16.0	Streak	Central streak on thorax; homozygous lethal
215	<i>thv</i>	16.0±	thick-veins	Veins thick, irregular
216	<i>cl</i>	16.5	clot	Eye color maroon, close to sepia ( <i>se</i> ); male sterile
217	<i>pi</i>	17.0±	ped	Facets jumbled
218	<i>Sp</i>	22.0	Sternopleural	Extra sternopleural bristles; homozygous lethal
219	<i>spd</i>	22.3±	spade	Wings shortened, broad
220	<i>gt-4</i>	24.0	giant-4	Giant fly
221	<i>d</i>	31.0	dachs	Tarsi 4-jointed, venation shifted
222	<i>fy</i>	33.0±	fuzzy	Thoracic hairs fuzzy
223	<i>fol</i>	39.0±	folded wings	Wings folded; overlap
224	<i>da</i>	39.3±	daughterless	Homozygous female produces no daughters
225	<i>J</i>	41.0	Jammed	Wing narrow strip
226	<i>M(2)S11</i>	43.0±	Minute (2) Schultz' 11	Slight minute
227	<i>ab</i>	44.0	abrupt	Shortened L5 vein, scutellars few
228	<i>oph</i>	45.0±	ophthalmopedia	Eyes kidney-shaped or with appendage
229	<i>rk</i>	46.0±	ricketts	Segments of legs flattened and bent
230	<i>l(2)bs<sup>3</sup>-d</i>	46.0±	lethal (2) with <i>bs<sup>3</sup>-d</i>	Lethal
231	<i>M(2)e</i>	46.0±	Minute (2) e	Medium minute
232	<i>b</i>	48.5	black	Body, legs, veins black
233	<i>j</i>	48.7	jaunty	Wings upturned
234	<i>el</i>	50.0	elbow	Wings bent, alulae and balancers small
235	<i>lm</i>	50.0±	limited	Sternites small; female sterile
236	<i>M(2)S13</i>	50.0±	Minute (2) Schultz' 13	Strong minute
237	<i>l(2)H</i>	50.0±	lethal (2) Humphrey	Pupal semilethal
238	<i>Su-H</i>	50.5	Suppressor of Hairless	Homozygous lethal
239	<i>rd</i>	51.0	reduced	Bristles small, irregular; female sterile
240	<i>pu</i>	51.0±	pupal	Wings unexpanded

continued



## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Chromosome II [1]				
241	<i>bys</i>	52.0±	polychaetous	Extra and double bristles
242	<i>cr-u</i>	52.5±	cream-underscored	Specific dilutor of <i>w<sup>e</sup></i> and Pale; male sterile
243	<i>nub</i>	53.0	nubbin	Wings very small and thin with tendency to curve up or down
244	<i>ck</i>	53.0±	crinkled	Wings flimsy
245	<i>rdo</i>	53.0±	reduced ocelli	Ocelli reduced in size, color moved to region between ocelli
246	<i>l(2)Bld</i>	53.1	lethal (2) opposite T (1,2) Bld	Lethal
247	<i>M(2)S5</i>	53.5	Minute (2) Schultz' 5	Medium minute
248	<i>hk</i>	53.9	hook	Bristles bent or barbed
249	<i>bri</i>	54.3±	bright	Eye color bright red
250	<i>pr</i>	54.5	purple	Eye color purplish ruby
251	<i>rn</i>	54.5±	rotund	Wings round, tarsi 3-jointed; sterile
252	<i>rh</i>	54.7±	roughish	Eyes moderately rough
253	<i>Bl</i>	54.8	Bristle	Bristles short, beaded; homozygous semilethal
254	<i>Alu</i>	54.9	Alula	Alula fused to wing; wing warped
255	<i>Jag</i>	54.9	Jagged	Wings nicked, eyes rough
256	<i>lt</i>	55.0-	light	Eye color yellowish pink
257	<i>tri</i>	55.0±	trident	Thorax darkened
258	<i>M(2)D</i>	55.0± (Df)	Minute (2) D	Body color and bristles pale
259	<i>rl</i>	55.1-	rolled	Wing edges rolled, frayed
260	<i>M(2)S2</i>	55.1 (Df)	Minute (2) Schultz' 2	Minute type
261	<i>M(2)S4</i>	55.1 (Df)	Minute (2) Schultz' 4	Medium minute
262	<i>M(2)S8</i>	55.1 (Df)	Minute (2) Schultz' 8	Slight minute
263	<i>M(2)S10</i>	55.1 (Df)	Minute (2) Schultz' 10	Slight minute
264	<i>stw</i>	55.1	straw	Body, wings, bristles yellow
265	<i>blt</i>	55.2±	blot	Wings inflated, blackened
266	<i>Cu</i>	55.2±	Curl	Lateral compression and indentation-fold of unfolded imaginal wing
267	<i>tk</i>	55.3	thick	Legs, tarsi thickened; wings short
268	<i>pk</i>	55.3	prickle	Bristles, hairs irregular
269	<i>ap</i>	55.4	apterous	Wings, balancers missing
270	<i>msf</i>	55.6-	misformed	Eyes misformed, wings crumpled
271	<i>bur</i>	55.7±	burgundy	Dull, darkish-brown eye color
272	<i>ti</i>	55.9	tarsi irregular	Tarsal segments fused, eyes rough
273	<i>ltd</i>	56.0±	lightoid	Eye color translucent yellowish pink, ocelli colorless
274	<i>M(2)S12</i>	56.0±	Minute (2) Schultz' 12	Slight minute
275	<i>std</i>	56.5±	staroid	Eyes small, very rough; male sterile
276	<i>ta</i>	56.6±	tapered	Wings narrow and pointed, veins close
277	<i>dil</i>	57.0±	specific dilutor	Dilutor of <i>bw</i> and <i>w</i> alleles
278	<i>buo</i>	57.1	burnt orange	Eye color orange brown
279	<i>M(2)38b</i>	57.0±	Minute (2) 38b	Extreme minute
280	<i>cn</i>	57.5	cinnabar	Eye color bright scarlet, ocelli colorless
281	<i>puf</i>	58.0±	puff	Wings blistered
282	<i>blo</i>	58.5	bloated	Wings ballooned, extra veins
283	<i>smk</i>	58.6±	smoky	Body color dark
284	<i>Np</i>	58.7-60.2 (Df)	Notopleural	Bristles short, wings broad; homozygous lethal
285	<i>at</i>	60.1±	arctus oculus	Number of facets reduced
286	<i>arch</i>	60.5±	arch	Wings downcurved in both axes
287	<i>ad</i>	60.7	arcoid	Wings arched, broad, short; crossveins close
288	<i>chl</i>	60.8	chaetelle	Bristles very small, slight plexus
289	<i>whd</i>	61.0±	withered	Wings warped or shrunken
290	<i>tom</i>	61.5±	tomboy	Homozygous female with male-like pigmentation of posterior tergites
291	<i>en</i>	62.0	engrailed	Scutellar notch, broken veins, extra sex comb
292	<i>upw</i>	62.0±	upward	Wings upturned
293	<i>l(2)rn</i>	63.0±	lethal (2) with rotund	Lethal
294	<i>Bkd</i>	65.0±	Blackoid	Dark body color
295	<i>M(2)40c</i>	65.0±	Minute (2) 40c	Minute type
296	<i>po</i>	65.2	pale-ocelli	Ocelli nearly colorless
297	<i>sca</i>	66.7	scabrous	Eyes rough, some bristles missing
298	<i>vg</i>	67.0	vestigial	Wings, balancers vestigial
299	<i>l(2)C</i>	67.0	lethal (2) Curry	Lethal before pupal stage

continued

# Contrails

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part 1. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Chromosome II [1]			
300	<i>wx</i>	waxy	Wings heavy, waxy; male sterile
301	<i>UH20</i>	Upturned H20	Wings curled
302	<i>l(2)mr<sup>2</sup></i>	lethal (2) with morula <sup>2</sup>	Lethal
303	<i>Pfd</i>	Puffed	Wings puffed, divergent; homozygous lethal
304	<i>bat</i>	bat	Wings extended, bent back
305	<i>cg</i>	comb-gap	Sex combs large; gap in wing vein L4; female sterile
306	<i>dr</i>	droopy	Wings spread wide apart and drooping
307	<i>sf</i>	safranin	Eye color dark chocolate
308	<i>L</i>	Lobe	Eyes small, nicked at anterior edge
309	<i>kn</i>	knot	Veins L3 and L4 close; eyes oblique
310	<i>ch</i>	chubby	Larva, pupa, adult short
311	<i>dke</i>	dark eye	Eye color soft, dark, dull, with tiny fleck
312	<i>gp</i>	gap	Vein L4 broken
313	<i>c</i>	curved	Wings thin, spread, lifted, curved
314	<i>Wr</i>	Wrinkled	Wings wrinkled; suppresses <i>L</i>
315	<i>M(2)S7</i>	Minute (2) Schultz' 7	Strong minute
316	<i>pw-c</i>	pink-wing-c	Eye color dilute, wings short, blunt
317	<i>fr</i>	fringed	Wing margins ragged
318	<i>ff</i>	four-jointed	Tarsi 4-jointed; wings short
319	<i>rf</i>	roof wings	Wings drooped at sides
320	<i>wl</i>	welt	Eyes seamed, reduced
321	<i>abr</i>	abero	Abdominal bands irregular; wings frayed, eyes rough; female sterile
322	<i>nw</i>	narrow	Wings narrow
323	<i>I-f</i>	Intensifier of forked	Enhances <i>f</i>
324	<i>sm</i>	smooth	Abdomen hairless; sterile
325	<i>M(2)173</i>	Minute (2) 173	Moderate minute
326	<i>ky</i>	humpy	Thorax ridged, wings truncated
327	<i>l(2)Su-H</i>	lethal (2) from Suppressor of Hairless	Lethal
328	<i>a</i>	arc	Wings broad, bent down, crossveins closer
329	<i>M(2)l</i>	Minute (2) 1	Extreme minute
330	<i>px</i>	plexus	Network of extra veins
331	<i>pa</i>	patulous	Wings spread wide apart
332	<i>M(2)l<sup>2</sup></i>	Minute (2) 1 <sup>2</sup>	Slight minute
333	<i>hv</i>	heavy vein	Veins thick, posterior crossveins oblique
334	<i>l(2)bw</i>	lethal (2) brown	Probable deficiency; lethal
335	<i>bw</i>	brown	Eye color brownish to garnet
336	<i>mi</i>	minus	Bristles hairlike; body small; female sterile
337	<i>abb</i>	abbreviated	Bristles slightly reduced
338	<i>slt</i>	slight	Body small, bristles reduced; female sterile
339	<i>pd</i>	purpleoid	Eye color dark pink, like purple ( <i>pr</i> )
340	<i>ll</i>	lanceolate	Wings narrow, pointed
341	<i>mr</i>	morula	Eyes rough, bristles small; female sterile
342	<i>l(2)ax</i>	lethal (2) ax	Very early larval lethal
343	<i>sp</i>	speck	Black speck in wing axil; body color olive
344	<i>or</i>	orange	Bright orange eye color
345	<i>Px</i>	Plexate	Venation as in blistered mutation ( <i>bs</i> ); veins thickened, broken; homozygous lethal
346	<i>bs</i>	blistered	Wings blistered, small; extra veins
347	<i>Pin</i>	Pin	Thoracic bristles pinlike
348	<i>ba</i>	balloon	Wings inflated, extra veins
349	<i>M(2)33a</i>	Minute (2) 33a	Strong minute
Chromosome III [1]			
350	<i>ru</i>	roughoid	Eyes small, rough; erupted facets
351	<i>mp</i>	microptera	Wings small, ballooned; tarsi 4-jointed
352	<i>aa</i>	anarista	Aristae small, without branches
353	<i>ve</i>	veinlet	Longitudinal wing veins interrupted
354	<i>R</i>	Roughened	Eyes rough; homozygous semilethal
355	<i>rai</i>	raisin	Deep brown eye color
356	<i>ju</i>	javelin	Bristles and hairs cylindrical
357	<i>dv</i>	divergent	Wings spread

continued

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Chromosome III [1]			
358	<i>Me</i> 20.0±	Moire	Eye color brownish, 7 flecks; homozygous lethal
359	<i>Hn</i> 23.0	Henna	Eye color dull, dark; homozygous lethal
360	<i>be-3</i> 25.0±	benign tumor in 3	Nonlethal melanotic tumors
361	<i>se</i> 26.0	sepia	Eye color brownish red, darkening to black
362	<i>su-t</i> 26.0±	suppressor of tan	Converts <i>t</i> to +
363	<i>h</i> 26.5	hairy	Extra hairs on scutellars, veins, pleurae, and head
364	<i>abd</i> 27.0±	abdominal	Abdominal bands broken, etched
365	<i>rs</i> 35.0	rose	Eye color translucent pink
366	<i>eyg</i> 35.5	eye-gone	Eyes and head reduced
367	<i>gv</i> 36.2	grooved	Longitudinal medial groove in thorax
368	<i>cr-3</i> 36.5±	cream in 3	Specific dilutor of <i>w<sup>e</sup></i> eye color
369	<i>rt</i> 37.0±	rotated	Abdomen twisted counterclockwise
370	<i>app</i> 37.5	approximated	Crossveins close; tarsi 4-jointed
371	<i>pyd</i> 39.0±	polychaetoid	Extra bristles
372	<i>M(3)S37</i> 39.7±	Minute (3) Schultz' 37	Extreme minute
373	<i>H</i> 40.0	tilt	Wings spread, warped, with gap in vein L3
374	<i>M(3)33j</i> 40.2 (Df)	Minute (3) 33j	Medium minute
375	<i>M(3)h</i> 40.2	Minute (3) h	Medium minute; allele of <i>M(3)33j</i>
376	<i>M(3)y</i> 40.2	Minute (3) y	Medium minute; allele of <i>M(3)33j</i>
377	<i>vo-3</i> 40.4±	vortex in 3	Intensifier of <i>dp<sup>v</sup></i>
378	<i>D</i> 40.4+	Dichaete	Wings spread; homozygous lethal
379	<i>Ly</i> 40.5 (Df)	Lyra	Wings cut, narrow; homozygous lethal
380	<i>Gl</i> 41.4	Glued	Eyes small, facets rounded; homozygous lethal
381	<i>fz</i> 41.7±	frizzled	Thoracic hairs, bristles turn toward midline
382	<i>rp</i> 41.7±	rotated-penis	Male genitalia rotated; male sterile
383	<i>wk</i> 42.0±	weak	Bristles weak, irregular; body small
384	<i>Wi</i> 43.0	Washed eye	Modified <i>w</i> ; homozygous lethal
385	<i>th</i> 43.2	thread	Aristae threadlike, without branches
386	<i>mb</i> 43.4±	minusbar	Modified <i>B</i> to larger eye
387	<i>Cm</i> 43.5±	Crimp	Posterior wing edge crimped; homozygous lethal
388	<i>bul</i> 43.6	bulge	Eyes bulging, wings squared off
389	<i>M(3)S38</i> 44.0±	Minute (3) Schultz' 38	Strong minute
390	<i>st</i> 44.0	scarlet	Eye color scarlet, ocelli white
391	<i>tra</i> 45.0±	transformed	Transforms female to normal-appearing male
392	<i>cp</i> 45.3	clipped	Wing margins clipped
393	<i>mot-28</i> 46.0	mottled-28	Eyes mottled with brown
394	<i>W</i> 46.0	Wrinkled	Wings incompletely unfolded, pebbled
395	<i>as</i> 46.0±	ascute	Wings held downward
396	<i>je</i> 46.0±	jelly	Eye color dark pinkish
397	<i>Pdr</i> 46.0±	Purpleoider	Intensifier of <i>pd</i>
398	<i>in</i> 46.9	inturned	Thoracic bristles directed toward midline
399	<i>M(3)S39</i> 47.0±	Minute (3) Schultz' 39	Strong minute
400	<i>dn</i> 47.0±	doughnut	Eye of <i>se dn</i> with light central spot; male sterile
401	<i>ri</i> 47.1	radius incompletus	Vein L2 shows gap
402	<i>eg</i> 47.3	eagle	Wings spread, raised
403	<i>Dfd</i> 47.5	Deformed	Eyes small; homozygous lethal
404	<i>wp</i> 47.5	warped	Wings spread, doubly warped
405	<i>pb</i> 47.7	proboscipedia	Mouth parts footlike; adult lethal; female sterile
406	<i>p</i> 48.0	pink	Eye color dull ruby
407	<i>Bb</i> 48.0±	Bubble	Wings small, inflated; male sterile; homozygous female lethal
408	<i>bod</i> 48.3	bowed	Wings arched
409	<i>fel</i> 48.5	tetraletera	Wings haltere-like
410	<i>by</i> 48.7	blistery	Wings blistered distally
411	<i>M(3)S34</i> 49.0±	Minute (3) Schultz' 34	Slight minute
412	<i>ma</i> 49.7	maroon	Eye color dull ruby
413	<i>cu</i> 50.0	curled	Wings upcurved, body dark, postscutellars crossed
414	<i>M(3)S31</i> 50.0 (Df)	Minute (3) Schultz' 31	Medium minute
415	<i>mu</i> 50.0±	mussed	Wings thin, crumpled
416	<i>ry</i> 51.0±	rosy	Eye color deep ruby
417	<i>kar</i> 52.0	karmoisin	Eye color like scarlet mutation ( <i>st</i> ) but duller, ocelli colorless

continued



## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Chromosome III [1]				
418	<i>c3G</i>	55.0±	crossover suppressor in 3 of Gowen	Eliminates crossing over
419	<i>red</i>	55.5±	red	Red malpighian tubules
420	<i>jvl</i>	56.7	javelin-like	Bristles cylindrical, crooked
421	<i>cv-c</i>	57.9	crossveinless-c	Posterior crossvein absent or reduced
422	<i>Sb</i>	58.2	Stubble	Bristles short, thick; homozygous lethal
423	<i>ss</i>	58.5	spineless	Bristles very small
424	<i>bx</i>	58.8	bithorax (pseudo)	Balancers winglike; metathorax resembles mesothorax
425	<i>Rf</i>	59.0±	Roof	Wings drooping at sides
426	<i>cal</i>	59.5±	coal	Black body color, similar to <i>e</i> <sup>4</sup>
427	<i>fl</i>	59.9	fluted	Wings creased, darkish
428	<i>sr</i>	62.0	stripe	Dark dorsal stripe
429	<i>M(3)f</i>	62.4	Minute (3) f	Minute type
430	<i>gl</i>	63.1	glass	Eye color dilute, facets fused
431	<i>gl-l</i>	64.0±	glass-like	Eyes orange, rough, and small
432	<i>k</i>	64.0±	kidney	Eyes kidney-shaped
433	<i>M(3)S35</i>	64.0±	Minute (3) Schultz' 35	Extreme minute
434	<i>sed</i>	64.5±	sepioid	Eye color chocolate
435	<i>cv-d</i>	65.0±	crossveinless-d	Posterior crossvein absent or reduced
436	<i>Cur</i>	66.0±	Curl	Curly wings; homozygous lethal
437	<i>Dl</i>	66.2	Delta	Veins thick at margin; homozygous lethal
438	<i>H</i>	69.5	Hairless	Some bristles and hair missing; homozygous lethal
439	<i>e</i>	70.7	ebony	Body color black
440	<i>det</i>	72.5	detached	Crossveins broken, wings folded under
441	<i>cd</i>	75.7	cardinal	Eye color dull scarlet, ocelli white
442	<i>wo</i>	76.2	white ocelli	Ocelli colorless
443	<i>obt</i>	77.5±	obtuse	Wings short, blunt
444	<i>bar-3</i>	79.1	bar-3	Phenotype like <i>B/B</i>
445	<i>M(3)124</i>	79.7	Minute (3) 124	Strong minute; allele of <i>M(3)w</i>
446	<i>M(3)B</i>	79.7	Minute (3) Burkart	Moderate minute; allele of <i>M(3)w</i>
447	<i>M(3)B<sup>2</sup></i>	79.7	Minute (3) Bridges	Strong minute; allele of <i>M(3)w</i>
448	<i>M(3)w</i>	79.7	Minute (3) w	Strong minute
449	<i>l(3)a</i>	79.7	lethal (3) first found	Lethal; allele of <i>M(3)w</i>
450	<i>M(3)Fla</i>	80.0±	Minute (3) Florida	Strong minute; allele of <i>M(3)w</i>
451	<i>M(3)36e</i>	84.5	Minute (3) 36e	Medium minute
452	<i>M(3)be</i>	87.0±	Minute (3) beta	Medium minute
453	<i>mah</i>	88.0±	mahogany	Eye color brownish, darkening
454	<i>Pr</i>	90.0	Prickly	Bristles vestigial; homozygous semilethal
455	<i>M(3)j</i>	90.2	Minute (3) j	Extreme minute
456	<i>l(3)PR</i>	90.2	lethal with In(3R)P	Lethal; allele of <i>M(3)j</i>
457	<i>lx</i>	91.0±	taxi	Wings divergent
458	<i>ro</i>	91.1	rough	Eyes rough, small
459	<i>l(3)XaR</i>	91.8	lethal (3) XaR	Balancer of <i>T(2,3)Xa</i>
460	<i>cmp</i>	93.0±	crumpled	Wings smaller, crumpled
461	<i>Bd</i>	93.8	Beaded	Wing margins excised; homozygous lethal
462	<i>Pw</i>	94.1	Pointed-wing	Wings pointed at tip; homozygous lethal
463	<i>hf</i>	95.0±	brief	Body small, bristles minute-like; male sterile
464	<i>rsd</i>	95.4	raised	Wings rise straight up
465	<i>su<sup>B</sup>-pr</i>	95.5	suppressor of purple	Suppresses purple ( <i>pr</i> )
466	<i>ra</i>	97.3	rase	Bristles, hairs smaller, fewer
467	<i>Dp</i>	99.3±	Duplication	Similar to ultra bar
468	<i>ld</i>	100.0±	loboid	Eyes lobe-like
469	<i>ca</i>	100.7	claret	Eye color clear ruby
470	<i>M(3)l</i>	101.0	Minute (3) l	Medium minute
471	<i>bv</i>	104.3	brevis	Bristles short, stubby
472	<i>M(3)g</i>	106.2	Minute (3) g	Slight minute, requires <i>E-M(3)g</i>
Chromosome IV [40]				
473	<i>ci</i>	0	cubitus-interruptus	Vein L4 interrupted
474	<i>M-4</i>	0-0.2±	Minute-4	Medium minute; deficiency for <i>ci</i> , <i>ar</i> , <i>gul</i> , and <i>Scn</i>
475	<i>ar</i>	0-0.2	abdomen rotatum	Abdomen twisted clockwise
476	<i>gul</i>	0.2	grooveless	Scutellar groove diminished

continued



## 5. LINKAGE GROUPS: INVERTEBRATES

### Part I. FRUIT FLY

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Chromosome IV [40]				
477	<i>bt</i>	1.4	bent	Wings bent, legs knobby
478	<i>ey</i>	2.0	eyeless	Eyes small or absent
479	<i>sv</i>	3.0	shaven	Abdominal bristles fewer

*Contributors:* (a) Warren, Katherine Brehme, (b) Novitski, E.

*References:* [1] Bridges, C. B., and K. S. Brehme. 1944. Carnegie Inst. Wash. Publ. 552. [2] Buzzati-Traverso, A. 1948. *Drosophila Inform. Serv.* 22:66. [3] Chino, M. 1936. *Japan. J. Genet.* 12:205. [4] Chino, M. 1937. *Ibid.* 13:105. [5] Chino, M. 1939. *Drosophila Inform. Serv.* 11:32. [6] Christie, A. L. M. 1939. *J. Genet.* 39:58. [7] Clarke, J. M. 1951. *Drosophila Inform. Serv.* 25:94. [8] Clarke, J. M. 1952. *Ibid.* 26:87. [9] Demerec, M. 1954. *Ibid.* 28:93. [10] Donald, H. P. 1936. *J. Genet.* 33:105. [11] Fahny, M. B., and O. Fahny. 1957-60. *Drosophila Inform. Serv.* 31-34. [12] Ferry, R. M., R. C. Lancefield, and C. W. Metz. 1923. *J. Heredity* 14:373. [13] Gordon, C., H. Spurway, and P. A. R. Street. 1939. *J. Genet.* 38:45. [14] Haldane, J. B. S. 1945. *Drosophila Inform. Serv.* 19:56. [15] Jermyn, J. E., et al. 1943. *Ibid.* 17:52. [16] Kiiil, V. 1946. *Ibid.* 20:82. [17] Kikkawa, H. 1938. *Genetica (Haag)* 20:458. [18] Lamy, R. 1944. *Drosophila Inform. Serv.* 18:52. [19] Lancefield, D. E. 1922. *Genetics* 7:375. [20] Lancefield, R. C., and C. W. Metz. 1922. *Am. Naturalist* 56:211. [21] Mainx, F. 1949. *Drosophila Inform. Serv.* 23:78. [22] Mainx, F. 1950. *Ibid.* 24:77. [23] Milani, R. 1949. *Ibid.* 23:78. [24] Miller, D. D. 1954. *Ibid.* 28:100. [25] Moriwaki, D. 1935. *Genetica (Haag)* 17:41. [26] Moriwaki, D. 1938. *Japan. J. Genet.* 14.1: [27] Moriwaki, D. 1949. *Drosophila Inform. Serv.* 23:77. [28] Osima, T. 1940. *Cytologia (Tokyo)* 10:450. [29] Osima, T. 1940. *Drosophila Inform. Serv.* 13:55. [30] Patterson, J. T., and W. S. Stone. 1952. *Evolution in the genus Drosophila*. Macmillan, New York. [31] Spassky, B., S. Zimmering, and T. Dobzhansky. 1950. *Heredity* 4:189. [32] Spencer, W. P. 1935. *Drosophila Inform. Serv.* 4:48. [33] Spencer, W. P. 1944. *Ibid.* 18:51. [34] Spencer, W. P. 1949. *Genetics, paleontology, and evolution*. Princeton Univ. Press, Princeton, N. J. [35] Spiess, E. B. 1952. *Drosophila Inform. Serv.* 26:87. [36] Spurway, H. 1939. *Ibid.* 12:54. [37] Spurway, H. 1951. *Ibid.* 25:95. [38] Sturtevant, A. H. 1929. Carnegie Inst. Wash. Publ. 399. [39] Sturtevant, A. H. 1940. *Genetics* 25:343. [40] Sturtevant, A. H. 1951. *Proc. Natl. Acad. Sci. U. S.* 37:405. [41] Sturtevant, A. H., and E. Novitski. 1941. *Genetics* 26:517. [42] Sturtevant, A. H., and C. C. Tan. 1937. *J. Genet.* 34:415. [43] Warren, K. B. Unpublished. Natl. Institutes of Health, Bethesda, Md., 1962.

### Part II. PARASITIC WASP

*Habrobracon juglandis* has 10 pair of chromosomes; linkage groups have been found for 8 pair. **Linkage** (column B): slant line (/) indicates complete linkage.

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	(E)	
Linkage Group I					
1	<i>Sk</i>	<i>Sk--r</i>	12	Speckled	Bright red flecks of pigment in white eye
2	<i>r</i>	<i>r--gl</i>	13	reduced	Small wings; reduced, irregular venation
3	<i>gl</i>	<i>gl--x</i>	30	glass	Small eyes, lacking facet outlines
4	<i>X</i>	<i>X--fu</i>	10	Sex	9 alleles known (each consisting of many factors determining sex differences) that produce similar phenotypes in males and in females
5	<i>fu</i>	<i>fu--sb</i>	22	fused	Antennal segments fused; tarsal segments lacking or fused

*continued*

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part II. PARASITIC WASP

Gene Symbol	Linkage	Recombination Percentage	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	(E)	
Linkage Group I					
6	<i>sb</i>	<i>sb--bl</i>	42	stubby	Males with antennae 7-9 segments long; females with antennae 5-7 segments long
7	<i>bl</i>	<i>bl--le</i>	30	black	Body color black
8	<i>le</i>	<i>le--c</i>	12	lemon	Body color pale lemon yellow
9	<i>c</i>	<i>c--l</i>	14	cantaloupe	Eyes light pink, darken to deep red
10	<i>l</i>	<i>l--n</i>	3	long	Antennal segments elongated; leg segments longer and thinner than in wild type
11	<i>n</i>	<i>n--ho</i>	7	narrow	Narrow wings; cuts off irregular slices of costal and inner wing margins
12	<i>ho</i>	<i>ho--vl</i>	8	honey	Body lacks black pigment entirely
13	<i>vl</i>	<i>vl--ro</i>	15	veinless	Wing veins missing, except along costal margin
14	<i>ro</i>	<i>ro--bu</i>	12	rough	4th radius vein absent, adjacent veins roughened
15	<i>bu</i>	<i>bu--cr</i>	37	bulged	Eyes abnormally bulged transversely
16	<i>cr</i>	<i>cr--sl/co</i>	41	crescent	Eyes small; ocelli crescent-shaped, pigment reduced
17	<i>sl</i>	<i>sl/co--ct</i>	33	semilong	Antennal and leg segments lengthened
18	<i>co</i>			coalescent	Antennal segments coalescent
19	<i>ct</i>	<i>ct--rd</i>	32	cut	Outer wing margin indented or straightened, giving cut appearance
20	<i>rd</i>	<i>rd--gy</i>	37	red	Eye color varies from light red to dark red, almost black with temperature increase
21	<i>gy</i>	<i>gy--ac/el</i>	7	gynoid	Short antennae in male, resembling those in female; abdominal sclerites resemble those in female
22	<i>ac</i>	<i>ac/el--gy</i>	7	aciform	Terminal half of antennae very slender, needlelike
23	<i>el</i>			eyeless	Head malformed; eye rudiments present
Linkage Group II					
24	<i>k</i>	<i>k--dw</i>	28	kidney	Eyes kidney-shaped
25	<i>dw</i>	<i>dw--m</i>	5	dwindling	Irregularity and fusion of antennal segments
26	<i>m</i>	<i>m--o</i>	11	miniature	Reduced body size; semilethal: many die as pupae
27	<i>o</i>	<i>o--m</i>	11	orange	Eyes orange, varying to pink and red
Linkage Group III					
28	<i>bk</i>	<i>bk--wh/pl</i>	25	broken	Outer margin of primary wing broken and wings fragile
29	<i>wh</i>	<i>wh/pl--st</i>	9	white	White eye; ocelli colorless
30	<i>pl</i>			pellucid	Compound eyes semitransparent
31	<i>st</i>	<i>st--wh/pl</i>	9	stumpy	Extreme reduction of tarsal segments
Linkage Group IV					
32	<i>sv</i>	<i>sv--td</i>	23	shot-veins	Wing veins broken and distorted
33	<i>td</i>	<i>td--ma</i>	27	truncated	Wings extremely reduced, irregular in shape
34	<i>ma</i>	<i>ma--td</i>	27	maroon	Light ocelli; compound eyes deep reddish brown
Linkage Group V					
35	<i>wa</i>	<i>wa--br</i>	22	wavy	Wings shortened, costal margin wavy
36	<i>br</i>	<i>br--wa</i>	22	broad	Thorax abnormally broadened
Linkage Group VI					
37	<i>ta</i>	<i>ta--un<sup>2</sup></i>	40	tapering	Antennae deficient, with much fusion and irregularity of segments distally
38	<i>un<sup>2</sup></i>	<i>un<sup>2</sup>--ta</i>	40	undulating-2	Surface of wings in undulating waves
Linkage Group VII					
39	<i>pk</i>	<i>pk/ew<sup>3</sup></i>		pink	Compound eyes pink
40	<i>ew<sup>3</sup></i>			extended wings	Wings extended in active wasps
Linkage Group VIII					
41	<i>wt</i>	<i>wt--bf</i>	17	wet	Wing microchaetae very long and irregular, giving wet appearance
42	<i>bf</i>	<i>bf--wt</i>	17	black feet	Tarsi abnormally black

Contributors: (a) Whiting, P. W., (b) Novitski, E.

References: [1] Carson, H. L. 1941. Am. Naturalist 75:608. [2] Clark, A. M. 1942. J. Heredity 33:78.

continued

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part II. PARASITIC WASP

- [3] Clark, A. M. 1943. Proc. Penna. Acad. Sci. 17:47. [4] Hesel, E. D. 1944. Am. Naturalist 78:188.  
 [5] Martin, A., Jr. 1947. An introduction to the genetics of *Habrobracon juglandis* (Ashmead). Hobson Book Press, New York. [6] Martin, A., Jr. 1947. Proc. Penna. Acad. Sci. 21:32. [7] Martin, A., Jr. 1947. Ibid. 21:36. [8] Martin, A., Jr. 1948. Univ. Pittsburgh Bull. 44:1. [9] Torvik-Greb, M. 1935. Biol. Bull. 68:25.  
 [10] Whiting, P. W. 1943. Genetics 28:365. [11] Whiting, P. W. 1946. Ibid. 32:112. [12] Whiting, P. W. 1950. J. Heredity 41:71. [13] Whiting, P. W., and L. H. Benkert. 1934. Genetics 19:268. [14] Whiting, P. W., and A. R. Whiting. 1934. J. Genet. 29:311.

### Part III. SILKWORM

*Bombyx mori* has 28 pair of chromosomes; linkage groups have been found for 15 pair. **Gene Symbol** (column A):  $\underline{l}$  = lethal.

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Linkage Group I (Z Chromosome)			
1 <i>os</i>	0	sex-linked	Low translucency of larva
2 <i>Ge</i>	14.0	Giant egg	Length and width 1.26 and 1.11, respectively, times the normal egg
3 <i>e</i>	36.4	elongate	First and second abdominal segments of larva unusually elongated
4 <i>Vg</i>	38.7	Vestigial	Wings poorly developed
5 <i>od</i>	47.6	translucent	Skin of larva shows high translucency
Linkage Group II			
6 <i>P</i>	0	Plain	Full grown larva white; $+p, p^B, p^M, p^S, p^{Sa}$ , multiple or pseudoalleles of <i>p</i>
7 <i>S</i>	6.1	New striped	Dark stripe on larva; heterozygote almost as dark as homozygote
8 <i>Gr</i>	6.9	Gray egg	Milky white shell, dark serosa pigment
9 <i>Y</i>	25.6	Yellow blood	Deep yellow hemolymph in larva
10 <i>oa</i>	26.7	mottled translucent	Mottled translucency on larval skin
11 <i>Rc</i>	31.8	Rusy	Yellowish-brown cocoon, lighter inner layer
Linkage Group III			
12 <i>Ze</i>	0	Zebra	Black band on anterior end of each segment; pair of black spots on ventral side of each larval segment
13 <i>ap</i>	0	apodal	All thoracic legs rudimentary
14 <i>lem</i>	22.8	lemon	Greenish-yellow coloring over skin visible from 2nd instar
Linkage Group IV			
15 <i>L</i>	0	Multilunar	Pairs of large brownish or yellowish round spots on thoracic and abdominal segments
16 <i>sk</i>	25.8	stick	Larva body slender and hard
17 <i>Spc</i>	33.1	Speckle	Many dark spots on larval skin; female sterile
Linkage Group V			
18 <i>pe</i>	0	pink-eyed	White egg; pigment absent from serosa
19 <i>ok</i>	4.7	kinshiryu	High translucency of larva
20 <i>re</i>	31.7	red egg	Reddish-brown serosa
21 <i>oc</i>	40.8	chinese	High translucency of larva
Linkage Group VI			
22 <i>E</i>	0	Plain supernumerary legs	Supernumerary legs in 1st and 2nd abdominal segments of larva; $E^{Ca}, E^D, E^H, E^{Kp}, E^N$ , multiple or pseudoalleles of <i>E</i>
23 <i>Nc</i>	1.4	No crescent supernumerary legs	Crescents absent; supernumerary leg in the 2nd abdominal segment
24 $+M$	3.0	Tetra molting	Standard type, larva pupates after 4th molt; $M^2, M^5$ , multiple or pseudoalleles of $+M$
25 <i>b2</i>	8.0	brown egg-2	Grayish-brown pigment in serosa
26 <i>F</i>	13.6	Flesh	Cocoon color reddish-yellow or salmon color
27 $\underline{l-k}$	17.7	lethal-k	Embryo killed few days before hatching

continued

## 5. LINKAGE GROUPS: INVERTEBRATES

### Part III. SILKWORM

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Linkage Group VII				
28	<i>q</i>	0	quail	Larval body tinted reddish-purple and covered with shredlike lines
29	<i>Gb</i>	0.7	Green b	Greenish cocoon color
30	<i>obt</i>	21.0	<i>b<sub>g</sub></i> -mottled	Moderate translucency of larva; not lethal
Linkage Group VIII				
31	<i>ae</i>	0	amylase negative	Amylase in digestive fluid weak
32	<i>be</i>	1.1	amylase negative	Amylase in body fluid (hemolymph) weak
Linkage Group IX				
33	<i>I</i>	0	Yellow inhibitor	Suppression of yellow blood and yellow cocoon
34	<i>I-a</i>	5.9	Dominant chocolate	Similar to chocolate mutation ( <i>ch</i> ); head black
35	<i>bd</i>	6.7	dilute black	Whole larval body dilute black
36	<i>og</i>	7.4	giallo ascoli	High translucency; female almost sterile
Linkage Group X				
37	<i>w<sub>1</sub>(w-1)</i>	0	white egg 1	No pigment in serosa; white eyes in moth
38	<i>fl</i>	0+	wingless	Fore and hind wings absent in pupa and moth
39	<i>w<sub>2</sub>(w-2)</i>	3.4	white egg 2	Egg gradually changes from white to light reddish color; white eyes in moth
40	<i>w<sub>3</sub>(w-3)</i>	6.9	white egg 3	Light purplish-brown egg; black eyes in moth
Linkage Group XI				
41	<i>K</i>	0	Knobbed	Dermal protuberances appear on dorsal sides of several segments of larva, pupa and moth
42	<i>Bu</i>	5.5	Burnt	Larva skin from 2nd to 5th segments shows burnlike scar
43	<i>bp</i>	17.1	black pupa	Black pupae (2 strains)
44	<i>mp</i>	24.0	micropterous	Small wings
Linkage Group XII				
45	<i>Ng</i>	0	No glue	Eggs easily separated from papers because of poor development of mucous glands in female
46	<i>C</i>	14.0	Golden egg	Cocoon golden yellow outside, nearly white inside
47	<i>rd</i>	52.1	clumpy	Irregular egg shape and highly variable
Linkage Group XIII				
48	<i>ch</i>	0	chocolate	Newly hatched larva reddish-brown
49	<i>cf</i>	11.3	crayfish	Fore and hind wings swollen and protrude laterally from body in pupa
Linkage Group XIV				
50	<i>Di</i>	0	Dirty	Irregular black lines and dots cover dorsal surface of larva
51	<i>U</i>	2.7	Ursa	Dark brown pigments cover dorsal and lateral sides of larva
52	<i>odk</i>	10.7	mottled	Low translucency
Linkage Group XV				
53	<i>Se</i>	0	White side egg	Egg surface irregular and with many furrows
54	<i>Gc</i>	7.8	Green c	Green cocoon

Contributors: (a) Novitski, E., (b) Kikkawa, H.

References: [1] Tanaka, Y. 1953. *Advan. Genet.* 5:239. [2] Tazima, Y. 1957. *Proc. Intern. Genet. Symp.*, Japan, 1956, p. 280.



# Contrails

## 6. LINKAGE GROUPS: PLANTS

### Part I. NEUROSPORA CRASSA

Genes for *Neurospora crassa* are listed in order of locus on the chromosome; they proceed from left arm to right arm, with the CENTROMERE the dividing marker. A line under the symbol indicates that the exact position has not been determined. Bracket ( ) signifies no recombination between loci. Capital letters in Columns A and B do not indicate dominant genes. A stable diploid, necessary for testing dominance, has not been achieved with *Neurospora* to date.

Gene Symbol	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	
Linkage Group I <sup>1</sup>				
1	<i>fr</i>	Frost	Delicate branching and nonconidial aerial growth	31
2	<i>nit-2</i>	Nitrate-2	Does not reduce nitrate	33
3	<i>leu-3</i>	Leucine-3	Requires leucine	3,31
4	<i>leu-4</i>	Leucine-4	Requires leucine	13
5	<i>un(b39)</i>	Unknown (b39)	Grows at 25°C, not at 34°C; distal to <i>A/a</i> ; not tested for allelism with <i>un(55701)</i>	21
6	<i>un(55701)</i>	Unknown (55701)	Unknown requirement; strain <i>55701t</i> grows at 25°C, but not on minimal medium at 35°C	3,16,18
7	<i>A/a</i>	Sex	Mating type	3,31
8	<i>dot</i>	Dot	Restricted colonial growth (data scanty)	33
9	<i>pat</i>	Patch	Circadian rhythm of dense and sparse mycelial growth; location proximal to <i>A/a</i>	39
10	<i>phen</i>	Phenylalanine	Requires phenylalanine or leucine, or other aromatic amino acids; location uncertain with respect to <i>ad-5</i>	2
11	<i>ad-5</i>	Adenine-5	Requires adenine	3,31,33
12	<i>amyc</i>	Amycelial	Growth budding; forms dot-like colonies; location proximal to <i>ad-5</i>	18,31
13	<i>arg-1</i>	Arginine-1	Requires arginine; does not utilize ornithine or citrulline	31
14	<i>arg-3</i>	Arginine-3	Requires arginine or citrulline; does not utilize ornithine	3,31
15	<i>ti</i>	Tiny	Very restricted colonial growth	30,31
16	<i>suc</i>	Succinic	Requires succinic acid or metabolically related compounds; closely linked to centromere	3
17	<i>cyt-1</i>	Cytochrome-1	Slow growth; altered cytochrome system; shows 5% recombination with <i>suc</i>	25
18	<i>sn</i>	Snowflake	Conidating colonial; location between <i>arg-3</i> and <i>hist-2</i>	23
CENTROMERE				
19	<i>hist-2</i>	Histidine-2	Requires histidine	44
20	<i>rg</i>	Ragged	Poor conidiation; colonial growth	31
21	<i>lys-4</i>	Lysine-4	Requires lysine	3,31,44
22	<i>hist-3</i>	Histidine-3	Requires histidine	3,44
23	<i>ad-3</i>	Adenine-3	Requires adenine; accumulates purple pigment on limiting adenine	3,5,6,31,36
24	<i>cut</i>	Cut	Tube culture appears as if mycelia were cut off part way up slant; location between <i>hist-2</i> and <i>arg-6</i>	21
25	<i>nic-2</i>	Nicotinic-2	Requires nicotinamide; accumulates red-brown pigment in the medium	3,31
26	<i>cr</i>	Crisp	Early uniform conidiation	3,31
27	<i>m-1</i>	Modifier of <i>vis(3717)</i>	Location proximal to <i>vis(3717)</i>	17
28	<i>vis(3717)</i>	Visible (3717)	Semi-colonial growth (located on right arm)	3,17
29	<i>m-2</i>	Modifier of <i>vis(3717)</i>	Location distal to <i>vis(3717)</i>	17
30	<i>st</i>	Sticky	Can only be scored by direct comparison with wild type; location between <i>hist-2</i> and <i>thi-1</i>	31
31	<i>un(44409)</i>	Unknown (44409)	Unknown requirement; strain <i>44409t</i> grows at 25°C on complete medium, but not at 34°C	3,31
32	<i>slo</i>	Slow	Slow growth; location proximal to <i>thi-1</i>	31
33	<i>thi-1</i>	Thiamine-1	Requires thiamine	3,31
34	<i>mac</i>	Methionine-adenine-cysteine	Requires methionine; grows best on all 3 substances or complete medium; location between <i>thi-1</i> and <i>al-2</i>	31
35	<i>me-6</i> <sup>2</sup>	Methionine-6	Requires methionine	3,27,31
36	<i>csh</i>	Cushion	Restricted colonial growth; location between <i>thi-1</i> and <i>nit-1</i>	36
37	<i>nit-1</i> <sup>3</sup>	Nitrate-1	Does not reduce nitrate	3,31,40
38	<i>un(STL6)</i>	Unknown (STL6)	Suboptimal response to methionine; best scored at 35°C; fluffy morphology with late conidiation	31
39	<i>arg-6</i>	Arginine-6	Requires ornithine, citrulline, or arginine	3,31

<sup>1</sup>/1/ Other markers known to be in Linkage Group I but presumably lost: *pa*--pale--conidia clumped and pale in color; *nd*--natural death--growth ceases progressively when fungus is homocaryotic; *dir*--dirty--conidia misshapen and few, yellow exudate; *gap*--gap--conidia produced in upper part of culture tube. [3] <sup>2</sup>/2/ Called *me(35809)* in reference 3. <sup>3</sup>/3/ Called *n-nit* in references 3 and 40.

continued

## 6. LINKAGE GROUPS: PLANTS

### Part I. NEUROSPORA CRASSA

Gene Symbol	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)
Linkage Group I			
40 <i>T</i>	Tyrosinase thermo-stability	<i>T<sup>S</sup></i> and <i>T<sup>L</sup></i> govern tyrosinase thermostability; location between <i>hist-2</i> and <i>al-2</i>	15
41 <i>su-1-me</i>	Suppressor-1-methionine	Suppresses <i>me-2</i> and <i>me-7</i> ; shows 1% recombination with <i>al-2</i>	3
42 <i>al-2</i>	Albino-2	Albino	3,31
43 <i>aur</i>	Aurescent	White at first; later forms pigmented terminal conidia	3,31
44 <i>hs</i>	Homoserine	Requires homoserine	3,31
45 <i>can</i> <sup>a</sup>	Canavanine	Resistance to canavanine	3,31-33
46 <i>lys-3</i>	Lysine-3	Requires lysine	3,31,35
47 <i>nic-1</i> <sup>b</sup>	Nicotinic-1	Requires nicotinamide	35
48 <i>os</i>	Osmotic	Inhibited by high osmotic pressure; can be scored by appearance; conidia rare	31
49 <i>so</i>	Soft	Dense pigmented growth in lower part of slant	31
Linkage Group II			
50 <i>cfl</i>	Cauliflower	Dense conidiation in bunches at top of slant	33
51 <i>thr-2</i>	Threonine-2	Requires threonine; leaky	8,27,33
52 <i>thr-3</i>	Threonine-3	Requires threonine; extremely close to <i>thr-2</i> ; leaky	33
53 <i>bal</i>	Balloon	Restricted growth; hemispherical colony	14,31
CENTROMERE			
54 <i>da</i>	Dapple	Flecks of conidia on agar surface; location uncertain with respect to <i>bal</i> and centromere	33
55 <i>arg-5</i>	Arginine-5	Requires ornithine, citrulline, or arginine	3,31
56 <i>arom-3</i>	Aromatic-3	Requires phenylalanine, tyrosine, tryptophan, and <i>p</i> -aminobenzoic acid; does not use shikimic acid	14
57 <i>cpl</i>	Carpet	Flat growth on agar slants; location uncertain with respect to <i>arom-3</i>	33
58 <i>pe</i>	Peach	Peach-colored conidia (see line 64)	3
59 <i>su-pe</i>	Suppressor of microconidial	Suppresses microconidial action of <i>pe<sup>m</sup></i> in <i>col-1</i> , <i>pe<sup>m</sup></i> genotype; location 14-22 map units from <i>pe</i>	12
60 <i>arom-1</i>	Aromatic-1	Requires phenylalanine, tyrosine, tryptophan, and <i>p</i> -aminobenzoic acid; grows on shikimic acid	3
61 <i>arom-4</i>	Aromatic-4	Requires phenylalanine, tyrosine, tryptophan, and <i>p</i> -aminobenzoic acid; does not use shikimic acid	14
62 <i>ac-1</i>	Acetate-1	Requires acetate plus ethanol	3
63 <i>tu</i>	Tuft	Conidia in large clusters at top of culture; location between <i>pe</i> and <i>fl</i>	3
64 <i>fl</i>	Fluffy	No macroconidia; few or no microconidia; <i>pe fl</i> genotype forms abundant microconidia	3
65 <i>tryp-3</i> <sup>c</sup>	Tryptophan-3	Requires tryptophan; does not use indole	3
66 <i>het-2</i>	Heterocaryon formation	Determines heterocaryon compatibility; alleles <i>d</i> , <i>D</i>	11
Linkage Group III			
CENTROMERE			
67 <i>thi-4</i>	Thiamine-4	Requires thiamine; location uncertain with respect to <i>sc</i>	3
68 <i>thi-10</i>	Modifier of <i>thi-4</i>	In presence of <i>thi-10</i> , <i>thi-4</i> requires intact thiamine; may be allelic with <i>thi-4</i>	7
69 <i>sc</i>	Scumbo	Flat, irregular, concentric growth	3
70 <i>mel-3</i>	Melon-3	Forms hemispherical colony; location uncertain with respect to centromere and markers <i>thi-4</i> to <i>leu-1</i>	26
71 <i>ser-1</i>	Serine-1	Requires serine; can use glycine; very leaky	3
72 <i>prol-1</i>	Proline-1	Requires proline; will not use ornithine, citrulline, or arginine	3
73 <i>com</i>	Compact	Small colonies	31,34
74 <i>me-8</i>	Methionine-8	Requires methionine	28
75 <i>ad-1</i>	Adenine-4	Requires adenine; will not use hypoxanthine	3,34
76 <i>leu-1</i>	Leucine-1	Requires leucine	3,31
77 <i>su-mel-3</i>	Suppressor-melon-3	A loose colonial results from interaction of <i>mel-3</i> and <i>su-mel-3</i> ; location distal to <i>leu-1</i>	26
78 <i>hist-7</i>	Histidine-7	Requires histidine; located between <i>ad-1</i> and <i>tryp-1</i>	44
79 <i>ad-2</i>	Adenine-2	Requires adenine or hypoxanthine	3,33,34

<sup>a</sup>/ Called *r-can* in reference 3. <sup>b</sup>/ Also called *Q*. <sup>c</sup>/ Also called *td*.

continued

# Contrails

## 6. LINKAGE GROUPS: PLANTS

### Part I. NEUROSPORA CRASSA

Gene Symbol	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	
Linkage Group III				
80	<i>tryp-1</i>	Tryptophan-1	Requires indole or tryptophan; accumulates yellow pigment	3
81	<i>thi-2</i>	Thiamine-2	Requires intact thiamine	3,34
82	<i>ro-2</i>	Ropy-2	Cable-like aggregations of hyphae; location between <i>tryp-1</i> and <i>vel</i>	31,34
83	<i>vel</i>	Velvet	Soft, conidiating colonial	31,34
84	<i>tyr-1</i>	Tyrosine-1	Requires tyrosine; very leaky	3,34
Linkage Group IV				
CENTROMERE				
85	<i>pyr-1</i>	Pyrimidine-1	Requires pyrimidine; uridine and cytidine 10-60 times as active as uracil	3,22,24
86	<i>pdx-1</i>	Pyridoxine-1	Requires pyridoxine	3,22,24
87	<i>pdx-2</i>	Pyridoxine-2	Requires pyridoxine	3
88	<i>rib-2</i>	Riboflavin-2	Requires riboflavin; location between <i>pdx-1</i> and <i>pyr-3</i>	3,22
89	<i>arg-2</i>	Arginine-2	Requires arginine; also uses citrulline	3,24
90	<i>col-4<sup>a</sup></i>	Colonial-4	Colonial, macroconidial	3,22,24
91	<i>me-1</i>	Methionine-1	Requires methionine; location between <i>pdx-1</i> and <i>pyr-3</i>	3,28
92	<i>pyr-3</i>	Pyrimidine-3	Requires pyrimidine ( <i>see</i> line 85)	3,24
93	<i>pt</i>	Phenylalanine-tyrosine	Requires phenylalanine plus tyrosine; location uncertain with respect to <i>pyr-3</i> ; accumulates brown pigment in medium on aging; fluoresces under ultraviolet light	22
94	<i>tryp-4</i>	Tryptophan-4	Requires indole or tryptophan; location between <i>pdx-1</i> and <i>pan-1</i>	3,22
95	<i>leu(37501)</i>	Leucine (37501)	Requires leucine; location between <i>pdx-1</i> and <i>pan-1</i>	22
96	<i>ad-6</i>	Adenine-6	Requires adenine; location between <i>pdx-1</i> and <i>pan-1</i>	3,22
97	<i>me-2</i>	Methionine-2	Requires methionine; location between <i>tryp-4</i> and <i>pan-1</i>	3,22,28
98	<i>fld</i>	Fluffyoid	Aconidial; location proximal to <i>pan-1</i>	33
99	<i>thi-5</i>	Thiamine-5	Requires thiamine; shows 1% recombination with <i>pan-1</i>	33
100	<i>pan-1</i>	Pantothenic-1	Requires pantothenic acid	3,22
101	<i>ro-1</i>	Ropy-1	Cable-like aggregations of hyphae	22,31
102	<i>nit-3<sup>a</sup></i>	Nitrate-3	Does not reduce nitrate; shows 13% recombination with <i>pan-1</i>	16,33
103	<i>chol-1</i>	Choline-1	Requires choline; location between <i>ad-6</i> and <i>me-5</i>	3,22
104	<i>col-1</i>	Colonial-1	Colonial growth; distal to <i>chol-1</i>	3
105	<i>cot</i>	Colonial-temperature sensitive	Colonial growth at 34°C; may be allelic with <i>col-1</i>	22,24
106	<i>ol</i>	Oleic acid	Requires higher fatty acid; lauric or larger, or Tween 80	33
107	<i>le-1</i>	Ascospore lethal	Colonial growth; location distal to <i>cot</i>	26
108	<i>hist-4</i>	Histidine-4	Requires histidine	22
109	<i>me-5</i>	Methionine-5	Requires methionine	3,22,24,34
110	<i>pyr-2</i>	Pyrimidine-2	Requires pyrimidine ( <i>see</i> line 85)	3,22
111	<i>dn</i>	Dingy	Gray lumps, presumably microconidia, on agar slants	24
112	<i>mat</i>	Mat	Colonial; grows better on sucrose than on glycerol; location distal to <i>pyr-2</i>	22,31
Linkage Group V				
CENTROMERE				
113	<i>lys-1</i>	Lysine-1	Requires lysine; location uncertain with respect to centromere	3
114	<i>sh</i>	Shallow	Spreading morphological; hyphae on surface of agar slant; location uncertain with respect to <i>lys-1</i>	33
115	<i>iv-1</i>	Isoleucine-valine-1	Requires isoleucine and valine	3
116	<i>val</i>	Valine	Requires valine	43
117	<i>iv-2</i>	Isoleucine-valine-2	Requires isoleucine and valine; location uncertain with respect to <i>iv-1</i> and <i>val</i>	3
118	<i>lys-2</i>	Lysine-2	Requires lysine; location between <i>lys-1</i> and <i>hist-1</i>	42,43
119	<i>sp</i>	Spray	Aerial mycelium fans outwards	31,43
120	<i>arg-4</i>	Arginine-4	Requires arginine; also uses ornithine or citrulline; may be allelic with <i>arg-7</i> ; location between <i>sp</i> and <i>inos</i>	3,33,37
121	<i>arg-7</i>	Arginine-7	Requires arginine; also uses ornithine or citrulline; may be allelic with <i>arg-4</i>	3,33,37
122	<i>am</i>	Amination-deficient	Requires α-amino nitrogen; leaky	9,31

/a/ Called *co* in reference 24. /a/ Called *nitr* in reference 16.

continued



## 6. LINKAGE GROUPS: PLANTS

### Part I. NEUROSPORA CRASSA

Gene Symbol (A)	Mutation (B)	Phenotypic Expression (C)	Reference (D)	
Linkage Group V				
123	<i>i</i>	Enhancer of <i>am</i>	Inhibits growth of <i>am</i> on medium containing inorganic nitrogen plus glutamic acid; shows 8% recombination with <i>am</i>	10
124	<i>wa</i>	Washed	Thin, spreading surface growth and conidiation; location between <i>lys-2</i> and <i>inos</i>	33
125	<i>hist-1</i>	Histidine-1	Requires histidine	3,43
126	<i>inos</i>	Inositol	Requires inositol	3,43
127	<i>arg-8</i>	Arginine-8	Requires arginine; data scanty	33
128	<i>pab-1</i>	para-Aminobenzoic acid-1	Requires <i>p</i> -aminobenzoic acid	3,43
129	<i>me-3</i>	Methionine-3	Requires methionine	3,43
130	<i>bis</i>	Biscuit	Conidiating colonial	31,43
131	<i>ser-2</i>	Serine-2	Requires serine; very leaky	33
132	<i>ad-7</i>	Adenine-7	Requires adenine	42,43
133	<i>pab-2</i>	para-Aminobenzoic acid-2	Requires <i>p</i> -aminobenzoic acid	43
134	<i>asp</i>	Asparagine	Requires asparagine	3,43
135	<i>pl</i>	Plug	Dense hyphae filling tube	31,43
Linkage Group VI <sup>a</sup>				
136	<i>ad-8</i>	Adenine-8	Requires adenine; far out on left arm	19
137	<i>cyt-2</i>	Cytochrome-2	Slow growth; altered cytochrome system	38
138	<i>asco</i>	Ascospores colorless	Low germination; requires lysine; <i>lys-5(DS6-85)</i> is an allele of <i>asco</i>	38,41
139	<i>um(66204)</i>	Unknown (66204)	Strain 66204 <i>i</i> does not grow on minimal medium at 35°C	3,38
140	<i>cys-2</i>	Cysteine-2	Requires cysteine or methionine (alleles designated <i>cys-c</i> , <i>cys-f</i> in reference 39)	38,39
141	<i>cys-1</i>	Cysteine-1	Requires cysteine or methionine	3,38
142	<i>ylo</i>	Yellow	Yellow conidia	3,38
143	<i>ad-1</i>	Adenine-1	Requires adenine	3,4,38
CENTROMERE				
144	<i>pan-2</i>	Pantothenic-2	Requires pantothenic acid	4
145	<i>rib-1</i>	Riboflavin-1	Requires riboflavin at 35°C; location uncertain with respect to <i>pan-2</i> and <i>del</i>	3,38
146	<i>del</i>	Delicate	Growth less than that of wild type	20,31
147	<i>tryp-2</i>	Tryptophan-2	Requires anthranilic acid, indole, or tryptophan; leaky	3,38
Linkage Group VII				
148	<i>nic-3</i>	Nicotinic-3	Requires nicotinamide	31,33
CENTROMERE				
149	<i>sfo</i>	Sulfonamide	Requires sulfonamide; location uncertain with respect to centromere	3
150	<i>thi-3</i>	Thiamine-3	Requires thiamine; leaky; best scored on agar slants after several days	3,31
151	<i>bn</i>	Button	Colonial; nonconidiating	31
152	<i>me-9</i>	Methionine-9	Requires methionine; leaky	29
153	<i>me-7</i>	Methionine-7	Requires methionine; proximal to <i>arg-11</i>	33
154	<i>col-2</i>	Colonial-2	Colonial; nonconidiating	1,3,33
155	<i>col-3</i>	Colonial-3	Colonial; nonconidiating	1,3,33
156	<i>thr-1</i>	Threonine-1	Requires threonine; proximal to <i>arg-11</i>	33
157	<i>wc</i>	White collar	No carotenoids except at low temperature; proximal to <i>arg-11</i>	33
158	<i>for</i>	Formate	Requires formate or adenine plus methionine; proximal to <i>arg-11</i>	33
159	<i>arg-11</i>	Arginine-11	Requires arginine, adenine, and uridine	31
160	<i>arg-10</i>	Arginine-10	Requires arginine; does not use ornithine or citrulline	31
161	<i>nt</i>	Nicotinic-tryptophan	Requires nicotinamide or tryptophan	3,31
162	<i>sk</i>	Skin	Flat growth; nonconidiating	31

<sup>a</sup> Also known to be in Linkage Group VI but presumably lost: *phen(38602)*--phenylalanine (38602)--requires phenylalanine. [2,16]

Contributors: Barratt, R. W., and Strickland, W. N.

continued



## 6. LINKAGE GROUPS: PLANTS

### Part I. NEUROSPORA CRASSA

*References:* [1] Barratt, R. W., and L. Garnjobst. 1949. *Genetics* 34:351. [2] Barratt, R. W., and W. Ogata. 1954. *Am. J. Botany* 41:763. [3] Barratt, R. W., et al. 1954. *Advan. Genet.* 6:1. [4] Case, M. E., and N. H. Giles. 1958. *Proc. Natl. Acad. Sci. U.S.* 44:378. [5] de Serres, F. J. 1956. *Genetics* 41:668. [6] de Serres, F. J. Unpublished. Oak Ridge Natl. Laboratory, Biology Division, Oak Ridge, 1961. [7] Eberhart, B. M., and E. L. Tatum. 1959. *J. Gen. Microbiol.* 20:43. [8] Emerson, S. 1950. Cold Spring Harbor Symp. Quant. Biol. 14:40. [9] Fincham, J. R. S. 1954. *J. Gen. Microbiol.* 11:236. [10] Fincham, J. R. S., and J. A. Pateman. 1957. *J. Genet.* 55:456. [11] Garnjobst, L. 1953. *Am. J. Botany* 40:607. [12] Grigg, G. W. 1958. *J. Gen. Microbiol.* 19:15. [13] Gross, S. R. Unpublished. Duke Univ., Dept. Microbiology, Durham, 1961. [14] Gross, S. R., and A. Fein. 1960. *Genetics* 45:885. [15] Horowitz, N. H., and M. Fling. 1956. *Proc. Natl. Acad. Sci. U.S.* 42:498. [16] Houlahan, M. B., G. W. Beadle, and H. G. Calhoun. 1949. *Genetics* 34:493. [17] Howe, H. B. 1956. *Ibid.* 41:610. [18] Howe, H. B. Unpublished. Univ. Georgia, Dept. Bacteriology, Athens, 1961. [19] Ishikawa, T. 1960. *Genetics* 45:993. [20] Ishikawa, T. Unpublished. Yale Univ., New Haven, 1961. [21] Kuwana, H. 1960. *Japan. J. Genet.* 35:49. [22] Maling, B. D. 1959. *Genetics* 44:1215. [23] Mitchell, M. B. 1958. *Ibid.* 43:799. [24] Mitchell, M. B., and H. K. Mitchell. 1954. *Proc. Natl. Acad. Sci. U.S.* 40:436. [25] Mitchell, M. B., H. K. Mitchell, and A. Tissieres. 1953. *Ibid.* 39:606. [26] Murray, J. C. 1959-60. *Dissertation Abstr.* 20:3480. [27] Murray, N. E. 1960. *Heredity* 15:199. [28] Murray, N. E. 1960. *Ibid.* 15:207. [29] Murray, N. E. Unpublished. Stanford Univ., Dept. Biological Sciences, Stanford, 1961. [30] Newmeyer, D. Unpublished. Stanford Univ., Dept. Biological Sciences, Stanford, 1961. [31] Perkins, D. D. 1959. *Genetics* 44:1185. [32] Perkins, D. D. 1960. *Microbiol. Genet. Bull.* 17:17. [33] Perkins, D. D. Unpublished. Stanford Univ., Dept. Biological Sciences, Stanford, 1961. [34] Perkins, D. D., and C. Ishitani. 1959. *Genetics* 44:1209. [35] St. Lawrence, P. 1956. *Proc. Natl. Acad. Sci. U.S.* 42:189. [36] St. Lawrence, P. Unpublished. Univ. California, Dept. Genetics, Berkeley, 1961. [37] Srb, A. M. 1946. Ph. D. Thesis. Stanford Univ., Palo Alto. [38] Stadler, D. R. 1956. *Genetics* 41:528. [39] Stadler, D. R. 1959. *Ibid.* 44:647. [40] Stadler, D. R. 1959. *Nature* 184:170. [41] Stadler, D. R. 1959. *Proc. Natl. Acad. Sci. U.S.* 45:1625. [42] Strickland, W. N. Unpublished. Stanford Univ., Dept. Biological Sciences, Stanford, 1961. [43] Strickland, W. N., D. D. Perkins, and C. C. Veatch. 1959. *Genetics* 44:1221. [44] Webber, B. B., and M. E. Case. 1960. *Ibid.* 45:1605.

### Part II. CHLAMYDOMONAS REINHARDI

Genes for *Chlamydomonas reinhardtii* are listed in order of locus on the chromosome; they proceed from left arm to right arm, with the CENTROMERE the dividing marker. In those linkage groups with genes mapped in only one arm of the chromosome, the CENTROMERE is listed first. A line under the gene symbol indicates that the exact position has not been determined. Bracket ( ) signifies no recombination between loci. Lower case letters in column A do not necessarily indicate recessive genes.

Gene Symbol	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)
Linkage Group I			
1 <u>ac-209</u>	Acetate-209	Requires acetate	5
CENTROMERE			
2 <u>ac-76</u>	Acetate-76	Requires acetate	5
3 <u>arg-1</u>	Arginine-1	Requires arginine, citrulline, or ornithine	1-4
4 <u>ac-14</u>	Acetate-14	Requires acetate	3
5 <u>arg-2</u>	Arginine-2	Requires arginine; does not utilize citrulline or ornithine	2-4
6 <u>pab-2</u>	<i>p</i> -Aminobenzoic-2	Requires <i>p</i> -aminobenzoic acid	2-4
7 <u>ac-115</u>	Acetate-115	Requires acetate; does not fix CO <sub>2</sub> ; shows 0.5% recombination with <u>pab-2</u>	7
8 <u>pf-4</u>	Paralyzed-4	Cells with paralyzed flagella	5
9 <u>thi-3</u>	Thiamine-3	Requires thiamine or thiazole	2-4

*continued*

## 6. LINKAGE GROUPS: PLANTS

### Part II. CHLAMYDOMONAS REINHARDI

Gene Symbol	Mutation	Phenotypic Expression	Reference	
(A)	(B)	(C)	(D)	
Linkage Group II				
10	<i>thi-9</i>	Thiamine-9	Requires thiamine	5
CENTROMERE				
11	<i>ac-12</i>	Acetate-12	Requires acetate	3
12	<i>pf-12</i>	Paralyzed-12	Cells with paralyzed flagella	3
13	<i>pf-18</i>	Paralyzed-18	Cells with paralyzed flagella	3
14	<i>nic-2</i>	Nicotinic-2	Requires nicotinamide	3,4
Linkage Group III				
15	<i>pf-15</i>	Paralyzed-15	Cells with paralyzed flagella	3
16	<i>ac-28</i>	Acetate-28	Requires acetate; shows 0.6% recombination with <i>pf-15</i>	3
17	<i>pf-5</i>	Paralyzed-5	Cells with paralyzed flagella; shows 0.5% recombination with <i>pab-1</i>	5
18	<i>pab-1</i>	<i>p</i> -Aminobenzoic-1	Requires <i>p</i> -aminobenzoic acid	3,4
19	<i>ac-26</i>	Acetate-26	Requires acetate	5
CENTROMERE				
20	<i>ac-17</i>	Acetate-17	Requires acetate; closely linked to centromere	3
21	<i>ac-141</i>	Acetate-141	Requires acetate; does not fix CO <sub>2</sub>	7
22	<i>thi-2</i>	Thiamine-2	Requires thiamine or thiazole plus pyrimidine	3,4
Linkage Group IV				
23	<i>nic-11</i>	Nicotinic-11	Requires nicotinamide	3
CENTROMERE				
24	<i>thi-4</i>	Thiamine-4	Requires thiamine or thiazole	3,4
25	<i>pf-20</i>	Paralyzed-20	Cells with paralyzed flagella	3
26	<i>ac-55</i>	Acetate-55	Requires acetate; colonies yellow	5
Linkage Group V				
27	<i>ac-31</i>	Acetate-31	Colonies yellow	3
28	<i>thi-8</i>	Thiamine-8	Requires thiamine or pyrimidine	3
29	<i>ac-18</i>	Acetate-18	Requires acetate	5
CENTROMERE				
30	<i>pf-1</i>	Paralyzed-1	Cells with paralyzed flagella	3
Linkage Group VI				
31	<i>mt</i>	Mating type	Mating type plus or minus	3
32	<i>nic-7</i>	Nicotinic-7	Requires nicotinamide	3,4
33	<i>thi-10</i>	Thiamine-10	Requires thiamine	5
34	<i>ac-29</i>	Acetate-29	Colonies yellow	5
CENTROMERE				
35	<i>pf-14</i>	Paralyzed-14	Cells with paralyzed flagella	3
Linkage Group VII				
36	<i>ac-6</i>	Acetate-6	Requires acetate	5
CENTROMERE				
37	<i>pf-17</i>	Paralyzed-17	Cells with paralyzed flagella	3
38	<i>ac-1</i>	Acetate-1	Requires acetate; colonies almost white	3,4
39	<i>ac-22</i>	Acetate-22	Requires acetate; grows slowly; colonies pale green	5
40	<i>ac-5</i>	Acetate-5	Requires acetate; grows slowly; colonies pale yellow	5
Linkage Group VIII				
41	<i>pf-3</i>	Paralyzed-3	Cells with paralyzed flagella	5
42	<i>thi-1</i>	Thiamine-1	Requires intact thiamine	3,4
CENTROMERE				
43	<i>ac-157</i>	Acetate-157	Requires acetate	3,4
Linkage Group IX				
44	<i>ac-51</i>	Acetate-51	Requires acetate	3
45	<i>pf-16</i>	Paralyzed-16	Cells with paralyzed flagella	3
46	<i>sr-1a</i>	Streptomycin-1a	Resistance to streptomycin; allelic with <i>sr-7</i> <sup>1</sup>	5,8
CENTROMERE				
47	<i>ac-15</i>	Acetate-15	Requires acetate	3
48	<i>pf-13</i>	Paralyzed-13	Cells with paralyzed flagella	3

1/ Consult reference 8.

continued

*Centromere*  
**6. LINKAGE GROUPS: PLANTS**  
**Part II. CHLAMYDOMONAS REINHARDI**

Gene Symbol	Mutation	Phenotypic Expression	Reference
(A)	(B)	(C)	(D)
Linkage Group X			
CENTROMERE <sup>2</sup>			
49 <i>ac-16</i>	Acetate-16	Requires acetate; does not fix CO <sub>2</sub>	3,7
50 <i>pf-19</i>	Paralyzed-19	Cells with paralyzed flagella	3
51 <i>pf-6</i>	Paralyzed-6	Cells with paralyzed flagella	5
52 <i>nic-13</i>	Nicotinic-13	Requires nicotinamide	5
Linkage Group XI			
CENTROMERE			
53 <i>pf-2</i>	Paralyzed-2	Cells with paralyzed flagella	3
54 <i>ac-7</i>	Acetate-7	Requires acetate; colonies yellow-green	5
55 <i>ac-21</i>	Acetate-21	Requires acetate; does not fix CO <sub>2</sub>	3,6

*is* Sequence may be: CENTROMERE, *nic-13*, *pf-6*, *pf-19*, *ac-16*.

*Contributors:* Ebersold, W. T., and Levine, E. E.

*References:* [1] Ebersold, W. T. 1956. Am. J. Botany 43:408. [2] Ebersold, W. T., and R. P. Levine. 1959. Z. Vererbungslehre 90:74. [3] Ebersold, W. T., et al. 1962. Genetics 47:531. [4] Eversole, R. A. 1956. Am. J. Botany 43:404. [5] Levine, E. E. Unpublished. Harvard Univ., Cambridge, 1963. [6] Levine, R. P. 1960. Proc. Natl. Acad. Sci. U.S. 46:972. [7] Levine, R. P., and D. Volkmann. 1961. Biochem. Biophys. Res. Commun. 6:264. [8] Sager, R. 1954. Proc. Natl. Acad. Sci. U.S. 40:356.

**Part III. CORN**

The genes in each linkage group for *Zea mays* are carried by the corresponding chromosome, e.g., linkage group I, chromosome 1; linkage group II, chromosome 2, etc. Capital letters (columns A and C) indicate dominant genes.

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Linkage Group I			
1 <i>sr</i>	0	striate	Leaves striated
2 <i>ga<sub>6</sub></i>	15	gametophyte factor	Gametophyte viability
3 <i>zb<sub>4</sub></i>	21	zebra striping	Leaves with alternating transverse bands of green and whitish sectors
4 <i>ms<sub>17</sub></i>	25	male sterile	Male sterile
5 <i>ts<sub>2</sub></i>	27	tassel seed	Terminal inflorescence with pistillate flowers
6 <i>P</i>	28	Pericarp	Pericarp color
7 <i>zl</i>	30	zygotic lethal	Lethal zygote
8 <i>as</i>	53	asynaptic	Chromosomes unpaired at meiosis
9 <i>hm</i>	66	<i>Helminthosporium</i> susceptibility (recessive)	Susceptible to <i>Helminthosporium</i> infection
10 <i>br</i>	80	brachytic	Stalk has short internodes
11 <i>Vg</i>	84	Vestigial glumes	Glumes underdeveloped
12 <i>f<sub>1</sub></i>	85	fine striped	Fine striped, green and white leaves
13 <i>an<sub>1</sub></i>	107	anther ear	Stamens develop in pistillate inflorescence
14 <i>Kn</i>	128	Knotted leaves	Wartlike growths on leaves and stalk
15 <i>gs<sub>1</sub></i>	134	green striped	Leaves with light green stripes between vascular bundles
16 <i>TS<sub>6</sub></i>	157	Tassel seed	Terminal inflorescence with pistillate flowers
17 <i>bm<sub>2</sub></i>	161	brown midrib	Brown pigment in leaf midrib
Linkage Group II			
18 <i>ws<sub>3</sub></i>	0	white sheath	Leaf sheaths and stalk deficient in chlorophyll
19 <i>al</i>	4	albescens	Seedlings become whitish
20 <i>lg<sub>1</sub></i>	11	liguleless	Absence of ligule on leaves

*continued*



## 6. LINKAGE GROUPS: PLANTS

### Part III. CORN

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Linkage Group II				
21	<i>gl<sub>2</sub></i>	30	glossy seedling	Seedlings smooth and shining
22	<i>B</i>	49	Anthocyanin booster	Increases anthocyanin pigments
23	<i>sk</i>	56	silkless	Ears without silks
24	<i>fl<sub>1</sub></i>	68	floury endosperm	Endosperm powdery
25	<i>ts<sub>1</sub></i>	74	tassel seed	Terminal inflorescence with pistillate flowers
26	<i>v<sub>4</sub></i>	83	virescent	Young seedlings deficient in chlorophyll
27	<i>Ch</i>	138	Chocolate	Chocolate pericarp
Linkage Group III				
28	<i>cr<sub>1</sub></i>	0	crinkly leaves	Leaves crinkled
29	<i>d<sub>1</sub></i>	18	dwarf	Abnormally undersized
30	<i>rt</i>	32	rootless	Lacking roots
31	<i>lg<sub>3</sub></i>	35	Liguleless	Absence of ligule on leaves
32	<i>kg</i>	40	Ragged leaves	Leaves appear split and torn due to development of necrotic areas
33	<i>ts<sub>4</sub></i>	47	tassel seed	Terminal inflorescence with pistillate flowers
34	<i>ba<sub>1</sub></i>	64	barren stalk	No ear produced
35	<i>na<sub>1</sub></i>	75	nana (dwarf)	Abnormally undersized
36	<i>a<sub>1</sub></i>	103	anthocyanin	Anthocyanin pigments present
37	<i>sh<sub>2</sub></i>	103.0+	shrunk endosperm	Endosperm shrunken
38	<i>et</i>	115	etched endosperm	Endosperm etched
39	<i>ga<sub>7</sub></i>	121	gametophyte factor	Gametophyte viability
Linkage Group IV				
40	<i>de<sub>1</sub></i>	0	defective endosperm	Endosperm defective
41	<i>Ca<sub>1</sub></i>	35	Gametophyte factor	Gametophyte viability
42	<i>Ts<sub>5</sub></i>	56	Tassel seed	Terminal inflorescence with pistillate flowers
43	<i>sp<sub>1</sub></i>	66	small pollen	Pollen of small size
44	<i>su<sub>1</sub></i>	71	sugary endosperm	Endosperm sugary
45	<i>de<sub>16</sub></i>	74	defective endosperm	Endosperm defective
46	<i>zb<sub>6</sub></i>	84	zebra striping	Leaves with alternating transverse bands of green and whitish sectors
47	<i>Tu</i>	100	Tunicate (pod corn)	Enlarged glumes in male and female inflorescences
48	<i>j<sub>2</sub></i>	105	japonica striping	Leaves green and white striped
49	<i>gl<sub>3</sub></i>	111	glossy seedling	Seedlings smooth and shining
Linkage Group V				
50	<i>gl<sub>17</sub></i>	0	glossy seedling	Seedlings smooth and shining
51	<i>a<sub>2</sub></i>	1	anthocyanin	Anthocyanin pigment present
52	<i>bm<sub>1</sub></i>	7	brown midrib	Brown pigment in leaf midrib
53	<i>bt<sub>1</sub></i>	8	brittle endosperm	Endosperm brittle
54	<i>v<sub>3</sub></i>	11	virescent	Young seedlings deficient in chlorophyll
55	<i>bv</i>	13	breviis (dwarf) plant	Undersized
56	<i>pr</i>	32	red aleurone color	Aleurone red
57	<i>ys</i>	41	yellow stripe	Leaves green and yellow striped
58	<i>v<sub>2</sub></i>	73	virescent	Young seedlings deficient in chlorophyll
Linkage Group VI				
59	<i>po</i>	0	polymitotic	Spores undergo extra meiotic-like divisions; male sterile
60	<i>Y</i>	13	Yellow endosperm	Endosperm yellow
61	<i>bg<sub>11</sub></i>	33	pale green	Light green seedlings and plants
62	<i>Pl</i>	44	Purple plant	Plant purple
63	<i>Bh</i>	45	Blotched aleurone	Aleurone blotched
64	<i>sm</i>	54	salmon silk	Salmon-colored silk
65	<i>py</i>	64	pigmy	Dwarf plant
Linkage Group VII				
66	<i>o<sub>2</sub></i>	0	opaque endosperm	Endosperm opaque
67	<i>in</i>	4	intensifier of aleurone color	Color of aleurone intensified
68	<i>v<sub>5</sub></i>	8	virescent	Young seedlings deficient in chlorophyll
69	<i>ra<sub>1</sub></i>	22	ramosa	Branching of ear and tassel
70	<i>gl<sub>1</sub></i>	26	glossy seedling	Seedlings smooth and shining
71	<i>Tp</i>	36	Teopod	Plant with many tillers and narrow leaves; ears and tassels have enlarged bracts

continued



## 6. LINKAGE GROUPS: PLANTS

### Part III. CORN

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Linkage Group VII			
72	<i>sl</i>	40 slashed leaves	Leaves slashed
73	<i>ij</i>	42 iojap	Leaves green and white striped
74	<i>Bn</i>	60 Brown endosperm	Endosperm brown
75	<i>bd</i>	96 branched silkless	Ears branched without silks
Linkage Group VIII			
76	<i>v<sub>16</sub></i>	0 virescent	Young seedlings deficient in chlorophyll
77	<i>ms<sub>8</sub></i>	14 male sterile	Male sterile
78	<i>j<sub>1</sub></i>	28 japonica striping	Leaves green and white striped
Linkage Group IX			
79	<i>Dt</i>	0 Dotted	Controller of <i>a<sub>2</sub></i> mutability
80	<i>yg<sub>2</sub></i>	7 yellow-green	Seedlings and plants yellow-green
81	<i>C</i>	26 Aleurone color	Determines color of aleurone
82	<i>sh<sub>1</sub></i>	29 shrunken endosperm	Endosperm shrunken
83	<i>bz</i>	31 bronze	Aleurone and plant bronze
84	<i>bp</i>	44 brown pericarp	Brown pericarp
85	<i>wx</i>	59 waxy endosperm	Waxy endosperm
86	<i>pg<sub>12</sub></i>	66 pale green	Seedlings and plants light green
87	<i>v<sub>1</sub></i>	71 virescent	Young seedlings deficient in chlorophyll
88	<i>bk<sub>2</sub></i>	74 brittle stalk	Stalk brittle
89	<i>Wc</i>	106 White cap of endosperm	Cap of endosperm white
Linkage Group X			
90	<i>Rp</i>	0 Resistance to <i>Puccinia</i>	Resistance to <i>Puccinia</i> infection
91	<i>Og</i>	16 Old gold striping	Leaves green and yellow striped
92	<i>li</i>	28 lineate	Leaves with fine longitudinal striations
93	<i>l<sub>8</sub></i>	38 luteus seedling	Yellow seedlings
94	<i>gl</i>	43 golden	Plant golden
95	<i>R</i>	57 Aleurone and plant color	Determines color of aleurone and plant

Contributor: Rhoades, M. M.

References: [1] Burnham, C. R. 1947. Maize Genet. Coop. News Letter 21:36. [2] Burnham, C. R. 1955. Ibid. 29:51. [3] Rhoades, M. M. 1950. J. Heredity 41:59. [4] Rhoades, M. M. 1955. In G. F. Sprague, ed. Corn and corn improvement. Academic Press, New York. pp. 123-219.

### Part IV. TOMATO

Linkage groups for *Lycopersicon esculentum* do not correspond to similarly numbered chromosomes. Some linkage groups have not been assigned to a particular chromosome, while some chromosomes have not been given a linkage group number. Capital letters (columns A and C) indicate dominant genes.

Gene Symbol	Locus	Mutation	Phenotypic Expression
(A)	(B)	(C)	(D)
Linkage Group I (Chromosome 2)			
1	<i>dv</i>	0 dwarf virescent	Stunted plants
2	<i>m</i>	3 mottled leaves	Leaves and cotyledons mottled
3	<i>d</i>	5 dwarf plant	Plant dwarfed; leaves dark and rugose
4	<i>p</i>	9 peach	Peach or pubescent fruit
5	<i>op</i>	13 opaca	Yellow-green patches on leaves
6	<i>dil</i>	17 dilute	Leaves light green
7	<i>ps</i>	20 positional-sterile	Positional-sterile flowers; prevents normal opening of corolla
8	<i>ro</i>	22 rosette	Rosette; very short internodes, no flowers

continued

## 6. LINKAGE GROUPS: PLANTS

### Part IV. TOMATO

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Linkage Group I (Chromosome 2)				
9	<i>O</i>	29	Oval fruit	Spherical, oblate, and elongate fruit
10	<i>aw</i>	30	without anthocyanin	No anthocyanin; stem green, not purple
11	<i>suf</i>	33	sufflava	Uniform light-green leaves
12	<i>ms<sub>10</sub></i>	35	male sterile-10	Pale anthers; exerted pistils; sterile
13	<i>bk</i>	36	beaked fruit	Sharp-pointed protuberance on blossom-end of fruit
14	<i>Cu</i>	38	Curl	Veins and petiole greatly foreshortened, leaves curled
15	<i>Me</i>	39	Mouse ears	Leaves pinnately compound, segments clavate
16	<i>wv</i>	42	white virescent	Plant virescent
17	<i>Wo</i>	48	Woolly plant	Woolly leaves and stems
18	<i>s</i>	53	compound inflorescence	Inflorescence much-branched; greatly increased number of flowers
19	<i>ne</i>	60	necrotic leaves	Necrotic leaf spots; leaves slowly killed
20	<i>Lc</i>	67	Few fruit locules	Fruits with only 2 or 3 locules
Linkage Group II (Chromosome 3)				
21	<i>r</i>	0	yellow fruit flesh	Yellow flesh color
22	<i>wf</i>	15	white flower	White or tan corolla
Linkage Group III (Chromosome 1)				
23	<i>br</i>	0	brachytic	Brachytic plants with short internodes
24	<i>y</i>	30	colorless fruit skin	Clear colorless skin on fruit
25	<i>Cf<sub>1</sub></i>	65	<i>Cladosporium</i> resistance	Resistance to races 1 and 3 of <i>Cladosporium fulvum</i>
Linkage Group IV (Chromosome 6)				
26	<i>c</i>	0	potato leaf	Potato leaf; reduced number of leaf segments
27	<i>sp</i>	2	self-pruning (determinate habit)	Self-pruning or determinate stems
28	<i>md</i>	25	mottled-2	Small chlorotic spots on leaves
29	<i>Mi</i>	59	Nematode resistance	Resistance to <i>Meloidogyne incognita</i>
30	<i>yv</i>	60	yellow virescent	Leaves yellowish
31	<i>Cf<sub>2</sub></i>	61	<i>Cladosporium</i> resistance	Resistance to races 1-4 of <i>Cladosporium</i>
Linkage Group V				
32	<i>bi</i>	0	bifurcata inflorescence	Branched inflorescence
33	<i>f</i>	2	fasciated fruit	Fasciated or many-loculed fruits
34	<i>a</i>	29	anthocyanin absent	Anthocyaninless; stems and leaves green, never purple
35	<i>hl</i>	49	hairless	Hairless plants; no hairs on hypocotyl
36	<i>gh</i>	54	ghost plants	Chlorophyll-deficient plants
37	<i>j<sub>1</sub></i>	69	jointless pedicel	Jointless pedicels
38	<i>Cf<sub>3</sub></i>	86	<i>Cladosporium</i> resistance	Resistance to races 1-4 of <i>Cladosporium</i>
Linkage Groups VI and VIII (Chromosome 8)				
39	<i>al</i>	0	anthocyanin loser	Anthocyanin loser; purple stems become green in 10-21 days
40	<i>gf</i>	23	green flesh	Persistent chlorophyll in fruit
41	<i>dl</i>	40	dialytic stamens	Dialytic; stamens are not united in a tube
42	<i>bu</i>	49	bushy	Bushy stems; short internodes, long petioles
43	<i>ch</i>	65	chartreuse petals	Small, yellow-green corolla
44	<i>l<sub>1</sub></i>	76	lutescent foliage	Premature yellowing of leaves; yellowish unripe fruits
(Chromosome 9)				
45	<i>wd</i>	0	wilty dwarf	Wilty dwarf plants; grayish-green, droopy leaves
46	<i>ah</i>	1.5	Hoffman's anthocyaninless (recessive)	Green stems
Linkage Group VII (Chromosome 10)				
47	<i>pe</i>	0	sticky peel	Sticky fruit epidermis
48	<i>lg</i>	8	light green foliage	Light green foliage
49	<i>u</i>	43	uniform-ripening fruit	Uniform light-green color of unripe fruits; no dark shoulders
50	<i>H</i>	61	Hairs absent	Nonhairy or smooth stems; hypocotyl and growing point hairy
51	<i>nd</i>	72	netted	Leaves chlorotic with chlorophyll concentrated around veins
52	<i>l<sub>2</sub></i>	79	lutescent-2	Leaves turn yellow
53	<i>t</i>	92	tangerine fruit color	Flesh and stamens orange color
54	<i>Xa</i>	124	Xantha seedlings	Xanthophyllic or yellow leaves
55	<i>ag</i>	139	Andrus' green stem (recessive)	Green stem but purple cotyledons

continued

## 6. LINKAGE GROUPS: PLANTS

### Part IV. TOMATO

Gene Symbol	Locus	Mutation	Phenotypic Expression	
(A)	(B)	(C)	(D)	
Linkage Groups X and XII (Chromosome 7)				
56	<i>wt</i>	0	wilty foliage	Wilty leaflets; leaf margins curl adaxially
57	<i>tf</i>	15	irifoliate	Terminal leaflet tripartite
58	<i>n</i>	30	nipple-tipped fruit	Nipple tips on fruit
59	<i>mc</i>	46	macrocalyx	Sepals much enlarged
Linkage Group XI (Chromosome 4)				
60	<i>e</i>	0	entire leaves	Entire or broad leaflets, as in Vilmorin's potato leaf
61	<i>di</i>	20	divergens	Leaf and stem color gray-green
62	<i>w</i>	28	wiry foliage	Wiry, slender, straplike leaflets; dwarfed plants
Unnumbered Linkage Group A				
63	<i>rv</i>	0	reticulate virescent	Leaves virescent, veins prominent
64	<i>sf</i>	34	solanifolium	Potato-like leaves
Unnumbered Linkage Group B				
65	<i>La</i>	0	Lanceolate	Entire, small leaves
66	<i>na</i>	28	nana	Tiny plants with short leaves

Contributors: (a) Butler, L., (b) Rick, Charles M.

References: [1] Butler, L. 1960. Can. J. Botany 38:365. [2] Butler, L. 1960. Tomato Genet. Coop. Rept. 10:5. [3] Rick, C. M., and L. Butler. 1956. Advan. Genet. 8:267.

## 7. GENETIC CODE

The nucleotide compositions of RNA codewords have been obtained by directing amino acids into protein in *Escherichia coli* extracts with randomly ordered polyribonucleotides synthesized with polynucleotide phosphorylase. Nucleotide sequence in codewords is not known, thus the order of bases is arbitrary. The tentative summary of RNA codewords shown in the table are considered to be *potential* codewords; that is, they code for amino acids in cell-free systems, but all codewords may not be applicable to cells in vivo.

Amino Acid		RNA Codewords <sup>1</sup>				Amino Acid		RNA Codewords <sup>1</sup>			
(A)		(B)				(A)		(B)			
1	Alanine	CCG	UCG <sup>2</sup>	ACG <sup>2</sup>	CGA <sup>2</sup>	11	Leucine	UUG	UUC	UCC	UUA
2	Arginine	CGC	AGA	UGC <sup>2</sup>		12	Lysine	AAA	AAU		
3	Asparagine	ACA	AUA	ACU <sup>2</sup>		13	Methionine	UGA			
4	Aspartic acid	GUA	GCA <sup>2</sup>	GAA <sup>2</sup>		14	Phenylalanine	UUU	CUU		
5	Cysteine	UUG				15	Proline	CCC	CCU	CCA	CCG <sup>2</sup>
6	Glutamic acid	GAA	GAU <sup>2</sup>	GAC <sup>2</sup>		16	Serine	UCU	UCC	UCG <sup>2</sup>	ACG
7	Glutamine	AAC	AGA	AGU <sup>2</sup>		17	Threonine	CAC	CAA		
8	Glycine	UGG	AGG	CGG		18	Tryptophan	GGU			
9	Histidine	ACC	ACU <sup>2</sup>			19	Tyrosine	AUU			
10	Isoleucine	UAU	UAA			20	Valine	UGU	UGA <sup>2</sup>		

/1/ Summary as of September, 1963. /2/ Probable codeword.

Contributor: Nirenberg, Marshall W.

References: [1] Nirenberg, M. W., et al. 1963. Cold Spring Harbor Symp. Quant. Biol. 28:549. [2] Speyer, J., et al. 1963. Ibid, 28:559.



## 8. CELL TYPES: SEED PLANTS

Because only approximate boundaries can be drawn between certain cell types in seed plants, the data have been restricted to the more salient and histologically apparent characteristics of the more common cell types. Spores, gametes, and many of the specialized cells have been omitted.

Cell Type		Specification	Description
(A)	(B)	(C)	
1	Apical meristem	Origin	Lineal descendants of cells of embryo, except in adventitious shoots and roots
2		Site	Apices of vegetative shoots, developing inflorescences, and flowers; in root, beneath inner edge of root cap
3		Morphology	Cells polyhedral; primary wall thin or irregularly thickened; primary pit fields may be present; nucleus large, ovoid; cytoplasm vacuolated; mitochondria, plastid primordia, and storage products may be present
4		Function	Point of origin of primary meristematic tissues (protoderm, ground meristem, procambium) from which primary body of shoot and root develops; in shoot apex, gives rise to tissue of leaf primordia
5	Vascular cambium	Origin	Procambium, and parenchyma in interfascicular areas, cortex, and phloem
6		Site	Lateral in stem and root, between secondary xylem and secondary phloem
7		Morphology	Two types of cambial cells: elongated fusiform initials (from which tracheary elements, sieve elements, fibers, and vertical parenchyma are derived), and ray initials (from which vascular rays originate); cytoplasm in both cell types highly vacuolated; primary walls have conspicuous pit fields
8	Function	Produces secondary phloem and secondary xylem cells; growth in diameter of woody stems and roots results from tissue formation by cambial cells	
9	Phellogen (cork cambium)	Origin	In stems, first phellogen cells arise from cortex (most commonly from outermost layer), or from epidermis or phloem parenchyma cells; in roots, from pericycle
10		Site	Lateral in stem and root, between phellem and subjacent phelloderm, cortical or phloem tissue; also beneath surface exposed by abscission of organs (leaf scars) or beneath wounds
11		Morphology	Cells rectangular and radially flattened in transectional view, polygonal or nearly isodiametric in longisection; walls thin, cytoplasm vacuolated; may contain tannins and chloroplasts
12		Function	Produces phellem cells outwardly and in many cases phelloderm cells inwardly; forms complementary tissue of lenticels
13	Epidermis	Origin	Protoderm
14		Site	Surface layer of foliar and floral organs, young stems and roots
15		Morphology	Cells polygonal, elongated, or with undulated contour in surface view; variable in radial dimensions; walls thin or thick, the outer wall often thicker; primary pit fields present; walls typically cutinized (may be lignified or silicified), the outer wall covered by cuticle except in roots; plastids, anthocyanin pigments, and ergastic substances may occur
16	Function	Mechanical protection; restriction of transpiration; storage of water and metabolic products; photosynthesis; water absorption in roots; by division and dedifferentiation may contribute to origin of adventitious shoots and roots	
17	Guard cell	Origin	Typically originate in pairs by division of specific "mother cells" of the protoderm; (in many plants paired guard cells are flanked or surrounded by distinctive subsidiary cells)
18		Site	Epidermis of foliage leaves and young stems; occurs in the epidermis of various types of floral organs; absent from epidermis of roots
19		Morphology	Cells usually crescentic or kidney-shaped in surface view; walls unevenly thickened, cutinized and overlaid by cuticle; often with ridgelike extensions above and below pore; conspicuous starch-forming chloroplasts present; protoplast physiologically active in mature cells but rarely divides in response to wound or other stimuli
20		Function	A pair of guard cells and the intercellular space or pore between them form a stoma. Reversible changes in turgor of guard cells result in the opening or closure of the pore, permitting diffusion of gases through the epidermis.
21	Phellem (cork)	Origin	Phellogen; in stems of some monocotyledons, tangentially dividing cortical parenchyma cells produce irregular bands of suberized cells termed "storied cork"
22		Site	Peripheral regions of stem, root, and some fruits; occurs in bud scales and petioles; often produced as a result of wounding
23		Morphology	Cells rectangular in transection and radially flattened; irregular or rectangular in longisection; wall typically suberized and devoid of pits; nonliving at maturity; may contain resin and tannins
24	Function	Mechanical protection; restriction of transpiration	
25	Parenchyma	Origin	Ground meristem, procambium, vascular cambium, and phellogen
26		Site	Dominant cell type in cortex, pith, mesophyll, fleshy fruits, and endosperm of seeds; occurs in phloem and xylem as component of vertical strands and vascular rays
27		Morphology	Cells approximately tetrakaidecahedral to elongated, or stellately branched; primary walls thin or thick, often with conspicuous pit fields; thick, lignified secondary walls with pits common in secondary xylem; plastids and a large number of ergastic substances present

*continued*



## 8. CELL TYPES: SEED PLANTS

Cell Type	Specification	Description
(A)	(B)	(C)
28	Parenchyma Function	Photosynthesis; food and water storage; secretion and excretion; commonly the protoplast retains marked capacity for growth, division, and differentiation, and therefore is prominently concerned in wound healing, formation of callus tissue, and the origin of adventitious shoots and roots
29	Collenchyma Origin	Ground meristem
30	Site	Sole component of cylinders or strands of tissue in the subepidermal portions of stems, petioles, and larger veins of leaves; may occur in root cortex
31	Morphology	Cells relatively short and prismatic, or elongated with tapering ends; primary walls unevenly thickened and composed of cellulose, pectin, and a high percentage of water; primary pit fields present; chloroplasts common; collenchyma and cortical parenchyma cells frequently intergrade in form and structure
32	Function	Mechanical support for growing stems and leaves; protoplast retains capacity for growth, division, and differentiation
33	Sclereid Origin	Protoderm, ground meristem, phellogen, vascular cambium, and procambium; frequently arises by sclerosis of a fully developed parenchyma cell
34	Site	Common in seed coats and fruits; diffusely arranged (as idioblasts or as a component of cell clusters) in cortex, functioning phloem, outer bark, pith, and mesophyll; in leaves of some dicotyledons, sclereids are restricted to vein endings (terminal sclereids)
35	Morphology	Cells may be polyhedral, columnar, fusiform, filiform, irregularly lobed or branched; in some cases they intergrade in form with fibers; secondary wall thick and lignified (sometimes with imbedded crystals); pits usually simple, often ramiform; protoplast may be retained at maturity
36	Function	Produces hard, incompressible texture of many tissues
37	Fiber Origin	Protoderm, ground meristem, procambium, and vascular cambium
38	Site	Cortex, primary and secondary vascular tissues of stem and root; epidermis of some leaves; component of hypodermal strands or layers and of the sclerenchymatous sheaths of vascular bundles in many kinds of leaves; cells may occur as idioblasts
39	Morphology	Typical prosenchymatous cell, frequently elongated; secondary wall usually thick, often highly lignified; pits abundant or scarce, simple or with greatly reduced borders; protoplast usually absent at maturity; a living protoplast and various ergastic materials occur in septate fibers; in secondary xylem of dicotyledons, fibers and tracheids frequently intergrade in form and structure
40	Function	Mechanical support
41	Tracheid Origin	Procambium and vascular cambium
42	Site	Primary and secondary xylem; in a modified form, the distinctive cell type in transpiration tissue of gymnosperm leaves; in leaves of some angiosperms, tracheid-like cells, termed "storage tracheids," occur as idioblasts, as cells associated with veinlets, or as components of cell groups or layers; commonly formed in masses in cultures of callus tissue
43	Morphology	Cells imperforate, and typically elongated with blunt, tapering, or inclined ends; patterns of lignified secondary wall thickenings diversified and often intergrading (annular, helical, scalariform, reticulate, pitted); devoid of protoplast at maturity
44	Function	Conduction of water and mineral solutes; mechanical support
45	Vessel member Origin	Procambium and vascular cambium
46	Site	Primary and secondary xylem of most dicotyledons; absent from xylem of all gymnosperms except members of the Gnetales; in some monocotyledons, restricted to primary xylem of root
47	Morphology	Cell form ranges from elongate to drum-shaped, with inclined or transversely oriented end walls; a series of superposed vessel members constitutes a vessel; perforation plates usually restricted to end walls and are either simple (one perforation) or multiperforated (scalariform, reticulate, or forminate); secondary walls lignified (with same range of patterns as in tracheids); devoid of protoplast at maturity
48	Function	Conduction of water and mineral solutes; possibly furnishes mechanical support
49	Sieve cell Origin	Procambium and vascular cambium
50	Site	Primary and secondary phloem of gymnosperms
51	Morphology	Cells elongated, with overlapping inclined or tapering ends; sieve areas numerous, uniform, and relatively undifferentiated; a sieve area is a portion of the primary wall, traversed by strands of cytoplasm, each strand enclosed in a cylinder of callose; protoplast enucleate at maturity; usually associated with certain ray and phloem-parenchyma cells termed "albuminous cells"
52	Function	Conduction of organic solutes
53	Sieve-tube member Origin	Procambium and vascular cambium
54	Site	Primary and secondary phloem of angiosperms

*continued*

## 8. CELL TYPES: SEED PLANTS

Cell Type	Specification	Description	
(A)	(B)	(C)	
55	Sieve-tube member	Morphology	Cells elongated, with inclined or transverse end walls; a series of superposed sieve-tube members constitutes a sieve tube; end walls with highly specialized sieve areas termed sieve plates; sieve plates are either simple (one sieve area) or compound (several sieve areas in scalariform or reticulate arrangement); lateral walls usually bear less-differentiated sieve areas; protoplast enucleate at maturity; each sieve-tube member associated with one or more nucleate cells (companion cells) which are ontogenetically developed as daughter cells of the sieve-tube member
56		Function	Conduction of organic solutes
57	Laticifer	Origin	Nonarticulated laticifers originate from single cells in the embryo and grow intrusively throughout plant; articulated laticifers arise from interconnected series of cells in which partial or complete removal of certain walls occurs
58		Site	Cortex, secondary phloem and xylem, pith, mesophyll
59		Morphology	Nonarticulated type is either an unbranched tube or a profusely ramified system of nonseptate tubes; articulated laticifers may become joined by lateral anastomoses to form a complex network; primary walls nonlignified and often thick; both types contain latex and are multinucleate
60		Function	Probably largely excretory because of storage of such apparently nonfunctional metabolic products as resins and rubber; the role of laticifers as food-conducting or food-storing structures is doubtful

*Contributors:* (a) Foster, Adriance S., (b) Burns, George W., (c) Armer, Sister Joseph Marie

*References:* [1] Bailey, I. W. 1954. Contributions to plant anatomy. Chronica Botanica, Waltham, Mass. [2] Carlquist, S. 1961. Comparative plant anatomy. Holt, Rinehart, and Winston; New York. [3] Esau, K. 1953. Plant anatomy. J. Wiley, New York. [4] Foster, A. S. 1949. Practical plant anatomy. Ed. 2. Van Nostrand, Princeton. [5] Foster, A. S. 1956. Protoplasma 46:184. [6] Huber, B. 1961. Grundzüge der Pflanzenanatomie. J. Springer, Berlin. [7] Metcalfe, C. R., and L. Chalk. 1950. Anatomy of the dicotyledons. Clarendon Press, Oxford.

## 9. TISSUE GROWTH CHARACTERISTICS: MAMMALS

Tissue	Growth Feature	Growth Characteristics	
(A)	(B)	(C)	
1	Adrenal	Division	Rat (adult), mitosis: zona capsula and zona fasciculata, 0.13% each; zona glomerulosa, 0.17%; zona reticularis, 0.06%; total gland, 0.12%. For young rat, percentages higher than for adult. [12, 73]
2		Mode of growth	Rat: Principally by cell division; capsule may contribute. [12]
3		Life span	Rat: In cortex, uncertain; phagocytosis occurs in zona reticularis. [12]
4		Replacement	Rat: In cortex, cell division and migration of cells from superficial to deeper layers [12, 73].
5		Regenerative capacity	Man, guinea pig: In cortex after damage, repair by cell division (limited in adult) [48]. Active regeneration follows postnatal degeneration. Man, rat: Mitosis scant in medulla; postnatal growth due to cell enlargement. [73, 85]
6	Alimentary canal	Division	Cat, stomach: Dividing cells at base of foveolae [49]. Rat: Dividing cells in crypts of duodenum and ileum; make up 3% of all cells; cycle, 1.13 hours. [64]
7		Mode of growth	Cell division and differentiation in mucous membrane. Muscle, by combination of cell division and increase in cell size. [75]
8		Life span	Rabbit, oral mucosa: 5.1(4.5-5.7)/1,000 cells; diurnal variation, 3.8(3.6-4.0) and 7.2(6.6-7.8). Mitotic duration calculated as 64 minutes. Intermitotic period calculated as 208 hours. [55] Rat, small intestine: 60-70% of superficial epithelium shed per day. Cell lives 1.57 days in duodenum, 1.35 days in ileum. [64]
9		Replacement	Cat: Cells multiply in crypts in base of foveolae, move toward lumen and differentiate [49].
10		Regenerative capacity	Cat: Brunner's glands can undergo limited regeneration [43]. Stomach: Movement at 0.2-0.4 mm/hr; epithelium, after sudden loss, restored in a few hours. [49] Dog, stomach: After removal of mucous membrane, denuded area covered by growth of undifferentiated epithelium at rate of 2 mm/wk [41].

*continued*

*Continued*

## 9. TISSUE GROWTH CHARACTERISTICS: MAMMALS

Tissue	Growth Feature	Growth Characteristics
(A)	(B)	(C)
11 Blood erythrocytes	Division	Confined largely to erythroblasts in bone marrow. Man, dividing cells: 1.17-1.83%. [76, 81]
	Mode of growth	Rabbit, rat, chicken: In precursors, growth phase sharply separated from hemoglobin formation [86].
	Life span	Man: 120 days most generally accepted [24].
	Replacement	Man: 0.83% red cells replaced each day [24].
	15 Regenerative capacity	Man: Recovery in 50 days from 600-ml blood loss [45]. Rabbit: Recovery in 3 weeks from 30% blood loss [18]. Rat: Recovery in 7 days from 30% blood loss [18].
16 Blood granulocytes	Division	Man: Confined largely to myeloblasts (2.7%) and myelocytes (0.46%) in bone marrow [81].
	Mode of growth	Rabbit, rat, chicken: Growth stages sharply demarcated from stages of granule formation [86].
	Life span	Man: Neutrophils estimated at 70-80 hours; eosinophils, 10-14 days. [58] Cat: Neutrophils disappear from blood at rate of 881/cu mm/hr [62].
	19 Replacement	Cat: Replaced 1.5 times per day [62].
20 Blood lymphocytes	Division	In tissues of origin. Rat (3 months old), dividing cells: thymus, 0.22%; lymph nodes, 0.058%; lymph follicles in spleen, 0.058%. [5]
	Mode of growth	Rabbit, marrow: Derived from reticulum cells and may be converted to other forms (?) [70]. Cat: $35 \times 10^6$ kg/hr enter circulation [2]. Dog: $25 \times 10^6$ kg/hr enter circulation [92].
	Life span	Approximately 24 hours. Removed from blood by lungs (rat) [89], spleen (rabbit) [40], lymph glands (dog) [92], skin (rat) [6], intestine [51].
	23 Replacement	Replacement in blood stream 2.06 times per day [2, 92, 93].
	24 Regenerative capacity	Rabbit (young): Limited regeneration of lymph nodes after removal; mesenchyme cells form complete gland in 3-4 weeks. [46]
25 Blood platelets	Mode of growth	Evidence favors formation in marrow from cytoplasmic fragmentation of megakaryocytes [91].
	Life span	Cat: 2-4 days; utilization, 2,500 cu mm blood/hr. [63]
	27 Replacement	Dog: After loss, new formation begins in a few hours; normal level regained in 3-4 days. [87].
28 Brain and spinal cord	Division	Rat: Very rare but has been reported [4].
	Mode of growth	Growth of axones, myelination of fiber tracts. May not be completed until 18th year in man. [34]
	Life span	Coextensive with normal function [75].
	31 Replacement	Confined to neuroglia [75].
	32 Regenerative capacity	Cat: Preganglionic fibers in peripheral nerves regenerate in 36-61 days [56]; function restored in 44 days [47]. Very largely confined to neuroglia; some axone formation. Rabbit: Regeneration of motor axones occurs at 4 mm/da after latent period of $\pm 7$ days. [94]
33 Hair	Growth rate	Man: head, 2.7 mm/wk; arm, 1.5 mm/wk; face, 2.1-3.5 mm/wk. [38, 88]
	34 Life cycle	Depends on thickness. Man: head, 3 years; eyebrow, 120 days. [23] Rat: body, 35 days [23].
35 Replacement	By multiplication and differentiation of cells at bottom of hair follicles [38, 88].	
36 Heart	Mode of growth	Man (from birth to maturity): Diameter of muscle fibers increases 2.6 times (to $14 \mu$ ) [82]. Rabbit: Diameter of muscle fibers increases 2.6 times (to $19 \mu$ ) [82].
	Life span	Coextensive with normal function [75].
	38 Replacement	None [75].
	39 Regenerative capacity	Negligible; hypertrophy caused by increase in size of fibers. Man: normal diameter, $14 \mu$ ; hypertrophied, $25 \mu$ . [82] Rabbit: normal diameter, $19.2 \mu$ ; hypertrophied, $22.2 \mu$ . [82]
40 Hypophysis	Division	Observed in normal animals. Occurs in anterior lobe after injury. Division stimulated when endocrine disturbances develop. Activity noted after administration of estrogen, testosterone, etc. [25]
41 Kidney	Division	Rare after early postnatal life, except for regeneration in rodents [75].
	Mode of growth	Increase in size of cells and structures. Man (from birth to maturity): Diameter of glomerulus increases from $118 \mu$ to $240 \mu$ , diameter of proximal collecting tubules from $18-34 \mu$ to $40-64 \mu$ . [32, 59, 75] Rat: Early postnatal growth partially caused by peripheral, undifferentiated nephrogenic zone; number of nephrons doubles in first two weeks of life. [13, 75]
	Life span	Very largely coextensive with normal function [75].
	44 Replacement	By cell divisions in tubules [75].
45 Regenerative capacity	Rat: True regeneration observed; hypertrophy caused by enlargement of existing elements, but increased cell division demonstrated after unilateral nephrectomy. [25, 26]	

*continued*



*Continued*  
9. TISSUE GROWTH CHARACTERISTICS: MAMMALS

Tissue	Growth Feature	Growth Characteristics
(A)	(B)	(C)
46	Liver	Division
		Rat: Dividing cells rise from low values to 3.3% on 23rd day, then return to low [19, 66]. (Albino), mitosis: 0.005% [19].
47		Mode of growth
		Rat: In early life, considerable contribution from cell division; later from increase in cell size. [66]
48		Life span
		Rat: No figure available [66].
49		Replacement
		Rat: Dividing cells persist in small numbers in adult, presumably to replace cell loss [66].
50		Regenerative capacity
		Rat: Extirpation of two-thirds regenerated in 21 days, caused by cell division in parenchymal cells and, to a smaller extent, in duct cells; division also occurs in Kupffer's cells and connective tissue cells. [1]
51	Muscle, smooth	Division
		Fibers retain some power of division [16].
52		Mode of growth
		New formation possible from undifferentiated connective tissue cells [16].
53		Regenerative capacity
		Regeneration after injury limited; fibrous scar usual. [16]
54	Muscle, striped	Division
		Scant; confined to nuclei. Some amitotic division. [16]
55		Mode of growth
		Enlargement and possible splitting of fibers. In newborn, some continued formation from mesenchyme cells. Hypertrophy caused by increase of sarcoplasm in preexisting cells. [16]
56		Life span
		Coextensive with normal function [75].
57		Replacement
		Normally none [75].
58		Regenerative capacity
		Rabbit: Protoplasmic outgrowth from preexisting fibers in which nuclei may divide by amitosis. Outgrowth begins 3rd day after injury, progresses at 1-1.5 mm/da. New fibers 30% of normal diameter in 21 days; normal diameter in 4 months. No new formation from undifferentiated cells. [28, 31]
59	Nails	Mode of growth
		Man, thumbnail: Daily increase of 95 $\mu$ . (Fingernails grow 4 times as fast as toenails.) [30]
60	Ovary	Division
		Rat: Mitosis demonstrated in germinal epithelium, and during pregnancy in granulosa and theca interna cells [11, 37].
61		Mode of growth
		Rabbit: Follicles increase in size exponentially with time; completed in 11 days. [35, 36] Rat: Total number of oocytes in gland related to age; $\log(\text{no.}) = 4.561 - 0.476 \log(\text{age in days})$ . [50, 67]
62		Life span
		Fertility loss of shed ova: ferret, 30 hours; guinea pig, 26 hours; rabbit, 12 hours. [53]
63		Regenerative capacity
		Mouse: No regeneration of ova; marked regenerative activity of germinal epithelium after hormonal stimulation. [65]
64	Pancreas	Mode of growth
		Man: Decrease in relative amount of connective tissue after birth; adult proportions reached at 11th-16th year. [39]
65		Regenerative capacity
		Man, dog, guinea pig, rabbit: Limited amount after resection or duct ligation. Mitosis in acinar cells and duct epithelium, the latter giving rise to acinar and islet cells. [25]
66	Parathyroid	Division
		Mouse, dividing cells: 8 days, 0.07%; 18 days, 0.71%; 28 days, 0.01%. [44]
67		Mode of growth
		Multiplication of clear or stem cells which differentiate into "dark" cells [42, 44].
68		Regenerative capacity
		Man: Negligible. Hypertrophy and hyperplasia in chronic nephritis. [3]
69	Prostate gland	Division
		Rat, mitosis: Numerous up to 20th day, scant at 100 days [77].
70		Mode of growth
		Man: During last months of pregnancy, fetal prostate shows proliferation and alteration of tubule epithelium, squamous metaplasia, cystic dilation of tubules, hyperplasia and dilation of ejaculatory ducts. These persist for 1-4 weeks after birth, then undergo gradual regression. [7, 25] Man, rat: Growth affected by hormonal activity [7, 77, 78]. Rat: By cell proliferation up to 20 days, then by increase in cell size and diameter of acini; from 11 to 55 days, height of cell increases from 18 $\mu$ to 34 $\mu$ , and diameter of acini from 43 $\mu$ to 170 $\mu$ . [77]
71		Regenerative capacity
		Man: Regeneration very rare after surgical removal [25].
72	Salivary gland	Division
		Rat, mitosis per 1,000 acinar cells: parotid, 1.02 $\pm$ 0.23; sublingual, 0.70 $\pm$ 0.23; submandibular, 0.64 $\pm$ 0.17. Mitosis per 1,000 tubular cells: submandibular, 0.44 $\pm$ 0.12. [27]
73		Life span
		Rat, acinar cells: parotid, 41 days; sublingual, 60 days; submandibular, 65 days. Tubular cells: submandibular, 95 days. [27]
74		Replacement
		Rat: By division of acinar or tubular cells [27, 72].
75		Regenerative capacity
		Rat: Slight regeneration by proliferation of acinar cells and of intercalated ducts [27, 72].

*continued*



9. TISSUE GROWTH CHARACTERISTICS: MAMMALS

Tissue	Growth Feature	Growth Characteristics
(A)	(B)	(C)
76	Skin	Division Mouse (newborn): 2-4% nucleated cells in mitosis [20, 33]. (8-16 weeks old): Mitotic cycle, 30.2±12 minutes [60]. (Adult): 2-8 dividing cells/cm length of 7-μ ear section [20, 33]. Mouse, rat: Occurs in varying proportions in stratum spinosum. Number of divisions varies with time of day, carbohydrate metabolism, hormonal stimulation, etc. [54] Rat (250 grams): Mitosis in the planta averages 5.24% for 24 hours at 27°C [84].
77	Mode of growth	Division of cells in deeper layers, followed by differentiation [75].
78	Life span	Rabbit, rat: Variable; dependent on rate of shedding of keratinized cells. [55, 84] Rat, plantar epidermis cells: 19.1 days, of which 16.9 days are spent in basal layer and 2.2 days in stratum granulosum [84]. Mitotic index for corneal epithelium, 0.4% [22].
79	Replacement	Mouse: Adjustment between cell loss and cell formation [21].
80	Regenerative capacity	Man: High capacity for regeneration and repair [29]; re-epithelization of wounds by proliferation and migration of adjacent cells [15, 28, 71]; regeneration possible from hair follicles [15]. Man, rat: Growth of granulation tissue checked by epithelium overgrowth [14, 15].
81	Submaxillary gland	Regenerative capacity Rat: Regeneration by division of acinar cells in 1st week, followed by proliferation of ducts with acini formed from terminal portions [25].
82	Testis	Division Rat (albino): Cycle of spermatogonial division, 48 minutes; spermatogenic wave lasts 4 days; spermatogenic cycle takes 16 days. [80]
83	Life span	Sperm survival time: man, 48-72 hours; horse, 12 hours; mouse, 13.5 hours; rabbit, 96 hours; bat, possibly through winter. [8, 9, 52, 83]
84	Regenerative capacity	Possible after slight injury. Rabbit: One day in abdominal cavity causes loss of spermatogenic elements; restoration in 2 weeks. [10]
85	Thymus	Division Mitosis: rabbit (adult), 0.52%; rat (3 month old), 0.22%. [5]
86	Mode of growth	Develops from pharyngeal epithelium [42]. Man: Increases from birth to puberty, decreases in later life. Weight at birth, 12-15 g; at puberty, 30-40 g; at 60 years, 10-15 g. [16]
87	Thyroid	Division Guinea pig: 100-125 dividing cells recorded in whole gland [17].
88	Mode of growth	Develops from pharyngeal epithelium [42]. Dog: Proliferation of interfollicular cells [95]. Rabbit: New follicles by budding [90].
89	Regenerative capacity	Man, dog, guinea pig: After partial removal, hypertrophy of remainder preventable by iodine administration [68, 69, 79].
90	Uterus	Mode of growth Man, menstruation: In proliferative stage, mitosis in endometrium rises to 0.56%, and nuclei of muscle fibers increase in size; regression in secretory phase. [74] Epithelial cells spread from gland tubules over connective tissue of endometrium, covering denuded surface by 7th day [25]. Pregnancy: Enlargement of preexisting epithelial cells, glands, and muscle fibers; cell division of epithelial and muscle elements; new muscle fibers may form from undifferentiated cells. [16, 61]
91	Regenerative capacity	Mouse: Experimental incision healed in 48 hours without scar [57].

Contributors: (a) O'Connor, R. J., (b) Glucksmann, A., (c) Hewitt, Harold B.

References: [1] Abercrombie, M., and R. D. Harkness. 1951. Proc. Roy. Soc. (London), B, 138:544. [2] Adams, W. S., R. H. Saunders, and J. S. Lawrence. 1945. Am. J. Physiol. 144:297. [3] Albright, F. 1933. New Engl. J. Med. 209:476. [4] Allen, E. 1912. J. Comp. Neurol. 22:547. [5] Andreasen, E., and S. Christensen. 1949. Anat. Record 103:401. [6] Andrew, W., and N. V. Andrew. 1949. Ibid. 104:217. [7] Andrews, G. S. 1951. J. Anat. 85:44. [8] Arey, L. B. 1954. Developmental anatomy. Ed. 6. W. B. Saunders, Philadelphia. [9] Asdell, S. A. 1946. Patterns of mammalian reproduction. Comstock, Ithaca. [10] Asdell, S. A., and G. W. Salisbury. 1941. Anat. Record 80:145. [11] Bassett, D. L. 1949. Ibid. 103:597. [12] Baxter, J. S. 1946. J. Anat. 80:139. [13] Baxter, J. S., and J. M. Yoffey. 1948. Ibid. 82:189. [14] Bentley, F. H. 1936. Ibid. 70:498. [15] Bishop, G. H. 1945. Am. J. Anat. 76:153. [16] Bloom, W., and D. W. Fawcett, ed. 1962. Maximow's Textbook of histology. Ed. 8. W. B. Saunders, Philadelphia. [17] Blumenthal, H. T. 1950. Growth 14:231. [18] Boycott, A. E. 1933. Trans. Roy. Soc. Trop. Med. Hyg. 27:529. [19] Brues, A. M., and B. B. Marble. 1937. J. Exptl. Med. 65:15. [20] Bullough, W. S. 1952. Biol. Rev. Cambridge Phil. Soc. 27:133. [21] Bullough, W. S., and F. J. Ebling. 1952. J. Anat. 86:29. [22] Buschke, W., J. S. Friedenwald, and W. Fleischmann. 1943. Bull. Johns

*continued*

9. TISSUE GROWTH CHARACTERISTICS: MAMMALS

Hopkins Hosp. 73:143. [23] Butcher, E. O. 1935. Anat. Record 61:5. [24] Callender, S. T., E. O. Powell, and L. J. Witts. 1945. J. Pathol. Bacteriol. 57:129. [25] Cameron, G. R. 1952. Pathology of the cell. Oliver and Boyd, Edinburgh. [26] Carnot, P., and R. Ray. 1938. Compt. Rend. Soc. Biol. 128:641. [27] Cherry, C. P., and A. Glucksman. 1959. Brit. J. Radio. 32:596. [28] Clark, W. E. leG. 1946. J. Anat. 80:24. [29] Clark, W. E. leG. 1958. The tissues of the body. Ed. 4. Oxford Univ. Press, London. [30] Clark, W. E. leG., and L. H. D. Buxton. 1938. Brit. J. Dermatol. Syphilis 50:221. [31] Clark, W. E. leG., and H. S. Wajda. 1947. J. Anat. 81:56. [32] Corning, H. K. 1925. Lehrbuch der Entwicklungsgeschichte des Menschen. Bergmann, Munich. [33] Cowdry, E. V., and H. C. Thompson. 1944. Anat. Record 38:403. [34] Davies, D. V., and F. Davies, ed. 1962. Gray's Anatomy. Ed. 33. Longmans, Green; London. [35] Desaive, P. 1947. Arch. Biol. (Paris) 58:331. [36] Desaive, P. 1948. Ibid. 59:34. [37] Dornfeld, E. J., and J. H. Berrian. 1951. Anat. Record 109:129. [38] Eaton, P., and M. W. Eaton. 1937. Science 86:354. [39] Emery, J. L. 1951. J. Anat. 85:159. [40] Farr, R. S. 1951. Anat. Record 109:515. [41] Ferguson, A. N. 1928. Am. J. Anat. 42:403. [42] Finerty, J. C., and E. V. Cowdry. 1960. A textbook of histology. Ed. 5. Lea and Febiger, Philadelphia. [43] Florey, H. W., and H. E. Harding. 1935. J. Pathol. Bacteriol. 40:211. [44] Foster, C. L. 1946. J. Anat. 80:171. [45] Fowler, W. M., and A. P. Barer. 1942. J. Am. Med. Assoc. 118:421. [46] Furuta, W. J. 1947. Am. J. Anat. 80:437. [47] Gibson, W. C. 1940. J. Neurophysiol. 3:237. [48] Graham, G. S. 1916. J. Med. Res. 34:241. [49] Grant, R. 1945. Anat. Record 91:175. [50] Green, S. H., A. M. Mandl, and S. Zuckerman. 1951. J. Anat. 85:325. [51] Haden, R. L. 1946. Principles of hematology. Ed. 3. Lea and Febiger, Philadelphia. [52] Hamilton, W. J., J. D. Boyd, and H. W. Mossman. 1962. Human embryology. Ed. 3. W. Heffer, Cambridge, England. [53] Hammond, J. 1941. Biol. Rev. Cambridge Phil. Soc. 16:165. [54] Hanson, J. 1947. J. Anat. 81:174. [55] Henry, J. L., et al. 1952. Arch. Pathol. 54:281. [56] Hinsey, J. C., R. A. Phillips, and K. Hare. 1939. Am. J. Physiol. 126:534. [57] Hooker, C. W. 1941. Anat. Record 79:211. [58] Jeanneret, H., and R. Fischer. 1941. Schweiz. Med. Wochschr. 22:204. [59] Kittleson, J. A. 1917. Anat. Record 13:385. [60] Knowlton, N. P., and W. R. Widner. 1950. Cancer Res. 10:59. [61] Krichesky, E. 1942. Anat. Record 82:551. [62] Lawrence, J. S., D. M. Ervin, and R. M. Wetrich. 1945. Am. J. Physiol. 144:284. [63] Lawrence, J. S., and W. N. Valentine. 1947. Blood 2:40. [64] Leblond, C. P., and C. E. Stevens. 1948. Anat. Record 100:357. [65] Li, M. H., and W. U. Gardner. 1947. Cancer Res. 7:549. [66] McKellar, M. 1949. Am. J. Anat. 85:263. [67] Mandl, A. M., and S. Zuckerman. 1951. J. Endocrinol. 7:190. [68] Marine, D. 1926. Arch. Pathol. Lab. Med. 2:829. [69] Marine, D., and C. H. Lenhart. 1909. Arch. Internal Med. 4:253. [70] Medawar, J. 1940. Brit. J. Exptl. Pathol. 21:205. [71] Medawar, P. B. 1945. Brit. Med. Bull. 3:70. [72] Millstein, B. B. 1950. Brit. J. Exptl. Pathol. 31:644. [73] Mitchell, R. M. 1948. Anat. Record 101:161. [74] Novak, E., and J. D. Woodruff. 1962. Gynecologic and obstetric pathology. Ed. 5. W. B. Saunders, Philadelphia. [75] O'Connor, R. J. Unpublished. Westminster Medical School, London, 1952. [76] Picena, J. P. 1937. Rev. Med. Rosario 27:1167. [77] Price, D. 1936. Am. J. Anat. 60:79. [78] Price, D. 1947. Physiol. Zool. 20:213. [79] Rienhoff, W. F. 1931. Medicine 10:257. [80] Roosen-Runge, E. C. 1951. Arch. J. Anat. 88:163. [81] Segerdahl, E. 1935. Acta Med. Scand., Suppl. 64. [82] Shipley, R. A., L. J. Shipley, and J. T. Wearn. 1937. J. Exptl. Med. 65:29. [83] Snell, G. D. 1941. Biology of the laboratory mouse. Blakiston, Philadelphia. [84] Storey, W. F., and C. P. Leblond. 1951. Ann. N. Y. Acad. Sci. 53:537. [85] Swinyard, C. A. 1943. Anat. Record 87:141. [86] Thorell, B. 1947. Acta Med. Scand., Suppl. 200. [87] Tocantins, L. M. 1936. Arch. Pathol. 21:69. [88] Trotter, M. 1923. Arch. Dermatol. Syphilol. 7:93. [89] Weisberger, A. R., et al. 1951. Blood 6:916. [90] Williams, R. G. 1939. Anat. Record 73:307. [91] Wintrobe, M. M. 1961. Clinical hematology. Ed. 5. Lea and Febiger, Philadelphia. p. 278. [92] Yoffey, J. M. 1933. J. Anat. 67:250. [93] Yoffey, J. M. 1936. Ibid. 70:507. [94] Young, J. Z. 1942. Physiol. Rev. 22:318. [95] Zechel, G. 1931. Surg. Gynecol. Obstet. 52:228.

## 10. CELL DIVISION FREQUENCY: MICROORGANISMS

For information on additional species, consult reference 1, Part I.

### Part I. PROTOZOA

Class	Species	Culture Medium	Temp. °C	Cell Divisions per Day	Refer-ence
(A)	(B)	(C)	(D)	(E)	(F)
1 Ciliata	<i>Didinium nasutum</i>	Hopkins' medium + <i>Paramecium</i>	21	3.6	2
2	<i>Paramecium aurelia</i>	Lettuce + bacteria	20±	0.72	6
3		Lettuce + bacteria	28	2.02	6
4	<i>Stentor coeruleus</i>	Peters' medium + ciliates	19	0.6-0.9	4
5		Modified Peters' medium + ciliates	18-20	0.7-2.1	4
6		Hetherington's medium + <i>Blepharisma</i>	22	0.65	3
7 Mastigophora	<i>Chilomonas paramecium</i>	Sodium acetate + mineral salts	24	3.5	5
8	<i>Euglena gracilis</i>	Wheat infusion	25	3.5	7

Contributor: Richards, Oscar W.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Beers, C. D. 1929. Am. Naturalist 63:125. [3] Gerstein, J. 1937. Proc. Soc. Exptl. Biol. Med. 37:210. [4] Hetherington, A. 1932. Arch. Protistenk. 76:118. [5] Mast, S. O., and D. M. Pace. 1933. Protoplasma 20:326. [6] Phelps, A. 1934. Arch. Protistenk. 82:134. [7] Sweet, H. E. 1939. Physiol. Zool. 12:173.

### Part II. VIRUSES AND BACTERIA

Culture Medium (column B): MS = mineral salts; AM = allantoic membrane.

	Species	Culture Medium	Temp. °C	Generation Time min	Refer-ence
	(A)	(B)	(C)	(D)	(E)
Vira <sup>1</sup>					
1	Influenza A (PR-8)	AM, chick embryo	37	330-510	10
2	Influenza A (5 strains)	AM, chick embryo	37	300-360	14
3	Influenza B (3 strains)	AM, chick embryo	37	480-600	14
4	Swine influenza	AM, chick embryo	37	360	14
Bacteriophyta <sup>2</sup>					
5	<i>Aerobacter aerogenes</i>	Broth or milk	37	18	16
6		Glucose + peptone	37	17.2-17.4	16
7		Peptone	37	22-30	16
8		Synthetic	37	29-44	9
9	<i>Azotobacter chroococcum</i>	Glucose broth	.....	27-39	16
10		MS + sugar	25-30	240-348	16
11		Sugar + urea	28	74	4
12	<i>Bacillus subtilis</i>	Glucose broth	.....	26-32	15
13	<i>Clostridium botulinum</i>	Glucose broth	37	35	23
14	<i>Corynebacterium diphtheriae</i>	Serum + glucose broth	37	34	16
15	<i>Diplococcus pneumoniae I</i>	Broth	37	24.5	2
16		Serum	37	29	1
17		Serum + broth	37	20.5	6
18	<i>D. pneumoniae II</i>	Broth	37	33	6
19		Glucose broth	37	30	15
20		Serum + broth	37	23	2

<sup>1/1</sup> Generation time: the time required for infected cells to release new virus. <sup>1/2</sup> Generation time: the average interval between cell divisions.

continued



## 10. CELL DIVISION FREQUENCY: MICROORGANISMS

### Part II. VIRUSES AND BACTERIA

	Species	Culture Medium	Temp. °C	Generation Time min	Refer- ence
	(A)	(B)	(C)	(D)	(E)
	Bacteriophyta <sup>2</sup>				
21	<i>Erwinia carotovora</i>	Broth	37	57	16
22		Glucose broth	37	42	16
23	<i>Escherichia coli</i>	Broth	37	16.5-17.0	22
24		Lactose broth	37	16	21
25		Milk	37	12.5	11
26	<i>Lactobacillus acidophilus</i>	Milk	37	66-87	16
27	<i>Mycobacterium tuberculosis</i> <sup>3</sup>	Synthetic	37	792-932	25
28	<i>Proteus vulgaris</i>	Broth	37	21.5	16
29		Peptone + phosphate	37	40	7
30	<i>Pseudomonas pyocyanea</i>	Broth	37	34	16
31		Glucose broth	37	31	16
32		Lactose broth	37	34	16
33	<i>Rhizobium leguminosarum</i>	MS + yeast + mannitol	25	79-187	5
34	<i>Salmonella typhosa</i>	Bile + pus	37	24.5	19
35		Broth	37	23.5	17
36		Glucose broth	37	29	8
37		Glucose + peptone	37	33	18
38	<i>Shigella dysenteriae</i>	Milk	37	23	13
39		Peptone + phosphate	37	37	7
40	<i>Staphylococcus aureus</i>	Broth	37	27	12
41		Glucose broth	37	32	16
42	<i>Streptococcus lactis</i>	Glucose milk	37	26	16
43		Lactose broth	30	48	16
44		Milk	37	26	20
45		Peptone milk	37	37	16
46	<i>Vibrio comma</i>	Broth	37	21.2-38.0	3
47	<i>Xanthomonas campestris</i>	Broth	23-25	165	24
48		Glucose broth	25	74	16

/2/ Generation time: the average interval between cell divisions. /3/ Human strain H-37.

Contributors: (a) Duca, Charles J., (b) Duggan, T. L.

References: [1] Barber, M. A. 1919. J. Exptl. Med. 30:569. [2] Blake, G. F. 1917. Ibid. 26:563. [3] Buchner, H., K. Longard, and G. Riedlin. 1887. Zentr. Bakteriolog. Parasitenk. 2:1. [4] Burk, D., and H. Lineweaver. 1930. J. Bacteriol. 19:389. [5] Cameron, G. M., and J. M. Sherman. 1935. Ibid. 30:647. [6] Chesney, M. M. 1916. J. Exptl. Med. 24:387. [7] Cohen, B., and W. M. Clark. 1917. J. Bacteriol. 4:409. [8] Coulter, C. B., and M. L. Isaacs. 1929. J. Exptl. Med. 49:711. [9] Dean, A. C. R., and C. Hinshelwood. 1951. J. Chem. Soc., p. 1457. [10] Fazekas de St. Groth, S. 1952. J. Immunol. 69:155. [11] Frazier, W. C., and E. O. Whittier. 1931. J. Bacteriol. 21:239. [12] Graham-Smith, C. S. 1920. J. Hyg. 19:133. [13] Heinemann, D. G., and T. H. Glenn. 1908. J. Infect. Diseases 5:534. [14] Henle, W., and E. S. Rosenberg. 1949. J. Exptl. Med. 89:279. [15] Lord, F. T., and R. N. Nye. 1919. Ibid. 30:389. [16] Mason, M. M. 1935. J. Bacteriol. 29:103. [17] Muller, M. 1895. Z. Hyg. Infektionskrankh. 20:245. [18] Penfold, W. J., and D. Norris. 1912. J. Hyg. 12:527. [19] Pies, W. 1907. Arch. Hyg. Bakteriolog. 62:107. [20] Rogers, L. A., and G. L. Greenbank. 1930. J. Bacteriol. 19:181. [21] Saito, K. 1907. Arch. Hyg. Bakteriolog. 63:215. [22] Sherman, J. M., and J. E. Holm. 1922. J. Bacteriol. 7:465. [23] Wagner, E., K. F. Mayer, and C. C. Dozier. 1925. Ibid. 10:321. [24] Wolf, F. A., and A. C. Foster. 1921. N. Carolina Agr. Expt. Sta. Tech. Bull. 20. [25] Youmans, G. P., and A. S. Youmans. 1950. J. Bacteriol. 60:569.



## 11. ORGANIC COMPOUNDS AFFECTING CELL DIVISION

Substance	Organism	Tissue	Effect	Reference
(A)	(B)	(C)	(D)	(E)
Interphase				
1 Acridines	<i>Gallus domesticus</i>	Chicken fibroblast	Initiation of prophase inhibited	37
2 Auxin	Spermatophyta	Flowering-plant cambium	Shortened	58
3 Azaguanine	<i>Mus sp.</i>	Mouse tumor	Initiation of prophase inhibited	66
4 Cortisone	<i>Allium cepa</i>	Onion root	Differentiated cell nuclei induced to divide	19
5 Dyes	<i>Rana sp.</i>	Frog sperm	Initiation of prophase inhibited	5
6 Folic acid antagonists	<i>Mus sp.</i>	Mouse intestine	Pycnosis from preprophase damage	23,24
7 Glucose	<i>Mus sp.</i>	Mouse epidermis	Shortened	8
8 Hydroquinone	<i>Mus sp.</i>	Mouse intestine	Pycnosis from preprophase damage	23,24
9 Hypoxanthine	<i>Gallus domesticus</i>	Chicken osteoblast	Shortened	31
10 Indoleacetic acid	<i>Allium cepa</i>	Onion root	Differentiated cell nuclei induced to divide	19
11	<i>Nicotiana tabacum</i>	Tobacco pith	Chromosome doubling within nucleus	43
12 Naphthaleneacetic acid	<i>Phaseolus sp.</i>	Bean internodes	Differentiated cell nuclei induced to divide	19
13 Neotetrazolium	<i>Allium cepa</i>	Onion root	Pycnosis from preprophase damage	56
14 Nitrogen mustard	<i>Rattus sp.</i>	Rat corneal epithelium	Initiation of prophase inhibited	26
15 Phenylacetic acid	<i>Allium cepa</i>	Onion root	Initiation of prophase inhibited	15
16 Trypaflavine	<i>Allium cepa</i>	Onion root	Destruction of interphase nucleus	52
17 Urethan	<i>Mus sp.</i>	Mouse intestine	Pycnosis from preprophase damage	23,24
Prophase				
18 Acridines	<i>Allium cepa</i>	Onion root	Reversion to interphase	16
19 Aureomycin	<i>Allium cepa</i>	Onion root	Membrane dissolution delayed	64
20 Dichlorophenoxyacetic acid	<i>Allium cepa</i>	Onion root	Blocked	17
21 Glutathione	<i>Amoeba proteus</i>	Whole organism	Accelerated	10
22 Nitrophenols	<i>Arbacia punctulata</i>	Sea-urchin egg	Blocked	11
23 Protoanemonin	<i>Zea mays</i>	Corn root	Blocked	25
24 Purines	<i>Arbacia punctulata</i>	Sea-urchin egg	Reversion to interphase	14
25 Tropolones	<i>Tradescantia sp.</i>	Spiderwort stamen hair	Precocious chromosome split	63
26 Trypan blue	<i>Oryctolagus cuniculus</i>	Rabbit fibroblast	Spindle formation slowed	61
27 Urethan	<i>Oryctolagus cuniculus</i>	Rabbit fibroblast	Accelerated	7
Metaphase				
28 Alcohol	<i>Allium cepa</i>	Onion root	Nucleolus neoformation	59
29 Colchicine	<i>Allium cepa</i>	Onion root	Monopolar mitotic figure induced	27
30 DDT	<i>Allium cepa</i>	Onion root	Nucleolus neoformation	59
31 Diethylbromacetyl carbamide	<i>Allium cepa</i>	Onion root	Multipolar spindle induced	49
32 Diphenyl	<i>Triticum sp.</i>	Wheat root	Spindle rotation	28
33 Endothal	<i>Pisum sativum</i>	Pea root	Chromosome noncongregation	65
34 Ethylmercuric phosphate	<i>Zea mays</i>	Corn seedling	Multipolar spindle induced	55
35 Indoleacetic acid	<i>Triticum sp.</i>	Wheat root	Spindle rotation	9
36 Methyl naphthohydroquinone diacetate	<i>Allium cepa</i>	Onion root	Multipolar spindle induced	46
37 Methyl naphthoquinone	<i>Allium cepa</i>	Onion root	Abnormal chromosome orientation	46
38 Narcotics	<i>Allium cepa</i>	Onion root	Monopolar mitotic figure induced	49
39 Phenylurethan	<i>Arbacia punctulata</i>	Sea-urchin egg	Monopolar mitotic figure induced	50
40 Streptomycin	<i>Allium cepa</i>	Onion root	Reversion to interphase	64
41 Testosterone, estrone	<i>Oryctolagus cuniculus</i>	Rabbit fibroblast	Abnormal chromosome orientation	60
42 Thallium acetate	<i>Allium cepa</i>	Onion root	Chromosome noncongregation	2
Anaphase				
43 Caffeine	<i>Allium cepa</i>	Onion root	Incomplete chromosome separation	42
44 Ryanodine	<i>Echinarachnius parma</i>	Sand-dollar egg	Incomplete chromosome separation	13
45 Trypaflavine	<i>Oryctolagus cuniculus</i>	Rabbit fibroblast	Incomplete chromosome separation	6

continued

## II. ORGANIC COMPOUNDS AFFECTING CELL DIVISION

Substance	Organism	Tissue	Effect	Ref-er-ence	
(A)	(B)	(C)	(D)	(E)	
Telophase					
46	Aureomycin	<i>Gallus domesticus</i>	Chicken fibroblast	Cytoplasmic division suppressed	33
47	Caffeine	<i>Allium cepa</i>	Onion root	Cytoplasmic division suppressed	29
48	Carbamates	<i>Lytechinus variegatus</i>	Sea-urchin egg	Cytoplasmic division suppressed	12
49		<i>Tripneustes esculentus</i>	Sea-urchin egg	Cytoplasmic division suppressed	12
50	Chloroacetophenone	<i>Gallus domesticus</i>	Chicken osteoblast	Nuclear reconstruction retarded	32
51	Nicotine	<i>Nicotiana tabacum</i>	Tobacco anther	Cytoplasmic division suppressed	35
52		<i>Pisum</i> sp.	Pea seedling	Spindle remnant persists	44
53	Quinone	<i>Tubifex</i> sp.	Oligochaete-worm egg	Cytoplasmic division suppressed	38
54	Rotenone	<i>Arbacia punctulata</i>	Sea-urchin egg	Cytoplasmic division suppressed	53
55	Sulfanilamide	<i>Allium cepa</i>	Onion root	Cytoplasmic division suppressed	4
56	Sulfhydryl compounds	Saccharomycetaceae	Yeasts	Cytoplasmic division augmented	45
57	Theobromine	<i>Allium cepa</i>	Onion root	Cytoplasmic division suppressed	29
58	Thiourea	<i>Gallus domesticus</i>	Chicken fibroblast	Nuclear reconstruction retarded	54
Not Confined to One Phase					
59	Acenaphthene	<i>Allium cepa</i>	Onion root	Chromosome breaks; centromere misdivision	18
60	Acridines	<i>Allium cepa</i>	Onion root	Chromosome breaks; pseudochiasmata	16,18
61	Aminoacridine	<i>Allium cepa</i>	Onion root	Chromosome breaks	24
62	Aminobenzoate	<i>Allium cepa</i>	Onion root	Chromosome breaks	24
63	Ammonia	<i>Tradescantia</i> sp.	Spiderwort stamen hair	Chromosome dispersion and despiralization	62
64	Ammonium thiocyanate	<i>Impatiens</i> sp.	Snapweed pollen mother cell	Chromosome dispersion and despiralization	36
65	Antibiotics	<i>Allium cepa</i>	Onion root	Chromosome reduction induced	64
66	<i>n</i> -Butyl gallate	<i>Allium cepa</i>	Onion root	Chromosome adhesion	40
67	Coumarin	<i>Allium cepa</i>	Onion root	Chromosome breaks; pseudochiasmata	48
68	Cysteine	Protozoa	Protozoan	Chromosome reduction induced	57
69	Dyes	<i>Allium cepa</i>	Onion root	Chromosome adhesion	3
70	Epoxides	<i>Vicia faba</i>	Broad-bean root	Chromosome rearrangements	41
71	Ethoxycaffeine	<i>Allium cepa</i>	Onion root	Chromosome rearrangements	34
72	Mustards	<i>Allium cepa</i>	Onion root	Chromosome rearrangements	20
73		<i>Tradescantia</i> sp.	Spiderwort pollen mother cell	Centromere misdivision	20
74	Nucleic acid	<i>Allium cepa</i>	Onion root	Chromosome reduction induced	1
75	Phenols	<i>Allium cepa</i>	Onion root	Chromosome breaks	39
76	Sulfhydryl	<i>Clymenella torquata</i>	Bamboo-worm regenerating tissue	Chromosomes widened	30
77	Sulfoxide	<i>Clymenella torquata</i>	Bamboo-worm regenerating tissue	Chromosomes widened	30
78	Uracil	<i>Allium cepa</i>	Onion root	Chromosome breaks	22
79	Urea	<i>Drosophila</i> sp.	Fruit-fly salivary glands	Chromosome dispersion and despiralization	51
80	Urethan	<i>Paeonia tenuifolia</i>	Fernleaf-peony bud	Chromosome rearrangements	47
81		<i>Vicia faba</i>	Broad-bean root	Chromosome breaks	21

Contributors: (a) Cornman, Ivor, (b) D'Amato, Francesco

References: [1] Allen, N. S., G. B. Wilson, and S. Powell. 1950. *J. Heredity* 41:159. [2] Avanzi, S. 1956. *Caryologia* 9:131. [3] Battaglia, E. 1950. *Ibid.* 2:223. [4] Bauch, R. 1949. *Pharmazie* 4:1. [5] Briggs, R. 1952. *J. Gen. Physiol.* 35:761. [6] Bucher, O. 1939. *Z. Zellforsch. Mikroskop. Anat.* 29:283. [7] Bucher, O. 1949. *Helv. Physiol. Pharmacol. Acta* 7:37. [8] Bullough, W. S. 1952. *Biol. Rev. Cambridge Phil. Soc.* 27:133. [9] Burström, H. 1942. *Lantbruks-Hogskol. Ann.* 10:209. [10] Chalkley, H. W. 1951. *Ann. N. Y. Acad. Sci.* 51:1303. [11] Clowes, G. H. A. 1951. *Ibid.* 51:1409. [12] Cornman, I. 1950. *J. Natl. Cancer Inst.* 10:1123. [13] Cornman, I. 1951. *Exptl. Cell Res.* 2:256. [14] Cornman, I. Unpublished. Hazleton Laboratories,

*continued*

## 11. ORGANIC COMPOUNDS AFFECTING CELL DIVISION

- Falls Church, Va., 1953. [15] D'Amato, F. 1949. *Caryologia* 1:109. [16] D'Amato, F. 1950. *Ibid.* 2:229. [17] D'Amato, F. 1950. *Protoplasma* 39:423. [18] D'Amato, F. 1950. *Pubbl. Staz. Zool. Napoli* 22(Suppl.):158. [19] D'Amato, F. 1952. *Caryologia* 4:311. [20] Darlington, C. D., and P. C. Koller. 1947. *Heredity* 1:187. [21] Deufel, J. 1951. *Chromosoma* 4:239. [22] Deysson, M. 1952. *Compt. Rend.* 234:650. [23] Dustin, P., Jr. 1947. *Nature* 159:794. [24] Dustin, P., Jr. 1950. *Compt. Rend. Soc. Biol.* 144:1297. [25] Erickson, R. O., and G. U. Rosen. 1949. *Am. J. Botany* 35:317. [26] Friedenwald, J. S. 1951. *Ann. N. Y. Acad. Sci.* 51:1432. [27] Gaulden, M. E., and J. G. Carlson. 1951. *Exptl. Cell Res.* 2:416. [28] Gavaudan, P. 1942. *Compt. Rend. Soc. Biol.* 136:419. [29] Gosselin, A. 1940. *Compt. Rend.* 210:544. [30] Hammett, F. S. 1934. *Protoplasma* 22:173. [31] Hopkins, F. G., and I. Simon-Reuss. 1944. *Proc. Roy. Soc. (London), B*, 132:253. [32] Hughes, A. F. W. 1950. *Quart. J. Microscop. Sci.* 91:251. [33] Keilova-Rodova, H. 1950. *Experientia* 6:428. [34] Kihlman, B. 1950. *Exptl. Cell Res.* 1:135. [35] Kostoff, D. 1931. *Bull. Soc. Bot. Bulgar.* 4:87. [36] Kuwada, Y., N. Shinke, and G. Oura. 1938. *Z. Wiss. Mikroskopie* 55:8. [37] Lasnitzki, I., and J. H. Wilkinson. 1948. *Brit. J. Cancer* 2:369. [38] Lehmann, F.-E. 1949. *Exptl. Cell Res., Suppl.* 1:156. [39] Levan, A., and J. H. Tjio. 1948. *Hereditas* 34:453. [40] Lopane, F. 1950. *Caryologia* 2:143. [41] Loveless, A. 1951. *Nature* 167:338. [42] Mangenot, G., and S. Carpentier. 1944. *Compt. Rend. Soc. Biol.* 138:232. [43] Naylor, J., G. Sandor, and S. Skoog. 1954. *Physiol. Plantarum* 7:25. [44] Nemeč, B. 1929. *Protoplasma* 7:99. [45] Nickerson, W. J., and J. W. van Rij. 1949. *Biochim. Biophys. Acta* 3:461. [46] Nybom, N., and B. Knutsson. 1947. *Hereditas* 33:220. [47] Oehlkers, F., and H. Marquardt. 1950. *Z. Induktive Abstammungs- Vererbungslehre* 83:299. [48] Östergren, G. 1948. *Botan. Notiser* (4):376. [49] Östergren, G. 1950. *Hereditas* 36:371. [50] Painter, T. S. 1918. *J. Exptl. Zool.* 24:445. [51] Painter, T. S. 1944. *Ibid.* 96:53. [52] Resende, F. 1951. *Bol. Soc. Port. Cienc. Nat.* 18:182. [53] Rogers, E. F., and I. Cornman. 1951. *Biol. Bull.* 101:227. [54] Rosin, A., E. Tenenbaum, and F. Doljanski. 1951. *Anat. Record* 111:239. [55] Sass, J. E. 1938. *Am. J. Botany* 25:624. [56] Sonnenblick, B. P., W. Antopol, and L. Goldman. 1950. *Trans. N. Y. Acad. Sci., Ser. 2*, 12:161. [57] Straub, J. 1951. *Biol. Zentr.* 70:24. [58] Thimann, K. V. 1938. *Physiol. Rev.* 18:524. [59] Vaarama, A. 1947. *Hereditas* 33:191. [60] Von Möllendorff, W. 1939. *Z. Zellforsch. Mikroskop. Anat.* 29:706. [61] Von Möllendorff, W., and M. Ostrouch. 1939. *Ibid.* 29:323. [62] Wada, B. 1937. *Cytologia (Tokyo), Fujii Jubilaei Vol. (2)*:785. [63] Wada, B. 1952. *Cytologia (Tokyo)* 17:14. [64] Wilson, G. B., and C. C. Bowen. 1951. *J. Heredity* 42:251. [65] Wilson, S. M., A. Daniel, and G. B. Wilson. 1956. *Ibid.* 47:151. [66] Woodside, G. L., et al. 1951. *Anat. Record* 111:501.



# *Contrails*

## II. REPRODUCTION

### 12. PROPAGATION: MAMMALS

For additional information, consult references 2 and 6. **Type of Estrus** (column E): P = polyestrous; M = monoestrous. Values in parentheses are ranges, estimate "c" or "d" unless otherwise indicated (cf. Introduction).

Species	Common Name	Age at Puberty	Breeding Season	Estrus		Gestation Period da	Young per Litter	Reference
				Type	Cycle da			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 <i>Homo sapiens</i>	Man	♀ 13.5 (11-16) <sup>b</sup> yr	All year	P	28.4 (24-33) <sup>b</sup>	278 <sup>1</sup> (253-303) <sup>b</sup>	1 <sup>2</sup>	3,17,37, 47,50,51, 64,81
2 <i>Balaenoptera physalus</i>	Finback whale	3 yr	Nov-Mar <sup>3</sup> , June-Aug <sup>4</sup>	...	.....	360	1	56,67,84
3 <i>Bos taurus</i>	Cattle	6-10 mo	All year	P	(14-23)	284(210-335)	Usually 1	49,54,66
4 <i>Camelus bactrianus</i>	Bactrian camel	.....	All year	P	(10-20)	(389-410)	1	8,12,65
5 <i>Canis familiaris</i>	Dog	6-8 mo	Spring- autumn	M	9	63(53-71)	7(1-22)	31,54,58
6 <i>Capra hircus</i>	Goat	8 mo	Sept-winter	P	21	151(135-160)	(1-5)	4,5
7 <i>Cavia porcellus</i>	Guinea pig	55-70 da	All year	P	(16-19)	68(58-75)	3(1-8)	11,46,83
8 <i>Dasyfus novemcinctus</i>	Nine-banded armadillo	1 yr	June-Aug	...	.....	(210-240)	4	75
9 <i>Didelphis marsupialis virginiana</i>	Virginia opossum	♂ 8 mo, ♀ 6 mo	Jan-Oct	P	28	(12.5-13.0)	9(5-13)	41-43
10 <i>Elephas maximus</i>	Asiatic elephant	8-16 yr	.....	P	.....	624(510-730)	Usually 1	18,27,28, 63,71
11 <i>Equus caballus</i>	Horse	1 yr	All year <sup>5</sup>	P	(10-37)	336(264-420)	Usually 1	54
12 <i>Erinaceus europaeus</i>	European hedgehog	2nd yr	Mar-Sept	M	.....	(35-49)	5(3-7)	22,45
13 <i>Felis catus</i>	Cat	6-15 mo	Feb-July	P <sup>8</sup>	(15-28)	63(52-69)	4	9,33,35, 53,73
14 <i>Macaca mulatta</i>	Rhesus monkey	♂ 3-4 yr, ♀ 1.5-2.5 yr	All year	P	28	168(144-194)	1	1,21,44, 77-80
15 <i>Mesocricetus auratus</i>	Golden hamster	5-8 wk	All year	P	4	16(15-18)	(1-12)	15,23,70
16 <i>Mus musculus</i>	House mouse	35 da	All year	P	4	(19-31)	6(1-12)	25,62
17 <i>Mustela vison</i>	Mink	1 yr	Mar-Apr	P <sup>6</sup>	(8-9)	53(39-76)	(4-10)	13,29,76
18 <i>Myotis lucifugus</i>	Little brown bat	♂, 2nd summer; ♀, end 1st summer	Autumn, spring	M <sup>7</sup>	.....	(50-60)	1	34,36,38, 59
19 <i>Ondatra zibethica</i>	Muskrat	1 yr	Apr-Oct	P	(3-5)	30(19-42)	7(1-11)	57,72
20 <i>Ornithorhynchus anatinus</i>	Platypus	1-2 yr	July-Oct	M	60	12 (incubation)	Usually 2	19,30
21 <i>Oryctolagus cuniculus</i>	European rabbit	5.5-8.5 mo	All year	P <sup>5</sup>	.....	31(30-35)	8(1-13)	7,39,61
22 <i>Ovis aries</i>	Sheep	7-8 mo	Sept-late winter <sup>9</sup>	P	(14-20)	151(144-152)	(1-4)	54,66
23 <i>Phoca vitulina</i>	Harbor seal	5-6 yr	June-Aug <sup>9</sup> , Sept <sup>10</sup>	M	.....	270	1	26,40
24 <i>Phocaena phocaena</i>	Harbor porpoise	14 mo	July-Aug	...	.....	(300-330)	1	48,60,68
25 <i>Procyon lotor</i>	Raccoon	♂ 2 yr, ♀ 1 yr	Jan-June	P	.....	63(60-73)	4(1-6)	10,74
26 <i>Rattus norvegicus</i>	Norway rat	40-60 da	All year	P	(4-5)	21	(6-9)	6
27 <i>Sciurus carolinensis</i>	Gray squirrel	1-2 yr	Dec-Aug	...	.....	44	4(1-6)	24,32,52
28 <i>Sorex araneus</i>	European shrew	2nd yr	Mar-Sept	P	.....	(13-19)	7	14
29 <i>Sus scrofa</i>	Swine	7(5-8) mo	All year	P	(18-24)	114(101-130)	9(6-15)	49,54,55, 66,69
30 <i>Tamias striatus</i>	Eastern chipmunk	2.5-3.0 mo	Mar-July	P	.....	31	(3-6)	20,33

/1/ From first day of last menses; 268(250-285)<sup>c</sup> days after rise in basal body temperature [82]. /2/ Multiple pregnancies (mainly twins) = 1.0-1.5% of total births [16]. /3/ Northern hemisphere. /4/ Southern hemisphere. /5/ Mainly spring-autumn. /6/ Induced ovulation. /7/ Ovulation in spring. /8/ Coarse-wooled breeds only; fine-wooled breeds, all year. [6] /9/ Atlantic. /10/ Pacific.

Contributors: (a) Asdell, S. A., (b) Blandau, Richard J., (c) Tietze, Christopher, (d) Van Wagenen, Gertrude

continued

## 12. PROPAGATION: MAMMALS

- References:* [1] Allen, E. 1927. *Contrib. Embryol. Carnegie Inst. Wash.* 19:1. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. *Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C.* [3] Arey, L. B. 1939. *Am. J. Obstet. Gynecol.* 37:12. [4] Asdell, S. A. 1926. *J. Agr. Sci.* 16:602. [5] Asdell, S. A. 1929. *Ibid.* 19:382. [6] Asdell, S. A. 1946. *Patterns of mammalian reproduction. Comstock, Ithaca.* [7] Asdell, S. A., and J. Hammond. 1933. *Am. J. Physiol.* 103:600. [8] Barmincev, J. 1939. *Konevodstvo* 1:42. [9] Beard, J. 1897. *The span of gestation and the cause of birth. G. Fischer, Jena.* [10] Bissonnette, T. H., and A. G. Csech. 1937. *Proc. Roy. Soc. (London), B*, 122:246. [11] Blandau, R. J., and W. C. Young. 1939. *Am. J. Anat.* 64:381. [12] Bosaev, J. 1938. *Konevodstvo* 4:44. [13] Bowness, E. R. 1942. *Can. Silver Fox Fur* 8:12. [14] Brambell, F. W. R. 1935. *Phil. Trans. Roy. Soc. London, B*, 225:1. [15] Bruce, H. M., and E. Hindle. 1934. *Proc. Zool. Soc. London*, p. 361. [16] Bunle, H. 1954. *Le mouvement naturel de la population dans le monde de 1906 à 1936. Institut National d'Études Démographiques, Paris.* [17] Burger, K., and I. Korompai. 1939. *Zentr. Gynaekol.* 63:1290. [18] Burne, E. C. 1943. *Proc. Zool. Soc. London, A*, 113:27. [19] Burrell, H. 1927. *The platypus. Angus and Robertson, Sydney.* [20] Burt, W. H. 1940. *Univ. Mich. Misc. Publ.* 45. [21] Corner, G. W. 1923. *Contrib. Embryol. Carnegie Inst. Wash.* 15:73. [22] Deanesly, R. 1934. *Phil. Trans. Roy. Soc. London, B*, 223:239. [23] Deanesly, R. 1938. *Proc. Zool. Soc. London, A*, 108:31. [24] Deanesly, R., and A. S. Parkes. 1933. *Phil. Trans. Roy. Soc. London, B*, 222:47. [25] Enzmann, E. V., N. R. Saphir, and G. Pincus. 1932. *Anat. Record* 54:325. [26] Fisher, H. D. 1954. *Nature* 173:877. [27] Flower, W. H., and R. Lydekker. 1891. *An introduction to the study of mammals living and extinct. A. and C. Black, London.* [28] Foot, A. E. 1935. *J. Bombay Nat. Hist. Soc.* 38:392. [29] Fritz, B. 1937. *Deut. Pelztierzucht*, 12:128. [30] Gatenby, J. B. 1922. *Quart. J. Microscop. Sci.* 66:475. [31] Gerlinger, H. 1925. *Le cycle sexuel chez la femelle des mammifères. Strasbourg.* [32] Goodrum, P. D. 1940. *Texas Agr. Expt. Sta. Bull.* 591. [33] Greulich, W. W. 1934. *Anat. Record* 58:217. [34] Griffin, D. R. 1940. *J. Mammal.* 21:181. [35] Gros, G. 1936. *Thesis. Univ. Algiers.* [36] Guthrie, M. J. 1933. *J. Mammal.* 14:199. [37] Haman, J. O. 1942. *Am. J. Obstet. Gynecol.* 43:870. [38] Hamilton, W. J., Jr. 1943. *The mammals of eastern United States. Comstock, Ithaca.* [39] Hammond, J., and F. H. A. Marshall. 1925. *Reproduction in the rabbit. Oliver and Boyd, Edinburgh.* [40] Harrison, R. J. 1960. *Mammalia* 24:374. [41] Hartman, C. G. 1922. *Smithsonian Inst. Ann. Rept.* 1921, p. 347. [42] Hartman, C. G. 1923. *Anat. Record* 32:353. [43] Hartman, C. G. 1928. *J. Morphol. Physiol.* 46:143. [44] Hartman, C. G. 1932. *Contrib. Embryol. Carnegie Inst. Wash.* 23:1. [45] Herter, K. 1933. *Z. Saeugetierk.* 8:195. [46] Ibsen, H. L. 1928. *J. Exptl. Zool.* 51:51. [47] Israel, S. 1959. *J. Obstet. Gynaecol. Brit. Empire* 66:311. [48] Jennison, G. 1927. *Natural history: animals. A. and C. Black, London.* app., p. 1. [49] Kenneth, J. H. 1947. *Imp. Bur. Animal Breeding Genet. Tech. Commun.* 5. [50] Kinsey, A. C., et al. 1953. *Sexual behavior in the human female. W. B. Saunders, Philadelphia.* [51] Lenner, A. 1944. *Acta Obstet. Gynecol. Scand.* 24:113. [52] Leopold, A. 1933. *Game management. Scribner's Sons, New York.* [53] Liche, H. 1939. *Nature* 143:900. [54] Lush, J. L. 1945. *Animal breeding plans. Iowa State College Press, Ames.* [55] McKenzie, F. F., and J. C. Miller. 1930. *Missouri Agr. Expt. Sta. Res. Bull.* 285:43. [56] Mackintosh, N. A. 1942-43. *Discovery Rept.* 22:197. [57] McLeod, J. A., and G. F. Bondar. 1952. *Can. J. Zool.* 30:243. [58] Marshall, F. H. A., and W. A. Jolly. 1905. *Phil. Trans. Roy. Soc. London, B*, 198:99. [59] Miller, R. E. 1939. *J. Morphol.* 64:267. [60] Møhl-Hansen, U. 1954. *Dansk Naturhist. Foren. Videnskab. Medd.*, 1, 116:369. [61] Nachtsheim, H. 1935. *Z. Zuecht.*, B, 33:343. [62] Parkes, A. S. 1926-27. *Brit. J. Exptl. Biol.* 4:93. [63] Parkes, A. S., ed. 1952-60. *Marshall's Physiology of reproduction. Ed. 3. Longmans, Green; London.* [64] Peters, H., and S. M. Shrikande. 1957. *Fertility Sterility* 8:355. [65] Pocock, R. I. 1910. *Proc. Zool. Soc. London*, p. 840. [66] Rice, V. A., et al. 1957. *Breeding and improvement of farm animals. Ed. 5. McGraw-Hill, New York.* [67] Ruud, J. T. 1945. *Hvalradets Skrifter Norske Videnskaps-Akad. Oslo* 29. [68] Scheffer, V. B., and J. W. Slepp. 1948. *Am. Midland Naturalist* 39:257. [69] Schmidt, J., E. Lamprecht, and H. Staubesand. 1936. *Z. Tierzucht. Zuechtungsbiol.* 36:55. [70] Sheehan, J. F., and J. A. Bruner. 1945. *Turttox News* 23:65. [71] Shortridge, G. C. 1934. *The mammals of South West Africa. W. Heinemann, London.* [72] Smith, F. R. 1938. *U. S. Dept. Agr.*

*continued*



## 12. PROPAGATION: MAMMALS

Circ. 474. [73] Soame, E. B. H. 1936. *Fur Feather* 96:10. [74] Stuewer, F. W. 1943. *J. Wildlife Management* 7:60. [75] Talmage, R. V., and G. D. Buchanan. 1954. *Rice Inst. Pam.* 41:109. [76] U. S. Fish and Wildlife Service. 1941. *Am. Fur Breeder* 14(4):6. [77] Van Wagenen, G. 1950. The care and breeding of laboratory animals. J. Wiley, New York. [78] Van Wagenen, G. 1952. *Anat. Record* 112:436. [79] Van Wagenen, G. 1958. In W. F. Windle, ed. *Neurological and psychological deficits of asphyxia neonatorum*. C. C. Thomas, Springfield, Ill. p. 274. [80] Van Wagenen, G., and M. E. Simpson. 1954. *Anat. Record* 118:231. [81] Wilson, D. C., and I. Sutherland. 1950. *Brit. Med. J.* 2:862. [82] Wislocki, G. B. 1930. *Contrib. Embryol. Carnegie Inst. Wash.* 22:173. [83] Young, W. C., et al. 1939. *J. Comp. Psychol.* 27:49. [84] Zenkavic, B. A. 1935. *Dokl. Akad. Nauk SSSR* 2:337.

## 13. PROPAGATION: BIRDS

For information on additional species, consult reference 1, Part I.

### Part I. NEST BUILDING, INCUBATION, AND PARENTAL CARE OF YOUNG

Symbols in columns C, D, F, G, J, and K range from 0 (no time given to specified activity) to +++ (one sex does all the work).

Family	Representative Genera	Nest Building		Incubation				Feeding and Care of Young			
		♂	♀	Duration da	Parental Activity		Duration <sup>3</sup> da	♂	♀	Trips <sup>5</sup> per Hour <sup>4</sup>	
					♂	♀					Attentive Period <sup>1</sup>
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
1 Anatidae (ducks, geese, swans)	<i>Anas, Anser, Cygnus</i>	+	+++	21-35	+	+++	4.0-23.7 hr	0	+	+++	.....
2 Cathartidae (vultures)	<i>Gyps</i>	...	....	39-56	++	++	.....	56-70	++	++	.....
3 Columbidae (pigeons, doves)	<i>Columba, Zenaidura</i>	++	++	11-19	++	++	4-20 hr	10-35	++	++	0.2-1.0
4 Corvidae (crows)	<i>Corvus, Pica</i>	++	++	16-20	+	+++	35-150 min	20-38	+++	+++	1.7-4.0
5 Fringillidae (finches, sparrows)	<i>Melospiza, Passer, Serinus</i>	+	+++	11-14	+	+++	14.6 min to continuous	8-17	+++	+++	1.3-21.3
6 Laridae (gulls)	<i>Larus</i>	++	++	20-34	++	++	0.5-24.0 hr	0-several	++	++	0.2-7.0
7 Meleagridae (turkeys)	<i>Meleagris</i>	0	++++	28	0	++++	23-24 hr	0	0	++++	.....
8 Phasianidae (quail, pheasant)	<i>Colinus, Phasianus</i>	++	++	21-28	+	+++	7-23+ hr	0	++	++	.....
9 Spheniscidae (penguins)	<i>Aptenodytes</i>	++	++	38-56	++	++	1-5+ da	56-112	++	++	1-2/da
10 Struthionidae (ostriches)	<i>Struthio</i>	++++	0	42	+++	+	9-15 hr	0	+++	+++	.....
11 Sturnidae (starlings)	<i>Sturnus</i>	++	++	12	++	++	20 min	20-21	++	++	19-22
12 Trochilidae (hummingbirds)	<i>Archilochus</i>	0	++++	15-17	0	++++	5-99 min	19-25	0	++++	1.1-3.3
13 Troglodytidae (wrens)	<i>Troglodytes</i>	++	++	13-19	0	++++	12-86 min	13-22	++	++	5.6-19.2
14 Turdidae (thrushes)	<i>Turdus</i>	+	+++	12-16	+	+++	12-120 min	12-18	++	++	5.5-38.5

/1/ Time parent sits on eggs at one sitting. /2/ Nestling period, time from hatching until young birds leave nest. /3/ Range of averages for various nests. /4/ Unless otherwise stated.

Contributors: (a) Kendeigh, S. Charles, (b) Skutch, Alexander F.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. *Growth, including reproduction and morphological development*. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Kendeigh, S. C. 1952. *Parental care and its evolution in birds*. Univ. Illinois Press, Urbana. [3] Skutch, A. F. Unpublished. San Isidro del General, Costa Rica, 1961.

*continued*

## 13. PROPAGATION: BIRDS

### Part II. CLUTCH SIZE

Requirements for selection of data: (i) at least 25 clutches recorded within less than 10 years and within a limited geographical area, (ii) proof that clutches were complete when counted, and (iii) proof that only one female laid in nest.

	Species	Common Name	Location	Clutches no.	Eggs/Clutch		Observer and Year
					Mean	Standard Deviation	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	<i>Anas platyrhynchos</i>	Mallard duck	California	178	9.2	....	Miller, 1954
2			California	108	8.5	....	Hunt, 1955
3	<i>Larus argentatus</i>	Herring gull	New Brunswick	1,011	2.38	0.71	Paynter, 1949
4			Holland	217	2.91	0.34	Paludan, 1951
5	<i>Melospiza melodia</i>	Song sparrow	Ohio	210	4.07	0.81	Nice, 1937
6	<i>Perdix perdix</i>	European partridge	England	4,051	14.6	2.38	Lack, 1947
7	<i>Phasianus colchicus</i>	Ring-necked pheasant	Iowa	60	8.7	....	Kozicky, 1956
8			Pennsylvania	157	10.6	3.18	Randall, 1941
9	<i>Pica nuttalli</i>	Yellow-billed magpie	California	70	6.5	0.92	Linsdale, 1937
10	<i>Sturnus vulgaris</i>	Starling	Maryland	101	4.54	1.15	McAtee, 1940
11			England	105	4.85	1.08	Lack, 1948
12			Germany	95	4.44	0.99	Berndt, 1939
13			Holland	1,785	5.14	1.11	Lack, 1948
14	<i>Troglodytes aedon</i>	House wren	Maryland	98	5.46	1.10	McAtee, 1940
15	<i>Turdus migratorius</i>	American robin	New York	127	3.39	0.62	Howell, 1942

Contributor: Davis, David E.

References: [1] Davis, D. E. 1955. In A. Wolfson, ed. Recent studies in avian biology. Univ. Illinois Press, Urbana. p. 264. [2] Davis, D. E. 1960. In H. S. Mosby, ed. Manual of game investigational techniques. Edwards Brothers, Ann Arbor, Mich. sect. 19, p. 4.

### Part III. HATCHING SUCCESS: PRECOICIAL SPECIES

Only studies that recorded at least 50 nests have been included.

	Species (Common Name)	Location	Nests		Eggs		Observer and Year
			no.	Successful %	no.	Hatched %	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	<i>Anas platyrhynchos</i>	California	161	38.5	616	83.4	Anderson, 1957
2	(mallard duck)	California	60	52	417	49.4	Earl, 1950
3		California	209	85.2	1,622	91.4	Miller, 1954
4		Montana	....	....	1,793	71.2	Girard, 1941
5		Utah	185	59	1,582	60	Williams, 1937
6	<i>Colinus virginianus</i>	Georgia-Florida	602	36	.....	....	Stoddard, 1931
7	(bobwhite quail)	Texas	189	46	.....	....	Lehmann, 1946
8		Texas	59	62.9	.....	....	Parmalee, 1955
9		Wisconsin	53	50.9	.....	....	Errington, 1933
10	<i>Perdix perdix</i> (European partridge)	England	....	....	57,202	90.4	Lack, 1947
11		England	7,251	78	.....	....	Middleton, 1935
12		England	4,090	....	59,825	93	Middleton, 1935
13		Michigan	143	32	.....	....	Yeatter, 1934
14		Washington	113	37.1	.....	....	Knott, 1943
15		Wisconsin	435	32	.....	....	McCabe, 1946
16	<i>Phasianus colchicus</i>	Colorado	333	65	.....	....	Yeager, 1951
17	(ring-necked pheasant)	Iowa	533	25.5	1,319	83	Baskett, 1947
18		Iowa	64	....	723	82.3	Hamerstrom, 1936
19		Michigan	193	35	.....	....	English, 1946
20		Minnesota	241	28.6	.....	....	Erickson, 1951
21		Ohio	563	58	.....	....	Leedy, 1945
22		Ohio	358	72	.....	....	Strode, 1946
23		Ontario	230	32.1	777	73.5	Ball, 1952

continued

**13. PROPAGATION: BIRDS**

**Part III. HATCHING SUCCESS: PRECOICIAL SPECIES**

	Species (Common Name)	Location	Nests		Eggs		Observer and Year
			no.	Successful %	no.	Hatched %	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
24	<i>Phasianus colchicus</i>	Oregon	145	44.8	.....	....	Eklund, 1942
25	(ring-necked pheasant)	Pennsylvania	310	20.3	.....	....	Randall, 1940
26		Utah	149	36	.....	....	Rasmussen, 1945
27		Washington	63	27	.....	....	Buss, 1950
28		Wisconsin	126	71.3	1,000	78.9	Errington, 1937

*Contributors:* (a) Davis, David E., (b) Nice, Margaret Morse

*References:* [1] Davis, D. E. 1960. In H. S. Mosby, ed. Manual of game investigational techniques. Edwards Brothers, Ann Arbor, Mich. sect. 19, p. 1. [2] Hickey, J. J. 1955. In A. Wolfson, ed. Recent studies in avian biology. Univ. Illinois Press, Urbana. p. 326.

**Part IV. HATCHING AND FLEDGING SUCCESS: ALTRICIAL SPECIES**

For additional information, consult reference 10.

	Species (Common Name)	Years Observed	Nests no.	Hatched		Fledged		Refer- ence	
				no.	%	no.	%		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
<b>Hole-Nesting Species</b>									
1	<i>Passer domesticus</i> (house sparrow)	6	....	114	....	....	97	78.5	6
2	<i>Sturnus vulgaris</i> (starling)	...	....	10,557	....	....	7,923	75.1	5
3		6	....	472	....	....	410	84.5	6
4	<i>Troglodytes aedon</i> (house wren)	19	1,056	6,673	5,576	82.3	5,351	79.0	3
5		3	34	211	135	64.0	118	55.2	4
6		6	....	469	....	....	339	83.7	6
7		21	64	333	199	59.7	161	48.3	11
<b>Open-Nest Species</b>									
8	<i>Melospiza melodia</i> (song sparrow)	...	77	585	389	66.5	243	41.5	9
9		...	30	321	147	45.8	80	24.9	
10	<i>Turdus migratorius</i> (American robin)	...	78	259	157	60.6	131	54.4	2
11		...	86	548	316	57.8	246	44.9	12
12	<i>Zenaidura macroura</i> (mourning dove)	...	130	500	...	....	213	42.6	8
13		...	2,043	8,018	4,379	54.6	3,734	46.6	7
14		...	142	398	310	77.8	274	68.8	1

*Contributor:* Nice, Margaret Morse

*References:* [1] Cowan, J. B. 1952. Calif. Fish Game 38:505. [2] Howell, J. C. 1942. Am. Midland Naturalist 28:529. [3] Kendeigh, S. C. 1942. J. Wildlife Management 6:19. [4] Kuerzi, R. G. 1941. Proc. Linnaean Soc. N. Y. 52-53:1. [5] Lack, D. 1948. Evolution 2:95. [6] McAtee, W. L. 1940. Auk 57:333. [7] McClure, H. E. 1946. Ibid. 63:24. [8] Nice, M. M. 1931. Univ. Oklahoma Biol. Survey 3(1):99. [9] Nice, M. M. 1937. Trans. Linnaean Soc. N. Y. 4:143. [10] Nice, M. M. 1957. Auk 74:305. [11] Walkinshaw, L. H. 1941. Wilson Bull. 53:1. [12] Young, H. 1955. Am. Midland Naturalist 53:329.

## 14. PROPAGATION: REPTILES

For information on additional species, consult reference 1. The manner of fertilization for all reptiles is internal (by copulation). **Gestation Time** (column E) = period from copulation to parturition (ovoviviparous reptiles); **Incubation Time** = period from laying to hatching of eggs (oviparous reptiles). **Manner of Birth** (column F): Ovo = ovoviviparous; O = oviparous. Values in parentheses are ranges, estimate "c" or "d" (cf. Introduction).

	Species	Common Name	Age at Sexual Maturity <sup>1</sup> yr	Breeding Season <sup>2</sup>	Gestation or Incubation Time <sup>3</sup>	Manner of Birth	Brood or Clutch <sup>4</sup>		Reference
							Size <sup>5</sup>	no./yr	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
1	<i>Alligator mississippiensis</i>	American alligator	(5-10)	Jan-Sept	(56-66) da	O	(29-88)	1	9,11
2	<i>Ancistrodon contortrix mokeson</i>	Northern U.S. copperhead	(2-4)	Apr-May	142 da	Ovo	(1-14)	1	4,5,14
3	<i>Anguis fragilis</i>	Slow worm	(3-4)	May-June	3 mo	Ovo	(7-19)	1	13,14
4	<i>Anolis carolinensis</i>	American "chameleon"	♂ 2, ♀ 1	Apr-Aug	(6-7) wk	O	(8-10)	1	6,15
5	<i>Caretta caretta</i>	Loggerhead turtle	.....	Mar-July	(31-65) da	O	(120-130)	(2-3)	2
6	<i>Chelydra serpentina</i>	Snapping turtle	.....	Apr-Nov	(81-90) da	O	25(8-80)	(1-2)	2
7	<i>Coluber constrictor</i>	American black snake	.....	May-June	(1-2) mo	O	(15-25)	1	14
8	<i>Crotalus viridis</i>	Prairie rattlesnake	(3-4)	Apr-June	(4-5) mo	Ovo	(3-13)	0.5	8,10,17
9	<i>Eumeces fasciatus</i>	Five-lined skink	<2	May-June	(4-9) wk	O	(2-18)	1	3,15
10	<i>Heloderma suspectum</i>	Gila monster	.....	.....	1 mo	O	(5-13)	1	16
11	<i>Malaclemys terrapin</i>	Diamondback terrapin	(5-6)	Spring	3 mo	O	8	(1-3)	7
12	<i>Natrix erythrogaster</i>	Copper-bellied water snake	.....	Apr-May	.....	Ovo	(8-27)	1	14
13	<i>Phrynosoma cornutum</i>	Horned lizard	.....	Apr-May	(39-47) da	O	(23-37)	1	15
14	<i>Pseudemys floridana peninsularis</i>	Peninsular turtle	.....	Nov-June	5 mo	O	(12-29)	2	2
15	<i>Sceloporus graciosus</i>	Sagebrush lizard	.....	Apr-May	62 da	O	(2-7)	1	15
16	<i>Sternotherus odoratus</i>	Musk turtle	♂(2-3), ♀(9-11)	Apr-Oct	(60-75) da	O	(1-5)	(1-2)	2,12
17	<i>Terrapene carolina</i>	Box turtle	(4-5)	Apr-May	88(70-114) da	O	(2-7)	1	2
18	<i>Thamnophis sirtalis</i>	Common garter snake	(2-3)	Mar-May and fall	(87-116) da	Ovo	28(6-51)	(1-2)	14

/1/ Males in some species mature before females. /2/ Varies with geographical location. /3/ Actual values expressed in days; approximations, in weeks or months. /4/ Brood = young produced at one time; clutch = eggs laid at one time. /5/ Number of eggs or young.

Contributors: (a) Altland, Paul D., (b) Fitch, Henry S., (c) Tanner, Vasco M.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Carr, A. 1952. Handbook of turtles. Comstock, Ithaca. [3] Fitch, H. S. 1954. Univ. Kansas Museum Nat. Hist. Publ. 8:145. [4] Fitch, H. S. 1956. Ibid. 8:269. [5] Fitch, H. S. 1960. Ibid. 13:272. [6] Hamlett, G. W. D. 1952. Copeia, p. 183. [7] Hildebrande, C. F. 1929. U. S. Fish Wildlife Serv. Fishery Bull. 45:25. [8] Klauber, L. M. 1936. Trans. San Diego Soc. Nat. Hist. 8:20. [9] McIlhenny, E. A. 1935. The alligator's life history. Christopher, Boston. [10] Rahn, H. 1942. Copeia, p. 233. [11] Reese, A. M. 1915. The alligator and its allies. G. P. Putnam, New York. [12] Risley, P. L. 1933. Papers Mich. Acad. Sci. 17:685. [13] Sanders, E. 1943. A beast book for the pocket. Oxford Univ. Press, London. [14] Schmidt, K. P., and D. D. Davis. 1941. Field book of snakes of the United States and Canada. G. P. Putnam, New York. [15] Smith, H. M. 1946. Handbook of lizards. Comstock, Ithaca. [16] Van Denburgh, J. 1922. The reptiles of western North America. California Academy of Sciences, San Francisco. v. 1. [17] Woodbury, A. M., et al. 1951. Herpetologica 7:24.



## 15. PROPAGATION: AMPHIBIANS

For information on additional species, consult reference 1. The rate of growth and development of amphibians is influenced by temperature, moisture, and light to a much greater degree than is the rate of growth and development of homoiotherms. Manner of development is oviparous. **Fertilization** (column D): Ext = external; Int = internal. **Parental Care** (column G): 0 = none; ♀ = female guards or transports eggs; ♂ = male guards or transports eggs.

Species	Common Name	Breeding Season	Fertilization	Eggs or Young per Brood	Egg Development	Parental Care	Form at Hatching	Period of Growth			Reference
								Egg da	Larva da	Sexual Maturity	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
1 <i>Ambystoma maculatum</i>	Spotted salamander	Mar-Apr	Int <sup>1</sup>	Up to 250	In water	0	Larva	31-54	61-110	2 yr	3
2 <i>A. tigrinum</i>	Tiger salamander	Jan-Mar	Int <sup>1</sup>	23-110	In water	0	Larva	23-30	180+	1 yr	3,11
3 <i>Amphiuma tri-dactylum</i>	Three-toed amphiuma	Jan-May <sup>2</sup> ; May-June <sup>3</sup>	Int <sup>4</sup>	42-131	In water	♀	Adult		2-3 yr		5
4 <i>Bufo americanus</i>	American toad	Mar-July	Ext	4,000-20,600	.....	...	.....	3-12	30-65	2-3 yr	4,7,11,14
5 <i>Cryptobranchus alleganiensis</i>	Hellbender	Aug-Dec	Ext	220-450	In water	♂	Larva	68-84	550-700	5-6 yr	2,3
6 <i>Hyla regilla</i>	Pacific tree frog	Jan-May	Ext	730-1,250	.....	...	.....	6-14	50-80	2 yr	8,12,14
7 <i>Necturus maculosus</i>	Mud puppy	Sept-Nov <sup>2</sup> ; May-June <sup>5</sup>	Int <sup>4</sup>	18-180	In water	...	Larva	38-63		5 yr	2
8 <i>Pipa pipa</i>	Surinam toad	.....	Ext	.....	In water (back of ♀)	♀	Adult	.....	.....	.....	6
9 <i>Rana catesbeiana</i>	American bullfrog	Feb-Aug	Ext	10,000-25,000	In water	0	Embryo	4-5	365-730	3-4 yr	12-14
10 <i>R. pipiens</i>	Leopard frog	Feb-Dec	Ext	3,500-6,500	In water	0	Embryo	9-20	60-80	3 yr	12-14
11 <i>Triturus viridescens</i>	Common newt	Apr-June	Int <sup>1</sup>	200-376	In water	0	Larva	20-35	80+	2 yr	3,11
12 <i>Xenopus laevis</i>	Clawed frog	Sept-Oct	Ext	<100-1,000	In water	0	Embryo	3	35-300	♂ <sup>1</sup> / <sub>2</sub> , ♀ <sup>2</sup> yr	9,10

/1/ Spermatophore laid by male and picked up by female. /2/ Mating season. /3/ Time of oviposition. /4/ Spermatophore deposited by male in female cloaca.

**Contributors:** (a) Cagle, Fred R., (b) Blair, Albert P., (c) Tanner, Vasco M., (d) Fitch, Henry S.

**References:** [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Bishop, S. C. 1941. N. Y. State Museum Bull. 324. [3] Bishop, S. C. 1943. Handbook of salamanders. Comstock, Ithaca. [4] Bragg, A. N. 1940. Am. Midland Naturalist 24:322. [5] Cagle, F. R. 1948. Ecology 29(4):479. [6] Dunn, E. R. 1942. Bull. Museum Comp. Zool. Harvard Univ. 91(6):440. [7] Hamilton, J. W., Jr. 1934. Copeia, p. 88. [8] Livezey, R. L., and A. H. Wright. 1947. Am. Midland Naturalist 37(1):179. [9] Noble, G. K. 1931. The biology of the amphibia. McGraw-Hill, New York. [10] Orton, G. L. 1949. Ann. Carnegie Museum 31:257. [11] Smith, H. M. 1950. Univ. Kansas Museum Nat. Hist. Misc. Publ. 2. [12] Stebbins, R. C. 1951. Amphibians of western North America. Univ. California Press, Berkeley. [13] Wright, A. H. 1932. Life histories of the frog of Okefinokee Swamp, Georgia. Macmillan, New York. [14] Wright, A. H., and A. A. Wright. 1949. Handbook of frogs and toads. Comstock, Ithaca.

## 16. PROPAGATION: FISHES

For additional information, consult references 2,13,21,25,31-33,38 Spawning activities vary with species, locale, and water temperature. The number of eggs produced varies (from a few to millions) with the species. The number of eggs may also differ greatly within a single species, depending chiefly on the size of the female. **Water** (column D): F = fresh; S = salt; B = brackish; L = lacustrine; V = fluviatile; (a) = anadromous. **Fertilization** (column E): Ext = external; Int = internal. **Egg Type** (column F): D = demersal; PB = pelagic or buoyant eggs. **Development or Birth** (column G): O = oviparous; Ovo = ovoviviparous; V = viviparous. **Parental Care** (column I): 0 = none; ♂ = male guards nest or eggs; C = eggs covered by gravel or sand.

Species	Common Name	Spawning		Ferti- li- za- tion	Egg Type	De- vel- op- ment or Birth	Eggs or Young per Spawning Period	Pa- ren- tal Care	Refer- ence	
		Season	Water							
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
Pisces										
1	<i>Acipenser fulvescens</i>	Lake sturgeon	Spring, summer	F, V, L	Ext	D	O	182,000-1,000,000	0	11,20,24
2	<i>Amia calva</i>	Bowfin	Spring	F, V, L	Ext	D	O	23,000-64,000	♂	1,12,24
3	<i>Anguilla rostrata</i>	American freshwater eel	Winter	S	Ext	PB	O	5,000,000-20,000,000	...	6,22
4	<i>Carassius auratus</i>	Goldfish	Spring	F, L	Ext	D	O	3,000	0	24,40
5	<i>Clupea harengus</i>	Atlantic herring	Spring- autumn	S	Ext	D	O	20,000-40,000	0	6,22
6	<i>Coregonus clupeaformis</i>	North American lake whitefish	Autumn	F, V, L	Ext	D	O	10,000-75,000	0	12,14, 24,34
7	<i>Cyprinus carpio</i>	Carp	Spring, summer	F, L	Ext	D	O	500,000-2,000,000	0	18
8	<i>Esox lucius</i>	Northern pike	Spring	F, V, L	Ext	D	O	10,000-100,000	0	1,8,9,34
9	<i>Fundulus heteroclitus</i>	Mummichog	Spring, summer	F, V, B	Ext	D	O	460	0	22
10	<i>Gadus morhua</i>	Atlantic cod	Winter, spring	S	Ext	PB	O	3,000,000-9,000,000	0	6
11	<i>Hippocampus hudsonius</i>	Atlantic sea horse	.....	S	Ext	...	O	150	♂	6,17,35
12	<i>Hippoglossus hippoglossus</i>	Atlantic halibut	Spring, summer	S	Ext	PB	O	2,182,773	...	6,19
13	<i>Ictalurus punctatus</i>	Channel catfish	Spring, summer	F, V	Ext	D	O	3,000-20,000	♂	20
14	<i>Lepisosteus osseus</i>	Longnose gar	Spring	F, V, L	Ext	D	O	6,200-77,156	...	23,26,34
15	<i>Lepomis macrochirus</i>	Bluegill	Spring, summer	F, V, L	Ext	D	O	4,670-61,815	♂	20
16	<i>Melanogrammus aeglefinus</i>	Haddock	Winter, spring	S	Ext	PB	O	169,000-1,839,581	0	6
17	<i>Micropterus salmoides</i>	Largemouth black bass	Spring, summer	F, V, L	Ext	D	O	2,000-26,000	♂	1,12
18	<i>Osmerus mordax</i>	American smelt	Spring	F, V	Ext	D	O	To 50,000	0	7,14,34
19	<i>Perca flavescens</i>	Yellow perch	Spring	F, V, L	Ext	D	O	10,000-40,000	0	1,20
20	<i>Polyodon spathula</i>	Paddlefish	Spring, winter	F, V	Ext	D	O	140,000	0	12,30, 36,37
21	<i>Pomoxis annularis</i>	White crappie	Spring, summer	F, V, L	Ext	D	O	2,900-14,750	♂	20
22	<i>Pseudopleuronectes americanus</i>	Winter flounder	Winter, spring	S	Ext	D	O	500,000-1,500,000	...	6
23	<i>Salmo salar</i>	Atlantic salmon	Spring	F, V (a)	Ext	D	O	7,000	C	15,29,36
24	<i>S. trutta</i>	Brown trout	Autumn, winter	F, V	Ext	D	O	200-6,000	C	20
25	<i>Salvelinus fontinalis</i>	Eastern brook trout	Autumn	F, V	Ext	D	O	200-2,500	C	20
26	<i>Scomber scombrus</i>	Atlantic mack- erel	Spring, summer	S	Ext	PB	O	41,000-546,000	0	28
Chondrichthyes										
27	<i>Dasyatis americana</i>	Southern sting- ray	.....	S	Int	...	Ovo	3-5	0	5
28	<i>Raja erinacea</i>	Little skate	All year	S	Int	D	O	6	0	5
29	<i>Sphyrna zygaena</i>	Hammerhead shark	Summer	S	Int	...	V	29-37	0	3,4
30	<i>Squalus acanthias</i>	Atlantic spiny dogfish	All year	S	Int	...	Ovo	2-11	0	4

continued

## 16. PROPAGATION: FISHES

Species	Common Name	Spawning		Ferti- li- za- tion	Egg Type	De- vel- op- ment or Birth	Eggs or Young per Spawning Period	Pa- ren- tal Care	Refer- ence	
		Season	Water							
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
Agnatha										
31	<i>Lampetra lamottei</i>	American brook lamprey	Spring	F, V	Ext	D	O	1,085-3,648	0	10,24,27
32	<i>Myxine glutinosa</i>	Atlantic hagfish	All year	S	....	D	O	19-30	0	4
33	<i>Petromyzon marinus</i>	Sea lamprey	Spring	F, V (a)	Ext	D	O	13,000-259,000	0	16,39,41

*Contributors:* (a) Migdalski, Edward C., (b) Katz, Max, (c) Carlander, Kenneth D.

*References:* [1] Adams, C. C., and T. L. Hankenson. 1928. Roosevelt Wild Life Ann. 1:241. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Baughman, J. L., and S. Springer. 1950. Am. Midland Naturalist 44:96. [4] Bigelow, H. B., and W. C. Schroeder. 1948. Fishes of the western North Atlantic. Yale Univ. Sears Foundation for Marine Research, New Haven. pt. 1. [5] Bigelow, H. B., and W. C. Schroeder. 1953. Ibid. pt. 2. [6] Bigelow, H. B., and W. C. Schroeder. 1953. U. S. Fish Wildlife Serv. Fishery Bull. 74. [7] Breder, C. M., Jr. 1948. Field book of marine fishes of the Atlantic Coast. G. P. Putnam's Sons, New York. [8] Carbine, W. F. 1942. Trans. Am. Fisheries Soc. 71:149. [9] Carbine, W. F. 1943. Papers Mich. Acad. Sci. 29:123. [10] Carlander, K. D. 1950. Handbook of freshwater fishery biology. W. C. Brown, Dubuque, Iowa. [11] Cuerrier, J. P. Unpublished, 1949. [12] Eddy, S., and T. Surber. 1947. Northern fishes. Univ. Minnesota Press, Minneapolis. [13] Eddy, S., and T. Surber. 1960. Ibid. Rev. ed. C. T. Branford, Newton Centre, Mass. [14] Everhart, H. W. 1950. Fishes of Maine. Maine Dept. Inland Fisheries and Game, Augusta. [15] Forbes, S. A., and R. E. Richardson. 1920. The fishes of Illinois. Ed. 2. Illinois Natural History Society, Danville. [16] Gage, S. H. 1893. In Wilder quarter century book. Comstock, Ithaca. p. 421. [17] Gill, T. 1905. Proc. U. S. Natl. Museum 28:805. [18] Gill, T. 1905-07. Smithsonian Inst. Misc. Collections 48:195. [19] Goode, G. B., et al. 1884. The food fishes of the United States. U. S. Commission of Fish and Fisheries, Washington, D. C. sect. 1(3). [20] Harlan, J. R., and E. B. Speaker. 1951. Iowa fish and fishing. Iowa State Conservation Commission, Des Moines. [21] Harlan, J. R., and E. B. Speaker. 1956. Ibid. Rev. ed. [22] Hildebrand, S. F., and W. C. Schroeder. 1927. U. S. Bur. Fisheries Bull. 43(1). [23] Holloway, A. 1954. J. Wildlife Management 18:438. [24] Hubbs, C. L., and K. F. Lagler. 1947. Cranbrook Inst. Sci. Bull. 26. [25] Hubbs, C. L., and K. F. Lagler. 1957. Fishes of the Great Lakes region. Cranbrook Institute of Science, Bloomfield Hills, Mich. [26] Knapp, F. T. 1953. Fishes found in the fresh waters of Texas. Ragland and Litho Print, Brunswick, Georgia. [27] Legendre, V. 1954. The freshwater fishes of Quebec. Société Canadienne d'Écologie, Quebec. v. 1. [28] MacCay, C. 1929. Bull. Boston Soc. Nat. Hist. 53. [29] MacFarland, W. L. 1925. Salmon of the Atlantic. Parke, Austin, and Lipscomb; New York. [30] Meyer, F. P. 1960. Ph.D. Thesis. Iowa State Univ., Iowa City. [31] Migdalski, E. C. 1958. Salt water game fishes--Atlantic and Pacific. Ronald Press, New York. [32] Migdalski, E. C. 1962. Fresh water sport fishes of North America. Ronald Press, New York. [33] Perlmutter, A. 1961. Guide to marine fishes. New York Univ. Press, New York. [34] Raney, E. C. 1951. In A. J. McClane, ed. Wise fisherman's encyclopedia. W. H. Wise, New York. p. 647. [35] Ryder, J. A. 1881. Bull. U. S. Fisheries Comm. 1:191. [36] Scott, W. B. 1954. Freshwater fishes of eastern Canada. Univ. Toronto Press, Ontario. [37] Thompson, D. H. 1933. Copeia, p. 33. [38] Trautman, M. B. 1957. The fishes of Ohio. Ohio State Univ. Press, Columbus. [39] Vladykov, V. D. 1951. Can. Fish Culturist 10:1. [40] Watson, F. R., and F. Perry. 1948. Fishponds and home aquaria. Collingridge, London. [41] Wigley, R. L. 1959. U. S. Fish Wildlife Serv. Fishery Bull. 59:561.

## 17. PROPAGATION: AQUATIC INVERTEBRATES

For information on additional species, consult reference 2. Breeding habits of invertebrates may vary with changes in location, temperature, light, and, for marine forms, with changes in salinity. **Type of Sexuality** (column F): D = dioecious (unisexual); M = monoecious (bisexual or hermaphroditic). **Dimorphism** (column G): + = sexual dimorphism; - = no sexual dimorphism. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Class and Species (Common Name)	Distribution <sup>1</sup>	Sexual Maturity		Breeding Season	Sexuality		Eggs or Young per Brood	Refer- ence
		Age at Onset	Size <sup>2</sup> mm		Type	Di- mor- phism		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
<b>Echinodermata<sup>3</sup></b>								
<b>Asteroida</b>								
1 <i>Asterias forbesi</i> (starfish)	Mexico to Maine (Long Island Sound)	(1-2) yr	(60-210) <sup>4</sup>	July- Oct	D <sup>5</sup>	-	Several thousand	1,9,10, 19
<b>Arthropoda</b>								
<b>Merostomata<sup>3</sup></b>								
2 <i>Limulus polyphemus</i> (king crab)	Yucatan to Nova Scotia (Delaware Bay)	(9-11) yr	♂(178-258), ♀(243-351) <sup>6</sup>	May- June	D	+	3,000	29
<b>Crustacea<sup>7</sup></b>								
3 <i>Callinectes sapidus</i> (blue crab)	Uruguay to Nova Scotia (Chesapeake Bay)	♂♀ 13 mo	♂(135-215), ♀(134-185)	July- Aug	D	+	1,750,000	8,22
4 <i>Cyclops viridis</i> (cyclops)	U.S. and Europe (Germany)	♂(41-132), ♀(36-128) da	♀(1.5-5.0) <sup>8</sup>	All year	D	+	75(20-160) <sup>9</sup>	6,34,35
5 <i>Daphnia longispina</i> (water flea)	Asia, Europe, N. America (Florida)	♀(75-86) hr	♂ 1.2, ♀ 1.9	All year except winter	D <sup>10</sup>	+	28(4-35)	3,4,6, 15
6 <i>Homarus ameri- canus</i> (American lobster)	Newfoundland to N. Carolina	(4-5) yr	♂(170-600), ♀(180-480)	July- Sept	D	+	8,500 <sup>11</sup>	14,32
7 <i>Orconectes im- munis</i> (crayfish)	Mississippi River & Great Lakes drain- age (New York State)	♂♀ 15 mo	♂(40-60), ♀(44-90)	June- Oct	D	+	102(84-195)	24,26, 30
<b>Mollusca<sup>3</sup></b>								
<b>Bivalvia</b>								
8 <i>Crassostrea vir- ginica</i> (eastern oyster)	Texas to Canada (Delaware Bay)	1 yr	(25-50)	June- Aug	M <sup>6</sup>	-	(500,000- 1,000,000)	16,18, 29
9 <i>Mercenaria mer- cenaria</i> (quahog)	Nova Scotia to Yuca- tan (Baja Califor- nia)	(1-2) yr	(50-70)	July- Aug	M <sup>6</sup>	-	ca. 1,000,000	7,17
10 <i>Mytilus edulis</i> (mussel)	Worldwide	(1-2) yr	.....	May- Sept	D	...	.....	18
11 <i>Pecten irradians</i> (scallop)	Maine to Mexico	12 mo	78	.....	M	-	.....	11
<b>Gastropoda</b>								
12 <i>Busycon canalicu- latum</i> (whelk)	Cape Cod to Mexico	.....	.....	.....	D	+	(360-6,240)	20,27, 28
13 <i>Helix pomatia</i> (land snail)	Europe and U.S.	(33-39) mo	.....	May- July	M <sup>12</sup>	-	(40-200)	5,21,25, 27,28
14 <i>Littorina littorea</i> (periwinkle)	N. Atlantic to Flor- ida	.....	.....	.....	D	+	(1-3)	33
15 <i>Lymnaea stagnalis</i> (freshwater snail)	Worldwide (Wiscon- sin)	(4-14) mo	(50-60)	July- Oct	M <sup>12</sup>	-	6,000	23,28
<b>Polyplacophora</b>								
16 <i>Ischnochiton mag- dalenensis</i> (chiton)	California to Mexico	2 yr	(35-36)	.....	D	...	57,970	12,13, 27,28, 31

/1/ Habitat of animals to which data apply is given in parentheses. /2/ Greatest dimension. /3/ All species listed are oviparous. /4/ Size is dependent on food supply. /5/ Protandrous hermaphrodite: male organs appear first, later replaced by female organs. /6/ Prosomal width. /7/ All species listed are ovigerous. /8/ Male is smaller. /9/ In early summer. /10/ Parthenogenetic reproduction during most of season. /11/ Under present fisheries conditions, average is difficult to obtain because females are not permitted to attain maximum egg-laying age. /12/ Cross-fertilization.

continued



## 17. PROPAGATION: AQUATIC INVERTEBRATES

*Contributors:* (a) Carriker, Melbourne R., (b) Abbott, R. Tucker, (c) Lochhead, John H., (d) Aldrich, Frederick A., (e) Loosanoff, Victor L., (f) Sellmer, George P., (g) Shuster, Carl N., Jr.

- References:* [1] Aldrich, F. A., and M. L. Aldrich. 1955. *Notulae Naturae Acad. Nat. Sci. Phila.* 276:1. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. *Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C.* [3] Banta, A. M. 1939. *Carnegie Inst. Wash. Publ.* 513:60. [4] Banta, A. M., and L. A. Brown. 1939. *Ibid.* 513:106. [5] Boycott, A. E. 1934. *J. Ecol.* 22(1):1. [6] Brown, F. A., Jr., ed. 1950. *Selected invertebrate types. J. Wiley, New York.* [7] Carriker, M. R. 1961. *J. Elisha Mitchell Sci. Soc.* 77(2):168. [8] Churchill, E. P. 1919. *U. S. Bur. Fisheries Bull.* 36:93. [9] Coe, W. R. 1912. *Conn. State Geol. Nat. Hist. Survey Bull.* 19:1. [10] Galtsoff, P. S., and V. L. Loosanoff. 1939. *U. S. Bur. Fisheries Bull.* 49:75. [11] Gutsell, J. S. 1931. *Ibid.* 46:569. [12] Heath, H. 1899. *Zool. Jahrb. Abt. Anat. Ontog. Tiere* 12:567. [13] Heath, H. 1906. *Zool. Anz.* 29:390. [14] Herrick, F. H. 1911. *U. S. Bur. Fisheries Bull.* 29:149. [15] Ingle, L., T. R. Wood, and A. M. Banta. 1937. *J. Exptl. Zool.* 76:325. [16] Korringa, P. 1952. *Quart. Rev. Biol.* 27:266. [17] Loosanoff, V. L. 1937. *Biol. Bull.* 72(3):389. [18] Loosanoff, V. L. Unpublished. U. S. Dept. of Interior, Milford, Conn., 1952. [19] Loosanoff, V. L., J. B. Engle, and C. A. Nomejko. 1955. *Biol. Bull.* 109(1):75. [20] Magalhaes, H. 1948. *Ecol. Monographs* 18(3):377. [21] Meisenheimer, J. 1907. *Zool. Jahrb. Abt. System. Oekol. Geog. Tiere* 25:461. [22] Newcombe, C. L., F. Campbell, and A. M. Eckstine. 1949. *Growth* 13:71. [23] Noland, L. E., and M. R. Carriker. 1946. *Am. Midland Naturalist* 36(2):467. [24] Pearse, A. S. 1909. *Am. Naturalist* 43:746. [25] Pelseneer, P. 1935. *Essai d'éthologie zoologiques d'après l'étude des mollusques. Bruselles.* [26] Pennak, R. W. 1953. *Fresh-water invertebrates of the United States. Ronald Press, New York.* [27] Pratt, H. S. 1948. *A manual of the common invertebrate animals. Blakiston, Philadelphia.* [28] Rogers, J. E. 1951. *The shell book. C. T. Branford, Boston.* [29] Shuster, C. N., Jr. Unpublished. Univ. Delaware, Newark, 1955. [30] Tack, P. I. 1941. *Am. Midland Naturalist* 25:420. [31] Taki, I. 1940. *Proc. Pacific Sci. Congr. Pacific Sci. Assoc.* 3:487. [32] Templeman, W. 1940. *Newfoundland Dept. Nat. Resources Serv. Bull. (Fisheries)* 15. [33] Thorson, G. 1946. *Medd. Komm. Danmarks Fisk-Havun. Plankton* 4(1). [34] Walter, E. 1922. *Zool. Jahrb. Abt. System. Oekol. Geog. Tiere* 44:375. [35] Yeatman, H. C. 1944. *Am. Midland Naturalist* 32:1.

## 18. PROPAGATION AND METAMORPHOSIS: INSECTS

For information on additional species, consult reference 2. Duration of stages varies with season, geographical area, and climate. **Type of Metamorphosis** (column B): C = complete (having internal development of wings until pupal stage); I = incomplete (having external development of wings).

	Species (Common Name)	Type of Meta- mor- phosis	Eggs per Female	Duration of Stage, da			Over- winter- ing Stage	Genera- tions per Season	Reference	
				Egg	Larva or Nymph	Pupa				Adult
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1	<i>Aedes aegypti</i> (yellow-fever mosquito)	C	.....	2-365	6	2-3	15-60	All stages <sup>1</sup>	.....	37
2	<i>Apanteles militaris</i> (parasitic wasp)	C	Hundreds	5-6	10-12	8-10	15-20	Larva in host body	Several	41
3	<i>Apis mellifera</i> (honeybee)	C	.....	3 <sup>2</sup>	8 <sup>2</sup>	9 <sup>2</sup>	35-40 <sup>2</sup>	Adult	See Fn. 3	16,39
4	<i>Bombyx mori</i> (silkworm)	C	300-400	9-12	21-25	14-21	2-3	.....	1-2	3,25,29
5	<i>Cochliomyia hominivorax</i> (screw-worm)	C	100-300	1-2	4-5	5-40	5-30	None	2-12	11
6	<i>Ctenocephalides felis</i> (cat flea)	C	200-400	2-4	8-24	5-7	50-200	Pupa	ca. 10	43
7	<i>Drosophila melanogaster</i> (fruit fly)	C	100	<1	3-11	2-8	14	Larva, adult	5-6	33

<sup>1/1</sup> Insect breeds all year regardless of season. <sup>2/2</sup> Worker bees only. <sup>3/3</sup> Not applicable to individual bee since colony is functioning unit. Queen reproduces and can survive for several years; workers do not reproduce.

*continued*

## 18. PROPAGATION AND METAMORPHOSIS: INSECTS

	Species (Common Name)	Type of Meta- mor- phosis	Eggs per Female	Duration of Stage, da				Over- winter- ing Stage	Genera- tions per Season	Reference
				Egg	Larva or Nymph	Pupa	Adult			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
8	<i>Ephestia kuehniella</i> (Mediterranean flour moth)	C	116-700	>3	40	5-7	3-4	.....	>6	24
9	<i>Heliothis armigera</i> (corn earworm)	C	1,000	2-8	13-28	14	12	Pupa	1-7	4,10,22,29,31,32,34
10	<i>Leptinotarsa decemlineata</i> (Colorado potato beetle)	C	>500	4-9	10-21	5-10	.....	Adult	1-3	9,12
11	<i>Magicicada septendecim</i> (periodical cicada)	I	400-600	42-49	13 or 17 yr	.....	30-40	Nymph	See Fn. 4	18,29,42
12	<i>Melanoplus mexicanus</i> (migratory grasshopper)	I	300-400	90- 120	40-60	.....	>30	Egg	1	4,29,30
13	<i>Musca domestica</i> (housefly)	C	75-200	1-3	4-10	4-18	10-50	All stages <sup>1</sup>	4-18	5,7,13,20,21,23,27,28,44
14	<i>Pediculus</i> sp. (louse)	I	50-300	5-21	7-10	None	10-30	All stages <sup>1</sup>	10-12	8,19
15	<i>Periplaneta americana</i> (American cockroach)	I	200- 1,000	32-58	200-550	.....	371- 441	All stages <sup>1</sup>	1 or less	14,26,29
16	<i>Pieris rapae</i> (imported cabbage-worm)	C	200-500	7	14	7-14	.....	Pupa	3-6	45
17	<i>Popillia japonica</i> (Japanese beetle)	C	40-60	14	275-300	8-20	30-45	Larva	1	6,17,29,36
18	<i>Stagmomantis carolina</i> (Carolina mantis)	I	75-300	210- 300	45-75	.....	20-60	Egg	1	35
19	<i>Tenebrio molitor</i> (yellow mealworm)	C	276	12-16	>600	18-20	60-90	Larva	<1	24
20	<i>Thermobia domestica</i> (firebrat)	I	.....	12-13	60-120	.....	1-2.5 yr	.....	2-5	1,29,40
21	<i>Tribolium confusum</i> (confused flour beetle)	C	300-400	4-14	>22	5-18	1,000	Adult	5-6	15,24,29,38

/1/ Insect breeds all year regardless of season. /4/ 13 or 17 years per generation.

Contributors: (a) Knipling, E. F., and Bishopp, Fred C., (b) Cutkomp, Laurence K., (c) Ritcher, Paul O., (d) Haeussler, G. J., (e) Lindquist, A. W., (f) Oman, Paul W., (g) Porter, B. A.

References: [1] Adams, J. A. 1933. Proc. Iowa Acad. Sci. 40:217. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Anonymous. 1947. Gen. Headquarters Supreme Commander Allied Powers Rept. 76. [4] Baker, W. A. Unpublished. U. S. Dept. of Agriculture, Washington, D. C., 1954. [5] Bishopp, F. C., W. E. Dove, and D. C. Parman. 1915. J. Econ. Entomol. 8:54. [6] Britten, W. E., and J. P. Johnson. 1938. Conn. Agr. Expt. Sta. New Haven Bull. 411. [7] Bucher, G. E., J. W. M. Cameron, and A. S. West, Jr. 1948. Can. J. Res., D. 26:57. [8] Buxton, P. A. 1939. The louse. E. Arnold, London. [9] Chittenden, F. H. 1907. U. S. Bur. Entomol. Circ. 87. [10] Ditman, L. P., and E. N. Cory. 1931. Maryland Agr. Expt. Sta. Bull. 328:443. [11] Dove, W. E., and D. C. Parman. 1935. J. Econ. Entomol. 28:765. [12] Dudley, J. E., Jr., B. J. Landis, and W. A. Shands. 1952. U. S. Dept. Agr. Farmers' Bull. 2040. [13] Feldman-Muhsam, B. 1944. Bull. Entomol. Res. 35:53. [14] Gould, G. E., and H. O. Deay. 1940. Purdue Univ. Agr. Expt. Sta. Bull. 451. [15] Gray, H. E. 1946. Ph.D. Thesis. Univ. Minnesota, Minneapolis. [16] Grout, R. A. 1949. The hive and the honey bee. Dadant, Hamilton, Ill. [17] Hadley, C. H., and I. M. Hawley. 1934. U. S. Dept. Agr. Circ. 332. [18] Haseman, L. 1915. Missouri Univ. Agr. Expt. Sta. Bull. 137. [19] Herms, W. B., and M. T. James. 1961. Medical entomology. Macmillan, New York. [20] Hewitt, C. G. 1914. The housefly. Cambridge Univ. Press, London. [21] Howard, L. O., and F. C. Bishopp. 1924. U. S. Dept. Agr. Farmers' Bull. 1408. [22] Isely, D. 1935. Arkansas Univ. (Fayetteville) Agr. Expt. Sta. Bull. 320. [23] Larsen, E. B., and M. Thomsen. 1940. Dansk Naturhist. Foren. Videnskab. Medd. 104. [24] Latta, R. Unpublished. U. S. Dept. of Agriculture, Washington, D. C. [25] Leggett, W. F. 1949. The story of silk. Lifetime Editions, New York. [26] Mallis, A. 1945. Handbook of pest control.

continued

## 18. PROPAGATION AND METAMORPHOSIS: INSECTS

Univ. California Buildings and Grounds Dept., Los Angeles. [27] Matheson, R. 1950. Medical entomology. Comstock, Ithaca. [28] Matthyse, J. G. 1945. J. Econ. Entomol. 39:743. [29] Metcalf, C. L., W. P. Flint, and R. L. Metcalf. 1951. Destructive and useful insects. McGraw-Hill, New York. [30] Parker, J. R. 1957. U. S. Dept. Agr. Farmers' Bull. 2064. [31] Phillips, W. J., and G. W. Barber. 1929. Virginia Agr. Expt. Sta. Tech. Bull. 40. [32] Phillips, W. J., and M. K. Kenneth. 1923. U. S. Dept. Agr. Farmers' Bull. 1310. [33] Powsner, L. 1955. Physiol. Zool. 8:474. [34] Quaintance, A. L., and C. T. Brues. 1905. U. S. Dept. Agr. Bull. 50. [35] Rau, P., and N. Rau. 1913. Trans. Acad. Sci. St. Louis 22:1. [36] Schread, J. C. 1947. Conn. Agr. Expt. Sta. New Haven Bull. 505. [37] Shannon, R. C., and P. Putnam. 1934. Proc. Entomol. Soc. Wash. 36:185. [38] Shepard, H. H. 1943. Publ. Am. Assoc. Advan. Sci. 20:40. [39] Snodgrass, R. E. 1925. Anatomy and physiology of the honey bee. McGraw-Hill, New York. [40] Sweetman, H. L. 1938. Ecol. Monographs 8:285. [41] Tower, D. G. 1915. J. Agr. Res. 5:495. [42] U. S. Department of Agriculture. 1953. U. S. Dept. Agr. Leaflet 340. [43] U. S. Department of Agriculture. 1955. Ibid. 392. [44] West, L. S. 1951. The housefly. Comstock, Ithaca. [45] Wilson, H. F. 1919. Wisconsin Univ. Agr. Expt. Sta. Res. Bull. 45.

## 19. PROPAGATION AND DEVELOPMENT: INVERTEBRATES

### Part I. METAZOA

**Fertilization** (column C): Ext = external fertilization of egg; Int = internal fertilization of egg. **Sex** (column F): D = dioecious; M = monoecious.

	Phylum and Class	Genus	Fertilization	Zygote	Development	Adult		Reference
						Sex	Form	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Chordata Cephalochordata <sup>1</sup>	<i>Branchiostoma</i>	Ext	Free	Direct	D	Amphioxus	3
2	Thaliacea	<i>Salpa</i>	Int	Placental	Direct	M <sup>2</sup>	Tunicate	3
3	Ascidiacea	<i>Ciona</i>	Ext	Free-floating	Appendicularia larva	M <sup>2</sup>	Sea squirt	3
4	Enteropneusta	<i>Saccoglossus</i>	Ext	Free	Free larva, gradual change to adult	D	Acorn worm	6
5	Echinodermata Ophiuroidea	<i>Ophiopholis</i>	Ext	Free	Dipleurula → ophiopluteus	D	Brittle star	9
6	Asteroidea	<i>Asterias</i>	Ext	Free	Dipleurula → bipinnaria → brachiolaria	D	Starfish	1
7	Echinoidea	<i>Arbacia</i>	Ext	Free	Dipleurula → echinopluteus	D	Sea urchin	1
8	Holothuroidea	<i>Cucumaria</i>	Ext	Free	Dipleurula → modified auricularia	D <sup>3</sup>	Sea cucumber	1
9	Crinoidea	<i>Antedon</i>	Ext	Attached to pinnules	Dipleurula → ciliated larva → stalked crinoid	D	Feather star	3
10	Arthropoda Pycnogonida	<i>Nymphon</i>	Ext	Carried by male	Direct	D	Sea spider	3
11	Arachnida	<i>Centrurus</i>	Int	In female	Direct	D	Scorpion	3
12		<i>Ixodes</i>	Int	In sticky secretion	Larva (nymphlike)	D	Tick	3
13		<i>Pardosa</i>	Int	In cocoon	Direct	D	Spider	3
14	Merostomata	<i>Limulus</i>	Ext	In beach nests	Trilobite larva	D	King crab	3
15	Crustacea	<i>Cambarus</i>	Int	Fastened to swimmerets	Direct	D	Crayfish	14
16		<i>Eubranchipus</i>	Int	In shell	Metanauplius larva	D	Fairy shrimp	1
17	Insecta	<i>Apis</i>	Int	Laid in hive	Larva → pupa	D	Honeybee	3
18		<i>Ephemera</i>	Int	Laid in water	Aquatic nymph	D	Mayfly	3
19		<i>Melolontha</i>	Int	Laid in ground	Grub → pupa	D	June beetle	3
20		<i>Pieris</i>	Int	Laid on plants	Caterpillar → pupa	D	Cabbage butterfly	3
21		<i>Romalea</i>	Int	Laid in ground	Nymph stages	D	Grasshopper	11

<sup>1</sup>/ Subphylum. <sup>2</sup>/ Protogynous. <sup>3</sup>/ Also reproduce asexually.

continued



## 19. PROPAGATION AND DEVELOPMENT: INVERTEBRATES

### Part I. METAZOA

	Phylum and Class	Genus	Fertilization	Zygote	Development	Adult		Reference
						Sex	Form	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
22	Arthropoda Chilopoda	<i>Lithobius</i>	Int	Laid in ground	Direct	D	Centipede	3
23	Diplopoda	<i>Julus</i>	Int	Laid in ground	Direct	D	Millipede	3
24	Onychophora	<i>Peripatus</i>	Int	In parent	Direct	D	Peripatus	1
	Annelida							
25	Hirudinea	<i>Hirudo</i>	Int	In capsule	Direct	M <sup>4</sup>	Leech	1
26	Oligochaeta	<i>Lumbricus</i>	Int	In capsule	Direct	M <sup>4</sup>	Earthworm	1
27	Polychaeta	<i>Nereis</i>	Ext	Free	Trochophore larva	D	Clam worm	5
	Mollusca							
28	Cephalopoda	<i>Loligo</i>	Int	Encased in sticky secretion	Direct	D	Squid	3
29	Bivalvia	<i>Anodonta</i>	Int	In gills of parent	Glochidium parasitic on fish gill	M	Mussel	1
30		<i>Mercenaria</i>	Ext	Free	Trochophore larva → veliger larva → pediveliger larva	D	Quahog	4, 12
31	Gastropoda	<i>Buccinum</i>	Int	In capsule	Trochophore larva → veliger larva	D	Whelk	1
32		<i>Helix</i>	Int	In ground	Direct	M <sup>4</sup>	Land snail	3
33	Aplousophora	<i>Neomenia</i>	Ext	Free	Trochophore larva	M	Solenogaster	11
34	Polyplacophora	<i>Ischnochiton</i>	Ext	Free	Trochophore larva	D	Chiton	3
35	Brachiopoda	<i>Lingula</i>	Ext	Free	Trochophore larva	D	Brachiopod	1
36	Phoronida	<i>Phoronis</i>	Ext or in coelom	Attached to adult tentacles	Actinotrocha larva	M	Phoronid	11
	Polyzoa							
37	Gymnolaemata	<i>Bugula</i>	Int	In body of parent	Trochophore larva	M <sup>5</sup>	Colony	1
38	Phylactolaemata	<i>Pectinatella</i>	Int	In body of parent	Ciliated hollow larva gemmates	M <sup>5</sup>	Colony	1
39	Acanthocephala	<i>Macracanthorhynchus</i>	Int	In capsule	Acanthor → acanthella in beetle larva	D	Thorny-headed worm (in swine)	7, 8
	Aschelminthes							
40	Nematoda	<i>Ascaris</i>	Int	In shell	To juvenile stage in open; completion in host	D	Large roundworm (in mammals)	7, 8
41	Nematomorpha	<i>Gordius</i>	Int	Laid in strings	Larva free, then invades arthropod and develops to juvenile	D	Horsehair worm	7, 8
42	Rotifera	<i>Philodina</i>	None	None	Direct	F <sup>5</sup>	Rotifer	7, 8
	Nemertina							
43	Anopla	<i>Cerebratulus</i>	Ext	Free	Coeloblastula → pilidium	D <sup>5</sup>	Ribbon worm	7, 8
	Platyhelminthes							
44	Cestoda	<i>Taenia</i>	Int	In capsule	Oncosphere → hexacanth → cysticercus (all development in mammals)	M <sup>5</sup>	Tapeworm (in mammals)	13
45	Trematoda	<i>Fasciola</i>	Int	In capsule	Miracidium → sporocyst → rediae → cercariae <sup>7</sup>	M <sup>4</sup>	Liver fluke	7, 8
46	Turbellaria	<i>Dugesia</i>	Int	In capsule	Direct	M <sup>4</sup>	Planarian	7, 8
	Ctenophora							
47	Nuda	<i>Beroe</i>	Ext	Free	Cydippid larva	M	Comb jelly	7, 8
	Cnidaria							
48	Anthozoa	<i>Metridium</i>	Ext	Free	Planula	D <sup>3</sup>	Sea anemone	3
49	Scyphozoa	<i>Aurelia</i>	Int	In folds of oral lobes	Planula → scyphistoma → ephyrae	D	Scyphomedusa	3
50	Hydrozoa	<i>Obelia</i>	Ext	Free	Planula → colony → medusa buds	D	Marine hydra	3
	Porifera							
51	Calcarea	<i>Scypha</i>	Int	In mesenchyme	Amphiblastula	M	Calcareous sponge	2

/3/ Also reproduce asexually. /4/ Cross-fertilization. /5/ Self-fertilization. /6/ Parthenogenetic female. /7/ Miracidia are free; sporocysts, rediae, and cercariae develop in snails; cercariae leave snails and are picked up by ruminant from water or grass.

continued



## 19. PROPAGATION AND DEVELOPMENT: INVERTEBRATES

### Part I. METAZOA

	Phylum and Class	Genus	Fertilization	Zygote	Development	Adult		Reference
						Sex	Form	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
52	Mesozoa	<i>Dicyema</i>	Int	In body	Infusoriform larva→unknown stage→nematogen→rhombogen	M	Infusorigen	10

Contributors: (a) Brown, Relis B., (b) Lochhead, John H., (c) Carriker, Melbourne R., (d) Shuster, Carl N., Jr.

References: [1] Borradaile, L. A., and G. A. Kerkut. 1958. The invertebrata. Ed. 3. Cambridge Univ. Press, London. [2] Brien, P. 1943. Bull. Musee Roy. Hist. Nat. Belg. (Brussels) 19(16):1. [3] Bullough, W. S. 1951. Practical invertebrate anatomy. Macmillan, New York. [4] Carriker, M. R. 1961. J. Elisha Mitchell Sci. Soc. 77(2):168. [5] Dales, R. P. 1951. J. Marine Biol. Assoc. U. K. 29:321. [6] Dawydoff, C. 1948. In P.-P. Grassé, ed. Traité de zoologie. G. Masson, Paris. v. 11, p. 367. [7] Hyman, L. H. 1940. The invertebrates. McGraw-Hill, New York. v. 1. [8] Hyman, L. H. 1951. Ibid. v. 2, 3. [9] Hyman, L. H. 1955. Ibid. v. 4. [10] McConnaughey, B. H. 1951. Univ. Calif. (Berkeley) Publ. Zool. 55(4):295. [11] Parker, T. J., and W. A. Haswell. 1940. A textbook of ecology. Ed. 6. Macmillan, New York. v. 1. [12] Pierce, M. E. 1950. In F. A. Brown, Jr., ed. Selected invertebrate types. J. Wiley, New York. p. 318. [13] Wardle, R. A., and J. A. McLeod. 1952. The zoology of tapeworms. Univ. Minnesota Press, Minneapolis. [14] Wolcott, R. H. 1946. Animal biology. Ed. 3. McGraw-Hill, New York.

### Part II. PROTOZOA

	Class and Genus	Gametocytes	Gametes	Zygote	Development	Adult Form	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
	Ciliata						
1	<i>Paramecium</i>	Conjugating pair	Pieces of micronuclei		Binary fission; each conjugant divides into four	Ciliate	3,4
2	<i>Podophrya</i>	Conjugating pair	Pieces of micronuclei		Budding of ciliated larva; each conjugant divides into four	Suctorian	2-4
	Sporozoa						
3	<i>Eimeria</i>	Megagametocyte, microgametocyte	One egg, many sperm	Encysted	Multiple fission; forms 4 smaller cysts, each with 2 sporozoites	Trophozoite	1,3,4
4	<i>Monocystis</i>	Megagametocyte, microgametocyte	Multiple fission into eggs and sperm	Encysted	Multiple fission into 8 sporozoites	Trophozoite	1,3,4
5	<i>Plasmodium</i>	Megagametocyte, microgametocyte	One egg, several sperm	Ookinete	Multiple fission into many sporozoites	Trophozoite	1,3,4
	Rhizopoda						
6	<i>Amoeba</i>				Binary fission	Amoeba	3,4
7	<i>Endamoeba</i>				Binary fission	Endamoeba	3,4
8	<i>Patellina</i> <sup>1</sup>	Meiosis and mitosis into 16 gamonts	Each gamont forms 8 gametes		Two nuclear divisions without cell division	Schizont	5
	Mastigophora						
9	<i>Astasia</i>				Binary fission	Flagellate	3,4
10	<i>Trypanosoma</i>				Binary and multiple fission	<i>Trypanosoma</i> in vertebrates; <i>Leishmania</i> in invertebrates	3,4

<sup>1</sup>/ Foraminifer.

continued

## 19. PROPAGATION AND DEVELOPMENT: INVERTEBRATES

### Part II. PROTOZOA

	Class and Genus	Gametocytes	Gametes	Zygote	Development	Adult Form	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
11	Mastigophora <i>Volvox</i>	Megagametocyte, microgametocyte	Egg, sperm	Encysted	Budding, forming daughter colonies; a sphere of flagellated cells	Colony	3,4

Contributor: Brown, Relis B.

References: [1] Bullough, W. S. 1951. Practical invertebrate anatomy. Macmillan, New York. [2] Burbanck, W. D. 1950. In F. A. Brown, Jr., ed. Selected invertebrate types. J. Wiley, New York, p. 72. [3] Hyman, L. H. 1940. The invertebrates. McGraw-Hill, New York, v. 1. [4] Hyman, L. H. 1951. Ibid. v. 2, 3. [5] Le Calvez, J. 1950. Zool. Exptl. Gen. 87:211.

## 20. BREEDING SYSTEMS: ANGIOSPERMS

For information on additional species, consult reference 1. Species are those of economic importance. The systems listed indicate the usual breeding classification for a species; where variability exists within a species, only the predominant system (enclosed in parentheses) is given. **System** (column C): SC-S = self-compatible (predominantly self-fertilized), with no inbreeding degeneration; SC-M = self-compatible, monoecious (staminate and pistillate flowers borne on same plant), but rarely self-fertilized under conditions of open pollination; SC-O = self-compatible, intermediate (between SC-S and SC-M), with perhaps a predominance of outcrossing; SI = self-incompatible (sterile to own pollen); D = dioecious (staminate and pistillate flowers borne on separate plants). All data are from reference 2, unless otherwise indicated.

	Species	Common Name	System		Species	Common Name	System
	(A)	(B)	(C)		(A)	(B)	(C)
Monocotyledoneae							
1	<i>Allium cepa</i>	Garden onion	(SC-O) <sup>1</sup>	27	<i>Fragaria virginiana</i> <sup>2</sup>	Virginia strawberry	D*
2	<i>Asparagus officinalis</i> <sup>2</sup>	Garden asparagus	D	28	<i>Fraxinus</i> spp.	Ash	SC-O
3	<i>Avena sativa</i>	Common oat	SC-S	29	<i>Glycine soja</i>	Soybean	(SC-S)
4	<i>Hordeum vulgare</i>	Barley	SC-S	30	<i>Gossypium</i> spp.	Cotton	SC-S
5	<i>Iris</i> spp.	Iris	SC-O	31	<i>Helianthus annuus</i>	Common sunflower	SI
6	<i>Lilium regale</i>	Regal lily	(SC-O)	32	<i>Ilex</i> spp. <sup>3</sup>	Holly	D
7	<i>Oryza sativa</i>	Rice	SC-S	33	<i>Ipomoea batatas</i>	Sweet potato	SI
8	<i>Phleum pratense</i>	Timothy	SI	34	<i>Juglans</i> spp.	Walnut	SC-M
9	<i>Phoenix dactylifera</i> <sup>2</sup>	Date palm	D	35	<i>Lactuca sativa</i>	Lettuce	SC-S
10	<i>Triticum</i> spp.	Wheat	SC-S	36	<i>Lycopersicon esculentum</i>	Tomato	SC-O
11	<i>Yucca</i> spp.	Yucca	SI	37	<i>Malus pumila</i>	Common apple	SI
12	<i>Zea mays</i>	Corn	SC-M	38	<i>Medicago sativa</i>	Alfalfa	SI
Dicotyledoneae							
13	<i>Acer</i> spp.	Maple	SC-O	39	<i>Nicotiana tabacum</i>	Common tobacco	SC-O
14	<i>Alnus</i> spp.	Alder	SC-M	40	<i>Pastinaca sativa</i>	Parsnip	SC-O
15	<i>Beta vulgaris</i>	Beet	SI	41	<i>Persea americana</i>	American avocado	SC-O
16	<i>Betula</i> spp.	Birch	SC-M	42	<i>Phaseolus vulgaris</i>	Kidney bean	SC-S
17	<i>Capsicum frutescens</i>	Bush red pepper	SC-O	43	<i>Pisum sativum</i>	Garden pea	SC-S
18	<i>Carya</i> spp.	Hickory	SC-M	44	<i>Populus</i> spp. <sup>3</sup>	Poplar	D
19	<i>Catalpa speciosa</i>	Northern catalpa	SC-O	45	<i>Prunus domestica</i>	Garden plum	(SI)
20	<i>Cinchona</i> spp.	Cinchona	SI	46	<i>P. persica</i>	Peach	SC-O
21	<i>Citrus</i> spp.	Citrus	(SC-O) <sup>1</sup>	47	<i>Pyrus communis</i>	Pear	SI
22	Cucurbitaceae species	Cultivated species of gourds	SC-M	48	<i>Quercus</i> spp.	Oak	SC-M
23	<i>Daucus carota</i>	Carrot	SC-O	49	<i>Raphanus sativus</i>	Garden radish	SI
24	<i>Digitalis purpurea</i>	Common foxglove	SC-O	50	<i>Ribes</i> spp.	Currant	SC-O
25	<i>Fagopyrum esculentum</i>	Buckwheat	SI <sup>3</sup>	51	<i>Rosa</i> spp.	Roses, most species	SC-O
26	<i>Fagus</i> spp.	Beech	SC-M	52	<i>Salix</i> spp. <sup>3</sup>	Willow	D
				53	<i>Solanum tuberosum</i>	Potato	SC-O
				54	<i>Trifolium hybridum</i>	Alsike clover	SI
				55	<i>Ulmus</i> spp.	Elm	SC-M
				56	<i>Vicia faba</i>	Broad bean	SC-O
				57	<i>Vitis vinifera</i>	European grape	SC-O

/1/ Some are apomictic (reproduction without fertilization). /2/ Information from reference 3. /3/ Heterostyled, i.e., stigma and stamens inserted at different levels. /4/ Cultivated forms are selected intersexes.

continued

**20. BREEDING SYSTEMS: ANGIOSPERMS**

Contributor: Bateman, Angus J.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] East, E. M. 1940. Proc. Am. Phil. Soc. 82:449. [3] Yampolsky, C., and H. Yampolsky. 1922. Bibliotheca Genet. 3:1.

**21. PROPAGATION METHODS: CULTIVATED PLANTS**

For information on additional species, consult reference 3. The methods listed for a genus are those most widely used in cultivation, but not all species of the genus can be propagated by each method. Horticultural varieties are not propagated by seed, as the new plants from seed may vary considerably from the parent plant. When propagation by seed is employed, the seed of species having no apparent rest period may be sown in the spring, while the seed of species having a definite rest period should be artificially stratified or sown in the autumn. Preferred Time (column C): spr = spring; sum = summer; aut = autumn; win = winter.

Species (Common Name)	Method	Preferred Time	Refer- ence	Species (Common Name)	Method	Preferred Time	Refer- ence
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
<b>Gymnospermae</b>				<b>Angiospermae (Dicotyledoneae)</b>			
1 <i>Abies</i> spp. (fir)	Seed	Aut or spr	19,20,48	35 <i>Iris germanica</i> (German iris)	Rhizome di- vision	Aut	22
2	Veneer graft	Win	31,32	36 <i>Lilium</i> spp. (lily)	Seed	Aut	5
3 <i>Cupressus</i> spp. (cypress)	Seed	Aut or spr	32,39,48	37	Scales	Sum	22
4	Cutting <sup>1</sup>	Sum	32,39,48	38	Stem bulbils	Sum	20
5	Veneer graft	Sum	32,39,48	39 <i>Phoenix dactyli- fera</i> (date palm)	Shoots	Spr	22
6 <i>Ginkgo biloba</i> (ginkgo)	Seed	Spr	48	<b>Angiospermae (Dicotyledoneae)</b>			
7	Cutting <sup>1</sup>	Spr	41	40 <i>Acer</i> spp. (maple)	Seed	Aut-spr	31,32,39
8	Whip graft	Early spr	32	41	Cutting <sup>2</sup>	Win	32,35
9	Air layering	Spr	44,45	42	Shield bud	Late sum	37
10 <i>Juniperus</i> spp. (juniper)	Seed	Spr or aut	32,48	43	Side graft	Win	24,32
11	Cutting <sup>2</sup>	Late win	32,48	44	Air layering	Spr	44,45
12	Cutting <sup>1</sup>	Late sum	32,48	45 <i>Alnus</i> spp. (alder)	Seed	Aut	32,39
13	Veneer or side graft	Win	32,48	46	Whip graft	Win	32
14	Simple layer- ing	Spr	48	47 <i>Betula</i> spp. (birch)	Seed	Spr or aut	21,32,39
15 <i>Larix</i> spp. (larch)	Seed	Aut or spr	32,39,48	48	Cleft graft	Early spr	17,32
16	Whip graft	Spr	32	49	Air layering	Spr	44,45
17 <i>Picea</i> spp. (spruce)	Seed	Aut or spr	32,39,48	50 <i>Carya</i> spp. (hick- ory)	Veneer graft	Win-spr	30
18	Cutting <sup>1</sup>	Sum	32	51 <i>C. illinoensis</i> (pe- can)	Patch bud	Spr	1
19	Veneer or side graft	Aut-win	24	52 <i>Catalpa</i> spp. (ca- talpa)	Seed	Spr	16,39
20 <i>Pinus</i> spp. (pine)	Seed	Aut or spr	26	53 <i>Chrysanthemum</i> spp. (chrysan- themum)	Cutting <sup>3</sup>	Yr-round	25
21	Veneer graft	Spr	26	54	Rough divi- sion	Aut	22
22	Air layering	Spr-aut	26	55 <i>Citrus hybrida</i> (citrus)	Inverted T- bud	Spr-sum	42
23 <i>Taxus</i> spp. (yew)	Seed	Aut or spr	28,32	56 <i>Cornus</i> spp. (dog- wood)	Seed	Aut or spr	13,32,39
24	Cutting <sup>2</sup>	Aut-win	9,32,48	57	Cutting <sup>3</sup>	Sum-aut	17,32
25	Cutting <sup>1</sup>	Sum	9,32,48	58	Simple layer- ing	Spr	32
26	Air layering	Aut or spr	44,45	59	Air layering	Spr	44,45
27 <i>Thuja</i> spp. (ar- borvitae)	Seed	Aut or spr	4,32	60 <i>Fagus</i> spp. (beech)	Seed	Aut or spr	32
28	Cutting <sup>2</sup>	Aut-win	32,48	61	Whip or cleft graft	Early spr	32
29	Cutting <sup>1</sup>	Sum	32,48	62 <i>Fragaria</i> spp. (strawberry)	Runners	Spr	12
30	Veneer or side graft	Aut-win	20	63 <i>Fraxinus</i> spp. (ash)	Seed	Aut or spr	32,36,39
31 <i>Tsuga</i> spp. (hem- lock)	Seed	Aut or spr	32,39,48	64 <i>Ilex</i> spp. (holly)	Cutting <sup>1</sup>	Sum	22,48
32	Cutting <sup>1</sup>	Sum	32	65	Shield bud	Spr	32
33	Air layering	Early spr	44,45	<b>Angiospermae (Monocotyledoneae)</b>			
34 <i>Gladiolus hortu- lanus</i> (horticul- tural gladiolus)	Cormels	Aut-spr	15				

1/ Semihardwood. 2/ Hardwood. 3/ Softwood

continued



**21. PROPAGATION METHODS: CULTIVATED PLANTS**

Species (Common Name)	Method	Preferred Time	Refer- ence	Species (Common Name)	Method	Preferred Time	Refer- ence		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
Angiospermae (Dicotyledoneae)				85	<i>Pyrus communis</i>	Shield bud	Sum	14	
66	<i>Ilex</i> spp. (holly)	Simple layering	Spr-sum	32	86	<i>Pyrus communis</i> (pear)	Whip graft	Win	38
67		Air layering	Spr	44,45	87	<i>Quercus</i> spp. (oak)	Seed	Aut or spr	18,32,39
68	<i>Juglans</i> spp. (walnut)	Veneer graft	Early spr	10	88		Side graft	Early spr	32,48
69		Patch bud	Spr-sum	10	89	<i>Rhododendron</i>	Seed	Spr	27,39,49
70	<i>Magnolia</i> spp. (magnolia)	Seed	Spr or aut	2,32,39	90	spp. (rhododendron)	Cutting <sup>1</sup>	Mid-sum	7,33,49
71		Cutting <sup>1</sup>	Aut	32	91		Veneer graft	Win	23,32,49
72		Side graft	Sum	6,32	92		Simple layering	Spr-sum	17,32,49
73		Air layering	Spr	44,45	93		Air layering	Spr-sum	11
74	<i>Malus pumila</i> (common apple)	Shield bud	Spr-sum	1	94	<i>Ribes sativum</i>	Cutting <sup>2</sup>	Aut-win	40
75	<i>Persea americana</i> (American avocado)	Patch bud	Spr	1	95	(common red currant)	Mound layering	Sum	1
76		Whip graft	Win	1	96	<i>Rosa</i> spp. (rose)	Cutting <sup>1</sup>	Late spr	8,32,46,47
77	<i>Populus</i> spp. (poplar)	Cutting <sup>2</sup>	Aut	32	97		Shield bud	Spr-sum	29,38,46,47
78		Root graft	Early win	32	98		Air layering	Spr-sum	44,45
79		Air layering	Early spr	44,45	99	<i>Ulmus</i> spp. (elm)	Seed	Aut or spr	32,39
80	<i>Prunus amygdalus</i> (almond)	Shield bud	Sum	1	100		Shoots	Early spr	17,32
81	<i>P. domestica</i> (garden plum)	Seed	Spr	1	101		Whip graft	Aut	17,32
82		Shield bud	Spr-sum	17	102	<i>Vitis</i> spp. (grape)	Cutting <sup>2</sup>	Win	1,43
83		Top graft	Win	1	103		Whip graft	Late win	1
84	<i>P. persica</i> (peach)	Shield bud	June-sum	17	104		Chip bud	Spr	34

/1/ Semihardwood. /2/ Hardwood.

Contributors: (a) Mahlstedt, John P., (b) Wyman, Donald, (c) Brase, Karl D., (d) Slate, George L.

References: [1] Adriance, G. W., and F. R. Brison. 1939. Propagation of horticultural plants. McGraw-Hill, New York. [2] Afanasiev, M. 1937. N. Y. State Agr. Expt. Sta. (Ithaca) Mem. 208. [3] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [4] Barton, L. V. 1930. Am. J. Botany 17:88. [5] Barton, L. V. 1936. Contrib. Boyce Thompson Inst. 8:297. [6] Beilman, A. P. 1932. Bull. Missouri Botan. Garden 20. [7] Bridgers, B. T. 1952. Quart. Bull. Am. Rhododendron Soc. 6:184. [8] Buck, G. J. Unpublished. Iowa State College Dept. of Horticulture, Ames, 1951. [9] Chadwick, L. C. 1933. N. Y. State Agr. Expt. Sta. (Ithaca) Bull. 571. [10] Chase, S. 1947. Proc. Am. Soc. Hort. Sci. 49:175. [11] Creech, J. 1950. Natl. Hort. Mag. 29:114. [12] Darrow, G. M. 1929. U. S. Dept. Agr. Tech. Bull. 122. [13] Davis, O. H. 1926. Florists Exchange Hort. Trade World 63:917. [14] Day, L. H. 1947. Calif. Univ. Agr. Expt. Sta. Bull. 700. [15] Denny, F. E. 1930. Contrib. Boyce Thompson Inst. 2:523. [16] Engstrom, H. E., and J. H. Stoeckeler. 1941. U. S. Dept. Agr. Misc. Publ. 434. [17] Gardner, F. E. 1932. Maryland Agr. Expt. Sta. Bull. 335. [18] Gardner, R. C. B. 1937. Quart. J. Forestry 31:32. [19] Griffiths, D. 1920. Flower Grower 7(12):199. [20] Hottes, A. C. 1922. Practical plant propagation. A. T. De La Mare, New York. [21] Joseph, H. C. 1929. Botan. Gaz. 87:127. [22] Kains, M. G., and L. M. McQuesten. 1949. Propagation of plants. Orange Judd, New York. [23] Mallinson, J. W. 1926. Florists Exchange Hort. Trade World 61:749. [24] Mallinson, J. W. 1926. Ibid. 61:1139. [25] Maxon, M. A., B. S. Pickett, and H. W. Richey. 1940. Iowa Agr. Expt. Sta. Bull. 280. [26] Mergen, F., and H. Rossoll. 1954. South-eastern Forest Expt. Sta. Paper 46. [27] Morrison, B. Y. 1929. U. S. Dept. Agr. Circ. 68. [28] Nichols, G. E. 1924. Ecology 15:364. [29] Post, K. C. 1950. Florist crop production and marketing. Orange Judd, New York. [30] Reed, C. A. 1926. U. S. Dept. Agr. Farmers' Bull. 1501. [31] Roe, E. T. 1941. J. Forestry 39:413. [32] Sheat, W. G. 1948. Propagation of trees, shrubs and conifers. Macmillan, London. [33] Skinner, H. T. 1937. Proc. Am. Soc. Hort. Sci. 35:830. [34] Slate, G. L., and K. D. Brase. Unpublished. Cornell Univ., Geneva, 1953. [35] Snow, A. G. 1941. J. Forestry 39:395. [36] Steinbauer, G. P. 1937. Plant Physiol. 12:813.

continued



## 21. PROPAGATION METHODS: CULTIVATED PLANTS

[37] Stewart, L. B. 1924. Trans. Proc. Botan. Soc. Edinburgh 29:43. [38] Tukey, H. B., and K. D. Brase. 1934. N. Y. State Agr. Expt. Sta. (Ithaca) Bull. 649. [39] U. S. Forest Service. 1948. U. S. Dept. Agr. Misc. Publ. 654. [40] Van der Lek, H. A. A. 1930. Rept. Proc. Intern. Hort. Congr., 9th, p. 66. [41] Wardin, R. W. Unpublished. Iowa State Teachers College, Cedar Falls, 1953. [42] Webber, H. J. 1920. Calif. Univ. Agr. Expt. Sta. Bull. 317. [43] Winkler, A. J. 1927. Hilgardia 2:230. [44] Wyman, D. 1951. Arnoldia 11(7). [45] Wyman, D. 1951. Ibid. 11(8). [46] Yerkes, G. E. 1928. Propagation of roses. U. S. Dept. of Agriculture, Bureau of Plant Industry, Washington, D. C. [47] Yerkes, G. E. 1939. Rose propagation by cuttings. U. S. Dept. of Agriculture, Bureau of Plant Industry, Washington, D. C. [48] Yerkes, G. E. 1945. U. S. Dept. Agr. Farmers' Bull. 1657. [49] Yerkes, G. E., and B. Y. Morrison. 1925. Propagation of rhododendrons and azaleas. U. S. Dept. of Agriculture, Bureau of Plant Industry, Washington, D. C.

## 22. SEED GERMINATION: HERBACEOUS PLANTS

For information on additional species, consult reference 1.

	Species	Common Name	Substrate	Germination		Special Requirements
				Temp. <sup>1</sup> °C	Time <sup>2</sup> da	
	(A)	(B)	(C)	(D)	(E)	(F)
<b>Monocotyledoneae</b>						
1	<i>Allium cepa</i>	Garden onion	Between blotters or toweling	20	6-10	None
2	<i>Asparagus officinalis</i>	Garden asparagus	Between blotters; between folded paper toweling; soil or sand	20-30	7-21	None
3	<i>Avena sativa</i>	Common oat	Between folded paper toweling; soil or sand	15	5-10	Prechill at 5° or 10°C for 5 days
4	<i>Hordeum vulgare</i>	Barley	Between folded paper toweling; soil or sand	15	4-7	Prechill at 5° or 10°C for 5 days
5	<i>Oryza sativa</i>	Rice	Between blotters; between folded paper toweling; soil or sand	20-30	5-14	None
6	<i>Phleum pratense</i>	Timothy	Closed petri dish with cotton, blotter, or filter paper; top of blotters	20-30	5-10	Light; prechill at 5° or 10°C for 5 days
7	<i>Poa pratensis</i>	Kentucky bluegrass	Closed petri dish with cotton, blotter, or filter paper	10-30	10-28	Light; 0.1% KNO <sub>3</sub> solution; prechill dormant seeds at 10°C for 5 days
8	<i>Triticum</i> spp.	Wheat	Between folded paper toweling; soil or sand	15	4-7	Prechill at 5° or 10° C for 5 days
9	<i>Zea mays</i>	Corn	Rolled towel; soil or sand	25	4-7	None
<b>Dicotyledoneae</b>						
10	<i>Antirrhinum</i> spp.	Snapdragon	Closed petri dish with cotton, blotter, or filter paper	20-30	5-12	Light; fresh and hybrid seed may require prechilling at 3° or 5°C for 10-20 days
11	<i>Beta vulgaris</i>	Beet	Between blotters; soil or sand	20-30	3-14	Soak in water for 2 hours; rinse, blot surface dry
12	<i>Capsicum</i> spp.	Pepper	Top of blotters	20-30	6-14	Light; 0.2% KNO <sub>3</sub> solution
13	<i>Chrysanthemum maximum</i>	Pyrenees chrysanthemum	Top of blotters	20-30	8	Light
14	<i>Cucumis sativus</i>	Cucumber	Between folded paper toweling; soil or sand; between blotters	20-30	3-7	None
15	<i>Cucurbita</i> spp.	Gourd	Between folded paper toweling; soil or sand	20-30	4-7	Keep substrate drier than for other seeds
16	<i>Daucus carota</i>	Carrot	Between blotters	20-30	6-21	None
17	<i>Fagopyrum esculentum</i>	Buckwheat	Between blotters; between folded paper toweling	20-30	3-6	None

1/ Ranges indicate that a daily fluctuating temperature is preferred for germination: 16 hours at the lower temperature and 8 hours at the higher temperature. 2/ Maximum germination is usually obtained during the given time limit; for hard-coated seeds an additional 5 days is recommended.

*continued*

## 22. SEED GERMINATION: HERBACEOUS PLANTS

	Species	Common Name	Substrate	Germination		Special Requirements
				Temp. <sup>1</sup> °C	Time <sup>2</sup> da	
	(A)	(B)	(C)	(D)	(E)	(F)
Dicotyledoneae						
18	<i>Glycine soja</i>	Soybean	Rolled towel; soil or sand	20-30	5-8	None
19	<i>Gossypium</i> spp.	Cotton	Rolled towel; soil or sand	20-30	4-12	None
20	<i>Helianthus annuus</i>	Common sunflower	Between folded paper toweling; between blotters	20-30	3-7	None
21	<i>Lactuca sativa</i>	Lettuce	Closed petri dish with cotton, blotter or filter paper	20	7	Light; prechill at 10°C for 3 days, or test at 15°C
22	<i>Lycopersicon esculentum</i>	Tomato	Between blotters or toweling; closed petri dish with cotton, blotter, or filter paper	20-30	5-14	Dormant seed: light; 0.2% KNO <sub>3</sub> solution
23	<i>Medicago sativa</i>	Alfalfa	Between blotters; soil or sand	20	4-7	None
24	<i>Nicotiana tabacum</i>	Tobacco	Closed petri dish with cotton, blotter, or filter paper; top of blotters	20-30	7-14	Light
25	<i>Pastinaca sativa</i>	Parsnip	Between blotters	20-30	6-28	None
26	<i>Phaseolus vulgaris</i>	Kidney bean	Rolled towel; soil or sand	20-30	5-8	None
27	<i>Pisum sativum</i>	Garden pea	Rolled towel; soil or sand	20	5-8	None
28	<i>Raphanus sativus</i>	Garden radish	Between blotters	20	4-6	None
29	<i>Rheum rhabonticum</i>	Garden rhubarb	Top of blotters; top of soil	20-30	7-21	Light
30	<i>Solanum melongena</i>	Eggplant	Closed petri dish with cotton, blotter or filter paper; top of blotters	20-30	7-14	Dormant seed: light; 0.2% KNO <sub>3</sub> solution
31	<i>Trifolium hybridum</i>	Alsike clover	Between blotters; soil or sand	20	3-7	Dormant seed: 15°C
32	<i>Vicia faba</i>	Broad bean	Soil or sand; creped cellulose paper wadding	20	4-14	Prechill at 10°C for 3 days

<sup>1</sup>/ Ranges indicate that a daily fluctuating temperature is preferred for germination: 16 hours at the lower temperature and 8 hours at the higher temperature. <sup>2</sup>/ Maximum germination is usually obtained during the given time limit; for hard-coated seeds an additional 5 days is recommended.

**Contributors:** Justice, O. L., and Rollin, S. F.

**References:** [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Justice, O. L., et al. 1960. Proc. Assoc. Offic. Seed Analysts 49(2):21.

## 23. SEED GERMINATION: FOREST TREES, NORTH AMERICAN

For information on additional species, consult reference 1. **Dormancy** (column C): E = embryo; SC = seed coat. **Storage Method** (column D): D = dry; M = moist; C = in sealed containers. **Pretreatment Method** (column G): P = stratify in moist peat; S = stratify in moist sand; H<sub>2</sub>SO<sub>4</sub> = soak in concentrated sulfuric acid.

	Species (Common Name)	Seed-bearing Age <sup>1</sup> yr	Dor- man- cy <sup>2</sup>	Storage			Pretreatment			Sowing to Full Germination, da		Germination			Ref- er- ence
				Meth- od	°C	Inter- val <sup>3</sup>	Meth- od	°C	Dura- tion	Pre- treated Seed	Un- treated Seed	°C <sup>4</sup>	Field %	Lab %	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)
Gymnospermae															
1	<i>Abies concolor</i> (white fir)	40-200	E <sup>5</sup>	D, C	2-4	>3 yr	S	5	60-90 da	30	>100	20-30	15	34	10
2	<i>Cupressus arizoni- ca</i> (Arizona cy- press)	.....	E?	D, C	5	10 yr	S	5	60 da	30	75	20-30	15	26	10

<sup>1</sup>/ Age of most abundant production. <sup>2</sup>/ Dormancy may be general, variable (dormant and nondormant seeds in same sample), occasional, or rare; type is general unless otherwise indicated. <sup>3</sup>/ Without serious loss in viability. <sup>4</sup>/ Lower limit of range is night temperature, upper limit is day temperature. <sup>5</sup>/ Variable.

*continued*

## 23. SEED GERMINATION: FOREST TREES, NORTH AMERICAN

Species (Common Name)	Seed-bearing Age <sup>1</sup> yr	Dor- man- cy <sup>2</sup>	Storage			Pretreatment			Sowing to Full Germination, da		Germination			Ref- er- ence	
			Meth- od	°C	Inter- val <sup>3</sup>	Meth- od	°C	Dura- tion	Pre- treated Seed	Un- treated Seed	°C <sup>4</sup>	Field %	Lab %		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	
Gymnospermae															
3	<i>Juniperus virginiana</i> (eastern red cedar)	10-175	E+SC	D, C	-7	>2 yr	H <sub>2</sub> SO <sub>4</sub>	....	30 min <sup>6</sup>	20-30	>180	10-25	30	42	2,10
4	<i>Larix occidentalis</i> (western larch)	40-100	E <sup>5</sup>	D, C	5	>2 yr	S	5	30 da	20-30	60	20-30	20	27	3,10
5	<i>Picea glauca</i> (white spruce)	30->100	E	M <sup>7</sup> , C	2-5 <sup>3</sup>	10 yr	S	5	60-90 da <sup>9</sup>	30-45	50-140	20-30	35	50	5,6,8-10
6	<i>Pinus strobus</i> (eastern white pine)	15-250	E	M <sup>10</sup> , C	0-5	>10 yr	Por S	10	30 da <sup>11</sup>	30-40	60-100	20-30	50	64	8-10
7	<i>Sequoia gigantea</i> (giant sequoia)	50->300	E <sup>5</sup>	D, C	-3 to -1	8-24 yr	None	....	.....	.....	40-60	15-20	15	25	9-11
8	<i>Taxodium distichum</i> (bald cypress)	.....	E+SC <sup>7</sup>	D, C	5	>1 yr	Por S	5	30-60 da	30-50	60-110	20-30	8	12	10
9	<i>Thuja occidentalis</i> (northern white cedar)	20->100	E <sup>5</sup>	D, C	2-5	5 yr	S	0-10	30-60 da	30	50	20-30	30	46	10
10	<i>Tsuga canadensis</i> (eastern hemlock)	30->400	E <sup>5</sup>	D, C	2-5	4 yr	S	5	60-120 da	60	200	20-30	20	38	10
Angiospermae (Dicotyledoneae)															
11	<i>Acer saccharinum</i> (silver maple)	35-	None	M, C	5	2 yr	None	....	.....	.....	20-30	25-30	18	76	10
12	<i>Alnus rubra</i> (red alder)	10-100	E?	D, C	0-5	>1 yr	S	5	30-60 da	30-40	60	20-30	14	27	9,10
13	<i>Betula lenta</i> (sweet birch)	40-	E?	D	5-7	18 mo	Por S	0-5	40-70 da	30	90	15-32	15	43	10
14	<i>Carya illinoensis</i> (pecan)	20-300	E	C <sup>12</sup>	5	3->5 yr	Por S	2-7	30-90 da	45-60	200-300	20-30	50	50	10
15	<i>Catalpa speciosa</i> (northern catalpa)	20-	None?	D, C	0-5	>2 yr	None	....	.....	.....	60	20-30	70	75	10
16	<i>Fagus grandifolia</i> (American beech)	40-100	E	D, C	1-5	>6 mo	S	5	90 da	60	150-160	20-30	40	85	10
17	<i>Fraxinus americana</i> (white ash)	20-175	E	D, C	2-5	>3 yr	Por S	5	60-90 da	40-60	>60	20-30	20	38	9,10
18	<i>Juglans nigra</i> (black walnut)	12-130	E+SC <sup>7</sup>	D, C	2-5	>1 yr	Por S	1-10	60-120 da	15-40	100-300	20-30	55	75	4,10
19	<i>Populus tremuloides</i> (quaking aspen)	20->70	None	D, C	5	>1 yr	None	....	.....	.....	7	20-30	50	59	9,10
20	<i>Prunus serotina</i> (black cherry)	10-180	E+SC <sup>7</sup>	C	5	>2 yr	Por S	5	90-120 da	30	>190	15-30	30	63	7,9,10
21	<i>Quercus alba</i> (white oak)	20-300	None	D, C <sup>13</sup>	0-2	>1 yr	None	....	.....	.....	30-50	20-30	66	78	10
22	<i>Ulmus americana</i> (American elm)	15-300	E <sup>5</sup>	D, C	5	>2 yr	S	5	60 da	15-60	90	20-30	15	63	9,10

/1/ Age of most abundant production. /2/ Dormancy may be general, variable (dormant and nondormant seeds in same sample), occasional, or rare; type is general unless otherwise indicated. /3/ Without serious loss in viability. /4/ Lower limit of range is night temperature, upper limit is day temperature. /5/ Variable. /6/ Or stratify in moist soil for 30 days at 25°C, then stratify for 90 days at 5°C. /7/ Moisture content, 5%. /8/ Or -15° to -5°C. /9/ Or soak in water for 14 days at 5°C. /10/ Moisture content, 6%. /11/ Or soak in water for 7-10 days at 5°C. /12/ Relative humidity, 90%. /13/ Relative humidity, 80-90%.

Contributor: Rudolf, Paul O.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Barton, L. V. 1951. Contrib. Boyce Thompson Inst. 16:387. [3] Boe, K. N. 1958. U. S. Dept. Agr. Forest Serv. Intermountain Forest

continued

**23. SEED GERMINATION: FOREST TREES, NORTH AMERICAN**


Range Expt. Sta. Misc. Publ. 16. [4] Brinkman, K. A. 1957. U. S. Dept. Agr. Forest Serv. Central States Forest Expt. Sta. Misc. Release 22. [5] Cram, W. H. 1951. Forestry Chron. 27:349. [6] Crossley, D. I., and L. Skov. 1951. Can. Dept. Resources Develop. Silvicultural Leaflet 59. [7] Hough, A. F. 1960. U. S. Dept. Agr. Forest Serv. Northeastern Forest Expt. Sta. Paper 139. [8] Rudolph, P. O. 1950. J. Forestry 48:31. [9] Rudolph, P. O. Unpublished. U. S. Dept. of Agriculture Forest Service, St. Paul, Minn., 1961. [10] Rudolph, P. O., et al. 1948. U. S. Dept. Agr. Misc. Publ. 654. [11] Schubert, G. H. 1957. U. S. Dept. Agr. Forest Serv. Calif. Forest Range Expt. Sta. Tech. Paper 20.



## III. DEVELOPMENT AND GROWTH

### 24. EARLY PRENATAL DEVELOPMENT: MAN


Diagrams show the development of germ layers and their derivatives, without regard to size relationships between stages.



*Contributors:* (a) Hertig, Arthur T., (b) Patten, Bradley M.

*References:* [1] Hertig, A. T., et al. 1954. *Contrib. Embryol. Carnegie Inst. Wash.* 35:199. [2] Patten, B. M. 1953. *Human embryology*. Ed. 2. Blakiston, New York.

Adapted from B. M. Patten, *Human Embryology*, Blakiston,




/1/ By cleavage divisions. /2/ By hollowing and expansion. /3/ Extraembryonic. /4/ Not present in all mammals. from endoderm, remainder from mesoderm. /5/ Lower part. /10/ Between visceral arches I-IV. /11/ Embryonic of peritoneum. /14/ Embryonic structure (disappears). /15/ Epithelial portion of structures from ectoderm, ectoderm along with neural crest cells. /18/ Probably in part only; remainder of pia, and all of dura and arachnoid, be derived from ectodermal placodes. /20/ Derivation from neural crest still disputed. /21/ Pars anterior, pars from chick only). /24/ Not all from splanchnic mesoderm; some from somatic mesoderm or head mesenchyme.

Contributors: (a) Patten, Bradley M., (b) Reyer, Randall W., (c) Hertig, Arthur T., (d) Arey, Leslie B.

## DERIVATIVES: EUTHERIAN MAMMALS

New York, 1953. Broken line indicates disputed origin.



/5/ In primates only. /6/ In man. /7/ Probably not in primates. /8/ Epithelial and glandular portion of structure; may form nucleus pulposus of intervertebral discs. /12/ Upper part. /13/ Or from splanchnic mesoderm remainder from mesoderm. /18/ In part only. /17/ In rostral areas of head, some mesenchyme comes in from the from mesenchyme. /19/ Evidence from lower vertebrates indicates that a portion of these ganglia and nerves may tuberalis, and pars intermedia of hypophysis. /22/ In region of somites only. /23/ Ventral and lateral part (evidence

## 26. TIME VARIATIONS IN DEVELOPMENTAL STAGES: MAMMALS AND BIRDS

In mammals, hours and days were counted from time of fertilization. In birds, one day was allowed for intrauterine development and added to the incubation age (lines 4-10). Sex differentiation most often occurs at Stage 30 but varies between Stages 28 (rat) and possibly 35 (opossum).

Standard Stages (Witschi)		Identification of Stages	Man	Ham-ster	Monkey, rhesus	Opos-sum	Rabbit	Rat	Sheep	Swine	Chick	Hawk	Spar-row
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	
1	2	2 cells	38 hr	16 hr	24 hr	40 hr	8 hr	24 hr	30 hr	30 hr	3 hr	.....	.....
2	3	4 cells	48 hr	40 hr	36 hr	56 hr	11 hr	48 hr	34 hr	40 hr	3 1/4 hr	.....	.....
3	7-8	Beginning of implantation	6 1/2 da	4 1/2 da	9 da	6 da	7 da	6 da	10 da	.....	.....	.....	.....
4	12	Primitive streak	19 da	6 1/2-7 da	19 da	6 1/2 da	6 1/2 da	8 1/2 da	13 da	11 da	1 1/2 da	.....	1 1/2 da
5	16	13-20 somite embryo	27 da	8 da	25 da	9 da	9 da	10 1/2 da	17 da	16 da	3 da	4 1/2 da	2 1/2 da
6	18	Formation of tail bud	29 da	8 1/2 da	26 da	9 1/2 da	9 1/2 da	11 1/2 da	18 da	17 da	3 1/4 da	.....	3 1/4 da
7	25	End of embryonic period	36 da	9 da	28 da	10 da	10 da	12 1/2 da	21 da	20 da	5 da	9 da	5 da
8	33-34	End of metamorphosis	60 da	13 1/2 da	40 da	12 da	14 da	16 1/2 da	32 da	35 da	8 1/2 da	13 1/2 da	8 da
9	35	Closed eyelids	70 da	.....	48 da	12 1/2 da	19 da	18 da	42 da	50 da	13 da	23 da	11 da
10	36	Open eyelids	140 da	.....	.....	72 da <sup>1</sup>	42 da <sup>2</sup>	38 da <sup>3</sup>	84 da	90 da	21 da	.....	20 da <sup>4</sup>
Birth or hatching													
11	Age		267 da	16 da	164 da	12 1/2 da	32 da	22 da	150 da	112 da	22 da	36 da	14 da
12	Standard stages (Witschi)		36	35	36	35	35+	35	36	36	36	35-36	35
13	Weight		3.2 kg	2.2 g	450 g	0.13 g	57 g	4.5 g	5 kg	2.5 kg	34 g	12 g	1.7 g
14	Weight relative to mother		5.5%	2.3%	7.5%	0.01%	3%	2.25%	8%	2.5%	3%	.....	6%

<sup>1</sup>/ 60 days after birth. <sup>2</sup>/ 10 days after birth. <sup>3</sup>/ 16 days after birth. <sup>4</sup>/ 6 days after hatching.

Contributor: Witschi, Emil

Reference: Witschi, E. 1956. Development of vertebrates. W. B. Saunders, Philadelphia.

## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

For information on development of tissues and organs, consult reference 1, Part I.

### Part I. MAN

**Age** (column C) = fertilization or ovulation age, usually calculated from last menstruation minus 14 days. **Size** (column D) = greatest diameter, or crown-to-rump length (approximate chorionic size is given in parentheses). **Identification** (column E) is for standard stages, and therefore Streeter's horizons are not always comparable to the information given.

Standard Stages (Witschi)	Streeter's Horizons	Age da	Size mm	Identification of Stages	Reference
(A)	(B)	(C)	(D)	(E)	(F)
Cleavage and Blastula					
1	I	1	0.125	1 cell; in the tubes of the oviducts	19,24
2	II	2	0.115	2 cells; in the tubes of the oviducts	7,24
3	II		0.115	4 cells; in the tubes of the oviducts	7,24
4	II	3	0.100	8-12 cells (morula), entering the uterus	7
5	III	4	0.101	Early blastocyst (58 cells), in lumen of uterus	7
6	III	5	0.095	Free blastocyst (107 cells), in lumen of uterus	7
7	III			Blastocyst beginning of implantation	7
Gastrula					
8	IV	7-8	0.05 (0.3)	Bilaminar disc (embryoblast) amniotic cavity	6,28
9	V	9	0.1 (0.5)	Embryonic disc and exoembryonic envelopes; exoembryonic mesoderm	6
10	VI	11-13	0.15 (1.0)	Beginning primitive streak; yolk sac; exocoelom	5,28
11	VII	14-17	0.3 (2.5)	Median primitive streak; syn- and cyto-trophoblast	11,16,28
Primitive Streak					
12	VIII	19	0.7 (8.0)	Complete primitive streak; chorionic villi	15,16,28

*continued*



**27. CHARACTERIZATION OF DEVELOPMENTAL STAGES**

**Part I. MAN**

Standard Stages (Witschi)	Streeter's Horizons	Age da	Size mm	Identification of Stages	Reference
(A)	(B)	(C)	(D)	(E)	(F)
<b>Neurula</b>					
13	IX	20	1.5 (12.0)	Presomite neurula; spreading neural plate	9,22
14	X	21	2 (13)	Occipital somites 1-4; neural folds; invagination canal	10,17,23,25
15	X	24	2.8 (16.0)	Cervical somites 5-12; neural tube starts forming	2,10,16,18,21,22
16	XI	27	3.3 (22.0)	Thoracal somites 13-20; 2 visceral arches; upper and lower neuropores; germ cells start leaving yolk sac	8,26-28
17	XII	28	3.5	Thoracal somites 21-24; 3 visceral arches; oral membrane ruptures	3,14,26
<b>Tail-Bud Embryo</b>					
18	XII	29	3.8 (24.0)	Lumbar somites 25-27; oral membrane resorbed; germ cells in hindgut and ventral mesentery	20,26,27
19	XII	30	4 (25)	Lumbar somites 28,29; appendicular ridges	26,28
20	XIII	31	4.3 (26.0)	Sacral somites 30-32; arm and leg buds appear; germ cell migration reaches borders of mesonephric ridges	26-28
21	XIII	32	4.6 (27.0)	Sacral somites 33,34; 4 visceral arches	26
22	XIII	33	4.8 (28.0)	Caudal somites 35,36; otic vesicles detach	13,26
23	XIII	34	5 (29)	Caudal somites 37; slender yolk stalk	16,26
24	XIII	35	5.4 (30.0)	Caudal somites 38; lens placode; germ cells from hindgut to median mesonephric ridges	26,27
<b>Complete Embryo</b>					
25	XIV	35-37	6 (28-35)	End of somite formation; arm and leg buds fully formed; regression of tail bud; germ cells in genital ridges, end of migration	4,12,26,28
<b>Metamorphosing Embryo</b>					
26	XV	38	8 (32)	Differentiation of hand plate; beginning of umbilical hernia	12,26
27	XVI	40	8-10 (35)	Visceral arches III and IV disappear under cervical fold and operculum; eyes pigmented; yolk sac separates from gut	26
28	XVII	42	12	Pentadactyl rudiments; closing of cervical sinus	26
29	XVII	44	12.5-14.0 (40.0)	Median processes of maxillaries advancing; chorionic villi longer where umbilicus attaches; cartilage formation in vertebrae	20,23,26,28
30	XVIII	46	14.6	Premaxillary processes; beginning sexual differentiation of gonads	20,27,30
31	XVIII	48	15.6	Closing of facial clefts; hands and feet lateral to body wall	26,28
32	XIX	50	17	Phalanges, first links; hands, far apart, bending over heart; first ossification centers in mandible and clavicle	26
33	XX-XXII	56	22-25 (47)	Closed facial clefts; auricles rising; large umbilical hernia; arms and feet growing, fingers from left and right touch nose	26,28
<b>Fetus</b>					
34	XXIII+	56-70	26-45	1st fetal stage: growth of eyelids; gut withdrawal from hernia; palatine raphe; differentiation of male and female external genitalia	12,16,26,28,29
35		70-140	45-180	2nd fetal stage: periderm sealed eyelids; ossification of vertebral column begins; first ovocytes in ovaries; hair follicles; disc placenta	20,28,30
36		140-266	180-340	3rd fetal stage: resorption of periderm; cornification and separation of eyelids; lanugo; uterovaginal differentiation	20,28,29

Contributor: Witschi, Emil

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Corner, G. W. 1929. Contrib. Embryol. Carnegie Inst. Wash. 20:81. [3] Davis, C. L. 1923. Ibid. 15:1. [4] Hamilton, W. J.,

*continued*

**27. CHARACTERIZATION OF DEVELOPMENTAL STAGES**

**Part I. MAN**

J. D. Boyd, and H. W. Mossman. 1952. Human embryology. Ed. 2. Williams and Wilkins, Baltimore. p. 87.

[5] Hertig, A. T., and J. Rock. 1941. Contrib. Embryol. Carnegie Inst. Wash. 29:127. [6] Hertig, A. T., and J. Rock. 1945. Ibid. 31:65. [7] Hertig, A. T., et al. 1954. Ibid. 35:199. [8] Heuser, C. H. 1930. Ibid. 22:135. [9] Heuser, C. H. 1932. Ibid. 23:251. [10] Heuser, C. H., and G. W. Corner. 1957. Ibid. 36:29. [11] Heuser, C. H., J. Rock, and A. T. Hertig. 1945. Ibid. 31:85. [12] His, W. 1880-85. Anatomie menschlicher Embryonen; atlas. Vogel, Leipzig. [13] Ingalls, N. W. 1907. Arch. Mikroskop. Anat. Entwicklungsmech. 70:506. [14] Johnson, F. P. 1917. Contrib. Embryol. Carnegie Inst. Wash. 6:125. [15] Jones, H. O., and J. I. Brewer. 1941. Ibid. 29:157. [16] Keibel, F. 1910. In F. Keibel and F. P. Mall, ed. Manual of human embryology. J. B. Lippincott, Philadelphia. v. 1, pp. 59-90. [17] Ludwig, E. 1928. Morphol. Jahrb. 59:41. [18] Ludwig, E. 1929. Compt. Rend. Assoc. Anat. 24me Reunion (Bordeaux), p. 580. [19] Mankin, M. F., and J. Rock. 1948. Am. J. Obstet. Gynecol. 55:440. [20] Patten, B. M. 1953. Human embryology. Ed. 2. Blakiston, Philadelphia. ch. 5, 7. [21] Payne, F. 1925. Contrib. Embryol. Carnegie Inst. Wash. 16:115. [22] Politzer, G., and F. Hann. 1935. Z. Anat. Entwicklungsgeschichte 104:670. [23] Sensenig, E. C. 1957. Contrib. Embryol. Carnegie Inst. Wash. 36:141. [24] Shettles, L. B. 1953. Am. J. Obstet. Gynecol. 66:235. [25] Sternberg, H. 1927. Z. Anat. Entwicklungsgeschichte 82:747. [26] Streeter, G. L. 1951. Contrib. Embryol. Carnegie Inst. Wash., Embryol. Reprint Vol. 2. [27] Witschi, E. 1948. Contrib. Embryol. Carnegie Inst. Wash. 32:67. [28] Witschi, E. 1956. Development of vertebrates. W. B. Saunders, Philadelphia. [29] Witschi, E. 1959. Ann. N. Y. Acad. Sci. 75:412. [30] Witschi, E. 1962. In H. G. Grady and D. E. Smith, ed. The ovary. Williams and Wilkins, Baltimore. p. 1.

**Part II. RAT**

**Age** (column B) = days after fertilization, calculated from copulation age minus 8 hours (corresponding ages of mouse embryos of the same stage, based chiefly on references 7 and 8, are given in parentheses). **Size** (column C) = largest dimension of embryo in natural position (largest and smallest dimensions of blastocysts and chorionic vesicles are given in parentheses).

Standard Stages (Witschi)	Age da	Size mm	Identification of Stages
(A)	(B)	(C)	(D)
<b>Cleavage and Blastula</b>			
1	1	0.07	1 cell (in oviduct)
2	2 (1)	0.08 x 0.06	2 cells (in oviduct)
3	3		4 cells (in oviduct)
4	3.25 (2)	0.08 x 0.05	8-12 cells (in oviduct)
5	3.5	0.08 x 0.04	Morula (in uterus)
6	4	(0.08 x 0.03)	Early blastocyst (in uterus)
7	5 (4)	(0.12 x 0.05)	Free blastocyst (in uterus)
<b>Gastrula</b>			
8	6 (4.5)	(0.28 x 0.07)	Implanting blastocyst, with trophoblastic cone and inner cell mass; outgrowth of endoderm (hypoblast)
9	6.75 (5)		Diplo-trophoblast; inner cell mass (pendant), covered with endoderm
10	7.25 (5.5)	(0.3 x 0.1)	Near complete implantation; pendant begins differentiation into embryonic and extraembryonic parts
11	7.75 (6.5)	(0.5 x 0.1)	Completion of implantation; primary amniotic cyst; ectoplacental cone
<b>Primitive Streak</b>			
12	8.5 (7)	(1.04 x 0.26)	Connecting ecto-chorionic and amniotic cavities; rudiments of amniotic folds; primitive streak; start of 3rd layer formation; blastemas of heart and pericardium

*continued*

## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part II. RAT

Standard Stages (Witschi)	Age da	Size mm	Identification of Stages	
(A)	(B)	(C)	(D)	
Neurula				
13	13	9 (7.5)	1.0 (1.40 x 0.45)	Presomite neurula; fusion of chorioamniotic folds, chorioamniotic stalk; neural plate; embryo bent dorsally; bud of allantoic stalk
14	14	9.5 (7.75)	1.5 (1.8 x 1.1)	Somites 1-4 (occipital); pendant with 3 cavities: ectochorionic cyst, exocoelom, and amniotic cavity; ectochorionic cyst collapsing; allantoic stalk projects into exocoelom; embryo bent dorsally
15	15	10 (8.0 x 8.5)	2	Somites 5-12 (cervical); 1st visceral arch; ectochorionic cyst fused with ectoplacenta and allantoic stalk; regression of peripheral (distal) yolk sac and trophoblast (diplotrophoblast); Reichert's membrane; gonias in endoderm; embryo bent dorsally
16	16	10.5 (8.5 x 9.0)	2.4 (2.2 x 3.4)	Somites 13-20 (upper thoracic); 2 visceral arches; disc and yolk-sac placentas; appendicular folds; embryo reverses, curves ventrally
17	17	11 (9.5)	3.3	Somites 21-25 (lower thoracic); yolk stalk closes at level of 15th somite; primary gonias in mesentery; primitive streak disappears; tail bud becomes organized; arm and leg buds recognizable
Tail-Bud Embryo				
18	18	11.5 (10)	3.8	Somites 26-28 (upper lumbar); 3 visceral arches; arm buds recognizable
19	19	11.75 (10.25)	4.2	Somites 29-31 (lower lumbar); visceral arches I-IV; cervical folds; appendicular folds and buds
20	20	11.875	5 (4.7 x 5.2)	Somites 32, 33 (upper sacral)
21	21	12	5.1	Somites 34, 35 (lower sacral); deep cervical sinuses
22	22	12.125 (10.5)	5.2	Somite 36 (1st caudal); olfactory pits
23	23	12.25	5.6 (4.5 x 5.8)	Somites 37, 38 (caudal); start of umbilical herniation
24	24	12.375	6	Somites 39, 40 (caudal)
Complete Embryo				
25	25	12.5 (11)	6.2	Somites 41, 42 (caudal); occipital somites dispersing; 4 visceral arches; deep cervical sinuses; arm buds at somite levels 8-14, about as high as long; leg buds at somite levels 28-31, smaller; body forms a spiral of about 1½ turns, the left face and trunk applied to yolk sac, the right side turned toward placenta; tail and allantoic stalk rise to the placenta
Metamorphosing Embryo				
26	26	12.75	7	Somites 43-45 (caudal); mandibular, maxillary, and frontonasal processes; cervical sinuses closing; mammary welts; differentiation of hand plates; arm buds vascularized, brachial nerves entering; beginning of umbilical hernia
27	27	13 (12)	8	Somites 46-48 (caudal); prominent facial processes and clefts; nose-snout projecting; cervical sinuses closed; primordia of mammary glands; round hand plates and foot plates; larger umbilical hernia
28	28	13.5 (12.5)	8.5	Somites 49-51 (caudal); 1st visceral cleft transforms into external ear duct; precartilaginous condensations in hand plates
29	29	14	9.5	Somites 52-55 (caudal); auricular hillocks on visceral arches I and II
30	30	14.5 (13)	10.5	Somites 56-60 (caudal); body uncoils; mandibular precartilage; nearly round opening of external ear duct; pleuroperitoneal canal has become very narrow
31	31	15	12	Somites 61-63 (caudal); facial clefts closed; pleuroperitoneal canal closed; complete diaphragm
32	32	15.5 (14.5)	14.2 (14.3 x 8.0)	Somite 64 (caudal); pinna turns forward; maximal size of umbilical hernia
33	33	16 (15)	15.5	Somite 65 (usually this is last caudal); snout lifts off chest; last stage of metamorphosis
Fetus				
34	34	17-18 (16.0-16.5)	16-20	1st fetal stage: rapid growth of eyelids (eyes entirely covered at end of 18th day); palate complete; pinna covers ear duct; umbilical hernia withdraws

*continued*

## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part II. RAT

Standard Stages (Witschi)	Age da	Size mm	Identification of Stages	
(A)	(B)	(C)	(D)	
Fetus				
35	35 ante-natal	19-22 (17-19)	20-40	2nd fetal stage: sealed eyelids; fetal membranes and placentas reach peak of development; tail grows to 10 mm; birth occurs (22nd day in rat, 19th day in mouse)
36	35 post-natal	1-16 (1-20) post-partum	40-100 <sup>1</sup>	After birth, fetus becomes a breathing and suckling nestling <sup>2</sup> ; during 1st 16 days (22-38 days total age), eyelids remain sealed and external ear ducts plugged with periderm
37	36 post-natal	17+ (21+) post-partum	100+ <sup>1</sup>	Periderm seals of ears and eyelids vanish; active feeding begins within next 3 days and weaning after 1 week (total weaning age, 45-48 days for rats and mice)

/1/ Body length from nose to root of tail. During preimplantation stages, mouse development gains a lead of 1½ to 2 days and maintains it until birth; its nestling period is correspondingly longer so that the average weaning age is nearly the same in the two species. /2/ Developmentally, nestling period belongs to second fetal stage.

Contributor: Witschi, Emil

References: [1] Butcher, E. O. 1929. Am. J. Anat. 44:381. [2] Henneberg, B. 1937. Normentafel zur Entwicklungsgeschichte der Wanderratte (*Rattus norvegicus* Erxleben). G. Fischer, Jena. [3] Huber, G. C. 1915. J. Morphol. 26:1. [4] Long, J. A., and P. L. Burlingame. 1938. Univ. Calif. (Berkeley) Publ. Zool. 43:143. [5] MacDowell, E. C., E. Allen, and C. G. MacDowell. 1927. J. Gen. Physiol. 11:57. [6] Nicholas, J. S., and D. Rudnick. 1938. J. Exptl. Zool. 78:205. [7] Otis, E. M., and R. Brent. 1954. Anat. Record 120:33. [8] Snell, G. D. 1941. The early embryology of the mouse. Blakiston, Philadelphia. pp. 1-54. [9] Witschi, E. 1956. The development of vertebrates. W. B. Saunders, Philadelphia.

### Part III. SWINE

Size (column C) = greatest length, neck (spine) length, or crown-to-rump length of embryo.

Standard Stages (Witschi)	Age da	Size mm	Identification of Stages	Reference	
(A)	(B)	(C)	(D)	(E)	
Cleavage and Blastula					
1	1	0.11-0.14	1 cell	11	
2	2	1.0-1.5	2 cells	9,11	
3	3	2	4 cells; passes into uterus	4,5,9,11	
4	4	3	8-12 cells	4,5,9,11	
5	5	3.5	16 cells (morula)	9,11	
6	6	4.75	Blastocyst	6,9	
7	7	5-7	Late blastocyst still free in uterus	5,11	
Gastrula					
8	8	0.49-1.36 <sup>1</sup>	Bilaminar disc begins elongation	4,9-11	
9	9	2.5-3.0 <sup>1</sup>	Proliferation of mesoderm	4,9,11	
10	10	8-9	Beginning primitive streak	9-11	
11	11	10	Medium primitive streak	9,11	
Primitive Streak					
12	12	11-12	10-65 <sup>1</sup>	Completed primitive streak; notochord; becomes attached to endometrium	5,9,11

/1/ Extraembryonic length.

continued



**27. CHARACTERIZATION OF DEVELOPMENTAL STAGES**

**Part III. SWINE**

Standard Stages (Witschi)	Age da	Size mm	Identification of Stages	Reference
(A)	(B)	(C)	(D)	(E)
<b>Neurula</b>				
13	13		Presomite neurula	11
14	14-15	2.5-3.0	Occipital somites 1-4; 1st somite not delimited anteriorly	2,7,10, 11
15	15-16	3.2-5.2	Cervical somites 5-12	1,7
<b>Tail-Bud Embryo</b>				
16	15-17	5.2-6.5	Thoracic somites 13-20; spiral torsion; heart bulge	1,7
17	17	4.9	Thoracic somites 21-24	1,7
18	16.5-18.0	4.5	Thoracic somites 25, 26; head and tail meet; anterior limb bud	1,7
19	16.5-17.5	3.6	Lumbar somites 27-29; hindlimb bud	1,7
20	17.5	6.8	Lumbar somites 30, 31; spiraling completed	1,7
21	17.5	5.2	Lumbar somites 32, 33; uncoiling; mandibular and maxillary processes	7
22	19	5.8-8.0	Sacral somites 34, 35	7
23	20	6.4	Sacral somites 36, 37	7
24	20		Caudal somites 38-40	7
<b>Embryo</b>				
25	20	8.0-8.6	Caudal somites 41-43	7
26	20-21	9-10	Caudal somites 44-46; beginning of umbilical hernia	7,9
27	21-22	11	Caudal somites 47-49	7,9
28	22	11.6-14.4	Caudal somites 50-52; end of somite formation; cervical sinus closing; hand plate	3,7
29	22	16.4-18.6	Cervical sinus closed; lateral palatine processes; pentadactyl rudiments	7,9
30	28	19.4-24.0	Median (premaxillary) palatine processes; sex differentiation; eyelids and plica semilunaris	7,9
31	30	25	Facial clefts closing; palate developing	9
32	32.5	26.5-29.5	Phalanges 3 and 4 most prominent; fusion of palatine processes	3,9
33	34.5	35	Facial clefts closed; palate completed	7,11
<b>Fetus</b>				
34	36-50	35-55	1st fetal stage: growth of eyelids, gut withdrawal from umbilical cord	3,8
35	50-90	55-130	2nd fetal stage: sealed eyelids	3,8
36	90-113 <sup>a</sup>	130-280	3rd fetal stage: separation of eyelids	3,8

<sup>a</sup>/ Duration of pregnancy is usually given as 110-116 days, with extreme deviations for certain breeds. Young are born with open eyelids and open external ear ducts.

*Contributor:* Kemp, Norman E.

*References:* [1] Boyden, E. A. 1936. A laboratory atlas of the 13-mm. pig embryo. Wistar Institute Press, Philadelphia. [2] Boyden, E. A. 1940. Contrib. Embryol. Carnegie Inst. Wash. 28:157. [3] Carey, E. J. 1922. J. Morphol. 37:1. [4] Green, W. W., and L. M. Winters. 1946. Ibid. 78:305. [5] Heuser, C. H. 1927. Contrib. Embryol. Carnegie Inst. Wash. 19:229. [6] Heuser, C. H., and G. L. Streeter. 1929. Ibid. 20:1. [7] Keibel, F. 1897. Normentafeln zur Entwicklungsgeschichte der Wirbelthiere. G. Fischer, Jena. pt. 1. [8] MacCallum, J. B. 1901. Bull. Johns Hopkins Hosp. 12:102. [9] Patten, B. M. 1948. Embryology of the pig. Ed. 3. Blakiston, New York. [10] Streeter, G. L. 1927. Contrib. Embryol. Carnegie Inst. Wash. 19:73. [11] Waterman, A. J. 1948. A laboratory manual of comparative vertebrate embryology. H. Holt, New York.

*continued*

## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part IV. CHICK

Data adapted from Hamilton [2]. Times at which stages occur are approximate and are based on incubation temperature of 38°C.

Standard Stages (Witschi)	Chick Stages <sup>1</sup>	Age	Identification of Stages	
(A)	(B)	(C)	(D)	
Before Laying				
1	3, 4	Early cleavage	3.5-4.5 hr <sup>a</sup>	Shell membrane of egg formed in isthmus of oviduct
2	5, 6	During cleavage		Germ wall formed from marginal periblast
3	7	Late cleavage	4.5-24.0 hr <sup>a</sup>	Shell of egg formed in uterus
After Laying				
4	8, 9	1		Preprimitive streak (embryonic shield)
5	10	2	6-7 hr	Initial primitive streak, 0.3-0.5 mm long
6	11	3	12-13 hr	Intermediate primitive streak
7	12	4	18-19 hr	Definitive primitive streak, $\approx$ 1.88 mm long
8	13a	5	19-22 hr	Head process (notochord)
9	13b	6	23-25 hr	Head fold
10	14a	7	23-26 hr	1 somite; neural folds
11	14b	7 to 8-	ca. 23-26 hr	1-3 somites; coelom
12	14c	8	26-29 hr	4 somites; blood islands
13	15a	9	29-33 hr	7 somites; primary optic vesicles
14	15b	9+ to 10-	ca. 33 hr	8-9 somites; anterior amniotic fold
15	15c	10	33-38 hr	10 somites; 3 primary brain vesicles
16	16a	11	40-45 hr	13 somites; 5 neuromeres of hindbrain
17	16b	12	45-49 hr	16 somites; telencephalon
18	16c	13	48-52 hr	19 somites; atrioventricular canal
19	17a	13+ to 14-	ca. 50-52 hr	20-21 somites; tail bud
20	17b	14	50-53 hr	22 somites; trunk flexure; visceral arches I and II, clefts 1 and 2
21	17c	14+ to 15-	ca. 50-54 hr	23 somites; premandibular head cavities
22	17d	15	50-55 hr	24-27 somites; visceral arch III, cleft 3
23	18	16	51-56 hr	26-28 somites; wing bud; posterior amniotic fold
24	19	17	52-64 hr	29-32 somites; leg bud; epiphysis
25	20	18	3 da	30-36 somites extending beyond level of leg bud; allantois
26	21	19	3.0-3.5 da	37-40 somites extending into tail; maxillary process
27	22	20	3.0-3.5 da	40-43 somites; rotation completed; eye pigment
28	23	21	3.5 da	43-44 somites; visceral arch IV, cleft 4
29	24	22	3.5-4.0 da	Somites extend to tip of tail
30	25	23	4 da	Dorsal contour from hindbrain to tail is a curved line
31	26	24	4.5 da	Toe plate
32	27	25	4.5-5.0 da	Elbow and knee joints
33	28	26	5 da	1st 3 toes
34	29	27	5.0-5.5 da	Beak
35	30	28	5.5-6.0 da	3 digits, 4 toes
36	31	29	6.0-6.5 da	Rudiment of 5th toe
37	32	30	6.5-7.0 da	Feather germs; scleral papillae; egg tooth
38	33a	31	7.0-7.5 da	Web between 1st and 2nd digits
39	33b	32	7.5 da	Anterior tip of mandible has reached beak
40	34a	33	7.5-8.0 da	Web on radial margin of wing and 1st digit
41	34b	34	8 da	Nictitating membrane
42	34c	35	8.5-9.0 da	Phalanges in toes
43	34d	36	10 da	Length of 3rd toe from tip to middle of metatarsal joint = $5.4 \pm 0.3$ mm; length of beak from anterior angle of nostril to tip of bill = 2.5 mm; primordium of comb; labial groove; uropygial gland
44	34e	37	11 da	Length of 3rd toe = $7.4 \pm 0.3$ mm; length of beak = 3.0 mm
45	34f	38	12 da	Length of 3rd toe = $8.4 \pm 0.3$ mm; length of beak = 3.1 mm
46	35a	39	13 da	Length of 3rd toe = $9.8 \pm 0.3$ mm; length of beak = 3.5 mm
47	35b	40	14 da	Length of beak = 4.0 mm; length of 3rd toe = $12.7 \pm 0.5$ mm
48	35c	41	15 da	Length of beak from anterior angle of nostril to tip of upper bill = 4.5 mm; length of 3rd toe = $14.9 \pm 0.8$ mm
49	35d	42	16 da	Length of beak = 4.8 mm; length of 3rd toe = $16.7 \pm 0.8$ mm
50	35e	43	17 da	Length of beak = 5.0 mm; length of 3rd toe = $18.6 \pm 0.8$ mm

<sup>1/</sup> As described by Hamburger and Hamilton [1]. <sup>2/</sup> After ovulation.

*continued*

## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part IV. CHICK

Standard Stages (Witschi)	Chick Stages <sup>1</sup>	Age	Identification of Stages	
(A)	(B)	(C)	(D)	
After Laying				
51	35f	44	18 da	Length of beak = 5.7 mm; length of 3rd toe = 20.4±0.8 mm
52	36a	45	19-20 da	Yolk sac half-enclosed in body cavity; chorioallantoic membrane contains less blood and is "sticky" in living embryo
53	36b	46	20-21 da	Newly hatched chick

/1/ As described by Hamburger and Hamilton [1].

Contributor: Hamilton, Howard L.

References: [1] Hamburger, V., and H. L. Hamilton. 1951. J. Morphol. 88:49. [2] Hamilton, H. L., ed. 1952. Lillie's Development of the chick. Ed. 3. H. Holt, New York.

### Part V. FROG

Data are principally for *Rana pipiens*. At a given stage, age and size can be expected to vary widely with differences in geographic strains and culture conditions. **Frog Stages** (column B), designated by Arabic numerals, are for the embryo at 18°C and are adapted from Shumway [18]; those designated by Roman numerals are for the larva at 20°C and are adapted from Taylor and Kollros [22].

Standard Stages (Witschi)	Frog Stages	Age <sup>1</sup>	Size mm	Identification of Stages	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Cleavage and Blastula						
1	0	1	0	1.5-2.0	Unfertilized egg	18
2	1	2	1 hr	1.5-2.0	Fertilized egg; gray crescent	18
3	2	3	3.5 hr		2 cells	18
4	3	4	4.5 hr		4 cells	18
5	4	5	5.7 hr		8 cells	18
6	5	6	6.5 hr		16 cells	18
7	6	7	7.5 hr		32 cells	18
8	7a	8	16 hr		Middle blastula	18
9	7b	9	21 hr		Late blastula	18
Gastrula						
10	8	10	26 hr		Early gastrula; dorsal lip stage	18
11	9, 10	11	34 hr		Middle gastrula; blastopore C- or U-shaped	18
12	11	12	42 hr		Late gastrula; yolk plug; primitive gut	16,18
Neurula						
13	12	13	50 hr		Early neurula; medullary plate defined	10,16,18
14	13	14	62 hr		Midneurula; well-defined neural folds approaching each other; oral plate; anal pit; postanal gut	9,18
15	14, 15	15	67 hr		Late neurula; neural folds touch each other over most of their length; neurenteric canal; embryo rotates in jelly	9,18
16	16	16	72 hr	3	Neural tube, ectoderm fused over tube; oral sucker	18,19
Tail-Bud Embryo						
17	17	17	84 hr	3.5	Tail bud; nasal pit; dorsal aorta	10,14,18
18	18	18	96 hr	4	Muscular response to stimulation of myotome; lens placode	18,19
19	19	19	118 hr	5	Heart beats; pronephros functional; Rohon-Beard cells; thyroid evagination	12,14,15, 18
20	20	20	140 hr	6	Embryo hatched; gill circulation; lens vesicle	18,19

/1/ Comparable ages in hours at 20°C for frog stages 2 through 20: 0.5, 2.3, 3.2, 4.0, 4.8, 5.6, 7, 17, 22, 28, 30, 38, 43, 49, 52, 61, 76, 88, and 96, respectively.

continued



## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part V. FROG

Standard Stages (Witschi)	Frog Stages	Age	Size mm	Identification of Stages	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Tail-Bud Embryo						
21	21	162 hr	7	Mouth open; free-swimming; cornea becoming transparent; olfactory nerve; 2 rudiments of ventral pancreas; lung rudiments	14,18	
22	22	192 hr	8	Circulation in tail fin; cartilaginous trabeculae; 2 gill slits are perforate; trabeculae carneae	14	
23	23	216 hr	8-9	Opercular folds and labial teeth appear; spontaneous respiratory activity of mouth begins; basal plate	6,14,18	
24	24	240 hr	9-10	Operculum closed on right side; adrenal cortex rudiment; respiratory rhythm begins	6,18,20	
25	25a	284 hr	10-11	Operculum closed except for spiracle; rods and cones; germinal ridge; sucker regressed; rudiments of mesonephric tubules	4,16,18,19	
Tadpole to Adult (Metamorphosis)						
26	25b	I	3 da	13	Feeding begins; rudiments of adrenal medulla and of hindlimb appear	17,22
27	25c	II	6 da	17	Lagena; neural lobe of hypophysis	1,23
28	25d	III	11 da	23	Limb bud of equal length and diameter; lateral motor column	3,22
29	26a	IV	19 da	33	Ovarial sac; cartilage in synotic tectum	4,14
30	26b	V	23 da	39	Limb bud twice as long as it is broad; distal half of bud is bent ventrad	22
31	26c	VI	26 da	43	Flattened paddle at distal end of limb bud; scapular cartilage; gonads distinguishable	14,22,24
32	27a	VII	31 da	50	Foot paddle indented between toes 4 and 5	22
33	27b	VIII	34 da	53	Urinary bladder rudiment; measurable thyroid hormone output	2,12,14
34	27c	IX	36 da	56	Separation of fat body from gonad; spontaneous limb twitches	14,22
35	27d	X	40 da	58	Indentations delimit toe margins; rudiments of fungiform papillae of tongue	7,22
36	28a	XI	43 da	61	Margin of 5th toe web directed toward toe 2	22
37	28b	XII	47 da	64	Margin of 5th toe web directed toward toe 1	22
38	28c	XIII	52 da	67	Margin of 5th toe web directed toward prehallux	22
39	28d	XIV	58 da	70	Rudiments of harderian glands; rudiments of skin glands	13,14
40	29a	XV	62 da	72	1st toe pads; hindlimbs take part in swimming	12,22
41	29b	XVI	64 da	73	Nictitating membrane a low fold anterior to eyeball	12
42	29c	XVII	67 da	73	Some skin glands patent; peritoneal thickening presages oviduct	4,13
43	30a	XVIII	70 da	74 <sup>a</sup>	Cloacal tail piece resorbed; corneal reflex	11,22
44	30b	XIX	72 da	73	Tail regression begins; skin windows form	5,22
45	31	XX	74 da	70	Skin windows perforate; forelimbs emerge; oral beaks lost	5,22
46	32a	XXI	76 da	63	Upper lid forms; 1st molt	5,8
47	32b	XXII	79 da	44	Conjunctival sac complete; lateral lines regressing	8,22
48	33a	XXIII	81 da	33	Labial fringes completely lost; vasa efferentia	22,24
49	33b	XXIV	84 da	26	Tympanic membrane outlined; tail stub = 1-2 mm	22
50	33c	XXV	88 da	25 <sup>a</sup>	Tail stub fully resorbed; oviduct extends nearly to cloaca	4,22
51	Juvenile	Juvenile	90+ da	25-70	Fully metamorphosed; gonads immature; urostyle	21,22
52	Adult	Adult	1-3 yr	60-110	Sexually mature	22

/a/ Maximum size highly variable; tadpoles over 100 mm long have been collected. /s/ Size upon completion of metamorphosis highly variable, ranging from 16 to 30 mm.

Contributor: Kollros, Jerry J.

References: [1] Atwell, W. J. 1918. Anat. Record 15:73. [2] Barch, S.H. 1953. Physiol. Zool. 26:223. [3] Beaudoin, A. R. 1955. Anat. Record 121:81. [4] Christensen, K. 1930. Am. J. Anat. 45:159. [5] Etkin, W. 1932. Physiol. Zool. 5:275. [6] Fribourgh, J. H. 1949. M.S. Thesis. State Univ. Iowa, Iowa City. [7] Helff, O. M., and M. C. Mellicker. 1941. Am. J. Anat. 68:371. [8] Holbert, M. 1952. M.S. Thesis. State Univ. Iowa, Iowa City. [9] Huettner, A. F. 1949. Fundamentals of comparative embryology of the vertebrates. Macmillan, New York. [10] Knouff, R. A. 1935. J. Comp. Neurol. 62:17. [11] Kollros, J. J. 1942. J. Exptl. Zool. 89:37. [12] Kollros, J. J. Unpublished. State Univ. Iowa, Iowa City, 1956. [13] Kollros, J. J., and J. C. Kaltenbach. 1952. Physiol. Zool. 25:163. [14] Kopsch, F. 1952. Die Entwicklung des braunen Grasfrosches *Rana fusca*

continued



## 27. CHARACTERIZATION OF DEVELOPMENTAL STAGES

### Part V. FROG

Roesel. G. Thieme, Stuttgart. [15] Rappaport, R., Jr. 1955. J. Exptl. Zool. 128:481. [16] Rugh, R. 1951. The frog: its reproduction and development. Blakiston, Philadelphia. [17] Segal, S. 1953. Anat. Record 115:205. [18] Shumway, W. 1940. Ibid. 78:139. [19] Shumway, W. 1942. Ibid. 83:309. [20] Stenger, A. H., and H. A. Charipper. 1946. J. Morphol. 78:27. [21] Stokeley, P. S., and J. C. List. 1955. Trans. Am. Microscop. Soc. 74:112. [22] Taylor, A. C., and J. J. Kollros. 1946. Anat. Record 94:7. [23] Villy, F. 1890. Quart. J. Microscop. Sci. 30:522. [24] Witschi, E. 1929. J. Exptl. Zool. 52:235.

### Part VI. SALMONID FISHES

**Age** (columns B-D) = days after fertilization (differences in the data of Witschi [W.], Pasteels [P.], and Lagler [L.] are due to the fact that speed of development varies widely according to temperature and other environmental factors). **Size** (column E) = length of embryo or fry (diameter of blastopore is given in parentheses). Size and Identification of Stages were compiled mostly from data in the references of Kopsch [2] and Witschi [12]. For information on development of other teleost species, consult references 1, 4, 5, 9-11.

Standard Stages (Witschi)	Age da			Size mm	Identification of Stages
	W. <sup>1</sup>	P.	L.		
(A)	(B)	(C)	(D)	(E)	(F)
Cleavage					
1	1				1 cell, fertilized
2	2	0.5		0.5	2 cells
3	3				4 cells
4	4			1	8 cells
5	5				16-32 cells
6	6	1			Blastodisc up to 500 cells
7	7	4		5 (1.5)	Disc blastula, hypoderm largely syncytial; blastocoele
Gastrula					
8	8		3	(1.6)	Start of invagination around dorsal blastopore lip; prospective prechordal mesoblast
9	9		3.2	(1.8)	Prospective upper notochord invaginated, small gastrocoele
10	10	5	3.5	(2)	Prospective mid-notochord and first somites invaginating
11	11		3.7	0.4 (2.2)	Prospective trunk notochord and somites invaginating; primitive node forming
Primitive Streak					
12	12		4	12 0.7 (2.5)	Prospective lower notochord and median neuroblast form primitive node (axial rudiment); prospective lateral neuroblasts and left and right mesoblasts meet in germ wall (lateral folds of primitive streak)
Neurula					
13	13	10	5	12 1.0 (3.1)	Presomite neurula; lateral primitive folds start concurring
14	14		5.5	1.5 (3.7)	1-4 somites; neural plate narrowing, forming central solid cord of blastemic brain; upper notochord distinctly differentiated; hindgut vesicle
15	15	15	6	22 2.0 (4.5)	5-9 somites; forebrain and midbrain not separated, optic rudiments protruding; hindbrain narrow, imbedded between somitic mesoderm
16	16		6	68 2.5 (4.5)	10-15 somites; optic vesicles; forebrain and midbrain with slit-shaped cavity; otic vesicles with small cavity; conrescence constantly progresses, while yolk-sac epithelia spread over surface of yolk (in teleosts with small eggs, the yolk becomes much earlier engulfed by the yolk-sac epithelia; the yolk-sac blastopore may close even before any somites have become externally noticeable)
17	17	20	6.5	3.0 (3.5)	16-25 somites; indication of lens placodes over optic vesicles; 2 or 3 visceral arches externally recognizable

/1/ Approximates natural conditions near 5°C.

*continued*

**27. CHARACTERIZATION OF DEVELOPMENTAL STAGES**

**Part VI. SALMONID FISHES**

Standard Stages (Witschi)	Age da			Size mm	Identification of Stages
	W. <sup>1</sup>	P.	L.		
(A)	(B)	(C)	(D)	(E)	(F)
Tail-Bud Embryo					
18	18	7	83	3.5 (0.01)	26-30 somites; yolk-sac blastopore closes, embryo measures $\frac{1}{4}$ of entire egg circumference and is in full length attached to yolk-sac epithelium; brain cavity prolongating into upper spinal cord; flat lens placodes over optic vesicles; otic vesicles with small cavity that opens at surface; flat solid pharynx with 2 "pouches" reaching surface epithelium, 2nd one breaking through; nephrotomes in pronephric region begin to organize
19	19	30		4	31-40 somites; short, free tail bud; rhombencephalic roof becomes thin and broad; widening of ventricle
20	20		86	4.5	40-50 somites; optic vesicles start invagination; lens placodes thicken
21	21	40		5.2	50-55 somites; foregut with pharyngeal pouches; olfactory placodes thicken; deep optic cups; rudiments of plug-shaped lens; nephric blastema-nephrostomes; nephric ducts get lost in lower mesonephric blastema cords; primordial germ cells widely scattered on both sides in mesodermal blastema; hindgut vesicle
22	22	50		6	55-58 somites; 2nd and 3rd pharyngeal pouches open on surface; pectoral limb buds; olfactory placodes saucer-shaped; free lenses in eye cups; round otic vesicle develops a thick neural epithelium ventrally; acoustic ganglion. Embryo begins to separate from yolk sac, especially the forehead; liver diverticles forming; tubular midgut; anus forming; nephric tubules and corpuscles; nephric ducts with free ends left and right of rectum.
23	23	60	86	10	58-60 somites; all somites have formed; pectoral fins fairly large and differentiated; buds of pelvic fins are present but not externally noticeable; eyes now heavily pigmented; otic vesicles become labyrinths; cartilaginous skeleton of head and upper body; germ cells still free in peritoneum or mesenchyme near nephric ducts and blastema; renal corpuscles; nephric ducts unite caudal to rectum
24	24	70		15	Hatching fish (fry); pelvic fin bud below caudal end of yolk sac (about 30th somite); anus at about 40th somite; total number of somites varies normally from about 56 to 60; swim bladder has grown out from dorsal wall of esophagus; undifferentiated gonads; skin pigment appearing
Young Adult					
25	25	80		20+	General appearance as at preceding stage, but pelvic and all unpaired fins are now well-differentiated; skin heavily pigmented; yolk sac shrinks and eventually disappears

/1/ Approximates natural conditions near 5°C.

Contributor: Witschi, Emil

References: [1] Armstrong, P. B. 1963. Stages in the development of *Ictalurus nebulosis*. Syracuse Univ. Press, Syracuse, N. Y. [2] Kopsch, F. 1898. Arch. Mikroskop. Anat. Entwicklungsmech. 51:181. [3] Lagler, K. F. 1952. Freshwater fishery biology. W. C. Brown, Dubuque, Iowa. [4] Oppenheimer, J. M. 1937. Anat. Record 68:1. [5] Outram, D. 1957. Turtox News 35:16. [6] Pasteels, J. 1936. Arch. Biol. (Leige) 47:206. [7] Schmidt, J. 1921. Compt. Rend. Trav. Lab. Carlsberg 14(15):1. [8] Schmidt, J. 1921. Ibid. 14(16):1. [9] Solberg, A. N. 1938. Progressive Fish Culturist 40. [10] Swarup, H. 1958. J. Embryol. Exptl. Morphol. 6:373. [11] Wilson, H. V. 1889. Bull. U.S. Fisheries Comm. 9:209. [12] Witschi, E. 1956. The development of vertebrates. W. B. Saunders, Philadelphia.

# Contrails

## 28. GROWTH: MAMMALS

For information on other body measurements, consult reference 1, Part I.

### Part I. BODY WEIGHT AND HEIGHT: MAN

Subject	Date	Age yr	Males		Females		Reference	Subject	Date	Age yr	Males		Females		Reference
			Wt kg	Ht cm	Wt kg	Ht cm					Wt kg	Ht cm			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 African Pygmy	1935	Birth	3.6	45.9	3.9	47.0	4	63 Czecho-slovakian	1931	3	....	89.7	....	87.7	4
2		1	....	71.0	7.0	65.0		64		5	17.1	103.5	16.8	102.7	
3		5	13.5	100.4	13.5	98.1		65	1934	7	21.7	115.6	21.3	115.0	
4		7	17.6	103.9	....	102.8		66		9	26.0	125.9	25.8	124.9	
5		9	....	113.3	....	113.0		67		11	30.2	133.4	30.2	133.4	
6		11	23.0	....	18.5	....		68		13	35.5	142.9	38.0	144.7	
7 Argentine	1931	9	28.1	123	27.6	122	4	69	1929	15	52.1	156.0 <sup>1</sup>	....	....	
8		11	33.1	136	33.9	132		70		17	61.3	159.0 <sup>1</sup>	....	....	
9		13	40.6	147	41.9	145		71 Dane	1930	Birth	3.4	....	3.3	....	4
10		15	44.9	153	46.6	153		72		7	21.8	118.5	21.5	117.9	
11 Austrian	1932	1	10.3	76	9.8	75	4	73		9	26.2	127.9	26.2	127.5	
12		3	14.5	95	14.0	94		74		11	31.8	137.5	32.3	137.5	
13		5	18.5	108	18.0	107		75		13	38.1	146.5	40.8	148.6	
14		7	22.5	119	22.0	118		76		15	49.2	160.1	49.6	158.0	
15		9	26.7	128	26.0	127		77 Frenchman	1935	Birth	3.1	49.9	3.1	49.2	4
16		11	31.8	137	31.5	138		78		5	18.4	107.9	17.8	106.9	
17		13	38.5	148	38.5	149		79		7	22.2	117.7	21.2	116.4	
18		15	48.0	....	47.5	....		80		9	27.0	127.9	26.8	127.7	
19 Briton	1935	Birth	3.4	51.2	3.3	50.8	4	81		11	32.6	137.8	33.9	138.6	
20		1	9.3	70.8	8.8	69.8		82 German	1928	Birth	3.5	51.0	5.3	50.5	4
21		3	13.7	96.3	13.2	96.0		83		1	9.7	75.0	9.5	74.5	
22		5	15.5	108.2 <sup>1</sup>	15.9	107.9 <sup>1</sup>	4	84		3	14.1	93.0	13.7	92.5	
23		7	19.8	118.1 <sup>1</sup>	19.3	117.9 <sup>1</sup>		85		5	18.6	107.0	17.8	106.5	
24		9	23.4	128.0 <sup>1</sup>	22.0	127.8 <sup>1</sup>		86		7	22.2	119.5	21.6	118.5	
25		11	26.8	136.7 <sup>1</sup>	27.2	137.9 <sup>1</sup>		87		9	26.7	129.0	25.9	128.5	
26		13	32.9	148.1 <sup>1</sup>	32.4	149.6 <sup>1</sup>		88		11	31.5	137.5	31.4	139.0	
27		14	42.7	155.2	45.5	154.4	3	89		13	37.5	147.0	40.2	151.0	
28		15	45.9	158.5	47.7	156.0		90		15	48.1	160.5	49.5	160.0	
29		16	50.9	163.8	50.0	157.5		91		17	59.8	170.5	56.3	164.0	
30		17	54.5	167.4	51.4	157.7		92		19	64.5	174.0	59.0	165.0	
31		18	56.4	168.4	51.8	158.0		93 Japanese	1949	Birth	3.1	50.2	3.0	49.3	8
32		20	59.5	169.9	52.7	158.2		94	1960	1	10.3	77.9	9.6	76.0	2
33		22	61.8	170.9	53.2	158.5		95		2	12.2	85.7	11.5	84.3	
34 Canadian	1953	2	13.6	88.1	12.7	85.3	5	96		3	14.0	93.4	13.6	92.6	
35		3	14.5	93.0	14.0	91.4		97		4	15.5	99.6	15.0	98.5	
36		4	16.8	99.6	16.3	99.6		98		5	17.1	104.7	16.5	104.0	
37		5	18.1	106.4	18.6	106.2		99		6	19.0	110.8	18.4	109.7	
38		6	20.8	113.3	19.9	112.3		100		7	21.0	116.7	20.6	115.5	
39		7	22.7	119.4	22.2	118.1		101		8	22.9	121.5	22.6	120.6	
40		8	25.8	124.7	25.8	124.2		102		9	25.6	126.6	25.1	126.0	
41		9	28.5	130.8	28.1	129.5		103		10	27.6	130.8	28.1	131.7	
42		10	31.7	135.9	31.3	135.4		104		11	30.5	135.9	32.1	138.0	
43		11	34.9	140.7	34.9	140.5		105		12	34.2	141.0	36.5	143.7	
44		12	38.1	145.8	41.7	147.8		106		13	39.1	147.6	40.7	147.9	
45		13	42.6	150.6	46.2	153.4		107		14	43.9	153.6	44.4	149.4	
46		14	48.9	158.0	48.5	155.7		108		15	49.4	158.7	47.9	151.5	
47		15	53.9	164.3	50.7	158.0		109		16	52.8	159.9	48.6	152.1	
48		16-17	61.6	169.4	54.4	158.7		110		17	54.9	163.2	50.1	151.8	
49		18-19	65.2	172.7	56.2	159.0		111		18	56.0	162.9	49.8	152.4	
50		20-24	69.8	172.5	56.2	159.5		112		19	55.4	163.2	50.6	152.7	
51 Chinese	1935	Birth	3.1	48.2	3.0	48.2	4	113 Roumanian	1937	5	17.5	104.5	17.0	103.7	4
52		1	....	73.5	....	71.4		114		7	21.2	115.8	20.6	115.4	
53		3	....	92.0	....	89.7		115		9	24.6	123.0	23.8	122.8	
54		5	14.9	104.4	14.0	108.0		116		11	28.7	131.2	28.7	131.6	
55		7	18.4	114.6	20.1	117.2		117		13	33.8	139.1	34.7	140.2	
56		9	22.2	123.7	23.4	126.6		118		15	39.8	146.8	41.8	148.1	
57		11	26.5	131.3	28.7	135.4		119 Russian	1935	Birth	3.4	48.6	3.3	48.6	4
58		13	32.3	141.1	37.0	145.7		120	1931	9	....	125.3	....	124.5	
59		15	41.7	152.7	45.0	150.0		121 Spaniard	1934	7	21.6	117.0	21.5	116.0	4
60		17	48.3	162.4	47.5	154.3		122		9	24.7	123.0	26.0	124.6	
61		19	52.6	165.0	44.4	152.0		123		11	29.0	132.7	31.5	134.0	
62		21	53.8	166.0	49.0	156.0		124		13	35.7	139.6	38.0	143.7	

/1/ Date of measurement, 1933.

continued

**28. GROWTH: MAMMALS**

**Part I. BODY WEIGHT AND HEIGHT: MAN**

Subject	Date	Age yr	Males		Females		Reference	Subject	Date	Age yr	Males		Females		Reference		
			Wt kg	Ht cm	Wt kg	Ht cm					Wt kg	Ht cm	Wt kg	Ht cm			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)		
125	Swiss	1935	Birth	3.3	.....	3.1	.....	4									
126		1933	7	22	118	19	110		155	U. S.	1925	11	....	140.9	....	141.2	4
127			9	26	126	23	118		156	Negro		12	37.0	144.8	40.0	149.0	
128			11	30	136	30.5	136		157			13	43.9	150.1	43.9	153.7	
129			13	36	146	40	147		158			14	48.0	156.5	47.7	154.7	
130			15	47	157	44.5	152		159			15	53.0	161.0	53.0	158.1	
	U. S.								160			16	57.0	163.6	57.0	157.8	
131	Navaho	1936	6	20.2	116.1	19.0	113.4	4	161			17	....	167.0	....	158.5	
132	Indian		7	21.7	120.8	21.2	119.0		162			18	....	170.1	....	159.2	
133			8	24.0	125.7	23.1	123.9		163			19	....	172.0	....	159.9	
134			9	26.2	131.1	25.6	129.9		164	White	1934-	Birth	3.5	50.8	3.4	50.0	7
135			10	28.4	135.3	28.3	134.9		165		59	1	10.5	75.4	9.5	74.4	
136			11	30.4	138.9	31.2	140.2		166			2	12.7	87.6	12.3	86.6	
137			12	32.8	143.0	34.6	143.9		167			3	14.5	96.0	14.1	95.3	
138			13	36.6	149.2	39.0	150.2		168			4	16.8	103.6	16.4	103.1	
139			14	41.2	153.8	43.6	153.1		169			5	19.1	111.0	18.6	111.3	
140			15	45.3	160.5	47.7	155.3		170			6	21.4	117.1	20.5	116.1	
141			16	50.0	165.5	51.1	156.3		171			7	24.5	122.4	22.7	121.7	
142			17	54.0	167.5	52.3	157.4		172			8	27.3	128.0	26.4	127.8	
143			18	56.3	169.5	54.0	157.5		173			9	30.0	134.1	29.1	132.3	
144	Negro	1961	Birth	3.2	49.6	3.3	49.6	6	174			10	33.2	138.4	32.7	138.7	
145			1	10.4	76.1	9.9	74.9		175			11	37.3	144.3	37.3	145.0	
146		1925	2	....	89.1	....	85.9	4	176			12	39.5	148.1	42.3	151.4	
147			3	....	95.2	....	95.8		177			13	45.0	154.2	46.4	156.0	
148			4	....	102.0	....	100.6		178			14	51.4	161.5	50.9	159.5	
149			5	....	110.0	....	109.9		179			15	58.2	168.4	53.2	161.0	
150			6	....	116.2	....	116.2		180			16	62.3	172.0	54.5	162.3	
151			7	....	120.3	....	120.8		181			17	65.0	173.5	55.5	162.8	
152			8	....	125.8	....	124.6		182			18	67.7	174.0	55.9	162.8	
153			9	....	130.8	....	131.3		183			19	69.5	174.2	56.4	162.8	
154			10	....	135.3	....	135.2		184			20-24	71.8	174.5	56.8	162.6	

*Contributors:* (a) Krogman, W. M., (b) Arimoto, Kunitaro, (c) Pett, L. Bradley, (d) Scott, Roland B., (e) Damon, Albert

*References:* [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Arimoto, K. Unpublished. Natl. Institute of Nutrition, Tokyo, 1960. [3] Kemsley, W. F. F. 1950. Ann. Eugenics 15:161. [4] Krogman, W. M. 1941. Tabulae Biologicae 20. [5] Pett, L. B. 1954. Survey, 1953. Nutrition Div., Dept. of Natl. Health and Welfare, Ottawa, Canada. [6] Scott, R. B., et al. Unpublished. Freedmen's Hospital, Washington, D. C. 1962. [7] Stoudt, H. W., A. Damon, and R. A. McFarland. 1960. Human Biol. 32:331. [8] Yanagi, K., et al. 1949. Studies on dietary allowances for Japanese. Natl. Council of Food and Nutrition, Tokyo. p. 1.

**Part II. BODY WEIGHT: RODENTS**

Species	Subjects (Age)	Weight g	Reference	Species	Subjects (Age)	Weight g	Reference
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
<i>Cavia porcellus</i> (guinea pig) <sup>1</sup>				<i>Cavia porcellus</i> (guinea pig) <sup>1</sup>			
1 Inbred line	112♂ (birth)	77(55-99)	6	6 Inbred line	112♂ (83 da)	381(239-523)	6
2 no. 2	112♂ (13 da)	127(87-167)		7 no. 2	112♂ (113 da)	474(336-612)	
3	112♂ (23 da)	165(107-223)		8	112♂ (143 da)	540(406-674)	
4	112♂ (33 da)	192(114-270)		9	112♂ (173 da)	585(455-715)	
5	112♂ (53 da)	266(166-366)		10	112♂ (233 da)	648(516-780)	

/-/ Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction).

*continued*



## 28. GROWTH: MAMMALS

### Part II. BODY WEIGHT: RODENTS

Species				Species				
Subjects (Age)	Weight g	Reference	Subjects (Age)	Weight g	Reference			
(A)	(B)	(C)	(A)	(B)	(C)	(D)		
<i>Cavia porcellus</i> (guinea pig) <sup>1</sup>			<i>Mus musculus</i> (house mouse)					
11 Inbred line no. 2	112♂ (293 da)	689(555-823)	6	71 Piebald,	81♂ (28 wk)	33.07(23.70-45.45)	5	
12	112♂ (353 da)	709(581-837)		72	78♀ (28 wk)	33.00(21.43-46.97)		
13	112♂ (413 da)	729(597-861)		73	81♂ (32 wk)	34.07(22.68-46.67)		
14	112♂ (473 da)	744(612-876)		74	78♀ (32 wk)	34.52(22.42-49.85)		
15	112♂ (533 da)	759(631-887)		75	81♂ (36 wk)	35.49(24.53-50.93)		
16	112♂ (593 da)	764(642-886)		76	78♀ (36 wk)	36.38(22.58-50.68)		
17	112♂ (653 da)	775(647-903)		77	White <sup>1</sup>	50♂ (3 wk)	8.16(7.34-8.98)	7
18	112♂ (713 da)	778(656-900)		78		50♂ (4 wk)	12.44(11.58-13.30)	
19	Random-bred line	68♂ (birth)	94(56-132)	79		50♂ (5 wk)	17.1(16.2-18.0)	
20		68♂ (13 da)	168(102-232)	80		50♂ (6 wk)	19.56(18.66-20.46)	
21		68♂ (23 da)	227(143-311)	81		50♂ (7 wk)	21.06(20.00-22.12)	
22		68♂ (33 da)	268(172-364)	82		50♂ (8 wk)	22.22(20.96-23.48)	
23		68♂ (53 da)	345(211-479)	83		43♂ (12 wk)	25.27(24.19-26.35)	
24		68♂ (83 da)	456(280-632)	84		42♂ (16 wk)	27.19(25.69-28.69)	
25		68♂ (113 da)	571(389-753)	85		42♂ (20 wk)	27.81(26.39-29.23)	
26		68♂ (143 da)	658(460-856)	86		38♂ (24 wk)	27.58(26.36-28.80)	
27		68♂ (173 da)	719(525-913)	87		24♂ (28 wk)	28.04(26.38-29.70)	
28		68♂ (233 da)	813(599-1027)	88		22♂ (32 wk)	29.36(28.22-30.50)	
29		68♂ (293 da)	872(668-1076)					
30		68♂ (353 da)	910(686-1134)					
31		68♂ (413 da)	938(714-1162)	89	Long-Evans, mixed	30-50♂ (birth)	6.12	3, 4
32		68♂ (473 da)	952(742-1162)	90		30-50♀ (birth)	5.75	
33		68♂ (533 da)	985(777-1193)	91		30-50♂ (3 wk)	40	
34		68♂ (593 da)	992(774-1210)	92		30-50♀ (3 wk)	39	
35		68♂ (653 da)	1001(805-1197)	93		30-50♂ (4 wk)	56	
36		68♂ (713 da)	1022(780-1264)	94		30-50♀ (4 wk)	52	
37	<i>Mesocricetus auratus</i> (golden hamster) <sup>2</sup>	111 (3 da)	3.3(2.0-5.0)	2		30-50♂ (5 wk)	92	
38		111 (7 da)	7(4-10)			30-50♀ (5 wk)	84	
39		111 (18 da)	23(9-33)	96		30-50♂ (6 wk)	125	
40		111 (25 da)	40(17-51)	97		30-50♀ (6 wk)	105	
41		111 (45 da)	79(65-88)	98		30-50♂ (7 wk)	155	
42		111 (90 da)	103(94-112)	99		30-50♀ (7 wk)	123	
43		111 (180 da)	112(103-121)	100		30-50♂ (8 wk)	185	
44		111 (730 da)	105(97-113)	101		30-50♀ (8 wk)	140	
				102		30-50♂ (8 wk)	221	
				103		30-50♀ (10 wk)	167	
				104		30-50♂ (10 wk)	259	
				105		30-50♀ (12 wk)	178	
				106		30-50♂ (12 wk)	295	
				107		30-50♀ (15 wk)	198	
				108		30-50♂ (15 wk)	350	
				109		30-50♀ (20 wk)	223	
				110		30-50♂ (20 wk)	405	
				111		30-50♀ (30 wk)	250	
				112		30-50♂ (30 wk)	434	
				113		30-50♀ (40 wk)	265	
				114		30-50♂ (40 wk)	470	
				115		30-50♀ (52 wk)	276	
				116				
				117	Sherman, albino, small <sup>1</sup>	27♂ (birth)	5.5(4.9-6.1)	8
				118		39♀ (birth)	5.52(4.80-6.20)	
				119		27♂ (1 wk)	13.4(10.0-16.5)	
				120		39♀ (1 wk)	13.1(10.5-16.0)	
				121		27♂ (2 wk)	25.3(20.5-30.0)	
				122		39♀ (2 wk)	24.9(20.0-30.0)	
				123		27♂ (3 wk)	40.4(32.0-48.5)	
				124		39♀ (3 wk)	36.8(30.5-43.0)	
				125		27♂ (4 wk)	60.7(50.5-70.5)	
				126		39♀ (4 wk)	55.6(48.0-63.0)	
				127		27♂ (5 wk)	89.6(76.0-104.0)	
				128		39♀ (5 wk)	77.5(68.0-87.0)	
				129		27♂ (6 wk)	121(106-136)	
				130		39♀ (6 wk)	100(89-111)	

<sup>1</sup>/ Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction). <sup>2</sup>/ Values in parentheses (column C) are ranges, estimate "c" (cf. Introduction).

continued

## 28. GROWTH: MAMMALS

### Part II. BODY WEIGHT: RODENTS

Species	Subjects (Age)	Weight g	Reference	Species	Subjects (Age)	Weight g	Reference	
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	
<i>Rattus norvegicus</i> (Norway rat)				<i>Rattus norvegicus</i> (Norway rat)				
131	Sherman, 27♂ (7 wk)	150(124-175)	8	181	Wistar, al- 30-50♂ (5 wk)	84	3,4	
132	albino, 39♀ (7 wk)	116(106-127)		182	bino 30-50♀ (5 wk)	79		
133	small <sup>1</sup> 27♂ (8 wk)	177(149-205)		183	30-50♂ (6 wk)	110		
134	39♀ (8 wk)	130(122-137)		184	30-50♀ (6 wk)	97		
135	27♂ (10 wk)	222(191-254)		185	30-50♂ (7 wk)	141		
136	39♀ (10 wk)	154(145-163)		186	30-50♀ (7 wk)	116		
137	27♂ (12 wk)	252(213-291)		187	30-50♂ (8 wk)	170		
138	39♀ (12 wk)	169(159-179)		188	30-50♀ (8 wk)	128		
139	27♂ (15 wk)	285(241-329)		189	30-50♂ (10 wk)	200		
140	39♀ (15 wk)	185(165-205)		190	30-50♀ (10 wk)	147		
141	27♂ (20 wk)	326(278-373)		191	30-50♂ (12 wk)	225		
142	39♀ (20 wk)	202(178-225)		192	30-50♀ (12 wk)	165		
143	27♂ (30 wk)	376(335-417)		193	30-50♂ (15 wk)	251		
144	39♀ (30 wk)	230(205-255)		194	30-50♀ (15 wk)	180		
145	27♂ (40 wk)	240(215-265)		195	30-50♂ (20 wk)	280		
146	39♀ (52 wk)	248(221-275)		196	30-50♀ (20 wk)	200		
147	Sherman, 26♂ (birth)	6.08(4.90-7.70)	8	197	30-50♂ (30 wk)	322		
148	albino, 38♀ (birth)	5.75(4.90-6.60)		198	30-50♀ (30 wk)	228		
149	large <sup>1</sup> 26♂ (1 wk)	17.5(12.5-22.5)		199	30-50♂ (40 wk)	342		
150	38♀ (1 wk)	16.2(13.0-19.0)		200	30-50♀ (40 wk)	245		
151	26♂ (2 wk)	36.9(28.5-45.0)		201	30-50♂ (52 wk)	364		
152	38♀ (2 wk)	33.5(27.5-39.5)		202	30-50♀ (52 wk)	243		
153	26♂ (3 wk)	59.6(48.0-71.0)		203	Wild <sup>e</sup> ♂ (40 da)	85(67-112)		1
154	38♀ (3 wk)	53(43-63)		204	♀ (40 da)	104(79-142)		
155	26♂ (4 wk)	92.9(78.0-108.0)		205	♂ (60 da)	170(127-218)		
156	38♀ (4 wk)	79.5(68.0-91.0)		206	♀ (60 da)	152(120-200)		
157	26♂ (5 wk)	138(117-158)		207	♂ (80 da)	237(176-299)		
158	38♀ (5 wk)	113(99-128)		208	♀ (80 da)	194(149-249)		
159	26♂ (6 wk)	188(157-218)		209	♂ (100 da)	289(217-361)		
160	38♀ (6 wk)	147(128-166)		210	♀ (100 da)	230(178-291)		
161	26♂ (7 wk)	233(195-260)		211	♂ (120 da)	330(251-408)		
162	38♀ (7 wk)	172(149-195)		212	♀ (120 da)	260(203-327)		
163	26♂ (8 wk)	274(231-317)	213	♂ (160 da)	388(302-472)			
164	38♀ (8 wk)	196(169-222)	214	♀ (160 da)	311(245-383)			
165	26♂ (10 wk)	339(291-386)	215	♂ (200 da)	424(335-509)			
166	38♀ (10 wk)	227(199-256)	216	♀ (200 da)	348(276-423)			
167	26♂ (12 wk)	393(328-458)	217	♂ (240 da)	446(358-531)			
168	38♀ (12 wk)	251(232-280)	218	♀ (240 da)	376(300-452)			
169	26♂ (15 wk)	440(379-501)	219	♂ (280 da)	460(374-545)			
170	38♀ (15 wk)	274(238-310)	220	♀ (280 da)	397(319-473)			
171	26♂ (20 wk)	490(423-556)	221	♂ (320 da)	468(385-551)			
172	38♀ (20 wk)	303(270-335)	222	♀ (320 da)	413(333-488)			
173	26♂ (30 wk)	335(298-373)	223	♂ (360 da)	474(392-556)			
174	38♀ (40 wk)	358(311-404)	224	♀ (360 da)	424(344-497)			
175	Wistar, al- 30-50♂ (birth)	5.63	3,4	225	♂ (400 da)	477(397-558)		
176	bino 30-50♀ (birth)	5.3		226	♀ (400 da)	433(352-507)		
177	30-50♂ (3 wk)	43		227	♂ (440 da)	480(400-560)		
178	30-50♀ (3 wk)	41		228	♀ (440 da)	440(358-512)		
179	30-50♂ (4 wk)	52		229	♂ (480 da)	482(402-561)		
180	30-50♀ (4 wk)	55		230	♀ (480 da)	445(363-516)		

/1/ Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction). /3/ Values in parentheses (column C) are the 10th and 90th percentiles.

**Contributors:** (a) Calhoun, John B., (b) DuBois, R. Callery, (c) Latyszewski, M., (d) Mills, Clarence A., (e) Wright, Sewall, (f) Zucker, Lois M.

**References:** [1] Calhoun, J. B. Unpublished. Walter Reed Army Med. Center, Washington, D. C., 1952. [2] Dubois, R. C. Unpublished, 1950. [3] Freudenberg, C. B. 1932. Am. J. Anat. 50:293. [4] Freudenberg, C. B. 1933.

continued

## 28. GROWTH: MAMMALS

### Part II. BODY WEIGHT: RODENTS

Anat. Record 56:47. [5] Kopeć, S. 1930. Mem. Inst. Natl. Polon. Econ. Rurale 11:2. [6] McPhee, H. C., and O. N. Eaton. 1931. U. S. Dept. Agr. Tech. Bull. 222. [7] Ogle, C. 1934. Am. J. Physiol. 107:635. [8] Zucker, L. M., and T. F. Zucker. Unpublished. Columbia Univ., New York, 1952.

### Part III. BODY WEIGHT: MAMMALS OTHER THAN MAN AND RODENTS

Species	Subjects (Age)	Weight kg	Reference	Species	Subjects (Age)	Weight kg	Reference
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
<i>Bos taurus</i> (cattle)				<i>Bos taurus</i> (cattle)			
1 Ayrshire	58♂ (birth)	36.8	3	53 Jersey	95♂ (birth)	26.4	3
2	100♀ (birth)	35.9	2	54	105♀ (birth)	24.5	2
3	23♂ (3 mo)	74.5	3	55	57♂ (3 mo)	59.1	3
4	81♀ (3 mo)	84.1	2	56	82♀ (3 mo)	62.3	2
5	12♂ (6 mo)	140.9	3	57	31♂ (6 mo)	114.1	3
6	91♀ (6 mo)	155.9	2	58	97♀ (6 mo)	129.1	2
7	4♂ (9 mo)	205.9	3	59	8♂ (9 mo)	184.1	3
8	75♀ (9 mo)	222.7	2	60	76♀ (9 mo)	188.2	2
9	3♂ (12 mo)	289.1	3	61	89♀ (12 mo)	231.8	2
10	86♀ (12 mo)	266.4	2	62	78♀ (18 mo)	301.8	2
11	76♀ (18 mo)	354.5	2	63	80♀ (2 yr)	365.5	2
12	72♀ (2 yr)	448.6	2	64	47♀ (3 yr)	406.8	2
13	56♀ (3 yr)	501.4	2	65	37♀ (4 yr)	441.4	2
14	32♀ (4 yr)	525.9	2	66	22♀ (5 yr)	464.5	2
15	18♀ (5 yr)	550.9	2	67	7♀ (6 yr)	441.8	2
16	8♀ (6 yr)	543.2	2	68	4♀ (7 yr)	482.3	2
17	4♀ (7 yr)	589.5	2				
18 Guernsey	25♂ (birth)	35.5	3	69 <i>Canis familiaris</i> (dog)			
19	78♀ (birth)	32.7	2	Basenji <sup>1</sup>	23♂ (birth)	0.29(0.20-0.38)	12
20	14♂ (3 mo)	69.1	3	70	27♀ (birth)	0.27(0.22-0.32)	
21	59♀ (3 mo)	69.5	2	71	23♂ (1 wk)	0.49(0.27-0.71)	
22	8♂ (6 mo)	129.1	3	72	27♀ (1 wk)	0.43(0.23-0.63)	
23	69♀ (6 mo)	138.2	2	73	23♂ (2 wk)	0.73(0.45-1.01)	
24	3♂ (9 mo)	204.5	3	74	27♀ (2 wk)	0.65(0.41-0.89)	
25	56♀ (9 mo)	201.4	2	75	23♂ (4 wk)	1.12(0.72-1.52)	
26	68♀ (12 mo)	246.4	2	76	27♀ (4 wk)	0.96(0.60-1.32)	
27	54♀ (18 mo)	330.0	2	77	23♂ (6 wk)	1.51(0.85-2.17)	
28	58♀ (2 yr)	414.5	2	78	27♀ (6 wk)	1.37(0.87-1.87)	
29	34♀ (3 yr)	469.5	2	79	23♂ (8 wk)	2.30(1.16-3.44)	
30	21♀ (4 yr)	506.4	2	80	27♀ (8 wk)	2.13(1.39-2.87)	
31	12♀ (5 yr)	519.5	2	81	23♂ (10 wk)	3.29(1.69-4.89)	
32	2♀ (6 yr)	586.8	2	82	27♀ (10 wk)	2.96(1.87-4.05)	
33	2♀ (7 yr)	526.4	2	83	23♂ (12 wk)	4.49(2.55-6.43)	
34 Holstein	220♂ (birth)	45.5	3	84	27♀ (12 wk)	3.97(2.65-5.29)	
35	262♀ (birth)	40.0	2	85	23♂ (14 wk)	5.66(3.58-7.74)	
36	145♂ (3 mo)	99.1	3	86	27♀ (14 wk)	4.97(3.71-6.23)	
37	256♀ (3 mo)	97.7	2	87	23♂ (16 wk)	6.57(4.34-8.80)	
38	104♂ (6 mo)	190.5	3	88	27♀ (16 wk)	5.51(4.08-6.94)	
39	247♀ (6 mo)	181.4	2	89 Beagle <sup>1</sup>	39♂ (birth)	0.31(0.17-0.45)	12
40	65♂ (9 mo)	287.3	3	90	31♀ (birth)	0.30(0.21-0.39)	
41	244♀ (9 mo)	258.2	2	91	39♂ (1 wk)	0.55(0.37-0.73)	
42	25♂ (12 mo)	370.5	3	92	31♀ (1 wk)	0.52(0.32-0.72)	
43	242♀ (12 mo)	320.0	2	93	39♂ (2 wk)	0.80(0.35-1.35)	
44	4♂ (18 mo)	526.8	3	94	31♀ (2 wk)	0.77(0.43-1.11)	
45	233♀ (18 mo)	420.0	2	95	39♂ (4 wk)	1.30(0.52-2.08)	
46	2♂ (2 yr)	640.9	3	96	31♀ (4 wk)	1.26(0.77-1.75)	
47	215♀ (2 yr)	522.3	2	97	39♂ (6 wk)	2.05(1.24-2.86)	
48	158♀ (3 yr)	587.3	2	98	31♀ (6 wk)	1.82(1.14-2.50)	
49	110♀ (4 yr)	628.2	2	99	39♂ (8 wk)	2.95(1.62-4.28)	
50	77♀ (5 yr)	653.2	2	100	31♀ (8 wk)	2.63(1.67-3.59)	
51	53♀ (6 yr)	673.6	2	101	39♂ (10 wk)	3.81(1.95-5.67)	
52	34♀ (7 yr)	679.1	2	102	31♀ (10 wk)	3.36(2.15-4.57)	
				103	39♂ (12 wk)	4.80(2.60-7.00)	

/1/ Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction).

continued



## 28. GROWTH: MAMMALS

### Part III. BODY WEIGHT: MAMMALS OTHER THAN MAN AND RODENTS

Species	Subjects (Age)	Weight kg	Reference	Species	Subjects (Age)	Weight kg	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
<i>Canis familiaris</i> (dog)				<i>Canis familiaris</i> (dog)					
104	Beagle <sup>2</sup>	31♀ (12 wk)	4.34(2.87-5.81)	12	165	Shetland sheepdog <sup>1</sup>	14♀ (14 wk)	4.86(1.20-8.44)	12
105		39♂ (14 wk)	5.71(3.32-8.10)		166		15♂ (16 wk)	6.96(1.92-12.00)	
106		31♀ (14 wk)	5.10(3.55-6.65)		167		14♀ (16 wk)	5.67(1.48-9.86)	
107		39♂ (16 wk)	6.52(3.54-9.50)		168	Wirehaired fox terrier <sup>1</sup>	21♂ (birth)	0.19(0.10-0.28)	12
108		31♀ (16 wk)	5.75(3.64-7.86)		169		23♀ (birth)	0.19(0.14-0.24)	
109	Cocker spaniel <sup>1</sup>	31♂ (birth)	0.24(0.17-0.31)	12	170		21♂ (1 wk)	0.37(0.22-0.52)	
110		37♀ (birth)	0.24(0.15-0.33)		171		23♀ (1 wk)	0.38(0.23-0.54)	
111		31♂ (1 wk)	0.41(0.27-0.61)		172		21♂ (2 wk)	0.57(0.35-0.79)	
112		37♀ (1 wk)	0.41(0.24-0.58)		173		23♀ (2 wk)	0.56(0.35-0.77)	
113		31♂ (2 wk)	0.62(0.40-0.84)		174		21♂ (4 wk)	1.01(0.58-1.44)	
114		37♀ (2 wk)	0.63(0.41-0.85)		175		23♀ (4 wk)	0.96(0.63-1.29)	
115		31♂ (4 wk)	1.04(0.64-1.44)		176		21♂ (6 wk)	1.59(1.06-2.12)	
116		37♀ (4 wk)	1.05(0.66-1.44)		177		23♀ (6 wk)	1.48(0.92-2.04)	
117		31♂ (6 wk)	1.82(1.14-2.50)		178		21♂ (8 wk)	2.25(1.49-3.01)	
118		37♀ (6 wk)	1.74(0.91-2.57)		179		23♀ (8 wk)	2.10(1.24-2.96)	
119		31♂ (8 wk)	2.83(1.90-3.76)		180		21♂ (10 wk)	2.94(1.83-4.05)	
120		37♀ (8 wk)	2.56(1.86-3.26)		181		23♀ (10 wk)	2.71(1.58-3.84)	
121		31♂ (10 wk)	3.78(2.71-4.85)		182		21♂ (12 wk)	3.73(2.26-5.20)	
122		37♀ (10 wk)	3.39(2.50-4.28)		183		23♀ (12 wk)	3.42(2.22-4.62)	
123		31♂ (12 wk)	4.88(3.55-6.21)		184		21♂ (14 wk)	4.45(2.93-5.97)	
124		36♀ (12 wk)	4.27(3.33-5.21)		185		23♀ (14 wk)	4.02(2.66-5.38)	
125		31♂ (14 wk)	5.93(4.36-7.50)		186		21♂ (16 wk)	5.14(3.50-6.78)	
126		37♀ (14 wk)	5.08(3.90-6.26)		187		23♀ (16 wk)	4.59(3.23-5.95)	
127		31♂ (16 wk)	6.82(5.02-8.62)		<i>Capra hircus</i> (goat)				
128		37♀ (16 wk)	5.77(4.39-7.15)		188	Angora	♂ (birth)	3.03(2.08-3.97)	13
129	German shepherd <sup>2</sup>	22♂ (birth)	0.49(0.34-0.68)	8, 16	189		♀ (birth)	2.75(1.90-3.61)	
130		22♂ (1 wk)	0.87(0.57-1.02)		190		♂ (6 mo)	17.0(11.3-22.7)	
131		15♀ (1 wk)	0.50(0.34-0.64)		191		♀ (6 mo)	15.4(10.5-20.3)	
132		22♂ (2 wk)	1.43(1.14-1.70)		192		♂ (12 mo)	27.5(18.1-36.9)	
133		15♀ (2 wk)	0.89(0.57-1.02)		193		♀ (12 mo)	21.1(15.8-26.4)	
134		22♂ (4 wk)	2.95(2.39-3.52)		194		♂ (2 yr)	36.6(24.7-48.5)	
135		15♀ (4 wk)	2.02(1.48-2.27)		195		♀ (2 yr)	26.1(20.2-32.0)	
136		20♂ (6 wk)	5.00(3.86-5.91)		196		♂ (3 yr)	45.3(29.6-61.0)	
137		15♀ (6 wk)	3.77(3.30-4.09)		197		♀ (3 yr)	29.4(24.0-34.8)	
138		♂ (8 wk)	(7.0-16.0)		198		♂ (4 yr)	55.4(39.3-71.5)	
139		♀ (8 wk)	(8.0-18.0)		199		♀ (4 yr)	31.7(25.6-37.9)	
140		♂ (10 wk)	(9.5-22.5)		200		♂ (5 yr)	63.1	
141		♀ (10 wk)	(9.5-24.5)		201		♀ (5 yr)	33.2(26.5-39.9)	
142		♂ (12 wk)	(10.5-29.5)		202	British Alpine	>50♂♀ (birth)	3.75	9
143		♀ (12 wk)	(15.5-28.0)		203		>50♂♀ (1 mo)	9.55	
144		♂ (14 wk)	(12.0-37.0)		204		>50♂♀ (2 mo)	14.5	
145		♀ (14 wk)	(16.5-31.5)		205		>50♂♀ (4 mo)	25.9	
146		♂ (16 wk)	(13.5-43.0)		206		>50♂♀ (6 mo)	33.2	
147		♀ (16 wk)	(16.5-38.0)		207		>50♂♀ (12 mo)	49.1	
148	Shetland sheepdog <sup>1</sup>	15♂ (birth)	0.21(0.14-0.28)	12	208		>50♂♀ (18 mo)	62.0	
149		14♀ (birth)	0.20(0.11-0.29)		209		>50♂♀ (21 mo)	65.5	
150		15♂ (1 wk)	0.39(0.23-0.55)		210	Saanen	♂ (birth)	3.60(2.36-4.84)	4
151		14♀ (1 wk)	0.36(0.16-0.56)		211		♀ (birth)	3.14(1.64-4.64)	
152		15♂ (2 wk)	0.58(0.32-0.84)		212		♂ (1 mo)	7.17(4.75-9.59)	
153		14♀ (2 wk)	0.55(0.24-0.86)		213		♀ (1 mo)	6.71(4.11-9.31)	
154		15♂ (4 wk)	1.04(0.42-1.66)		214		♂ (2 mo)	11.3(7.5-15.1)	
155		14♀ (4 wk)	0.97(0.42-1.52)		215		♀ (2 mo)	11.0(7.3-14.7)	
156		15♂ (6 wk)	1.92(0.68-3.16)		216		♂ (3 mo)	15.0(9.2-20.7)	
157		14♀ (6 wk)	1.67(0.57-2.77)		217		♀ (3 mo)	14.6(9.1-20.1)	
158		15♂ (8 wk)	2.92(0.95-4.89)		218		♂ (6 mo)	24.6(14.9-34.5)	
159		14♀ (8 wk)	2.44(0.72-4.16)		219		♀ (6 mo)	24.5(15.4-33.5)	
160		15♂ (10 wk)	3.92(1.18-6.66)		220		♂ (9 mo)	30.6(20.8-40.3)	
161		14♀ (10 wk)	3.23(0.81-5.65)		221		♀ (9 mo)	29.9(18.9-40.8)	
162		15♂ (12 wk)	4.96(1.56-8.36)		222		♂ (12 mo)	40.7(28.1-54.3)	
163		14♀ (12 wk)	4.04(0.98-7.10)		223		♀ (12 mo)	35.3(21.6-49.1)	
164		15♂ (14 wk)	5.93(1.72-10.14)		224		♂ (18 mo)	52.2(34.7-69.6)	

/1/ Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction). /2/ Values in parentheses (column C) are ranges, estimate "c" (cf. Introduction), unless otherwise indicated.

continued



## 28. GROWTH: MAMMALS

### Part III. BODY WEIGHT: MAMMALS OTHER THAN MAN AND RODENTS

Species	Subjects (Age)	Weight kg	Reference	Species	Subjects (Age)	Weight kg	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
<i>Capra hircus</i> (goat) <sup>1</sup> Saanen	♀ (18 mo)	44.9(28.9-60.9)	4	285	<i>Felis catus</i> (cat) <sup>2</sup>	6♂ (8 wk)	0.714(0.559-0.820)	7	
	♂ (2 yr)	58.2(29.8-86.6)		286	6♀ (8 wk)	0.684(0.645-0.760)	7		
	♀ (2 yr)	53.7(34.1-73.4)		287	6♂ (9 wk)	0.811(0.589-0.963)	7		
	♂ (3 yr)	67.8(33.5-102.1)		288	6♀ (9 wk)	0.763(0.715-0.817)	7		
	♀ (3 yr)	58.4(32.9-83.9)	4	289	6♂ (10 wk)	1.006(0.872-1.159)	7		
	♂ (4 yr)	81.7		290	6♀ (10 wk)	0.891(0.790-1.032)	7		
	♀ (4 yr)	60.3(37.1-83.6)		291	6♂ (11 wk)	0.998(0.789-1.219)	7		
	♂ (5 yr)	76.7		292	6♀ (11 wk)	1.009(0.897-1.105)	7		
	♀ (5 yr)	70.1(37.5-102.7)		293	6♂ (12 wk)	1.280(1.200-1.347)	7		
	Toggenburg	♂ (birth)		3.49(2.31-4.67)	294	6♀ (12 wk)	1.011(0.902-1.216)	7	
		♀ (birth)		3.08(2.00-4.16)	295	6♂ (13 wk)	1.440(1.271-1.550)	7	
		♂ (1 mo)		6.76(4.48-9.04)	296	6♀ (13 wk)	1.202(1.024-1.361)	7	
		♀ (1 mo)		6.35(4.41-8.29)	297	52♂ (adult)	2.822(1.410-4.234) <sup>1</sup>	6	
		♂ (2 mo)		11.2(7.9-14.4)	298	52♀ (adult)	2.445(1.415-3.476) <sup>2</sup>	6	
		♀ (2 mo)		10.2(7.5-13.0)	11	299	<i>Macaca mulatta</i> (rhesus monkey) <sup>2</sup>	28♂ (birth)	0.49(0.39-0.67)
		♂ (3 mo)	15.0(10.1-19.9)	300		50♀ (birth)	0.47(0.33-0.64)		
		♀ (3 mo)	13.7(10.2-17.3)	301		28♂ (3 mo)	0.96(0.76-1.30)		
		♂ (6 mo)	23.3(16.8-29.8)	302		50♀ (3 mo)	0.92(0.54-1.16)		
		♀ (6 mo)	20.8(14.8-26.8)	303		28♂ (6 mo)	1.45(1.07-1.88)		
		♂ (9 mo)	27.2(18.7-35.7)	304		50♀ (6 mo)	1.42(0.89-1.80)		
		♀ (9 mo)	25.4(18.3-32.4)	305		28♂ (9 mo)	1.84(1.25-2.33)		
		♂ (12 mo)	34.8(25.8-43.8)	306		50♀ (9 mo)	1.82(1.18-2.28)		
		♀ (12 mo)	29.0(21.0-36.9)	307		28♂ (12 mo)	2.20(1.48-2.98)		
		♂ (18 mo)	42.6(31.9-53.3)	308		50♀ (12 mo)	2.19(1.45-2.68)		
		♀ (18 mo)	38.1(26.7-49.4)	309		22♂ (18 mo)	2.88(2.01-3.76)		
	♂ (2 yr)	47.9(30.9-65.0)	310	45♀ (18 mo)		2.83(1.86-3.44)			
	♀ (2 yr)	45.0(28.6-61.4)	311	17♂ (2 yr)		3.45(2.70-4.76)			
♂ (3 yr)	58.2(39.7-76.8)	312	43♀ (2 yr)	3.41(2.40-4.35)					
♀ (3 yr)	51.6(31.0-72.1)	313	12♂ (3 yr)	5.27(4.19-7.22)					
♂ (4 yr)	70.9(52.3-89.4)	314	34♀ (3 yr)	4.82(3.72-5.94)					
♀ (4 yr)	51.6(33.9-69.4)	315	10♂ (4 yr)	7.52(5.74-10.76)					
♂ (5 yr)	66.5(31.2-101.8)	316	31♀ (4 yr)	5.95(4.80-7.21)					
♀ (5 yr)	54.2(39.7-68.6)	317	9♂ (5 yr)	8.71(6.83-10.29)					
<i>Equus caballus</i> (horse), thoroughbred	18♂ (3.1 da)	52.45	318	28♀ (5 yr)	6.66(5.28-9.60)				
	19♀ (5.6 da)	54.32	319	7♂ (6 yr)	9.97(8.78-11.10)				
	3♂ (33.5 da)	93.89	320	25♀ (6 yr)	7.29(5.65-10.90)				
	4♀ (83 da)	116.77	321	6♂ (7 yr)	10.97(8.80-12.13)				
	8♂ (9 mo)	285.13	322	21♀ (7 yr)	8.01(6.31-12.20)				
	5♂ (yearling)	306.35	14	323	<i>Oryctolagus cuniculus</i> (European rabbit), New Zealand white <sup>2</sup>	♂♀ (birth)	0.065		
	1♀ (yearling)	354.00		324	♂♀ (7 da)	0.146			
	2♀ (12 mo)	380.11		325	♂♀ (14 da)	0.260			
	3♂ (2-3 yr)	433.92		326	♂♀ (21 da)	0.357			
	7♀ (2-3 yr)	408.50		327	♂♀ (28 da)	0.584			
11♂ <sup>3</sup> (4.3 yr)	445.76	328		♂♀ (35 da)	0.916				
<i>Felis catus</i> (cat) <sup>2</sup>	6♂ (birth)	0.098(0.083-0.107)		329	♂♀ (42 da)	1.25			
	6♀ (birth)	0.104(0.097-0.120)		330	♂♀ (49 da)	1.56			
	6♂ (1 wk)	0.129(0.083-0.196)		331	♂♀ (56 da)	1.75			
	6♀ (1 wk)	0.144(0.097-0.212)		332	♂ (8 wk)	1.95(1.60-2.30)			
	6♂ (2 wk)	0.213(0.146-0.282)	333	♀ (8 wk)	2.04(1.50-2.50)				
	6♀ (2 wk)	0.230(0.162-0.296)	334	♂ (10 wk)	2.32(2.00-2.60)				
	6♂ (3 wk)	0.297(0.259-0.365)	335	♀ (10 wk)	2.37(1.90-2.60)				
	6♀ (3 wk)	0.324(0.267-0.377)	336	♂ (12 wk)	2.67(2.30-3.00)				
	6♂ (4 wk)	0.364(0.266-0.487)	337	♀ (12 wk)	2.72(2.10-3.00)				
	6♀ (4 wk)	0.402(0.330-0.475)	338	♂ (14 wk)	2.98(2.50-3.30)				
	6♂ (5 wk)	0.446(0.346-0.578)	339	♀ (14 wk)	3.05(2.30-3.40)				
	6♀ (5 wk)	0.467(0.387-0.563)	340	♂ (16 wk)	3.13(2.60-3.50)				
	6♂ (6 wk)	0.541(0.420-0.625)	341	♀ (16 wk)	3.26(2.60-3.70)				
	6♀ (6 wk)	0.540(0.467-0.623)	342	♂ (18 wk)	3.3(2.8-3.7)				
6♂ (7 wk)	0.642(0.515-0.767)	343	♀ (18 wk)	3.49(2.90-4.00)					
6♀ (7 wk)	0.622(0.521-0.701)	344	♂ (20 wk)	3.45(2.80-3.90)					
			345	♀ (20 wk)	3.7(3.0-4.3)				

<sup>1/</sup> Values in parentheses (column C) are ranges, estimate "b" (cf. Introduction). <sup>2/</sup> Values in parentheses (column C) are ranges, estimate "c" (cf. Introduction), unless otherwise indicated. <sup>3/</sup> Geldings.

continued

## 28. GROWTH: MAMMALS

### Part III. BODY WEIGHT: MAMMALS OTHER THAN MAN AND RODENTS

Species	Subjects (Age)	Weight kg	Reference	Species	Subjects (Age)	Weight kg	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
346	<i>Oryctolagus cuniculus</i> (European rabbit), New Zealand white <sup>a</sup>	♂ (22 wk)	3.53(3.00-4.00)	14	383	<i>Ovis aries</i> (sheep) <sup>a</sup> Southdown	♀ (12 mo)	38.4; 40.8	10
347		♀ (22 wk)	3.85(3.30-4.40)		384	<i>Sus scrofa</i> (swine) <sup>b</sup> Berkshire	7 (birth)	1.84(1.34-2.17)	5
348		♂ (24 wk)	3.61(3.00-4.30)		385		7 (1 wk)	2.58(2.17-3.10)	
349		♀ (24 wk)	4.0(3.4-4.8)		386		7 (2 wk)	4.32(3.76-4.91)	
350		♂ (26 wk)	3.73(3.00-4.40)		387		7 (3 wk)	7.02(5.78-8.14)	
351		♀ (26 wk)	4.08(3.50-4.90)		388		7 (4 wk)	9.85(8.43-12.60)	
	<i>Ovis aries</i> (sheep) <sup>a</sup> Corriedale	♂ (birth)	3.7; 4.4	10	389		7 (5 wk)	14.02(12.30-17.00)	
353		♀ (birth)	3.3; 4.0		390		7 (6 wk)	18.31(16.14-21.36)	
354		♂ (3 mo)	20.5; 24.7		391		7 (7 wk)	23.16(21.02-27.04)	
355		♀ (3 mo)	17.8; 20.9		392	7 (8 wk)	27.45(24.32-31.36)		
356		♂ (6 mo)	32.9; 36.6		393	Duroc-Jersey	2 (birth)	(2.73-2.76)	5
357		♀ (6 mo)	26.9; 30.5		394		2 (1 wk)	(2.96-3.64)	
358		♂ (12 mo)	55.9; 56.0		395		2 (2 wk)	(4.75-5.74)	
359		♀ (12 mo)	43.3; 46.0		396		2 (3 wk)	(6.84-8.03)	
360	Hampshire	♂ (birth)	4.5; 5.1		397		2 (4 wk)	(8.55-9.12)	
361		♀ (birth)	4.1; 4.5		398		2 (5 wk)	(12.80-13.45)	
362		♂ (3 mo)	26.5; 30.1		399		2 (6 wk)	(16.50-17.45)	
363		♀ (3 mo)	23.1; 27.0		400		2 (7 wk)	(20.43-21.45)	
364		♂ (6 mo)	37.2; 42.2		401		2 (8 wk)	(23.86-24.55)	
365		♀ (6 mo)	32.6; 36.5		402		2 (9 wk)	(31.0-32.0)	
366		♂ (12 mo)	64.5; 66.3		403		2 (10 wk)	(36.8-38.0)	
367		♀ (12 mo)	54.2; 57.7		404		2 (11 wk)	(43.4-44.5)	
368	Shropshire	♂ (birth)	3.5; 4.4		405		2 (12 wk)	(50.0-51.5)	
369		♀ (birth)	3.2; 3.8		406	2 (13 wk)	(56.7-58.5)		
370		♂ (3 mo)	20.0; 22.4		407	Yorkshire	154 (birth)	1.23	1
371		♀ (3 mo)	17.5; 20.6		408		127 (8 wk)	11.8	
372		♂ (6 mo)	28.6; 30.4		409		191 (10 wk)	16.3	
373		♀ (6 mo)	25.7; 28.9		410		64 (12 wk)	20.4	
374	♂ (12 mo)	53.2; 55.1		411	142 (14 wk)		27.2		
375	♀ (12 mo)	42.8; 48.3		412	85 (16 wk)		35.8		
376	Southdown	♂ (birth)	3.6; 3.9		413		97 (18 wk)	46.2	
377		♀ (birth)	3.0; 3.6		414		61 (20 wk)	47.6	
378		♂ (3 mo)	19.0; 20.8		415		220 (22 wk)	65.2	
379		♀ (3 mo)	15.5; 18.4		416		80 (24 wk)	78.8	
380		♂ (6 mo)	26.4; 29.0		417	136 (26 wk)	79.3		
381		♀ (6 mo)	22.6; 25.4		418	64 (28 wk)	87.9		
382		♂ (12 mo)	43.3; 47.3						

<sup>a</sup>/ Values in parentheses (column C) are ranges, estimate "c" (cf. Introduction), unless otherwise indicated.

<sup>b</sup>/ First value (column C) is for twin-birth animals, second value is for single-birth animals.

**Contributors:** (a) Asdell, S. A., (b) Crown, R. M., and Rusoff, Louis Leon, (c) Eaton, Orson N., (d) Johnson, B. Connor, (e) Latimer, Homer B., (f) Light, Amos E., (g) Potts, Carl G., (h) Scott, J. P., (i) Shelton, Maurice, (j) Swett, Walter W., (k) Templeton, George S., (l) Van Wagenen, Gertrude, (m) Walker, Henry, (n) Weagley, John L.

**References:** [1] Crampton, E. W. 1939. *Sci. Agr.* 19:736. [2] Davis, H. P., and I. L. Hathaway. 1956. *Nebraska Univ. Agr. Expt. Sta. Res. Bull.* 179. [3] Davis, H. P., and I. L. Hathaway. 1959. *Ibid.* 189. [4] Eaton, O. N. Unpublished. U. S. Dept. of Agriculture, Beltsville, Md., 1952. [5] Johnson, B. C. Unpublished. Univ. Illinois College Agriculture, Urbana, 1952. [6] Latimer, H. B. 1936. *Am. J. Anat.* 58:329. [7] Latimer, H. B., and H. L. Ibsen. 1932. *Anat. Record* 52:1. [8] Light, A. E. Unpublished. Wellcome Research Laboratories, Tuckahoe, N. Y., 1954. [9] Plimpton, A. A. 1940. *Brit. Goat Soc. Year Book*, p. 24. [10] Potts, C. G. Unpublished. U. S. Dept. of Agriculture, Beltsville, Md., 1952. [11] Quiring, D. P. 1950. *Functional anatomy of the vertebrates*. McGraw-Hill, New York. [12] Scott, J. P. Unpublished. Roscoe B. Jackson Memorial Laboratory, Bar Harbor, Maine, 1961. [13] Shelton, M. Unpublished. Texas Agricultural Experiment Station, Sonora, 1961. [14] Templeton, G. S. Unpublished. U. S. Dept. of Agriculture, Fontana, Calif., 1955. [15] Van Wagenen, G., and H. R. Catchpole. 1956. *Am. J. Phys. Anthropol.*, N.S. 14:245. [16] Weagley, J. L. Unpublished, 1955.

## 29. GPOWTH: VERTEBRATES OTHER THAN MAMMALS

For information on organ weights of chicken, and for information on additional species of reptiles, amphibians and fishes, consult reference 2, Part I.

### Part I. BODY WEIGHT: BIRDS

	Species	Age	Weight kg		Refer- ence		Species	Age	Weight kg		Refer- ence
			Males	Females					Males	Females	
	(A)	(B)	(C)	(D)	(E)		(A)	(B)	(C)	(D)	(E)
1	<i>Anas platyrhynchos</i>	Hatched	0.059		5		<i>Gallus domesticus</i> (chicken)				
2	<i>domesticus</i> (Pekin duck) <sup>1</sup>	1 wk	0.150			60	White Leghorn	5 wk	0.354	0.367	6
3		2 wk	0.458			61		6 wk	0.449	0.436	
4		3 wk	0.744			62		7 wk	0.603	0.549	
5		4 wk	1.148			63		8 wk	0.689	0.640	
6		5 wk	1.506			64		9 wk	0.875	0.721	
7		6 wk	2.005			65		10 wk	0.944	0.776	
8		8 wk	2.758			66		12 wk	1.243	0.934	
9	<i>Anser anser</i> (gray-lag goose)	Hatched	.....	0.077	1						
10		1 wk	0.227	0.227		67		14 wk	.....	1.107	
11		2 wk	0.635	0.589		68		16 wk	.....	1.270	
12		3 wk	1.270	1.270		69		18 wk	.....	1.402	
13		4 wk	1.905	1.769		70		20 wk	.....	1.551	
14		5 wk	2.404	1.814		71	<i>Meleagris gallopavo</i> (turkey)				
15		6 wk	3.039	2.585		72	Beltville Small White	Hatched	0.045	0.045	6
16		8 wk	3.946	3.447		73		1 wk	0.095	0.086	
17		10 wk	4.264	3.719	74		2 wk	0.181	0.163		
18		12 wk	5.035	4.218	75		4 wk	0.472	0.404		
19		16 wk	5.352	4.672	76		6 wk	0.921	0.721		
20	<i>Colinus virginianus</i> (bobwhite quail) <sup>1</sup>	Hatched	0.004		3		8 wk	1.483	1.148		
21		1 wk	0.018			77		10 wk	2.205	1.674	
22		2 wk	0.027			78		12 wk	2.726	2.087	
23		3 wk	0.045			79		14 wk	3.357	2.608	
24		4 wk	0.063			80		16 wk	4.264	3.062	
25		5 wk	0.082			81		18 wk	4.704	3.357	
26		6 wk	0.095			82		20 wk	5.643	3.742	
27		8 wk	0.132			83		24 wk	7.438	4.382	
28		12 wk	0.159		84		26 wk	8.038	4.631		
29		16 wk	0.172		85		28 wk	9.008	4.740		
					86		30 wk	9.113	5.085		
30	<i>Gallus domesticus</i> (chicken) Cornish	Hatched	0.032	0.032	4	87	Broad-Breasted Bronze	Hatched	0.054	0.050	6
31		1 wk	0.059	0.059		88		1 wk	0.113	0.109	
32		2 wk	0.109	0.105		89		2 wk	0.204	0.195	
33		3 wk	0.182	0.172		90		4 wk	0.585	0.517	
34		4 wk	0.268	0.256		91		6 wk	1.252	0.998	
35		8 wk	0.727	0.636		92		8 wk	2.028	1.651	
36		12 wk	1.272	1.045		93		10 wk	2.939	2.354	
37		16 wk	1.727	1.318		94		12 wk	4.037	3.166	
38		20 wk	2.091	1.545	95		14 wk	4.922	3.715		
39	New Hampshire	Hatched	0.041	0.036	6	96		16 wk	6.214	4.604	
40		1 wk	0.086	0.082		97		18 wk	6.985	5.121	
41		2 wk	0.154	0.154		98		20 wk	8.328	5.851	
42		3 wk	0.272	0.250		99		22 wk	8.850	6.083	
43		4 wk	0.404	0.363		100		24 wk	10.614	6.836	
44		5 wk	0.563	0.504		101		26 wk	11.508	7.307	
45		6 wk	0.735	0.640		102		28 wk	12.633	7.625	
46		7 wk	0.934	0.807		103		36 wk	14.710	7.997	
47		8 wk	1.152	0.948	104		40 wk	14.814	8.437		
48		9 wk	1.325	1.107	105					7	
49		10 wk	1.628	1.284	106	Eastern wild	Hatched	0.04	0.04		
50		12 wk	1.849	1.551	107		2 wk	0.08	0.08		
51		14 wk	2.554	1.828	108		4 wk	0.28	0.25		
52		16 wk	2.994	2.019	109		6 wk	0.56	0.48		
53		18 wk	3.293	2.254	110		8 wk	0.85	0.69		
54		20 wk	3.375	2.309	111		10 wk	1.22	0.96		
55	White Leghorn	Hatched	0.036	0.036	112		12 wk	1.64	1.24		
56		1 wk	0.059	0.073	113		14 wk	2.10	1.58		
57		2 wk	0.123	0.118	114		16 wk	2.60	1.98		
58		3 wk	0.191	0.195	115		18 wk	3.32	2.52		
59		4 wk	0.268	0.272	116		20 wk	4.05	3.00		
					117		22 wk	4.62	3.32		
							24 wk	5.10	3.48		

<sup>1</sup>/ Values are for males and females combined.

continued



## 29. GROWTH: VERTEBRATES OTHER THAN MAMMALS

### Part I. BODY WEIGHT: BIRDS

	Species	Age	Weight kg		Reference		Species	Age	Weight kg		Reference
			Males	Females					Males	Females	
	(A)	(B)	(C)	(D)	(E)		(A)	(B)	(C)	(D)	(E)
118	<i>Meleagris gallopavo</i> (turkey) Eastern wild	26 wk	5.50	3.62	7	121	<i>Meleagris gallopavo</i> (turkey) Eastern wild	36 wk	6.26	3.91	7
119		28 wk	5.78	3.71		122		40 wk	6.35	3.96	
120		30 wk	5.95	3.77							

Contributors: (a) Johnson, Elton L., (b) Mosby, Henry S.

References: [1] Aitken, J. R. Unpublished. Central Experimental Farm, Ottawa, Canada, 1953. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Callenbach, E. W. Unpublished. Univ. Pennsylvania, Philadelphia, 1953. [4] Gilbreath, J. C., Jr., and C. W. Upp. 1952. Louisiana Agr. Expt. Sta. Tech. Bull. 464. [5] Heuser, G. F., et al. 1951. Poultry Sci. 30:672. [6] Johnson, E. L. Unpublished. Iowa State College, Ames, 1953. [7] Mosby, H. S., and C. O. Handley. 1943. The wild turkey in Virginia: its status, life history, and management. Virginia Commission of Game and Inland Fisheries, Richmond.

### Part II. BODY LENGTH: REPTILES AND AMPHIBIANS

Values give snout-to-vent length, unless otherwise indicated. Subjects (column B): GS = growing season. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Species (Common Name) [Location]	Subjects (Age)	Length mm	Reference		Species (Common Name) [Location]	Subjects (Age)	Length mm	Reference	
										(A)
Reptilia										
1	<i>Ancistrodon contortrix moreson</i> (northern U.S. copper-head) [Kansas]	(birth)	220(200-299)	8	29	<i>Crotalus viridis lutosus</i> (Great Basin rattle-snake) [Utah]	1♀ (3 yr)	685.8	16	
2		♂ (1 yr)	354(300-409)		30		2♂ (4 yr)	701.0(678-724)		
3		♀ (1 yr)	345(300-390)		31		2♀ (4 yr)	662.9(642-681)		
4		♂ (2 yr)	480(410-530)		32		2♂ (5 yr)	703.6(645-762)		
5		♀ (2 yr)	450(391-510)		33		2♂ (6 yr)	769.6(724-815)		
6		♂ (3 yr)	560(531-589)		34		1♀ (6 yr)	665.5		
7		♀ (3 yr)	538(511-565)		35		1♂ (8 yr)	909.3		
8		♂ (4 yr)	620(590-650)		36		2♀ (8 yr)	713.7(711-716)		
9		♀ (4 yr)	578(566-589)		37		1♂ (9 yr)	833.1		
10		♂ (5 yr)	668(651-684)		38		<i>Eumeces fasciatus</i> (five-lined skink) [Kansas]	(hatchling) (23-27)		6, 7
11		♀ (5 yr)	598(590-615)		39		1 (<2 wk)	27		
12		♂ (6 yr)	710(685-734)		40		1♂ (3 wk)	34		
13		♀ (6 yr)	626(616-635)		41		1 (1 mo)	36		
14		♂ (7 yr)	760(735-785)		42		1♂ (8.5 mo)	43		
15		♀ (7 yr)	643(636-650)		43		1♂ (9 mo)	46.5		
16		♂ (8+ yr)	>786		44		3♀ (9 mo)	49(46.0-50.5)		
17		♀ (8+ yr)	>651		45		1♂ (10 mo)	48		
18	<i>Anolis carolinensis</i> (American "chameleon") [Louisiana]	(hatchling)	(22-25)	9	46	1♀ (10 mo)	48	16		
19		(8 mo)	40		47	2♂ (11 mo)	58.7(52.5-65.0)			
20		(12 mo)	(35-45)		48	2♀ (11 mo)	52.7(51.0-54.5)			
21		♀ (18 mo)	(45-48)		49	1♂ (12 mo)	61			
22		(21 mo)	(50-52)		50	1♀ (12 mo)	59			
23		♂ (24 mo)	60		51	2♂ (13 mo)	60(56-64)			
24	<i>Crotalus viridis lutosus</i> (Great Basin rattle-snake) [Utah]	20♂ (1 yr)	457.2(365-498)	16	52	1♀ (13 mo)	64	16		
25		14♀ (1 yr)	449.5(419-503)		53	1♂ (14 mo)	66			
26		4♂ (2 yr)	556.8(492-627)		54	1♂ (21 mo)	72.5			
27		2♀ (2 yr)	553.7(530-574)		55	4♂ (22 mo)	69(67-73)			
28		6♂ (3 yr)	655.3(609-711)		56	3♀ (22 mo)	74			
				57	2♂ (24 mo)	71(70-72)				
				58	2♀ (26 mo)	71.5(69-74)				

continued



**29. GROWTH: VERTEBRATES OTHER THAN MAMMALS**

**Part II. BODY LENGTH: REPTILES AND AMPHIBIANS**

Species (Common Name) [Location]	Subjects (Age)	Length mm	Ref- er- ence	Species (Common Name) [Location]	Subjects (Age)	Length mm	Ref- er- ence	
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	
Reptilia								
59	<i>Eumeces fasci-</i>	2♂ (33 mo)	6, 7	116	<i>Terrapene or-</i>	59♀ (7 yr)	11	
60	<i>atus</i> (five-	2♀ (33 mo)		117	<i>nata</i> (ornate	30♂ (8 yr)		94(76-117)
61	lined skink)	1♂ (34 mo)		118	box turtle) <sup>1</sup>	102(82-118)		
62	[Kansas]	2♀ (34 mo)		119		102(81-125)		
63		1♀ (35 mo)		120		106(83-119)		
64		1♀ (37 mo)		121		107(89-129)		
65		2♂ (45 mo)		122		109(92-119)		
66		2♀ (45 mo)		123		111(94-135)		
67		1♂ (57 mo)		124		112(97-121)		
68		1♂ (>9 yr)		125		114(95-129)		
69	<i>Malaclemys</i>	12 (hatchling)	3	126		115(99-121)		
70	<i>terrapin pile-</i>	12 (end 1st GS)		127		118(111-131)		
71	<i>ata</i>	12 (end 2nd GS)		128	<i>Thamnophis</i>	7♀ (13 yr)	120(114-129)	
72	(Mississippi	12 (end 3rd GS)		129	<i>sirtalis</i> (com-	40♂ (newborn)	141.2(118-151)	
73	diamondback	10 (end 4th GS)		130	mon garter	38♀ (newborn)	139.5(117-151)	
74	terrapin) <sup>2</sup>	6 (end 5th GS)		131	snake)	♂ (1 yr)	350	
75	[Louisiana]	2 (end 6th GS)		132	[Michigan]	♀ (1 yr)	370	
76		57♂ (adult)		133		♂ (2 yr)	430	
77		2♀ (adult)		134		♀ (2 yr)	480	
78	<i>Natrix septem-</i>	12 (newborn)		15	135		480	
79	<i>vittata</i> (queen	12 (3 mo)	136			♂ (3 yr)	550	
80	snake) <sup>3</sup> [Ohio]	17 (1 yr)	137			♀ (3 yr)	550	
81		68♂ (>2 yr)	138			♂ (4 yr)	640	
82		58♀ (>2 yr)	139			♂ (5 yr)	580	
			140			♀ (5 yr)	670	
83	<i>Sternotherus odoratus</i> (musk turtle) <sup>3</sup>	2 broods	1, 5, 13	141		590		
	[Indiana]	(hatchling)		142		♂ (7 yr)	690	
84		2 broods (3 da)		143		♀ (7 yr)	600	
85		2 broods (5 da)		144		♂ (8 yr)	700	
86		2 broods (7 da)		145		♀ (8 yr)		
87		2 broods (9 da)						
88		2 broods (11 da)						
89		2 broods (14 da)						
90		2 broods (30 da)						
91	[Iowa]	4 (hatchling)						
92		3♀ (adult)						
93	[Michigan]	200 (hatchling)						
94		9 (6 mo)						
95		4 (1.5 yr)						
96		9 (2.5 yr)						
97		10 (3.5 yr)						
98		11 (4.5 yr)						
99		15 (5.5 yr)						
100		12 (6.5 yr)						
101		5 (7.5 yr)						
102		(8+ yr)						
103	<i>Terrapene or-</i>	46♂ (1 yr)	11	146	<i>Bufo valliceps</i>	16♂ ("August"	2	
104	<i>nata</i> (ornate	65♀ (1 yr)		147	(Mexican toad)	juvenile)		20.8(13-34)
105	box turtle) <sup>1</sup>	47♂ (2 yr)		148	[Texas]	5♀ ("August"		26(15-38)
106	[Kansas]	67♀ (2 yr)		149		juvenile)		
107		48♂ (3 yr)		150		1♀ (1 mo later)		53
108		66♀ (3 yr)		151		1♂ (2 mo later)		55
109		48♂ (4 yr)		152		8♂ (8 mo later)		68.2(61-78)
110		67♀ (4 yr)		153		5♂ (9 mo later)		72.6(64-77)
111		46♂ (5 yr)		154		2♀ (10 mo later)		94(93-95)
112		84(64-114)		155		5♂ (11 mo later)		78.2(71-86)
113		80(61-102)		156		2♀ (11 mo later)		99.5(97-102)
114		92(66-108)		157	<i>Hyla regilla</i>	6♂ (12 mo later)		79.8(75-88)
115		87(67-115)		158	(Pacific tree	3♀ (13 mo later)		82.5(70-104)
		97(70-114)		159	frog) [Oregon]	1,156 (just		13.8(12.1-15.3)
			160		transformed)			
			161		5 (2 wk)	19.8(18-21)		
			162		4 (3 wk)	22(19-25)		
			163		6 (4 wk)	20.3(17-23)		
			164		6 (5 wk)	21.9(20-24)		
			165		7 (6 wk)	21.9(20-24)		
			166		1 (7 wk)	23		
			167		1 (8 wk)	21		
			168		2♂ (9 mo)	31(29.4-32.6)		
			169	<i>Rana catesbei-</i>	38 (>2 yr)	38.6(37-40)		
			170	<i>ana</i> (American	(at transforma-	52(36-60)		
				bullfrog) <sup>2</sup>	tion)			
				[New York]	(8 mo)	55(39-64)		
					(9 mo)	62(45-70)		
					(10 mo)	73(46-82)		

1/ Plastron length. 2/ Total length. 3/ Carapace length.

continued

**29. GROWTH: VERTEBRATES OTHER THAN MAMMALS**

**Part II. BODY LENGTH: REPTILES AND AMPHIBIANS**

Species (Common Name) [Location]	Subjects (Age)	Length mm	Ref- er- ence	Species (Common Name) [Location]	Subjects (Age)	Length mm	Ref- er- ence	
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	
Amphibia				177	<i>R. pipiens</i> (leopard frog) [New York]	(at transfor- mation)	25(20.0-30.5)	14
171	<i>Rana catesbeiana</i> (American)	(11 mo)	74(47-94)	178		(1 mo)	33(28-39)	
172		(12 mo)	86(59-106)	179		(2 mo)	41(36-48)	
173	bullfrog <sup>2</sup>	(13 mo)	92(65-112)	180		(3 mo)	46(40-53)	
174	[New York]	(15 mo)	94(68-114)	181		♂ (15 mo)	(52-82)	
175		♂ (adult)	>85	182		♀ (15 mo)	(54.0-92.5)	
176		♀ (adult)	>90					

<sup>2</sup>/ Total length.

Contributor: Hardy, Ross

References: [1] Adler, K. K. 1960. Copeia, p. 156. [2] Blair, W. F. 1953. Ibid., p. 208. [3] Cagle, F. R. 1952. Ibid., p. 74. [4] Carpenter, C. C. 1952. Ibid., p. 237. [5] Dogge, C. H. 1956. Herpetologica 12:176. [6] Fitch, H. S. 1954. Univ. Kansas Publ. Museum Nat. Hist. 8(1):1. [7] Fitch, H. S. 1956. Herpetologica 12:328. [8] Fitch, H. S. 1960. Univ. Kansas Publ. Museum Nat. Hist. 13(4):85. [9] Hamlett, G. W. D. 1952. Copeia, p. 183. [10] Jameson, D. L. 1956. Ibid., p. 25. [11] Legler, J. M. 1960. Univ. Kansas Publ. Museum Nat. Hist. 11(10):529. [12] Martof, B. 1954. Copeia, p. 100. [13] Risley, P. L. 1932. Papers Mich. Acad. Sci. 17:685. [14] Ryan, R. A. 1953. Copeia, p. 73. [15] Wood, J. T., and W. E. Duellman. 1950. Am. Midland Naturalist 43:173. [16] Woodbury, A. M., F. LaM. Heyrend, and A. Call. 1951. Herpetologica 7(1):28.

**Part III. BODY LENGTH AND WEIGHT: FISHES**

Age (column B): Ages are completed years; Max. = age at maximum length and/or weight. Length measurements give total length--from tip of head (jaws closed) to tip of tail--unless otherwise indicated.

Species (Common Name)	Age	Length cm	Weight kg	Refer- ence	Species (Common Name)	Age	Length cm	Weight kg	Refer- ence		
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)		
Pisces					21	<i>Coregonus clupeaformis</i>	8 yr	58	2.27	5,19,32	
1	<i>Acipenser fulvescens</i>	1 yr	24	0.068	5,6,17,	22	(North American lake	10 yr	64	2.78	
2	(lake sturgeon) <sup>1</sup>	2 yr	30	0.136	31	23	whitefish)	Max.	71 <sup>1</sup>	4.88	
3		4 yr	45	0.39	24	<i>Cyprinus carpio</i> (carp)	1 yr	18	0.09	5,11,14	
4		6 yr	54	0.72	25		2 yr	31	0.45		
5		8 yr	60	1	26		4 yr	48	1.8		
6		10 yr	71	1.4	27		6 yr	53	2.5		
7		Max.	168	50	28		8 yr	58	3.2		
8	<i>Carassius auratus</i> (gold- fish)	1 yr	9	.....	29		10 yr	66	5.1		
9		2 yr	14	.....	30		Max.	127	37.88		
10	<i>Clupea pallasii</i> (Pacific herring) <sup>1</sup>	1 yr	6	.....	23	31	<i>Esox lucius</i> (northern pike)	1 yr	20	0.09	5,11
11		2 yr	14	.....	32		2 yr	38	0.27		
12		4 yr	21	.....	33		4 yr	61	1.1		
13		6 yr	24	.....	34		6 yr	79	2.1		
14		8 yr	26	.....	35		8 yr	97	2.95		
15		10 yr	28	.....	36		10 yr	107	4.5		
16		Max.	40	0.31	37	38	<i>Gadus morhua</i> (Atlantic cod)	1 yr	16	.....	14,27
17	<i>Coregonus clupeaformis</i>	1 yr	15	0.03	5,19,32	39		2 yr	41	.....	
18	(North American lake	2 yr	23	0.085	40		4 yr	64	.....		
19	whitefish)	4 yr	42	0.7	41		6 yr	81	.....		
20		6 yr	53	1.36	42		Max.	142	25		

<sup>1</sup>/ Fork length, measured from tip of snout to end of rays in center of caudal fin.

continued

## 29. GROWTH: VERTEBRATES OTHER THAN MAMMALS

### Part III. BODY LENGTH AND WEIGHT: FISHES

Species (Common Name)					Age	Length cm	Weight kg	Reference	Species (Common Name)					Age	Length cm	Weight kg	Reference
(A)					(B)	(C)	(D)	(E)	(A)					(B)	(C)	(D)	(E)
Pisces																	
43	<i>Ictalurus punctatus</i> (channel catfish)				1 yr	8	0.045	5,14	83	<i>Perca flavescens</i> (yellow perch)				10 yr	30	0.37	5,14,30
44					2 yr	15	0.135		84					Max.	41.9	1.913	
45					4 yr	30	0.23		85	<i>Polyodon spathula</i> (paddlefish) <sup>2</sup>				1 yr	25	0.077	1,5,10,
46					6 yr	41	0.68		86					2 yr	64	1.35	12
47					8 yr	53	1.63		87					4 yr	84	2.27	
48					10 yr	69	4.3		88					6 yr	97	3.4	
49					Max.	127	24.05		89					8 yr	102	5	
50	<i>Lepisosteus osseus</i> (longnose gar)				1 yr	16	.....	2,5,21,	90					10 yr	112	6.8	
51					2 yr	32	.....	22	91					Max.	188 <sup>3</sup>	74	
52					10 yr	102	.....		92	<i>Pomoxis annularis</i> (white crappie)				1 yr	7	0.006	5,16,26
53					Max.	160	18		93					2 yr	15	0.03	
54	<i>Lepomis macrochirus</i> (bluegill)				1 yr	5	0.005	5,13	94					4 yr	25	0.21	
55					2 yr	9	0.026		95					6 yr	32	0.45	
56					4 yr	16	0.07		96					8 yr	38	0.71	
57					6 yr	20	0.17		97					Max.	40	0.865	
58					8 yr	23	0.34		98	<i>Salmo salar</i> (Atlantic salmon) <sup>1</sup>				1 yr	4	0.011	5,7,14
59					10 yr	23	0.34		99					2 yr	10	0.033	
60					Max.	39	1.955		100					4 yr	76	4.54	
61	<i>Melanogrammus aeglefinus</i> (haddock) <sup>1</sup>				1 yr	20	.....	28,29	101					6 yr	107	16	
62					2 yr	30	0.29		102					Max.	120	47	
63					4 yr	45	0.9		103	<i>S. trutta</i> (brown trout)				1 yr	10	0.025	5,14,15
64					6 yr	55	1.53		104					2 yr	20	0.095	
65					8 yr	61	.....		105					4 yr	36	0.88	
66					Max.	90	.....		106					6 yr	56	1.8	
67	<i>Micropterus salmoides</i> (largemouth black bass)				1 yr	11	0.023	5,18	107					8 yr	64	4.26	
68					2 yr	20	0.12		108					Max.	120	18.5	
69					4 yr	34	0.57		109	<i>Salvelinus fontinalis</i> (eastern brook trout)				1 yr	10	0.025	5,8
70					6 yr	41	1.02		110					2 yr	16	0.06	
71					8 yr	46	1.36		111					4 yr	35	0.65	
72					10 yr	51	1.81		112					6 yr	53	1.59	
73					Max.	95	10.48		113					8 yr	56	.....	
74	<i>Osmerus mordax</i> (American smelt)				1 yr	14	0.023	4,5,25	114					Max.	80	6.58	
75					2 yr	18	0.036		115	<i>Thunnus thynnus</i> (bluefin tuna) <sup>1</sup>				1 yr	64	.....	20,24,
76					4 yr	25	0.11		116					2 yr	82	.....	34
77					Max.	36	0.141		117					4 yr	118	.....	
78	<i>Perca flavescens</i> (yellow perch)				1 yr	7	0.003	5,14,30	118					6 yr	153	.....	
79					2 yr	12	0.03		119					Max.	311	726	
80					4 yr	20	0.11		Agnatha								
81					6 yr	25	0.23		120	<i>Petromyzon marinus</i> (sea lamprey)				1 yr	3.8	.....	3,33
82					8 yr	27	0.285		121					2 yr	7.9	.....	
									122					4 yr	43	.....	
									123					Max.	84	1.14	

<sup>1</sup>/ Fork length, measured from tip of snout to end of rays in center of caudal fin. <sup>2</sup>/ Standard length, measured from tip of snout (upper jaw) to end of vertebral column. <sup>3</sup>/ Total length.

Contributor: Carlander, Kenneth D.

References: [1] Adams, L. A. 1942. Am. Midland Naturalist 28:617. [2] Allen, E. R. 1946. Fishes of Silver Springs, Florida. The author, Silver Springs. [3] Applegate, V. C. 1950. U. S. Fish Wildlife Serv. Spec. Sci. Rept. Fisheries 55. [4] Beckman, W. C. 1942. Copeia, p. 120. [5] Carlander, K. D. 1950. Handbook of freshwater fishery biology. W. C. Brown, Dubuque, Iowa. [6] Cuerrier, J. P. 1949. Chasse Peche (Montreal) 1:26. [7] Dixon, B. 1934. J. Conseil, Conseil Perm. Intern. Exploration Mer 9:66. [8] Eddy, S., and T. Surber. 1947. Northern fishes. Univ. Minnesota Press, Minneapolis. [9] Embury, G. C. 1915. Cornell Country Life Ser. 3:57. [10] Evermann, B. W., and E. L. Goldsborough. 1902. N. Y. State Fish Comm. Rept., 1901, p. 169. [11] Flower, S. S. 1935. Proc. Zool. Soc. London, p. 265. [12] Forbes, S. A., and R. E. Richardson. 1908. The fishes of Illinois. Natural History Survey of Illinois, Urbana. p. 16. [13] Ford, T. 1947. Alabama Conserv. 19:7. [14] Gabrielson, I. N., and F. R. LaMonte, ed. 1950. The fisherman's encyclopedia. Stackpole and Heck, New York.

continued







## 30. LIFE SPANS: ANIMALS

### Part I. VERTEBRATES

Species	Common Name	Recorded Maximum Life Span	Reference	Species	Common Name	Recorded Maximum Life Span	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
Aves				Pisces					
45	<i>Perdix perdix</i>	European partridge	>5 yr	19	77	<i>Cryptobranchus alleganiensis</i>	Hellbender	>28 yr 7 mo	17
46	<i>Phasianus colchicus</i>	Ring-necked pheasant	27 yr	14	78	<i>Hyla arborea</i>	Tree frog	>14 yr 1 mo	12,17
47	<i>Serinus canarius</i>	Canary	24 yr	26	79	<i>Necturus maculosus</i>	Mud puppy	>8 yr 10 mo	17
48	<i>Struthio camelus</i>	African ostrich	50 yr	26	80	<i>Pipa pipa</i>	Surinam toad	7 yr 10 mo <sup>a</sup>	17
49	<i>Sturnus vulgaris</i>	Starling	>15 yr 10 mo	38	81	<i>Rana catesbeiana</i>	American bullfrog	>15 yr 8 mo	17
50			8 yr <sup>a</sup>	30	82	<i>R. pipiens</i>	Leopard frog	>5 yr 11 mo	17
51	<i>Turdus migratorius</i>	American robin	12 yr 10 mo	38	83	<i>Triturus cristatus</i>	Crested newt	>4 yr 1 mo	12
52			12 yr 6 mo <sup>a</sup>	38	84	<i>T. viridescens</i>	Common newt	>2 yr 11 mo	12
52					85	<i>Xenopus laevis</i>	Clawed toad	15 yr	17
Reptilia				Pisces					
53	<i>Alligator mississippiensis</i>	American alligator	>56 yr <sup>a</sup>	18	86	<i>Acipenser fulvescens</i>	Lake sturgeon	152 yr <sup>a</sup>	2
54	<i>Ancistrodon contortrix mokesoni</i>	Northern U.S. copperhead	18 yr 6 mo <sup>a</sup>	41	87	<i>A. ruthenus</i>	Sterlet	>46 yr 1 mo	16
55	<i>Anguis fragilis</i>	Slowworm	32 yr	18	88	<i>Amia calva</i>	Bowfin	24 yr	16
56	<i>Anolis equestris</i>	Giant Cuban "chameleon"	3 yr 5 mo	6	89			30 yr <sup>a</sup>	5
57	<i>Caretta caretta</i>	Loggerhead turtle	33 yr	6	90	<i>Anguilla anguilla</i>	European freshwater eel	15 yr <sup>a</sup>	23
58	<i>Chalcides ocellatus</i>	Sand skink	9 yr 6 mo	18	91	<i>A. rostrata</i>	American freshwater eel	50 yr	11
59	<i>Chelydra serpentina</i>	Snapping turtle	20 yr	18	92	<i>Carassius auratus</i>	Goldfish	30 yr	11,16
60	<i>Coluber constrictor</i>	American black snake	5 yr 4 mo <sup>a</sup>	32	93	<i>Clupea harengus</i>	Atlantic herring	19 yr <sup>a</sup>	16
61	<i>Crotalus viridis helleri</i>	Southern Pacific rattlesnake	19 yr 5 mo	44	94	<i>Coregonus clupeaformis</i>	North American lake whitefish	12 yr	16
62	<i>Heloderma suspectum</i>	Gila monster	24 yr 7 mo	8	95	<i>Cyprinus carpio</i>	Carp	26 yr <sup>a</sup>	24
63	<i>Malaclemys terrapin centrata</i>	Southern diamondback terrapin	>21 yr <sup>a</sup>	18	96	<i>Electrophorus electricus</i>	Electric eel	47 yr	16
64	<i>Naja naja</i>	Indian cobra	12 yr 4 mo	6	97	<i>Esox lucius</i>	Northern pike	>11 yr 6 mo	11
65	<i>Natrix sipedon</i>	North American water snake	7 yr	18	98			10 yr	11
66	<i>Pseudemys scripta elegans</i>	Red-eared turtle	7 yr 1 mo	32	99			24 yr <sup>a</sup>	37
67	<i>Sceloporus graciosus</i>	Sagebrush lizard	8 yr	48	100	<i>Gadus morhua</i>	Atlantic cod	16 yr <sup>a</sup>	10
68	<i>Sphenodon punctatus</i>	Tuatara	>28 yr <sup>a</sup>	13	101	<i>Hippocampus hudsonius</i>	Atlantic sea horse	>4 yr 7 mo	11
69	<i>Sternotherus odoratus</i>	Musk turtle	53 yr 3 mo <sup>a</sup>	6	102	<i>Hippoglossus hippoglossus</i>	Atlantic halibut	40 yr <sup>a</sup>	16
70	<i>Terrapene carolina</i>	Box turtle	83-88 yr	18	103	<i>Ictalurus catus</i>	White catfish	>8 yr 1 mo	11
71	<i>Thamnophis sirtalis</i>	Common garter snake	6 yr	32	104	<i>I. punctatus</i>	Channel catfish	13 yr <sup>a</sup>	29
Amphibia				Pisces					
72	<i>Ambystoma maculatum</i>	Spotted salamander	>24 yr	17	105	<i>Lepidosiren paradoxa</i>	South American lungfish	>8 yr 3 mo	16
73	<i>A. tigrinum</i>	Tiger salamander	11 yr	17	106	<i>Lepisosteus osseus</i>	Longnose gar	24 yr	16
74	<i>Amphiuma means</i>	Two-toed amphiuma	>26 yr 9 mo	12,17	107			30 yr <sup>a</sup>	5
75	<i>Bufo americanus</i>	American toad	10-15 yr	12	108	<i>Lepomis cyanellus</i>	Green sunfish	>7 yr 6 mo <sup>a</sup>	11
76	<i>B. arenarum</i>	Sand toad	>7 yr 5 mo	12	109			9 yr <sup>a</sup>	3
					110	<i>Melanogrammus aeglefinus</i>	Haddock	15 yr <sup>a</sup>	16
					111	<i>Micropterus salmoides</i>	Largemouth black bass	>11 yr <sup>a</sup>	11
					112			16 yr <sup>a</sup>	25
					113	<i>Osmerus mordax</i>	American smelt	6 yr <sup>a</sup>	31
					114	<i>Perca fluviatilis</i>	European perch	>10 yr 8 mo	16
					115			10 yr <sup>a</sup>	43
					116	<i>Pleuronectes platessa</i>	European plaice	<30 yr <sup>a</sup>	16
					117	<i>Polyodon spathula</i>	Paddlefish	24 yr <sup>a</sup>	45
					118	<i>Pomoxis annularis</i>	White crappie	9 yr <sup>a</sup>	36
					119	<i>P. nigromaculatus</i>	Black crappie	12 yr	11

/<sup>a</sup>/ In natural habitat. /<sup>s</sup>/ Still alive at time of report.

continued

## 30. LIFE SPANS: ANIMALS

### Part I. VERTEBRATES

Species	Common Name	Recorded Maximum Life Span	Reference		Species	Common Name	Recorded Maximum Life Span	Reference	
(A)	(B)	(C)	(D)		(A)	(B)	(C)	(D)	
Pisces					Chondrichthyes				
120	<i>Protopterus annectans</i>	West African lungfish	18 yr	11,16	128	<i>Dasyatis pastinaca</i>	Stingray	>21 yr	16
121	<i>Salmo salar</i>	Atlantic salmon	13 yr <sup>a</sup>	16	129	<i>Raja maculata</i>	Skate	>5 yr 10 mo	16
122	<i>S. trutta</i>	Brown trout	10 yr	16	Agnatha				
123	<i>Salvelinus namaycush</i>	Lake trout	12 yr	16	130	<i>Lampetra fluviatilis</i>	River lamprey	<1 yr	11
124			41 yr <sup>a</sup>	47	131	<i>Petromyzon marinus</i>	Sea lamprey	7 yr <sup>a</sup>	51
125	<i>Scomber scombrus</i>	Atlantic mackerel	3-4 yr	11					
126			15 yr <sup>a</sup>	39					
127	<i>Thunnus thynnus</i>	Bluefin tuna	7 yr <sup>a</sup>	42					

<sup>a</sup>/ In natural habitat.

**Contributors:** (a) Rockstein, Morris, (b) Cole, LaMont C., (c) Manville, Richard H., (d) Tanner, Vasco M., (e) Shaw, Charles E., (f) Carlander, Kenneth D., (g) Fitch, John E.

**References:** [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Anderson, A. W., ed. 1954. Com. Fisheries Rev. 16(9):28. [3] Bailey, R. M., and K. F. Lagler. 1938. Papers Mich. Acad. Sci. 23:577. [4] Bourlière, F. 1954. The natural history of mammals. Knopf, New York. [5] Breder, C. M., Jr. 1936. Bull. N. Y. Zool. Soc. 39:116. [6] Conant, R., and R. J. Hudson. 1949. Herpetologica 5:1. [7] Crandall, L. S. Unpublished. New York Zoological Society, N. Y., 1952. [8] Crossman, A. M. 1956. Copeia (1):54. [9] Farris, E. J. 1950. The care and breeding of laboratory animals. J. Wiley, New York. [10] Fleming, A. M. 1960. J. Fisheries Res. Board Can. 17:775. [11] Flower, S. S. 1925. Proc. Zool. Soc. London, p. 247. [12] Flower, S. S. 1925. Ibid., p. 269. [13] Flower, S. S. 1925. Ibid., p. 911. [14] Flower, S. S. 1925. Ibid., p. 1365. [15] Flower, S. S. 1931. Ibid., p. 145. [16] Flower, S. S. 1935. Ibid., p. 265. [17] Flower, S. S. 1936. Ibid. (2):369. [18] Flower, S. S. 1937. Ibid., p. 1. [19] Flower, S. S. 1938. Ibid., A, 108:195. [20] Flower, S. S. 1947. Ibid. 117:680. [21] Griffith, J. Q., and E. J. Farris. 1942. The rat in laboratory investigation. J. B. Lippincott, Philadelphia. [22] Gruneberg, H. 1943. Genetics of the mouse. Cambridge Univ. Press, London. [23] Haempel, A., and E. Neresheimer. 1914. Z. Fischerei 14(4):265. [24] Hart, J. L. 1931. Contrib. Can. Biol. Fisheries 6:429. [25] Hile, R. 1931. Indiana Lakes Streams Invest. 2:9. [26] Korschelt, E. 1922. Lebensdauer Altern und Tod. G. Fischer, Jena. [27] Lankester, E. R. 1870. On comparative longevity in man and the lower animals. Macmillan, London. [28] Laws, R. M. 1955. Zoo Life 10:41. [29] Lewis, W. M. 1950. Iowa State Coll. J. Sci. 24:287. [30] Low, S. H. Unpublished. U.S. Fish and Wildlife Service, Patuxent Research Refuge Records, 1952. [31] McKenzie, R. A. 1958. J. Fisheries Res. Board Can. 15:1313. [32] Mann, W. M. 1934. Wild animals in and out of the zoo. Smithsonian Institution, New York. p. 338. [33] Manville, R. H. 1957. J. Mammal. 38:279. [34] Manville, R. H. 1958. Ibid. 39:582. [35] Manville, R. H. Unpublished. Fish and Wildlife Service, U.S. Natl. Museum, Washington, D. C., 1961. [36] Miller, R. B. 1949. Preliminary biological surveys of Alberta watersheds, 1947-49. Alberta Dept. of Lands and Forests, Edmonton. [37] Miller, R. B., and W. A. Kennedy. 1948. J. Fisheries Res. Board Can. 7:176. [38] Mitchell, P. C. 1911. Proc. Zool. Soc. London, p. 425. [39] Nedelec, C. 1958. Rev. Trav. Inst. Pech. Marit. 22(2):121. [40] Palmer, R. S. 1954. The mammal guide. Doubleday, Garden City, New York. [41] Perkins, C. B. 1955. Copeia (3):262. [42] Schaefer, M. B., and J. C. Marr. 1948. U.S. Bur. Fisheries Bull. 51:187. [43] Seemann, W. 1961. Z. Fischerei, N.F. 9(7-10):603. [44] Shaw, C. E. 1957. Copeia (4):310. [45] Shields, J. T. 1958. S. Dakota Dept. Game Fish Parks Mimeo Rept. D.-J. F-1-R-7. [46] Snell, G. D. 1941. Biology of the laboratory mouse. Blakiston, Philadelphia. [47] Sprules, W. M. 1952. J. Fisheries Res. Board Can. 9:1. [48] Stebbins, R. C. 1948. Copeia (1):20. [49] Todd, T. W. 1939.

continued

## 30. LIFE SPANS: ANIMALS

### Part I. VERTEBRATES

In E. V. Cowdry, ed. Problems of ageing. Williams and Wilkins, Baltimore, p. 71. [50] U.S. National Office of Vital Statistics. 1961. Vital statistics of the United States, 1959. U.S. Dept. of Health, Education, and Welfare, Washington, D. C. [51] Wigley, R. L. 1959. U.S. Fish Wildlife Serv. Fishery Bull. 59:561.

### Part II. INVERTEBRATES

Class and Species	Common Name	Recorded Maximum Life Span	Reference	Class and Species	Common Name	Recorded Maximum Life Span	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
Chordata									
1	Cephalochordata <sup>1</sup> <i>Branchiostoma lanceolatum</i>	Amphioxus	7 mo	6	28	Insecta <i>Tribolium</i> spp.	Flour beetle	3 yr	7
2	Ascidiacea <i>Ciona intestinalis</i>	Sea squirt	5 mo	11	29	Symphyla <i>Scutigera immaculata</i>	Garden symphylid	11-12 mo	7
Echinodermata				Annelida					
3	Asteroidea <i>Asterias rubens</i>	Starfish	>5 yr	11	30	Hirudinea <i>Hirudo medicinalis</i>	Medicinal leech	27 yr	9, 11
4	Holothuroidea <i>Cucumaria planci</i>	Sea cucumber	>10 yr	11	31	Oligochaeta <i>Lumbricus terrestris</i>	Earthworm	6 yr	19
Arthropoda				Polychaeta					
5	Arachnida <i>Dermacentor andersoni</i>	Rocky Mountain wood tick	3-4 yr	7	32	<i>Nereis irrorata</i>	Clam worm	2 yr	5
6	<i>Tegenaria derhami</i>	Spider	4 yr	15	Mollusca				
7	Crustacea <i>Astacus fluviatilis</i>	Crayfish	30 yr	11	33	Cephalopoda <i>Loligo pealeii</i>	Squid	3-4 yr <sup>2</sup>	4
8	<i>Callinectes sapidus</i>	Blue crab	3 yr <sup>2</sup>	3	34	Bivalvia <i>Mercenaria mercenaria</i>	Quahog	25-40 yr <sup>2</sup>	4
9	<i>Cyclops viridis</i>	Cyclops	10 mo <sup>3</sup>	20	35	<i>Mytilus edulis</i>	Mussel	8-10 yr	4, 17
10	<i>Daphnia magna</i>	Water flea	108 da	14	36	<i>Ostrea edulis</i>	Oyster	7-12 yr	17
11	<i>Homarus gammarus</i>	European lobster	33 yr	9	37	<i>Pecten maximus</i>	Scallop	22 yr <sup>2</sup>	4
12	Insecta <i>Aedes geniculatus</i>	Mosquito	1 yr 6 mo	2	38	<i>Teredo navalis</i>	Shipworm	2 yr <sup>2</sup>	4
13	<i>Apis mellifera</i>	Honeybee	5 yr	11	39	Gastropoda <i>Acmaea dorsuosa</i>	Limpet	15 yr <sup>2</sup>	4
14	Queen		6 mo	11	40	<i>Aplysia punctata</i>	Sea hare	1 yr <sup>2</sup>	4
15	Drone		11 mo	11	41	<i>Doris</i> spp.	Sea lemon	1 yr	11
16	Worker		6 mo	21	42	<i>Helix pomatia</i>	Land snail	18 yr <sup>4</sup>	11
17	<i>Cimex lectularius</i>	Bedbug (unfed)	6 mo	21	43	<i>Littorina littorea</i>	Periwinkle	20 yr	4, 9
18	<i>Drosophila melanogaster</i>	Fruit fly	46 da	16	44	<i>Lymnaea</i> spp.	Freshwater snail	4-5 yr	11
19	Normal		20 da	16	45	Polyplacophora <i>Ischnochiton magdalenensis</i>	Chiton	3-4 yr <sup>2</sup>	4
20	Vestigial		13 yr	11	Aschelminthes				
21	<i>Formica fusca</i>	Black ant (queen)	5 yr	11	46	Nematoda <i>Enterobius vermicularis</i>	Pinworm	2 mo	10
22	<i>F. sanguinea</i>	Red ant (worker)	2 yr	7	47	<i>Heterakis spumosa</i>	Rodent cecal worm	10 mo	18
23	<i>Lepisma saccharina</i>	Silverfish	17 yr <sup>3</sup>	11	48	<i>Loa loa</i>	African filarial worm	15 yr	18
24	<i>Magicicada septendecim</i>	Periodical cicada	8 yr	13	49	<i>Necator americanus</i>	Hookworm	12 yr	18
25	<i>Mantis religiosa</i>	Praying mantis	3 mo	7	50	<i>Rhabditis elegans</i>	Free-living roundworm	12 da	10
26	<i>Melanoplus differentialis</i>	Differential grasshopper	4-5 yr	11	51	<i>Trichinella spiralis</i>	Trichina worm	5 wk	18
27	<i>Melolontha vulgaris</i>	June beetle (larva)	76 da	9, 12	52	Adults (in guinea pig)		30 yr	9
28	<i>Musca domestica</i>	Housefly	4 yr 7 mo	8		Cysts			
29	<i>Periplaneta americana</i>	American cockroach							

<sup>1</sup>/ Subphylum. <sup>2</sup>/ In natural habitat. <sup>3</sup>/ Including developmental period. <sup>4</sup>/ Still alive at time of report.

continued



# Continails

## 30. LIFE SPANS: ANIMALS

### Part II. INVERTEBRATES

Class and Species	Common Name	Recorded Maximum Life Span	Reference	Class and Species	Common Name	Recorded Maximum Life Span	Reference	
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	
Aschelminthes				Cnidaria				
53	Nematoda <i>Wuchereria bancrofti</i>	Bancroft's filarial worm (in man)	17 yr	18	Anthozoa <i>Actinia equina</i>	Sea anemone	15 yr	9
	54	Rotifera <i>Asplanchna sieboldi</i>	Rotifer	3 wk	10	Hydrozoa <i>Hydra grisea</i>	Freshwater hydra	1-2 yr
Platyhelminthes				Porifera				
55	Cestoda <i>Diphyllobothrium latum</i>	Fish tapeworm (in man)	29 yr	18	Demospongiae <i>Hippospongia</i> spp.	Commercial sponge	50 yr	10
	56	<i>Moniezia expansa</i>	Sheep tapeworm	70 da	18	Calcarea <i>Grantia capillosa</i>	Marine sponge	3 mo
57	<i>Taenia saginata</i>	Beef tapeworm (in man)	35 yr	18	Protozoa			
58	Turbellaria <i>Planaria torva</i>	Flatworm	1 yr 2 mo	9, 11	Ciliata <i>Didinium nasutum</i>	Carnivorous ciliate (cysts)	10 yr	1

Contributors: (a) Cole, LaMont, C., (b) Rockstein, Morris, (c) Rehder, Harald A., (d) Hartman, Olga

References: [1] Brown, F. A., Jr. 1950. Selected invertebrate types. J. Wiley, New York. [2] Calvert, P. P. 1929. Proc. Am. Phil. Soc. 68:227. [3] Churchill, E. P. 1919. U.S. Bur. Fisheries Bull. 36:93. [4] Comfort, A. 1957. Proc. Malacol. Soc. (London) 32:219. [5] Durchon, M. 1948. Compt. Rend. 227:157. [6] Flower, S. S. 1925. Proc. Zool. Soc. London, p. 247. [7] Galtsoff, P. S. 1937. Culture methods of invertebrate animals. Comstock, Ithaca. [8] Haydak, M. H. Unpublished. Univ. Minnesota Experiment Station, St. Paul, 1952. [9] Heilbrunn, L. V. 1952. An outline of general physiology. W. B. Saunders, Philadelphia. [10] Hyman, L. H. 1940. The invertebrates. McGraw-Hill, New York. [11] Korschelt, E. 1922. Lebensdauer Altern und Tod. G. Fischer, Jena. [12] Korschelt, E. 1927. Tabulae Biologicae 4:346. [13] Lankester, E. K. 1870. On comparative longevity in man and the lower animals. Macmillan, London. [14] MacArthur, J. W., and W. H. T. Baillie 1929. J. Exptl. Zool. 53:221. [15] McCook, H. C. 1887. Proc. Acad. Nat. Sci. Phila. 39:369. [16] Pearl, R. 1928. Quart. Rev. Biol. 3:391. [17] Pelseneer, P. 1935. Acad. Roy. Belg. Classe Sci. Mem. Publ. Fond. Agathon de Potter 1:617. [18] Sandground, J. H. 1936. J. Parasitol. 22:464. [19] Stephenson, J. 1930. The Oligochaeta. Oxford Univ. Press, New York. [20] Walter, E. 1922. Zool. Jahrb. Abt. System. Oekol. Geog. Tiere 44:375. [21] Weismann, A. 1882. Über die Dauer des Lebens. G. Fischer, Jena.

### 31. DEVELOPMENT AND LIFE SPANS: FOREST TREES, NORTH AMERICAN

For information on additional species, consult reference 1. Values are approximate, as great variation exists within species.

Species	Common Name	Age at First Flowering yr	Trunk Diameter <sup>1</sup> at Maturity, ft		Height at Maturity, ft		Relative Growth Rate	Life Span <sup>2</sup> yr	
			Average	Maximum	Average	Maximum			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
Gymnospermae									
1	<i>Abies concolor</i>	White fir	30-40	3-4	6	120-150	200	Moderate	150-400
2	<i>Cupressus arizonica</i>	Arizona cypress	.....	1-2.5	5	50-60	90	Slow	100-300
3	<i>Juniperus virginiana</i>	Eastern red cedar	10-15	1-2	4	40-50	100	Slow	150-300
4	<i>Larix occidentalis</i>	Western larch	20-40	3-4	8	140-180	210	Slow	300-600
5	<i>Picea glauca</i>	White spruce	10-15	1.5-2	4	60-70	120	Slow	150-350

/1/ Measurements at breast height (4.5 ft). /2/ Age at natural death.

continued



## 31. DEVELOPMENT AND LIFE SPANS: FOREST TREES, NORTH AMERICAN

Species	Common Name	Age at First Flowering yr	Trunk Diameter <sup>1</sup> at Maturity, ft		Height at Maturity, ft		Relative Growth Rate	Life Span <sup>2</sup> yr	
			Average	Maximum	Average	Maximum			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
Gymnospermae									
6	<i>Pinus strobus</i>	Eastern white pine	10	2-4	6	80-120	220	Rapid	300-500
7	<i>Sequoia gigantea</i>	Giant sequoia	60	10-15	38	250-280	350	Rapid	2,000-3,000
8	<i>Taxodium distichum</i>	Bald cypress	.....	2-5	12	80-120	150	Slow	600-1,200
9	<i>Taxus brevifolia</i>	Pacific yew	.....	1-1.5	2	20-40	65	Slow	250-350
10	<i>Thuja occidentalis</i>	Northern white cedar	30	2-3	6	30-50	125	Slow	300-400
11	<i>Tsuga canadensis</i>	Eastern hemlock	30	2-3	6	60-80	160	Slow	300-600
Angiospermae									
12	<i>Acer saccharinum</i>	Silver maple	.....	2-3	7	60-80	120	Rapid	50-125
13	<i>Alnus rubra</i>	Red alder	10	1-3	5	80-100	130	Rapid	60-100
14	<i>Betula lenta</i>	Sweet birch	40	1-2	5	50-60	80	Moderate	150-250
15	<i>Carya illinoensis</i>	Pecan	10	2-4	6	90-120	180	Moderate	300
16	<i>Catalpa speciosa</i>	Northern catalpa	10	1-3	5	30-60	120	Rapid	100
17	<i>Cornus florida</i>	Flowering dogwood	5	0.5-1	1.5	20-40	50	Slow	125
18	<i>Fagus grandifolia</i>	American beech	40	1-3	4	70-100	120	Slow	300-400
19	<i>Fraxinus americana</i>	White ash	20	2-3	6	60-80	125	Rapid	260-300
20	<i>Ilex opaca</i>	American holly	5	1-2	4	40-50	140	Slow	100-150
21	<i>Juglans nigra</i>	Black walnut	12	2-3	7	50-90	150	Rapid	150-250
22	<i>Magnolia grandiflora</i>	Southern magnolia	.....	2-3	4.5	60-80	135	Moderate	80-120
23	<i>Populus tremuloides</i>	Quaking aspen	5-20	1-2	4.5	40-60	120	Very rapid	70-100
24	<i>Prunus serotina</i>	Black cherry	10-15	1.5-3	5	50-60	100	Rapid	100-200
25	<i>Quercus alba</i>	White oak	20	2.5-4	8	80-100	150	Slow	300-600
26	<i>Salix nigra</i>	Black willow	10	1-2	6	30-40	120	Rapid	50-125
27	<i>Ulmus americana</i>	American elm	15	2-4	11	80-100	120	Rapid	150-300

<sup>1</sup>/ Measurements at breast height (4.5 ft). <sup>2</sup>/ Age at natural death.

Contributors: (a) Little, Elbert L., Jr., (b) Harrar, E. S.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Collingwood, G. H., and W. D. Brush. 1947. Knowing your trees. American Forestry Association, Washington, D. C. [3] Harlow, W. M., and E. S. Harrar. 1958. Textbook of dendrology covering the important forest trees of the United States and Canada. Ed. 4. McGraw-Hill, New York. [4] Little, E. L., Jr. Unpublished. U. S. Dept. of Agriculture Forest Service, Washington, D. C., 1953. [5] Preston, R. J., Jr. 1948. North American trees (exclusive of Mexico and tropical United States). Iowa State College Press, Ames. [6] Righter, F. I. 1937. J. Forestry 37:935. [7] Sudworth, G. B. 1908. Forest trees of the Pacific slope. U.S. Dept. of Agriculture Forest Service, Washington, D. C. [8] U.S. Department of Agriculture Forest Service. 1948. U.S. Dept. Agr. Misc. Publ. 654.

## 32. LIFE SPANS: SEEDS

For information on additional species, consult reference 2, Part I.

### Part I. IN AIR-DRY STORAGE

Species	Common Name	Storage Conditions	Viability %		Life Span	Reference	
			Initial	Final			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
1	<i>Abies procera</i>	Noble fir	-5°C	...	18	5 yr	14
2	<i>Acer saccharinum</i>	Silver maple	10°C; over water	...	95	6 mo	15
3	<i>Allium cepa</i>	Garden onion	5°-10°C; 6.4% moisture content; sealed	94	89	13 yr	8
4	<i>Avena sativa</i>	Common oat	20°-30°C	98	91	13 yr	10
5	<i>Capsicum frutescens</i>	Bush red pepper	5°C; 10.4% moisture content	73	74	6 yr	4

continued

*Contrails*  
32. LIFE SPANS: SEEDS

Part I. IN AIR-DRY STORAGE

	Species	Common Name	Storage Conditions	Viability %		Life Span	Reference
				Initial	Final		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
6	<i>Daucus carota</i>	Carrot	20°-30°C; 10.7% moisture content	67	52	6 yr	4
7	<i>Fraxinus pennsylvanica</i>	Green ash	5°C; 7.6% moisture content; sealed	68	39	8 yr	7
8	<i>Glycine soja</i>	Soybean	.....	100	48	8 yr	17
9	<i>Gossypium</i> spp.	Cotton	7% moisture content; sealed	95	80	1 yr	11
10	<i>Hordeum vulgare</i>	Barley	.....	100	76	10 yr	17
11	<i>Lactuca sativa</i>	Lettuce	0°C; 50% relative humidity	98	99	8 mo	13
12	<i>Lilium regale</i>	Regal lily	5°C; 4.5% moisture content; sealed	90	94	6 yr	6
13	<i>Lycopersicon esculentum</i>	Tomato	0°C; over CaCl <sub>2</sub>	75	61	312 da	18
14	<i>Medicago sativa</i>	Alfalfa	23°C; 15% relative humidity	100	98	6.5 yr	1
15	<i>Nelumbium nelumbo</i>	Hindu lotus	.....	...	...	150 yr	19
16	<i>Phaseolus vulgaris</i>	Kidney bean	23°C; 15% relative humidity	92	23	6 yr	1
17	<i>Phleum pratense</i>	Timothy	20°-30°C	97	56	10 yr	10
18	<i>Pinus caribaea</i>	Slash pine	5°C	78	49	7 yr	3
19	<i>Populus tremuloides</i>	Quaking aspen	20°-30°C	100	45	8 wk	16
20	<i>Prunus americana</i>	American plum	.....	100	48	46 mo	12
21	<i>Solanum tuberosum</i>	Potato	0°C; sealed	45	87	12 yr	9
22	<i>Trifolium pratense</i>	Red clover	.....	...	1	100 yr	19
23	<i>Triticum aestivum</i>	Wheat	.....	100	91	10 yr	17
24	<i>Ulmus americana</i>	American elm	5°C; 7% moisture content; sealed	82	88	16 mo	5
25	<i>Zea mays</i>	Corn	.....	92	79	9 yr	17

Contributor: Steinbauer, George P.

References: [1] Akamine, E. K. 1943. Hawaii Agr. Expt. Sta. Bull. 90. [2] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Barton, L. V. 1935. Contrib. Boyce Thompson Inst. 7:379. [4] Barton, L. V. 1939. Ibid. 10:205. [5] Barton, L. V. 1939. Ibid. 10:221. [6] Barton, L. B. 1939. Ibid. 10:399. [7] Barton, L. V. 1945. Ibid. 13:427. [8] Brown, E. 1939. Science 89:292. [9] Clark, C. F. 1940. Am. Potato J. 17:147. [10] Eastham, A. 1914. Agr. Gaz. Can. 1:634. [11] Flores, F. B. 1938. Philippine J. Agr. 9:347. [12] Giersbach, J., and W. Crocker. 1932. Contrib. Boyce Thompson Inst. 4:39. [13] Griffiths, A. E. 1942. Cornell Univ. Agr. Expt. Sta. Mem. 245. [14] Isaac, L. A. 1934. Ecology 15:216. [15] Jones, H. A. 1920. Botan. Gaz. 69:127. [16] Moss, E. H. 1938. Ibid. 99:529. [17] Robertson, D. W., and A. M. Lute. 1933. J. Agr. Res. 46:455. [18] San Pedro, A. V. 1936. Philippine Agriculturist 24:649. [19] Youngman, B. J. 1951. Kew Bull. Roy. Botan. Gardens, p. 423.

Part II. UNDISTURBED IN SOIL

For additional information on life spans of seeds buried in soil, consult reference 1. All seeds were buried at a depth of eight inches.

	Species	Common Name	Viability %		Life Span yr	Reference
			Initial	Final		
	(A)	(B)	(C)	(D)	(E)	(F)
1	<i>Avena fatua</i>	Wild oat	70	9	1	3,6
2	<i>Beta vulgaris</i> <sup>1</sup>	Beet	153	1	10	3,6
3	<i>Chrysanthemum leucanthemum</i>	Oxeye daisy	96	4	30	2
4	<i>Fraxinus americana</i>	White ash	50	4	6	3,6
5	<i>Helianthus annuus</i>	Common sunflower	100	44	1	2
6	<i>Medicago sativa</i>	Alfalfa	85	1	6	3,6
7	<i>Nelumbium nelumbo</i> <sup>2</sup>	Hindu lotus	...	...	1,040	4,5
8	<i>Nicotiana tabacum</i>	Common tobacco	89	13	30	3,6

<sup>1/1</sup> Seed balls rather than seeds, which accounts for the 153% initial viability. <sup>2/2</sup> In peat bed.

continued

# Contrails

## 32. LIFE SPANS: SEEDS

### Part II. UNDISTURBED IN SOIL

	Species	Common Name	Viability %		Life Span yr	Reference
			Initial	Final		
	(A)	(B)	(C)	(D)	(E)	(F)
9	<i>Oenothera biennis</i> <sup>a</sup>	Common evening primrose	...	32	30	2, 3, 6
10	<i>Phleum pratense</i>	Timothy	...	1	21	3, 6
11	<i>Poa pratensis</i>	Kentucky bluegrass	91	1	30	3, 6
12	<i>Solanum nigrum</i>	Black nightshade	98	82	30	3, 6
13	<i>Trifolium pratense</i>	Red clover	90	1	30	3, 6

/a/ For additional data on seeds of this species buried for 70 years, consult reference 2.

Contributor: Steinbauer, George P.

References: [1] Crocker, W. 1938. Botan. Rev. 4:235. [2] Darlington, H. T. 1951. Am. J. Botany 38:379.  
 [3] Duvel, J. W. T. 1905. U.S. Dept. Agr. Bur. Plant Ind. Bull. 83. [4] Libby, W. F. 1951. Science 114:291.  
 [5] Ohga, I. 1926. Am. J. Botany 13:754. [6] Toole, E. H., and E. Brown. 1946. J. Agr. Res. 72:201.

### Part III. AT VARIOUS TEMPERATURES

Seeds were stored in sealed containers. Median life span is number of years for 50% seed survival; maximum life span is for a single seed.

	Species	Common Name	Moisture Content <sup>1</sup> %	Life Span (Years) at						Reference
				24°C		5°C		-4°C		
				Median	Maximum	Median	Maximum	Median	Maximum	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
1	<i>Abies procera</i>	Noble fir	11	<1	1	1	>10	...	>16	9
2	<i>Allium cepa</i>	Garden onion	6	11	14	...	...	>20	>28	1,3,9,10
3	<i>Capsicum frutescens</i>	Bush red pepper	5	8	12	...	...	>20	>28	1,3,9,10
4	<i>Cinchona ledgeriana</i>	Ledger-bark cinchona	6	2	4	7	8	>9	>17	7,10
5	<i>Citrus limon</i>	Lemon	56	<1	<1	>1	>1	<1	<1	5
6	<i>Daucus carota</i>	Carrot	5	16	>20	...	...	>20	>28	1,3,9,10
7	<i>Fraxinus pennsylvanica</i>	Green ash	7	2	5	8	<9	...	...	6
8	<i>Gladiolus</i> spp.	Gladiolus	8	6	10	7	8	>10	>20	9,10
9	<i>Gossypium</i> spp.	Cotton	5	1	8	>13	>13	>13	>13	9
10	<i>Lactuca sativa</i>	Lettuce	4	13	15	...	...	>20	>28	1,3,9,10
11	<i>Lilium regale</i>	Regal lily	5	8	11	13	14	>17	>17	8,9
12	<i>Lycopersicon esculentum</i>	Tomato	5	17	>20	...	...	>20	>28	1,3,9,10
13	<i>Picea abies</i>	Norway spruce	5	...	...	...	...	17	>17	2,9
14	<i>Pinus caribaea</i>	Slash pine	Air-dry	4	8	8	>8	>10	>10	2,9
15	<i>Solanum melongena</i>	Eggplant	5	18	>20	...	...	>20	>28	1,3,9,10
16	<i>Ulmus americana</i>	American elm	7	2	4	8	10	15	>15	4,9

/1/ At time of storage.

Contributor: Barton, Lela V.

References: [1] Barton, L. V. 1935. Contrib. Boyce Thompson Inst. 7:323. [2] Barton, L. V. 1935. Ibid. 7:379.  
 [3] Barton, L. V. 1939. Ibid. 10:205. [4] Barton, L. V. 1939. Ibid. 10:221. [5] Barton, L. V. 1943. Ibid. 13:47.  
 [6] Barton, L. V. 1945. Ibid. 13:427. [7] Barton, L. V. 1947. Ibid. 15:1. [8] Barton, L. V. 1948. Boyce Thompson Inst. Plant Res. Professional Paper 2(6):45. [9] Barton, L. V. 1953. Contrib. Boyce Thompson Inst. 17:87. [10] Barton, L. V. Unpublished. Boyce Thompson Institute, Yonkers, N. Y., 1961.

# Contrails

## 33. LIFE SPANS: POLLEN

For information on additional species, consult reference 1. Lyophilized (columns C and D) = freeze-drying followed by storage in nitrogen in an otherwise uncontrolled environment. **Temp.** (column C): "17.5; 21" indicates that tests were conducted at temperatures averaging 17.5°C during the winter and 21°C during the summer. **Relative Humidity** (column D): 0 indicates lack of humidity due to storage over concentrated sulfuric acid in desiccator, unless otherwise specified; A = air; A-D = air-dry; Un = uncontrolled. **Viability** (column E) was based on germination tests made on artificial media at age given in column F.

	Species	Common Name	Storage Conditions		Viability %	Life Span da	Reference
			Temp. °C	Relative Humidity %			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
Gymnospermae							
1	<i>Ginkgo biloba</i>	Ginkgo	7	0 <sup>1</sup>	35-45	730	30
2	<i>Picea abies</i>	Norway spruce	2	10-75	48	365	13
3	<i>Pinus strobus</i>	Eastern white pine	0-4	50 <sup>2</sup>	91	413	7
4	<i>Tsuga canadensis</i>	Eastern hemlock	1-16	10-50	70-90	365	28
Angiospermae (Monocotyledoneae)							
5	<i>Allium cepa</i>	Garden onion	Lyophilized		22	191	15
6	<i>Gladiolus hybrida</i>	Hybrid gladiolus	10	50	30	102 <sup>3</sup>	23
7	<i>Hordeum vulgare</i>	Barley	2.2 <sup>4</sup>	.....	9.5-40.5	19-26 <sup>5</sup>	27
8	<i>Iris graminea</i>	Grass iris	17.5; 21	0	.....	57	26
9	<i>Lilium regale</i>	Regal lily	-20	A-D	65	161	25
10	<i>Phoenix dactylifera</i>	Date palm	-13 <sup>6</sup>	.....	60-70	365 <sup>5</sup>	6
11	<i>Poa compressa</i>	Canada bluegrass	17.5; 21	0-90	.....	1	26
12	<i>Tradescantia virginiana</i>	Virginia spiderwort	17.5; 21	0	.....	40	26
13	<i>Triticum aestivum</i>	Wheat	16-18	Humid	.....	0.5	8
14	<i>Zea mays</i>	Corn	5-10	50-80	70	3 <sup>5</sup>	17
Angiospermae (Dicotyledoneae)							
15	<i>Acer sp.</i>	Maple	17-22	A-D	.....	18	12
16	<i>Alnus glutinosa</i>	European alder	17.5; 21	0	.....	53	26
17	<i>Antirrhinum majus</i>	Snapdragon	10-22	.....	Poor	670	17
18	<i>Betula lutea</i>	Yellow birch	Room	25	3	30	13
19	<i>Carya illinoensis</i>	Pecan	5	Moist	40	4	33
20	<i>Cinchona ledgeriana</i>	Ledger-bark cinchona	10	35-50	5-10	365	24
21	<i>Citrus sp.</i>	Citrus	2	25	63	550	16
22	<i>Cornus mas</i>	Cornelian cherry dogwood	17.5; 21	30	.....	74	26
23	<i>Cucumis melo</i>	Muskmelon	-18	.....	98	30	9
24	<i>Cucurbita moschata</i>	Cushaw	-17	Un	98	30 <sup>6</sup>	10
25	<i>Digitalis purpurea</i>	Common foxglove	0-4	0	.....	172	26
26	<i>Fagus sylvatica</i>	European beech	24	A-D	.....	41	18
27	<i>Fragaria spp.</i>	Strawberry	24	A-D	.....	>16	5
28	<i>Gossypium barbadense</i>	Pima cotton	4.4-10.0	.....	64	4 <sup>5</sup>	11
29	<i>Ipomoea batatas</i>	Sweet potato	Lyophilized		76 <sup>7</sup>	354	15
30	<i>Juglans sieboldiana</i>	Siebold walnut	0	40-60	12	253	4
31	<i>Lycopersicon esculentum</i>	Tomato	-190	.....	.....	1,095	31
32	<i>Malus pumila</i>	Common apple	2-8	50	20	1,460	20
33	<i>Medicago sativa</i>	Alfalfa	Room	0 <sup>1</sup>	.....	180 <sup>5</sup>	2
34	<i>Nicotiana sylvestris</i>	Tobacco	17-22	.....	.....	205 <sup>8</sup>	12
35	<i>Oenothera biennis</i>	Common evening primrose	17.5; 21	A; 30	.....	8	26
36	<i>Persea americana</i> <sup>9</sup>	American avocado	15	0 <sup>1</sup>	.....	153	29
37	<i>Pisum sativum</i>	Garden pea	-5	35	.....	450	32
38	<i>Populus suaveolens</i>	Mongolian poplar	-3 to +3	A-D	.....	45	3
39	<i>Prunus amygdalus</i>	Almond	-18	.....	24	1,130	9
40	<i>P. domestica</i>	Garden plum	2-8	50	20	1,278	20
41	<i>P. persica</i>	Peach	2-8	50	1-20	1,095	20
42	<i>Pyrus communis</i>	Pear	2-8	50	20	1,278	20
43	<i>Quercus coccinea</i>	Scarlet oak	2	25-35	46	365	13
44	<i>Rhododendron spp.</i>	Rhododendron	-190	.....	.....	730-1,095	31
45	<i>Ribes glutinosum</i>	Nutmeg currant	17-22	0-27.2	.....	117	12
46	<i>Salix gracistyla</i>	Big catkin willow	10	A-D	1	105	21
47	<i>Solanum tuberosum</i>	Potato	-30 to -20	Un	.....	365 <sup>5</sup>	14

<sup>1</sup>/ Storage over calcium chloride in desiccator. <sup>2</sup>/ Humidified, at 75% relative humidity and 4°C, for 12 hours after storage. <sup>3</sup>/ Data recorded on the basis of seed set per capsule. <sup>4</sup>/ Flower spike or flower cut in early morning and kept under refrigeration. <sup>5</sup>/ Data recorded on the basis of seed or fruit set. <sup>6</sup>/ Pollen mixed with diluent and then stored in sealed or stoppered vials. <sup>7</sup>/ Peroxidase test. <sup>8</sup>/ Pollen sealed in ampules with CO<sub>2</sub> and stored under reduced pressure. <sup>9</sup>/ Horticultural varieties.

continued



## 33. LIFE SPANS: POLLEN

Species	Common Name	Storage Conditions		Viability %	Life Span da	Reference	
		Temp. °C	Relative Humidity %				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Angiospermae (Dicotyledoneae)							
48	<i>Trifolium hybridum</i>	Alsike clover	24	A-D	.....	12	19
49	<i>Vicia faba</i>	Broad bean	17.5; 21	0	.....	21	26
50	<i>Vitis</i> spp. <sup>9</sup>	Grape	-12	28	21	1,460	22

<sup>9</sup> / Horticultural varieties.

Contributors: (a) Pfeiffer, Norma E., (b) Hesse, Claron O.

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1962. Growth, including reproduction and morphological development. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Anonymous. 1960. Crops Soils 13(1):19. [3] Bogdanov, P. L. 1935. Sov. Botan. 1:98. [4] Cox, L. G. 1943. Northern Nut Growers Assoc. Ann. Rept. 34:58. [5] Crandall, C. S. 1912. Proc. Am. Soc. Hort. Sci. 9:121. [6] Crawford, C. L. 1937. Ibid. 35:91. [7] Duffield, J. W., and A. G. Snow, Jr. 1941. Am. J. Botany 28:175. [8] Firbas, H. 1922. Z. Pflanzenzuecht. 8:70. [9] Griggs, W. H., G. H. Vansell, and B. T. Iwakiri. 1953. Calif. Agr. 7:12. [10] Griggs, W. H., G. H. Vansell, and J. F. Reinhardt. 1950. J. Econ. Entomol. 43:549. [11] Harrison, G. J., and H. J. Fulton. 1934. J. Agr. Res. 49:891. [12] Holman, R. M., and F. Brubaker. 1926. Univ. Calif. (Berkeley) Publ. Botany 13:179. [13] Johnson, L. P. V. 1943. Can. J. Res., C, 21:332. [14] King, J. R. 1955. Am. Potato J. 32:460. [15] King, J. R. 1961. Econ. Botany 15(1):91. [16] King, J. R., and C. O. Hesse. 1938. Proc. Am. Soc. Hort. Sci. 36:310. [17] Knowlton, H. E. 1922. Cornell Univ. Agr. Expt. Sta. Mem. 52:747. [18] Mangin, L. 1886. Bull. Soc. Botan. France 33:337. [19] Molisch, H. 1893. Sitzber. Akad. Wiss. Wien Math. Naturw. Kl., I, 102:423. [20] Nebel, B. R. 1939. Proc. Am. Soc. Hort. Sci. 37:130. [21] Nohara, S. 1922. Japan. J. Botany 1:1. [22] Olmo, H. P. 1942. Proc. Am. Soc. Hort. Sci. 41:219. [23] Pfeiffer, N. E. 1939. Contrib. Boyce Thompson Inst. 10:429. [24] Pfeiffer, N. E. 1944. Ibid. 13:281. [25] Pfeiffer, N. E. 1955. Ibid. 18:153. [26] Pfundt, M. 1909. Jahrb. Wiss. Botan. 47:1. [27] Pope, M. M. 1939. J. Agr. Res. 59:453. [28] Santamour, F. S., Jr., and H. Nienstaedt. 1956. J. Forestry 54:269. [29] Schroeder, C. A. 1942. Proc. Am. Soc. Hort. Sci. 41:181. [30] Tulecke, W. R. 1954. Bull. Torrey Botan. Club 81:509. [31] Visser, T. 1955. Mededel. Landbouwhogeschool Wageningen 55:1. [32] Warnock, S. J., and D. J. Hagedorn. 1956. Agron. J. 48:347. [33] Woodroof, J. G. 1930. J. Agr. Res. 40:1059.

## 34. GROWTH RATES: PLANT TISSUES

Nutrient fluids used in plant tissue cultures are chiefly composed of chemicals. **Culture Medium** (column C): GA = Gautheret's agar; CM = coconut milk (liquid endosperm of *Cocos nucifera*); IAA = indoleacetic acid; HA = Hildebrandt's agar; WL = White's liquid; WA = White's agar; WAS = modified White's agar for sunflower tissues; HAS = modified Hildebrandt's agar for sunflower tissues; WAT = modified White's agar for tobacco tissues; BL = Bonner's liquid; BNA = Burkholder-Nickell agar. **Relative Increase** (column F):  $W_1/W_0$  = final weight divided by initial weight. **Relative Growth Rate** (column G): r = instantaneous growth rate expressed as percent increase/day.

Species (Common Name)	Tissue	Culture Medium	Growth Period da	Initial Weight mg	Relative Increase $W_1/W_0$	Relative Growth Rate 100 r	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 <i>Amorphophallus rivieri</i> (devil's-tongue)	Tuber	GA + 15% CM	30	250	8	6.92	13
2 <i>Brassica campestris</i> (bird rape)	Root	GA	60	100	4.1	2.35 <sup>1</sup>	6
3		GA + 0.3 mg/liter IAA	60	100	5.6	2.87	

<sup>1</sup> / Growth had ceased by fourth subculture, but tissue survived for another year.

continued

## 34. GROWTH RATES: PLANT TISSUES

	Species (Common Name)	Tissue	Culture Medium	Growth Period da	Initial Weight mg	Relative Increase W <sub>1</sub> /W <sub>0</sub>	Relative Growth Rate 100 r	Refer- ence			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)			
4	<i>Chrysanthemum frutescens</i> (marguerite chrysanthemum)	Crown gall	HA	42	35	3.71	3.12	9			
5	<i>Cichorium intybus</i> (chicory)	Tuber	GA	60	100	3.7	2.18	6			
6			GA + 0.3 mg/liter IAA	60	100	5.9	2.96				
7	<i>Daucus carota</i> (carrot)	Root	WL	21	15	11.1	11.4	12			
8			Root cambium	GA + 0.1 mg/liter IAA	18	125	3.56	7.04	4		
9					304	0.5	300,000	4.14	5		
10					0-7	170	2	9.9	7		
11					7-15	340	1.61	6	7		
12					15-21	548	1.72	9	7		
13					21-41	941	1.97	3.3	7		
14					41-67	1,852	2.13	2.7	7		
15					67	170	20.32	4.5	7		
16					Vascular cambium	GA	60	100	5	2.68	6
17							GA + 0.3 mg/liter IAA	60	100	7.1	
18			Root phloem	WL + 15% CM	0-2	3.98	1.15	7	2		
19					2-4	4.6	1.2	9.1			
20					4-6	5.55	1.58	22.9			
21					6-8	8.75	1.72	27.1			
22	8-10	15.05			1.98	34.2					
23	10-12	29.8			1.9	32.1					
24	12-14	56.6			1.63	22.4					
25	14-16	92.15			1.56	22.2					
26	16-20	139			1.41	8.6					
27	20-24	196.9			1.19	4.3					
28		24	3.98	59	16.98						
29	<i>Helianthus annuus</i> (common sunflower)	Crown gall <sup>2</sup>	WA <sup>3</sup>	42	25	15.6	6.54	10			
30			WA	42	25	5.2	3.92	10			
31			WAS	42	25	13.6	6.21	10			
32			HAS <sup>4</sup>	42	25	80	10.41	20			
33	<i>H. tuberosus</i> (Jerusalem artichoke sunflower)	Tuber	GA	60	100	1.95 <sup>5</sup>	1.11	6			
34			GA + 0.3 mg/liter IAA	60	100	5.4	2.81	6			
35			GA	35	283	1.08	0.22	3			
36			GA + 0.3 mg/liter IAA	35	247	1.86	1.77	3			
37			GA + 100% CM	35	237	4.36	4.21	3			
38		GA + 25% CM	35	245	3.03	3.17	3				
39		Crown gall <sup>2</sup>	GA	60	100	5.1	2.72	6			
40	GA + 0.3 mg/liter IAA			60	100	4.5	2.51				
41	<i>Nicotiana glauca</i> x <i>N. langsdorffii</i> <sup>6</sup> (tobacco)	Stem	WA	35	19.1	6.1	5.17	1			
42				42	6.8	14.2	6.32				
43			WAT	42	25	4.84	3.75	11			
44				42	25	7.80	4.89				
45	<i>Pisum sativum</i> (garden pea)	Root callus	BL + 1 g/liter yeast extract + 1 x 10 <sup>-6</sup> M 2,4-D	56	90	10.9	4.26	23			
46	<i>Rosa</i> sp. (rose)	Stem	WL + 6 mg/liter 2,4-D + 0.1% each yeast and malt extracts	14	.....	19.9	19.3	18			
47	<i>Rumex acetosa</i> (garden sorrel)	Root tumor <sup>7</sup>	BNA + 0.4 mg/liter thiamine	21	.....	12	11.8	16			
48			BNA	25-34	.....	6.0	19.9	17			
49				40	.....	16	6.92				
50			BNA + 30% CM	21	.....	11.1	11.5	15			
51	<i>Scorzonera hispanica</i> (black salsify ser-pentroot)	Root	GA	60	100	2.53	1.57	8			
52			GA + 1.0 mg/liter IAA	60	100	10.86	3.98				
53		Crown gall <sup>2</sup>	GA	60	100	8.85	3.63	8			
54				GA + 1.0 mg/liter IAA	60	100	9.20		3.70		
55	<i>Solanum tuberosum</i> (potato)	Tuber	WA + 6% CM and 18 mg/liter 2,4-D	35	3	54.7	11.4	21			
56	<i>Tagetes erecta</i> (Aztec marigold)	Stem	HAS	35	20	55	11.44	14			
57			HAS	42	35	25	7.65	9			
58		Crown gall	HAS + 0.5% dulcitol	42	35	28.7	7.98				
59				HAS + 0.5% methanol	42	35	30.6	8.14			

<sup>2</sup>/ Free from inducing microorganism (*Agrobacterium tumefaciens*). <sup>3</sup>/ Optimal growth at sucrose concentration of 1% at pH 5, 26°C. <sup>4</sup>/ Nitrate present as optimum of 0.016 M instead of 0.0038 M. <sup>5</sup>/ Growth during first subculture; dead after third subculture. <sup>6</sup>/ One of a number of tobacco hybrids forming spontaneous tumors at certain stages of development. <sup>7</sup>/ Wound tumor disease of sorrel and other plants, resulting from infection by *Aureogenus magnivena*.

continued

**34. GROWTH RATES: PLANT TISSUES**

	Species (Common Name)	Tissue	Culture Medium	Growth Period da	Initial Weight mg	Relative Increase W <sub>1</sub> /W <sub>0</sub>	Relative Growth Rate 100 r	Ref- er- ence
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
60	<i>Taxus brevifolia</i> (Pacific yew)	Pollen	WA + 15% CM and 0.6 ppm 2,4-D	28		3	3.92	24
61	<i>Vinca rosea</i> (Madagas- car periwinkle)	Crown gall	HA	42	35	12.7	6.05	9
62	<i>Vitis vinifera</i> (Euro- pean grape)	Stem	HA <sup>a</sup>	42	30	40	8.78	19
63		Gall <sup>a</sup>		42	30	30	8.10	
64	<i>Zea mays</i> (corn)	Endosperm	WA + 1.5 x 10 <sup>-2</sup> M asparagine	25	120	3.6	5.12	22

<sup>a</sup>/ Plus 3.0 g/liter casein hydrolysate, 0.1 mg/liter each NAA and kinetin, 40 mg/liter adenine. <sup>s</sup>/ Induced by *Phylloxera vestatrix*.

*Contributor:* Caplin, Samuel M.

*References:* [1] Caplin, S. M. 1947. *Botan. Gaz.* 108:379. [2] Caplin, S. M., and F. C. Steward. 1949. *Nature* 163:920. [3] Duhamet, L., and J. Magrou. 1949. *Compt. Rend.* 229:1353. [4] Gautheret, R. J. 1939. *Ibid.* 208:1340. [5] Gautheret, R. J. 1942. *Titres et travaux scientifiques.* Jouve, Paris. [6] Gautheret, R. J. 1947. *Compt. Rend. Soc. Biol.* 141:475. [7] Gautheret, R. J. 1947. *Rev. Gen. Botan.* 54:5. [8] Gautheret, R. J. 1948. *Compt. Rend. Soc. Biol.* 142:774. [9] Hildebrandt, A. C., and A. J. Riker. 1949. *Am. J. Botany* 36:74. [10] Hildebrandt, A. C., A. J. Riker, and B. M. Duggar. 1945. *Ibid.* 32:357. [11] Hildebrandt, A. C., A. J. Riker, and B. M. Duggar. 1946. *Ibid.* 33:591. [12] Melchers, G., and U. Engelmann. 1955. *Max Planck Inst. Publ. Biol. (Tuebingen)* 20:564. [13] Morel, G. 1950. *Compt. Rend.* 230:1099. [14] Muir, W. H., A. C. Hildebrandt, and A. J. Riker. 1958. *Am. J. Botany* 45:589. [15] Nickell, L. G. 1950. *Botan. Gaz.* 112:225. [16] Nickell, L. G. 1952. *Bull. Torrey Botan. Club* 79:427. [17] Nickell, L. G., and P. R. Burkholder. 1949. *Am. J. Botany* 37:538. [18] Nickell, L. G., and W. Tulecke. 1959. *Science* 130:863. [19] Pelet, F., et al. 1960. *Am. J. Botany* 47:186. [20] Riker, A. J., and A. E. Gutsche. 1948. *Ibid.* 35:227. [21] Steward, F. C., and S. M. Caplin. 1951. *Science* 113:518. [22] Straus, J. 1960. *Am. J. Botany* 47:641. [23] Torrey, J. G., and Y. Shigemura. 1957. *Ibid.* 44:325. [24] Tulecke, W. 1959. *Bull. Torrey Botan. Club* 86:283.

# *Contrails*




# Contrails

## IV. MORPHOLOGY

### 35. BODY COMPOSITION WITH INCREASING WEIGHT AND AGE: MAN

Method of determination: *Interstitial fluid* calculated by subtracting plasma volume from thiocyanate diffusion space.<sup>1</sup> *Intracellular fluid* calculated by subtracting thiocyanate diffusion space<sup>2</sup> from total body water. *Intercellular solids*<sup>2</sup> estimated by the method of McCance [2] (intracellular water = 67% intracellular mass). *Bone*<sup>3</sup> estimated from the data of Iob and Swanson [1], Mitchell [3], Shohl [4], and Widdowson [5]. *Fat*<sup>2</sup> estimated by difference between total body weight and all other components.



[1] The use of thiocyanate diffusion space as a measure of extracellular fluid is based on the probability that in normal persons the thiocyanate and other similar diffusion spaces are in fairly constant proportion in that nebulous entity, extracellular fluid, and that therefore the rate of change of the curves should not vary (although the absolute values may). [2] Because of the lack of data, values indicate order of magnitude only.

*Contributors:* Henschel, Austin; Bass, David E.; and Wedgwood, Ralph J.

*References:* [1] Iob, V., and W. W. Swanson. 1934. *Am. J. Diseases Children* 47:302. [2] Ling, W. S. M., and H. Sprinz. 1948. *Am. J. Med. Sci.* 215:555. [3] Mitchell, H. H., T. S. Hamilton, F. R. Steggerda, and H. W. Bean. 1945. *J. Biol. Chem.* 158:625. [4] Shohl, A. T. 1939. *Mineral metabolism*. Reinhold, New York. [5] Widdowson, E. M., R. A. McCance, and C. M. Spray. 1951. *Clin. Sci.* 10:113.

## 36. BODY SURFACE AREA: MAMMALS

### Part I. SURFACE AREA FOR KNOWN WEIGHT AND HEIGHT: MAN

Values are square meters of body surface area and were derived by the Sendroy and Cecchini method.

	Height cm	Weight, kg																		
		5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95
1	20	0.18																		
2	30	0.20	0.35																	
3	40	0.23	0.36																	
4	50	0.26	0.38																	
5	60	0.29	0.41	0.54																
6	70	0.33	0.44	0.57																
7	80	0.37	0.48	0.60	0.68															
8	90	0.42	0.52	0.63	0.72	0.80														
9	100	0.48	0.57	0.67	0.76	0.84	0.92													
10	110	0.55	0.64	0.72	0.80	0.88	0.96	1.04	1.11											
11	120	0.62	0.69	0.77	0.85	0.93	1.01	1.08	1.15	1.23	1.30	1.37	1.44							
12	130		0.76	0.83	0.91	0.98	1.05	1.12	1.20	1.27	1.34	1.42	1.48	1.54	1.61	1.68	1.74	1.81	1.87	
13	140			0.89	0.97	1.03	1.10	1.17	1.25	1.32	1.39	1.46	1.52	1.58	1.65	1.72	1.78	1.84	1.90	1.97
14	150				1.03	1.09	1.16	1.23	1.30	1.37	1.44	1.50	1.57	1.63	1.70	1.76	1.82	1.88	1.94	2.01
15	160					1.15	1.22	1.29	1.36	1.43	1.49	1.55	1.62	1.68	1.75	1.81	1.86	1.92	1.98	2.05
16	170						1.28	1.35	1.42	1.48	1.54	1.61	1.67	1.73	1.80	1.86	1.91	1.97	2.03	2.09
17	180							1.42	1.48	1.54	1.60	1.67	1.73	1.79	1.85	1.91	1.96	2.02	2.08	2.14
18	190								1.55	1.61	1.67	1.73	1.79	1.85	1.91	1.96	2.02	2.07	2.13	2.18
19	200									1.74	1.80	1.85	1.91	1.96	2.02	2.07	2.13	2.18	2.24	2.24
20	210										1.92	1.97	2.02	2.07	2.13	2.18	2.24	2.30	2.36	2.30
21	220											2.08	2.13	2.18	2.24	2.30	2.36	2.42	2.48	2.36
22	230												2.25	2.31	2.36	2.42	2.48	2.54	2.60	2.42
23	240																			2.48

	Height cm	Weight, kg																		
		100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	
13	140	2.03	2.10	2.17	2.23															
14	150	2.07	2.14	2.21	2.27	2.33														
15	160	2.12	2.18	2.24	2.30	2.36	2.42	2.47	2.53	2.58	2.63	2.69	2.74	2.80	2.86	2.91	2.96	3.01	3.06	
16	170	2.16	2.22	2.28	2.33	2.39	2.45	2.51	2.56	2.62	2.67	2.73	2.78	2.83	2.89	2.94	2.99	3.04	3.09	
17	180	2.20	2.26	2.32	2.38	2.43	2.49	2.54	2.60	2.66	2.71	2.77	2.83	2.88	2.93	2.98	3.03	3.08	3.08	
18	190	2.24	2.31	2.36	2.42	2.48	2.53	2.59	2.64	2.70	2.75	2.81	2.87	2.92	2.97	3.03	3.08	3.08	3.08	
19	200	2.30	2.35	2.41	2.47	2.53	2.58	2.63	2.69	2.74	2.80	2.86	2.92	2.97	3.02	3.07	3.07	3.07	3.07	
20	210	2.35	2.41	2.47	2.53	2.58	2.63	2.68	2.74	2.80	2.86	2.92	2.97	3.02	3.07	3.07	3.07	3.07	3.07	
21	220	2.41	2.47	2.53	2.58	2.63	2.69	2.75	2.81	2.87	2.92	2.97	3.03	3.08	3.08	3.08	3.08	3.08	3.08	
22	230	2.47	2.53	2.58	2.64	2.70	2.76	2.82	2.87	2.93	2.98	3.03	3.08	3.08	3.08	3.08	3.08	3.08	3.08	
23	240	2.54	2.60	2.65	2.71	2.77	2.83	2.88	2.93	2.98	3.04	3.09	3.09	3.09	3.09	3.09	3.09	3.09	3.09	
24	250			2.73	2.78	2.84	2.90	2.95	3.00	3.06	3.11	3.16	3.21	3.26	3.31	3.36	3.41	3.46	3.51	
25	260				2.93	2.97	3.02	3.07	3.12	3.17	3.22	3.27	3.32	3.37	3.42	3.47	3.52	3.57	3.62	

Contributor: Sendroy, Julius, Jr.

Reference: Sendroy, J., Jr., and L. P. Cecchini. 1954. J. Appl. Physiol. 7:1.

### Part II. CONSTANTS FOR USE IN SURFACE AREA FORMULA: MAMMALS

K-values were derived from surface area values taken from extensive literature sources, using the formula

$$K = \frac{\text{area (sq cm)}}{\text{weight}^{2/3} \text{ (g)}} \quad \text{Method (column D): } P = \text{perimeter; } S = \text{skinning; } T = \text{triangulation; } M = \text{mold; } C = \text{paper cover;}$$

I = surface integrator. Values in parentheses are ranges, estimate "c" for body weight (column E) and "d" for K-value (column F) (cf. Introduction).

	Species	Common Name	Subjects no.	Method	Body Weight g	K-Value Constant	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	<i>Balaenoptera physalus</i>	Finback whale	3	P	160,000(115,000-220,000)	8.3(7.5-8.9)	19
2			1	P	43,000,000	11.1	19

continued

**36. BODY SURFACE AREA: MAMMALS**

**Part II. CONSTANTS FOR USE IN SURFACE AREA FORMULA: MAMMALS**

	Species	Common Name	Subjects no.	Method	Body Weight g	K-Value Constant	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
3	<i>Bos taurus</i> (Hereford-Shorthorn)	Cattle	15	S	476,000(208,000-762,000)	9.3(8.1-10.8)	25
4			15 <sup>1</sup>	S	375,000(163,000-641,000)	11.0(9.0-13.8)	25
5			10 <sup>1,2</sup>	S	241,000(89,000-407,000)	9.9(9.3-10.5)	28
6			11 <sup>1,3</sup>	S	315,000(78,000-493,000)	9.4(8.8-10.0)	28
7			7 <sup>2,4</sup>	S	695,000(476,000-815,000)	7.6(7.3-7.9)	28
8	<i>Canis familiaris</i>	Dog	6	S	1,070(130-3,650)	10.1(9.3-11.0)	27
9			1	S	1,080	11.0	10
10			8	S & P	12,700(3,200-29,800)	11.6(10.2-12.5)	23
11			2	T	9,500(8,900-10,100)	9.9(9.85-9.90)	6
12			7	M	14,310(3,390-32,640)	11.2(10.3-12.1)	5
13			1	C	27,000	12.3	12
14	<i>Capra hircus</i>	Goat	1	T	15,100	10.5	6
15	<i>Cavia</i> spp.	Guinea pig	6	S	206(123-269)	9.5(8.4-10.8)	21
16			3	S	157(123-191)	10.4(10.1-10.8)	21
17			3	S	256(235-269)	8.6(8.4-8.9)	21
18			3	S	373(148-650)	9.6(9.0-9.9)	8
19			13 <sup>5</sup>	S	323(160-810)	8.9(7.9-9.6)	14
20			2	T	400(380-420)	7.1	6
21	<i>Didelphis</i> sp.	Opossum	4	S	1,200(100-1,300)	11.3(10.5-11.8)	11
22	<i>Equus caballus</i>	Horse	8	S	(47,000-555,000)	10.5	26
23			11	I	(70,000-750,000)	(8.2-10.3)	3
24	<i>Erinaceus europaeus</i>	European hedgehog	1	S	200	7.5	10
25	<i>Felis catus</i>	Cat	3	S	708(219-1,389)	10.7(9.5-11.9)	27
26			2	S	100(84-116)	10.0(9.9-10.0)	27
27			2	T	1,550(1,500-1,600)	8.7(8.6-8.9)	6
28	<i>Macaca mulatta</i>	Rhesus monkey	6	M	2,670(800-6,600)	11.8(10.8-13.2)	2
29	<i>Mus musculus</i> (albino)	House mouse	12	S	16(10-22)	11.4(9.7-13.3)	21
30			11	S	15(6-27)	7.9	9
31			3	S	16(11-20)	10.5(10.4-10.5)	8
32			64 <sup>5</sup>	S	13	6.9	21,24
33			13	M	(16-25)	9.0(8.4-9.4)	1
34	<i>Myotis lucifugus</i>	Little brown bat	2	S	8.3(5.0-11.6)	44.5(44.0-45.0)	20
35	<i>Oryctolagus cuniculus</i>	European rabbit	3 <sup>5</sup>	S	32(26-40)	8.5	9
36			3 <sup>5</sup>	S	560(70-925)	9.7	9
37			2	T	1,130(1,120-1,140)	10.0(9.0-11.0)	6
38	<i>Ovis aries</i>	Sheep	8	S	(21,800-29,100)	10.7	16
39			15	S	(3,780-50,400)	9.1	16
40			14	S	(23,600-37,700)	8.5	18
41			115	I	(2,200-68,000)	8.3	22
42	<i>Rattus norvegicus</i> (albino)	Norway rat	62	S	176(25-461)	11.4(9.6-13.0)	4
43			22	S	197(65-335)	10.5(9.0-12.7)	17
44			14	S	133(70-310)	11.6(10.9-12.1)	2
45			5	S	80(50-129)	9.9(9.6-10.4)	13
46			5	S	42(35-53)	10.5(10.1-10.8)	2
47			2	T	170(164-177)	7.15	6
48			72	M	(19-418)	9.0	15
49			56	M	125(24-366)	7.5(6.6-8.3)	7
50	14 <sup>5</sup>	M	95(22-164)	7.6(7.3-8.8)	7		
51	<i>Sorex cinereus</i>	Gray shrew	1	S	3.5	8.0	20
52	<i>Sus scrofa</i>	Swine	7	S	48,300(1,100-123,000)	9.9(8.6-12.4)	25
53			16	I	(25,000-330,000)	9.0	3
54			1	T	40,110	15.3	6

/1/ Empty weight. /2/ Thin. /3/ Medium. /4/ Fat. /5/ Starved. /6/ Surface area of one side of ear only.

Contributors: Morrison, Peter R., and Meyer, Marion P.

References: [1] Benedict, F. G. 1932. Yale J. Biol. Med. 4:385. [2] Benedict, F. G. 1934. Ergeb. Physiol. Exptl. Pharmacol. 36:300. [3] Brody, S., J. E. Comfort, and J. S. Matthews. 1928. Missouri Univ. Agr. Expt. Sta. Bull. 115. [4] Carman, G. G., and H. H. Mitchell. 1926. Am. J. Physiol. 76:380. [5] Cowgill, G. R., and D. L. Drabkin. 1927. Ibid. 81:36. [6] Custor, J. 1873. Arch. Anat. Physiol., Physiol. Abt., p. 478. [7] Diack, S. L. 1930. J. Nutr. 3:289. [8] Dreyer, G., and W. Ray. 1912. Phil. Trans. Roy. Soc. London, B, 202:191. [9] Giaja, J.



*continued*

## 36. BODY SURFACE AREA: MAMMALS

### Part II. CONSTANTS FOR USE IN SURFACE AREA FORMULA: MAMMALS

1925. Ann. Physiol. Physicochim. Biol. 1:597. [10] Giaja, J., and B. Males. 1928. Ibid. 4:884. [11] Gley, E., and A. O. De Almeida. 1924. Compt. Rend. Soc. Biol. 90:467. [12] Hecker, C. 1894. Z. Veterinaerk. 6:97. [13] Hill, A. V., and A. M. Hill. 1913. J. Physiol. (London) 46:81. [14] Kettner, H. 1909. Arch. Anat. Physiol., Physiol. Abt., p. 447. [15] Lee, M. O., and E. Clark. 1929. Am. J. Physiol. 89:24. [16] Lines, E. W., and A. W. Pierce. 1931. Australia Council Sci. Ind. Res. Bull. 55:21. [17] Mardones, G. 1931. Compt. Rend. Soc. Biol. 108:118. [18] Mitchell, H. H. 1928. Illinois Agr. Expt. Sta. Ann. Rept. 317:155. [19] Parry, D. A. 1949. Quart. J. Microscop. Sci. 90:13. [20] Pearson, O. P. 1947. Ecology 28:127. [21] Pfandler, M. 1916. Z. Kinderheilk. 14:69. [22] Ritzman, E. G., and N. F. Colovos. 1930. New Hampshire Univ. Agr. Expt. Sta. Circ. 32. [23] Rubner M. 1883. Z. Biol. 19:553. [24] Rubner, M. 1902. Die Gesetze des Energieverbrauches bei der Ernährung. F. Deuticke, Leipzig. [25] Seuffert, R. W., R. Giese, and R. Meyer. 1926. Beitr. Physiol. 3:203. [26] Seuffert, R. W., and F. Hertel. 1925. Z. Biol. 82:7. [27] Thomas, D. 1911. Arch. Anat. Physiol., Physiol. Abt., p. 9. [28] Trowbridge, P., C. Moulton, and L. Haigh. 1915. Missouri Univ. Agr. Expt. Sta. Res. Bull. 18.

### 33. THE BRAIN: MAN



*continued*



## 37. BRAIN: MAN

*Contributors:* (a) von Bonin, Gerhardt, (b) Bartelmez, George W.

*References:* [1] Bailey, P., and G. von Bonin. 1951. The isocortex of man. Univ. Illinois Press, Urbana.

[2] Ranson, S. W., and S. L. Clark. 1959. The anatomy of the nervous system. Ed. 10. W. B. Saunders, Philadelphia.

### Part I. REGIONS AND FUNCTIONS

Region	General Functions	Sub-regions	Specific Functions	
(A)	(B)	(C)	(D)	
P R O S E P H A L O N	Telencephalon			
	1 Cerebral cortex	Highest level of integration; symbolism, memory, forecasting	Prefrontal, frontal, parietal, occipital, temporal	Association: autonomic, general motor, sensory, visual, auditory.
	2 Rhinencephalon	Olfaction, "visceral brain," emotion	Olfactory bulb, cortex; amygdala; limbic cortex of hippocampus; fornix; mammillary body	Association: autonomic-visceral integration; olfaction.
3	Corpus striatum	Smoothing of motor behavior; inhibition of posture, movement patterns, extrapyramidal relay	Caudate nucleus, putamen, globus pallidus	Motor relay (globus pallidus) back to cortex into thalamus.
Diencephalon				
4	Epithalamus	"Drive"	Habenular nucleus	
5	Thalamus	Sensory relay to cortex; thalamocortical circuits	Anterior, midline, medial, lateral, posterior, pulvinar, ventral	Cortical relay nuclei: anterior, emotion (?); medial and ventral, sensory; pulvinar, gnostic and practic. Unspecific intralaminar nuclei. Basal ganglia relay ventrolateral.
6	Metathalamus		Medial, lateral geniculate	Medial: acoustic to supratemporal plane. Lateral: visual to striate area.
7	Subthalamus	Motor, extrapyramidal	Subthalamic nucleus	Integrative facilitation and inhibition.
8	Hypothalamus	Principal forebrain center for integration of visceral functions involving autonomic nervous system <sup>1</sup>	Anterior, middle, lateral, posterior	Generally anterior part, trophotropic (parasympathetic); posterior part, ergotropic (orthosympathetic).
9	MESENCEPHALON	Postural reflexes; nuclei for cranial nerves	Superior colliculus, inferior colliculus, substantia nigra, red nucleus, tegmentum, reticular formation (part of), basis pedunculi, nucleus nerves III, IV, V (part of)	Relay for visual reflexes (protective), auditory reflexes, extrapyramidal junction of striatal and cortical influences; contributes to righting reflexes, tracts; facilitatory and inhibitory influences on motor performance.
Metencephalon				
10	Cerebellum	Maintenance of posture; equilibrium; coordination, smoothing of complex movements	Corpus cerebelli, anterior and posterior lobe; flocculonodular lobe	Paleocerebellum: equilibration, maintenance of posture. Neocerebellum: postural reflexes, stabilizing, smoothing more complex movements initiated in cortex, facilitation of posture change.
11	Pons		Pontine nucleus, reticular formation, cerebellar peduncles, tracts, nucleus nerve V (part of)	Relay between cerebro-cerebellar, motor inhibitory areas.
Myelencephalon				
12	Medulla oblongata	Reflex center for cardiac vasomotor, vomiting, deglutition, respiratory, gustatory, facial reflexes	Nucleus nerves V (part of), IX, XI, XII; inferior olivary; tracts; reticular nuclei of medullary tegmentum	Posture.

<sup>1/</sup> Energy and water exchange, sexual function, sleep, vasomotor.

*continued*

# Contrails

## 37. BRAIN: MAN

### Part I. REGIONS AND FUNCTIONS










*Contributors:* (a) Stevenson, James A. F., (b) von Bonin, Gerhardt

*References:* [1] Barr, M. L. Unpublished. Univ. Western Ontario Dept. of Anatomy, London, Canada, 1953.

[2] Fulton, J. F. 1949. Physiology of the nervous system. Ed. 3. Oxford Univ. Press, New York. [3] Ranson, S. W., and S. L. Clark. 1959. The anatomy of the nervous system. Ed. 10. W. B. Saunders, Philadelphia.

### Part II. CORTICAL CEREBRAL REGIONS AND FUNCTIONS




**Area** (column C): numbers indicate areas of Brodmann.

Gross Region	Location	Area	Principal Connected Pathways and Areas	Function
(A)	(B)	(C)	(D)	(E)
1 Occipital (striate cortex)		17	Optic radiation; lateral geniculate body of the thalamus; optic tract; chiasm; optic nerve; retina	Vision
2 Occipital (parastriate) and parietal (preoccipital)		18,19	Superior colliculus of brain stem; areas 37, 8, 23, 21 of Brodmann; opposite hemisphere (areas 18, 19)	Visual elaboration
3 Temporal (Heschl's gyrus)		41	Auditory radiation; medial geniculate body; inferior colliculus; nucleus of lateral lemniscus; superior olive; dorsal and ventral cochlear nucleus in brain stem; cochlear nerve; spiral ganglion; hair cells in organ of Corti	Hearing (auditosensory)
4 Temporal		42,22		Hearing (audiotpsychic)
5 Temporal (piriform area)		28,34	Lateral root; olfactory tract; olfactory bulb; fila olfactoria	Olfaction
6 Parietal (post central convolution)		3,1,2	Connected through posteroventral nuclei with mesial and trigeminal fillets and spinothalamic tract; dorsal root; sensory root ganglion; peripheral sensory nerves	Somatic sensation
7 Parietal (superior lobule)		5,7 <sub>a</sub>	Dorsolateral thalamus; association fibers connecting areas 1, 44, 19, 8, 46 of Brodmann	Sensory elaboration (motor skills)
8 Frontal (pons triangularis and opercularis); dominant hemisphere; Broca's area		44,45	Receives afferents from area 3 of Brodmann; sends impulses to areas 4 and 5	Speech (motor)
9 Parietal (lower lobule); dominant hemisphere; Wernicke's area		40,39	Association fibers connecting areas 1, 42, 19 of Brodmann	Speech (sensory) (Gnosia)

*continued*

## 37. BRAIN: MAN

### Part II. CORTICAL CEREBRAL REGIONS AND FUNCTIONS

Gross Region	Location	Area	Principal Connected Pathways and Areas	Function
(A)	(B)	(C)	(D)	(E)
10 Temporal		21,22	Frontal and parietal association areas (areas 11, 12, 13, 39, 40 of Brodmann)	Memory
11 Frontal		9,10,11, 12,13, 14,24	Orbitofrontal cortex (areas 9,10,11,12 of Brodmann) and posterior and medial orbital gyri (areas 13 and 14 of Walker) are connected to dorsomedial nuclei of thalamus which has hypothalamic connections; the anterior cingulate gyrus (area 24) is projection area of anterior nuclei of thalamus	Projection of conscious thought
12 Frontal (precentral area, motor and premotor)		4,6	Internal capsule, pyramidal decussation, corticospinal tracts, anterior horn cells, motor roots; connected by way of lateroventral nucleus and the superior cerebellar peduncle with spino-cerebellar afferents	Motion (motor activity)

*Contributor:* Raaf, John

*References:* [1] Fulton, J. F. 1949. Physiology of the nervous system. Ed. 3. Oxford Univ. Press, New York. p. 370. [2] Kluver, H., and P. Bucy. 1939. Arch. Neurol. Psychiat. 42:979. [3] Penfield, W., and T. Rasmussen. 1950. The cerebral cortex of man. Macmillan, New York. [4] Ranson, S. W., and S. L. Clark. 1959. The anatomy of the nervous system. Ed. 10. W. B. Saunders, Philadelphia. [5] Spiegel, E. A. 1934. Arch. Neurol. Psychiat. 31:469. [6] von Bonin, G. 1950. Essay on the cerebral cortex. C. C. Thomas, Springfield, Ill.

### Part III. NUCLEI OF METATHALAMUS AND DORSAL THALAMUS

**Function** (column B): A = associational; D = diffuse associational; I = internal relay; E = external relay; U = unclassified or doubtful.

Nucleus	Function	Afferents	Efferents	Reference
(A)	(B)	(C)	(D)	(E)
1 Medialis dorsalis	A	Principal: some from medioventral region (hypothalamus)	Frontal granular cortex.	2,4,16,19
2 Lateralis posterior	A	Principal: parietal cortex. Diffuse: sensorimotor, limbic, auditory, visual, frontal cortexes. From nuclei 5-9 in this table.	Parietal field.	2,4,10,13, 16,18, 19
3 Lateralis anterior	A	Principal: parietal cortex. Diffuse: sensorimotor, auditory, visual cortexes. From nuclei 5-9 in this table.		
4 Pulvinar	A	Principal: parietal association. Diffuse: frontal, sensorimotor, limbic, auditory, visual cortexes. Subcortical: ventral posterior nucleus of thalamus. From nuclei 5-9 in this table.	Posterior parietal and temporal cortex, superior parietal lobule, posterior Sylvian region, supramarginal gyrus, temporo-occipital region, cortical area 18, superior colliculus and pretectum, posterior parietal, temporal and occipital (parasensory) fields.	2,4,10,11, 13,16, 18-20
5 Centrum medianum	D	Subcortical: reticular formation; other intralaminar <sup>1</sup> thalamic nuclei	Subcortical: caudate nucleus, corpus striatum.	2,4,7,8, 12,16-18

<sup>1</sup>/ Denotes nuclei not only adjacent to internal medullary lamina of thalamus but also others (such as midline nuclei, reticularis anterior and ventralis anterior) which, when stimulated, can evoke cortical and intrathalamic recruiting responses.

*continued*

## 37. BRAIN: MAN

### Part III. NUCLEI OF METATHALAMUS AND DORSAL THALAMUS

Nucleus	Function	Afferents	Efferents	Reference
(A)	(B)	(C)	(D)	(E)
6 Centralis medialis	D	Reticular formation; intralaminar thalamic nuclei	Cortical associational areas: frontal, cingulate, orbital caudate nucleus.	2,4,7,8,16-18
7 Centralis lateralis				
8 Ventralis anterior				
9 Reticularis anterior	D	Reticular formation; other intralaminar thalamic nuclei		
10 Anterodorsalis	I	Corresponding opposite nucleus	Retrosplenial region	2,4,16,19,20
11 Anteromedialis	I	Mammillothalamic tract; limbic cortex. From nuclei 5-9 in this table.	Anterior gyrus cinguli	2,4,13,16,18,20
12 Anteroventralis	I	Subcortical: from nuclei 5-9 in this table; mammillothalamic tract. Cortical: gyrus cinguli.	Posterior gyrus cinguli	2,4,13,16,18,19
13 Ventral anterior	I	Principal: globus pallidus. Diffuse: orbital, parietal, frontal, sensorimotor, limbic cortices. From nuclei 5-9 in this table.	Globus pallidus; prefrontal cortex	2,4,5,9,13,16,18
14 Ventralis lateralis	I	Principal: superior cerebellar peduncle	Precentral motor cortex, areas 4 and 6. Brachium conjunctivum.	2,4,9,16,19,20
15 Ventral medial	I		Globus pallidus; lateral frontal cortex	2,4,9,16
16 Medial geniculate body	E	Inferior colliculus and parabrachial body; auditory cortex	Auditory cortex; parvocellular and magnocellular to temporal cortex in lower wall of Sylvian fissure	2,4,15,16
17 Lateral geniculate body	E	Ganglion cell layer of retina		2,4,16
18 Ventral postero-lateral	E	Medial lemniscus; spinothalamic tract; sensorimotor cortex	Sensorimotor cortex	2-4,6,13,14,16
19 Ventral postero-medial	E	Trigeminal lemniscus; trigeminal thalamic tract	Sensorimotor cortex	2,4,16
20 Reticularis lateralis (posterior)	U	Unclassified or doubtful	Cortex	1,2,4,16,20
21 Midline <sup>a</sup>	D or U	Spinothalamic tract; medial lemniscus	Hypothalamus; basal ganglia; lateral thalamus nuclei	2,4,16-18,20

/z/ Midline nuclei: rhomboideus, reuniens, paracentralis, parafascicularis, paraventricularis (anterior and posterior), parataenialis.

*Contributors:* (a) Niemer, William T., (b) von Bonin, Gerhardt

*References:* [1] Chow, K. L. 1952. *J. Comp. Neurol.* 97:37. [2] Crosby, E. C., T. Humphrey, and E. W. Lauer. 1962. *Correlative anatomy of the nervous system.* Macmillan, New York. [3] Dusser de Barenne, J. G., and W. S. McCulloch. 1938. *J. Neurophysiol.* 1:176. [4] Fields, J., ed. 1959-61. *Handbook of physiology.* American Physiological Society, Washington, D. C. sect. 1, v. 1-3. [5] Freeman, W., and J. W. Watts. 1947. *J. Comp. Neurol.* 86:65. [6] Getz, B. 1952. *Acta Anat.* 16:271. [7] Jasper, H. H., et al., ed. 1958. *Reticular formation of the brain.* Little and Brown, Boston. [8] Johnson, F. H. 1953. *Anat. Record* 115:327. [9] Krieg, W. J. S. 1953. *Functional neuroanatomy.* Ed. 2. Blakiston, New York. [10] Le Gros Clark, W. E. 1932. *Brain* 55:406. [11] Le Gros Clark, W. E., and D. W. C. Northfield. 1937. *Ibid.* 60:126. [12] McLardy, T. 1948. *Ibid.* 71:290. [13] Niemer, W. T., and J. Jimenez-Castellanos. 1950. *J. Comp. Neurol.* 93:101. [14] Rasmussen, A. T. 1948. *J. Comp. Neurol.* 88:411. [15] Rundles, R. W., and J. W. Papez. 1938. *Ibid.* 68:267. [16] Sheer, D. E., ed. 1961. *Electrical stimulation of the brain.* Univ. Texas Press, Austin. [17] Starzl, T. E., and H. W. Magoun. 1951. *J. Neurophysiol.* 14:133. [18] Starzl, T. E., and D. G. Whitlock. 1952. *Ibid.* 15:449. [19] von Bonin, G. Unpublished. Univ. Illinois College Medicine, Chicago, 1956. [20] Walker, A. E. 1938. *The primate thalamus.* Univ. Chicago Press, Chicago.

*continued*



# Contrails

## 37. BRAIN: MAN

### Part IV. TRACTS

*Abbreviations:* C = cervical; S = sacral; T = thoracic.

Tract	Origin	Termination	Pathway	Function
(A)	(B)	(C)	(D)	(E)
1 Allen's fasciculus <sup>1</sup>	Solitary nucleus	Ventral column: C 3-T 6	Lateral funiculus along dorso-lateral edge of ventral column	Associated with respiratory control
2 Arcuate fasciculus	Cerebral cortex of basal frontal lobes	Cortex of temporal, lower parietal, and occipital regions	Through base of angular gyrus into interior and middle frontal gyri; in parietal and frontal opercula over upper border of insula	Association bundle
3 Central acoustic	Contralateral and homolateral cochlear nucleus and olivary complex; nuclei of central acoustic tract	Medial geniculate body, inferior colliculus	With lateral lemniscus to ventrolateral surface of inferior colliculus; runs in brachium of inferior colliculus	Auditory
4 Corticospinal, lateral	Cerebral cortex	Contralateral anterior horn of spinal cord	Internal capsule; cerebral peduncle; decussation of pyramids, lateral corticospinal tract	Motor
5 Corticospinal, ventral	Cerebral cortex	Ipsilateral anterior horn of cord	Internal capsule; peduncle, ventral corticospinal tract about anterior median fissure of cord	Motor
6 Cuneate fasciculus	Dorsal root ganglia of C 1-T 6	Cuneate nucleus	Lateral portion of dorsal funiculus above T 6	Proprioception; discriminative touch (upper extremities)
7 Dorsolateral fasciculus (Lissauer)	Dorsal root fibers, fibers from substantia gelatinosa and dorsal horn of gray matter	Substantia gelatinosa (Rolando), nucleus proprius within 2-4 segments	Dorsolateral to substantia gelatinosa of dorsal horn of gray matter	Pain, temperature, and some tactile
8 Fastigiobulbar fasciculus	Fastigial nuclei	Reticular formation of medulla	Mingled with uncinata fasciculus of Russel in juxtarestiform body	Cerebellar inhibitory path from vermis bulbar reticular substance
9 Frontal fasciculus, superior (Burdach)	Basal frontal regions of cortex	Temporal, lower parietal, and occipital cortex	Dorsal to insula	Association fibers
10 Frontopontine fasciculus	Posterior part of superior and middle frontal gyri bordering on precentral gyrus	Pontine nuclei	Extreme medial and lateral portions of cerebral peduncle	Cortical relay fibers to middle lobe of cerebellum
11 Fasciculus gracilis (Goll)	Dorsal root ganglia of T 6-S 5	Nucleus gracilis	Medial portion of dorsal funiculus	Proprioception; discriminative touch (lower extremities)
12 Habenulopeduncular fasciculus <sup>2</sup>	Habenular nuclei	Interpeduncular nucleus	Runs ventrolaterally; arches beneath centrum medianum through medial border of red nucleus and along midventral line	
13 Habenulotegmental	Habenular nuclei	Dorsal tegmental nucleus		
14 Hypothalamic, descending <sup>3</sup>	Dorsomedial nuclei, posterior and lateral hypothalamic areas, perifornical areas	Autonomic cells in medulla oblongata and intermediolateral cell column of cord	Descends between mammillary body and red nucleus to ipsilateral reticular formation; dorsal to substantia nigra, then to region of vestibular fiber complex	
15 Interfascicular fasciculus (Schultze)	Descending fibers of posterior funiculus	Nucleus proprius of gray matter	Between, and mingled with, fasciculus gracilis and cuneatus	Reflex collaterals
16 Interstitiospinal (Cajal)	Mesencephalon, region of posterior commissure	Intermediolateral cell column of cord (C 8-T 1)	Descends along ventromedian sulcus of cervical cord	
17 Laterocerebellar	Lateral reticular nucleus of medulla	Cerebellum	Runs with external arcuate fibers	

/1/ Solitariospinal. /2/ Retroflex bundle of Meynert. /3/ Not yet demonstrated anatomically.

*continued*

# Contrails

## 37. BRAIN: MAN

### Part IV. TRACTS

	Tract	Origin	Termination	Pathway	Function
	(A)	(B)	(C)	(D)	(E)
18	Longitudinal fasciculus, dorsal (Schlitz)	Hypothalamus and dorsal tegmental nucleus	Somato-motor and autonomic motor nuclei of brain stem	Continue into ventral fasciculus proprius of cord	
19	Longitudinal fasciculus, inferior	Occipital cortex	Temporal cortex		Association fibers
20	Longitudinal fasciculus, medial	Mesencephalon, at level of posterior commissural nuclei	Through medulla, continued in cord as sulcomarginal fasciculus of Marie	Bilateral, about midventral line between central gray matter and tectospinal fasciculus (in medulla)	Contains vestibular reflex fibers, fibers from reticular nuclei, abducens nucleus
21	Mammillary fasciculus	Medial mammillary nuclei	Anterior thalamic nuclei and tegmentum	Runs dorsally from mammillary bodies, then bifurcates; one limb continues dorsally, the other caudally	
22	Mammillotegmental fasciculus	Medial and lateral mammillary nuclei	Tegmentum	Initially a component of the principal mammillary fasciculus which it leaves to run caudally	
23	Mammillothalamic fasciculus (Vicq d'Azyr)	Medial and lateral mammillary nuclei	Anterior nuclear mass of thalamus	Dorsorostrally through medial thalamic wall	
24	Olfactory	Olfactory bulb	In three striae: medial, intermediate, and lateral olfactory	Lies between gyrus rectus and medial orbital gyrus, covering olfactory sulcus	Olfaction
25	Olivocochlear bundle	Nuclei of olivary complex	Organ of Corti	Via auditory nerve	Inhibition of neural response from organ of Corti
26	Optic	Ganglion cells of retina	Lateral geniculate and/or superior colliculus	Tract encircles thalamus ventrally	Vision
27	Perpendicular fasciculus	Inferior parietal lobule	Fusiform gyrus	Obliquely dorsomedial and ventrolateral	Association
28	Probst's	Sensory cells associated with nucleus of cranial nerve V in area of cerebral aqueduct or lateral reticular nucleus of mesencephalon	Intercalated nucleus and dorsal vagal nucleus	Ventrolateral to solitary fasciculus, dorsomedial to nucleus of spinal tract of cranial nerve V	Appears to link various parts of trigeminal system; relating masticatory movements with salivation
29	Reticulospinal fasciculus, lateral, direct	Reticular substance	Gray matter of cord	Lies in region of overlap of lateral cortico- and rubro-spinal tracts	Extrapyramidal aspects of motor function
30	Reticulospinal fasciculus, ventral, crossed	Reticular substance	Gray matter of cord	Ventrolateral to ventral cortico-spinal fibers	Extrapyramidal aspects of motor function
31	Rubrospinal	Red nucleus	Gray matter of cord	Ventromedial to, and overlapping, lateral corticospinal tract	Extrapyramidal function
32	Septomarginal fasciculus	Collaterals of dorsal funicular fibers (fasciculus cuneatus)	Gray matter of cord	Along posterior median sulcus in middle of posterior funiculus	Proprioceptive reflex connections
33	Solitary fasciculus	Fibers from cranial nerves VII, IX, and X	Solitary nucleus	Dorsomedial to spinal root of cranial nerve V; extends from level of medullary stria to caudal end of medulla	Visceral afferent; oral and concerned with taste
34	Spinal trigeminal	Cells of trigeminal nerve	Nucleus of spinal trigeminal tract	Ventromedial to restiform body; in position of Lissauer's zone	Pain, temperature, and some tactile from face

*continued*

# Contrails

## 37. BRAIN: MAN

### Part IV. TRACTS

Tract	Origin	Termination	Pathway	Function
(A)	(B)	(C)	(D)	(E)
35 Spinocerebellar fasciculus, ventral	Border cells about medial border of central lateral ventral column, cells about dorsal nucleus of Clark	Vermis of anterior lobe of cerebellum	On periphery of cord ventral to dorsal spinocerebellar, and lateral to lateral spinothalamic tract	
36 Spinocerebellar fasciculus, dorsal (Flechsig)	Dorsal horn and dorsal nucleus of Clark	Cortex of anterior cerebellar lobe, uvula and pyramis of vermis	On periphery of lateral funiculus of cord	
37 Spinoolivary (Helweg)	Dorsal horn of spinal cord	Inferior olive	Ventral and superficial part of lateral funiculus between lateral and ventral spinothalamic tracts	
38 Spinothalamic, ventral <sup>4</sup>	Dorsal horn of gray matter (substantia gelatinosa)	Posterolateral ventral nucleus of thalamus	Lateral funiculus just medial to ventral spinocerebellar tract	Pain and temperature from extremities and trunk
39 Spinothalamic, ventral	Proper nucleus of contralateral dorsal horn	Posterolateral ventral nucleus of thalamus	Lateral portion of ventral funiculus	Light touch
40 Subcallosal fasciculus	Frontal cortex	Striate body	Dorsal to caudate nucleus below radiation of corpus callosum	
41 Tectospinal (Lowenthal)	Superior colliculus	Spinal gray matter	Ventromedial portion of ventral funiculus	Head and shoulder reflex movements to light and sound
42 Tegmental fasciculus, central	Variable composition		In reticular formation of medulla lateral to crossed vestibulospinal tract	
43 Tegmental fasciculus, central (part of)	Central gray matter of cerebral aqueduct	About sac of inferior olive	Flattened bundle running through tegmentum, oblique to horizontal plane, ventrolateral to medial longitudinal fasciculus	
44 Thalamic fasciculus	Fibers of brachium conjunctivum going to thalamus through tegmental field of Forel; fibers from globus pallidus	Thalamus; nucleus (ventralic anterior and ventralic lateralis)	Ventrally through Forel's field H <sub>1</sub> , between mammillothalamic tract and lenticular fasciculus	
45 Vestibulofastigial	Vestibular nuclei and spinal nucleus of cranial nerve V	Fastigial nucleus and vermian cortex	Between restiform body and periventricular gray matter in juxtarestiform body	
46 Vestibulofloculonodular	Cells of origin in vestibular ganglion (Scarpa)	Cortex of flocculus, nodulus of cerebellum		Vestibular
47 Vestibuloglobose	Vestibular nuclei and spinal nucleus of cranial nerve V	Globose nucleus and vermian cortex	Between restiform body and periventricular gray matter in juxtarestiform body	
48 Vestibulospinal, crossed, ventral	Medial, inferior, and spinal vestibular nuclei	Gray matter of cord	Medial portion of ventral funiculus in sulcomarginal fasciculus of Marie	Vestibular reflex
49 Vestibulospinal, direct, lateral	Lateral vestibular nucleus	Gray matter of cord	Ventrolateral part of ventral funiculus and ventromedial portion of lateral funiculus	Vestibular reflex

<sup>4</sup>/ Lemniscæ.

*Contributors:* (a) Sutin, Jerome, (b) Campbell, Berry

*References:* [1] Crosby, E. C., T. Humphrey, and E. W. Lauer. 1962. Correlative anatomy of the nervous system. Macmillan, New York. [2] Mettler, F. A. 1942. Neuroanatomy. C. V. Mosby, St. Louis. [3] Riley, H. A. 1960. An atlas of the basal ganglia, brain stem, and spinal cord. Rev. ed. S. Hafner, New York.




## 38. AUTONOMIC NERVOUS SYSTEM: MAN

Both sympathetic and parasympathetic pathways contain afferent and efferent fibers, and the pathways are associated generally with the same viscera or vessels. Afferent sympathetic nerve cells are located in dorsal spinal nerve root ganglia. Their peripheral processes run in visceral and vascular nerves; their central processes enter the cord through dorsal nerve roots. Segmental numbers of nerve roots transmitting afferents from any viscus correspond usually to segments containing preganglionic neurones for that structure. Afferent parasympathetic nerve cells lie in root ganglia of cranial nerves V, VII, IX, and X, which are homologues of dorsal spinal root ganglia. Their peripheral processes run in the nerves indicated and carry impulses from glands and vessels; their central processes enter the brain stem. Afferent parasympathetic fibers from pelvic structures (cervix uteri, base of bladder, prostate, and rectum) are carried in the pelvic splanchnic nerves to the cord. Autonomic afferents participate in segmental reflex arcs. Relays also ascend in tracts in the anterolateral and posterior white columns of the cord to the brain stem and hypothalamus, where they may end or are further relayed to cortical levels, e.g., in the frontal lobes of the brain.

*Contributor:* Mitchell, G. A. G.

*References:* [1] Mitchell, G. A. G. 1953. Anatomy of the nervous system. E. and S. Livingstone, Edinburgh. [2] Mitchell, G. A. G. 1956. Cardiovascular innervation. E. and S. Livingstone, Edinburgh. [3] White, J. C., and W. H. Sweet. 1955. Pain: its mechanisms and neurosurgical control. C. C. Thomas, Springfield, Ill.



*Contributors:* Magoun, Horace, W., and Bridgman, Charles F.

*continued*



## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part I. SYMPATHETIC CONNECTIONS

**Cell Body** (column C): boldface type indicates cell bodies agreed upon by most investigators. **Pathway** (column D): WRC = white ramus communicans; ST = sympathetic trunk.

	Organ	Effector	Preganglionic Neurones		Postganglionic Neurones		Actions	Reference
			Cell Body	Pathway	Cell Body	Distribution		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Eye	Dilator pupillae	<b>T 1, 2, 3<sup>1</sup></b>	WRC, ST	Superior cervical ganglion	Internal carotid plexus → short and long ciliary nerves	Dilation	12,13
2			C 8, <b>T 1, 2</b>	WRC, ST	Superior cervical ganglion	Internal carotid plexus → short and long ciliary nerves	Dilation	11
3		Muscle orbitalis	<b>T 1, 2, 3</b>	WRC, ST	Superior cervical ganglion	Internal carotid plexus → ophthalmic nerve	?	12
4	Lacrimal gland	Blood vessels	<b>T 1, 2, 3</b>	WRC, ST	Superior cervical ganglion	Internal carotid plexus and branches	Vasoconstriction	8,9
5	Heart	Mostly ventricular muscle, some atrial	<b>T 1-5</b>	WRC, ST	Superior, middle, and inferior cervical ganglia; stellate and upper 4 or 5 thoracic ganglia	Superior, middle, and inferior cervical and thoracic cardiac nerves → cardiac plexus	Augmentation, acceleration	7,15
6	Coronary arteries	Smooth muscle	<b>T 1-4 or 5</b>	WRC, ST	Superior, middle, and inferior cervical ganglia; stellate and upper 3 or 4 thoracic ganglia	Superior, middle, and inferior cervical and thoracic cardiac nerves → cardiac plexus → coronary plexuses	Vasodilation	5,14
7	Blood vessels	Meningeal arteries	<b>T 1, 2</b>	WRC, ST	Superior cervical and internal carotid ganglia	External carotid and vertebral plexuses → meningeal arteries	Vasoconstriction	16
8		Cerebral arteries	<b>T 1, 2</b>	WRC, ST	Superior cervical ganglion	Internal carotid plexus and branches	Vasoconstriction	
9		Vertebral system of brain	<b>T 1, 2</b>	WRC, ST	Stellate ganglion	Vertebral plexus	Vasoconstriction	
10	Blood vessels, sweat glands, and piloerection muscles	Head, neck	<b>T 1, 2, 3<sup>1</sup></b>	WRC, ST	Superior cervical ganglion, internal carotid ganglia	Gray rami → cervical plexus, and certain cranial nerves and perivascular plexuses	Vasoconstriction	5,14
11		Upper limb	<b>T 1, 2-9, 10</b>	WRC, ST	Superior, middle, and inferior cervical ganglia; stellate ganglion	Gray rami → brachial plexus and subclavian artery and branches	Sweating, piloerection, vasoconstriction	13
12		Thoracic and upper abdominal wall	<b>T 2-10</b>	WRC, ST	Middle and inferior cervical ganglia; stellate and upper thoracic ganglia	Gray rami → intercostal nerves	Sweating, piloerection, vasoconstriction	3
13	Lower limb and trunk	<b>T 6-L 2</b>	WRC, ST	Lumbar 1-4; sacral 1-3	Gray rami → lumbar and sacral nerves	Sweating, piloerection, vasoconstriction	3	
14	Suprarenal medulla	Cells of medulla	<b>T 5-9, 10-L 1, 2</b>	WRC, ST (splanchnic)	Cells of suprarenal medulla	No postganglionic pathway	Secretion	5,14
15	Lung	Trachea and bronchi	<b>T 2-4</b>	WRC, ST	Inferior cervical, stellate, and upper 4 thoracic ganglia	Tracheal and bronchial nerves	Tracheal and bronchial dilation	2,11
16		Blood vessels	<b>T 2-4</b>	WRC, ST	Inferior cervical, stellate, and upper 4 thoracic ganglia	Pulmonary nerves and plexuses	Vasoconstriction	11

<sup>1</sup>/ Occasionally C 8 and T 4.

*continued*

## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part I. SYMPATHETIC CONNECTIONS

	Organ	Effector	Preganglionic Neurons		Postganglionic Neurons		Actions	Reference
			Cell Body	Pathway	Cell Body	Distribution		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
17	Submaxillary and sublingual glands	Gland and vessels	T 1, 2, 3	WRC, ST	Superior cervical ganglion	External carotid plexus and branches	Vasoconstriction, weak secretion	1,8
18	Parotid gland	Gland and vessels	T 1, 2, 3	WRC, ST	Superior cervical ganglion	External carotid plexus and branches	Vasoconstriction, weak secretion ?	1,8
19	Lower esophagus	Smooth muscle	T 1-3, 4-6	WRC, ST (greater splanchnic)	Stellate, upper thoracic, and celiac ganglia	Esophageal rami and plexus, left gastric and phrenic nerves	Inhibits peristalsis	6
20	Cardiac sphincter	Smooth muscle	T 1-3, 4-6	WRC, ST (greater splanchnic)	Stellate, upper thoracic, and celiac ganglia	Esophageal rami and plexus, left gastric and phrenic nerves	Contraction	6
21	Stomach	Smooth muscle and gland	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac ganglion	Accompanies gastric and gastroepiploic arteries	Inhibits peristalsis	5,14
22		Pyloric sphincter	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac ganglion	Accompanies right gastric artery	Contraction	
23		Blood vessels	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac ganglion	Accompanies gastric and gastroepiploic arteries	Vasoconstriction	
24	Pancreas	Gland	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac ganglion	Accompanies pancreatic arteries	Secretion ?	4
25		Blood vessels	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac ganglion	Accompanies pancreatic arteries	Vasoconstriction	
26	Liver	Blood vessels	T 5, 6, 7-10	WRC, ST (splanchnic)	Celiac ganglion	Periarterial plexus of hypogastric artery	Vasoconstriction	5,14
27	Gallbladder	Smooth muscle	T 5, 6, 7-10	WRC, ST (splanchnic)	Celiac ganglion	Periarterial plexus of hepatic and cystic arteries	Relaxation	5,14
28		Sphincter of common bile duct	T 5, 6, 7-10	WRC, ST (thoracic splanchnic)	Celiac ganglion	Periarterial plexuses	Contraction	
29	Small intestine and proximal colon	Smooth muscle and glands	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac and superior mesenteric ganglia	Celiac and superior mesenteric rami	Secretion (?), inhibition	5,14
30	Ileocecal sphincter	Blood vessels and sphincter muscle	T 5, 6-10, 11	WRC, ST (thoracic splanchnic)	Celiac and superior mesenteric ganglia	Celiac and superior mesenteric rami	Vasoconstriction, contraction	5,14
31	Distal colon and rectum	Smooth muscle	T 12, L 1, 2, 3	WRC, ST (lumbar splanchnic)	Inferior mesenteric ganglion	Plexus of inferior mesenteric artery	Contraction, inhibition	5,14
32		Blood vessels	T 12, L 1, 2, 3	WRC, ST (lumbar splanchnic)	Inferior mesenteric ganglion	Plexus of inferior mesenteric artery	Vasoconstriction	
33	Internal sphincter ani	Blood vessels and sphincter muscle	L 1, 2, 3	WRC, ST (lumbar splanchnic)	Inferior mesenteric ganglion	Superior hypogastric plexus	Vasoconstriction, contraction	5,14
34	Kidney	Blood vessels and smooth muscle	T 5 - L 2	WRC, ST (thoracic splanchnic)	Aorticorenal or renal ganglion	Renal plexus	Vasomotor changes	5,14
35	Ureter	Blood vessels and smooth muscle	T 5 - L 2	WRC, ST (thoracic and lumbar splanchnic)	Aorticorenal or renal and inferior hypogastric ganglia	Renal plexus and hypogastric nerves	Rhythmic contraction	5,14
36	Bladder	Detrusor vesicae	L 1-3	WRC, ST (hypogastric nerves, inferior hypogastric plexus)	Vesical plexus, intramural ganglia	Intramural plexus	Relaxation	9

*continued*

### 38. AUTONOMIC NERVOUS SYSTEM: MAN

#### Part I. SYMPATHETIC CONNECTIONS

Organ	Effector	Preganglionic Neurones		Postganglionic Neurones		Actions	Reference
		Cell Body	Pathway	Cell Body	Distribution		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
37 Bladder	Sphincter vesicae	L 1-3	WRC, ST (hypogastric nerves, inferior hypogastric plexus)	Vesical plexus	Intramural plexus	Constriction for ejaculation	10
38 Urethra	Compressor urethrae	L 1-3	WRC, ST (hypogastric nerves, inferior hypogastric plexus)	Vesical plexus, intramural ganglia	Prostatic plexus	Contraction	5,14
39 Prostate gland	Smooth muscle	T 12 - L 2	WRC, ST (hypogastric nerves, inferior hypogastric plexus)	Vesical and prostatic plexuses	Prostatic plexus	Contraction, ejaculation	5,14
40 Seminal vesicle and vas deferens	Smooth muscle	T 12 - L 2	WRC, ST (hypogastric nerves, inferior hypogastric plexus)	Vesical plexus	Plexuses of seminal vesicles and vasa deferentia	Contraction, ejaculation	5,14
41 Testis	Blood vessels	T 6-12	WRC, ST	Probably lower thoracic ganglia of sympathetic trunk	Thoracic splanchnic nerves → aortic and renal plexuses → spermatic plexus	Vasoconstriction	5,14
42 Corpora cavernosa	Blood vessels	T 12 - L 2	WRC, ST	Lumbar and sacral ganglia of sympathetic trunk	Inferior hypogastric plexus → prostatic plexus → cavernous plexus	Vasoconstriction	5,14
43 Clitoris, labia minora	Blood vessels	T 12 - L 2	WRC, ST	Lumbar and sacral ganglia of sympathetic trunk	Inferior hypogastric plexus → vaginal plexus → cavernous plexus	Vasoconstriction	5,14
44 Vagina	Smooth muscle	T 12 - L 2	WRC, ST (hypogastric nerves)	Inferior hypogastric and uterovaginal plexuses	Vaginal plexus	Contraction	5,14
45 Ovary and uterine tube	Vascular bed and stroma	T 6-12	WRC, ST	Probably lower thoracic ganglia of sympathetic trunk	Thoracic splanchnic nerves → aortic and renal plexuses → ovarian plexus	Vasoconstriction, contraction	5,14

*Contributors:* (a) Rogers, William M., (b) Mitchell, G. A. G.

*References:* [1] Babkin, B. P. 1950. Secretory mechanism of the digestive glands. Ed. 2. P. B. Hoeber, New York. [2] Dixon, W. E., and F. Ransom. 1912. J. Physiol. (London) 45:413. [3] Foerster, O., et al. 1936. In O. Bumke and O. Foerster, ed. Handbuch der Neurologie. J. Springer, Berlin. Bd. 5. [4] Fulton, J. F., et al., ed. 1955. Howell's Textbook of physiology. Ed. 17. W. B. Saunders, Philadelphia. p. 1013. [5] Gaskell, J. F. 1886. J. Physiol. (London) 7:1. [6] Knight, G. C. 1934. Brit. J. Surg. 22:155. [7] Kuntz, A. 1953. The autonomic nervous system. Ed. 4. Lea and Febiger, Philadelphia. [8] Langley, J. N. 1898. In E. A. Schäfer, ed. Textbook of physiology. Y. J. Pentland, Edinburgh. v. 1, p. 475. [9] Langley, J. N. 1900. Ibid. v. 2, p. 616. [10] Langworthy, O. R., L. C. Kolb, and L. G. Lewis. 1940. Physiology of micturition. Williams and Wilkins, Baltimore. [11] Müller, L. R. 1931. Lebensnerven und Lebenstrieb. J. Springer, Berlin. Aufl. 3. [12] Rasmussen, A. T. 1952. The principal nervous pathways. Ed. 4. Macmillan, New York. [13] Ray, B. S., J. C. Hinsey, and W. A. Geohegan. 1943. Ann. Surg. 118:647. [14] Rogers, W. M. Unpublished. Columbia Univ., New York, 1953. [15] Saccamanno, G. 1943. J. Compt. Neurol. 79:355. [16] White, J. C., R. H. Smithwick, and F. A. Simeone. 1952. The autonomic nervous system. Ed. 3. Macmillan, New York.

*continued*

### 38. AUTONOMIC NERVOUS SYSTEM: MAN

#### Part II. PARASYMPATHETIC CONNECTIONS

Cell Body (column C): Boldface type indicates cell bodies agreed upon by most investigators.

	Organ (A)	Effector (B)	Preganglionic Neurons		Distribution of Postganglionic Neurons (E)	Actions (F)	Reference (G)
			Cell Body (C)	Pathway (D)			
1	Eye	Sphincter pupillae	Accessory (autonomic) oculomotor nucleus	Oculomotor nerve → motor root → ciliary ganglion	Ciliary ganglion → short ciliary nerves	Constriction of pupil	3,7
2		Ciliary muscle	Accessory (autonomic) oculomotor nucleus	Oculomotor nerve → motor root → ciliary ganglion	Ciliary ganglion → short ciliary nerves	Accommodation	12
3	Lacrimal gland	Gland cells	Nucleus salivatorius superior	Nervus intermedius → greater petrosal nerve → nerve of pterygoid canal	Pterygopalatine ganglion → zygomatic nerve → lacrimal nerve	Secretion	12
4	Heart	Sinoatrial node; atrioventricular node, conduction system; cardiac muscle	Dorsal motor nucleus of vagus	Superior, inferior, and thoracic cardiac rami of vagus	Intrinsic cardiac ganglia	Inhibitory; decreases heart rate	5
5	Coronary arteries	Smooth muscle	Dorsal motor nucleus of vagus	Superior, inferior, and thoracic cardiac rami of vagus	Intrinsic cardiac ganglia, adventitia and branches of coronary artery	Constriction	13
6	Blood vessels	Smooth muscle	Demonstrated only for brain, meninges, face, most glands, and pelvic and genital organs			Vasoconstriction	12
7	Suprarenal medulla	None demonstrated					12
8	Spleen	None demonstrated					12
9	Lung and trachea	Smooth muscle of trachea, bronchial tree	Dorsal motor nucleus of vagus	Vagal branches → anterior and posterior pulmonary, and cardiac plexuses	Intramural tracheal and bronchial ganglia	Tracheal and bronchial constriction	12
10		Tracheal and bronchial glands	Dorsal motor nucleus of vagus	Vagal branches → anterior and posterior pulmonary, and cardiac plexuses	Intramural tracheal and bronchial ganglia	Secretion	
11	Submaxillary and sublingual glands	Gland cells and blood vessels	Nucleus salivatorius superior	Nervus intermedius → 7th nerve → chorda tympani → lingual nerve	Submaxillary ganglion → submaxillary and sublingual branches	Secretion, vasodilation	6
12	Parotid gland	Gland cells and blood vessels	Nucleus salivatorius inferior	9th nerve → tympanic nerve → lesser petrosal nerve	Otic ganglion → auriculotemporal nerve	Secretion, vasodilation	12
13	Lower esophagus	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → esophageal plexus	Myenteric ganglionated plexus	Increases tonus; peristalsis	12
14		Cardiac sphincter	Dorsal motor nucleus of vagus	Vagus nerve → esophageal plexus	Myenteric ganglionated plexus	Relaxation	
15	Stomach	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → gastric branches	Myenteric plexus	Contraction; increases tonus; peristalsis	2
16		Gastric glands	Dorsal motor nucleus of vagus	Vagus nerve → gastric branches	Submucosal plexus	Secretion	11
17	Pyloric sphincter	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → pyloric branches	Myenteric plexus	Inhibitory; diminishes tonus	12
18	Pancreas	Gland cells and blood vessels	Dorsal motor nucleus of vagus	Vagus nerve → pancreatic branches	Pancreatic ganglia	Secretion, vasodilation	12

*continued*



## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part II. PARASYMPATHETIC CONNECTIONS

	Organ	Effector	Preganglionic Neurones		Distribution of Postganglionic Neurones	Actions	Reference	
			Cell Body	Pathway				
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	
19	Liver	Hepatic cells, blood vessels, ducts	Dorsal motor nucleus of vagus	Vagus nerve → hepatic branches	Intramural ganglia	Secretion?	12	
20	Gallbladder	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → hepatic and cystic plexuses	Intramural ganglia	Emptying of gallbladder	5	
21	Biliary tree	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → hepatic and gastroduodenal plexuses	Intramural ganglia	Elevates pressure of bile ducts	5	
22	Sphincter of common bile ducts	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → hepatic and gastroduodenal plexuses	Intramural ganglia	Relaxation	5	
23	Small intestine	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → celiac and superior mesenteric branches	Myenteric plexus	Contraction, peristalsis	12	
24		Gland	Dorsal motor nucleus of vagus	Vagus nerve → celiac and superior mesenteric branches	Submucosal plexus	Secretion		
25	Ileocecal sphincter	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → celiac and superior mesenteric branches	Myenteric plexus	Inhibitory; diminishes tonus	12	
26	Proximal colon	Smooth muscle	Dorsal motor nucleus of vagus	Vagus nerve → celiac and superior mesenteric branches	Myenteric ganglionated plexus	Contraction, peristalsis	12	
27		Glands	Dorsal motor nucleus of vagus	Vagus nerve → celiac and superior mesenteric branches	Submucosal ganglionated plexus	Secretion		
28	Distal colon	Smooth muscle	S 2-4	Pelvic splanchnic nerves → hypogastric nerves → inferior mesenteric plexus	Myenteric plexus	Contraction, vasodilation; peristalsis	1	
29		Glands	S 2-4	Pelvic splanchnic nerves → hypogastric nerves → inferior mesenteric plexus	Submucosal plexus	Secretion		
30		Internal sphincter ani	S 2-4	Pelvic splanchnic nerves → inferior hypogastric plexus	Myenteric plexus	Inhibition, vasodilation		
31	Kidney	None demonstrated						12
32	Ureter	Smooth muscle	S 2-4	Pelvic splanchnic nerves → hypogastric nerves	Intramural ganglia	Contraction	9	
33	Bladder	Detrusor vesicae	S 2, 3, 4	Pelvic splanchnic nerves (nervi erigentes)	Inferior hypogastric plexus → vesical and intramural ganglia	Contraction	8, 10	
34		Sphincter vesicae	S 2, 3, 4	Pelvic splanchnic nerves (nervi erigentes)	Inferior hypogastric plexus → vesical and intramural ganglia	Relaxation	8	
35	Urethra	Smooth muscle	S 2, 3, 4	Pelvic splanchnic nerves	Inferior hypogastric plexus → vesical plexus → prostatic plexus	Control of sphincter	8	
36	Prostate gland	Smooth muscle	S 2, 3, 4	Pelvic splanchnic nerves	Inferior hypogastric plexus → vesical plexus → prostatic plexus	?	12	
37	Seminal vesicle and vas deferens	Smooth muscle	S 2, 3, 4	Pelvic splanchnic nerves	Inferior hypogastric and vesical plexuses	?	12	
38	Testis	None demonstrated						12

*continued*

## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part II. PARASYMPATHETIC CONNECTIONS

	Organ	Effector	Preganglionic Neurones		Distribution of Postganglionic Neurones	Actions	Reference	
			Cell Body	Pathway				
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	
39	Corpora cavernosa	Erectile tissue	S 2, 3, 4	Pelvic splanchnic nerves	Inferior hypogastric plexus → cavernous plexus	Vasodilation, erection	10	
40	Clitoris and labia minora	Erectile tissue	S 2, 3, 4	Pelvic splanchnic nerves	Inferior hypogastric plexus → uterovaginal plexus	Vasodilation, erection	10	
41	Vagina	None demonstrated						4
42	Cervix of uterus	Smooth muscle	S 2-4	Pelvic splanchnic nerves	Inferior hypogastric plexus	?	12	
43	Uterine tube	None demonstrated						12
44	Ovary	None demonstrated						12
45	Thyroid gland	No true secretory fibers present						12
46	Face	Blood vessels	Nucleus salivatorius superior ?	Nervus intermedius → 7th nerve	Vascular rami from 7th nerve to external carotid artery and its branches	Vasodilation, blushing	12	
47	Mucosal glands of palate	Gland cells	Nucleus salivatorius superior	Nervus intermedius → great superficial petrosal nerve → vidian nerve	Pterygopalatine ganglion and palatine nerves	Secretion	12	

*Contributors:* (a) Rogers, William M., (b) Mitchell, G. A. G.

*References:* [1] Alvarez, W. C. 1929. J. Am. Med. Assoc. 92:1231. [2] Cannon, W. B. 1911. Am. J. Physiol. 29:250. [3] Edinger, L. 1911. Vorlesungen über den Bau den nervösen Centralorgane des Menschen und der Thiere. F. C. W. Vogel, Leipzig. Aufl. 8, Bd. 1. [4] Gunn, J., and K. J. Franklin. 1920. Proc. Roy. Soc. (London), B, 94:197. [5] Kuntz, A. 1953. The autonomic nervous system. Ed. 4. Lea and Febiger, Philadelphia. [6] Langley, J. N. 1898. In E. A. Schäfer, ed. Textbook of physiology. Y. J. Pentland, Edinburgh. v. 1, p. 475. [7] Langley, J. N., and H. E. Anderson. 1892. J. Physiol. (London) 13:504. [8] Langworthy, O. R., L. C. Kolb, and L. G. Lewis. 1940. Physiology of micturition. Williams and Wilkins, Baltimore. [9] Mitchell, G. A. G. Unpublished. Univ. Manchester, London, 1963. [10] Müller, L. R. 1931. Lebensnerven und Lebenstrieb. J. Springer, Berlin. Aufl. 3. [11] Pavloff, I. P. 1910. The work of the digestive glands. Ed. 2. C. Griffin, London. [12] Rogers, W. M. Unpublished. Columbia Univ., New York, 1953. [13] Woollard, H. H. 1926. J. Anat. 60:345.

### Part III. GANGLIA

*Division* (column B): S = sympathetic; P = parasympathetic.

	Ganglion	Division	Location	Preganglionic Connections	Postganglionic Distribution
	(A)	(B)	(C)	(D)	(E)
1	Aorticorenal	S	Root of renal artery	Thoracic splanchnic nerves	Renal and aortic plexuses
2	Bronchial	P	Bronchial plexuses	Vagus nerves	Bronchial musculature and glands
3	Cardiac	P	Cardiac plexus	Vagus nerves	Heart, coronary vessels, pulmonary plexuses
4	Celiac	S	Celiac plexus	Thoracic splanchnic nerves	Abdominal viscera and blood vessels
5	Cervical sympathetic inferior	S	Sympathetic trunk, level of vertebra T 1	Upper thoracic nerves	Brachial plexus, inferior cervical cardiac nerve, common carotid plexus, vertebral nerves

*continued*

## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part III. GANGLIA

Ganglion	Division	Location	Preganglionic Connections	Postganglionic Distribution	
(A)	(B)	(C)	(D)	(E)	
6	Cervical sympathetic Middle	S	Sympathetic trunk, level of vertebra C 6	Upper thoracic nerves	Middle cervical cardiac nerve, sympathetic roots of nerves C 5, 6
7	Superior	S	Sympathetic trunk opposite 2nd, 3rd, 4th cervical vertebrae	Upper thoracic nerves	Internal, external carotid nerves, sympathetic roots of nerves C 1-3, superior cervical cardiac nerves
8	Cervical of uterus	P,S	Inferior hypogastric plexus adjacent to cervix of uterus	Pelvic splanchnic nerves, inferior hypogastric plexus	Uterus and vagina
9	Ciliary	P	Orbit, between optic nerve and lateral rectus muscle	Oculomotor	Short ciliary nerves
10	Enteric	P	Wall of enteric canal	Vagus and pelvic splanchnic nerves	Enteric muscles and glands
11	Impar (coccygeal)	S	Ventral surface of coccyx	Lumbar spinal nerves	Caudal spinal nerves
12	Intermediate	S	In relation to ventral nerve roots and communicating rami	Ventral nerve roots	Primary ventral rami of spinal nerves, visceral nerves
13	Lingual	P	Tongue	Facial and chorda tympani	Lingual glands
14	Mesenteric Inferior	S	Adjacent to inferior mesenteric artery	Lumbar splanchnic nerves	Inferior mesenteric and superior hypogastric plexuses
15	Superior	S	Adjacent to root of superior mesenteric artery	Thoracic splanchnic nerves	Superior mesenteric, aortic, and renal plexuses
16	Myenteric	P	Between longitudinal and circular enteric muscle layers	Vagus and pelvic splanchnic nerves	Enteric muscles
17	Otic	P	Medial to mandibular nerve, just below foramen ovale	Glossopharyngeal nerve	Supplies auriculotemporal nerve with fibers to parotid gland
18	Pelvic	P,S	Adjacent to pelvic viscera	Pelvic splanchnic nerves, inferior hypogastric plexus	Pelvic viscera and blood vessels, and vessels in external genitalia
19	Pterygopalatine <sup>1</sup>	P	Pterygopalatine fossa	Facial nerve	Pharyngeal, palatine, nasal, and orbital nerves
20	Pulmonary	P	Pulmonary plexuses	Vagus nerves	Bronchial plexuses
21	Splanchnic	S	Posterior mediastinum in relation to thoracic splanchnic nerves	Thoracic splanchnic nerves	Celiac plexus
22	Submandibular	P	Between lingual nerve and duct of submandibular gland	Chorda tympani (through facial)	Submandibular, sublingual, and lingual glands
23	Submucous	P	Submucosa of enteric canal	Vagus and pelvic splanchnic nerves	Enteric muscles and glands
24	Sympathetic trunk	S	Ventrolateral to vertebral column	Spinal nerves T 1 - L 2	Sympathetic roots of spinal nerves, cephalic sympathetic, cardiac and splanchnic nerves
25	Terminale	P	Adjacent to olfactory tract	Nervus terminalis	Anterior cerebral artery, vomeronasal organ, nasal mucosa
26	Tracheal	P	Tracheal wall	Vagus nerves	Tracheal and bronchial plexuses
27	Vertebral	S	Sympathetic trunk, level of vertebra C 8	Upper thoracic nerves	Ansa subclavia, inferior cervical cardiac nerve, vertebral nerves, sympathetic roots of nerves C 6, 7, 8

/1/ Supplied by greater petrosal through facial nerve.

*Contributors:* (a) Kuntz, Albert, (b) Mitchell, G. A. G.

*Reference:* Kuntz, A. 1953. The autonomic nervous system. Ed. 4. Lea and Febiger, Philadelphia.

*continued*



## 38. AUTONOMIC NERVOUS SYSTEM: MAN

### Part IV. PLEXUSES

Division (column B): S = sympathetic; P = parasympathetic.

	Plexus	Division	Origin	Distribution
	(A)	(B)	(C)	(D)
1	Aortic	S	Sympathetic trunks	Aorta and proximal portions of its branches
2	Cardiac	P,S	Cervical and thoracic sympathetic cardiac nerves, branches of vagus nerves	Heart, coronary vessels, anterior pulmonary plexuses
	Superficial			
3	Deep	P,S	Right superior, middle and inferior cervical and thoracic sympathetic cardiac nerves; all cardiac branches of right vagus, left middle and inferior cervical and thoracic cardiac nerves, superior cervical and cardiac branches of left vagus	Heart, coronary vessels, anterior pulmonary plexuses
4	Carotid Common	S	Sympathetic trunk	Common carotid artery
5	External	S	Superior cervical ganglion, common and external carotid plexuses	External carotid artery and its branches
6	Internal	S	Superior cervical ganglion, common and internal carotid plexuses	Internal carotid artery and its branches, caroticotympanic and deep petrosal nerves, cavernous plexus
7	Cavernous	S	Internal carotid plexus	Cavernous sinus, oculomotor, trochlear and ophthalmic nerves, ciliary ganglion, hypophysis cerebri
8	Celiac	S	Intrinsic ganglia, splanchnic nerves	Celiac artery and its branches
9	Colic	S	Inferior mesenteric plexus	Colon and rectum
10	Duodenal	S	Gastroduodenal, superior mesenteric, and pancreatic plexuses	Duodenum and pancreas
11	Enteric	P	Vagus, pelvic splanchnic nerves, intrinsic ganglia; esophageal, celiac, mesenteric, and pelvic plexuses	Enteric canal
12	External carotid	S	External carotid plexus	External carotid artery, submandibular and otic ganglia
13	Hepatic	P,S	Vagus nerves and celiac plexus	Biliary system, hepatic blood vessels
14	Hypogastric	S	Celiac, aortic, inferior mesenteric plexuses	Inferior hypogastric plexus
15	Spermatic	S	Aortic and renal plexuses	Spermatic artery, spermatic cord, testis
16	Mesenteric Inferior	S	Celiac plexus, lumbar splanchnic nerves	Inferior mesenteric artery and its branches
17	Superior	S	Celiac plexus	Superior mesenteric artery and its branches
18	Meningeal, middle	S	External carotid plexus	Middle meningeal artery
19	Ovarian	S	Aortic and renal plexuses	Ovarian artery, ovary
20	Pancreatic	P,S	Vagus nerves, intrinsic ganglia, sympathetic trunks	Pancreas, pancreatic ducts and vessels
21	Pelvic	P,S	Inferior hypogastric plexus, pelvic splanchnic nerves, intrinsic ganglia	Pelvic viscera and blood vessels, and vessels of erectile tissue
22	Pulmonary	P,S	Vagus nerves, intrinsic ganglia, sympathetic trunks	Bronchial plexuses, pulmonary vessels
23	Phrenic	S	Celiac plexus, lesser and least splanchnic nerves	Renal blood vessels; supplies suprarenal gland, diaphragm, esophagus, inferior vena cava
24	Prostatic	P,S	Inferior hypogastric plexus	Prostate gland
25	Renal	S	Celiac plexus, thoracic splanchnic nerves	Renal blood vessels
26	Sigmoid	S	Inferior mesenteric plexus	Sigmoid colon
27	Splenic	S	Celiac plexus	Spleen, pancreas, stomach
28	Suprarenal	S	Celiac plexus	Suprarenal artery and gland
29	Tympanic	P,S	Internal carotid nerve, ramus from geniculate ganglion	Tympanum, mastoid cells, auditory tube
30	Uterine	P,S	Inferior hypogastric plexus	Uterus
31	Vaginal	P,S	Inferior hypogastric plexus	Vagina
32	Vesical	P,S	Inferior hypogastric plexus	Urinary bladder

Contributors: (a) Kuntz, Albert, (b) Mitchell, G. A. G.

Reference: Kuntz, A. 1953. The autonomic nervous system. Ed. 4. Lea and Febiger, Philadelphia.



## 39. DIGESTIVE ENZYMES: VERTEBRATES

**Tissue or Secretion** (column D): T = tissue; S = secretion. **Symbols** (columns E-S): + = present; - = absent; ± = doubtful. For information on monkey, consult references 3, 17, 28; for mouse, references 18, 23.

	Species	Common Name	Organ	Tissue or Secretion	Enzyme													Reference							
					Amylase (Diastase)	Carbonic Anhydrase	Elastase	Enterokinase	Erepsin, Peptidase	Invertase (Saccharase)	Lipase, Esterases	Maltase	Pepsin	Phosphatase	Ribonuclease	Rennin (Chymosin)	Trypsin, Other Non-acid Proteases		Urease	β-D-Galactosidase					
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)					
1	<i>Homo sapiens</i>	Man	Salivary gland	T						+										5					
2				S	+					+						+						17			
3					Esophagus	T						+										5			
4					Stomach	T						+			+							K,5;M,N,17;R,10			
5				S									+			+			- <sup>1</sup>				K,24;M,17;P,11		
6					Pancreas	T					+	+											24		
7				S		+					+	+	+									+		E,L,Q,4;I,K,24	
8					Small intestine	T	+				+											+		E,17;I,24;N,23,26	
9				S		±					+	+	+	±											E,24;H-K,31
10					Cecum & colon	T																+		26	
11	<i>Bos taurus</i>	Cattle (cow)	Salivary gland	T							+										+	8			
12				S	+								+											E,8;K,17	
13					Esophagus	T										-								24	
14					Stomach	T	+						+						+					E,R,26;K,17;P,5	
15				S			±									+			± <sup>2</sup>						M,4;P,17
16					Pancreas	T	+						+										+		E,24;K,17;O,Q,5
17		S	+																			+		24	
18			Small intestine	T																	+		M,24;N,15		
19		S								+														17	
20	<i>Capra hircus</i>	Goat	Salivary gland	S	-																		24		
21					T																				26
22				S																			+		26
23					Small intestine	S	±				+	-	±	±									-		E,I-K,Q,31;H,17
24	<i>Ovis aries</i>	Sheep	Salivary gland	T							+											+	8		
25					S	-								+											17
26					Esophagus	T										-									24
27					Stomach	T							+						+						K,17;M,R,26;P,5
28				T		+								+						+					
29				S	+																		+		E,17;Q,26
30			Small intestine	S						+													17		
31			Cecum & colon	T								-											24		
32	<i>Sus scrofa</i>	Swine	Salivary gland	T							+											+	8		
33					S	+																			17
34					Esophagus	T										-									24
35					Stomach	T							+											+	L,M,17;K,13;R,10
36				S		-																			
37					Pancreas	T	+				+	+	+						+				+		E,I,K,Q,17;G,20;K,14;N,18;O,25
38		S	+							+	+											+		E,Q,24;G,20	
39			Small intestine	T						+	-		+										H,25;I,31;K,17;M,24;N,26		
40		S		+							+	±	±									-		E,J,K,Q,31;H,17	
41	<i>Equus caballus</i>	Horse	Salivary gland	S	±						+												E,24;K,26		
42					T									+											17

/1/ In adult. /2/ Only in young.

continued

*Contrails*  
39. DIGESTIVE ENZYMES: VERTEBRATES

Species	Common Name	Organ	Tissue or Secretion	Enzyme													Reference		
				Amylase (Diastase)	Carbonic Anhydrase	Elastase	Enterokinase	Erepsin, Peptidase	Invertase (Saccharase)	Lipase, Esterases	Maltase	Pepsin	Phosphatase	Ribonuclease	Rennin (Chymosin)	Trypsin, Other Non-acid Proteases		Urease	$\beta$ -D-Galactosidase
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)
43	<i>Equus caballus</i>	Horse	Small intestine	T															24
44				S <sup>a</sup>	+			+	+	-	+								E, J-L, Q, 1; H, 17
45			Cecum & colon	T	+														29
46	<i>Felis catus</i>	Cat	Salivary gland	T															26
47				S	-														17
48			Stomach	T		+					+								F, 4; K, 17; N, 26; R, 10
49				S															4
50			Pancreas	T		+			+		+								F, L, 4; I, 17; N, 26
51				S	+						+								4
52			Small intestine	T		+		+	±										F, 4; H, I, Q, 17; N, 23, 26
53				S	+			+	±	±	-								E, I, J, Q, 31; H, 17; K, 26
54			Cecum & colon	T							+								K, 22; N, 26
55			Colon	S							-								22
56	<i>Canis familiaris</i>	Dog	Salivary gland	T		+					+								F, 6; K, N, S, 8
57				S	+						+								E, 8; K, 17
58			Esophagus	T															24
59			Stomach	T		+					+								F, 4; K, 24; R, 10
60				S	±						+								E, K, 24; M, P, 3
61			Pancreas	T	+		+		+		+								E, I, K, Q, 4; G, 7; O, 25
62				S	+				+		+								4
63			Small intestine	T				+	+										H, 4; I, K, 24; N, 26
64				S	+			+	+	±	±								E, J, K, Q, 31; H, I, 4
65			Cecum & colon	S	±			-	±		+								E, K, 24; H, I, 19
66	<i>Cavia porcellus</i>	Guinea pig	Salivary gland	S	+														17
67			Stomach	T		+					+								F, 18; K, 17
68			Pancreas	T		+	+		+										F, 18; G, 9; I, 17
69				S							+								24
70			Small intestine	T					+										I, 17; N, 23
71	<i>Rattus spp.</i>	Rat	Salivary gland	T							+								8
72				S	+														27
73			Stomach	T	+	+					+								E, 21; F, 4; K, 17; M, 26; R, 10
74				S															16
75			Pancreas	T	+	+	+				+								E, F, K, Q, 4; G, 9
76				S	+														4
77			Small intestine	T	+	+		+	+			+							E, I, L, 30; F, 4; H, N, 23, 26
78			Colon	T	+														21
79	<i>Oryctolagus cuniculus</i>	Rabbit	Salivary gland	T							+								8
80				S	+														8
81			Stomach	T		+					+		+	+					F, 4; K, 17; M, 24; N, 26; R, 10
82				S							+								24
83			Pancreas	T		+			+										F, 4; I, 17
84				S	+						+								4

/<sub>2</sub>/ Only in young. /<sub>3</sub>/ Lactase also present [1].

continued

39. DIGESTIVE ENZYMES: VERTEBRATES

Species	Common Name	Organ	Tissue or Secretion	Enzyme														Reference			
				Amylase (Diastase)	Carbonic Anhydrase	Elastase	Enterokinase	Erepsin, Peptidase	Invertase (Saccharase)	Lipase, Esterases	Maltase	Pepsin	Phosphatase	Ribonuclease	Rennin (Chymosin)	Trypsin, Other Non-acid Proteases	Urease		β-D-Galactosidase		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)		
85	<i>Oryctolagus cuniculus</i>	Rabbit	Small intestine	T	+			+				-	+						F,4;I,M,24;N,26		
86			Cecum & colon	T	+			+	-		±									E,I,K,Q,31;H,17	
87				T								+							26		
88	<i>Gallus domesticus</i>	Chicken	Salivary gland	S	+														17		
89			Stomach	T								+	+							K,17;M,5	
90			Pancreas	T	+																17
91			Small intestine	T	+										+						E,17;N,23
92	<i>Rana</i> spp.	Frog	Salivary gland	S	+														17		
93			Esophagus	T									+							24	
94			Stomach	T										+					+	M,17;R,10	
95				S										+							17
96			Pancreas	T						+			+						+		I,17;L,Q,26
97			Small intestine	T						+			-								I,17;L,26

Contributor: Hollander, Franklin

References: [1] Alexander, F., and A. K. Chowdhury. 1958. *Nature* 181:190. [2] Aqvist, S. E. G., and C. B. Anfinsen. 1959. *J. Biol. Chem.* 234:1112. [3] Babkin, B. P. 1929. *Die äussere Sekretion der Verdauungsdrüsen*. Ed. 2. J. Springer, Berlin. [4] Babkin, B. P. 1950. *Secretory mechanism of the digestive glands*. Ed. 2. P. B. Hoeber, New York. [5] Boyer, P. D., H. Lardy, and K. Myrbäck, ed. 1959-63. *The enzymes*. Ed. 2. Academic Press, New York. [6] Brusilow, S. W., and C. L. Diaz. 1962. *Am. J. Physiol.* 202:158. [7] Carter, A. E. 1956. *Science* 123:669. [8] Chauncey, H. H., and G. Quintarelli. 1961. *Am. J. Anat.* 108:263. [9] Cohen, H., H. Megel, and W. Kleinberg. 1958. *Proc. Soc. Exptl. Biol. Med.* 97:8. [10] Conway, E. J. 1953. *The biochemistry of gastric acid secretion*. C. C. Thomas, Springfield, Ill. [11] Dotti, L. B., and I. S. Kleiner. 1942. *Am. J. Physiol.* 138:557. [12] Dukes, H. H. 1955. *The physiology of domestic animals*. Ed. 7. Comstock, Ithaca. [13] Evans, R. A., and D. A. Stansfield. 1961. *Nature* 190:1110. [14] Gjessing, E., and J. C. Hartnett. 1960. *Federation Proc.* 19:49. [15] Harris, E. S., et al. 1952. *Proc. Soc. Exptl. Biol. Med.* 81:593. [16] Hirschowitz, B. I., and W. G. Underhill. 1959. *Am. J. Physiol.* 196:837. [17] Koningsberger, V. J., E. J. Slijper, and H. J. Vonk, ed. 1946. *Tabulae Biologicae* 21(1). [18] Kurata, Y. 1953. *Stain Technol.* 28:231. [19] Kuvavaeva, I. B. 1957. *Fiziol. Zh. SSSR* 43:311. [20] Lewis, U. J., D. E. Williams, and N. G. Brink. 1956. *J. Biol. Chem.* 222:705. [21] McGeachin, R. L., and K. F. Norwood, Jr. 1959. *Am. J. Physiol.* 196:972. [22] Martin, B. F. 1959. *Nature* 183:1464. [23] Moog, F. 1962. *Federation Proc.* 21:51. [24] Oppenheimer, C. 1925-26. *Die Fermente und ihre Wirkungen*. Ed. 5. G. Thieme, Leipzig. v. 1, 2. [25] Oppenheimer, C., and L. Pincussen, ed. 1929. *Ibid.* G. Thieme, Leipzig. v. 3. [26] Oppenheimer, C. 1935-39. *Ibid.* W. Junk, Haag. suppl. [27] Schneyer, L. H., and C. A. Schneyer. 1956. *Federation Proc.* 15:164. [28] Smith, G. P., and F. P. Brooks. 1959. *Ibid.* 18:147. [29] Sym, E. A., W. Stankiewicz, and F. Zielinski. 1939. *Enzymologia* 6:113. [30] Van Genderen, H., and C. Engel. 1938. *Ibid.* 5:71. [31] Wright, R. D., et al. 1940. *Quart. J. Exptl. Physiol.* 30:73.

For a comprehensive review of the

Component (A)	Mammalia			Aves (E)
	<i>Homo sapiens</i> (B)	<i>Bos taurus</i> (C)	<i>Canis familiaris</i> (D)	
1 Sinus venosus	Sinus venosus completely incorporated into right atrium <sup>2</sup> . Sinoatrial node same as in <i>Bos</i> and <i>Canis</i> .	Incorporated into right atrium. Sinoatrial node, or "pacemaker," is specialized tissue in atrial region (was part of sinus in embryo).		Largely included in right atrium.
2 Sinoatrial junction	Vestige of embryonic sinoatrial junction found in adult right atrium.	Right valve of embryonic sinus becomes valve of inferior vena cava and of coronary sinus.		Embryonic right sinoatrial valve becomes valve of inferior vena cava.
3 Atrium	Complete interatrial septum except for small remnant of foramen ovale. Sinoatrial and atrioventricular nodes connected by typical cardiac fibers of right atrium.	Complete interatrial septum. Histological characteristics of sinoatrial and atrioventricular nodal fibers unusually clear.	Complete interatrial septum. Sinoatrial node near opening of superior vena cava, atrioventricular node near opening of coronary sinus.	Complete interatrial septum. Extensive distribution of specialized conduction fibers.
4 Atrioventricular junction	Atrioventricular ring of dense connective tissue. Right atrioventricular valve is tricuspid, left valve is bicuspid. Atrioventricular bundle of His present from atrioventricular node to ventricles. In the adult, usually no muscular atrioventricular connections other than the bundle of His [9].	Connective tissue of atrioventricular ring (annulus fibrosus) contains bone. Atrioventricular bundle is usually distinct. Right atrioventricular valve usually tricuspid, left valve usually bicuspid.	Atrioventricular valves the same as in <i>Bos</i> . Ring of connective tissue separates atrial and ventricular muscle, except for atrioventricular bundle of His.	Atrial and ventricular muscles separated by ring of connective tissue, except for atrioventricular bundles. Right atrioventricular valve is large and muscular, left valve is bicuspid.
5 Ventricle	Complete interventricular septum, membranous in uppermost reaches.	Complete septum. Histological characteristics of Purkinje (conduction) fibers distinct.	Complete ventricular septum.	Complete ventricular septum.
6 Conus arteriosus	Proximal portion incorporated in right ventricle, distal incorporated in left aortic and pulmonary trunks.			Proximal portion incorporated in right ventricle, distal incorporated in right aortic and pulmonary trunks.
7 Truncus arteriosus <sup>6</sup>	Divided into aortic and pulmonary trunks. Semilunar valves at embryonic juncture of conus and truncus.			

/1/ Data for Crocodylia from White [13]. /2/ Data from Favaro [3]. /3/ Some authorities are of the opinion that a specialized conduction expanded portion of the ventral aorta lying within the pericardial cavity; also known as the bulbus arteriosus, or

*Contributors:* (a) Copenhaver, Wilfred M., (b) Andrew, Warren, (c) Monie, I. W., (d) White, Fred N., (e) Grodziński,

*References:* [1] Arey, L. B. 1954. *Developmental anatomy*. Ed. 6. W. B. Saunders, Philadelphia. [2] Davies, F., *Klassen und Ordnungen des Thier-Reichs*. C. F. Winter, Leipzig. Bd. 6. [4] Goodrich, E. S. 1930. *Studies on the*



**CIRCULATORY SYSTEM: VERTEBRATES**

**HEART**

subject, consult references 4, 8, 10.

Reptilia <sup>1</sup>	Amphibia	Pisces and Chondrichthyes	Agnatha <sup>2</sup>	
(F)	(G)	(H)	(I)	
Partly incorporated into right atrium, and is distinct only internally. Chelonia: Origin of beat is dependent on intrinsic ganglia [6].	Salientia: Separate chamber and relatively smaller than in Caudata. Caudata: Thin-walled, triangular-shaped chamber, shifted toward right side.	Smooth, thin-walled chamber into which systemic veins open. Cardiac muscle has high intrinsic contraction rate and acts as the "pacemaker" (although contraction is myogenic in origin, rate of beat is under nervous control, being depressed by vagal stimulation and accelerated by sympathetic stimulation).	Thin-walled, elongated sac or tube into which systemic veins open.	1
Approximately the same as in Pisces.		Sinoatrial opening guarded by pair of valves. Cardiac muscle continuous from sinus to atrium.	Sinoatrial opening guarded by pair of valves.	2
Same as in Salientia, but interatrial septum contributes to valves dividing atrioventricular opening in two.	Salientia: Complete interatrial septum. Caudata: Incomplete septum partially dividing chamber bilaterally, with sinus venosus opening into right atrium and single pulmonary vein into left atrium.	Thin, reticulate-walled chamber; no division into right and left sides, no pulmonary veins. Dipnoi: Partial septation. Pulmonary vein enters to left and sinus venosus to right of septum.	Single muscular sac. Atrium lateral to ventricle.	3
Crocodylia: Atrioventricular ring contains cartilage extending into base of right atrioventricular valve and similar valve of right aortic arch. Chelonia: Continuity of muscle, as in teleosts. Presence of "Purkinje-like" fibers controversial. <sup>4</sup>	Continuity of atrial and ventricular muscles, as in Pisces. Salientia: Four valve cusps. Caudata: Atrioventricular valve of two or four muscular thickenings.	Atrioventricular valve composed of two cusps. Cardiac muscle of atrium continuous with that of ventricle around entire circumference of atrioventricular junction. Dipnoi: Fibrocartilaginous plug regulates blood flow.	Atrioventricular channel connects both vesicles. Two valves present.	4
Serpentes, Sauria, Chelonia: Ventricle partially divided by incomplete septum. Crocodylia: Complete ventricular septum containing cartilage.	No ventricular septum.	Thick-walled chamber; network of muscular trabeculae; no division into right and left sides. Dipnoi: Septum present.	Thick-walled muscular sac with smooth internal surface.	5
Conus incorporated in ventricle and in arterial trunks.	Semilunar valves; also a "spiral valve" coursing lengthwise.	Semilunar valves present. Pisces: Small. Dipnoi: Divided into dorsal and ventral channels. Chondrichthyes: Relatively large.	Absent, unless two valves may be regarded as remnants.	6
Crocodylia: Right and left systemics connected by foramen of Panizza. Chelonia: Divided into right and left systemic and pulmonary trunks.	Salientia: Divided both internally and externally into right and left channels. Caudata: Divided internally into right and left channels.	Pisces: Enlarged in species having reduced conus in shape of bulbus arteriosus. Dipnoi: Divided into three paired channels.	Part bordering on ventricle enlarged to form bulbus arteriosus.	7

small part of the sinus venosus is also incorporated in the left atrium [5]. /4/ The view that "Purkinje-like" fibers system of Purkinje fibers is neomorphic for birds and mammals is supported by Davies and Francis [2]. /s/ An aortic sac [1, 7, 11].

Z., (f) DeGaris, Charles F., (g) Moog, Florence, (h) Ballard, W. W.

and E. T. B. Francis. 1946. Biol. Rev. Cambridge Phil. Soc. 21:173. [3] Favaro, G. 1901. In H. G. Bronn, ed. structure and development of vertebrates. Macmillan, London. [5] Hamilton, W. J., J. D. Boyd, and H. W. Mossman.

*continued*

1962. Human embryology. Ed. 3. W. Heffer, Cambridge, England. [6] Heinbecker, P., and G. H. Bishop. 1935. [8] Kingsley, J. S. 1926. Outlines of comparative anatomy of vertebrates. Ed. 3. Blakiston, Philadelphia. [9] Kistin, New York. [11] Patten, B. M. 1953. Human embryology. Ed. 3. Blakiston, New York. [12] Robb, J. S. 1953.

Part II. BLOOD

Vessel	Mammalia	Aves	Reptilia
(A)	(B)	(C)	(D)
<b>Main Longitudinal Vessels</b>			
1 Aorta and caudal artery	Continuous vessels located ventral to the axial skeleton, extending from the region of the heart to the tip of the tail. Caudal artery enclosed in the hemal channel.		
2 Aortic arches	Left aortic arch only.	Right aortic arch only.	One pair of aortic arches.
3 Carotid arteries	Various connections of internal and external carotids with aortic arch: innominate artery, common carotid artery, carotid trunk.	Internal and external carotids originate from common carotid arteries, which may be symmetrical, asymmetrical, fused, or one may be obliterated.	Internal and external carotid arteries originate from common carotids. Many Serpentes: Only one common carotid. Crocodylia: Carotid duct absent.
4 Posterior cardinal veins	In all embryos, two posterior cardinal veins are present, being continuous with the caudal vein. No renal portal system. Posterior cardinals obliterated and replaced by supracardinal derivatives, the azygos and hemiazygos veins. Posterior vena cava is main trunk vein.	Rudimentary caudal and renal portal veins. Renal veins join femoral and internal iliac veins in kidney, which in turn join to form posterior vena cava. Posterior vena cava receives hepatic vein; no posterior cardinal veins persist.	Formation of kidneys causes great changes in distribution of posterior cardinals (renal portal system). Sauria: Caudal vein empties into two renal portal veins. Two hepatic veins join posterior vena cava. Crocodylia: Posterior vena cava emerges as single vessel from kidneys.
5 Anterior cardinal veins	Two anterior venae cavae, each consisting of an anterior cardinal vein, inferior jugular vein, and subclavian vein. Primates, Carnivora: Section of left vein, located between heart and anastomosis, disappears. Ungulata: Right vena cava more prominent than left because blood passes from left into right by anastomosis. Rodentia, Insectivora: Innominate veins of equal size.	Two anterior venae cavae, each consisting of an anterior cardinal vein, vertebral vein, and subclavian vein. Right cardinal vein more prominent than left. Inferior jugular vein absent.	Sauria: Similar in structure to <i>Sphenodon</i> . <i>Sphenodon</i> : Two anterior cardinal veins (internal jugular veins). Inferior jugular veins (external) reduced and replaced by tracheal vein; these veins join subclavian vein to form anterior vena cava.
<b>Main Segmental Vessels<sup>1</sup></b>			
6 Dorsal segmental vessels	In each myoseptum, one artery and one vein are present. Tributaries are ventral medullary arteries and veins, and muscular arteries and veins.		
7 Dorsal rami			Serpentes: Longitudinal dorsal vein present.
8 Ventral rami	Segmental vessels run from the main longitudinal vessels (aorta, cardinal veins) in horizontal myoseptum and reach skin. Tributaries are muscular arteries and veins. Some segmental become intercostal arteries and veins that surpass main vessels in size. In region of paired limbs, segmental vessels become subclavian and iliac arteries and veins.		

<sup>1</sup>/ In all embryos, dorsal, lateral, and ventral segmental arteries and veins are present. They persist at least

**CIRCULATORY SYSTEM: VERTEBRATES**

**HEART**

Am. J. Physiol. 114:212. [7] Hyman, L. H. 1947. Comparative vertebrate anatomy. Univ. Chicago Press, Chicago. A. D. 1949. Am. Heart J. 37:849. [10] Nelsen, O. E. 1953. Comparative embryology of the vertebrates. Blakiston, Am. J. Physiol. 172:7. [13] White, F. N. 1956. Anat. Record 125:417.

**VESSELS**

Amphibia (E)	Pisces and Chondrichthyes (F)	Agnatha (G)	
<b>Main Longitudinal Vessels</b>			
Continuous vessels located ventral to the axial skeleton, extending from the region of the heart to the tip of the tail. Caudal artery enclosed in the hemal channel; aorta similarly enclosed in <i>Acipenser</i> (sturgeon) only.			1
One to four pair of aortic arches.	Mainly four to five epibranchial arteries. Dipnoi: Pulmonary arteries present.	Six, seven, or more epibranchial arteries.	2
Internal and external carotid arteries. Salientia: Carotid duct absent.	Internal carotid arteries.	Petromyzones: Internal carotid artery dorsal, external carotid artery ventral. Myxini: Internal and external carotid arteries.	3
In all embryos, two posterior cardinal veins are present, being continuous with the caudal vein. Formation of kidneys causes great changes in distribution of posterior cardinals (renal portal system). Posterior cardinals rudimentary. Well-developed posterior vena cava drains kidneys. Caudata: Renal portal blood derived from iliac and caudal veins.	Pisces: Posterior cardinals often asymmetrical but receive renal and hepatic veins. Renal portal blood derived from caudal vein and/or segmental veins; caudal vein often leads directly to posterior cardinal. Dipnoi: Caudal vein, renal portal veins, asymmetric posterior cardinal vein, and a posterior vena cava. Chondrichthyes: Caudal vein empties into renal portal veins; hepatic veins empty into posterior cardinal veins, which are sinus-like distentions.	Renal portal system absent. Petromyzones: Same as in embryo. Myxini: Same as in embryo, but right posterior cardinal vein much thinner than left.	4
Salientia: Anterior cardinal veins (internal jugulars) and inferior jugular veins fuse into one short trunk on each side and enter sinus venosus by intermediation of anterior vena cava (cranial). Caudata: Two anterior cardinal veins (internal jugulars), two inferior jugular veins.	Pisces: Two cardinal veins. Mainly one internal jugular vein joining sinus venosus. Dipnoi: Two cardinal and two jugular veins. Chondrichthyes: Anterior cardinal veins (internal jugulars) are sinus-like distentions. Inferior jugular veins open into duct of Cuvier.	Petromyzones: Anterior cardinal veins unite to form common trunk and open into sinus venosus. Inferior jugular vein (ventral) present. Myxini: Left anterior cardinal vein (internal jugular) and inferior jugular vein (ventral) join sinus venosus. Right anterior cardinal vein opens into cor portale of liver.	5
<b>Main Segmental Vessels<sup>1</sup></b>			
Arteries and veins alternate in successive segments; their dorsal tips join to form the longitudinal dorsal trunk, and may persist in adults. Tributaries of dorsal segmental vessels are ventral medullary arteries and veins, and muscular arteries and veins.			6
Deep pinnal arteries and veins are segmental vessels, supplying dorsal fins.			7
Segmental vessels run from the main longitudinal vessels (aorta, cardinal veins) in horizontal myoseptum and reach skin. Tributaries are muscular arteries and veins. Some segmental become intercostal arteries and veins that surpass main vessels in size. In region of paired fins and limbs, segmental vessels become subclavian and iliac arteries and veins.			8

partially in adults and give rise to some longitudinal as well as other important stems.

*continued*

Vessel	Mammalia	Aves	Reptilia
(A)	(B)	(C)	(D)
Main Segmental Vessels <sup>1</sup>			
Dorsal segmental vessels			
9 Lateral cutaneous vein <sup>2</sup>			Sauria: Lateral cutaneous vein probably present in all species.
10 Abdominal vein <sup>2</sup>	Only anterior root of abdominal vein persists and is represented by umbilical vein in the embryo.		Abdominal vein, double or single, connects renal portal with hepatic portal system.
11 Epigastric arteries and veins <sup>2</sup>	Two distinct trunks running close to each other are located on the inner surface of the abdominal wall between longitudinal abdominal and segmental trunk muscles. Probably absent in birds.		
12 Lateral segmental vessels	Paired lateral vessels supply pronephros, mesonephros, and, when present, metanephros. Their number is much reduced in higher forms when forming renal and gonadal arteries.		
13 Ventral segmental vessels	In all embryos, ventral segmental vessels originate from main longitudinal vessels (aorta, caudal artery, caudal vein). In trunk, only arteries (mesenteric) originate from main longitudinal vessels, and in tail both arteries and veins. Paired arteries frequently unite to form a single median vessel.		
14 Arteries	Celiac artery, anterior mesenteric artery (cranial or superior), posterior mesenteric artery (caudal or inferior).	Celiac artery, anterior mesenteric artery (superior), posterior mesenteric artery (inferior).	Serpentes: Many arteries (mesenteric) reach intestine from the aorta ( <i>Boa</i> ). Sauria: Gastric, celiomesenteric, and posterior mesenteric arteries. Crocodylia: Gastroesophageal, celiomesenteric, and mesenteric arteries. <i>Sphenodon</i> : Gastric, celiac, common mesenteric (anterior), and posterior mesenteric arteries.
15 Veins	Subintestinal vein, which gets some blood directly from caudal vein, develops with invasion of intestine by ventral segmental arteries. Subintestinal vein is prominent vessel in embryonic fish and amphibians; less significant in birds and reptiles; remnants found in <i>Didelphis</i> ; parallels ventral border of intestinal tube and enters right omphalomesenteric vein. Subintestinal and parts of right and left omphalomesenteric veins form the hepatic portal vein. Omphalomesenteric veins and umbilical vein participate in liver circulation; hepatic vein opens into posterior vena cava; subintestinal vein is incorporated into trunk of hepatic portal vein.		
Vessels of the Forelimb			
16 Arteries	Right subclavian artery arises from left aortic arch, chiefly as innominate artery, and left subclavian directly from arch or truncus communis. Axillary artery, brachial artery, median artery (main vessel in most mammals), ulnar artery (main vessel in Prosimii), radial artery, interosseal artery (main vessel in <i>Ornithorhynchus</i> ). Metacarpal arteries, dorsal and volar digital arteries.	Subclavian artery, with carotid artery, originates from right aortic arch (innominate artery). Axillary, brachial, ulnar (main vessel), interosseal, radial, and ulnar nerve arteries; metacarpal and digital arteries.	Sauria: Both subclavian arteries arise from right aortic arch. Axillary, brachial, interosseal (main vessel), ulnar, radial, and median arteries; metacarpal and digital arteries. Chelonia: Subclavian artery, with carotid artery, originates from innominate artery. Brachial artery replaced by lateral brachial artery. Two arterial arches of hand, dorsal and volar.

<sup>1/1</sup> In all embryos, dorsal, lateral, and ventral segmental arteries and veins are present. They persist at least originating from ventral rami of dorsal segmental vessels.



## CIRCULATORY SYSTEM: VERTEBRATES

### VESSELS

Amphibia (E)	Pisces and Chondrichthyes (F)	Agnatha (G)	
<b>Main Segmental Vessels<sup>1</sup></b>			
Gymnophiona, Caudata, Dipnoi, Chondrichthyes: Lateral cutaneous vein in lateral line groove, below skin from tail to region of forelimb. Originates from end tips of lateral segmental veins. Salientia: Great and small cutaneous vein.			9
Extends from region of cloaca to shoulder girdle, where it merges into sinus venosus or into hepatic portal system. Salientia, Caudata: A single median, ventral abdominal vein. Chondrichthyes: Two lateral abdominal veins, one on each side of body wall.			10
Two distinct trunks running close to each other are located on the inner surface of the abdominal wall between longitudinal abdominal and segmental trunk muscles.		Myxini: Netlike track.	11
Paired lateral vessels supply pronephros and mesonephros.			12
In all embryos, ventral segmental vessels originate from main longitudinal vessels (aorta, caudal artery, caudal vein). In trunk, only arteries (mesenteric) originate from main longitudinal vessels, and in tail both arteries and veins. Paired arteries frequently unite to form a single median vessel.			
Salientia: Only celiomesenteric artery. Caudata: Celiac artery and about 13 mesenteric arteries, almost segmental in arrangement, distributed to intestine ( <i>Siren</i> ). Segmental arteries fuse into complex vessels ( <i>Menobranchus</i> , <i>Cryptobranchus</i> , <i>Salamandra</i> ).	Pisces: Mainly one celiomesenteric artery. Dipnoi: Celiac artery, two or three mesenteric arteries. Chondrichthyes: Celiac artery, two or three mesenteric arteries.	Petromyzones: Only one artery persists as celiomesenteric artery. Myxini: Mesenteric arteries (approximately 35 in number) distributed to intestine as segmental vessels.	14
Subintestinal vein, which gets some blood directly from caudal vein, develops with invasion of intestine by ventral segmental arteries. Subintestinal vein is prominent vessel in embryonic fish and amphibians; parallels ventral border of intestinal tube and enters right omphalomesenteric vein. Subintestinal and parts of right and left omphalomesenteric veins form the hepatic portal vein; vitelline vein (omphalomesenteric) may participate in liver circulation. Hepatic vein opens into posterior vena cava. Subintestinal vein is incorporated in trunk of hepatic portal vein.		Petromyzones: Subintestinal vein well-developed. Myxini: Subintestinal vein reduced.	15
<b>Vessels of the Forelimb</b>			
Salientia: Subclavian arteries originate from aortic arch, brachial, and deep brachial arteries. Other vessels same as in Caudata. Caudata: Subclavian artery arises from median aorta. Brachial, interosseal (main vessel), radial (radiomarginal), and ulnar (ulnomarginal) arteries. Dorsal arterial arch of hand, metacarpal and digital arteries.	Pisces: Subclavian artery originates from median aorta. Basal arteries, interrarial arteries. Chondrichthyes: Subclavian artery arises from median aorta. Lateral and medial pterygial artery, adradial arteries.	Forelimbs absent.	16

partially in adults and give rise to some longitudinal as well as other important stems. /2/ Longitudinal vessels

*continued*

Vessel	Mammalia	Aves	Reptilia
(A)	(B)	(C)	(D)
Vessels of the Forelimb			
17 Veins	Comparative anatomy of these veins awaits revision: digital veins, metacarpal veins, volar and dorsal venous arch, basilic vein (ulnomarginal), cephalic vein (radiomarginal), ulnar vein, radial vein, medial vein, brachial vein, axillary vein, subclavian vein. (The above list valid for five-fingered appendage.)	Metacarpal veins; basilic vein (main vessel), ulnar vein, interosseal vein, radial vein (radiomarginal), brachial vein. Subclavian vein enters anterior vena cava.	Sauria: Digital veins, dorsal venous arch of hand, radial vein (main vessel, marginal radial vein in embryos), ulnar vein, interosseal vein, brachial vein and lateral brachial vein. Lateral cutaneous vein empties into axillary vein. Subclavian vein enters internal jugular vein. <i>Alligator, Emys</i> : Only one brachial vein.

*Contributors*: (a) Grodziński, Z., (b) Monie, I. W., (c) DeGaris, Charles F., (d) Moog, Florence, (e) Ballard, W. W.

*References*: [1] Brash, J. C., ed. 1951. *Cunningham's Textbook of anatomy*. Ed. 9. Oxford Univ. Press, London. 1933. In L. Bolk, ed. *Handbuch der vergleichenden Anatomie der Wirbeltiere*. Urban and Schwarzenberg, Berlin. [5] Górkiewicz, C. 1947. *Bull. Intern. Acad. Sci. Cracovie*, p. 241. [6] Grodziński, Z. 1926. *Ibid.*, p. 955. B, 1:110. [9] Grodziński, Z. 1933. *Bull. Intern. Acad. Sci. Cracovie*, pp. 243, 259, 321. [10] Grodziński, Z. 1938. 1946. *Bull. Intern. Acad. Sci. Cracovie*, pp. 1, 22. [12] Grodziński, Z. 1948. *Ibid.*, p. 61. [13] Hafferl, A. 1933. p. 563. [14] Nelsen, O. E. 1953. *Comparative embryology of the vertebrates*. Blakiston, New York. [15] Sikorowa, 1947. *Bull. Intern. Acad. Sci. Cracovie*. p. 145. [18] Weidenreich, F. 1933. In L. Bolk, ed. *Handbuch der*

Part III.

Component	Mammalia	Aves	Reptilia
(A)	(B)	(C)	(D)
1 Lymph hearts	Absent even in embryos.	In all embryos only one pair of posterior hearts, located in region between pelvis and femur. Hearts persist in some adults ( <i>Alca, Anser, Casuaris, Podiceps, Struthio</i> ).	Only posterior hearts. Serpentes: Two elongated vesicles, each surrounded by bifurcated transverse processes of four-to-five caudal vertebrae. Sauria: Two ovoid vesicles attached to both ends of transverse process of first caudal vertebra. Crocodylia, Chelonia: Spherical in shape.
2 Lymph sacs (other than hearts), lymph sinuses	Jugular and iliac lymph sacs in embryos only.		Serpentes: Mandibular sinus. Sauria: Retrocardial, axillary, jugular, tracheal, and thyroidal sinuses. Crocodylia: Absent. Chelonia: Jugular cistern.
3 Lymph nodes	Many lymph nodes: approximately 465 in man, 300 in cattle, 60 in dog, 8,000 in horse.	Microscopically discernible nodes in walls of lymph vessels. Anseriformes: Two cervicothoracic and two lumbar lymph glands macroscopically visible.	Chelonia: Small nodes in lower eyelid.

/1/ Cole [7] and Favaro [8] regard these sacs as venous hearts.

## CIRCULATORY SYSTEM: VERTEBRATES

### VESSELS

Amphibia	Pisces and Chondrichthyes	Agnatha	
(E)	(F)	(G)	
Vessels of the Forelimb			
<p><b>Salientia:</b> Digital veins, dorsal venous arch of hand, ulnar vein (ulnomarginal), radial vein (radiomarginal), interosseal vein. Brachial vein and great cutaneous vein unite to join subclavian vein which enters anterior vena cava (cranial). Deep brachial vein continues as subscapular vein which, together with internal jugular vein, forms brachiocephalic vein.</p> <p><b>Caudata:</b> Digital veins, interosseal vein (main vessel), ulnar vein (ulnomarginal), radial vein (radiomarginal), brachial vein. Subclavian vein, together with lateral cutaneous vein, enters sinus venosus.</p>	<p><b>Pisces:</b> Interradial veins, basal vein. Subclavian vein enters duct of Cuvier or posterior cardinal vein. Left subclavian joins abdominal vein and epigastric vein, right subclavian joins only epigastric vein (<i>Salmo</i>).</p> <p><b>Chondrichthyes:</b> Aduial veins, lateral and medial pterygial vein. Subclavian vein fuses with epigastric vein and enters duct of Cuvier.</p>	<p>Forelimbs absent.</p>	17

- [2] Francis, E. T. B. 1934. The anatomy of the salamander. Oxford Univ. Press, London. [3] Gelderen, C. A. v. 6, p. 685. [4] Goodrich, E. S. 1930. Studies on the structure and development of vertebrates. Macmillan, London. [7] Grodziński, Z. 1928. Ibid., p. 417. [8] Grodziński, Z. 1928. Mem. Acad. Polon. Sci. Classe Sci. Math. Nat., In H. G. Bronn, ed. Klassen und Ordnungen des Thier-Reichs. C. F. Winter, Leipzig. Bd. 6. [11] Grodziński, Z. In L. Bolk, ed. Handbuch der vergleichenden Anatomie der Wirbeltiere. Urban and Schwarzenberg, Berlin. v. 6, L. 1947. Bull. Intern. Acad. Sci. Cracovie, p. 299. [16] Stephen, F. 1954. Traite Zool. 12:854. [17] Szarski, H. vergleichenden Anatomie der Wirbeltiere. Urban and Schwarzenberg, Berlin. v. 6, p. 375.

### LYMPHATICS

Amphibia	Pisces and Chondrichthyes	Agnatha	
(E)	(F)	(G)	
<p><b>Gymnophiona:</b> Approximately 100 spherical vesicles in trunk and tail, beneath skin in lateral line groove.</p> <p><b>Salientia:</b> One pair of anterior and one-to-four pair of posterior hearts.</p> <p><b>Caudata:</b> Ten to twenty rounded vesicles on each side of trunk, located as in <i>Gymnophiona</i>.</p>	<p><b>Pisces:</b> Two elongated vesicles joined by a channel and located at base of tail. Vesicles not pulsating in ganoid fishes.</p> <p><b>Chondrichthyes:</b> Absent.</p>	<p><b>Petromyzones:</b> Absent.</p> <p><b>Myxini:</b> One pair of pulsating sacs located in tail<sup>1</sup>.</p>	1
<p><b>Salientia:</b> In tadpoles, mandibular, circumoral, pericardial, and temporal sinuses; in adults, several subcutaneous sacs.</p> <p><b>Caudata:</b> Orbital sinus, sinus lymphaticus cordis, axillary sinus in larvae only.</p>	<p><b>Pisces:</b> Pectoral pinneal, orbital, cephalic, occipital, and lateral sinuses.</p> <p><b>Chondrichthyes:</b> Absent.</p>	<p><b>Petromyzones:</b> Supralabial, orbital, ocular ring, and deep labial sinuses.</p> <p><b>Myxini:</b> Three subcutaneous sacs underlie entire skin.</p>	2
			3

continued

## 40. COMPARATIVE ANATOMY OF THE

Part III.

Component	Mammalia	Aves	Reptilia
(A)	(B)	(C)	(D)
4 Subvertebral lymphatic trunks (thoracic duct)	One or two trunks associated with aorta and caudal artery.	Two trunks located on both sides of aorta.	Serpentes, Sauria: Sinus surrounds aorta. Crocodylia: Two slender trunks. Chelonia: Two trunks in tail, single in body cavity and bifurcated anteriorly.
5 Cisterna chyli in lumbar area	Always present. Great variation in shape and size.		Serpentes, Crocodylia: Absent. Sauria, Chelonia: Present.
6 Connections with veins	In most mammals thoracic duct connects with left anterior cardinal vein, in some with right, in few with both.	With anterior cardinal veins.	With anterior cardinal veins.
7 Jugular lymphatic trunk	Irregular lymph vessels join anterior cardinal veins separately or by way of subvertebral trunks.	Each of two trunks joins corresponding subvertebral trunk.	Sauria: Two trunks connected with subvertebral trunks by way of jugular sinus. Crocodylia: Two trunks enter corresponding anterior cardinal veins. Chelonia: Two trunks enter jugular cistern which connects both subvertebral trunks.
8 Lateral longitudinal lymphatic trunks	In adult, present only in tail. In embryos, transitory thoracic part observed.		Serpentes: Lateral trunk reaches maxillary sinus. Sauria: From tip of tail to forelimb. Caudal part enters lymph heart, thoracic part enters axillary sinus. Crocodylia: Only caudal part present. Chelonia: Thoracic part well-developed.
9 Longitudinal lymphatic trunks (other than lateral)			

*Contributor:* Grodziński, Z.

*References:* [1] Allen, W. F. 1906. Proc. Wash. Acad. Sci. 8:41. [2] Allen, W. F. 1908. Am. J. Anat. 8:49. Rindes. A. Hirschwald, Berlin. [5] Baum, H. 1928. Das Lymphgefäßsystem des Pferdes. J. Springer, Berlin. Trans. Roy. Soc. Edinburgh 54:309. [8] Favaro, G. 1905. Atti Reale Ist. Veneto Sci. Lettere Arti 65:195. [9] Glaser, p. 433. [11] Grodziński, Z. 1932. Ibid., p. 221. [12] Hoyer, H. 1905. Anat. Anz. 27:50. [13] Hoyer, H. 1908. Polon. Sci. Classe Sci. Math. Nat., B, 1:205. [16] Kampmeier, O. 1925-26. J. Morphol. 41:95. [17] Kihara, R., 1911. Anat. Anz. 40:469. [20] Panizza, B. 1833. Sopra il sistema linfatico dei rettili. Bizzoni, Pavia. [21] Retzius, Morphol. Jahrb. 58:209. [24] Tretjakoff, D. 1930. Ibid. 64:133. [25] Weidenreich, F., H. Baum, and A. Trautmann. v. 6, p. 745.



## CIRCULATORY SYSTEM: VERTEBRATES

### LYMPHATICS

Amphibia (E)	Pisces and Chondrichthyes (F)	Agnatha (G)	
Gymnophiona: One sinus-like extended trunk accompanies aorta. Salientia: Two trunks. Caudata: One or two trunks.	Two slender trunks located on both sides of aorta and caudal artery.	Petromyzones: Unpaired sinus-like trunk beneath aorta and cardinal veins. Myxini: Two wide trunks located on both sides of aorta; fused into wide sinus in liver region.	4
Gymnophiona, Salientia: Absent. Caudata: Present. Extends to thoracic region.			5
Salientia: With anterior lymph hearts. Caudata: With anterior cardinal veins.	Pisces: With anterior cardinal veins. Chondrichthyes: With posterior cardinal vein at point where subclavian artery crosses.	Petromyzones: Numerous connections with both posterior cardinal veins. Myxini: With anterior cardinal vein.	6
Salientia: Two short trunks connecting head sinuses with anterior lymph hearts. Caudata: Two trunks connected with corresponding subvertebral trunks.	Pisces: Two sinus-like distended vessels connected with corresponding subvertebral trunks. Chondrichthyes: Two trunks connected with corresponding anterior cardinal veins.	Petromyzones: Seven peribranchial sinuses, each connected with anterior cardinal vein. Myxini: Absent.	7
Gymnophiona, Caudata: In lateral line groove from base of tail to head. Opens into lymph hearts. Salientia: In tadpoles: from base of tail to anterior lymph heart, disappearing during metamorphosis.	Pisces: Below skin in lateral line groove from base of tail fin to head; opens into some of head sinuses or directly into duct of Cuvier. Chondrichthyes: Absent.		8
Salientia: Dorsal and ventral trunks only in fin of tadpoles. Caudata: Dorsal trunk unpaired, located in dorsal midline of tail and body. Ventral trunk unpaired, located in ventral midline of tail. Abdominal trunk paired in wall of abdomen.	Pisces: Dorsal trunk unpaired, located in dorsal midline of tail and body, from base of tail fin to head; ventral trunk unpaired, located in ventral midline of tail and in middle of abdominal wall. Spinal trunk dorsal to spinal cord. Chondrichthyes: Absent.		9

[3] Allen, W. F. 1913-14. Quart. J. Microscop. Sci. 59(2):309. [4] Baum, H. 1912. Das Lymphgefäßsystem des [6] Clark, E. R., and E. L. Clark. 1920. Contrib. Embryol. Carnegie Inst. Wash. 9:447. [7] Cole, F. J. 1925. G. 1933. Z. Anat. Entwicklungsgeschichte 100:433. [10] Grodziński, Z. 1929. Bull. Intern. Acad. Sci. Cracovie, Bull. Intern. Acad. Sci. Cracovie, p. 451. [14] Hoyer, H. 1928. Ibid., p. 79. [15] Hoyer, H. 1934. Mem. Acad. and E. Naito. 1933. Folia Anat. Japon. 11:405. [18] Marcus, H. 1908. Morphol. Jahrb. 38:590. [19] Mozejko, B. G. 1890. Biol. Untersuch. (Stockholm) 1:20. [22] Sabin, F. R. 1909. Am. J. Anat. 9:43. [23] Tretjakoff, D. 1927. 1933. In L. Bolk, ed. Handbuch der vergleichenden Anatomie der Wirbeltiere. Urban and Schwarzenberg, Berlin.

## 41. COMPARATIVE ANATOMY OF THE

Gland or Tissue	Mammalia	Aves	Reptilia	Amphibia
(A)	(B)	(C)	(D)	(E)
1 Hypophysis Adenohy- pophysis Pars dis- talis	Situating anteriorly and ventrally ("anterior lobe" of older terminology) [18,20].	Large. Situated ventrally or anteroventrally to neurohypophysis. Histologically distinguishable into cephalic and caudal regions. [40]	Well-developed. Situated ventrally and often posteriorly. [40]	Situating at posterior end of gland [32].
2 Pars inter- media	Between pars distalis and neural lobe. Often separated from pars distalis by remains of hypophyseal cleft. Occasionally absent. [18,20]	Absent.	Variable in development, usually with hypophyseal cleft separating it from pars distalis [40].	Situating anterodorsally to pars distalis [32].
3 Pars tu- beralis	Surrounds infundibular stem, forming bed for primary plexus of hypophyseal portal system which vascularizes pars distalis [18,20].	Consists of (i) pars tuberalis proper, a layer of cells on the ventral surface of the diencephalon and within the pia mater; (ii) a portal zone of cell cords and blood vessels, connecting pars tuberalis proper with (iii) pars tuberalis interna which is fused with the pars distalis. [40]	Reduced or absent [40]. Serpentes: Absent [40]. Sauria: Distinguishable only as a few cells in floor of brain [40].	Salientia: Usually reduced (perhaps sometimes absent). Sauria: Distinguishable. Forms two small plates on ventral surface of tuber cinereum. [32] Caudata: Represented by lateral lobes connected with pars distalis [32].
4 Neurohy- pophysis	Divisible into two regions, pars nervosa and median eminence, each with separate vascularization. Pars nervosa usually globular; formed from distal extremity of infundibulum; may or may not contain an extension of the third ventricle. Median eminence is an ill-defined region of infundibular stalk on floor of diencephalon, usually anterior to pars nervosa and always connected to pars distalis by portal blood vessels. Both parts of neurohypophysis contain endings of neurosecretory neurones from the hypothalamus. [18,40]	Distinct neural lobe, or pars nervosa, lying posteriorly and carried by an infundibular stem. A median eminence develops from floor of third ventricle, or infundibular stem anterior to pars nervosa, from which it is not sharply delimited; covered by capillary net of hypophyseal portal system. [40]	Well-defined neural lobe [40]. Serpentes: Compact organ [40]. <i>Sphenodon</i> , some Sauria, Chelonia: Thin-walled sac [40]. A median eminence is differentiated out of infundibular floor. Ranges from a simple form, differing little from neural lobe, to a thickened structure with capillaries of the hypophyseal portal system buried in it. [18,40]	A true neural lobe, or pars nervosa, with independent blood supply, is present as a posterodorsal thickening of the infundibular process. Relatively large in terrestrial species, smaller in more aquatic ones. [18] <i>Rana</i> : A median eminence with portal vessels [18].

## ENDOCRINE SYSTEM: VERTEBRATES

Pisces		Chondrichthyes	Agnatha		
Crossopterygii	Neopterygii and Palaeopterygii		Myxini	Petromyzones	
(F)	(G)	(H)	(I)	(J)	
<p><i>Protopterus</i>: Situated ventrally. Separated from pars intermedia by large hypophyseal cleft. [41]</p> <p><i>Latimeria</i>: has form of elongated cord [30].</p>	<p>Probably represented by rostral zone and proximal zone of pars distalis. Rostral zone is follicular in lower forms such as <i>Acipenser</i>, <i>Amia</i>, and <i>Lepisosteus</i>. [32,33]</p> <p><i>Polypterus</i>: Hypophyseal duct remains open in adult [32,33].</p>	<p>Probably represented by tonguelike rostral lobe; also by a ventral lobe, variable in shape and size, which is peculiar to the group. [10,33]</p>	<p>Rostral zone (irregular cell cords), histologically differentiated from proximal zone (broad cell cords). Both separated from brain by layer of connective tissue. [31,38]</p>	<p>Not divided into regions. Separated from brain by layer of connective tissue. May be penetrated by small diverticula from nasopharyngeal duct. [31,38]</p>	1
<p><i>Protopterus</i>: Dorsal to hypophyseal cleft; tube-shaped lobules with wide cavities. Closely attached to infundibular process lying rostrally to pars intermedia. [41]</p>	<p>Probably represented by "pars intermedia," tissue most intimately associated with neurohypophysis [18,32,33].</p>	<p>Interdigitates with infundibular process to form neurointermediate lobe [10,33].</p>	<p>Probably represented by most posterior part of adeno-hypophysis. Long and thin, and in close contact with neurohypophysis along entire length. [31,38]</p>	<p>Not visibly differentiated.</p>	2
<p><i>Protopterus</i>: Possibly represented by rostral part of pars distalis [41].</p>	<p>Probably not represented.</p>	<p>Not distinguishable.</p>	<p>Not distinguishable.</p>	<p>Not distinguishable.</p>	3
<p><i>Protopterus</i>: Gives rise to distinctly paired infundibular process, as branches of the hypothalamus that intermingle with lobules of the pars intermedia. Median eminence is believed to be represented by an area in the floor of the third ventricle, just rostral to the anterior tip of the adeno-hypophysis; surface is indented by capillaries that communicate with sinusoids of adjacent adeno-hypophysis. [41]</p>	<p>Extensions of the floor of the hypothalamus interdigitate closely with the pars intermedia. Contains ends of neurosecretory fibers and accumulated neurosecretion originating mainly in the nucleus preopticus. Also interdigitates with adeno-hypophysis to a lesser extent. [24,32,33]</p>	<p>Infundibular process interdigitates with pars intermedia. Diffuse structure formed by endings of neurosecretory fibers originating in nucleus preopticus. [10,33] Median eminence on ventral surface of hypothalamus, with portal vessels vascularizing pars distalis [29].</p>	<p>Formed by thickened floor of diencephalon, particularly posterior part [31,38].</p>	<p>Caudally-directed, saclike projection of the hypothalamus represents infundibular process; thin-walled and unpaired, with many terminations of neurosecretory fibers. No contact with adeno-hypophysis. Well-vascularized, with a folded surface. Contains neurosecretion. [31,38]</p>	4

*continued*

## 41. COMPARATIVE ANATOMY OF THE

Gland or Tissue	Mammalia	Aves	Reptilia	Amphibia
(A)	(B)	(C)	(D)	(E)
5 Thyroid	Two lateral lobes and a median isthmus; lateral lobes at sides of trachea, cranial ends at level of caudal edge of cricoid cartilage.	One pair of glands, situated at boundary of neck and thoracic cavity [7].	Single lobulate organ, lying close to trachea [9]. <i>Chrysemys</i> : Flattened mass ventral to truncus arteriosus [9].	Paired, widely separated lobes, closely associated with wings of hyoid cartilage (in almost all Amphibia).
6 Parathyroid	Usually two pairs, situated along dorsal-lateral border of the thyroid as anterior and posterior pairs. Accessory glands common along carotid arteries and in mediastinum. [19] Man: Two to six may be present. Anterior pair may be imbedded in thyroid. [19] Cat, dog: Anterior pair may be imbedded in thyroid [19]. Rat: Only posterior pair present. Visible on surface of thyroid. [19]	One or two pairs, developing from third and fourth branchial pouches. Situated posterior to thyroid glands. [7]	Usually two pairs, situated in neck; posterior and lateral to thyroid.	<i>Rana</i> : Two pairs, against jugular veins [13]. <i>Salamandra</i> : One pair of spherical bodies, derived from ventral ends of third and fourth branchial clefts. Widely separated from thyroids; lateral to arterial arches. Absent in some perennibranchiates ( <i>Ambystoma</i> , <i>Necturus</i> ). [13] Sauria: Situated at level of vagus ganglion [39].
7 Ultimobranchial bodies	Fused with thyroid tissue. Sometimes found aberrant in lateral neck tissues. [17]	Closely associated with, and may surround, parathyroid glands. Cell strands with some colloid-filled vesicles. [39]	Paired; groups of vesicles and strands, varying in position [39]. Serpentes: Situated against thyroid [39]. Reportedly absent in some snakes [17]. Sauria: Situated at level of vagus ganglion [39].	Gymnophiona, Saliencia: Bilateral, at level of truncus arteriosus; vesicular. [13] <i>Salamandra</i> : Small single body near heart [13].
8 Adrenals	One pair, ovoid-shaped [23]. Eutheria: Situated immediately anterior to kidneys, with which adrenals may be in contact [23]. Eutheria, Metatheria: Adrenocortical tissue forms cortex, chromaffin tissue forms medulla [23]. Prototheria: Chromaffin tissue interdigitates with adrenocortical tissue [23].	One pair (occasionally fused), situated at anterior end of kidneys wholly or partly covered by gonads. Adrenocortical tissue forms anastomosing cords, with chromaffin tissue irregularly distributed in its meshes. [21]	One pair, elongated in shape. Adrenocortical tissue forms anastomosing cords; chromaffin tissue intermingled and also forming a peripheral layer. [23,40] Serpentes, Sauria: Anterior to kidneys [23, 40]. Crocodylia: Situated between, and anterior to, kidneys [23,40]. Chelonia: On ventral surface of kidneys. May fuse. [23,40]	<i>Salamandra</i> : Series of orange patches along ventromesial border of kidneys. May extend forward, in association with sympathetic ganglia, to subclavian artery. Brighter and more conspicuous in breeding season. [13,21] <i>Siren</i> : Situated between, and in front of, kidneys [13,21]. Chromaffin cells interspersed with cords of adrenocortical cells [13,21].

/1/ Corpuscles of Stannius, one or more of which lie in or on posterior kidney, were at one time regarded as



## ENDOCRINE SYSTEM: VERTEBRATES

Pisces		Chondrichthyes	Agnatha		
Crossopterygii	Neopterygii and Palaeopterygii		Myxini	Petromyzones	
(F)	(G)	(H)	(I)	(J)	
	Diffuse follicles around ventral aorta [28]. <i>Scarus</i> : Follicles compact and encapsulated [28].	Compact organ lying at, or ventral to, anterior end of ventral aorta, between coracomandibular and coracohyoid muscles.	Elongated follicles extend from second to fifth branchial pouch, dorsal to medioventral cartilage of branchial basket [12].	Separate, large follicles scattered in loose connective tissue along most of ventral surface of pharynx [37].	5
	Reportedly absent.	Reportedly absent.	Reportedly absent.	Reportedly absent.	6
	Vesicular, glandlike bodies, often on left side only, or fused to median body, above pericardial wall. Parathyroid-like function has been suggested. [11,17,39]	Vesicles with colloid contents on left side only. Dorsal to pericardium (suprapericardial body). [8,39]	Reportedly absent [39].	Reportedly absent [39].	7
Exact limits of adrenocortical tissue doubtful. Probably present as small groups of cells between ventral margins of kidneys, and in walls of capillaries supplying posterior cardinal veins. [15,23] Chromaffin tissue present as groups of cells distributed along posterior cardinal veins. Also associated with intercostal arteries. [23]	Adrenocortical tissue present as areas of cells in the (usually lymphoid) pronephros. May be scattered ( <i>Salmo</i> ), or arranged as layers of cells closely associated with cardinal veins. [1,23] Chromaffin tissue mainly in pronephros, as scattered cells or islets near or in walls of cardinal veins and their branches. [1,23]	<i>Dogfish</i> : Adrenocortical tissue present as elongated body between kidneys and close to cardinal vein, with additional small groups of cells anteriorly [21,23]. <i>Raja</i> : Adrenocortical tissue a horseshoe-shaped body, usually asymmetrical, with the two limbs extending anteriorly against inner sides of kidneys [21,23]. <i>Torpedo</i> : Adrenocortical tissue an oval body lying posteriorly on left kidney [21,23]. Chromaffin tissue present as paired series of suprarenal bodies along length of body cavity. Situated above cardinal veins and kidneys, or imbedded in kidneys. Variable shape, tending to be distributed in relation to segmental arteries. Some large bodies lie anteriorly. [21]	Adrenocortical tissue present as scattered islets of cells against ventral and lateral walls of cardinal veins in trunk and tail; also as groups of cells in pronephros and around kidney vessels. [16] Chromaffin tissue probably represented by groups of cells distributed along main arteries and veins in trunk [14].	8	

adrenocortical tissue, but this now seems doubtful [17].

continued

41. COMPARATIVE ANATOMY OF THE

Gland or Tissue	Mammalia	Aves	Reptilia	Amphibia
(A)	(B)	(C)	(D)	(E)
9 Pancreatic tissue	Islets of Langerhans; small masses of tissue in pancreas. Principal cell types are $\alpha$ and $\beta$ .	Islets of Langerhans in pancreas. More $\alpha$ than $\beta$ cells. [17]	Islets of Langerhans in pancreas. More $\alpha$ than $\beta$ cells. [17] Serpentes: Large aggregations of islets macroscopically visible at splenic end [5].	Islet tissue distributed throughout organ, ranging from single cells in exocrine acini to groups of cells [4, 17]. Caudata: $\beta$ cells predominate [17].
10 Gonadal tissue	In testis, consists of interstitial (Leydig) cells between seminiferous tubules [32]. In ovary, probably represented by theca interna of graafian follicle and/or by interstitial cells of stroma. Progesterone-secreting cells of corpus luteum derived from granulosa cells of follicle, with perhaps a contribution from theca interna. [32]	Interstitial cells in testis [7].	Interstitial cells between seminiferous tubules of testis [27].	Interstitial cells in testis, conspicuous at breeding season.
11 Urohypophysis (Urophysis)				

*Contributors:* (a) Barrington, E. J. W., (b) Gorbman, Aubrey.

*References:* [1] Baecker, R. 1928. Z. Mikroskop. Anat. Forsch. 15:204. [2] Barrington, E. J. W. 1942. J. Exptl. Biol. 92:205. [5] Barrington, E. J. W. 1953. Ibid. 94:281. [6] Barrington, E. J. W. 1963. Introduction to general morphology. W. E. 1917. J. Morphol. 28:369. [9] Charipper, H. A. 1929. Anat. Record 44:117. [10] Dodd, J. M., P. J. [12] Fontaine, M., et al. 1952. Ann. Endocrinol. (Paris) 13:55. [13] Francis, E. T. B. 1934. The anatomy of the pituitary gland. J. Anat. 88:225. [16] Giacomini, E. 1902. Monit. Zool. Ital. 13:143. [17] Gorbman, A., and J. Anat. 88:225. [19] Greep, R. O. 1948. In G. Pincus and K. V. Thimann, ed. The hormones. Academic Press. [21] Hartman, F. A., and K. A. Brownell. 1949. The adrenal gland. Lea and Febiger, Philadelphia. [22] Holmgren, [24] Kerr, T. 1949. Proc. Zool. Soc. London 118:973. [25] Marshall, A. J., and B. Lofts. 1956. Nature 177:704. 1957. Ibid. 98:89. [28] Matthews, S. A. 1948. Anat. Record 101:251. [29] Meurling, P. 1960. Nature 187:336. Anat. 51:97. [32] Parkes, A. S., ed. 1952-65. Marshall's Physiology of reproduction. Ed. 3. Longmans, Green; pituitary gland of fishes. New York Zoological Society, New York. [34] Robertson, O. H. 1958. U. S. Fish Wildlife [37] Tong, W., et al. 1961. Biochim. Biophys. Acta 52:299. [38] Van de Kamer, J. C., and A. F. Schreurs. [40] Wingstrand, K. G. 1951. The structure and development of the avian pituitary. C. W. K. Gleerup; Lund,

**ENDOCRINE SYSTEM: VERTEBRATES**

Pisces		Chondrichthyes	Agnatha		
Crossopterygii	Neopterygii and Palaeopterygii		Myxini	Petromyzones	
(F)	(G)	(H)	(I)	(J)	
	Organ usually diffuse. Islet tissue may be scattered but is often one large islet situated near spleen ( <i>Lophius</i> ).	Islet tissue not well-identified. <i>Carcharhinus, Mustelus, Dasyatis</i> : Islet cells form a second epithelial layer around duct cells of pancreas. [36] <i>Raja</i> : Solid cords of islet cells contiguous to ducts of pancreas [36]. <i>Squalus acanthias</i> : Small islands of solid cell cords which are frequently separated from duct system of pancreas [36].	Islet tissue probably represented in ammocoete larva by groups of cells in wall of intestine adjacent to bile duct, and in adult by groups of cell cords in anterior wall of intestine [2,6].	Islet tissue possibly represented by cluster of cell cords in wall of intestine around bile duct [3,6].	9
	<i>Gasterosteus, Oncorhynchus</i> : Typical interstitial cells between tubules of testis [25,26]. <i>Esox</i> : Lobule boundary cells at periphery of testis lobules [25,26]. <i>Salmo gairdneri</i> : Both above types of endocrine tissue present [34].	<i>Chimaera, Scyliorhinus</i> : Probably interstitial cells between tubules of testis [25].		Interstitial cells reported in testis.	10
	Neurosecretory cells at hind end of spinal cord [17,22,35].	Neurosecretory cells (Dahlgren cells) at hind end of spinal cord [17].			11

Biol. 19:45. [3] Barrington, E. J. W. 1945. Quart. J. Microscop. Sci. 85:391. [4] Barrington, E. J. W. 1951. and comparative endocrinology. Clarendon Press, Oxford. [7] Benoit, J. 1950. Traite Zool. 15:290. [8] Camp, Evannett, and C. K. Goddard. 1960. Symp. London Zool. Soc. 1:77. [11] Eggert, B. 1938. Endokrinologie 20:1. salamander. Clarendon Press, Oxford. [14] Gaskell, J. F. 1912. J. Physiol. (London) 44:59. [15] Gérard, P. H. A. Bern. 1962. A textbook of comparative endocrinology. J. Wiley, New York. [18] Green, J. D. 1951. Am. New York. v. 1, p. 255. [20] Harris, G. W. 1955. Neural control of the pituitary gland. E. Arnold, London. U. 1959. Anat. Record. 135:51. [23] Jones, I. C. 1957. The adrenal cortex. Cambridge Univ. Press, London. [26] Marshall, A. J., and B. Lofts. 1957. Quart. J. Microscop. Sci. 98:79. [27] Marshall, A. J., and F. M. Woolf. [30] Millot, J., and J. Anthony. 1955. Compt. Rend. 241:114. [31] Olsson, R. 1959. Z. Zellforsch. Mikroskop. London, v. 1, 2; Little and Brown, Boston, v. 3. [33] Pickford, G. E., and J. W. Atz. 1957. The physiology of the Serv. Fishery Bull. 58:9. [35] Sano, Y. 1961. Ergeb. Biol. 24:191. [36] Thomas, T. B. 1940. Anat. Record 76:1. 1959. Z. Zellforsch. Mikroskop. Anat. 49:605. [39] Watzka, M. 1933. Z. Mikroskop. Anat. Forsch. 34:485. Sweden. [41] Wingstrand, K. G. 1956. Dansk. Naturhist. Foren. Videnskab. Medd. 118:193.

## 42. COMPARATIVE ANATOMY OF

Names of bones are those used in mammalian anatomy. Alternate names used in human anatomical terminology  
 pn = prenasal; var = variable; inn = innominate; px = proximally; d = distally.

### Part I. AXIAL

Bone	Primates					Artiodactyla		Perisso-	Sirenia	Probos-
	<i>Homo sapiens</i> (man)	<i>Pan troglodytes</i> (chimpanzee)	<i>Macaca mulatta</i> (rhesus monkey)	<i>Alouatta balzabul</i> (howler monkey)	<i>Lemur macaco</i> (lemur)	<i>Bos taurus</i> (cattle)	<i>Sus scrofa</i> (swine)	<i>Equus caballus</i> (horse)	<i>Trichechus</i> spp. (manatee)	<i>Elephas maximus</i> (Asiatic elephant)
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
<b>Skull</b>										
1 Occipital	1	1	1	1	1	1	1	1	1	1
2 Parietal	1 pr	1 pr	1 pr, f	1 pr, f	1 pr	1 pr, f	1 pr, f	1 pr	1 pr	1 pr
3 Ethmoid	1	1	1	1	1	1	1	1	1	1
4 Turbinal	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	1 pr	1+ pr, ru
5 Interparietal	0	0	0	0	0	0	0	0	0	0
6 Frontal	1 pr, f	1 pr, f	1 pr, f	1 pr, f	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
7 Nasal	1 pr	1 pr, f	1 pr, f	1 pr	1 pr	1 pr	1 pr	1 pr	0 or ru	1 pr
8 Lacrimal	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	ru	1 pr
9 Temporal										
10 Periotic (petrosal)	} f	} f	} f	} f	} f	} f	1 pr	} f	} 1 pr	} 1 pr
11 Tympanic							1 pr			
12 Squamosal							1 pr			
13 Premaxilla	} f	1 pr, f	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
14 Maxilla		1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
15 Zygomatic	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
16 Vomer	1	1	1	1	1	1	1	1	1	1
17 Palatine	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	f <sup>a</sup>	1 pr
18 Mandible	1 pr, f	1 pr, f	1 pr, f	1 pr, f	1 pr, f	1 pr	1 pr, f	1 pr, f	1 pr	1 pr, f
19 Hyoid										
20 Basi-	1	1	1	1	1	1	1	1	1	1
21 Stylo-	0	0	0	0	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
22 Epi-	0	0	0	0	1 pr	1 pr	0	1 pr	0	0
23 Cerato-	0	0	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	0	0
24 Thyro-	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
<b>Vertebrae</b>										
25 Cervical	7	7	7	7	7	7	7	7	6	7
26 Thoracic	12	13	12-13	12-13	12	13-14	14-15	18-19	17-18	19-20
27 Lumbar	5	4	7	6	6	6	5-7	6	2	3-4
28 Sacral	5	4-5	3	3	3	5	4	5	1 ru	4
29 Caudal	4	4-5	20+	27+	25-29	16-21	20-26	15-21	25+	31+
<b>Vertebral Ribs</b>										
30 "True" ribs	7 pr	7 pr	8 pr	5 pr	7 pr	7-8 pr	7 pr	8 pr	0 or 1 pr	} 19-20 pr
31 "False" ribs	5 pr	6 pr	4 pr	7-8 pr	5 pr	5-7 pr	7-8 pr	10-11 pr	16-18 pr	
<b>Sternum</b>										
32 Manubrium	1	1	1	1	1	1	1	1	} 1	1
33 Sternebrae	1	1	5	4	4	5	4	5		
Xiphisternum	1	1	1	1	1	1	1	1		1
34 Sternal ribs										

/1/ The extreme diversification in the skeletal system of the Edentata has not been included. /a/ Fused with

Contributors: Moyer, Elizabeth K., and Kaliszewski, Barbara Freeman



## THE SKELETAL SYSTEM: MAMMALS

(*Nomina Anatomica*, 1955) appear in parentheses in column A. *Abbreviations*: pr = pair; f = fused; ru = rudimentary;

### SKELETON

Carnivora			Cetacea		Rodentia		Lago- morpha	Eden- tata <sup>1</sup>	Chiro- ptera	Insec- tivora	Marsu- pialia	Monotremata		
<i>Canis familiaris</i> (dog)	<i>Felis catus</i> (cat)	<i>Phoca vitulina</i> (harbor seal)	<i>Balaenoptera physalus</i> (finback whale)	<i>Phocaena phocaena</i> (harbor porpoise)	<i>Cavia tschudi pallidior</i> (guinea pig)	<i>Rattus norvegicus</i> (Norway rat)	<i>Lepus americanus virginianus</i> (varying hare)	<i>Dasyurus novemcinctus</i> (nine-banded armadillo)	<i>Eptesicus fuscus</i> (big brown bat)	<i>Sorex cinereus</i> (gray shrew)	<i>Didelphis marsupialis virginiana</i> (Virginia opossum)	<i>Ornithorhynchus</i> spp. (platypus)	<i>Tachylossus</i> spp. (spiny anteater)	
(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	(W)	(X)	(Y)	
Skull														
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr, f	1 pr, f	1 pr	1 pr, f	1 pr, f	2
1	1	1	1	1	1	1	1	1	1	1	1	1	1	3
2 pr	2 pr	2 pr	ru	0	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	2 pr	4
1	1	1	0	1	0	1	0	1	0	0	0	0	0	5
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr, f	1 pr, f	1 pr, f	6
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr, pn	1 pr, f	7
1 pr	1 pr	1 pr	1 pr	0	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	8
} f	} f	} f	} 1 pr	} 1 pr	} 1 pr	} 1 pr	} 1 pr	} f	1 pr	1 pr	1 pr	1 pr	1 pr	9
			1 pr	1 pr	1 pr	1 pr	1 pr		1 pr	1 pr	1 pr	1 pr	1 pr	10
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	0	1 pr	1 pr	1 pr	1 pr	11
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	12
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	0	1 pr	1 pr	1 pr	13
1	1	1	1	1	1	1	1	1	1	1	1	1	1	14
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr, f	1 pr	15
1 pr	1 pr	1 pr	1 pr	1 pr, f	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	16
1	1	1	1	1	1	1	1	1	1	1	1	1	1	17
1	1	1	1	1	1	1	1	1	1	1	1	1	1	18
1 pr	1 pr	1 pr	1 pr	1 pr	0	0	0	1 pr	1 pr	1 pr	0	0	0	19
1 pr	1 pr	1 pr	0	0	0	0	0	1 pr	1 pr	1 pr	0	1 pr	1 pr	20
1 pr	1 pr	1 pr	0	0	0	0	0	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	21
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	22
Vertebrae														
7	7	7	7 f	7 f	7	7	7	7, 2+3 f	7	7	7	7	7	23
13	13	14-15	14	13	13	13	12	10	11-12	13	13	17	16	24
7	7	5	15	17	5-6	6-7	7	4-5	5-6	6	6	2	3	25
3	2-3	4-5	0	0	2	3-4	2-3	7	3		2	2	3-4	26
var	18±	9-14	24±	25±	8±	27±	var	27±	9±		19-35	20-23	12-13	27
Vertebral Ribs														
9 pr	9 pr	10 pr	1 pr	4-6 pr <sup>3</sup>	6 pr	7 pr	7 pr	5 pr	7 pr	} 13 pr	7 pr	6 pr	6 pr	28
4 pr	4 pr	5 pr	13 pr	7-9 pr	7 pr	6 pr	5 pr	5 pr	4 pr		6 pr	11 pr	10 pr	29
Sternum														
1	1	1	} 1	} 1	1	1	1	1	1	1	1	1	1	30
8	6	6			3-4	3-5	3-4	4	} f	5	4	3	3f	31
1	1	1			1	1	1	1		1	1	0		32
								4 pr	7 pr		7 pr	6 pr	6 pr	33

sphenoid. /s/ Ossified intermedial ribs are present at the ventral ends of the first six pair of vertebral ribs.

continued

## 42. COMPARATIVE ANATOMY OF

### Part I. AXIAL

References: [1] Adams, L. A., and S. Eddy. 1949. Comparative Anatomy. J. Wiley, New York. pp. 132-241. [3] International Anatomical Nomenclature Committee. 1955. Nomina anatomica. Spottiswoode, Ballantyne; London. 1962. The vertebrate body. Ed. 3. W. B. Saunders, Philadelphia. [6] Simpson, G. G. 1945. Bull. Am. Museum Saunders, Philadelphia. pp. 20-253.

### Part II. APPENDICULAR

Bone	Primates					Artiodactyla		Perisso-	Sirenia	Probos-
	<i>Homo sapiens</i> (man)	<i>Pan troglodytes</i> (chimpanzee)	<i>Macaca mulatta</i> (rhesus monkey)	<i>Alouatta balzabul</i> (howler monkey)	<i>Lemur macaco</i> (lemur)	<i>Bos taurus</i> (cattle)	<i>Sus scrofa</i> (swine)	<i>Equus caballus</i> (horse)	<i>Trichechus</i> spp. (manatee)	<i>Elephas maximus</i> (Asiatic elephant)
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
<b>Pectoral Girdle</b>										
1	Interclavicle	0	0	0	0	0	0	0	0	0
2	Coracoid	0	0	0	0	0	0	0	0	0
3	Anterior	0	0	0	0	0	0	0	0	0
4	Clavicle	1 pr	1 pr	1 pr	1 pr	1 pr	0	0	0	0
5	Scapula	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr
<b>Upper Extremity<sup>a</sup></b>										
6	Humerus	1	1	1	1	1	1	1	1	1
7	Radius	1	1	1	1	1	1	1	} f, px	1
8	Ulna	1	1	1	1	1	ru	1		ru
9	Carpus									
10	Scaphoid	1	1	1	1	1	1	1	1	1
11	Lunate	1	1	1	1	1	1	1	1	1
12	Cuneiform (triquetral)	1	1	1	1	1	1	1	1	1
13	Pisiform	1	1	1	1	1	1	1	0	1
14	Centrale	0	0	1	1	0	0	0	0	0
15	Trapezium	1	1	1	1	1	0	0	0 or 1	1
16	Trapezoid	1	1	1	1	1	} 1	1	1	1
17	Magnum (capitate)	1	1	1	1	1		1	1	1
18	Unciform (hamate)	1	1	1	1	1	1	1	1	1
19	Metacarpus	5	5	5	5	5			5	5
20	1st						0	0		
21	2nd						ru	1	ru	
22	3rd						} 1	1	1	
23	4th							1	ru	
24	5th						ru	1	0	
<b>Phalanges</b>										
25	1st digit	2	2	2	2	2	0	0	0	2
26	2nd digit	3	3	3	3	3	0	3	0	3
27	3rd digit	3	3	3	3	3	3	3	3	3
28	4th digit	3	3	3	3	3	3	3	0	3
29	5th digit	3	3	3	3	3	0	3	0	3

/1/ The extreme diversification in the skeletal system of the Edentata has not been included. /2/ Single extremity. maining four, varying numbers of phalanges become ossified.

## THE SKELETAL SYSTEM: MAMMALS

### SKELETON

- [2] Flower, W. H. 1885. An introduction to the osteology of the Mammalia. Ed. 3. Macmillan, London.  
 [4] Parker, T. J., and W. A. Haswell. 1940. A textbook of zoology. Ed. 6. Macmillan, London. [5] Romer, A. S. Nat. Hist. 85:1. [7] Sisson, S., and J. D. Grossman. 1953. The anatomy of the domestic animals. Ed. 4. W. B.

### SKELETON

Carnivora			Cetacea		Rodentia		Lago- morpha	Eden- tata <sup>1</sup>	Chiro- ptera	Insec- tivora	Marsu- pialia	Monotremata		
<i>Canis familiaris</i> (dog)	<i>Felis catus</i> (cat)	<i>Phoca vitulina</i> (harbor seal)	<i>Balaenoptera physalus</i> (finback whale)	<i>Phocaena phocaena</i> (harbor porpoise)	<i>Cavia tschudi palidior</i> (guinea pig)	<i>Rattus norvegicus</i> (Norway rat)	<i>Lepus americanus virginianus</i> (varying hare)	<i>Dasyurus novemcinctus</i> (nine-banded armadillo)	<i>Eptesicus fuscus</i> (big brown bat)	<i>Sorex cinereus</i> (gray shrew)	<i>Didelphis marsupialis virginiana</i> (Virginia opossum)	<i>Ornithorhynchus</i> spp. (platypus)	<i>Tachyglossus</i> spp. (spiny anteater)	
(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	(W)	(X)	(Y)	
Pectoral Girdle														
0	0	0	0	0	0	0	0	0	0	0	0	1	1	1
0	0	0	0	0	0	0	0	0	0	0	0	1 pr	1 pr	2
0	0	0	0	0	0	0	0	0	0	0	0	1 pr	1 pr	3
ru	ru	0	0	0	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	4
1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	1 pr	5
Upper Extremity <sup>2</sup>														
1	1	1	1	1	1	1	1	1	1	1	1	1	1	6
1	1	1	1	1	1	1	1	1	1	1	1	1	1	7
1	1	1	1	1	1	1	1	1	ru or 0	1	1	1	1	8
			5 <sup>3</sup>											9
}1	}1	}1		1	}1	}1	}1	1	}1	1	1	}1	}1	10
				1				1		1	1			11
1	1	1		1	1	1	1	1	1	1	1	1	1	12
1	1	1		0	1	1	1	1	1	1	1	1	1	13
0	0	0		0	0	0	1	0	0	0	0	0	0	14
1	1	1		0	1	1	1	1	1	1	1	1	1	15
1	1	1		1	1	1	1	1	1	1	1	1	1	16
1	1	1		0 or 1	1	1	1	1	1	1	1	1	1	17
1	1	1		1	1	1	1	1	1	1	1	1	1	18
ru	ru	5		5	5		5	5	5	5	5	5	5	19
1	1		0			ru								20
1	1		1			1								21
1	1		1			1								22
1	1		1			1								23
1	1		1			1								24
2	2	2	0	ru <sup>4</sup>	2	2	2	2	2	2	2	2	2	25
3	3	3	3		3	3	3	3	2	3	3	3	3	26
3	3	3	5		3	3	3	3	3	3	3	3	3	27
3	3	3	5		3	3	3	3	2	3	3	3	3	28
3	3	3	3		3	3	3	3	2	3	3	3	3	29

/3/ Homologies difficult. /4/ *P. phocaena* has five digits. The first is rudimentary and cartilaginous. In the re-

continued

## 42. COMPARATIVE ANATOMY OF

### Part II. APPENDICULAR

Bone	Primates					Artiodactyla		Perisso- dactyla	Sirenia	Probos- cida	
	<i>Homo sapiens</i> (man)	<i>Pan troglodytes</i> (chimpanzee)	<i>Macaca mulatta</i> (rhesus monkey)	<i>Alouatta balzabul</i> (howler monkey)	<i>Lemur macaco</i> (lemur)	<i>Bos taurus</i> (cattle)	<i>Sus scrofa</i> (swine)	<i>Equus caballus</i> (horse)	<i>Trichechus</i> spp. (manatee)	<i>Elephas maximus</i> (Asiatic elephant)	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
<b>Pelvic Girdle</b>											
30 Epipubic	0	0	0	0	0	0	0	0	0	0	
31 Ilium	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	ru	} 1 pr, inn	
32 Ischium									0		
33 Pubis									0		
<b>Lower Extremity<sup>a</sup></b>											
34 Femur	1	1	1	1	1	1	1	1	0	1	
35 Patella	1	1	1	1	1	1	1	1	0	1	
36 Tibia	1	1	1	1	1	1	1	1	0	1	
37 Fibula	1	1	1	1	1	ru	1	ru	0	1	
Tarsus											
38 Astragalus (talus)	1	1	1	1	1	1	1	1	0	1	
39 Calcaneus	1	1	1	1	1	1	1	1	0	1	
40 Navicular	1	1	1	1	1	} 1	1	1	0	1	
Cuneiform											
41 Internal	1	1	1	1	1		1	} 1	} 1	0	1
42 Middle	1	1	1	1	1	1	1			0	1
43 External	1	1	1	1	1	1	1	1	0	1	
44 Cuboid	1	1	1	1	1	1	1	1	0	1	
Metatarsus											
45 1st	5	5	5	5	5	0	0	0	0	5	
46 2nd						0	1	ru			
47 3rd						} 1	1	1			
48 4th							1	ru			
49 5th						0	1	0			
Phalanges											
51 1st digit	2	2	2	2	2	0	0	0	0	1	
52 2nd digit	3	3	3	3	3	0	3	0	0	3	
53 3rd digit	3	3	3	3	3	3	3	3	0	3	
54 4th digit	3	3	3	3	3	3	3	0	0	3	
55 5th digit	3	3	3	3	3	0	3	0	0	1	

/1/ The extreme diversification in the skeletal system of the Edentata has not been included. /2/ Single extremity.

**Contributors:** Moyer, Elizabeth K., and Kaliszewski, Barbara Freeman

**References:** [1] Adams, L. A., and S. Eddy. 1949. Comparative Anatomy. J. Wiley, New York. pp. 132-241.  
 [3] International Anatomical Nomenclature Committee. 1955. Nomina anatomica. Spottiswoode, Ballantyne; London.  
 1962. The vertebrate body. Ed. 3. W. B. Saunders, Philadelphia. [6] Simpson, G. G. 1945. Bull. Am. Museum  
 Saunders, Philadelphia. pp. 20-253.



## THE SKELETAL SYSTEM: MAMMALS

### SKELETON

Carnivora			Cetacea		Rodentia		Lago- morpha	Eden- tata <sup>1</sup>	Chirop- tera	Insec- tivora	Marsu- pialia	Monotremata		
<i>Canis familiaris</i> (dog)	<i>Felis catus</i> (cat)	<i>Phoca vitulina</i> (harbor seal)	<i>Balaenoptera physalus</i> (finback whale)	<i>Phocaena phocaena</i> (harbor porpoise)	<i>Cavia tschudi pallidor</i> (guinea pig)	<i>Rattus norvegicus</i> (Norway rat)	<i>Lepus americanus virginianus</i> (varying hare)	<i>Dasybus novemcinctus</i> (nine-banded armadillo)	<i>Eptesicus fuscus</i> (big brown bat)	<i>Sorex cinereus</i> (gray shrew)	<i>Didelphis marsupialis virginiana</i> (Virginia opossum)	<i>Ornithorhynchus</i> spp. (platypus)	<i>Tachyglossus</i> spp. (spiny anteater)	
(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	(W)	(X)	(Y)	
Pelvic Girdle														
0	0	0	0	0	0	0	0	0	0	0	1 pr	1 pr	1 pr	30
} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	0	0	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} f <sup>s</sup> , inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	} 1 pr, inn	31
			ru	ru										32
			0	0										33
Lower Extremity <sup>2</sup>														
1	1	1	0	0	1	1	1	1	1	1	1	1	1	34
1	1	1	0	0	1	1	1	1	1	1	0	1	1	35
} f, d?	} 1	} f, px	0	0	1	1	1	} f, px	1	1	1	1	1	36
			0	0	1	1	1		1	1	1	37		
1	1	1	0	0	1	1	1	1	1	1	1	1	1	38
1	1	1	0	0	1	1	1	1	1	1	1	1	1	39
1	1	1	0	0	1	1	1	1	1	1	1	1	1	40
1	1	1	0	0	1	1	1	1	1	1	1	1	1	41
1	1	1	0	0	1	1	1	1	1	1	1	1	1	42
1	1	1	0	0	1	1	1 or 0	1	1	1	1	1	1	43
1	1	1	0	0	1	1	1	1	1	1	1	1	1	44
ru	ru	5	0	0	5	5	0	5	5	5	5	5	5	45
1	1						1							46
1	1						1							47
1	1						1							48
1	1						1							49
1	1						1							50
ru	ru	2	0	0	2	2	0	2	2	2	2	2	2	51
3	3	3	0	0	3	3	3	3	3	3	3	3	3	52
3	3	3	0	0	3	3	3	3	3	3	3	3	3	53
3	3	3	0	0	3	3	3	3	3	3	3	3	3	54
3	3	3	0	0	3	3	3	3	3	3	3	3	3	55

/E/ Fused with sacrum.

[2] Flower, W. H. 1885. An introduction to the osteology of the Mammalia. Ed. 3. Macmillan, London.

[4] Parker, T. J., and W. A. Haswell. 1940. A textbook of zoology. Ed. 6. Macmillan, London. [5] Romer, A. S.

Nat. Hist. 85:1. [7] Sisson, S., and J. D. Grossman. 1953. The anatomy of the domestic animals. Ed. 4. W. B.

# *Contrails*

## V. NUTRITION AND DIGESTION

### 43. NUTRIENTS: CHEMICAL ELEMENTS

Accumulation in the tissues of an organism is not alone sufficient evidence that an element is required. *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species or strains, but not to all forms studied. R and r = required; H = not required; u = utilized as effectively as, replaces wholly, or is interchangeable with, another element; u< = can partially replace, or spare the use of, another element; s = stimulates growth or other processes; a = accumulates in the tissues; c = commonly present at similar concentrations in the food and tissues, but requirement is uncertain.

Nutrient (Symbol)	Vertebrata	Invertebrata			Phyto-flagellata (green) <sup>2</sup>	Algae	Bacterio-phyta	Fungi		Spermato-phyta
		Insecta	Other	Protozoa <sup>1</sup>				Saccharo-mycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
1 Aluminum (Al)	H	H	.....	R	H	H	H, u< <sup>3</sup>	H	H, s	r, s, a
2 Arsenic (As)	H	H	a	H	H	H	H	H	H	H
3 Boron (B)	H	H	H	H	r?	r	r	H	H, r?	R
4 Bromine (Br)	u<	H	r? <sup>4</sup> , a?	H	H	H, a	H	H	H	H
5 Calcium (Ca)	R	r, c	r	R	r	R	r, s	r, u <sup>6</sup>	r, s, a	R
6 Carbon (C) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
7 Chlorine (Cl)	R	r, c	r	H	H	a	H	H	H	r, s
8 Chromium (Cr)	r?	H	.....	H	H	H	u <sup>6</sup>	H	r, s	H
9 Cobalt (Co)	r	r?, s?	.....	r	r	r	r	u <sup>7</sup>	r, a	r
10 Copper (Cu)	r	r, c	r	...	...	r	r, u< <sup>3</sup>	r	R	R
11 Fluorine (F)	H, s	H	a	H	H	H	H	H	H	H, a
12 Gallium (Ga)	H	H	H	H	H	H	H	H	r, s	r
13 Hydrogen (H) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
14 Iodine (I)	R	H	r?, a	R	H	H, a	H, s	H, s	H	H, s
15 Iron (Fe)	R	r	r	R	R	R	r, u< <sup>3</sup>	R	R	R
16 Magnesium (Mg)	R	r	R	R	R	R	r	R	R	R
17 Manganese (Mn)	R	r, c, a	r <sup>8</sup>	r	r	r	r, u <sup>6</sup>	r, u <sup>7</sup>	R, u<	R
18 Molybdenum (Mo)	r <sup>9</sup>	.....	.....	H	H	r <sup>10</sup>	r <sup>10</sup>	r	r <sup>10</sup> , s	R, a
19 Nitrogen (N) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
20 Oxygen (O) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
21 Phosphorus (P) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
22 Potassium (K)	R	r, c, a	R	R	R	R, a	r	r	r	R
23 Rubidium (Rb)	H	H	.....	H	H	H	u <sup>11</sup>	H	H	H
24 Selenium (Se)	r	H	.....	H	H	H	H	H	H	H, a
25 Silicon (Si)	H	H	r	r	r	r	H	H	H	r
26 Sodium (Na)	R	r?, c	.....	r	H	r, a	r	H	H	r?, s, a
27 Strontium (Sr)	u<	H	.....	H	H	H, u <sup>6</sup>	u <sup>12</sup>	H	s, u<	H
28 Sulfur (S) <sup>5</sup>	R	R	R	R	R	R	R	R	R	R
29 Tungsten (W)	H	H	.....	H	H	H	H	H	s	H
30 Vanadium (V)	H	H	r <sup>13</sup>	H, s?	H	r	u< <sup>14</sup>	H	H, s	H
31 Zinc (Zn)	R	r	r?, a	r	r	r	r, u <sup>6</sup>	r	r, s	R
Reference	1,4,9,11,15, 18,21,25,30, 35,37-39,41, 42,46-48,52, 53,56	11,19, 27,28, 58	3,6,7, 11,14, 20,25, 26	10,11,22, 29,36	10,11,22, 29,36	8,11, 22,33, 45,50,54, 49	11,17,43- 55	5,11,13,44, 45,54,57	5,11, 12,16, 24,34, 44,45, 51,54	2,5,11,23, 26,31,32, 35,40,54

/<sup>1</sup>/ Including the colorless Phytoflagellata. /<sup>2</sup>/ Also Dinoflagellata and Chrysoomonadina. /<sup>3</sup>/ Replaces or spares manganese or chromium in *Aerobacter aerogenes*. /<sup>4</sup>/ Occurs as dibromotyrosine in scleroprotein of certain corals. /<sup>5</sup>/ Universal constituent of protoplasm. /<sup>6</sup>/ Replaces, or is interchangeable with, manganese in *Aerobacter aerogenes*. /<sup>7</sup>/ Replaces, or is interchangeable with, calcium in yeast cocarboxylase. /<sup>8</sup>/ In blood respiratory pigment of *Pinna squamosa* (mollusk). /<sup>9</sup>/ Xanthine oxidase factor. /<sup>10</sup>/ Required for NO<sub>3</sub><sup>-</sup> utilization by some fungi and some algae; required for nitrogen fixation by some bacteria and algae. /<sup>11</sup>/ Replaces, or is interchangeable with, calcium in some bacteria. /<sup>12</sup>/ Replaces, or is interchangeable with, calcium in *Azotobacter*. /<sup>13</sup>/ In blood pigment of certain tunicates (Chordata). /<sup>14</sup>/ Replaces or spares molybdenum in nitrogen fixation.

*Contributors:* (a) Cantino, Edward C., (b) Fogg, G. E., (c) Gordon, Harold Thomas, (d) Hansard, Sam L., (e) Haskins, R. H., (f) House, Howard L., (g) Koser, Stewart A., (h) Kratzer, F. H., (i) Loefer, John B., (j) Pelletier, Réal L., (k) Purvis, E. R., (l) Rusoff, Louis Leon, (m) Schaefer, Arnold Edward, (n) Shorb, Mary S., (o) Turrell, Franklin M., (p) Van Wagtenonk, W. J., (q) Zipkin, Isadore

*continued*

43. NUTRIENTS: CHEMICAL ELEMENTS

- References:* [1] American Medical Association Council on Foods and Nutrition. 1951. Handbook of nutrition; symposium. Ed. 2. Blakiston, New York. [2] Arnon, D. I. 1951. Mineral nutrition of plants. Univ. Wisconsin Press, Madison. [3] Baldwin, E. 1948. An introduction to comparative biochemistry. Ed. 3. Cambridge Univ. Press, London. [4] Bohstedt, G. 1942. *J. Dairy Sci.* 25:441. [5] Bonner, J. 1950. Plant biochemistry. Academic Press, New York. [6] Bourne, G. H., and G. W. Kidder, ed. 1953. Biochemistry and physiology of nutrition. Academic Press, New York. [7] Buchsbaum, R. 1948. Animals without backbones. Rev. ed. Univ. Chicago Press, Chicago. [8] Burlew, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [9] Caldecott, R. S., and L. A. Snyder, ed. 1960. Radioisotopes in the biosphere. Univ. Minnesota Press, Minneapolis. [10] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [11] Chilean Nitrate Educational Bureau, Inc. 1948. Bibliography of literature on the minor elements. New York. [12] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [13] Dunn, G. G. 1952. *Wallerstein Lab. Commun.* 15(48):61. [14] Everett, M. R. 1946. Medical biochemistry. Ed. 2. P. B. Hoeber, New York. [15] Follis, R. H. 1948. The pathology of nutritional disease. C. C. Thomas, Springfield, Ill. [16] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [17] Gale, E. F. 1951. The chemical activities of bacteria. Ed. 3. Academic Press, New York. [18] Gamble, J. L. 1954. Chemical anatomy, physiology and pathology of extracellular fluid. Ed. 6. Harvard Univ. Press, Cambridge. [19] Gilmour, D. 1960. Biochemistry of insects. Academic Press, New York. [20] Gortner, R. A., and W. A. Gortner, ed. 1949. Outlines of biochemistry. Ed. 3. J. Wiley, New York. [21] Griffith, J. Q., and E. J. Farris, ed. 1949. The rat in laboratory investigation. Ed. 2. J. B. Lippincott, Philadelphia. [22] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [23] Hambidge, G., ed. 1941. Hunger signs in crops. American Society of Agronomy and National Fertilizer Association, Washington, D. C. [24] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [25] Heilbrunn, L. V. 1952. An outline of general physiology. Ed. 3. W. B. Saunders, Philadelphia. [26] Hoagland, D. R. 1948. Mineral nutrition of plants. *Chronica Botanica*, Waltham, Mass. [27] House, H. L. 1961. *Ann. Rev. Entomol.* 6:13. [28] House, H. L. 1962. *Ann. Rev. Biochem.* 31:653. [29] Hutner, S. H., et al. 1950. *Proc. Am. Phil. Soc.* 94:152. [30] Jolliffe, N., et al. 1962. Clinical nutrition. Ed. 2. P. B. Hoeber, New York. [31] Kostychev, S. P. 1931. Chemical plant physiology. Blakiston, Philadelphia. [32] Kraus, E. J. 1939. *Yearbook Agr. (U.S. Dept. Agr.)*, p. 405. [33] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [34] Lilly, V. G., and H. L. Barnett. 1951. Physiology of the fungi. McGraw-Hill, New York. [35] Loosli, J. K. 1950-51. In D. E. H. Frear, ed. *Agricultural chemistry*. Van Nostrand, New York. v. 1, p. 615. [36] Lwoff, A., ed. 1951. Biochemistry and physiology of protozoa. Academic Press, New York. [37] McCollum, E. V., E. Orent-Keiles, and H. G. Day. 1939. The newer knowledge of nutrition. Ed. 5. Macmillan, New York. [38] Madsen, L. L. 1942. *Yearbook Agr. (U.S. Dept. Agr.)*, p. 323. [39] Maynard, L. A., and J. K. Loosli. 1962. Animal nutrition. Ed. 5. McGraw-Hill, New York. [40] Miller, E. C. 1938. Plant physiology. Ed. 2. McGraw-Hill, New York. [41] Mitchell, H. H., and F. J. McClure. 1937. *Bull. Natl. Res. Council (U.S.)* 99. [42] Monier-Williams, G. W. 1949. Trace elements in food. J. Wiley, New York. [43] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [44] Prescott, S. C., and G. C. Dunn. 1959. Industrial microbiology. Ed. 3. McGraw-Hill, New York. [45] Sarles, W. B., et al. 1956. Microbiology, general and applied. Ed. 2. Harper and Row, New York. [46] Schaible, P. J. 1941. *Poultry Sci.* 20:278. [47] Sherman, H. C. 1952. Chemistry of food and nutrition. Ed. 8. Macmillan, New York. [48] Shohl, A. T. 1939. Mineral metabolism. Reinhold, New York. [49] Smith, G. M. 1951. Manual of phycology. *Chronica Botanica*, Waltham, Mass. [50] Stephenson, M. 1949. Bacterial metabolism. Ed. 3. Longmans, Green; New York. [51] Steward, F. C., ed. 1963. Plant physiology. Academic Press, New York. v. 3. [52] Sumner, J. B., and G. F. Somers. 1953. Chemistry and methods of enzymes. Ed. 3. Academic Press, New York. [53] Underwood, E. J. 1962. Trace elements in human and animal nutrition. Academic Press, New York. [54] Wallace, T. 1950. Trace elements in plant physiology. *Chronica Botanica*, Waltham, Mass. [55] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [56] West, E. S., and W. R. Todd. 1961. Textbook of biochemistry. Ed. 3. Macmillan, New York. [57] Wickersham, L. J. 1951. *U.S. Dept. Agr. Tech. Bull.* 1029. [58] Wigglesworth, V. B. 1961. The principles of insect physiology. Ed. 4. Methuen, London.



## 44. NUTRIENTS: LIPIDS

**Abbreviations:** Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required; R̄ = not required; rm = required by one or more mutants; u = utilized; ū = utilized as effectively as a related substance; s = stimulates growth or other processes; i = inhibits growth or other processes.

Nutrient (A)	Verte- brata (B)	Invertebrata		Phyto- flagellata (green) <sup>e</sup> (E)	Algae (F)	Bacterio- phyta (G)	Fungi		Sperma- tophyta (J)
		Insecta (C)	Protozoa <sup>1</sup> (D)				Saccharo- mycetaceae (H)	Other (I)	
<b>Sterols</b>									
1 Cholesterol	R̄	R <sup>3</sup>	r <sup>4</sup>	R̄	R̄	u	u	r <sup>5</sup> , s	R̄
2 7-Dehydrocholesterol	u	ū	R̄, ū <sup>6</sup>	R̄	R̄	R̄	R̄	r <sup>5</sup>	R̄
3 Ergosterol acetate	s <sup>7</sup>	....	R̄, ū <sup>6</sup>	R̄	R̄	R̄	R̄	R̄	R̄
4 Ergosterol	u	ū	R̄, ū <sup>6</sup>	R̄	R̄	R̄	u, s	s	R̄
5 Stigmasterol	R̄, s	ū	r <sup>8</sup>	R̄	R̄	R̄	R̄	r <sup>5</sup>	R̄
<b>Long-Chain Fatty Acids and Their Derivatives</b>									
6 Arachidonic acid	r, ū	s	....	R̄	R̄	....	R̄	.....	R̄
7 Linoleic acid	r, ū	r	ū	R̄	R̄	s, i	R̄	r, rm	R̄
8 Linolenic acid	r, ū	ū	....	R̄	R̄	....	R̄	r	R̄
9 Oleic acid	R̄	r	r, s, i	R̄	R̄	r, s, i	R̄	r, rm, s	R̄
10 Palmitic acid	....	....	r <sup>9</sup>	...	...	R	...	.....	...
11 Myrj G 2144 <sup>10</sup>	R̄	R̄	R̄, s	R̄	R̄	R̄	R̄	R̄	R̄
12 Tween 80, 85 <sup>11</sup>	R̄	R̄	R̄, s	R̄	R̄	R̄, s, i	R̄	R̄, s	R̄
<b>Phospholipids</b>									
13 Lecithin <sup>12</sup>	R̄	ū, s	s	R̄	R̄	R̄	R̄	R̄	R̄
Reference	7,14,21	5,9,27	2,6,12,13, 17,22,26	2,6,13,17, 18	1,11,18, 24	4,8,19,20, 25,26	4,8,19,25, 26	3,4,8,10, 19,25,26	15,16,23

<sup>1/2</sup> Including the colorless Phytoflagellata. <sup>2/3</sup> Also Dinoflagellata and Chrysomonadina. <sup>3/4</sup> Several insect species utilize various sterols as effectively as cholesterol. <sup>4/5</sup> Required by various *Trichomonas* species. <sup>5/6</sup> Required by *Labyrinthula vitella pacifica* only, which also requires fucosterol, campesterol, β-sitosterol, cionosterol, brassicasterol, and poriferasterol. <sup>6/7</sup> Utilized in place of cholesterol by *Trichomonas*. <sup>7/8</sup> Relieves stiffness syndrome of guinea pig. <sup>8/9</sup> Required by *Paramecium aurelia*; the organism's requirement for sitosterol has also been noted. <sup>9/10</sup> Required by *Trichomonas gallinae*. <sup>10/11</sup> A synthetic detergent (polyoxalkaline derivative of oleic acid). <sup>11/12</sup> Synthetic detergents (sorbitan esters of fatty acids, e.g., oleic). <sup>12/13</sup> A poorly defined, complex mixture of di-esters of a-glycerophosphoryl choline, with many unsaturated fatty acids and other substances (especially amino acids).

**Contributors:** (a) Briggs, George M., (b) Cantino, Edward C., (c) Gordon, Harold Thomas, (d) Haskins, R. H., (e) House, Howard L., (f) Loeffler, John B., (g) Pelletier, Réal L., (h) Shorb, Mary S., (i) Turrell, Franklin M., (j) Van Wagtenonk, W. J.

**References:** [1] Burlew, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [2] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [3] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [4] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [5] Gilmour, D. 1961. Biochemistry of insects. Academic Press, New York. [6] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [7] Hassinen, J. B., et al. 1950. Arch. Biochem. 25:91. [8] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [9] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [10] Hutner, S. H., and G. G. Holz, Jr. 1962. Ann. Rev. Microbiol. 16:189. [11] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [12] Lund, P. G., and M. S. Shorb. 1962. J. Protozool. 9:151. [13] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [14] Maynard, L. A., and J. K. Loosli. 1962. Animal nutrition. Ed. 5. McGraw-Hill, New York. [15] Meyer, B. S., and D. B. Anderson. 1952. Plant physiology. Ed. 2. Van Nostrand, New York. [16] Miller, E. C. 1938. Plant physiology. Ed. 2. McGraw-Hill, New York. [17] Miner, R. W. 1953. Ann. N. Y. Acad. Sci. 56:815. [18] Myers, J. 1951. Ann. Rev. Microbiol. 5:157. [19] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [20] Power, D. H., and M. J. Pelczar, Jr. 1959. J. Bacteriol. 77:789. [21] Rosenberg, H. R. 1945. Chemistry and physiology of the vitamins. Rev. ed. Interscience, New York.

continued

44. NUTRIENTS: LIPIDS

- [22] Shorb, M. S., and P. G. Lund. 1959. *J. Protozool.* 6:122. [23] Skoog, F., ed. 1951. Plant growth substances. Univ. Wisconsin Press, Madison. [24] Smith, G. M. 1951. Manual of phycology. *Chronica Botanica*, Waltham, Mass. [25] Stephenson, M. 1949. Bacterial metabolism. Ed. 3. Longmans, Green; New York. [26] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [27] Wigglesworth, V. B. 1961. The principles of insect physiology. Ed. 4. Methuen, London.

45. NUTRIENTS: PROTEINS, PEPTIDES, AND AMINO ACIDS

Amino acids not known to be required from the environment have been omitted (e.g., hydroxyproline, iodogorgoic acid, norleucine, ornithine, thyroxine). No distinction has been made between dextro- and levo-isomers, although the levo-isomers are usually nutritionally superior. In most studies with multicellular animals, microorganisms were present and in some instances supplied one or more amino acids. When requirement and utilization are noted together, at least one member of the group of organisms requires the amino acid specifically for energy or synthesis of other compounds, and other members, although not requiring it, utilize it as a general nitrogen source. *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required; rm = required by one or more mutants; R̄ = not required; U and u = utilized as a source of nitrogen and/or carbon although not a specific requirement; ū = replaces effectively one or more of the other amino acids, one of the interchangeable series being required in the diet; ū = not utilized; s = stimulates growth or other processes; \* = serves as a complete nitrogen source; \*\* = serves as the simplest complete nitrogen source.

Nutrient	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Organic nitrogen (per se)	R, U*	R, U*	r, U*	r, u*	R̄, u*	r, u* <sup>3</sup>	r, rm, u* <sup>3</sup>	r, rm, u* <sup>3</sup>	R̄, u*
2 Proteins (per se)	R̄ <sup>4</sup> , s, U*	R̄ <sup>4</sup> , s, U*	r <sup>5</sup> , u*	u	R̄	R̄, u*	R̄, u*	R̄, u*	R̄
3 Polypeptides <sup>6</sup> , peptones	R̄, U	R̄	r, u*	u* <sup>7</sup>	R̄, u*	r, u* <sup>8</sup>	R̄, u*	R̄, u*	R̄, u*
4 Amino acids	R, U**	R, U**	r <sup>9</sup> , u*	r, u <sup>7</sup> , u* <sup>10</sup>	R̄, u* <sup>11</sup>	r <sup>12</sup> , rm, u*	rm, u* <sup>13</sup>	r, rm, u*	R̄, u* <sup>14</sup>
5 Alanine	R̄, U	r, ū	u	R̄, u*	R̄, u*	r, u*	u*	u*, u <sup>15</sup>	R̄, u*
6 Arginine	r <sup>16</sup> , U	R <sup>17</sup>	r, s, u	R̄, u*	R̄, u*	r, rm, u*	rm, u*	r, rm, u*	R̄, u*
7 Aspartic acid	R̄, U	r <sup>18</sup> , ū	u, s	r, u*	R̄, u*	r, u*	u*	u*, u <sup>15</sup>	R̄, u*
8 Citrulline	R̄, ū, U	u, u <sup>17</sup>	u	.....	R̄, u*	r, u*	.....	rm, u*	R̄
9 Cysteine	R̄, U	r, u	u	.....	R̄, u*	r, u*	u*	r, rm, u*	R̄, u*
10 Cystine	R̄, U	r <sup>19</sup> , ū	u	.....	R̄	r, rm, u*	u*	rm, u*	R̄, u*
11 Glutamic acid	s, U	r <sup>20</sup>	u, s	R̄, u*	R̄, u*	r, u*	u*	r, rm, u <sup>15</sup> , u*	R̄, u*
12 Glycine	r <sup>20</sup> , U	r	r, u	R̄, u*	R̄, u*	r, u*	u*	rm, u*	R̄, u*

/1/ Including the colorless Phytoflagellata. /2/ Also Dinoflagellata and Chrysomonadina. /3/ Most species grow better on organic than on inorganic nitrogen. /4/ On the assumption that suitable amino acid combinations can replace complete proteins. /5/ Many species require living prey. /6/ See streptogenin (Table 48). /7/ Photoautotrophs, growing in the dark, require polypeptides, peptones, or amino acids. /8/ Some bacteria directly assimilate entire peptides, polypeptides, and low-molecular-weight proteins. /9/ *Glaucoma scintillans*, *Herpetomonas culicidarum*, *Tetrahymena geleii*, and *Tritrichomonas foetus* require arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, tryptophan, and valine; in addition, *G. scintillans* and *T. foetus* require glycine, proline, serine, and threonine, and *H. culicidarum* requires tyrosine. *Paramecium aurelia* requires arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tryptophan, and tyrosine. *Trichomonas gallinae* requires proline. *Tetrahymena geleii* needs no other carbon source. /10/ Various species and varieties differ markedly in utilization. /11/ Based mainly on *Chlorella pyrenoidosa*. /12/ Species differ widely in requirement. /13/ Amino acid mixtures are superior to NH<sub>4</sub><sup>+</sup> as a nitrogen source for some fungi (e.g., *Saccharomyces cerevisiae*, *S. carlsbergensis*). /14/ Several tested intact plants (tomato, tobacco, clover, pea, orchid embryo, young orchid) grew on single amino acids as the sole nitrogen source. Growth attained on some amino acids is superior to that achieved with NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup> as the nitrogen source; on other amino acids the effect is inferior. Some plants grow less well on amino acids than on inorganic nitrogen. Species differ markedly in amino acid utilization. /15/ Interchangeable for a *Neurospora* mutant. /16/ Required by rat and chicken. /17/ Arginine and citrulline are interchangeable for some insects; citrulline can partly replace arginine in *Drosophila melanogaster*. /18/ *Phormia regina* seems to require aspartic acid or glutamic acid, but not both. /19/ Some species apparently require either cystine or methionine, but not both, and proline. *Blattella germanica* requires neither cystine nor methionine in the presence of inorganic sulfates; threonine and tryptophan possibly are not required. /20/ Required by chick for rapid growth.

*continued*

**45. NUTRIENTS: PROTEINS, PEPTIDES, AND AMINO ACIDS**

Nutrient	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Amino acids									
13 Histidine	R <sup>21</sup>	R	r, s, u	r, u*	R, u*	r, rm, u*	r, u*	r, rm, u*, s	R, u*
14 Isoleucine	R	R	r, u	.....	R, u*	r, rm, u*	rm, u*	rm, u*	R, u*
15 Leucine	R	R, u <sup>1</sup>	r, u	R, u*	R, u	r, rm, u*	rm, u*	r, rm, u*	R, u*
16 Lysine	R	R	r, u	R, u*	R, u	r, rm, u*	u*	rm, u*, s	R, u*
17 Methionine	R <sup>22</sup>	R <sup>19</sup>	r, u	r	R, u	r, rm, u*	r, u*	r, rm, u*	R, u*
18 Phenylalanine	R <sup>22</sup>	R, u <sup>1</sup>	r, u	R, u*	R, u	r, rm, u*	rm, u*	rm, u*, s	R, u*
19 Proline	r, U	r <sup>19</sup>	r, u	R, u*	R, u	r, u*	u*	rm, u*, s	R, u*
20 Serine	R, U	r	r, s, u	R, u*	R, u	r, u*	u*	rm, u*	R, u*
21 Threonine	R	R <sup>19</sup> , u <sup>1</sup>	r, u	.....	R, u	r, u*	u*	rm, u*	R, u*
22 Tryptophan <sup>23</sup>	R	R <sup>19</sup> , s, u <sup>1</sup>	r, u	R, u	R, u*	r, rm, u*	u*	r, rm, u*, s	R, u*
23 Tyrosine	R, U	u <sup>24</sup>	r, u	R, u	.....	r, rm, u*	u*	rm, u*, s	R, u*
24 Valine	R	R, u <sup>1</sup>	r, s, u	R, u*	R, u*	r, rm, u*	u*	rm, u*	R, u*
Reference	1,5,11, 28,34, 39,44, 45,67, 80,95	36,48, 49,96	10,19,27, 41-43,51, 52,58,59, 69,75,76, 81,92	10,19,21- 23,40-42, 55,58- 57-59,69, 79,83,92	3,7,9,31, 55,58- 60,69, 71,72, 77,83	14,20,74, 84,87,94	8,16-18,24-26, 29,30,37,38, 46,47,53,54, 61-66,70,74, 78,87,90,91,94	4,12,13,32, 33,46,56,74, 87,89,94	2,6,15, 35,50,68, 73,82,85, 86,88,93

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>2/2</sup> Also Dinoflagellata and Chrysomonadina. <sup>3/3</sup> Some species apparently require either cystine or methionine, but not both, and proline. *Blattella germanica* requires neither cystine nor methionine in the presence of inorganic sulfates; threonine and tryptophan possibly are not required. <sup>4/4</sup> Histidine is not required to maintain nitrogen balance in adult man. <sup>5/5</sup> The amount of methionine required by man depends on the amount of cystine in the diet, and the amount of phenylalanine required depends on the amount of tyrosine in the diet. <sup>6/6</sup> Precursor of niacin; spares niacin for some organisms. <sup>7/7</sup> May replace phenylalanine in certain insects but not in others.

**Contributors:** (a) Briggs, George M., (b) Cantino, Edward C., (c) Fogg, G. E., (d) Gordon, Harold Thomas, (e) House, Howard L., (f) Koser, Stewart A., (g) Kratzer, F. H., (h) Loefer, John B., (i) Pelletier, Réal L., (j) Purvis, E. R., (k) Shorb, Mary S., (l) Van Wagtenonk, W. J.

**References:** [1] Anderson, R. J. 1949. Essentials of physiological chemistry. J. Wiley, New York. [2] Audus, L. J., and J. H. Quastel. 1947. Nature 160:222. [3] Bold, H. C. 1942. Botan. Rev. 8:70. [4] Bonner, D. 1946. Cold Spring Harbor Symp. Quant. Biol. 9:14. [5] Bourne, G. H., and G. W. Kidder, ed. 1953. Biochemistry and physiology of nutrition. Academic Press, New York. [6] Brigham, R. O. 1917. Soil Sci. 3:155. [7] Brunel, J. B., et al. 1950. Culturing of algae; symposium. C. F. Kettering Foundation, Dayton, Ohio. [8] Brunton, T. L., and A. Macfadyen. 1889. Proc. Roy. Soc. (London) 46:542. [9] Burlew, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [10] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [11] Cannon, P. R. 1948. Federation Proc. 7:391. [12] Cantino, E. C., and G. Turian. 1959. Ann. Rev. Microbiol. 13:97. [13] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [14] Cowperthwaite, J., et al. 1953. Ann. N. Y. Acad. Sci. 56:972. [15] Crowther, E. M. 1925. J. Agr. Sci. 15:300. [16] Desnuelle, P. 1939. Enzymologia 6:242. [17] Desnuelle, P. 1940. Ibid. 6:387. [18] Desnuelle, P., and C. Fromageot. 1939. Ibid. 6:80. [19] Doyle, W. L. 1942. Biol. Rev. Cambridge Phil. Soc. 18:119. [20] Dunn, M. S. 1949. Physiol. Rev. 29:219. [21] Dusi, H. 1933. Ann. Inst. Pasteur 50:550. [22] Dusi, H. 1933. Ibid. 50:840. [23] Dusi, H. 1936. Arch. Zool. Exptl. Gen. 78:133. [24] Ehrlich, F. 1907. Ber. Deut. Chem. Ges. 40:1027. [25] Ehrlich, F. 1911. Ibid. 44:139. [26] Ehrlich, F. 1912. Ibid. 45:883. [27] Elliott, A. M. 1949. Physiol. Zool. 22:337. [28] Ewing, W. R. 1951. Poultry nutrition. The author, South Pasadena, Calif. [29] Fildes, P. 1941. Brit. J. Exptl. Pathol. 22:293. [30] Fildes, P. 1945. Ibid. 26:416. [31] Fogg, G. E. 1953. The metabolism of algae. Methuen, London. [32] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [33] Fries, N. 1961. In W. Ruhland, ed. Encyclopedia of plant physiology. J. Springer, Berlin. v. 14, p. 33. [34] Fruton, J. S., and S. Simmonds. 1958. General biochemistry. Ed. 2. J. Wiley, New York.

continued



**45. NUTRIENTS: PROTEINS, PEPTIDES, AND AMINO ACIDS**

- [35] Ghosh, B. P., and R. H. Burris. 1950. *Soil Sci.* 70:187. [36] Gilmour, D. 1961. *Biochemistry of insects*. Academic Press, New York. [37] Gorbach, G. 1930. *Arch. Mikrobiol.* 1:537. [38] Gorini, C., et al. 1932. *Z. Physiol. Chem.* 205:133. [39] Gortner, R. A., and W. A. Gortner, ed. 1949. *Outlines of biochemistry*. Ed. 3. J. Wiley, New York. [40] Hall, R. P. 1939. *Quart. Rev. Biol.* 14:1. [41] Hall, R. P. 1943. *Vitamins Hormones* 1:249. [42] Hall, R. P. 1953. *Protozoology*. Prentice-Hall, New York. [43] Hall, R. P., and H. W. Schoendorn. 1939. *Physiol. Zool.* 12:201. [44] Harrow, B. 1962. *Textbook of biochemistry*. Ed. 8. W. B. Saunders, Philadelphia. [45] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1954. *Practical physiological chemistry*. Ed. 13. Blakiston, New York. [46] Hawker, L. E. 1950. *Physiology of fungi*. Univ. London Press, London. [47] Hills, G. M. 1940. *Biochem. J.* 34:1057. [48] House, H. L. 1961. *Ann. Rev. Entomol.* 6:13. [49] House, H. L. 1962. *Ann. Rev. Biochem.* 31:653. [50] Hutchinson, H. B., and N. H. J. Miller. 1911. *Centr. Bakteriell. Parasitenk.*, II, 30:513. [51] Kidder, G. W. 1951. *Ann. Rev. Microbiol.* 5:139. [52] Kidder, G. W., and V. C. Dewey. 1945. *Arch. Biochem.* 6:425. [53] Knight, B. C. J. G. 1945. *Vitamins Hormones* 3:105. [54] Kohn, H. E., and J. S. Harris. 1942. *J. Bacteriol.* 44:717. [55] Lewin, R. A., ed. 1962. *Physiology and biochemistry of algae*. Academic Press, New York. [56] Lilly, V. G., and H. L. Barnett. 1951. *Physiology of the fungi*. McGraw-Hill, New York. [57] Lwoff, A. 1938. *Arch. Protistenk.* 90:194. [58] Lwoff, A. 1943. *L'évolution physiologique*. Hermann, Paris. [59] Lwoff, A., ed. 1951-55. *Biochemistry and physiology of protozoa*. Academic Press, New York. [60] Mainx, F. 1929. *Tabulae Biologicae* 5:1. [61] Maschmann, E. 1937. *Biochem. Z.* 294:1. [62] Maschmann, E. 1937-38. *Ibid.* 295:1. [63] Maschmann, E. 1937-38. *Ibid.* 295:351. [64] Maschmann, E. 1937-38. *Ibid.* 295:391. [65] Maschmann, E. 1937-38. *Ibid.* 295:400. [66] Maschmann, E. 1938-39. *Ibid.* 300:89. [67] Maynard, L. A., and J. K. Loosli. 1962. *Animal nutrition*. Ed. 5. McGraw-Hill, New York. [68] Miller, E. C. 1938. *Plant physiology*. Ed. 2. McGraw-Hill, New York. [69] Miner, R. W. 1953. *Ann. N. Y. Acad. Sci.* 56:815. [70] Monod, J. 1958. *Recherches sur la croissance des cultures bactériennes*. Hermann, Paris. [71] Myers, J. 1944. *Plant Physiol.* 19:579. [72] Myers, J. 1951. *Ann. Rev. Microbiol.* 5:157. [73] Nightingale, G. T. 1947. *Botan. Rev.* 3:85. [74] Porter, J. R. 1946. *Bacterial chemistry and physiology*. J. Wiley, New York. [75] Pringsheim, E. G. 1937. *Nature* 139:196. [76] Pringsheim, E. G. 1937. *Planta* 27:61. [77] Pringsheim, E. G. 1946. *Pure cultures of algae*. Cambridge Univ. Press, London. [78] Regneroy, D. C. 1944. *J. Biol. Chem.* 154:151. [79] Reinhardt, K. 1950. *Arch. Mikrobiol.* 15:270. [80] Rose, W. C. Unpublished. Univ. Illinois Dept. Chemistry, Urbana, 1954. [81] Scheer, B. T. 1948. *Comparative physiology*. J. Wiley, New York. [82] Skoog, F., ed. 1951. *Plant growth substances*. Univ. Wisconsin Press, Madison. [83] Smith, G. M. 1951. *Manual of phycology*. *Chronica Botanica*, Waltham, Mass. [84] Snell, E. E. 1945. *Advan. Protein Chem.* 2:85. [85] Spoerl, E. 1948. *Am. J. Botany* 35:88. [86] Steinberg, R. A. 1947. *J. Agr. Res.* 75:81. [87] Stephenson, M. 1949. *Bacterial metabolism*. Ed. 2. Longmans, Green; New York. [88] Tanaka, I. 1931. *Japan. J. Botany* 5:323. [89] Tatum, E. L. 1944. *Ann. Rev. Biochem.* 13:667. [90] Tatum, E. L., and D. Bonnel. 1944. *Proc. Natl. Acad. Sci. U.S.* 30:30. [91] Thorne, R. S. W. 1937. *J. Inst. Brewing* 43:288. [92] Trager, W. 1941. *Physiol. Rev.* 21:1. [93] Virtanen, A. I., and H. Linkola. 1946. *Nature* 158:515. [94] Werkman, C. H., and P. W. Wilson, ed. 1951. *Bacterial physiology*. Academic Press, New York. [95] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York. [96] Wigglesworth, V. B. 1961. *The principles of insect physiology*. Ed. 4. Methuen, London.



## 46. NUTRIENTS: PURINES AND PYRIMIDINES

Purine and pyrimidine compounds are essential components of the nucleic acids. Inability to synthesize these compounds makes it necessary for many organisms to obtain them from the environment or diet. For some organisms the requirement is for a specific compound, or compounds; for others any one of an interchangeable series of compounds satisfies the need. *Abbreviations:* Capital letter indicates data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R = not required; r = required; rm = required by one or more mutants; u = utilized;  $\bar{u}$  = not utilized;  $\bar{r}$  = utilized as effectively as, or is interchangeable with, one or more related compounds, the presence of at least one of the series being required; u< = partially replaces, or spares the use of, one or more required, or interchangeably required, compounds; s = stimulates growth or other processes; i = inhibits growth or other processes.

Nutrient	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Purine compounds	R, s	r	r, s	R	R	r, s, u	s, u	rm, s, u	R
2 Adenine	R, s	r, s	r <sup>3</sup> , u< <sup>3</sup>	R, u< <sup>4</sup>	R	$\bar{r}$ <sup>5</sup>	r <sup>6</sup>	r, rm, u, s	R
3 Adenosine	R	$\bar{r}$ , s	R, u< <sup>3</sup>	R	R	$\bar{r}$ <sup>5</sup>	R	R, u	R
4 Adenosine triphosphate	R	$\bar{r}$	R, s	R	R	$\bar{r}$ <sup>5</sup>	R	R	R
5 Adenylic acid	R, s	$\bar{r}$ , s	r, u< <sup>3</sup>	R	R	$\bar{r}$ <sup>5</sup>	R	R, u	R
6 Guanine	R	r	r <sup>7</sup>	s	R	$\bar{r}$ <sup>5, 8, 9, s</sup>	R	r, rm, s	R
7 Guanosine	R	$\bar{r}$ , s	r <sup>7</sup> , $\bar{r}$ , u< <sup>9</sup> , i	R	R	$\bar{r}$ <sup>5</sup>	R	R, u	R
8 Guanylic acid	R	$\bar{r}$ , s	$\bar{r}$ , s	R	R	$\bar{r}$ <sup>5</sup>	R	R, u	R
9 Hypoxanthine	R	u	$\bar{r}$ , u< <sup>3</sup>	s	R	$\bar{r}$ <sup>5, s</sup>	R	rm, u, s	R
10 Xanthine	R	s	r <sup>7, 10</sup> , $\bar{u}$	R	R	$\bar{r}$ <sup>5</sup>	R	R, u	R
11 Others	...	....	s <sup>11</sup>	.....	...	.....	...	u	R
12 Pyrimidine compounds	R, s	r	r, s	r	r	r, s, u	r, s	r, rm, s	R
13 Cytidine	R	R	$\bar{r}$ , i	R	R	$\bar{r}$ <sup>12</sup>	R	R, rm, s	R
14 Cytidylic acid	R	R	$\bar{r}$ , s	R	R	$\bar{r}$ <sup>12</sup>	R	R, rm, s <sup>13</sup>	R
15 Cytosine	R	R	r <sup>7</sup> , $\bar{u}$ , i	R	R	$\bar{r}$ <sup>5</sup>	R	rm, u	R
16 Orotic acid	R, s	R	R, $\bar{u}$	R	R	r <sup>14</sup> , s	R	rm, s <sup>13</sup> , u	R
17 Pyrimidine	R, $\bar{u}$	R	r, u <sup>15</sup>	r, u <sup>15</sup>	r <sup>15</sup>	r, u <sup>15</sup>	r, s <sup>15</sup>	r, rm, u <sup>15</sup>	R
18 Thymine	R, s	r, s	r <sup>7</sup> , u< <sup>16</sup>	R, u< <sup>4</sup>	R	$\bar{r}$ <sup>17</sup> , u< <sup>18</sup>	R	R, u	R
19 Thymidine	R, s	$\bar{r}$	u< <sup>16</sup>	R	R	u< <sup>18</sup>	R	R	R
20 Uracil	R	r	r, $\bar{r}$ <sup>19</sup>	r <sup>7</sup>	R	$\bar{r}$ <sup>12, s</sup>	R	rm, s <sup>13</sup>	R
21 Uridine	R	R	$\bar{r}$	R	R	$\bar{r}$ <sup>12, s</sup>	R	R, rm, s	R
22 Uridylic acid	R	R	$\bar{r}$	R	R	$\bar{r}$ <sup>12, s</sup>	R	R, rm, s <sup>13</sup>	R
Reference	1,8,12, 15,16	11,18, 19,21	6,13,22-24, 27-29,42	6,13,14, 21,28-30	2,4,5,21,26, 30,33,36,40	8,10,17, 21,31,35, 41,44,45	8,10,17,21, 31,35,44-46	7,8,10,17, 21,31,34,35, 37,44,45	3,9,20,25, 32,38,39, 43

<sup>1/</sup> Including the colorless Phytoflagellata. <sup>2/</sup> Also Dinoflagellata and Chrysoomonadina. <sup>3/</sup> Spares, but cannot replace, guanine, guanosine, and guanylic acid in *Tetrahymena*. <sup>4/</sup> Spares and, when given with amino acids, substitutes for *p*-aminobenzoic and pteroylglutamic acids in *Euglena gracilis*. <sup>5/</sup> Items 2-10 are variously interchangeable for different bacteria, but at least one must be present (e.g., *Streptococcus pyogenes* requires at least one of adenine, adenosine, adenylic acid, guanine, guanosine, guanylic acid, xanthine); requirement is relieved by CO<sub>2</sub> in high concentration. *S. lactis* requires adenine, guanine, hypoxanthine, or xanthine. *Bacillus megaterium* requires adenosine for spore germination. <sup>6/</sup> Required by *Saccharomyces octosporus* on certain media. <sup>7/</sup> Required by *Tetrahymena*, but is replaceable by guanosine or guanylic acid. <sup>8/</sup> *Lactobacillus plantarum* and *Leuconostoc mesenteroides* require cytosine or guanine (interchangeable). <sup>9/</sup> Spares pteroylglutamic acid in *Herpetomonas culicidarum*. <sup>10/</sup> In vitro studies indicate possibility that *Plasmodium* requires xanthine (and also cytosine, pyrimidine, uracil, adenine, and guanosine). <sup>11/</sup> Methyl purines (theobromine, theophylline, caffeine) stimulate some ciliates and Suctoria. <sup>12/</sup> At least one required, but interchangeable, for several species (e.g., *Clostridium tetani*, *Haemophilus parainfluenzae*). <sup>13/</sup> Stimulates mutants of *Neurospora*. <sup>14/</sup> Required by *Lactobacillus bulgaricus* 09. <sup>15/</sup> Pyrimidine moiety of thiamine. <sup>16/</sup> Spares pteroylglutamic acid in *Tetrahymena*. <sup>17/</sup> *Clostridium tetani* requires either adenine or hypoxanthine (interchangeable). <sup>18/</sup> Spares pteroylglutamic acid in *Streptococcus lactis*; thymine + thymidine replaces pteroylglutamic acid in *S. lactis*. <sup>19/</sup> Replaces, or is interchangeable with, cytidine, cytidylic acid, uridine, or uridylic acid, in *Tetrahymena*.

*Contributors:* (a) Briggs, George M., (b) Cantino, Edward C., (c) Fogg, G. E., (d) Gordon, Harold Thomas, (e) House, Howard L., (f) Koser, Stewart A., (g) Loefer, John B., (h) Shorb, Mary S., (i) Turrell, Franklin M., (j) Van Wagtenonk, W. J.

*References:* [1] Anderson, R. J. 1949. Essentials of physiological chemistry. J. Wiley, New York. [2] Bold, H. C. 1942. Bot. Rev. 8:70. [3] Bonner, D. M., and J. Bonner. 1940. Am. J. Botany 27:38. [4] Brunel, J. B., et al. 1950. Culturing of algae; symposium. C. F. Kettering Foundation, Dayton, Ohio. [5] Burlew, J. S. 1953.

continued

### 46. NUTRIENTS: PURINES AND PYRIMIDINES

Carnegie Inst. Wash. Publ. 600. [6] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [7] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [8] Davidson, J. N. 1960. Biochemistry of the nucleic acids. Ed. 4. J. Wiley, New York. [9] Deysson, M. 1953. Bull. Soc. Botan. France 100:14. [10] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [11] Gilmour, D. 1961. Biochemistry of insects. Academic Press, New York. [12] Gortner, R. A., and W. A. Gortner, ed. 1949. Outlines of biochemistry. Ed. 3. J. Wiley, New York. [13] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [14] Hamilton, L. 1953. Ann. N. Y. Acad. Sci. 56:961. [15] Harrow, B. 1962. Textbook of biochemistry. Ed. 8. W. B. Saunders, Philadelphia. [16] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1954. Practical physiological chemistry. Ed. 13. Blakiston, New York. [17] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [18] House, H. L. 1961. Ann. Rev. Entomol. 6:13. [19] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [20] Jones, R. F., and H. G. Baker. 1943. Ann. Botany (London) 7:379. [21] Kalckar, H. M. 1951. Symp. Soc. Exptl. Biol. 1:38. [22] Kidder, G. W. 1947. Ann. N. Y. Acad. Sci. 49:99. [23] Kidder, G. W. 1951. Ann. Rev. Microbiol. 5:139. [24] Kidder, G. W., and V. C. Dewey. 1948. Proc. Natl. Acad. Sci. U.S. 34:566. [25] Kotsovsky, D. 1942. Biol. Muenschen 11:276. [26] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [27] Lilly, D. M., F. J. Sterbenz, and V. Tarantola. 1953. Proc. Soc. Exptl. Biol. Med. 83:434. [28] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [29] Miner, R. W. 1953. Ann. N. Y. Acad. Sci. 56:815. [30] Myers, J. 1951. Ann. Rev. Microbiol. 5:157. [31] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [32] Porutskii, G. V. 1949. Dokl. Akad. Nauk SSSR 64:103. [33] Pringsheim, E. G. 1946. Pure cultures of algae. Cambridge Univ. Press, London. [34] Robbins, W. J., and V. Kavanagh. 1942. Botan. Rev. 8:411. [35] Robinson, F. A. 1951. The vitamin B complex. J. Wiley, New York. [36] Sager, R., and S. Granick. 1953. Ann. N. Y. Acad. Sci. 56:831. [37] Schopfer, W. H. 1943. Plants and vitamins. Chronica Botanica, Waltham, Mass. [38] Schreiner, O., and J. Skinner. 1917. U.S. Dept. Agr. Bur. Soils Bull. 87. [39] Skoog, F., ed. 1951. Plant growth substances. Univ. Wisconsin Press, Madison. [40] Smith, G. M. 1951. Manual of phycology. Chronica Botanica, Waltham, Mass. [41] Snell, E. E., and H. K. Mitchell. 1942. Arch. Biochem. 1:93. [42] Sprince, H., et al. 1953. Ann. N. Y. Acad. Sci. 56:1016. [43] Steinberg, R. A. 1947. J. Agr. Res. 75:81. [44] Stephenson, M. 1949. Bacterial metabolism. Longmans, Green; New York. [45] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [46] Williams, R. J. 1941. Biol. Rev. Cambridge Phil. Soc. 16:49.

### 47. NUTRIENTS: VITAMINS AND RELATED COMPOUNDS

It is probable that many vitamins and related compounds are indispensable participants in the metabolic activities of all living substances, and are in this sense universally "required." Data, however, have been limited to the presently known requirement, or non-requirement, for compounds obtained from the external environment or in the diet, and do not include metabolically essential compounds provided to an organism by associated microorganisms (e.g., by the intestinal flora of mammals). *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required; R̄ = not required; rm = required by one or more mutants; u = utilized; ū = utilized as effectively as a related vitamin or compound; u> = utilized more effectively than a related vitamin or compound; u< = utilized less effectively than a related vitamin; † = not utilized in place of the related vitamin; s = stimulates growth or other processes; i = inhibits growth or other processes.

Compound	Vertebrata	Invertebrata		Phyto- flagellata (green) <sup>2</sup>	Algae	Bacterio- phyta	Fungi		Sperma- tophyta
		Insecta	Protozoa <sup>1</sup>				Saccharo- mycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
<b>Vitamins</b>									
1 Vitamin A	R	R̄	r?	R̄	R̄	R̄	R̄	R̄	R̄, s
2 p-Aminobenzoic acid	R, s	R̄	r	R̄	R̄	r	r	r, rm, s	R̄

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>2/2</sup> Also Dinoflagellata and Chrysomonadina.

*continued*

*Continued*

**47. NUTRIENTS: VITAMINS AND RELATED COMPOUNDS**

Compound	Vertebrata	Invertebrata		Phyto- flagellata (green) <sup>2</sup>	Algae	Bacterio- phyta	Fungi		Sperma- tophyta
		Insecta	Protozoa <sup>1</sup>				Saccharo- mycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
<b>Vitamins</b>									
3 Ascorbic acid	r	r	r, s	H	r?, s	H, s	H, s	H, s	H, s
4 Biotin	r	r	r, s	H	r	r	r, s	r, rm, s	H
5 Choline group <sup>3</sup>	r	r	H	H	H	r	H	r, rm	H, s
6 Cobalamin <sup>4</sup>	r	r?, s	r	r	r	r <sup>5</sup>	H	H	H, s
7 Vitamin D <sup>5</sup>	r	H	H	H	H	H	H	H, s	H, s
8 Vitamin E	r	H, s	r <sup>7</sup>	H	H	H	H	H	H
9 Inositol <sup>8</sup>	r?	r	H	H	H	r?, s	r, s	r, rm, s	H, s
10 Vitamin K	r	H	r	H	H	r	H	H	H
11 Nicotinic acid	r	r	r	H	H	r, s	r	r, rm, s	H, s?
12 Nicotinamide	UF	UF	UF	H	H	r, UF, s	UF	r, rm, s	H, s?
13 Pantothenic acid	R	r	r	H	r	r, s	r	r, rm, s	H, s
14 Pteroylglutamic acid <sup>9</sup>	R	r	r	H	H	r <sup>10</sup>	H	H, s	H
15 Pyridoxal <sup>11</sup>	UF	UF	UF	H	H	r, u>	UF, s	UF	H
16 Pyridoxamine <sup>11</sup>	UF	UF	UF	H	H	r, u>	UF, s	UF	H
17 Pyridoxine	R	r	r	H	H, s	r, s	r, s	r, rm, s	H, s
18 Riboflavin	R	r	r	H	H	r, s	H	r, rm	H
19 Thiamine	R	r	r	r	r	r, s	r, i <sup>12</sup>	r, rm, s, i <sup>13</sup>	H, s
<b>Compounds Chemically Related to Vitamins</b>									
20 β-Alanine	H, s	H	H, u	H	H	H, UF	H, UF	rm, UF	H
21 Biocytin	H	H	H	H	H	u	H	H, UF	H
22 β-Carotene <sup>14</sup>	H, u<	r	H	H	H	H	H	H	H
23 Coenzyme A	H, u<	H, UF	H	H	H	r, u>	H, s	H	H
24 5,6-Dimethylbenzimidazole	H, u<	H	H	H, u<	H	H	H	H	H
25 Desthiobiotin	H	H	H	H	H	H, u>	H	H, UF	H
26 Diphosphopyridine nucleotide	H, UF?	H, UF?	H	H	H	r, UF	H, UF	r, u>, s	H
27 Diphosphothiamine	H, UF	H, UF	H, UF	H, UF	H	r, u>	H, UF	H, u<	H
28 Folic acid conjugates <sup>15</sup>	H, UF	H, UF	H, UF	H	H	H, UF	H	H	H
29 Folinic acid <sup>16</sup>	H, u<	H, UF	H, UF	H	H	r	H	H	H
30 Hesperidin <sup>17</sup>	H, s	H	H	H	H	H	H	H	H, s
31 o-Heterobiotin	H	H	H	H	H	H, u<, u	H	H	H
32 LBF <sup>18</sup>	H	H	H	H	H	r, u>	H	H	H
33 Lipothiamide	H, UF	H	H	H	H	H	H	H	H
34 Lyxoflavin	H, s, u<	H	H	H	H	H, u< <sup>19</sup>	H	H	H
35 Oxybiotin	H	H	H	H	H	H, u<	H	H, s	H
36 Pantoic acid	H, u	H	H	H	H	H, u	H, u	H	H
37 Pantothenic acid conjugate	H	H	H	H	H	r?, u>	H	H	H
38 Pimelic acid	H, u	H	H, u<	H	H	H, UF?	H	H, s	H
39 Pseudovitamin B <sub>12</sub>	H, u	H	H	H, UF	H	H, UF	H	H	H
40 Pteric acid	H, u	H	H, UF, u	H	H	H, u<	H	H	H
41 Pyridoxal-PO <sub>4</sub>	H, UF	H, UF	H	H	H	r, u>	H, u	H, UF	H
42 Pyridoxamine-PO <sub>4</sub>	H, UF	H, UF	H, UF	H	H	r, u>	H, u	H, UF	H
43 Pyrimidine	H, u	H	r, u <sup>20,21</sup>	r, u <sup>20,21</sup>	r <sup>20,21</sup>	r, u <sup>20,21</sup>	r, s <sup>20,21</sup>	r, rm, u <sup>20,21</sup>	H
44 Rhizopterine	H, u	H	H	H	H	H, UF	H	H	H
45 α-Ribazole	H, s	H	H	H, u<	H	H	H	H	H

/1/ Including the colorless Phytoflagellata. /2/ Also Dinoflagellata and Chrysomonadina. /3/ Includes choline, betaine and other methyl donors. /4/ Generic term; includes cyanocobalamin, hydroxocobalamin, vitamins B<sub>12</sub>, B<sub>12a</sub>, and B<sub>12b</sub>. /5/ For some species or strains, thymine desoxyriboside, hypoxanthine, adenine, or guanine may substitute in certain media. /6/ D<sub>2</sub> active for mammals only; D<sub>3</sub> active for all and required by chicken. /7/ Required by a variant of *Trichomonas gallinae*. /8/ Of doubtful status as a vitamin. /9/ Folic acid, folacin. /10/ For some species or strains, pteric acid, or thymine + thymidine, will substitute. /11/ Member of the pyridoxine group (vitamin B<sub>6</sub>). /12/ Inhibits growth of *Saccharomyces cerevisiae* strains. /13/ Inhibits growth of *Rhizopus nigricans*. /14/ And other carotenoid precursors of vitamin A. /15/ Di-, tri- and hepta-glutamates of pteroylglutamic acid. /16/ The citrovorum factor; required by *Leuconostoc citrovorum*. /17/ Hesperidin, rutin and citrin = vitamin P series. /18/ *Lactobacillus bulgaricus* factor = pantetheine (thiol form), and pantethine (disulfide form). /19/ Replaces riboflavin in *Lactobacillus lactis*. /20/ Thiamine or pyrimidine moiety (thiazole moiety is synthesized). /21/ Thiamine or pyrimidine + thiamine moieties (pyrimidine and thiazole moieties combine to give thiamine).

*continued*



## 47. NUTRIENTS: VITAMINS AND RELATED COMPOUNDS

Compound	Vertebrata	Invertebrata		Phyto- flagellata (green) <sup>2</sup>	Algae	Bacterio- phyta	Fungi		Sperma- tophyta
		Insecta	Protozoa <sup>1</sup>				Saccharo- mycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Compounds Chemically Related to Vitamins									
46	Rutin <sup>17</sup>	R, s	R	R	R	R	R	R, u <sup>22</sup>	R
47	Thiazole	R	R	r, u <sup>21,23</sup>	r, u <sup>21,23</sup>	r <sup>21,23</sup>	r, u <sup>21,23</sup>	r, s <sup>21,23</sup>	r, rm, u <sup>21,23</sup>
48	Triphosphopyridine nucleotide	R, u <sup>24</sup>	R	R	R	R	R, u <sup>24</sup>	R, u <sup>24</sup>	r, u <sup>24</sup> , s
49	Xanthopterin	r <sup>24</sup> , u <sup>24</sup> , s	R	R	R	R	R, u <sup>24</sup>	R	R
Reference	1-6,12-16, 29,32-34, 38,40-44, 52	6,9,19, 23,24, 40,48, 50,52	6,8,11,20, 21,25,28, 30,40,52	6,8,20, 21,28,30, 31,48	6,7, 27,30, 31,46	35,36,40, 45,47,49, 52	6,17,22,26, 36,47,49,51	6,10,17,18, 22,26,36, 37,39,45,49	4,6,45

/1/ Including the colorless Phytoflagellata. /2/ Also Dinoflagellata and Chryomonadina. /17/ Hesperidin, rutin and citrin = vitamin P series. /21/ Thiamine or pyrimidine + thiamine moieties (pyrimidine and thiazole moieties combine to give thiamine). /22/ Utilized by *Aspergillus flavus*, *Polyporus versicolor*. /23/ Thiamine or thiazole moiety (pyrimidine moiety is synthesized). /24/ More active than pteroylglutamic (folic) acid in relieving anemia of Chinook salmon.

**Contributors:** (a) Briggs, George M., (b) Cantino, Edward C., (c) Fogg, G. E., (d) Gordon, Harold Thomas, (e) Hansard, Sam L., (f) Haskins, R. H., (g) House, Howard L., (h) Koser, Stewart A., (i) Loefer, John B., (j) Pelletier, Réal L., (k) Rusoff, Louis Leon, (l) Schaefer, Arnold Edward, (m) Shorb, Mary S., (n) Turrell, Franklin M., (o) Van Wagtenonk, W. J.

**References:** [1] American Medical Association Council on Pharmacy and Chemistry and Council on Foods. 1939. The vitamins; symposium. Chicago. [2] Association of Vitamin Chemists, Inc. 1951. Methods of vitamin assay. Ed. 2. Interscience, New York. [3] Baumann, C. A. 1953. Ann. Rev. Biochem. 22:527. [4] Bessey, O. A., H. J. Lowe, and L. Salomon. 1953. Ibid. 22:545. [5] Bourne, G. H., and G. W. Kidder, ed. 1953. Biochemistry and physiology of nutrition. Academic Press, New York. [6] Briggs, G. M. Unpublished. Biochemistry and Nutrition Laboratory, Natl. Institutes of Health, Bethesda, Md., 1953. [7] Burlew, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [8] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [9] Chauvin, R. W. 1949. Physiologie de l'insecte. W. Junk, Haag. [10] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [11] Cooperman, J. M., et al. 1952. J. Nutr. 46:467. [12] Cowgill, G. R. 1934. The vitamin B requirement of man. Yale Univ. Press, New Haven. [13] Deuel, H. J. 1951-57. The lipids; their chemistry and biochemistry. Interscience, New York. [14] Eddy, W. H. 1949. Vitaminology. Williams and Wilkins, Baltimore. [15] Eddy, W. H., and G. Dalldorf. 1944. The avitaminoses. Ed. 3. Williams and Wilkins, Baltimore. [16] Ewing, W. R. 1951. Poultry nutrition. Ed. 4. The author, South Pasadena, Calif. [17] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [18] Fries, N. 1946. Trans. Brit. Mycol. Soc. 30:118. [19] Gilmore, D. 1961. Biochemistry of insects. Academic Press, New York. [20] Hall, R. P. 1943. Vitamins Hormones 1:249. [21] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [22] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [23] House, H. L. 1961. Ann. Rev. Entomol. 6:13. [24] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [25] Kidder, G. W. 1951. Ann. Rev. Microbiol. 5:139. [26] Knight, B. C. J. G. 1945. Vitamins Hormones 3:108. [27] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [28] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [29] Maynard, L. A., and J. K. Loosli. 1962. Animal nutrition. Ed. 5. McGraw-Hill, New York. [30] Miner, R. W. 1953. Ann. N. Y. Acad. Sci. 56:815. [31] Myers, J. 1951. Ann. Rev. Microbiol. 5:157. [32] National Research Council Committee on Animal Nutrition. 1953-62. Recommended nutrient allowances. Washington, D. C. nos. 1-10. [33] National Research Council Food and Nutrition Board. 1948. Nutr. Rev. 6:319. [34] Nutrition Foundation, Inc. 1953. Present knowledge in nutrition. New York. [35] Peterson, W. H., and M. S. Peterson. 1945. Bacteriol. Rev. 9:49. [36] Porter, J. R. 1946. Bacterial

continued



## 47. NUTRIENTS: VITAMINS AND RELATED COMPOUNDS

chemistry and physiology. J. Wiley, New York. [37] Raper, J. R., and A. J. Haagen-Smit. 1942. J. Biol. Chem. 143:311. [38] Reed, C. I., H. C. Struck, and I. E. Steck. 1939. Vitamin D. Univ. Chicago Press, Chicago. [39] Robbins, W. J., and V. Kavanagh. 1942. Bot. Rev. 8:411. [40] Robinson, F. A. 1951. The vitamin B complex. J. Wiley, New York. [41] Rosenberg, H. R. 1945. Chemistry and physiology of the vitamins. Rev. ed. Interscience, New York. [42] Sherman, H. C. 1943. The science of nutrition. Columbia Univ. Press, New York. [43] Sherman, H. C. 1952. Chemistry of food and nutrition. Ed. 8. Macmillan, New York. [44] Sherman, H. C., and S. L. Smith. 1931. The vitamins. Ed. 2. Chemical Catalogue, New York. [45] Skoog, F., ed. 1951. Plant growth substances. Univ. Wisconsin Press, Madison. [46] Smith, G. M. 1951. Manual of phycology. Chronica Botanica, Waltham, Mass. [47] Stephenson, M. 1949. Bacterial metabolism. Ed. 3. Longmans, Green; New York. [48] Trager, W. 1947. Biol. Rev. Cambridge Phil. Soc. 22:148. [49] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [50] Wigglesworth, V. B. 1961. The principles of insect physiology. Ed. 4. Methuen, London. [51] Williams, R. J. 1941. Biol. Rev. Cambridge Phil. Soc. 16:49. [52] Williams, R. J., et al. 1950. The biochemistry of B vitamins. Reinhold, New York.

## 48. NUTRIENTS: MISCELLANEOUS GROWTH FACTORS

Many of the compounds listed are utilized by some organisms only for their carbon, nitrogen, and/or hydrogen content (e.g., CO<sub>2</sub>, glutamine, and asparagine). *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required; H = not required; rm = required by one or more mutants; H = replaces effectively, or is utilized interchangeably with, one or more substances, but one of the interchangeable substances must be present; s = stimulates growth or other processes; i = inhibits growth or other processes.

Nutrient	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Adenylthiomethylpentose <sup>3</sup>	H, s	H	H	H	H	H	H	H	H
2 Anthranilic acid <sup>4</sup>	H, s	H	H	H	H	u <sup>5</sup>	H	r, rm <sup>6</sup>	H
3 Antibiotics <sup>5</sup>	H, s	H	H, s	H	H	r <sup>7</sup>	H	H, s	H, s
4 Asparagine	H	H	H	H	H	r <sup>8</sup>	H, s, i	H	H
5 Bifidus factor	H	H	H	H	H	r	H	H	H
6 Carbon dioxide	H	H	r <sup>9</sup>	R <sup>10</sup> , s	R	r <sup>11</sup> , s	.....	r <sup>11</sup> , s, i	R
7 Carnitine <sup>12</sup>	H	r <sup>13</sup>	H	H	H	H	H	H	H
8 Coprogen <sup>14</sup>	H	H	H	H	H	H	H	r	H
9 N-D-Glucosylglycine ester	H	H	H	H	H	r	H	H	H
10 Glutamine	H	H	H	H	H	r <sup>15</sup> , s	s, i	H, s, i	H
11 Glutathione	H	s <sup>16</sup>	H	H	H	r <sup>16</sup>	.....	H	H
12 Guanidine	H	H	r <sup>17</sup>	H	H	H	.....	H	H
13 Indole-3-acetic acid <sup>18</sup>	H, s?	H	H <sup>19</sup> , i	H, s <sup>19</sup>	H, s	H, s	H	r?, s	H, s
14 Hematin <sup>20</sup>	H	r	r	H	H	r	H	r	H
15 β-Hydroxybenzoic acid	H	H	H	H	H	rm	H	rm	H
16 Krebs cycle intermediates <sup>21</sup>	H	H	H, s <sup>22</sup>	r <sup>22</sup> , s	H	r	H	r	H

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>2/2</sup> Also Dinoflagellata and Chrysomonadina. <sup>3/3</sup> "Vitamin L<sub>2</sub>," a purine nucleoside. <sup>4/4</sup> "Vitamin L<sub>1</sub>," precursor of nicotinic acid. <sup>5/5</sup> Substitutes for tryptophan and/or indole. <sup>6/6</sup> Aureomycin, penicillin, streptomycin, bacitracin, neomycin, and other anti-infectious substances (e.g., arsenicals and sulfonamides which in small amounts may stimulate growth). <sup>7/7</sup> Required by "dependent" mutants. <sup>8/8</sup> Required as growth factor and not replaceable by aspartic or glutamic acids. <sup>9/9</sup> Required by some colorless Phytoflagellata. <sup>10/10</sup> Required although another carbon source is available, particularly in darkness. Functions in metabolism of C<sub>3</sub> and C<sub>4</sub> compounds. <sup>11/11</sup> Required in higher-than-atmospheric concentrations by some species. <sup>12/12</sup> α-Hydroxy-γ-aminobutyric acid. <sup>13/13</sup> Required by *Tenebrio molitor*; interchangeable with γ-amino-β-hydroxybutyric acid. <sup>14/14</sup> Fe-containing molecule of unknown structure; not a heme compound. <sup>15/15</sup> Favors larval growth of *Drosophila* and *Aedes aegypti*. <sup>16/16</sup> Required by *Neisseria gonorrhoeae*. <sup>17/17</sup> Possibly required in vitro by *Plasmodium*; not required by *Tetrahymena*. <sup>18/18</sup> And related auxins (plant hormones). <sup>19/19</sup> Ineffective for *Astasia* (colorless counterpart of *Euglena*); stimulates growth of *Euglena gracilis* (green Phytoflagellata). <sup>20/20</sup> Also hemin, protohemin, protoporphyrin, and several other porphyrins. <sup>21/21</sup> Acetate, citrate, fumarate, α-ketoglutarate, oxalacetate, succinate, cis-aconitate, isocitrate, malate, and oxalosuccinate. <sup>22/22</sup> Several intermediates are utilized for growth by the "acetate" flagellates; acetate is utilized by most species. There is wide variation among species with respect to utilization or availability of individual Krebs intermediates and related compounds, such as pyruvate.

continued

**48. NUTRIENTS: MISCELLANEOUS GROWTH FACTORS**

Nutrient	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>1</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>2</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
17 Mucin	H, s	H	H	H	H	r <sup>23</sup>	H	H	H
18 Mycobactin <sup>24</sup>	H	H	H	H	H	r <sup>26</sup>	H	H	H
19 Putrescine	H	H	H	H	H	r, u <sup>26</sup>	H	rm	H
20 Quinic acid <sup>27</sup>	H	u <sup>28</sup>	H	H	H	rm	H	rm	H
21 Shikimic acid <sup>27</sup>	H	u <sup>28</sup>	H	H	H	rm	H	rm	H
22 Spermidine <sup>29</sup>	H	H	H	H	H	r, u <sup>26</sup>	H	H	H
23 Strepogenin <sup>30</sup>	H, s?	H	H	H	H	r, rm, s	H	H, s?	H
24 Thioctic acid <sup>31</sup>	H	H	r <sup>32</sup>	H	H	r	H	H	H
25 Unidentified factors <sup>33</sup>	r	r	r	.....	....	r	.....	r	....
Reference	5,45	15,16, 22,39	2,6,8,11, 18,24,25, 31,33	6,11,18, 23,31,33	3,12,14, 29,31, 34,38,41	10,21,27, 35-37, 42-44	1,13,19,20, 28,43,44	1,7,9, 13,17, 20,30	4,26,32, 40

<sup>1</sup>/ Including the colorless Phytoflagellata. <sup>2</sup>/ Also Dinoflagellata and Chrysomonadina. <sup>3</sup>/ Required by *Corynebacterium diphtheriae*. <sup>4</sup>/ C47H75ON5. <sup>5</sup>/ Required by *Mycobacterium paratuberculosis*. <sup>6</sup>/ Interchangeable with spermidine in some species. <sup>7</sup>/ Probable precursors of aromatic amino acids. <sup>8</sup>/ Spares or replaces phenylalanine or tyrosine in *Blattella germanica*. <sup>9</sup>/ Also agmatine and spermine. <sup>10</sup>/ Also D-alanyl-histidine, amino-n-butyryl-L-histidine, carnosine, and various tyrosine peptides. <sup>11</sup>/ Protogen or α-lipoic acid. Exists in tissues as lipothiamide which catalyzes the oxidation of pyruvate and α-ketoglutarate. <sup>12</sup>/ Required by *Tetrahymena geleii* (8 strains), and *T. vorax* (2 strains); spared but not replaced by acetate; required (?) by *Pervanema tri-chophorum*. <sup>13</sup>/ Tissue extracts and unknown substances or complexes in living tissue or protoplasm.

**Contributors:** (a) Briggs, George M., (b) Cantino, Edward C., (c) Fogg, G. E., (d) Gordon, Harold Thomas, (e) House, Howard L., (f) Koser, Stewart A., (g) Loeffler, John B., (h) Pelletier, Réal L., (i) Shorb, Mary S., (j) Turrell, Franklin M., (k) Van Wagtenonk, W. J.

**References:** [1] Archibald, R. M., and F. Reiss. 1950. Ann. N. Y. Acad. Sci. 50:1388. [2] Bessey, O. A., H. J. Lowe, and L. L. Salomon. 1953. Ann. Rev. Biochem. 22:545. [3] Bold, H. C. 1942. Botan. Rev. 8:69. [4] Bonner, J., and A. W. Galston. 1952. Principles of plant physiology. W. H. Freeman, San Francisco. [5] Bourne, G. H., and G. W. Kidder, ed. 1953. Biochemistry and physiology of nutrition. Academic Press, New York. [6] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [7] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [8] Cowperthwaite, J., et al. 1953. Ann. N. Y. Acad. Sci. 56:972. [9] Davis, B. D. 1950. Nature 166:1120. [10] Davis, B. D. 1951. J. Biol. Chem. 191:315. [11] Doyle, W. L. 1943. Biol. Rev. Cambridge Phil. Soc. 18:119. [12] Fogg, G. E. 1953. The metabolism of algae. Methuen, London. [13] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [14] Ghosh, B. P., and R. H. Burris. 1950. Soil Sci. 70(3):187. [15] Gilmour, D. 1960. Biochemistry of insects. Academic Press, New York. [16] Gordon, H. T. Unpublished. Univ. California Dept. Entomology, Berkeley, 1953. [17] Gordon, M., et al. 1950. Proc. Natl. Acad. Sci. U.S. 36:427. [18] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [19] Hartelius, V. 1946. Compt. Rend. Trav. Lab. Carlsberg, Ser. Physiol., 24:185. [20] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [21] Herbst, E. J., and E. E. Snell. 1948. J. Biol. Chem. 176:989. [22] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [23] Hutner, S. H., L. Provasoli, and J. Filfus. 1953. Ann. N. Y. Acad. Sci. 56:852. [24] Kidder, G. W. 1947. Ibid. 49:99. [25] Kidder, G. W. 1951. Ann. Rev. Microbiol. 5:139. [26] Klosa, J. 1952. Naturwissenschaften 39:405. [27] Knight, B. C. J. G. 1945. Vitamins Hormones 3:108. [28] Krebs, H. A. 1943. Ann. Rev. Biochem. 12:529. [29] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [30] Lilly, V. G., and H. L. Barnett. 1951. Physiology of the fungi. McGraw-Hill, New York. [31] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [32] Miller, E. C. 1938. Plant physiology. Ed. 2. McGraw-Hill, New York. [33] Miner, R. W. 1953. Ann. N. Y. Acad. Sci. 56:815. [34] Myers, J. 1951. Ann. Rev. Microbiol. 5:157. [35] Peterson, W. H. 1941. Biol. Symp. 5:31. [36] Peterson, W. H., and M. S. Peterson. 1945. Bacteriol. Rev. 9:49. [37] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley,

*continued*

## 48. NUTRIENTS: MISCELLANEOUS GROWTH FACTORS

New York. [38] Sager, R., and S. Granick. 1953. *Ann. N. Y. Acad. Sci.* 56:831. [39] Schultz, J., et al. 1946. *Anat. Record* 96:540. [40] Skoog, F., ed. 1951. *Plant growth substances*. Univ. Wisconsin Press, Madison. [41] Smith, G. M. 1951. *Manual of phycology*. Chronica Botanica, Waltham, Mass. [42] Snell, E. E. 1949. *Ann. Rev. Microbiol.* 3:97. [43] Stephenson, M. 1949. *Bacterial metabolism*. Longmans, Green; New York. [44] Werkman, C. H., and P. W. Wilson, ed. 1951. *Bacterial physiology*. Academic Press, New York. [45] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York.

## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part I. CARBON SOURCES

*Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required;  $\bar{R}$  = not required; U and u = utilized;  $\bar{U}$  = not utilized; \* = may serve as sole or partial carbon source; \*\* = simplest carbon source.

Carbon Source (A)	Verte- brata (B)	Invertebrata		Phyto- flagellata (green) <sup>a</sup> (E)	Algae (F)	Bacterio- phyta (G)	Fungi		Sperma- tophyta (J)
		Insecta (C)	Protozoa <sup>1</sup> (D)				Saccharo- mycetaceae (H)	Other (I)	
<b>Inorganic Carbon</b>									
1 Carbon, amorphous, C	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	u**?	$\bar{U}$	u?	$\bar{R}$
2 Carbon monoxide, CO	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$ ?	u <sup>3</sup>	$\bar{U}$	$\bar{U}$	$\bar{U}$
3 Carbon dioxide, CO <sub>2</sub>	U	U?	r, u* <sup>4</sup>	r, U**	R, U**	r, u**	...	r, U	R, U**
4 Bicarbonates, -HCO <sub>3</sub> <sup>-</sup>	U	U?	r, u* <sup>4</sup>	r, U**	U**?	r, u**	...	U	R, U**
5 Cyanogens, -CN <sup>-</sup>	....	....	....	....	....	u	...	u	?
<b>Organic Carbon</b>									
6 Alkanes, alkenes	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	u <sup>5</sup>	$\bar{U}$	u	$\bar{U}$
7 Alcohols	U	u	u	$\bar{R}$ , u	$\bar{R}$ , u	u	u	u*	u
8 Aldehydes	$\bar{R}$ , u	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}$	u	u	u	$\bar{R}$
9 Ketones	$\bar{R}$ , u	$\bar{R}$	u	$\bar{R}$	$\bar{R}$	u	...	$\bar{U}$	$\bar{R}$
10 Acids	r, U	r, U	u	$\bar{R}$ , u	$\bar{R}$ , u	u	u	u*	$\bar{R}$
11 Carbohydrates	U	r, U	r, u <sup>6</sup>	$\bar{R}$ , u	$\bar{R}$ , u	U*	U*	U*	U <sup>7</sup>
12 Glycosides	$\bar{R}$ , u	$\bar{R}$	$\bar{R}$	$\bar{R}$ , u	$\bar{R}$ , u	u	u	u	$\bar{R}$
13 Fats	U	r, U <sup>8</sup>	u	$\bar{R}$ , $\bar{U}$	$\bar{R}$ , $\bar{U}$	u	u	u	$\bar{R}$ , $\bar{U}$
14 Waxes	u	u	$\bar{R}$ , $\bar{U}$	$\bar{R}$ , $\bar{U}$	$\bar{R}$ , $\bar{U}$	u	...	u*	$\bar{R}$ , $\bar{U}$
15 Amino acids, peptides, proteins	R, U	u	u	$\bar{R}$ , u	$\bar{R}$ , u	u	u	u*	$\bar{R}$ , u
Reference	13,17, 19,25	5,9,26	2,6,12,16	2,6,12, 16,21	10,21	1,18,22, 24	1,4,7,8,11, 18,22,24	3,4,7, 11	14,15,20, 23

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>2/2</sup> Also Dinoflagellata and Chrysomonadina. <sup>3/3</sup> Utilized by *Carboxydomonas oligocarbophila* and *Sarcina barkeri*. <sup>4/4</sup> Utilized by *Tetrahymena geleii* (condensation with pyruvate, oxalacetate, succinate). Required by *Chilomonas paramecium* and *Astasia longa*, but not on acetate medium. Required by *Polytomella caeca* on media containing casein hydrolysate, and also on acetate medium. <sup>5/5</sup> Utilized by *Sarcina methanica*: ethane, propane, butane, hexane, propylene, butylene, and paraffin oils. Utilized by other species: gasoline, kerosene, mineral oils, paraffin wax, etc. <sup>6/6</sup> Carbohydrate carbon source not required by *Tetrahymena geleii* except when utilizing NH<sub>4</sub><sup>+</sup> in a medium low in amino acids. <sup>7/7</sup> Utilized by isolated tissues (e.g., roots, callus and tumor tissues, green plants in aseptic culture). <sup>8/8</sup> Insects do not require dietary fats other than specific lipids as growth substances and vitamins.

*Contributors:* (a) Cantino, Edward C., (b) Fogg, G. E., (c) Gordon, Harold Thomas, (d) House, Howard L., (e) Loefer, John B., (f) Pelletier, Réal L., (g) Rusoff, Louis Leon, (h) Turrell, Franklin M., (i) Van Wagtenonk, W. J.

*References:* [1] Buchanan, R. E., and E. I. Fulmer. 1928-30. *Physiology and biochemistry of bacteria*. Williams and Wilkins, Baltimore. [2] Calkins, G. N., and F. M. Summers, ed. 1941. *Protozoa in biological research*. Columbia Univ. Press, New York. [3] Cochrane, V. W. 1958. *Physiology of fungi*. J. Wiley, New York.

*continued*



## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part I. CARBON SOURCES

- [4] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [5] Gilmour, D. 1960. Biochemistry of insects. Academic Press, New York. [6] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [7] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [8] Henrici, A. T. 1947. Molds, yeasts, and actinomycetes. Ed. 2. J. Wiley, New York. [9] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [10] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [11] Lilly, V. G., and H. L. Barnett. 1951. Physiology of the fungi. McGraw-Hill, New York. [12] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [13] Maynard, L. A., and J. K. Loosli. 1962. Animal nutrition. Ed. 5. McGraw-Hill, New York. [14] Meyer, B. S., and D. B. Anderson. 1952. Plant physiology. Ed. 2. Van Nostrand, New York. [15] Miller, E. C. 1938. Plant physiology. Ed. 2. McGraw-Hill, New York. [16] Miner, R. W. 1953. Ann. N. Y. Acad. Sci. 56:815. [17] Peterson, W. H., and F. M. Strong. 1953. General biochemistry. Prentice-Hall, New York. [18] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [19] Sherman, H. C. 1952. Chemistry of food and nutrition. Ed. 8. Macmillan, New York. [20] Skoog, F., ed. 1951. Plant growth substances. Univ. Wisconsin Press, Madison. [21] Smith, G. M. 1951. Manual of phycology. Chronica Botanica, Waltham, Mass. [22] Stephenson, M. 1949. Bacterial metabolism. Ed. 3. Longmans, Green; New York. [23] Steward, F. C., ed. 1963. Plant physiology. Academic Press, New York. v. 3. [24] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [25] West, E. S., and W. R. Todd. 1961. Textbook of biochemistry. Ed. 3. Macmillan, New York. [26] Wigglesworth, V. B. 1961. The principles of insect physiology. Ed. 4. Methuen, London.

### Part II. NITROGEN SOURCES

Nitrogen is a universal requirement of living organisms. Although some utilize it as molecular nitrogen, most organisms require it in the form of compounds. *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required;  $\bar{R}$  = not required; rm = required by one or more mutants; U and u = utilized;  $\bar{U}$  and  $\bar{u}$  = not utilized; s = stimulates growth or other processes; \* = serves as adequate or partial nitrogen source; \*\* = simplest adequate nitrogen source.

Nitrogen Source	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>1</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Inorganic Nitrogen									
1 Nitrogen, molecular, N <sub>2</sub>	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{U}$	$\bar{R}, u^{**5}$	$u^{**4}$	$\bar{R}$	$\bar{R}, u^{?6}$	$u^5$
2 Ammonia, NH <sub>3</sub> ; ammonium, -NH <sub>4</sub> <sup>+</sup>	$\bar{R}, u^7$	$\bar{R}, u^7$	r, u <sup>**</sup>	$\bar{R}, U^{**}$	$\bar{R}, U^{**9}$	$\bar{R}, u^{*9}$	$\bar{R}, u^{**}$	rm, u <sup>**10c</sup>	$\bar{R}, U^{**}$
3 Hyponitrite, -HN <sub>2</sub> O <sub>2</sub> <sup>-</sup> or -N <sub>2</sub> O <sub>2</sub> <sup>=</sup>	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}, u^{11}$	.....	$\bar{R}, \bar{u}^{12}$	$\bar{R}, \bar{u}^{13}$
4 Nitrite, -NO <sub>2</sub> <sup>-</sup>	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}$	$\bar{R}, u^{*14}$	$\bar{R}, u^*$	$\bar{R}, u^*$	rm, u <sup>**15</sup>	$\bar{R}, u^{*16}$
5 Nitrate, -NO <sub>3</sub> <sup>-</sup>	$\bar{R}$	$\bar{R}$	$\bar{R}, u$	$\bar{R}, u^{*17}$	$\bar{R}, U^{*17}$	$\bar{R}, u^{*18}$	$\bar{R}, u^{*19}$	rm, u <sup>**20</sup>	$\bar{R}, U^*$

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>1/2</sup> Also Dinoflagellata and Chrysomonadina. <sup>1/3</sup> Utilized in nitrogen fixation by blue-green algae (Nostocaceae). Not utilized if hydrogen or carbon monoxide are present. <sup>1/4</sup> Utilized by nitrogen-fixing bacteria. <sup>1/5</sup> Evidence is conflicting for nitrogen fixation. <sup>1/6</sup> Utilized via symbiotic bacteria, as in root nodules of legumes. <sup>1/7</sup> Ruminants, and possibly other vertebrates, utilize dietary NH<sub>4</sub><sup>+</sup>. NH<sub>4</sub><sup>+</sup> originating as a metabolic intermediate is utilized in amino acid synthesis. <sup>1/8</sup> Utilized by *Chlorella* in preference to NO<sub>3</sub><sup>-</sup>. <sup>1/9</sup> Many bacteria require specific amino acids from the environment, but as a class they utilize NH<sub>4</sub><sup>+</sup> for synthesis of other amino acids. <sup>1/10</sup> Utilized by *Phycomyces blakesleeanus* (Mucorales). Probably utilized by all fungi. <sup>1/11</sup> Utilized by *Clostridium pasteurianum* but not for growth; not utilized for denitrification by *Pseudomonas stutzeri*. <sup>1/12</sup> Not utilized by *Aspergillus niger*. <sup>1/13</sup> Not utilized by *Nicotiana*. <sup>1/14</sup> Utilized poorly by *Chlorella pyrenoidosa*. <sup>1/15</sup> Utilized as sole nitrogen source by many fungi. <sup>1/16</sup> Toxic to many plants; poorly utilized by *Nicotiana*. <sup>1/17</sup> Utilized in light. <sup>1/18</sup> Not utilized by purple photosynthetic bacteria. <sup>1/19</sup> Poorly utilized by most yeasts. <sup>1/20</sup> Acts as sole hydrogen acceptor in anaerobic metabolism of *Aspergillus niger*. Required by some species when manitol is the carbon source. Some species require NO<sub>3</sub><sup>-</sup>.

continued



## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part II. NITROGEN SOURCES

Nitrogen Source (A)	Verte- brata (B)	Invertebrata		Phyto- flagellata (green) <sup>1</sup> (E)	Algae (F)	Bacterio- phyta (G)	Fungi		Sperma- tophyta (J)
		Insecta (C)	Protozoa <sup>1</sup> (D)				Saccharo- mycetaceae (H)	Other (I)	
<b>Inorganic Nitrogen</b>									
6 Nitrohydroxamate, -HN <sub>2</sub> O <sub>3</sub> <sup>-</sup>	R	R	R	R	R	R	R	R, u* <sup>21</sup>	R, u* <sup>22</sup>
7 Cyanide, -CN <sup>-</sup>	U <sup>23</sup>	U <sup>23</sup>	U <sup>23</sup>	R	R	R	R	R, u* <sup>24</sup>	U
8 Thiocyanate, -CNS <sup>-</sup>	U	U	U	R	R	R, u*	R, u	R, u?	U
9 Cyanamide, -NHCM <sup>-</sup>	U	U	U	R	R	R, u* <sup>25</sup>	R	u* <sup>25</sup>	U
<b>Organic Nitrogen</b>									
10 Oximino compounds <sup>26</sup> , RONH <sub>2</sub>	R	R	R	R	U <sup>27</sup>	R, u* <sup>28</sup>	R, u?	R, u*	R, U <sup>29</sup>
11 Amines <sup>30</sup> , RNH <sub>2</sub>	R	R	R	R	R, u* <sup>31</sup>	R, u*	R, u*	R, u*	R
12 Acid imides, (RCO) <sub>2</sub> NH	R	R	R	R	R	R, u*	R	R	R
13 Acid amides, RCONH <sub>2</sub>	R	R	R	R	R, u*	R, u*	r, u*	R, u* <sup>32</sup>	R, u* <sup>33</sup>
14 Urea, (NH <sub>2</sub> ) <sub>2</sub> CO	R, u <sup>34</sup>	R	R, u* <sup>35</sup>	R, u*	R, u*	R, u*	R, u*	R, u*	R, U*
15 Amino acids, RCH(NH <sub>2</sub> )COOH	R, U** <sup>7</sup>	R, U**	r, u*	r, u*	R, u* <sup>33</sup>	r <sup>3</sup> , U*, s	r, U*	r, rm, U*	R, u*
16 Peptides, polypeptides <sup>37</sup>	R, U*	R	r, u*	R, u*	R, u*	r, u*, s	R, u*	R, u*	R, u*
17 Proteins (per se)	R <sup>39</sup> , U*, s	R, U*	r, u*	R	R	R, u*	R	R, u*	R
18 Imidazole compounds	R	R	R	R	R	R, u*	R, u*	R, u*	R
19 Pyridine compounds	r	r	r	R	R	r, u*	r	R	R
20 Pyrimidine compounds	R	r	r, u*	r	r	R, u*, s	R, u*	R, u*, s	R
21 Purine compounds	R	R	r, u*	R	R	R, u*, s	R, u	rm, u*, s	R, u
22 Indole compounds	R	R	R	R, s	R, s	R, u*, s	R	rm, u*	R, s
Reference	1,3,13, 17,18, 50	12,20, 48	8,14-16, 23,26,27, 32,42	8,14,26, 27,32,34, 40,42	4,9,11, 21,22,25, 29,33,34, 37,38,44, 45	24,39,46, 49	2,5	7,10,19, 28,41	6,30,31, 35,36,43, 47,49

<sup>1/1</sup> Including the colorless Phytoflagellata. <sup>1/2</sup> Also Dinoflagellata and Chrysomonadina. <sup>1/3</sup> Ruminants, and possibly other vertebrates, utilize dietary NH<sub>4</sub><sup>+</sup>. NH<sub>4</sub><sup>+</sup> originating as a metabolic intermediate is utilized in amino acid synthesis. <sup>1/4</sup> Many bacteria require specific amino acids from the environment, but as a class they utilize NH<sub>4</sub><sup>+</sup> for synthesis of other amino acids. <sup>1/5</sup> Good nitrogen source for *Aspergillus niger*. <sup>1/6</sup> Good nitrogen source for *Nicotiana*. <sup>1/23</sup> Toxic. <sup>1/24</sup> Utilized by *Aspergillus niger* when nitrogen-starved. <sup>1/25</sup> Many species utilize cyanamide and derivatives. <sup>1/26</sup> Including hydroxylamine. <sup>1/27</sup> Hydroxylamine is toxic. <sup>1/28</sup> *Clostridium perfringens* utilizes hydroxylamine; *Nitrosomonas* utilizes a nontoxic concentration. <sup>1/29</sup> Hydroxylamine is poor nitrogen source. <sup>1/30</sup> Alkylamines, as methylamine (CH<sub>3</sub>NH<sub>2</sub>), and arylamines, as benzylamine (C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>NH<sub>2</sub>). <sup>1/31</sup> All organisms studied utilize glucosamine. <sup>1/32</sup> *Aspergillus* utilizes formamide and other acid amides; utilizes both amino and amide nitrogen of asparagine. <sup>1/33</sup> Some species utilize acetamide. <sup>1/34</sup> Utilized by ruminants via rumen microflora. <sup>1/35</sup> Utilized by *Astasia longa*. <sup>1/36</sup> L-Arginine, glutamine, cysteine and L-asparagine support more rapid growth of *Chlorella* than does NH<sub>4</sub><sup>+</sup>. <sup>1/37</sup> Including peptones, synthetic di- and tri-peptides, strepogenin, and glutathione. <sup>1/38</sup> Proteins, per se, are listed as "not required" for vertebrates, since only certain amino acids are required.

**Contributors:** (a) Cantino, Edward C., (b) Fogg, G. E., (c) Gordon, Harold Thomas, (d) House, Howard L., (e) Koser, Stewart A., (f) Loefer, John B., (g) Pelletier, Réal L., (h) Purvis, E. R., (i) Shorb, Mary S., (j) Turrell, Franklin M., (k) Van Wagtenonk, W. J.

**References:** [1] Anderson, R. J. 1949. Essentials of physiological chemistry. J. Wiley, New York. [2] Archibald, A. M., and F. Reiss. 1950. Ann. N. Y. Acad. Sci. 50:1388. [3] Bourne, G. H., and G. W. Kidder, ed. 1953. Biochemistry and physiology of nutrition. Academic Press, New York. [4] Burlaw, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [5] Chester, K. S. 1946. The nature and prevention of the cereal rusts as exemplified in the leaf rust of wheat. Chronica Botanica, Waltham, Mass. [6] Chibnall, A. C. 1939. Protein metabolism in the plant. Yale Univ. Press, New Haven. [7] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [8] Doyle, W. L. 1943. Biol. Rev. Cambridge Phil. Soc. 18:119. [9] Fogg, G. E. 1953. The metabolism of algae. Methuen, London. [10] Foster, J. W. 1949. Chemical activities of fungi. Academic Press,

continued

## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part II. NITROGEN SOURCES

New York. [11] Ghosh, B. P., and R. H. Burris. 1950. *Soil Sci.* 70:187. [12] Gilmour, D. 1960. *Biochemistry of insects*. Academic Press, New York. [13] Gortner, R. A., and W. A. Gortner, ed. 1949. *Outlines of biochemistry*. Ed. 3. J. Wiley, New York. [14] Hall, R. P. 1953. *Protozoology*. Prentice-Hall, New York. [15] Hanson, R. W. 1952. Ph.D. Thesis. Univ. California, Los Angeles. [16] Hanson, R. W., and T. L. Jahn. 1953. *Federation Proc.* 12:61 (Abstr. 195). [17] Harrow, B. 1962. *Textbook of biochemistry*. Ed. 8. W. B. Saunders, Philadelphia. [18] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1954. *Practical physiological chemistry*. Ed. 13. Blakiston, New York. [19] Hawker, L. E. 1950. *Physiology of fungi*. Univ. London Press, London. [20] House, H. L. 1962. *Ann. Rev. Biochem.* 31:653. [21] Ketchum, B. H. 1939. *Am. J. Botany* 26:399. [22] Ketchum, B. H., and A. C. Redfield. 1949. *J. Cellular Comp. Physiol.* 33:281. [23] Kidder, G. W. 1951. *Ann. Rev. Microbiol.* 5:139. [24] Knight, B. C. J. G. 1936. *Brit. Med. Res. Council Spec. Rept. Ser.* 210. [25] Lewin, R. A., ed. 1962. *Physiology and biochemistry of algae*. Academic Press, New York. [26] Lwoff, A. 1944. *L'évolution physiologique*. Hermann, Paris. [27] Lwoff, A., ed. 1951-55. *Biochemistry and physiology of protozoa*. Academic Press, New York. [28] Lilly, V. G., and H. L. Barnett. 1951. *Physiology of the fungi*. McGraw-Hill, New York. [29] Ludwig, C. A. 1938. *Am. J. Botany* 25:448. [30] Meyer, B. S., and D. B. Anderson. 1952. *Plant physiology*. Ed. 2. Van Nostrand, New York. [31] Miller, E. C. 1938. *Plant physiology*. Ed. 2. McGraw-Hill, New York. [32] Miner, R. W. 1953. *Ann. N. Y. Acad. Sci.* 56:815. [33] Myers, J. 1944. *Plant Physiol.* 19:579. [34] Myers, J. 1951. *Ann. Rev. Microbiol.* 5:157. [35] Nightingale, G. T. 1937. *Botan. Rev.* 3:85. [36] Nightingale, G. T. 1948. *Ibid.* 14:185. [37] Norris, L. C. Unpublished. Cornell Univ. Dept. Poultry Husbandry, Ithaca, 1953. [38] Pirson, A., and G. Wilhelmi. 1950. *Z. Naturforsch.* 5b:211. [39] Porter, J. R. 1946. *Bacterial chemistry and physiology*. J. Wiley, New York. [40] Reinhardt, K. 1950. *Arch. Mikrobiol.* 15:270. [41] Robbins, W. J. 1937. *Am. J. Botany* 24:243. [42] Scheer, B. T. 1948. *Comparative physiology*. J. Wiley, New York. [43] Skoog, F., ed. 1951. *Plant growth substances*. Univ. Wisconsin Press, Madison. [44] Smith, G. M. 1951. *Manual of phycology*. *Chronica Botanica*, Waltham, Mass. [45] Spohr, H. A., and H. W. Milner. 1949. *Plant Physiol.* 24:120. [46] Stephenson, M. 1949. *Bacterial metabolism*. Ed. 3. Longmans, Green; New York. [47] Steward, F. C., ed. 1963. *Plant physiology*. Academic Press, New York. v. 3. [48] Trager, W. 1941. *Physiol. Rev.* 21:1. [49] Werkman, C. H., and P. W. Wilson, ed. 1951. *Bacterial physiology*. Academic Press, New York. [50] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York.

### Part III. SULFUR SOURCES

Sulfur occurs in several amino acids: cysteine, cystine, and methionine. Methionine is required by all animal forms studied and may be essential in the protein structure of all living organisms. Sulfur occurs also in chondroitin sulfuric acid, a component of vertebrate connective tissue, and in sulfolipids found in the white matter of the brain and in other tissues. *Abbreviations:* Capital letters indicate data are pertinent to all organisms studied; lower case letters and symbols indicate data apply to one or more species, or strains, but not to all forms studied. R and r = required; R̄ = not required; rm = required by one or more mutants; U and u = utilized; Ū = not utilized.

Sulfur Source	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Inorganic Sulfur									
1 Sulfur <sup>3</sup> (elemental), S	R, u <sup>4</sup>	R̄	R̄	R̄	R̄, Ū	r <sup>5</sup>	u	u <sup>6</sup>	R̄, Ū
2 Sulfhydryl, -SH <sup>-</sup>	R̄	R̄	R̄	R̄	R̄	...	...	u	R̄

<sup>1/</sup> Including the colorless Phytoflagellata. <sup>2/</sup> Also Dinoflagellata and Chrysoomonadina. <sup>3/</sup> The substance alone, not in combination with other elements. <sup>4/</sup> In ruminants, bacteria build the element into amino acids (methionine, cystine). <sup>5/</sup> Utilized by *Thiobacillus thiooxidans*, *Desulfovibrio desulfuricans*, and *D. aestuarii* (Thiorhodaceae). <sup>6/</sup> Utilized by *Fusarium lini*.

*continued*

## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part III. SULFUR SOURCES

Sulfur Source	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>1</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Inorganic Sulfur									
3 Sulfide, -S <sup>=</sup>	R, U	R	R	R	R, u <sup>7</sup>	u <sup>9</sup>	...	u <sup>9</sup>	R
4 Bisulfite, -HSO <sub>3</sub> <sup>-</sup>	R, U	R	R	R	R	...	...	u <sup>10</sup>	R
5 Sulfite, -SO <sub>3</sub> <sup>=</sup>	R, U	R	R	R	R	u <sup>11</sup>	...	u <sup>10</sup>	R, u
6 Sulfate, -SO <sub>4</sub> <sup>=</sup>	R, u <sup>12</sup>	R	r	U	U	r	u	u <sup>10</sup>	U
7 Thiosulfate, -S <sub>2</sub> O <sub>3</sub> <sup>=</sup>	R, U	R	R	R	R	u <sup>13</sup>	...	rm, u	R, u
8 -S <sub>2</sub> O <sub>8</sub> <sup>=14</sup>	R	R	R	R	R	...	...	u?	R
9 Tetrathionate, -S <sub>4</sub> O <sub>6</sub> <sup>=</sup>	R	R	R	R	R	u	...	....	R
10 Sulfoxylate, -SOOH <sup>-</sup>	R	R	R	R	R	u	...	u <sup>15</sup>	R
11 -SO as sulfur hydrate, H <sub>2</sub> SO	R	R	R	R	R	u	...	....	R
12 Thiocyanate, -SCN <sup>-</sup>	R	R	R	R	R	u <sup>16</sup>	...	u	R
13 Persulfate	R	R	R	R	R	...	...	u <sup>10</sup>	R
Organic Sulfur <sup>17</sup>									
14 Cystathionine	u	u	R, u	R	R	u	...	u	R
15 Cysteine	U	u <sup>18</sup>	u	R, u	R, u <sup>19</sup>	r	u	u <sup>9</sup>	R, u
16 Cystine	U	u	u	R, u	R	r	u	u <sup>2</sup>	R, u
17 Homocysteine	u	u	R, u	R	R	u	...	U	R
18 Homocystine	u	u	R, u	R	R	u	...	U	R
19 Methionine	R	r <sup>20</sup>	r	r	R, U	r	u	U	R, u
20 Peptones	R, U	u	r	u	R, u	u	u	U	R, u
21 Biotin <sup>21</sup>	R?, U	r	r?	r	R	u	u	u	R
22 Coenzyme A	R, U	R	R	R	R	R	R	R	R
23 Glutathione ("G-SH") <sup>22</sup>	R, U	u	R, u	u	R	r	u	u	R
24 Thiamine <sup>23</sup>	R	r	r	r	u	r	r	U	R
25 Thiazole <sup>23</sup>	R	R	r <sup>24</sup> , u	r <sup>24</sup>	r	r <sup>24</sup>	r <sup>24</sup>	u? <sup>25</sup>	R
26 Thiocetic acid <sup>26</sup>	R	R	r <sup>27</sup>	R	R	r <sup>28</sup>	R	R	R
Miscellaneous									
27 Alkylsulfides, R-S-S-R	R	R	R	R	R	...	...	u <sup>29</sup>	R
28 Alkylsulfates, R-SO <sub>4</sub> -R	R	R	R	R	R	...	...	u	R
29 Alkylsulfonates, R-SO <sub>3</sub> -R	R	R	R	R	R	...	...	u	R
30 Dithionate	R	R	R	R	R	...	...	u <sup>30</sup>	R
31 Etheral sulfates	R	R	R	R	R	...	...	U	R
32 Sulfamate, -SO <sub>3</sub> -NH <sub>2</sub> <sup>=</sup>	R	R	R	R	R	...	...	u	R
33 Sulfonic acid amides	R	R	R	R	R	U	...	rm	R
34 Sulfoxides, R <sub>2</sub> SO	R	R	R	R	R	u	...	u	R
35 Taurine	R	R	R	R	R	U	...	u	R
36 Thioacetamide	R	R	R	R	R	u	...	u	R

/1/ Including the colorless Phytoflagellata. /2/ Also Dinoflagellata and Chrysomonadina. /7/ *Synechococcus*, grown in an atmosphere of nitrogen, utilizes Na<sub>2</sub>S (with reduction of CO<sub>2</sub>); *Oscillatoria* & *Pinnularia* reduce CO<sub>2</sub> with H<sub>2</sub>S, depositing sulfur in their cells; *Scenedesmus* also utilizes sulfide. H<sub>2</sub>S toxic to *Chlorella*. /8/ Utilized (e.g., by *Beggiatoa*, *Thiothrix*, *Thioploca*, *Thiobacillus*). /9/ Utilized by some aquatic fungi (e.g., members of Blastocladales and Saprolegniales which cannot utilize oxidized sulfur but require a reduced sulfur source: H<sub>2</sub>S, cysteine, cystine, methionine, thioacetate, thiocarbonate, thioglycolate, thiourea). /10/ Utilized by *Brevilegnea gracilis*; not utilized by many other Saprolegniaceae. /11/ Utilized by *Desulfovibrio desulfuricans* and *D. aestuarii*. /12/ Utilized for formation of chondroitin sulfate and heparin; utilized by laying hens via conversion to cystine. /13/ Utilized by *Thiobacillus novellus*, *Pseudomonas aeruginosa*, *P. fluorescens*, and others. /14/ Decomposes on contact with water. /15/ Inorganic sulfur, less oxidized than sulfinate, is not efficiently utilized by *Aspergillus niger*. /16/ *Thiobacillus thiooxidans* can utilize NH<sub>4</sub>SCN as the sole source of carbon, nitrogen and sulfur. /17/ Compound may be a sulfur source, or may be required for its molecular structure; not synthesized by the organism. /18/ Also utilize cysteic acid and isethionic acid. /19/ Utilized as nitrogen source (and sulfur source?) by *Chlorella pyrenoidosa*. /20/ Also utilize methionine sulfoxide, taurine. /21/ Numerous fungi, yeasts, bacteria, and most of the vertebrates and invertebrates studied, require biotin. The replacement of sulfur in the biotin molecule does not affect the activity for some bacteria. /22/ Complex of cysteine, glycine, and glutamic acid. /23/ Thiamine, containing pyrimidine and thiazole (the latter an imidazole ring with one carbon atom replaced by sulfur) is required by numerous organisms; probably also a sulfur source. /24/ Satisfies thiamine requirement for some (see Fn. 23); probably a sulfur source. /25/ Not utilized as a sulfur source by *Aspergillus niger* (cannot rupture the thiazole ring?). /26/ Protogen, or α-lipoic acid. /27/ Required by *Tetrahymena geleii* (8 strains), and *T. vorax* (2 strains). /28/ Required by *Streptococcus faecalis* for oxidation of pyruvate. /29/ Utilized by *Penicillium brevicaulis* and *Schizophyllum commune*. /30/ Not utilized by Saprolegniaceae.

continued



**49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR**

**Part III. SULFUR SOURCES**

Sulfur Source	Vertebrata	Invertebrata		Phytoflagellata (green) <sup>2</sup>	Algae	Bacteriophyta	Fungi		Spermatophyta
		Insecta	Protozoa <sup>2</sup>				Saccharomycetaceae	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Miscellaneous									
37 Thioacetate	⊘	⊘	⊘	⊘	⊘	u? <sup>31</sup>	...	r, u <sup>9</sup>	⊘
38 Thiocarbonate	⊘	⊘	⊘	⊘	⊘	...	...	r, u <sup>9</sup>	⊘
39 Thioglycolate	⊘	⊘	⊘	⊘	⊘	u? <sup>32</sup>	...	r, u <sup>9</sup>	⊘
40 Thiols, R-SH	⊘	⊘	⊘	⊘	⊘	...	...	u	⊘
41 Thiooxalate	⊘	⊘	⊘	⊘	⊘	...	...	u	⊘
42 Thiourea	⊘	⊘	⊘	⊘	⊘	u? <sup>33</sup>	u <sup>34</sup>	u <sup>9</sup>	⊘
Reference	3,26, 29,31, 41-44, 58	25,35, 59	12,27,33, 40	12,27,33, 40	11,22, 37,46, 47,51	8,10,21, 49,52,56, 57	50,54	13,14, 23,32, 38,39	1,2,4-7,9, 15-20,24,28, 30,34,36,45, 48,53,55,60

<sup>1/</sup> Including the colorless Phytoflagellata. <sup>2/</sup> Also Dinoflagellata and Chrysoomonadina. <sup>3/</sup> Utilized by some aquatic fungi (e.g., members of Blastocladales and Saprolegniales which cannot utilize oxidized sulfur but require a reduced sulfur source: H<sub>2</sub>S, cysteine, cystine, methionine, thioacetate, thiocarbonate, thioglycolate, thiourea). <sup>4/</sup> Not utilized as carbon source by many species or strains; improbable sulfur source. <sup>5/</sup> Surface active in culture media for many fastidious forms. Powerful reducing agent. Not utilized as a carbon source, and improbable as a sulfur source. <sup>6/</sup> Utilized as a nitrogen source by many bacteria; probable sulfur source (?). <sup>7/</sup> Utilized by *Torula monosa* and *T. dattila*.

**Contributors:** (a) Cantino, Edward C., (b) Fogg, G. E., (c) Gordon, Harold Thomas, (d) Loefer, John B., (e) Pelletier, Réal L., (f) Turrell, Franklin M., (g) Van Wagtenonk, W. J.

**References:** [1] Alway, F. J. 1940. J. Am. Soc. Agron. 32:913. [2] Ames, J. W., and G. E. Boltz. 1916. Ohio Agr. Expt. Sta. Res. Bull. 292:221. [3] Anderson, R. J. 1949. Essentials of physiological chemistry. J. Wiley, New York. [4] Balks, R. 1938. Forschungsdienst, Sonderh. 8:208. [5] Balks, R. 1939. Ernaehr. Pflanze 35:194. [6] Balks, R. 1942. Forschungsdienst, Sonderh. 16:217. [7] Barrien, B. S., and J. G. Wood. 1939. New Phytologist 38:257. [8] Bavendamm, W. 1924. Die farblosen und roten Schwefelbakterien des Süß- und Salzwassers. G. Fischer, Jena. [9] Bogdanov, S. 1903. Sel'skoe Khoz. Lyesov. 210:628. [10] Bunker, H. J. 1936. A review of the physiology and biochemistry of the sulfur bacteria. H. M. Stationery Office, London. [11] Burlew, J. S. 1953. Carnegie Inst. Wash. Publ. 600. [12] Calkins, G. N., and F. M. Summers, ed. 1941. Protozoa in biological research. Columbia Univ. Press, New York. [13] Cantino, E. C. 1950. Quart. Rev. Biol. 25:269. [14] Cochrane, V. W. 1958. Physiology of fungi. J. Wiley, New York. [15] Crocker, W. 1945. Soil Sci. 60:149. [16] Cultrera, R., and C. Vicini. 1941. Ann. Staz. Sper. Agrar. Modena 7:103. [17] Eaton, S. V. 1935. Botan. Gaz. 97:68. [18] Eaton, S. V. 1935. Trans. Illinois State Acad. Sci. 28:88. [19] Eaton, S. V. 1941. Botan. Gaz. 102:536. [20] Eaton, S. V. 1942. Ibid. 104:306. [21] Ellis, D. 1932. Sulfur bacteria. Longmans, Green; New York. [22] Fogg, G. E. 1953. The metabolism of algae. Methuen, London. [23] Foster, J. W. 1949. Chemical activities of fungi. Academic Press, New York. [24] Gilbert, F. A. 1951. Botan. Rev. 17:671. [25] Gilmour, D. 1960. Biochemistry of insects. Academic Press, New York. [26] Gortner, R. A., and W. A. Gortner, ed. 1949. Outlines of biochemistry. Ed. 3. J. Wiley, New York. [27] Hall, R. P. 1953. Protozoology. Prentice-Hall, New York. [28] Hambidge, G., ed. 1941. Hunger signs in crops. American Society of Agronomy and Natl. Fertilizer Association, Washington, D. C. [29] Harrow, B. 1962. Textbook of biochemistry. Ed. 8. W. B. Saunders, Philadelphia. [30] Hart, E. B., and W. E. Tottingham. 1915. J. Agr. Res. 5:233. [31] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1954. Practical physiological chemistry. Ed. 13. Blakiston, New York. [32] Hawker, L. E. 1950. Physiology of fungi. Univ. London Press, London. [33] Heilbrunn, L. V. 1952. An outline of general physiology. Ed. 3. W. B. Saunders, Philadelphia. [34] Heiserich, E. 1935. Z. Pflanzenernaehr. Dueng. Bodenk. 37:55. [35] House, H. L. 1962. Ann. Rev. Biochem. 31:653. [36] Krugel, C., C. Dreyspring, and F. Heinrich.

*continued*




## 49. NUTRIENTS: CARBON, NITROGEN, AND SULFUR

### Part III. SULFUR SOURCES

1938. Superphosphate 11:181. [37] Lewin, R. A., ed. 1962. Physiology and biochemistry of algae. Academic Press, New York. [38] Lilly, V. G., and H. L. Barnett. 1951. Physiology of the fungi. McGraw-Hill, New York. [39] Lin, C.-K. 1945. Am. J. Botany 32:296. [40] Lwoff, A., ed. 1951-55. Biochemistry and physiology of protozoa. Academic Press, New York. [41] Marston, H. R., and T. B. Robertson. 1928. Australia Council Sci. Ind. Res. Bull. 39:51. [42] Maurel, E. 1904. Compt. Rend. Soc. Biol. 56:796. [43] Maynard, L. A., and J. K. Loosli. 1962. Animal nutrition. Ed. 5. McGraw-Hill, New York. [44] Medvedeva, N. B. 1940. Akad. Nauk Ukr. SSR, p. 3. [45] Meyer, B. S., and D. B. Anderson. 1952. Plant physiology. Ed. 2. Van Nostrand, New York. [46] Myers, J. 1944. Plant Physiol. 19:579. [47] Myers, J. 1951. Ann. Rev. Microbiol. 5:157. [48] Olson, G. A., and J. L. St. John. 1921. Wash. State Univ. Agr. Expt. Sta. Bull. 165. [49] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [50] Schultz, A. S., and D. K. McManus. 1950. Arch. Biochem. 25:401. [51] Smith, G. M. 1951. Manual of phycology. Chronica Botanica, Waltham, Mass. [52] Stephenson, M. 1949. Bacterial metabolism. Ed. 3. Longmans, Green; New York. [53] Steward, F. C., ed. 1963. Plant physiology. Academic Press, New York. v. 3. [54] Sugata, H., and F. C. Koch. 1926. Plant Physiol. 1:337. [55] Valatx, A., and J. Dufrenoy. 1937. Compt. Rend. Congr. Chim. Ind., 17, Paris 1:494. [56] Van Niel, C. B. 1941. Advan. Enzymol. 1:263. [57] Werkman, C. H., and P. W. Wilson, ed. 1951. Bacterial physiology. Academic Press, New York. [58] West, E. S., and W. R. Todd. 1961. Textbook of biochemistry. Ed. 3. Macmillan, New York. [59] Wigglesworth, V. B. 1961. The principles of insect physiology. Ed. 4. Methuen, London. [60] Wood, J. G., and B. S. Barrien. 1939. New Phytologist 38:125.

## 50. PATHWAYS OF PROTEIN DIGESTION: MAN AND LABORATORY MAMMALS

Pepsin, trypsin, and chymotrypsin are endopeptidases, i.e., they hydrolyze peptide bonds in the interior of peptide chains as well as terminal bonds. Carboxypeptidase and leucine aminopeptidase are exopeptidases and can act only on terminal peptide bonds.




/1/ Pepsin hydrolyzes many types of peptide bonds but splits most rapidly those in which an aromatic amino acid provides the amino group. /2/ Trypsin hydrolyzes peptide bonds to which L-arginine, or L-lysine, contributes the carbonyl group. /3/ Chymotrypsin hydrolyzes many types of peptide bonds but splits most rapidly those in which an aromatic amino acid contributes the carbonyl group. /4/ Carboxypeptidase does not exhibit absolute specificity with respect to the terminal amino acid forming the bond being split; it acts most rapidly on those linkages in which aromatic amino acids are in the terminal position. The terminal amino acid must have a free carboxyl group. /5/ Amino acids and dipeptides enter the intestinal mucosal cells. Amino acids pass through unaltered--with a few exceptions, such as transamination of glutamic acid--and dipeptides are split to amino acids in the microvilli of the cell where the peptidases are localized. /6/ Only a few of the intestinal mucosal peptidases have been characterized. The best known is leucine aminopeptidase.

*Contributor:* Grossman, Morton I.

*Reference:* Fruton, J. S., and S. Simmonds. 1958. General biochemistry. Ed. 2. J. Wiley, New York.



## 52. PATHWAYS OF LIPID DIGESTION: MAN AND LABORATORY MAMMALS



<sup>1/1</sup> Pancreatic lipase acts preferentially on ester linkages at the terminal or 1 position of glycerol. Thus the major products of digestion are fatty acids and monoglycerides. <sup>2/2</sup> Bile salts, in their conjugated form, participate in at least three reactions during fat digestion and absorption: (i) as a cofactor for pancreatic lipase; (ii) to form micelles containing monoglyceride and fatty acid, as well as other lipids (these micelles are probably the form in which lipid is absorbed into the cell); (iii) as a cofactor for thiokinase in the intestinal mucosal cell. <sup>3/3</sup> Absorbed fatty acids go mainly into the triglycerides of chylomicrons, but small amounts are synthesized into cholesterol esters and phospholipids which also are constituents of chylomicrons. <sup>4/4</sup> Fatty acids with chain lengths shorter than ten carbon atoms are absorbed mainly into the portal blood, those with longer chain lengths mainly into the lymph.

*Contributor:* Grossman, Morton I.

*Reference:* Senior, J. R., and K. J. Isselbacher. 1962. J. Biol. Chem. 237:1454.

## 53. EXCRETION PRODUCTS: MAN

Values are based on "normal" dietary intake, including approximately 10 grams of nitrogen per day. In reducing values to mg/kg or µg/kg, a body weight of 70 kg was assumed, unless specific weight was reported in the literature. Values in parentheses are ranges, estimate "c" (cf. Introduction).

### Part I. URINE

Constituent	Amount Excreted per kg body wt per da	Reference	Constituent	Amount Excreted per kg body wt per da	Reference
(A)	(B)	(C)	(A)	(B)	(C)
General Chemical Components, mg			Amino acids		
1 Solids	860(780-1,000)	6,32,47,82	44 Citrulline	0.09(0-2.8)	103
2 Water	20,000 (7,000-42,000)	45	45 Cystine	1.7(1.0-2.6) <sup>2</sup>	85
Electrolytes, mg			46	1.3(0.6-1.9) <sup>3</sup>	75
3 Aluminum	0.0011 (0.0007-0.0016)	41	47 Glutamic acid	(3.7-5.0) <sup>2</sup>	9,33
4 Arsenic	0.00033(0-0.00130)	97	48	0.8(0-1.5) <sup>3</sup>	9,33
5 Bicarbonate	2.0(0.5-12.0)	27	49 Glycine	6.5 <sup>3</sup>	20
6 Bromine	(0.012-0.110)	12	50	2.2 <sup>3</sup>	9
7 Calcium	3.3(0.6-8.3)	32	51 Histidine	2.7(1.0-5.0) <sup>2</sup>	9,85,102
8 Chlorine	100(40-180)	32	52	2.0(1.2-2.7) <sup>3</sup>	9,102
9 Cobalt	0.00007 (0.00005-0.00012)	30	53 Hydroxyproline	0.02 <sup>2</sup>	9
10 Copper	0.0005 (0.0003-0.0007)	90	54 Isoleucine	0.2(0.1-0.3) <sup>2</sup>	20,75,102
11 Fluorine	0.022(0.007-0.100) <sup>1</sup>	32,51	55	0.08(0.04-0.20) <sup>3</sup>	20,75,102
12 Iodine	(0.0001-0.0070)	8	56 Leucine	0.30(0.22-0.45) <sup>2</sup>	20,75,102
13 Iron	0.007	21	57	0.13(0.05-0.17) <sup>3</sup>	20,75,102
14 Lead	0.00040 (0.00016-0.00110)	41	58 Lysine	0.80(0.48-1.70) <sup>2</sup>	20,75,102
15 Magnesium	1.35(0.42-2.40)	28	59	0.40(0.17-0.67) <sup>3</sup>	20,33,75
16 Manganese	(0.0001-0.0014)	41,42	60 Methionine	0.14(0.10-0.17) <sup>2</sup>	20,75,102
17 Mercury	(0.000007-0.000010)	81	61	0.05(0.03-0.10) <sup>3</sup>	20,75,102
18 Nickel	(0.002-0.004)	42	62 1-Methylhistidine	Trace <sup>4</sup>	83
19 Phosphorus			63 3-Methylhistidine	Trace <sup>4</sup>	83
20 Inorganic P	12(10-15)	93	64 Ornithine	0.15	9
21 Organic P	0.131(0.089-0.187)	65	65 Phenylalanine	0.30(0.21-0.54) <sup>2</sup>	9,20,102
22 Potassium	34(16-56)	32	66	0.17(0.09-0.23) <sup>3</sup>	9,20,102
23 Selenium	0.0005(0-0.0020)	80	67 Proline	0.61(0.30-0.90) <sup>2</sup>	102
24 Silicon	0.13(0.06-0.20)	5	68	0.12(0.03-0.20) <sup>3</sup>	102
25 Sodium	60(25-94)	32	69 Serine	0.6(0.5-0.7) <sup>2</sup>	9,94
26 Sulfur			70	0.3(0.2-0.5) <sup>3</sup>	9,94
27 Total S	16.0(5.1-20.6)	23	71 Taurine	(0.11-0.20)	103
28 Inorganic S	11.1(3.5-17.5)	23	72 Threonine	0.50(0.36-2.60) <sup>2</sup>	9,20,75,94,102
29 Ethereal S	0.95(0.56-1.40)	23	73	0.25(0.11-0.35) <sup>3</sup>	9,20,75,94,102
30 Neutral S	1.90(1.05-2.60)	23	74 Tryptophan	0.40(0.23-0.70) <sup>2</sup>	94,102
31 Tin	(0.00013-0.00025)	11,41	75	0.20(0.11-0.36) <sup>3</sup>	94,102
32 Zinc	0.018(0.011-0.033)	88	76 Tyrosine	0.70(0.44-0.82) <sup>2</sup>	85,94,102
Nitrogenous Substances, mg			77	0.20(0.15-0.30) <sup>3</sup>	85,94,102
31 Protein	(0.03-1.00)	69	78 Valine	0.30(0.25-0.42) <sup>2</sup>	20,75,94
32 Acetylkynurenine	0.03	7	79	0.09(0.04-0.18) <sup>3</sup>	20,75,94
33 Adenine	0.020(0.016-0.024)	98	80 o-Aminohippuric acid	0.06	7
34 Allantoin	0.17(0.14-0.21)	100	81 Anthranilic acid	(0.002-0.009)	1
35 Amino acids	(20-40) <sup>2</sup>	9	82 Bilirubin	0.70	58
36	(13-20) <sup>3</sup>	9	83 Carnosine (+ anserine)	(0.045-0.14)	13
37 Alanine	0.55 <sup>2</sup>	9	84 Coproporphyrin I & III	(0.00024-0.00400)	96
38 β-Alanine	(0.2-0.3)	13	85 Creatine	0.8(0-2.0)	89
39 Arginine	0.45(0.34-0.50) <sup>2</sup>	89	86 Creatinine	23(15-30)	89
40	0.16(0.07-0.30) <sup>3</sup>	89	87 1,3-Dimethyluric acid	Trace	17
41 Asparagine	0.77	78	88 Dopamine	(0.0041-0.0063)	3
42 Aspartic acid	1.7(1.2-2.7) <sup>2</sup>	102	89 Epinephrine	0.00014	92
43	0.04(0.01-0.07) <sup>3</sup>	102	90 Ergothionine	(1.7-4.0)	60
			91 Ethanolamine	(0.07-0.7)	49
			92 Guanidinoacetic acid	(0.2-0.5)	89
			93 Guanine	0.006(0.003-0.009)	98
			94 Hippuric acid	(7.0-18.0)	101
			95 Histamine	(0.00025-0.001)	39,71

<sup>1/</sup> Upper limit of range was obtained in an area of Texas where dental fluorosis is endemic. <sup>2/</sup> Total. <sup>3/</sup> Free. <sup>4/</sup> Determined by chromatography.

*continued*



## 53. EXCRETION PRODUCTS: MAN

### Part I. URINE

Constituent	Amount Excreted per kg body wt per da	Refer- ence	Constituent	Amount Excreted per kg body wt per da	Refer- ence		
(A)	(B)	(C)	(A)	(B)	(C)		
Nitrogenous Substances, mg							
96	3-Hydroxyanthranilic acid	(0.008-0.04)	1	144	Citric acid	(3-20)	84
97	<i>p</i> -Hydroxybenzylamine	0.002	39	145	Formic acid	0.8(0.4-2.0)	4
98	5-Hydroxyindoleacetic acid	(0.02-0.03)	101	146	Lactic acid	3(2-5)	26
99	3-Hydroxykynurenine	0.075	7	147	Oxalic acid	0.5(0.3-0.7)	32
100	8-Hydroxy-7-methylguanaine	0.020(0.016-0.030)	98	148	Arabinose	Trace <sup>4</sup>	87
101	<i>p</i> -Hydroxyphenylacetic acid	(0.2-1.2)	101	149	Deoxyribose	Trace <sup>4</sup>	87
102	Hypoxanthine	0.14(0.08-0.19)	98	150	Galactose	Trace <sup>5</sup>	87
103	Imidazole derivatives	(2-3)	44	151	Glucose	Trace <sup>4</sup>	87
104	Indoleacetic acid	(0.02-0.06)	101	152	Glucuronic acid	Trace <sup>4</sup>	87
105	Indoxylsulfuric acid <sup>6</sup>	1.0(0.5-2.0)	74	153	Lactose	Trace <sup>4</sup>	87
106	Kynurenic acid	0.03	7	154	Ribose	Trace <sup>4</sup>	87
107	Kynurenine	(0.023-0.078)	1	155	Ribulose	Trace <sup>4</sup>	25
108	Metanephrine	(0.002-0.006)	39	156	Xylose	Trace <sup>4</sup>	25
109	Methionine sulfoxide	(0-0.31)	9,103	157	Xylulose	Trace <sup>4</sup>	86
110	3-Methoxytyramine	(0-0.0005)	39	Vitamins and Related Compounds, µg			
111	7-Methylguanaine	0.09(0.08-0.11)	98	158	Vitamins A, D, K	(0-trace)	82
112	<i>N</i> <sup>2</sup> -Methylguanaine	0.007(0.006-0.009)	98	159	<i>p</i> -Aminobenzoic acid	(2-3)	16
113	1-Methylhypoxanthine	0.006(0.003-0.010)	98	160	Ascorbic acid	(100-400)	10
114	<i>N</i> -Methyl-2-pyridone-5-carboxamide	0.24	7	161	Biotin	0.5(0.2-1.0)	16
115	Norepinephrine	0.001	92	162	Choline	79(68-130)	36
116	Normetanephrine	(0.0002-0.0005)	39	163	Citrovorum factor	0.037(0.023-0.069)	68
117	Purine bases	(0.2-1.0)	32	164	Cyanocobalamin	0.00044	68
118	Serotonin	(0.00025-0.001)	39			(0.00023-0.00079)	
119	6-Succinopurine	0.014	99	165	Inositol	200	38
120	Theophylline	Trace	17	166	Nicotinic acid <sup>7</sup>	3.4(2.0-20.0)	16,37
121	Tryptamine	(0.0013-0.0028)	63	167	Nicotinamide <sup>8</sup>	20(10-50)	16,37
122	<i>m</i> -Tyramine	(0.001-0.0025)	39	168	Pantothenic acid	45(16-100)	16
123	<i>p</i> -Tyramine	(0.0005-0.0025)	39	169	Pteroylglutamic acid <sup>9</sup>	0.058(0.030-0.300)	16,68
124	Urea	(200-500)	32	170	Pyridoxine	(0.08-2.70)	35
125	Uric acid	2.0(0.8-3.0)	32	171	Riboflavin	12.4(2.0-24.0)	16,61
126	Urobilin	(0.143-1.857)	46	172	Thiamine	3.0(0.6-6.0)	16
127	Urobilinogen	(0.043-0.357)	46	173	Dehydroascorbic acid	(190-290)	10
128	Xanthine	0.09(0.07-0.12)	98	174	Dehydroascorbic + diketogulonic acid	230(0-1,280)	24
129	Xanthurenic acid	0.02	7	175	Diketogulonic acid	(140-190)	10
Nitrogen							
130	Total N	(130-300)	32	176	<i>N</i> -Methylnicotinamide	(40-600)	16,37
131	Amino acid N	(3-6)	31	177	Pyridoxal	1.0(0.7-5.3)	35,64
132	Ammonia N	(3-13)	95	178	Pyridoxamine	1.6(0.4-3.0)	64
133	Protein N	(0.0046-0.0180)	95	179	4-Pyridoxic acid	(9-160)	35,64
Lipids, Carbohydrates, Miscellaneous Organic Acids, mg							
134	Aconitic acid	Trace	29	180	Trigonelline	(30-300)	62
135	Cholesterol	(0-0.0714)	54	Hormones, µg			
136	Homovanillic acid	(0.065-0.110)	101	181	Aldosterone		
137	3-Methoxy, 4-hydroxymandelic acid	0.053	92		♂	0.05(0.01-0.13)	91
138	Reducing substances	(7-21)	6	182	♀	0.06(0.03-0.10)	91
139	Acetone bodies	0.20(0.03-0.30) <sup>2</sup>	77	Androgens			
140	Acetoacetic acid	0.04(0.03-0.06)	77	183	♂, 3-5 yr	210	19
141	Carbolic acid <sup>6</sup>	(0.2-0.6) <sup>2</sup>	15	184	♂, 20-40 yr	260(200-330)	19
142		(0-0.05) <sup>3</sup>	15	185	♂, 60+ yr	70(30-130)	19
143	Carbonic acid	2.7(2.1-3.3)	27	186	♀, 3-5 yr	50	19
				187	♀, 20-40 yr	200(180-210)	19
				188	♀, 60+ yr	40(15-130)	19
				Androsterone			
				189	♂	50(35-60)	73
				190	♀	60(50-80)	73
				Etiocolanolone			
				191	♂	60(40-70)	73
				192	♀	50(30-60)	73
				Estradiol			
				193	♀, follicular phase	0.03(0-0.05)	70

<sup>1</sup>/<sub>2</sub> Total. <sup>1</sup>/<sub>3</sub> Free. <sup>1</sup>/<sub>4</sub> Determined by chromatography. <sup>1</sup>/<sub>5</sub> Indican. <sup>1</sup>/<sub>6</sub> Phenol. <sup>1</sup>/<sub>7</sub> Niacin. <sup>1</sup>/<sub>8</sub> Niacinamide. <sup>1</sup>/<sub>9</sub> Folic acid.

continued

## 53. EXCRETION PRODUCTS: MAN

### Part I. URINE

Constituent	Amount Excreted per kg body wt per da	Refer- ence	Constituent	Amount Excreted per kg body wt per da	Refer- ence
(A)	(B)	(C)	(A)	(B)	(C)
<b>Hormones, µg</b>					
	<b>Estradiol</b>				
194	♀, luteal phase	0.10(0.07-0.17)	70		
195	♀, postmenopause	0.01(0-0.09)	50		
	<b>Estriol</b>				
196	♀, follicular phase	0.1(0-0.3)	70		
197	♀, luteal phase	0.40(0.13-1.30)	70		
198	♀, postmenopause	0.05(0-0.18)	50		
	<b>Estrone</b>				
199	♀, follicular phase	0.08(0.06-0.12)	70		
200	♀, luteal phase	0.20(0.17-0.40)	70		
201	♀, postmenopause	0.03(0-0.12)	50		
	<b>17-Hydroxysteroids</b>				
202	♂	80(40-170)	66,67		
203	♀	60(20-140)	66,67		
	<b>17-Ketogenic adreno- corticoids</b>				
204	♂	210(150-310)	18		
205	♀	180(120-300)	18		
206	α-Ketol steroids	260(130-470)	53		
	<b>Pregnanediol</b>				
207	♂	13(5-20)	43		
208	♀, follicular phase	18(13-25)	43		
209	♀, luteal phase	55(30-70)	43		
				<b>Enzymes</b>	
			210	Pregnanediol ♀, postmenopause	10(5-14) 43
			211	Pregnanetriol ♀, follicular phase	25 79
			212	♀, luteal phase	32 79
			213	♀, postmenopause	11 79
			214	Tetrahydrocortisol	24(8-50) 70
			215	Tetrahydrocortisone	54(20-120) 70
			216	Adrenocorticotropin	72
			217	Insulin	57
			218	Melanocyte-stimulating hormone	76
			219	Parathyroid hormone	14
				} Consult references	
			220	Acid phosphatase	2
			221	Amylase	52
			222	Cadaverinase	40
			223	Cathepsin	56
			224	β-Glucuronidase	55
			225	Histaminase	40
			226	Lipase	59
			227	Maltase	22
			228	Ribonuclease	48
			229	Uropepsinogen	34

*Contributor:* Van Pilsum, John F.

- References:* [1] Abul-Fadl, M. A. M., and A. S. Khalafallah. 1961. *Brit. J. Cancer* 15:479. [2] Aoyama, S. 1961. *Acta Schol. Med. Univ. Kioto* 37:203. [3] Barbeau, A., and T. L. Sourkes. 1961. *Rev. Can. Biol.* 20:197. [4] Benedict, E. M., and G. A. Harrop. 1922. *J. Biol. Chem.* 54:443. [5] Bloomfield, I. J., R. R. Sayers, and F. H. Goldman. 1932. *Public Health Rept. (U.S.)* 50:421. [6] Bradley, S. E. 1945. *Med. Clin. N. Am.* 29:1314. [7] Brown, R. R., M. J. Thornton, and J. M. Price. 1961. *J. Clin. Invest.* 40:617. [8] Bruger, M., J. W. Hinton, and W. G. Lough. 1941. *J. Lab. Clin. Med.* 26:1942. [9] Carsten, M. E. 1952. *J. Am. Chem. Soc.* 74:5954. [10] Chen, S. D., and C. Shuck. 1951. *J. Nutr.* 23:111. [11] Clark, G. W. 1926. *Univ. Calif. (Berkeley) Publ. Physiol.* 5(17):195. [12] Conway, E. J., and J. C. Flood. 1936. *Biochem. J.* 30:716. [13] Crokaert, R. 1953. *Ann. Soc. Roy. Sci. Med. Nat. Bruxelles* 6:157. [14] Davies, B. M. A. 1958. *J. Endocrinol.* 16:369. [15] Deichmann, W., and L. J. Schafer. 1942. *Am. J. Clin. Pathol.* 12:129. [16] Denko, C. W., et al. 1946. *Arch. Biochem.* 10:33. [17] Dikstein, S., F. Bergman, and M. Chaimovitz. 1958. *J. Biol. Chem.* 230:203. [18] Diszfalusy, E., et al. 1955. *Acta Endocrinol.* 18:356. [19] Dorfman, R. I., and R. A. Shipley. 1956. *The androgens.* J. Wiley, New York. [20] Dunn, M. S., et al. 1947. *Arch. Biochem.* 13:207. [21] Figueroa, W. G., et al. 1955. *J. Lab. Clin. Med.* 46:534. [22] Fleury, P. F., J. E. Courtois, and D. Ramon. 1951. *Bull. Soc. Chim. Biol.* 33:1762. [23] Folin, O. 1905. *Am. J. Physiol.* 13:45. [24] Freeman, J. T., R. Hafkesbring, and E. K. Caldwell. 1951. *Gastroenterology* 18:224. [25] Futterman, S., and J. H. Roe. 1955. *J. Biol. Chem.* 215:257. [26] Gambigliani-Zoccoli, A., et al. 1939. *Z. Klin. Med.* 135:457. [27] Gamble, J. L. 1954. *Chemical anatomy, physiology and pathology of extracellular fluid.* Ed. 6. Harvard Univ. Press, Cambridge. [28] Gwens, M. H. 1918. *J. Biol. Chem.* 34:119. [29] Halpern, M. N. 1960. *Clin. Chim. Acta* 5:264. [30] Harp, M. J., and F. I. Scoular. 1952. *J. Nutr.* 47:67. [31] Harrow, B., and A. Mazur. 1962. *Textbook of biochemistry.* Ed. 8. W. B. Saunders, Philadelphia. [32] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1947. *Practical physiological chemistry.* Ed. 12. Blakiston, Philadelphia. [33] Hier, S. W. 1948. *Trans. N. Y. Acad. Sci.* 10:200. [34] Hostrup, H., and P. Bastrup-Madsen. 1957. *Acta Med. Scand.* 158:193. [35] Johnson, B. C., T. S. Hamilton,

*continued*

**53. EXCRETION PRODUCTS: MAN**

**Part I. URINE**

and H. H. Mitchell. 1945. J. Biol. Chem. 158:619. [36] Johnson, B. C., T. S. Hamilton, and H. H. Mitchell. 1945. Ibid. 159:5. [37] Johnson, B. C., T. S. Hamilton, and H. H. Mitchell. 1945. Ibid. 159:231. [38] Johnson, B. C., H. H. Mitchell, and T. S. Hamilton. 1945. Ibid. 161:357. [39] Kakimoto, Y., and M. D. Armstrong. 1962. Ibid. 237:208. [40] Kapeller-Adler, R., and R. Renwick. 1956. Clin. Chim. Acta 1:197. [41] Kehoe, R. A., J. Cholak, and R. V. Story. 1940. J. Nutr. 19:579. [42] Kent, N. L., and R. A. McCance. 1941. Biochem. J. 35:877. [43] Klopper, A., E. A. Mitchie, and J. B. Brown. 1955. J. Endocrinol. 12:209. [44] Koessler, K. K., and M. T. Hanke. 1924. J. Biol. Chem. 59:803. [45] Kolmer, J. A., et al. 1951. Approved laboratory technique. Ed. 5. Appleton-Century-Crofts, New York. [46] Lemberg, R., and J. W. Legge. 1949. Hematin compounds and bile pigments. Interscience, New York. [47] Levinson, S. A., and R. P. MacFate. 1961. Clinical laboratory diagnosis. Ed. 6. Lea and Febiger, Philadelphia. [48] Levy, A. L., and A. Rottino. 1960. Clin. Chem. 6:43. [49] Luck, J. M., and A. Wilcox. 1953. J. Biol. Chem. 205:859. [50] McBride, J. M. 1957. J. Clin. Endocrinol. Metab. 17:1440. [51] McClure, F. J. 1944. Public Health Rept. (U.S.) 59:1575. [52] MacFate, R. P. 1961. Assoc. Clin. Scientists Proc. 2nd Appl. Seminar 2:14. [53] Marks, L. J., J. H. Leftin, and P. Leonard. 1957. J. Clin. Endocrinol. Metab. 17:407. [54] Mattice, M. R. 1936. Chemical procedures for clinical laboratories. Lea and Febiger, Philadelphia. [55] Melicow, M. M., A. C. Uson, and R. Lipton. 1961. J. Urol. 86:89. [56] Merten, R., and H. Wojta. 1954. Z. Ges. Exptl. Med. 123:315. [57] Mirsky, I. A., et al. 1948. J. Clin. Invest. 27:515. [58] Nauman, H. N. 1936. Biochem. J. 36:692. [59] Nothmann, M. M., J. H. Pratt, and A. D. Callow. 1955. Arch. Internal Med. 96:188. [60] Ohara, M., et al. 1952. Japan. J. Med. Sci. Biol. 5:259. [61] Oldham, H., B. B. Sheft, and T. Porter. 1950. J. Nutr. 41:231. [62] Perlzweig, W. A., H. P. Sarett, and L. H. Margoles. 1942. J. Am. Med. Assoc. 118:28. [63] Perry, T. L. 1962. Science 136:879. [64] Rabinowitz, J. C., and E. E. Snell. 1949. Proc. Soc. Exptl. Biol. Med. 70:235. [65] Rae, J. J. 1937. Biochem. J. 31:1622. [66] Reddy, W. J., D. Jenkins, and G. W. Thorn. 1952. Metab. Clin. Exptl. 1:511. [67] Reddy, W. J., et al. 1956. J. Clin. Endocrinol. Metab. 16:380. [68] Register, V. D., and H. P. Sarett. 1951. Proc. Soc. Exptl. Biol. Med. 77:837. [69] Rigas, D. A., and C. G. Heller. 1951. J. Clin. Invest. 30:853. [70] Romanoff, L. P., et al. 1957. J. Clin. Endocrinol. Metab. 17:777. [71] Rose, B., et al. 1951. Proc. Clin. ACTH Conf., 2nd, 1:519. [72] Rubin, B. L., R. I. Dorfman, and A. Dorfman. 1954. J. Clin. Endocrinol. 14:154. [73] Rubin, B. L., R. I. Dorfman, and G. Pincus. 1954. Recent Progr. Hormone Res. 9:213. [74] Sharlit, H. 1938. Arch. Pediat. 55:277. [75] Sheffner, A. L., J. B. Kirsner, and W. L. Palmer. 1948. J. Biol. Chem. 175:107. [76] Shizume, K., W. Mori, and A. B. Lerner. 1962. Gen. Comp. Endocrinol., Suppl. 1:110. [77] Stark, I. E., and M. Somogyi. 1943. J. Biol. Chem. 147:319. [78] Stein, W. H. 1953. J. Biol. Chem. 201:45. [79] Stern, M. I. 1957. J. Endocrinol. 16:180. [80] Sterner, J. H., and V. Lidfeldt. 1941. J. Pharmacol. Exptl. Therap. 73:205. [81] Stock, A. 1940. Biochem. Z. 304:73. [82] Sunderman, F. W., and F. Boerner. 1949. Normal values in clinical medicine. W. B. Saunders, Philadelphia. [83] Tallan, H. H., W. H. Stein, and S. Moore. 1954. J. Biol. Chem. 206:825. [84] Taussky, H. H. 1949. Ibid. 181:195. [85] Tompsett, S. L., and J. Fitzpatrick. 1950. Brit. J. Exptl. Pathol. 31:70. [86] Touster, O., R. M. Hutcheson, and V. H. Reynolds. 1954. J. Am. Chem. Soc. 76:5005. [87] Tower, D. B., E. L. Peters, and M. A. Pogorelskin. 1956. Neurology 6:37. [88] Tribble, H. M., and F. I. Scouler. 1954. J. Nutr. 52:210. [89] Van Pilsum, J. F., et al. 1956. J. Biol. Chem. 222:225. [90] Van Ravesteyn, A. H. 1944. Acta Med. Scand. 118:163. [91] Venning, E. H., I. Dyrenfurth, and C. J. P. Giroud. 1956. J. Clin. Endocrinol. Metab. 16:1326. [92] Voorhess, M. L., and L. I. Gardner. 1962. J. Clin. Endocrinol. Metab. 22:126. [93] Walker, B. S. 1931. J. Lab. Clin. Med. 17:347. [94] Wallraff, E. B., G. C. Brodie, and A. L. Borden. 1950. J. Clin. Invest. 29:1542. [95] Wang, C. C., et al. 1930. J. Nutr. 3:79. [96] Watson, C. J., et al. 1949. J. Clin. Invest. 28:447. [97] Webster, S. H. 1941. Public Health Rept. (U.S.) 56:1953. [98] Weissmann, B., P. A. Bromberg, and A. B. Gutman. 1957. J. Biol. Chem. 224:423. [99] Weissmann, B., and A. B. Gutman. 1957. Ibid. 229:239. [100] Wiechouski, W. 1909. Biochem. Z. 19:368. [101] Williams, C. M., and C. C. Sweeley. 1961. J. Clin. Endocrinol. Metab. 21:1500. [102] Woodson, H. W., et al. 1948. J. Biol. Chem. 172:613. [103] Young, M. K., et al. 1951. Texas Univ. Publ. 5109:189.

*continued*



*Contrails*  
53. EXCRETION PRODUCTS: MAN

Part II. FECES

Constituent			Constituent		
(A)	Amount Excreted per kg body wt per da (B)	Reference (C)	(A)	Amount Excreted per kg body wt per da (B)	Reference (C)
General Chemical Components, mg			Amino acids		
1 Solids	394(140-560)	26	26 Lysine	5.7(4.5-6.9) <sup>1</sup>	18
2 Water	(910-1,820)	21	27 Threonine	4.0(3.3-5.2) <sup>1</sup>	18
Electrolytes, mg			28 Valine	4.6(3.6-6.2) <sup>1</sup>	18
3 Aluminum	0.0006	10	29 Imidazole derivatives	(0-0.2)	14
4 Arsenic	0.033(0.001-0.116)	17	30 Purine bases	(2-3)	15
5 Calcium	(5-10)	19	31 Urobilinogen	(0.00057-0.00400)	24
6 Chlorine	(0.21-0.50)	1	Nitrogen		
7 Cobalt	(0.000002-0.000020)	8	32 Total N	(11.4-36.0)	9
8 Copper	0.027(0.023-0.037)	10	33 Ammonia N	(0.36-1.20)	16
9 Iron	120(65-208)	4	Lipids and Miscellaneous Organic Acids, mg		
10 Lead	0.0042	10	Fat		
11 Magnesium	2.5(1.510-3.185)	13	34 Total	56(30-100)	26
12 Manganese	(0.018-0.120)	10,11	35 Neutral	(10-45)	7
13 Mercury	0.00014	20	36 Unsaponifiable	33(22-38) <sup>2</sup>	25
14 Nickel	(0.0012-0.0025)	11	37 Fatty acids	16(4-38) <sup>1</sup>	7
15 Phosphorus	0.00986	4	38	30(4-64) <sup>3</sup>	7
16 Potassium	(0.00710-0.0200) <sup>1</sup>	3	39 Soaps	53(40-66) <sup>2</sup>	25
17 Silver	6.7	10	40 Carboic acid	(0-3) <sup>1</sup>	6
18 Sodium	0.0008	3	Vitamins and Related Compounds, µg		
19 Sulfur	1.7	3	41 β-Aminobenzoic acid	3.50(1.01-8.20)	5
20 Tin	2.0 <sup>1</sup>	3	42 Ascorbic acid	(60-70)	2
21 Zinc	(0.17-0.45)	3,10	43 Biotin	1.90(0.63-6.64)	5
Nitrogenous Substances, mg			44 Vitamin E	308(226-391)	12
Amino acids			45 Nicotinic acid	52(12-124)	5
22 Arginine	3.8(2.9-5.0) <sup>1</sup>	18	46 Pantothenic acid	31.40(3.85-63.40)	5
23 Histidine	1.7(1.4-2.1) <sup>1</sup>	18	47 Pteroylglutamic acid	4.3(1.8-7.7)	5
24 Isoleucine	4.3(3.3-5.5) <sup>1</sup>	18	48 Riboflavin	14.7(8.0-23.0)	5
25 Leucine	5.6(4.3-6.9) <sup>1</sup>	18	49 Thiamine	7.80(0.67-18.00)	5
			50 Carotene + xanthophyll	(20-600)	23
			51 Xanthophyll	(8-100)	23

/1/ Total. /2/ At age 8-12 years. /3/ Free.


Contributor: Van Pilsum, John F.

References: [1] Cammidge, P. J. 1914. The faeces of children and adults. W. Wood, New York. [2] Chinn, H., and C. J. Farmer. 1939. Proc. Soc. Exptl. Biol. Med. 41:561. [3] Clark, G. W. 1926. Univ. Calif. (Berkeley) Publ. Physiol. 5(17):195. [4] Daum, K., et al. 1951. J. Am. Dietet. Assoc. 27:475. [5] Denko, C. W., et al. 1946. Arch. Biochem. 10:33. [6] Folin, O., and W. Denis. 1916. J. Biol. Chem. 26:507. [7] Fowweather, F. S. 1926. Brit. J. Exptl. Pathol. 7:15. [8] Harp, M. J., and F. I. Scoular. 1952. J. Nutr. 47:67. [9] Hawk, P. B., B. L. Oser, and W. H. Summerson. 1947. Practical physiological chemistry. Ed. 12. Blakiston, Philadelphia. [10] Kehoe, R. A., J. Cholak, and R. V. Story. 1940. J. Nutr. 19:579. [11] Kent, N. L., and R. A. McCance. 1941. Biochem. J. 35:877. [12] Klatskin, G., and D. W. Molander. 1952. J. Lab. Clin. Med. 39:802. [13] Leichsenring, J. M., L. M. Norris, and S. A. Lamison. 1951. J. Nutr. 45:477. [14] Loeper, M., A. Lesurl, and A. Thomas. 1934. Bull. Soc. Chim. Biol. 16:385. [15] Mendel, L. B., and J. F. Lyman. 1910. J. Biol. Chem. 8:115. [16] Robinson, C. S. 1922. Ibid. 52:445. [17] Schwarz, L., and W. Deckert. 1931. Arch. Hyg. Bakteriol. 106:346. [18] Sheffner, A. L., J. B. Kirsner, and W. L. Palmer. 1948. J. Biol. Chem. 175:107. [19] Shohl, A. T. 1939. Mineral metabolism. Reinhold, New York. [20] Stock, A. 1940. Biochem. Z. 304:73. [21] Sunderman, F. W., and F. Boerner. 1949. Normal values in clinical medicine. W. B. Saunders, Philadelphia. [22] Tribble, H. M., and F. I. Scoular. 1954. J. Nutr. 52:210. [23] Wald, G., W. R. Carroll, and D. Sciarra. 1941. Science 94:95. [24] Watson, C. J. 1937. Arch. Internal Med. 59:196. [25] Williams, H. H., et al. 1943. J. Nutr. 25:379. [26] Wollaeger, E. E., M. W. Comfort, and A. E. Osterberg. 1947. Gastroenterology 9:272.



**VI. METABOLISM**

DIAGRAM FOR TABLE 54



<sup>1</sup>/ Percent absorption from oral administration has in some cases been rather arbitrarily classified, since the extent of absorption may depend on the amount administered and on the presence or absence of food residues in the digestive tract. <sup>2</sup>/ Some trace elements with no known function also included. <sup>3</sup>/ Primarily muscle, skin, and extracellular fluids. <sup>4</sup>/ Other than in the bile, or by a route not definitely established.

### 54. PATHWAYS OF MINERAL METABOLISM: LABORATORY MAMMALS

The course, or courses, of various ions during metabolism can be located in the diagram (at left) by tracing the combination of letters and numbers accompanying each ion (columns B and C below). Observations were made on a wide variety of mammalian species. Ions were administered in the form of simple soluble compounds or metallic oxides, unless otherwise specified. Underscoring indicates radioactive elements, or that data were obtained at least in part from studies using radioactive isotopes. Different isotopes of the same element may show different tissue predilections, but there is usually no difference in their absorption or route of excretion. "Plus" symbols indicate valence states to which data apply. **Other Known Pathways** (column C) are listed, so far as possible, in order of decreasing importance.

Ion	Principal Oral Pathways <sup>1</sup>	Other Known Pathways	Reference
(A)	(B)	(C)	(D)
Cations <sup>2</sup>			
1 <u>Actinium</u>	CE	PDFK(1, 2)	10
2 <u>Aluminum</u>	CE	PDH2, PDH3, PDFJ(2)	40,91,93
3 <u>Americium</u>	CE	PDH4, PDFK(2, 1), PDH2	88,102
4 <u>Antimony</u> <sup>+++</sup>	BE, BDH2	BDFK(2, 3), BDH4	31,34,40,93
5 <u>Arsenic</u> <sup>+++</sup>	BE, BDH2	BDFK(2, 3), BDH4	16,40,54,58,93
6 <u>Barium</u>	BE, BDH4	BDFK(1), BDH2	12,36,69,93,102
7 <u>Beryllium</u>	CE	PDFK(1, 3, 2), PDH2, PDH4	83,93
8 <u>Bismuth</u> <sup>+++</sup>	CE	PDFK(2), PDH4, PDH2	40,57,93
9 <u>Cadmium</u>	CE	PDFK(2, 3, 1), PDH4, PDH2	20,23,40,78,93,110
10 <u>Calcium</u>	BE, BDFJ(1)	BDH4, BDH3, BDH2, BDFJ(3)	3,19,40,75,92,97,103,110,111
11 <u>Cerium</u> <sup>3</sup>	CE	PDFK(2, 1), PDH3, PDH2	23,24,40,46,94,102
12 <u>Cesium</u>	ADH2	ADH3, ADH4, ADFJ(3)	36,52,71-74,90,93
13 <u>Chromium</u> <sup>+++</sup>	CE	PDH2, PDH4, PDFK(2, 1)	23,25,50,106
14 <u>Cobalt</u> <sup>++</sup>	BE, BDH2	BDH3, BDH4, BDFJ(2)	5,17,19,40,56,93,110
15 <u>Copper</u> <sup>++</sup>	BE, BDH3	BDH2, BDFJ(2)	19,40,64,93,110
16 <u>Curium</u>	Probably CE <sup>4</sup>	PDFK(2, 1), PDH4, PDH2	86, 102
17 <u>Dysprosium</u> <sup>3</sup>	.....	PDH3, PDFK(1)	24
18 <u>Erbium</u> <sup>3</sup>	.....	PDH3, PDFK(1, 2)	24
19 <u>Europium</u> <sup>3</sup>	CE	PDH3, PDH2, PDFK(1, 2)	24
20 <u>Francium</u>	Probably ADH2 <sup>4</sup>	PDFK(2, 3)	76
21 <u>Gadolinium</u> <sup>3</sup>	.....	PDH2, PDH3, PDFK(1, 2)	24
22 <u>Gallium</u>	CE	PDH2, PDFK(1, 2), PDH4	22,53,93
23 <u>Germanium</u>	ADH2	ADH4, ADFK(2)	93
24 <u>Gold</u> <sup>+++</sup>	CE	PDFK(2), PDH2, PDH4	34,40,79,93
25 <u>Hafnium</u> <sup>3</sup>	.....	PDFK(2, 1), PDH2, PDH4	47,93
26 <u>Holmium</u> <sup>3</sup>	.....	PDH3, PDFK(1, 2)	24
27 <u>Indium</u>	CE	PDFK(3, 1, 2), PDH4, PDH2	23,39
28 <u>Iridium</u>	CE	PDH2, PDH4, PDFK(3, 2, 1)	23
29 <u>Iron</u> <sup>++</sup>	CE	PDH4, PDFJ(2)	19,40,65,93,110,111
30 <u>Lanthanum</u> <sup>3</sup>	CE	PDH2, PDH3, PDFK(2, 1)	13,23,24,36,40,93,95,102
31 <u>Lead</u> <sup>++</sup>	BE, BDH4	BDH2, BDFK(1, 3, 2)	1,9,23,34,35,40,57,82,93
32 <u>Lithium</u>	ADH2	ADH4, ADFJ(3, 2)	30,35,81,93
33 <u>Lutetium</u> <sup>3</sup>	.....	PDH3, PDFK(1)	24
34 <u>Magnesium</u>	BE, BDH2	BDH3, BDFJ(1, 2, 3)	2,40,91,93,110
35 <u>Manganese</u> <sup>++</sup>	CE	PDH3, PDH4, PDFJ(2, 3)	8,19,28,40,60,66,93
36 <u>Mercury</u> <sup>++</sup>	BE, BDH2	BDFK(1, 2, 3), BDH3, BDH4	23,40,93
37 <u>Neodymium</u> <sup>3</sup>	Probably CE <sup>4</sup>	PDH2, PDFK(2, 1), PDH3	23,24,38,40,55
38 <u>Neptunium</u>	CE	PDFK(1)	36
39 <u>Nickel</u> <sup>++</sup>	BE, BDH2	BDH4, BDFJ(2)	5,35,40,93,98,108,110
40 <u>Niobium</u> <sup>3</sup>	CE	PDH2, PDH4, PDFK(2, 1, 3)	13,23,36,80,102
41 <u>Palladium</u>	Probably CE <sup>4</sup>	PDH2, PDH4, PDFK(2)	23,40,67
42 <u>Platinum</u>	CE	PDH2, PDH4, PDFK(2, 3, 1)	23,40
43 <u>Plutonium</u> <sup>3</sup>	CE	PDFK(1, 2), PDH4, PDH2	11,29,36,84,89,102,109
44 <u>Polonium</u>	CE	PDFK(2, 1, 3), PDH4, PDH2	27,44,93,96,102

<sup>1/2</sup> Ions may be assumed to follow the same pathways when given parenterally. <sup>1/3</sup> Because of inadequate information, no pathways have been listed for berkelium, californium, einsteinium, fermium, and mendelevium. <sup>1/3</sup> Usually given as soluble complex. <sup>1/4</sup> As judged from the position of the element in the periodic table, or on solubility at neutral pH values.

*continued*

## 54. PATHWAYS OF MINERAL METABOLISM: LABORATORY MAMMALS

	Ion	Principal Oral Pathways <sup>1</sup>	Other Known Pathways	Reference
	(A)	(B)	(C)	(D)
Cations				
45	Potassium	ADH2	ADFJ(3,2), ADH3, ADH4	18,19,40,71,73,103
46	Praseodymium <sup>a</sup>	CE	PDH2, PDFK(2,1), PDH3	24,36,40
47	Promethium	CE	PDH2, PDFK(2,1), PDH3	24,36,102
48	Protactinium	Probably CE <sup>4</sup>	PDFK(1)	88
49	Radium	BE, BDFK(1,2)	BDH4, BDH2	27,45,75,93,97,102
50	Radium D	Probably BE, BDH2 <sup>4</sup>	PDH2, PDH4, PDFK(1)	9,70
51	Rhodium	CE	PDFK(3,2,1), PDH2, PDH4	23
52	Rubidium	ADH2	ADFJ(3,2), ADH3, ADH4	68,71
53	Ruthenium	CE	PDH2, PDH4, PDFK(3,1,2)	23,100,101
54	Samarium <sup>3</sup>	CE	PDH2, PDFK(2,1), PDH3	23,40
55	Scandium	Probably CE <sup>4</sup>	CDH2, CDFJ(1,3)	4
56	Selenium	ADH2	ADGH1, ADFK(2,3), ADH4	61,62,93
57	Silver	CE	PDH3, PDFK(2,3), PDH2, CDGFK <sup>5</sup>	40,87,93,110
58	Sodium	ADH2	ADFJ(3,1,2), ADH3, ADH4	19,40,91,93
59	Strontium	BE, BDFK(1)	BDH2, BDH4, BDH3	3,36,45,75,92,102,103
60	Tantalum	CE	PDH2, PDH4, PDFK(2,1,3)	23,24
61	Technetium	.....	PDH2, PDH4, PDFK(3,2,1)	23
62	Tellurium <sup>+++</sup>	BE, BDGH2	BDGH3, BDGH1, BDGFK(2)	36,93,102
63	Terbium <sup>3</sup>	CE	PDH2, PDH3, PDFK(1,2)	24
64	Thallium	BE	BDH4, BDFK(3,1,2), BDH2	40,59,93
65	Thorium	CE	PDFK(2,1), PDH4, PDH2, PDH3	36,85,93,102
66	Thulium <sup>3</sup>	CE	PDH2, PDH3, PDFK(1)	24
67	Tin <sup>++</sup>	BE, BDH2	BDFJ(3,2), BDH4, BDH3	40,93,110
68	Tin <sup>++++</sup> <sup>2</sup>	BE, BDH2	BDH4, BDFJ(1,3,2)	23
69	Titanium	Probably CE <sup>4</sup>	CDFJ(2)	93
70	Uranium <sup>+++</sup>	.....	BDFK(2)	102
71	Uranium <sup>+++++</sup>	BE, BDH2	BDFK(1,2)	21,40,93,102
72	Ytterbium <sup>3</sup>	.....	PDH3, PDFK(1,2)	24
73	Yttrium	CE	PDFK(1,2), PDH2, PDH4	13,24,32,36,46,84,102,109
74	Zinc	CE	CDH4, CDFJ(2,3), CDH2	33,40,60,77,93,102-105,110
75	Zirconium	CE	PDFK(1,3)	13,36,80,102
Anions <sup>6</sup>				
76	Astatide	Probably A <sup>7</sup>	PDFK(3), PDH2, PDH4	37
77	Bicarbonate	ADH1, ADH2, ADH3	ADH4, ADFJ (all tissues)	93
78	Borate	ADH2	.....	40,93
79	Bromate	ADH2	ADG (to bromide)	41,93
80	Bromide	ADH2	ADH3, ADH4, ADFJ(3)	14,93
81	Chlorate	ADH2	.....	40,93
82	Chloride	ADH2	ADH3, ADH4, ADFJ(3,1,2)	91,93
83	Chromate	BDH2	BDH4, BDGH2, BDGH4, BDGFK(2)	40,63,93,106
84	Cyanide	ADH2	ADH1, ADG (to SCN <sup>-</sup> )	93
85	Ferrocyanide	ADH2	.....	48,49,93
86	Fluoride	BDH2	BDFJ(1,3), BDH4	60,91,93
87	Hypophosphite	ADH2	.....	93
88	Iodate	ADG (to iodide)	PDH2	93
89	Iodide	ADH2	ADFJ(3,2), ADH3	93
90	Molybdate	BDH2	BDFJ(2,3,1)	7,15,24,35,40,99,102
91	Nitrate	ADH2	ADFK(3)	93
92	Nitrite	ADG (to nitrate)	.....	93
93	Osmate	.....	PDH2, PDH4, PDFK(3,2,1)	23
94	Oxalate	ADH2	.....	93
95	Perchlorate	ADH2	.....	41
96	Permanganate	CE (reduced to MnO <sub>2</sub> )	.....	41
97	Perrhenate	CE	PDH2, PDFK(3) <sup>5</sup> , PDH4	23,43
98	Phosphate	BE, BDH2	BDFJ(1,2,3), BDH3, BDH4	19,93
99	Silicate	BE, BDH2	BDFJ(2)	34,35,42,91,93
100	Sulfate	BE, BDH2	BDH3, BDH4, BDG	19,26,51

<sup>1</sup>/ Ions may be assumed to follow the same pathways when given parenterally. <sup>2</sup>/ Usually given as soluble complex. <sup>3</sup>/ As judged from the position of the element in the periodic table, or on solubility at neutral pH values. <sup>4</sup>/ Skin. <sup>5</sup>/ Because of inadequate information, no pathways have been listed for cyanate, ferricyanide, and periodate. <sup>6</sup>/ As judged from solubility at neutral pH values.

*continued*



*Continued*

## 54. PATHWAYS OF MINERAL METABOLISM: LABORATORY MAMMALS

Ion	Principal Oral Pathways <sup>1</sup>	Other Known Pathways	Reference
(A)	(B)	(C)	(D)
Anions			
101 Sulfide	ADG (to sulfate)	ADH1	93
102 Thiocyanate	ADH2	ADFK(3), ADH3, ADH4	93
103 Thiosulfate	BE, BDG (to sulfate)	PDH2	93
104 Tungstate	BE, BDH2	BDFK(3, 1, 2), BDH4	24,40,107
105 Vanadate	ADH2	ADFJ(2), ADH4	6,40,93

<sup>1</sup>/ Ions may be assumed to follow the same pathways when given parenterally.

*Contributor:* McChesney, Evan W.

- References:* [1] Adam, K. R., and M. Weatherall. 1954. *J. Pharm. Pharmacol.* 6:403. [2] Aikawa, J. K., et al. 1959. *Am. J. Physiol.* 197:99. [3] Bauer, G. C. H., A. Carlsson, and B. Lindquist. 1955. *Acta Physiol. Scand.* 35:56. [4] Beck, G. 1948. *Mikrochemie Ver. Mikrochim. Acta* 34:62. [5] Bertrand, G., and M. Macheboeuf. 1925. *Bull. Soc. Chim. France* 37:934. [6] Boyd, T. C., and N. K. De. 1933. *Indian J. Med. Res.* 20:789. [7] Bruner, H. D. 1955. *Am. J. Physiol.* 183:600. [8] Bruner, H. D., J. D. Perkinson, and R. L. Hayes. 1953. *Federation Proc.* 12:305. [9] Calhoun, J. A., et al. 1954. *Arch. Ind. Hyg. Occupational Med.* 9:9. [10] Campbell, J. E., E. S. Robadjek, and D. S. Anthony. 1956. *Radiation Res.* 4:294. [11] Carritt, J., et al. 1947. *J. Biol. Chem.* 171:273. [12] Castagnou, R., C. Paoletti, and S. Larcebau. 1957. *Compt. Rend.* 244:2996. [13] Cochran, K. W., et al. 1950. *Arch. Ind. Hyg. Occupational Med.* 1:637. [14] Cole, B. T., and H. Patrick. 1958. *Arch. Biochem. Biophys.* 74:357. [15] Comar, C. L., L. Singer, and G. K. Davis. 1949. *J. Biol. Chem.* 180:913. [16] Crema, A. 1955. *Arch. Intern. Pharmacodyn.* 103:57. [17] Cuthbertson, W. F. J., A. A. Free, and D. M. Thornton. 1950. *Brit. J. Nutr.* 4:42. [18] Danowski, T. S., and J. R. Elkinton. 1951. *Pharmacol. Rev.* 3:42. [19] Davis, G. K., and J. K. Loosli. 1954. *Ann. Rev. Biochem.* 23:459. [20] Decker, C. F., R. U. Byerrum, and C. A. Hoppert. 1957. *Arch. Biochem. Biophys.* 66:140. [21] Dounce, A. L. 1949. In C. Voegtlin and H. C. Hodge, ed. *Natl. Nucl. Energy Ser. VI-1(2):951*. [22] Dudley, H. C., and H. H. Marrer, Jr. 1952. *J. Pharmacol. Exptl. Therap.* 106:129. [23] Durbin, P. W., K. G. Scott, and J. G. Hamilton. 1957. *Univ. Calif. (Berkeley) Publ. Pharmacol.* 3(1):1. [24] Durbin, P. W., et al. 1956. *Proc. Soc. Exptl. Biol. Med.* 91:78. [25] Edstrom, R. 1959. *Acta Psychiat. Neurol. Scand.* 34:26. [26] Everett, N. B., and B. S. Simmons. 1952. *Arch. Biochem. Biophys.* 35:152. [27] Fink, R. M., ed. 1950. *Natl. Nucl. Energy Ser. VI-3*. [28] Fore, H. H., and R. A. Morton. 1952. *Biochem. J.* 51:600. [29] Foreman, J. 1953. *J. Am. Pharm. Assoc. Sci. Ed.* 42:629. [30] Fox, H. M., and H. Ramage. 1931. *Proc. Roy. Soc. (London), B*, 108:157. [31] Gellhorn, A., N. A. Tupikova, and H. B. van Dyke. 1946. *J. Pharmacol. Exptl. Therap.* 87:169. [32] Gensicke, F., H. W. Nitschke, and E. Spode. 1963. *Fortschr. Gebiete Roentgenstrahlen Nuklearmed.* 98:338. [33] Gilbert, I. G. F., and D. M. Taylor. 1956. *Biochim. Biophys. Acta* 21:545. [34] Goodman, L. S., and A. Gilman. 1955. *The pharmacological basis of therapeutics*. Ed. 2. Macmillan, New York. [35] Guelbenzu, M. D., J. M. Lopez de Ancona, and A. Santos. 1951. *Rev. Espan. Fisiol.* 7:63. [36] Hamilton, J. G. 1950. *New Engl. J. Med.* 240:863. [37] Hamilton, J. G., et al. 1953. *Univ. Calif. (Berkeley) Publ. Pharmacol.* 2:283. [38] Hara, R. 1949. *Bull. Chem. Soc. Japan* 22:179,194,225. [39] Harrold, G. C., et al. 1943. *J. Ind. Hyg. Toxicol.* 25:233. [40] Heffter, A., ed. 1927-35. *Handbuch der experimentellen Pharmakologie*. J. Springer, Berlin, Bd. 3. [41] Heffter, A., ed. 1950. *Ibid.* Bd. 10. [42] Holt, P. F., D. M. Yates, and D. H. Tomlin. 1951. *Biochem. J.* 48:xliv. [43] Hurd, L. C., J. K. Colehour, and P. P. Cohen. 1933. *Proc. Soc. Exptl. Biol. Med.* 30:926. [44] Hursh, J. B. 1951. *J. Pharmacol. Exptl. Therap.* 103:451. [45] Hursh, J. B., et al. 1960. *Am. J. Physiol.* 199:513. [46] Jowsey, J., R. E. Rowland, and J. H. Marshall. 1958. *Radiation Res.* 8:490. [47] Kittle, C. F., et al. 1951. *Proc. Soc. Exptl. Biol. Med.* 76:278. [48] Kleeman, C. R., and F. H. Epstein. 1956. *Ibid.* 93:228. [49] Kleeman, C. R., et al. 1955. *Am. J. Physiol.* 182:548. [50] Kraintz, L., and R. V. Talmage. 1952. *Proc. Soc. Exptl. Biol. Med.* 81:490. [51] Kulwich, R., L. Struglia, and P. B. Pearson. 1957.

*continued*

*Continails*

**54. PATHWAYS OF MINERAL METABOLISM: LABORATORY MAMMALS**


- J. Nutr. 61:113. [52] Kurlyandskaya, E. B., N. L. Beloborodova, and E. F. Baranova. 1958. Chem. Abstr. 52:3078.
- [53] Lang, F. R. 1951. Ann. Internal Med. 35:1237. [54] Lang, H., Jr., P. C. Wallace, and J. G. Hamilton. 1950. Univ. Calif. (Berkeley) Publ. Pharmacol. 2:263. [55] Lass, A., E. Scharpff, and K. Wunderlich. 1955. Klin. Wochschr. 33:959. [56] Lee, C. C. 1953. Federation Proc. 12:84. [57] Lomholt, S. 1924. Biochem. J. 18:693.
- [58] Lowry, O. H., et al. 1942. J. Pharmacol. Exptl. Therap. 76:221. [59] Lund, A. 1956. Acta Pharmacol. Toxicol. 12:251. [60] McClure, F. J. 1949. Ann. Rev. Biochem. 18:335. [61] McConnell, K. P., and R. G. Martin. 1952. J. Biol. Chem. 194:183. [62] McConnell, K. P., and O. W. Portman. 1952. Ibid. 195:277. [63] MacKenzie, R. D., et al. 1959. Arch. Biochem. Biophys. 79:200. [64] Mahoney, J. P., et al. 1955. J. Lab. Clin. Med. 46:702.
- [65] Maynard, L. A., and S. E. Smith. 1947. Ann. Rev. Biochem. 16:273. [66] Maynard, L. S., and G. C. Cotzias. 1955. J. Biol. Chem. 214:489. [67] Meek, S. F., G. C. Harrold, and C. P. McCord. 1943. Ind. Med. 12:447.
- [68] Mendel, L. B., and O. E. Closson. 1906. Am. J. Physiol. 16:152. [69] Mendel, L. B., and D. F. Sicher. 1906. Ibid. 16:147. [70] Miwa, M., and H. Yamashita. 1938. Gann 32:395. [71] Mraz, F. R., and H. Patrick. 1957. Proc. Soc. Exptl. Biol. Med. 94:409. [72] Mraz, F. R., and H. Patrick. 1957. Arch. Biochem. Biophys. 71:121.
- [73] Mraz, F. R., and H. Patrick. 1957. J. Nutr. 61:535. [74] Mraz, F. R., et al. 1957. Arch. Biochem. Biophys. 66:177. [75] Norris, W. P., and W. Kiesleski. 1948. Cold Spring Harbor Symp. Quant. Biol. 13:164.
- [76] Perey, M., and A. Chevallier. 1951. Compt. Rend. Soc. Biol. 145:1205. [77] Perrault, M., and F. Chain. 1958. Presse Med. 66:1394. [78] Princi, F., and E. F. Geever. 1950. Arch. Ind. Hyg. Occupational Med. 1:651.
- [79] Rosenfeld, G. 1954. Arch. Biochem. Biophys. 48:84. [80] Sastry, B. V., R. L. Weiland, and C. O. T. Ball. 1963. Federation Proc. 22:540. [81] Schou, M. 1957. Pharmacol. Rev. 9:17. [82] Schubert, J., and M. R. White. 1952. J. Lab. Clin. Med. 39:260. [83] Schubert, J., M. R. White, and A. Lindenbaum. 1952. J. Biol. Chem. 196:279. [84] Schubert, J., et al. 1950. Ibid. 182:635. [85] Scott, J. K., W. F. Neuman, and J. F. Bonner. 1952. J. Pharmacol. Exptl. Therap. 106:286. [86] Scott, K. G., D. J. Axelrod, and J. G. Hamilton. 1949. J. Biol. Chem. 177:325. [87] Scott, K. G., and J. G. Hamilton. 1950. Univ. Calif. (Berkeley) Publ. Pharmacol. 2:241. [88] Scott, K. G., et al. 1948. J. Biol. Chem. 175:691. [89] Scott, K. G., et al. 1948. Ibid. 176:283. [90] Shapiro, R. 1956. Acta Radiol. 46:635. [91] Shohl, A. T. 1939. Mineral metabolism. Reinhold, New York. [92] Singer, L., et al. 1957. Arch. Biochem. Biophys. 66:404. [93] Sollman, T. 1957. A manual of pharmacology. W. B. Saunders, Philadelphia. [94] Spode, E., and F. Gensicke. 1958. Naturwissenschaften 45:117. [95] Spode, E., and F. Gensicke. 1958. Ibid. 45:135. [96] Stannard, J. N., and F. A. Smith. 1957. Univ. Rochester At. Energy Proj. Bull. UR-287. [97] Stover, B. J., D. R. Atherton, and J. S. Arnold. 1957. Proc. Soc. Exptl. Biol. Med. 94:269. [98] Tedeschi, R. E., and F. W. Sunderman. 1957. Arch. Ind. Hyg. Occupational Med. 16:486. [99] Ter Meulen, H. 1931. Rec. Trav. Chim. 50:491. [100] Thompson, R. C., and O. L. Hollis. 1956. Hanford At. Prod. Oper. Bull. HW-45546. [101] Thompson, R. C., et al. 1958. Am. J. Roentgenol. Radium Therapy Nucl. Med. 79:1026. [102] Tregubenko, I. P. 1961. Chem. Abstr. 55:1922. [103] Underwood, E. J. 1959. Ann. Rev. Biochem. 28:499. [104] Vallee, B. L. 1959. Physiol. Rev. 39:443. [105] Vallee, B. L., and R. G. Fluharty. 1947. J. Clin. Invest. 26:1199. [106] Visek, W. J., et al. 1953. Proc. Soc. Exptl. Biol. Med. 84:610. [107] Wase, A. W. 1956. Arch. Biochem. Biophys. 61:272. [108] Wase, A. W., D. M. Goss, and M. J. Boyd. 1954. Ibid. 51:1. [109] White, M. R., and J. Schubert. 1952. J. Pharmacol. Exptl. Therap. 104:317. [110] Widdowson, E. M., and R. A. McCance. 1944. Proc. Nutr. Soc. (Engl. Scot.) 1:220. [111] Widdowson, E. M., R. A. McCance, and C. M. Spray. 1951. Clin. Sci. 10:113.





## 56. PATHWAYS OF CARBOHYDRATE METABOLISM

The conversion of stored or ingested carbohydrate to pyruvate releases stored energy by means of anaerobic oxidation (glycolysis). Released energy is partly dissipated as heat and partly stored (temporarily) in the labile energy pool as high-energy phosphate ( $-PO_4$ ) with continuously available ADP (adenosine diphosphate) to form ATP (adenosine triphosphate). In the conversion of 1 mole of glucose (180 g), or of other monosaccharides, to 2 moles of pyruvate (174 g), 2 moles of ATP are converted to ADP and 4 moles of ATP are formed from ADP, making a net gain of 2 moles of ATP, or approximately 14 kilocalories of readily available energy. If glucose-6- $PO_4$  has come from the metabolic breakdown of glycogen, the cost is only 1 mole of ATP, making a net gain of 3 moles of ATP (approximately 21 kilocalories). The ATP is an immediate source of energy, the utilization of which (e.g., for muscular activity) is independent of oxygen supply. Aerobic oxidation of the reduced coenzymes formed in glycolysis and in the Krebs cycle yields considerable amounts of energy. The reactions may be summarized as follows:  $glucose + 2 ATP \rightarrow 2 pyruvate + 8 ATP$ ;  $2 pyruvate \rightarrow 2 acetyl CoA + 6 ATP$ ;  $2 acetyl CoA \rightarrow 4 CO_2 + 4 H_2O + 24 ATP$ ; a total of 38 ATP. The complete oxidation of glucose to  $CO_2$  and water releases 685.5 kilocalories of energy. The ADP system traps 266 ( $38 \times 7$ ) kilocalories of this energy, resulting in a possible storage of 39% ( $266/685$ ) of the energy in the form of ATP.



continued



## 56. PATHWAYS OF CARBOHYDRATE METABOLISM

/1/ Adenylic acid and  $PO_4$  required for activity in either direction. /2/ Digestion; glycogen and/or starch are hydrolyzed to glucose in intestinal lumen. /3/  $Mg^{++}$  required for this reaction. /4/ Cori ester. /5/ Robison ester. /6/ Hexokinase reaction assumed to be inhibited by growth hormone plus adrenal cortex hormone; inhibition by these substances is blocked by insulin, thus favoring conversion of glucose to glucose-6-phosphate. /7/ The reaction, glycogen to glucose-6- $PO_4$  to blood glucose, takes place in liver only; conversion of glucose to glucose-6- $PO_4$  to glycogen takes place in liver, muscle, and other tissues. /8/ Neuberger ester. /9/ In liver and muscle. /10/ In all tissues. /11/ Hardin-Young ester. /12/ This reaction (to left) causes each step in the conversion to pyruvate to be doubled quantitatively; thus, 1 mole of glucose gives rise to 2 moles of pyruvate. /13/ Hydrogen atoms released. /14/ Hydrogen enters into the reaction. /15/  $NAD^+$  acts as acceptor of released hydrogen atoms, becoming NADH in oxidative direction of the reaction. NADH gives up hydrogen atoms and becomes  $NAD^+$  in reverse direction. Hydrogen atoms accepted by  $NAD^+$  are passed on in turn to flavoprotein, cytochrome-c, cytochrome oxidase, and molecular  $O_2$ . If molecular  $O_2$  is not sufficiently available, hydrogen atoms may be passed from NADH to pyruvate-forming lactate. /16/ Inhibited by fluoride. /17/  $K^+$  also required. /18/ Thiamine pyrophosphate required as coenzyme. /19/ Pyruvate, followed by conversion to lactate when oxygen supply is deficient (*see* Fn. 15), ends glycolysis in animal tissues. If oxygen is available, pyruvate is oxidized via the Krebs cycle. /20/ End of fermentation of plant tissue. /21/ Uridine diphosphate glucose required as coenzyme. /22/ Inhibited by iodoacetate.

**Contributors:** (a) Bishop, David W., (b) Bonner, James F., (c) Van Bruggen, John T., (d) Roe, Joseph H.

**References:** [1] Axelrod, B. 1961. In D. M. Greenberg, ed. *Metabolic pathways*. Academic Press, New York. [2] Baldwin, E. 1957. *Dynamic aspects of biochemistry*. Ed. 3. Cambridge Univ. Press, New York. [3] Cori, G. T., et al. 1951. *Biochim. Biophys. Acta* 7:304. [4] Dickens, F. 1951. In J. B. Sumner and K. Myrbäck, ed. *The enzymes*. Academic Press, New York. v. 2, p. 624. [5] Lardy, H. A., ed. 1949. *Respiratory enzymes*. Burgess, Minneapolis. [6] Krebs, H. A. 1949. *Advan. Enzymol.* 3:191. [7] Sumner, J. B., and G. F. Somers. 1953. *Chemistry and methods of enzymes*. Ed. 3. Academic Press, New York. [8] Umbreit, W. W. 1952. *Metabolic maps*. Burgess, Minneapolis. [9] Werkman, C. H., and F. Schlenk. 1951. In C. H. Werkman and P. W. Wilson, ed. *Bacterial physiology*. Academic Press, New York. p. 281. [10] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York.

## 57. PATHWAYS OF AMINO ACID METABOLISM

Amino Acid	Product of Oxidative Deamination or Transamination	Product of Decarboxylation	Pathways and Products of Metabolism
(A)	(B)	(C)	(D)
1 L-Alanine	Pyruvic acid		
2 L-Arginine	$\alpha$ -Keto- $\delta$ -guanidovaleric acid	Agmatine	Arginine $\rightarrow$ ornithine + urea; arginine $\rightarrow$ citrulline + $NH_3$ ; arginine $\rightarrow$ agmatine + $CO_2$ ; arginine + glycine $\rightleftharpoons$ guanidoacetic acid + ornithine.
3 L-Asparagine	$\alpha$ -Ketosuccinamic acid		Asparagine $\rightleftharpoons$ aspartic acid + $NH_3$ ; asparagine $\rightarrow$ $\alpha$ -ketosuccinamic acid $\rightarrow$ $NH_3$ + oxalacetic acid.
4 L-Aspartic acid	Oxalacetic acid	$\alpha$ -Alanine, $\beta$ -alanine	Aspartic acid + carbamylphosphate $\rightarrow$ $PO_4^{=}$ + carbamylaspartic acid $\rightarrow$ pyrimidines; aspartic acid $\rightleftharpoons$ fumaric acid + $NH_3$ ; aspartic acid $\rightleftharpoons$ homoserine $\left\{ \begin{array}{l} \text{threonine} \\ \text{methionine} \end{array} \right.$
5 L-Citrulline	$\alpha$ -Keto- $\delta$ -carbamidovaleric acid		Citrulline + aspartic acid + ATP $\rightarrow$ AMP + PP + arginosuccinic acid $\rightleftharpoons$ arginine + fumaric acid; citrulline + $PO_4^{=}$ $\rightleftharpoons$ ornithine + carbamylphosphate; citrulline $\rightarrow$ carbamylphosphate + ADP $\rightleftharpoons$ $CO_2$ + $NH_3$ + ATP.
6 L-Cysteine & L-cystine	$\beta$ -Mercaptopyruvic acid		Cysteine $\rightarrow$ $\beta$ -mercaptopyruvic acid $\rightarrow$ pyruvic acid + S; cysteine $\rightarrow$ $H_2S$ + $NH_3$ + pyruvic acid; cysteine $\rightarrow$ cysteine sulfonic acid $\rightarrow$ (i) cysteic acid $\rightarrow$ taurine, (ii) hypotaurine, or (iii) via transamination $\rightarrow$ $\beta$ -sulfinylpyruvate $\rightarrow$ pyruvate + $SO_3$ . (2 cysteine $\rightleftharpoons$ cystine)
7 L-Glutamic acid	$\alpha$ -Ketoglutaric acid	$\gamma$ -Aminobutyric acid	Glutamic acid $\rightarrow$ $\gamma$ -aminobutyric acid + $CO_2$ . <i>See also</i> ornithine, proline, histidine, glutamine.
8 L-Glutamine	$\alpha$ -Ketoglutaramic acid		Glutamine $\rightleftharpoons$ glutamic acid + $NH_3$ ; glutamine $\rightarrow$ $\alpha$ -ketoglutaramic acid $\rightarrow$ $NH_3$ + $\alpha$ -ketoglutaric acid.

*continued*

**57. PATHWAYS OF AMINO ACID METABOLISM**


Amino Acid	Product of Oxidative Deamination or Transamination	Product of Decarboxylation	Pathways and Products of Metabolism
(A)	(B)	(C)	(D)
9 Glycine	Glyoxylic acid		Glycine + N <sup>5</sup> -10-methenyl-tetrahydrofolate ⇌ serine + tetrahydrofolate; glycine → glyoxylic acid → formate + CO <sub>2</sub> ; glycine + succinyl-CoA → δ-aminolevulinic acid → porphyrins; glycine + 5'-phospho-β-D-ribosamine → purines.
10 L-Histidine	β-Imidazolepyruvic acid	Histamine	Histidine → urocanic acid → glutamic acid + NH <sub>3</sub> + formate; histidine → carnosine; histidine → anserine; histidine → histamine → imidazoleacetic acid → NH <sub>3</sub> + formylaspartic acid.
11 L-Hydroxyproline	α-Keto-γ-hydroxy-δ-aminovaleric acid		Hydroxyproline → Δ <sup>1</sup> -proline-4-hydroxy-2-carboxylate → pyrrolidine-2-carboxylate; hydroxyproline → glutamate.
12 L-Isoleucine	d-α-Keto-β-methylvaleric acid		Isoleucine → α-keto-β-methylvaleric acid → CO <sub>2</sub> + α-methylbutyryl-CoA ⇌ tiglyl-CoA ⇌ α-methyl-β-hydroxybutyryl-CoA ⇌ α-methyl acetoacetyl-CoA ⇌ acetyl-CoA + propionyl-CoA.
13 L-Leucine	α-Ketoisocaproic acid		Leucine → α-ketoisocaproic acid → CO <sub>2</sub> + isovaleryl-CoA ⇌ seneciyl-CoA + CO <sub>2</sub> ⇌ β-methylglutaconyl-CoA ⇌ β-hydroxy-β-methyl glutaryl-CoA ⇌ acetoacetic acid + acetyl-CoA.
14 L-Lysine	α-Keto-ε-aminocaproic acid	Cadaverine	Lysine → α-keto-ε-aminocaproic acid → Δ <sup>1</sup> -piperidine-2-carboxylic acid → pipercolic acid → Δ <sup>6</sup> -piperidine-2-carboxylic acid → α-aminoadipic-ε-semialdehyde → L-α-aminoadipic acid → α-ketoadipic acid → glutamic acid → α-ketoglutaric acid → L-glutamic acid.
15 L-Methionine	α-Keto-γ-methylbutyric acid		Methionine → labile CH <sub>3</sub> + homocysteine → (i) homocysteic acid, (ii) H <sub>2</sub> S + NH <sub>3</sub> + α-ketobutyric acid, or (iii) serine → cystathionine → cysteine + NH <sub>3</sub> → α-ketobutyric acid.
16 L-Ornithine	Glutamic-γ-semialdehyde, or α-keto-δ-aminovaleric acid	Putrescine	Ornithine ⇌ proline; ornithine ⇌ glutamic acid; ornithine + carbamylphosphate → citrulline. <i>See also</i> citrulline.
17 L-Phenylalanine	Phenylpyruvic acid	Phenylethylamine	Phenylalanine → tyrosine; phenylalanine → phenylpyruvic acid → phenylacetic and phenyllactic acids.
18 L-Proline	Glutamic-γ-semialdehyde, or α-keto-δ-aminovaleric acid		Proline ⇌ ornithine; proline ⇌ glutamic acid; proline → hydroxyproline.
19 L-Serine	β-Hydroxyppyruvic acid	Ethanolamine	Serine → NH <sub>3</sub> + H <sub>2</sub> O + pyruvic acid; serine + indole-3-glycerol phosphate ⇌ tryptophan. <i>See also</i> glycine.
20 L-Threonine	d-α-Keto-β-hydroxybutyric acid		Threonine → NH <sub>3</sub> + H <sub>2</sub> O + α-ketobutyric acid; threonine → glycine + acetaldehyde; threonine → α-keto-β-hydroxybutyric acid; threonine → aminoacetone. <i>See also</i> aspartic acid.
21 L-Tryptophan	β-Indolepyruvic acid	Tryptamine	Tryptophan → formylkynurenine → formate + kynurenine → (i) kynurenic acid, (ii) anthranilic acid + alanine, or (iii) 3-hydroxykynurenine → 3-hydroxyanthranilic acid → 2-acrolein-3-aminofumaric acid → quinolinic acid → nicotinic acid ribonucleotide; tryptophan → 5-hydroxytryptophan → 5-hydroxytryptamine → 5-hydroxyindoleacetic acid.
22 L-Tyrosine	β-Hydroxyphenylpyruvic acid	Tyramine	Tyrosine → β-hydroxyphenylpyruvic acid → CO <sub>2</sub> + homogentisic acid + maleylacetoacetic acid → fumarylacetoacetic acid → fumaric acid + acetoacetic acid.
23 L-Valine	α-Ketoisovaleric acid		Valine → α-ketoisovaleric acid → CO <sub>2</sub> + isobutyryl-CoA ⇌ methacrylyl-CoA ⇌ β-hydroxybutyryl-CoA ⇌ β-hydroxyisobutyric acid ⇌ methylmalonic acid semialdehyde ⇌ β-aminoisobutyric acid.

*Contributors:* (a) Meister, Alton, (b) Sallach, H. J., (c) Elwyn, David H., (d) Richert, Dan A., (e) Turner, Robert A.

*References:* [1] McElroy, W. D., and B. Glass, ed. 1955. Amino acid metabolism. Johns Hopkins Press, Baltimore. [2] Meister, A. 1953. In G. H. Bourne and G. W. Kidder, ed. Biochemistry and physiology of nutrition. Academic Press, New York. v. 1, p. 187. [3] Meister, A. 1964. Biochemistry of the amino acids. Ed. 2. Academic Press, New York. [4] West, E. S., and W. R. Todd. 1961. Textbook of biochemistry. Ed. 3. Macmillan, New York.

## 58. PATHWAYS OF NUCLEOPROTEIN CATABOLISM

See Table 59 for detailed pathways of purine and pyrimidine nucleotide catabolism.



/1/ Catabolism of nucleoprotein, nucleic acid, and protein may take place in the alimentary canal or in the tissues. /2/ Little intestinal absorption. /3/ In the biosynthesis of nucleotides, inosinic acid is the precursor of adenylic and guanylic acids; in uricotelic species, a portion is a direct precursor of uric acid. /4/ Absorbed from the intestine. Purine nucleosides are split into purines and pentoses by purine nucleosidase present in tissues. /5/ D-Ribose and D-2-deoxyribose are the sugars typical of the two types of nucleic acids. /6/ Mammals do not require but can synthesize exogenous purines or pyrimidines from products of protein metabolism. /7/ Adenine and guanine are the major purines occurring in nucleic acids. /8/ The route adenine → hypoxanthine is of no importance in animals. Adenase is not found to any extent in mammals. /9/ NH<sub>3</sub>, as in the case of CO<sub>2</sub>, is also used to synthesize many tissue constituents; hence, it may enter into metabolic processes, be built into amino acids, incorporated into urea and excreted, or excreted as NH<sub>3</sub> across the kidney tubule. /10/ Excreted by pig and spider. /11/ Cytosine is converted to uracil at the nucleotide stage. Free cytosine is excreted unchanged. /12/ Excreted by most fishes, amphibians, and freshwater lamellibranchs. /13/ Excreted by mammals as the end product of amino acid metabolism. Excreted by some animals as the end product of purine and pyrimidine metabolism. /14/ Excreted by primates, some reptiles, and some insects as the end product of purine catabolism. Excreted by birds as the end product of protein, purine, and pyrimidine catabolism; no urea formation by birds. /15/ Excreted by most mammals, gastropods, and some insects. /16/ Excreted by some teleost fishes. /17/ May enter into metabolic processes, into the ornithine cycle and be incorporated into and excreted as urea, or be excreted as CO<sub>2</sub>. /18/ Urea formation in mammalian liver occurs via the ornithine cycle (Krebs-Henseleit cycle): ornithine → citrulline → arginine succinate → arginine → ornithine. CO<sub>2</sub> and NH<sub>3</sub> enter the cycle via carbamyl glutamic acid at ornithine; NH<sub>3</sub> enters the cycle via aspartic acid at citrulline. Arginine succinate is split to arginine and fumaric acid; arginine is then converted to ornithine with the release of urea. /19/ Via Krebs cycle. In the course of amino acid metabolism, prior to entry into the Krebs cycle, sulfur-containing amino acids lose their sulfur—usually in the form of SO<sub>4</sub>. /20/ Crustacea, gephyrean worms, and marine lamellibranchs do not excrete urea but break it down to, and excrete it as, CO<sub>2</sub> and NH<sub>3</sub>.


**Contributors:** (a) Brown, George B., (b) Elwyn, David H., (c) Richert, Dan A., (d) Bishop, David W.

**References:** [1] Baldwin, E. 1957. Dynamic aspects of biochemistry. Ed. 3. Cambridge Univ. Press, New York. [2] Chargaff, E., and J. N. Davidson, ed. 1955-60. The nucleic acids. Academic Press, New York. [3] Davidson, J. N. 1960. Biochemistry of the nucleic acids. Ed. 4. Methuen, London.



## 59. PATHWAYS OF PURINE AND PYRIMIDINE CATABOLISM

Nucleoproteins (see Table 58) generally are composed of basic proteins (histones or protamines) associated with nucleic acids. The nucleic acids are complex molecules (each composed of many nucleotide units) joined by phosphate sugar linkage. The nucleotides shown below are obtained by enzymic hydrolysis of nucleic acids, although several may be obtained from other sources. Each nucleotide is composed of a purine or pyrimidine base linked to a pentose sugar which, in turn, is linked to phosphate. In catabolism, the phosphate is removed by a nucleotide phosphatase to yield inorganic phosphate and a nucleoside; the nucleoside is cleaved by a nucleoside phosphorylase to give the free base plus ribose-1-phosphate. Also, as a catabolic step, hydrolytic deamination may occur at any level, although it is most frequent at the nucleoside level.




/1/ Purine. /2/ Pyrimidine. /3/ Demonstrated in yeast and *Escherichia coli*. /4/ End product of primates, birds, terrestrial reptiles, and most insects. /5/ End product of mammals other than primates. /6/ End product of some teleost fishes. /7/ End product of most fishes, amphibians, and freshwater lamellibranchs.

**Contributors:** (a) Barrett, Harold W., (b) Lansford, Edwin M., Jr., and Shive, William, (c) Zbarsky, S. H.

**References:** [1] Baldwin, E. 1957. Dynamic aspects of biochemistry. Ed. 3. Cambridge Univ. Press, New York. [2] Cantarow, A., and B. Schepartz. 1962. Biochemistry. Ed. 3. W. B. Saunders, Philadelphia. [3] Chargaff, E., and J. N. Davidson. 1955-60. The nucleic acids. Academic Press, New York. [4] Colowick, S. P., and N. O. Kaplan. 1955. Methods in enzymology. Academic Press, New York. v. 2. [5] Davidson, J. N. 1960. Biochemistry of the nucleic acids. Ed. 4. Methuen, London. [6] West, E. S., and W. R. Todd. 1961. Textbook of biochemistry. Ed. 3. Macmillan, New York. [7] White, A., et al. 1959. Principles of biochemistry. McGraw-Hill, New York.



## 60. METABOLIC INTERRELATIONSHIPS: CARBOHYDRATE, FAT, AND PROTEIN




/1/ Phosphorylation of hexose units in stored polysaccharides by phosphorylase and phosphate; phosphorylation of hexoses by hexokinase and ATP. /2/ Proteolysis by proteases in digestive tract or tissues. /3/ Lipase splits fat into fatty acids and glycerol. (Glycerol, via glycerol phosphate and dihydroxyacetone phosphate, enters the glycolytic cycle. Fatty acid then is acted upon by coenzyme A.) /4/ Glycolysis. /5/  $\beta$ -Oxidation. /6/ Oxidative deamination. /7/ Transamination. /8/ Oxidative decarboxylation. /9/ Carboxylation. /10/ Chain of electron-transmitting enzymes. /11/ High energy phosphorus.

**Contributors:** (a) Bonner, James F., and Saltman, Paul, (b) Van Bruggen, John T., (c) Elwyn, David H., (d) Welt, Isaac D.

**References:** [1] Baldwin, E. 1957. Dynamic aspects of biochemistry. Ed. 3. Cambridge Univ. Press, New York. [2] Fruton, J. S., and S. Simmonds. 1958. General biochemistry. Ed. 2. J. Wiley, New York. [3] Greenberg, D. M. 1960-61. Metabolic pathways. Academic Press, New York. [4] West, E. S., and W. R. Todd. 1961. Text-book of biochemistry. Ed. 3. Macmillan, New York. [5] White, A., et al. 1959. Principles of biochemistry. Ed. 2. McGraw-Hill, New York.

## 61. KREBS CYCLE

The Krebs cycle (tricarboxylic acid cycle) is a major pathway for the final aerobic oxidation of carbohydrates, fats, and proteins, which are channeled into the cycle via their two key metabolites, pyruvate and acetyl-CoA (active acetate). Each "revolution" of the cycle oxidizes acetate to  $\text{CO}_2$  and  $\text{H}_2\text{O}$ . One mole (59 g) of acetate thus oxidized releases approximately 200 kilocalories of energy. A portion of the released energy (approximately 144 kilocalories) enters the phosphate pool as ATP. Twelve moles of ATP are formed from ADP and  $\text{PO}_4$  (by energizing  $\text{PO}_4$  to  $-\text{PO}_4$ ). The remainder of the released energy appears as heat. Oxidation of 1 mole (87 g) of pyruvate, via acetyl-CoA, contributes a total of 14 moles of ATP to the energy pool. Heavy lines indicate main sequence of reactions.



*continued*

## 61. KREBS CYCLE


<sup>/1/</sup> Glucogenic amino acid precursors for pyruvate are alanine, glycine, serine, threonine, methionine, cysteine, valine. <sup>/2/</sup> Ketogenic amino acid precursors for acetyl-CoA are leucine, isoleucine, phenylalanine, tyrosine. <sup>/3/</sup> Aspartic acid occurs as a component of protein. <sup>/4/</sup> Glutamic acid occurs as a component of protein or may be formed from arginine, proline, hydroxyproline, histidine, ornithine. <sup>/5/</sup> Coenzyme A (ATP-pantoyl- $\beta$ -alanyl-thioethanolamine) and  $\alpha$ -lipoic acid required. <sup>/6/</sup> In the oxidative direction, NAD<sup>+</sup> (nicotinamide adenine dinucleotide) acts as hydrogen acceptor; in the reverse direction, NADH is hydrogen donor. <sup>/7/</sup> In the oxidative direction, NADP<sup>+</sup> (nicotinamide adenine dinucleotide phosphate) acts as hydrogen acceptor; in the reverse direction, NADPH is hydrogen donor. <sup>/8/</sup> Cocarboxylase (thiamine pyrophosphate) required as coenzyme for the carboxylase (A); also Mg<sup>++</sup> or Mn<sup>++</sup> required as activator for the enzyme. <sup>/9/</sup> Biotin required as coenzyme for decarboxylation. <sup>/10/</sup> Two moles of hydrogen enter into the reaction. <sup>/11/</sup> Two moles of hydrogen released and their electrons transferred to cytochrome. <sup>/12/</sup> Hydrogen atoms transferred to NAD<sup>+</sup> (or, in the case of isocitrate  $\rightarrow$  oxalsuccinate, to NADP<sup>+</sup>) and pass in turn to flavoprotein, cytochrome-c, cytochrome oxidase, and finally to combination with molecular oxygen. For each mole of hydrogen thus passed and finally oxidized, 1.5 moles of ATP are formed by the addition of energized phosphate (-PO<sub>4</sub>) to ADP. <sup>/13/</sup> CO<sub>2</sub> enters into the reaction. <sup>/14/</sup> CO<sub>2</sub> released. <sup>/15/</sup> H<sub>2</sub>O enters into the reaction. <sup>/16/</sup> H<sub>2</sub>O released. <sup>/17/</sup> NH<sub>3</sub> enters into the reaction by transamination. <sup>/18/</sup> NH<sub>3</sub> transferred from glutamate by transamination, then enters into Krebs cycle via  $\alpha$ -ketoglutarate. <sup>/19/</sup> Footnotes 5, 6, 8, 12, 14 apply to this reaction. <sup>/20/</sup> For details, see Table 62 on the cytochrome system.

**Contributors:** (a) Bishop, David W., (b) Bonner, James F., (c) Roe, Joseph H., (d) Welt, Isaac D.

**References:** [1] Artom, C. 1953. *Ann. Rev. Biochem.* 22:211. [2] Baldwin, E. 1957. *Dynamic aspects of biochemistry*. Ed. 3. Cambridge Univ. Press, New York. p. 415. [3] Black, K. 1952. *Ann. Rev. Biochem.* 21:273. [4] Dickens, F. 1951. In J. B. Sumner and K. Myrbäck, ed. *The enzymes*. Academic Press, New York. v. 2, p. 624. [5] Evans, E. A., Jr. 1944. *Ann. Rev. Biochem.* 13:187. [6] Frazer, A. C. 1952. *Ibid.* 21:245. [7] Fruton, J. S., and S. Simmonds. 1958. *General biochemistry*. Ed. 2. J. Wiley, New York. [8] Greenberg, D. M. 1960-61. *Metabolic pathways*. Academic Press, New York. [9] Krebs, H. A. 1943. *Advan. Enzymol.* 3:191. [10] Ochoa, S. 1951. *Physiol. Rev.* 31:56. [11] Ochoa, S., and J. R. Stern. 1952. *Ann. Rev. Biochem.* 21:547. [12] Potter, V. R., and C. Heidelberger. 1950. *Physiol. Rev.* 30:487. [13] Umbreit, W. W. 1952. *Metabolic maps*. Burgess, Minneapolis. p. 90. [14] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York. [15] White, A., et al. 1959. *Principles of biochemistry*. Ed. 2. McGraw-Hill, New York.

## 62. CYTOCHROME SYSTEM

The cytochromes (iron-containing compounds) in association with certain enzymes comprise the cytochrome system. The system operates as the final pathway by which an intermediate metabolite (substrate), under the influence of its specific dehydrogenase, releases hydrogen to the first member in a series of carriers for ultimate combination with oxygen to form water. Each step in the process involves both oxidation and reduction: the cytochrome system oxidizes the hydrogen of the substrate by removing electrons from it, thereby producing oxidized substrate and hydrogen ions. The system itself is reduced in the process and is finally oxidized by molecular oxygen. For each gram of hydrogen thus passed to NADH and finally oxidized, enough energy is produced to form 1.5 moles of ATP from ADP and PO<sub>4</sub>.



**Contributor:** Wainio, Walter W.

**References:** [1] Chance, B. 1961. In J. E. Falk, ed. *Haematin enzymes*. Pergamon Press, Oxford. p. 597. [2] Green, D. E. 1961. *Ciba Found. Symp. Quinones Electron Transport*, 1960, p. 130. [3] Okunuki, K., et al. 1958. *Proc. Intern. Symp. Enzyme Chem., Tokyo-Kyoto, 1957*, p. 264. [4] Slater, E. C. 1958. *Advan. Enzymol.* 20:147. [5] Slater, E. C. 1961. In J. E. Falk, ed. *Haematin enzymes*. Pergamon Press, Oxford. p. 575. [6] Wainio, W. W. 1953. *Rutgers Univ. Bur. Biol. Res. Symp. Some Conjugated Proteins*, p. 19. [7] Wainio, W. W. 1961. In J. E. Falk, ed. *Haematin enzymes*. Pergamon Press, Oxford. p. 281.

**63. PROPERTIES OF CYTOCHROMES: ANIMALS AND HIGHER PLANTS**

The cytochromes of animals and higher plants are intracellular chromoproteins which are entirely associated with lipoprotein structural elements of cytoplasm. The prosthetic group contains coordinated iron which may undergo alternate oxidation and reduction. **Physical and Chemical Properties** (column C): PG = prosthetic group; MW = molecular weight;  $E'_O$  = oxidation-reduction potential. **Spectral Characteristics** (columns D and E):  $\lambda$  maximum in  $m\mu$  = wave length of maximum absorption; figures in parentheses are  $E'_{1\text{cm}}^{m\mu}$ , i.e., extinction coefficients of millimolar solutions of 1-cm thickness.

Source of Pigments (A)	Cytochrome (B)	Physical and Chemical Properties (C)	Spectral Characteristics $\lambda$ maximum in $m\mu$		Remarks (F)	Reference (G)	
			Reduced (D)	Oxidized (E)			
<b>Animals</b>							
1	Mitochondria <sup>1</sup>	<i>a</i>	PG = hematin- <i>a</i> ; $E'_O = 0.29$ volts	603 452	590-600 418-420	Closely associated with cytochrome- <i>a</i> <sub>3</sub>	B,3,19,25,35; C,3,25;D,19; E,35;F,29
2		<i>a</i> <sub>3</sub>		604 448	590-600 418-420	Carbon monoxide- <i>a</i> <sub>3</sub> complex, when reduced, has absorption bands at 590-593 $m\mu$ and at 432 $m\mu$ .	B,D,19;E,35; F,6,19
3		<i>b</i>	PG = protohematin; $E'_O = -0.04$ volts	564((20.8) 530 430	416	May not be on direct electron transfer path	B,D,16,31;C, 3,16,31;E,16; F,29
4		<i>c</i>	MW = 12,200; PG = hematin- <i>c</i> ; Fe = 0.45%; heat stable at neutral pH; isoelectric point = pH 10.5-10.8; $E'_O = 0.255$ volts	550 (27.8) 520 415 316	530 408 355	Data refer to pigment purified from horse heart. Fe content may be increased by enrichment with iron-containing impurities.	B,11,13,17,18, 22,34;C,13, 17,18,34;D, E,11,22;F, 11,13,22,34
5		<i>c</i> <sub>1</sub>	PG = hematin- <i>c</i>	553-554 522-524 416-418	410	Identical with cytochrome- <i>e</i>	B,D,20,30;C, E,30;F,20,21
6	Endoplasmic reticulum <sup>2</sup>	<i>b</i> <sub>5</sub>	MW = 16,900; PG = protohematin; non-mitochondrial oxidation of NADH and NADPH; $E'_O = 0.12$ volts	556 (25.6) 526 423 320-340	500-580 413 355-370	Identical with cytochrome- <i>m</i> . Separate flavoproteins catalyze the reactions of NADH and of NADPH with cytochrome- <i>b</i> <sub>5</sub> in animal tissues.	B,D,E,33;C,2, 33;F,2,32
7	Intracellular <sup>3</sup>	<i>h</i>	MW = 18,500; PG = modified protohematin; Fe = 0.33%; heat stable at neutral pH; isoelectric point = <pH 4.3; $E'_O = 0.20$ volts	556 (22.7) 526.5 422	562 536 408	Occurs in land snails and other invertebrates. Helicorubin is probably a degraded form of cytochrome- <i>h</i> .	B,C,F,23,24; D,E,23,29
<b>Higher Plants</b>							
8	Mitochondria <sup>4</sup>	<i>a</i>	As found in animal mitochondria. Presence indicated by spectroscopic and spectrophotometric observations.				5,12,27,28
9		<i>a</i> <sub>3</sub>	As found in animal mitochondria. Presence indicated by spectroscopic and spectrophotometric observations.			Role in electron transport inferred from cyanide inhibition of respiration, and from shift in spectrum observed with carbon monoxide	B-E,5,12,27, 28;F,12,26, 28
10		<i>b</i>	As found in animal mitochondria. Presence indicated by spectroscopic and spectrophotometric observations.				5,12,27,28
11		<i>c</i>	As found in animal mitochondria. Presence indicated by spectroscopic and spectrophotometric observations.				5,10,12,27,28
12		<i>c</i> <sub>1</sub>	As found in animal mitochondria. Presence indicated by spectroscopic and spectrophotometric observations.			Present in high concentration in wheat roots	B-E,5,10,12, 27,28;F,28
13		<i>b</i> <sub>7</sub>	$E'_O = -0.03$	560 529		Observed in mitochondria from spadix of <i>Arum maculatum</i>	B,F,4;C,29;D, 4,29

/1/ From rat liver [7]. /2/ Found in microsomes [2, 32]. /3/ Intracellular localization unknown. /4/ From wheat roots.

*continued*



*Contributions*

**63. PROPERTIES OF CYTOCHROMES: ANIMALS AND HIGHER PLANTS**

Source of Pigments	Cytochrome	Physical and Chemical Properties	Spectral Characteristics		Remarks	Reference	
			$\lambda$ maximum in $m\mu$				
(A)	(B)	(C)	Reduced	Oxidized	(F)	(G)	
Higher Plants							
14	Endoplasmic reticulum <sup>5</sup>	$b_3$	PG = protohematin; non-mitochondrial oxidation of NADH and NADPH	559 525 425		Observed in microsomes from wheat roots and beet petiole. A pigment with a similar spectrum observed in autolysates of green leaves.	B,D,27,28;C,27;F,15,28
15	Chloroplasts <sup>6</sup>	$b_6$	PG = protohematin; $E'_0 = -0.06$ volts	563		Distinguished from cytochrome- $b$ by greater stability to organic solvents	B,F,14;C,29;D,14,29
16		$f$	MW = 110,000; PG = hematin- $c$ ; heat labile; $E'_0 = 0.365$ volts	555 526 421 330	410 350	Isolated from parsley. Localization in chloroplasts observed in etiolated leaves.	B,E,9;C,D,9,29;F,8,9


<sup>5/6</sup> Found in microsomes [27]. <sup>6/6</sup> Cytochrome- $b_6$  and cytochrome- $f$  probably are components of a system for oxidation of reduced lipoic acid formed during photosynthesis. The energy from this oxidation is used for generation of ATP from ADP and inorganic phosphate. [1]

*Contributors:* Morton, R. K., and Armstrong, J. M.

*References:* [1] Arnon, D. I. 1955. *Science* 122:9. [2] Ballie, M., and R. K. Morton. 1955. *Nature* 176:111. [3] Ball, E. 1938. *Biochem. Z.* 295:262. [4] Bendall, D. S., and R. Hill. 1956. *New Phytologist* 55:206. [5] Bhagvat, K., and R. Hill. 1951. *Ibid.* 50:112. [6] Chance, B. 1953. *J. Biol. Chem.* 202:397. [7] Chance, B., and G. R. Williams. 1955. *Ibid.* 217:395. [8] Davenport, H. E. 1952. *Nature* 170:1112. [9] Davenport, H. E., and R. Hill. 1952. *Proc. Roy. Soc. (London), B*, 139:327. [10] Goddard, D. R. 1944. *Am. J. Botany* 31:270. [11] Hagihara, B., et al. 1956. *Nature* 178:631. [12] Hartree, E. F. 1957. *Advan. Enzymol.* 18:1. [13] Henderson, R. W., and W. A. Rawlinson. 1956. *Biochem. J.* 62:21. [14] Hill, R. 1954. *Nature* 174:501. [15] Hill, R., and R. Scarisbrick. 1951. *New Phytologist* 50:98. [16] Hübscher, G., M. Kiese, and R. Nicholas. 1954. *Biochem. Z.* 325:223. [17] Keilin, D. 1929. *Proc. Roy. Soc. (London), B*, 104:206. [18] Keilin, D., and E. F. Hartree. 1937. *Ibid.*, B, 122:298. [19] Keilin, D., and E. F. Hartree. 1939. *Ibid.*, B, 127:167. [20] Keilin, D., and E. F. Hartree. 1949. *Nature* 164:254. [21] Keilin, D., and E. F. Hartree. 1955. *Ibid.* 176:200. [22] Keilin, D., and E. C. Slater. 1953. *Brit. Med. Bull.* 9:89. [23] Keilin, J. 1956. *Biochem. J.* 64:663. [24] Keilin, J. 1957. *Nature* 180:427. [25] Lemberg, R. 1953. *Ibid.* 172:619. [26] Lundegårdh, H. 1952. *Ibid.* 169:1088. [27] Martin, E. M., and R. K. Morton. 1955. *Ibid.* 176:113. [28] Martin, E. M., and R. K. Morton. 1957. *Biochem. J.* 65:404. [29] Neilands, J. B. 1958. *Ann. Rev. Biochem.* 27:455. [30] Okunuki, K., and E. Yakashiji. 1941. *Proc. Imp. Acad. (Tokyo)* 17:263. [31] Sekuzu, I., and K. Okunuki. 1956. *J. Biochem. (Tokyo)* 43:107. [32] Strittmatter, P., and E. Ball. 1954. *J. Cellular Comp. Physiol.* 43:57. [33] Strittmatter, P., and S. F. Velick. 1956. *J. Biol. Chem.* 221:253. [34] Tint, H., and W. Reiss. 1950. *Ibid.* 182:385. [35] Wainio, W. W., and S. J. Cooperstein. 1956. *Advan. Enzymol.* 17:347.

## 64. PATHWAYS OF BIOSYNTHESIS: PURINES

\* = isotopically labeled atom incorporated into next product.




**Contributors:** Lansford, Edwin M., Jr., and Shive, William

**References:** [1] Buchanan, J. M., et al. 1959. *J. Cellular Comp. Physiol.* 54:139. [2] Hartman, S. C., and J. M. Buchanan. 1959. *Ann. Rev. Biochem.* 28:365. [3] White, A., et al. 1959. *Principles of biochemistry*. Ed. 2. McGraw-Hill, New York. p. 582.

## 65. PATHWAYS OF BIOSYNTHESIS: PYRIMIDINES

### ENZYME KEY

ATP = adenosine triphosphate	PP = inorganic pyrophosphate
CTP = cytidine triphosphate	PRPP = phosphoribosylpyrophosphate
dCTP = deoxycytidine triphosphate	P-R = 5'-ribosylphosphate
UTP = uridine triphosphate	PPP-R = 5'-ribosyltriphosphate
dTTP = deoxythymidine triphosphate	P-dR = 5'-deoxyribosulphosphate
NAD <sup>+</sup> = nicotinamide adenine dinucleotide	THF = tetrahydrofolate coenzyme




<sup>/1/</sup> The mechanism for this reaction varies slightly with the enzyme source (animal or microbial). <sup>/2/</sup> One or more enzymic reactions not yet fully elucidated. <sup>/3/</sup> Reaction catalyzed by RNA polymerase. <sup>/4/</sup> Reaction catalyzed by DNA polymerase.

**Contributors:** Lansford, Edwin M., Jr., and Shive, William

**References:** [1] Buchanan, J. M., et al. 1959. *J. Cellular Comp. Physiol.* 54:139. [2] Hartman, S. C., and J. M. Buchanan. 1959. *Ann. Rev. Biochem.* 28:365. [3] White, A., et al. 1959. *Principles of biochemistry*. Ed. 2. McGraw-Hill, New York. p. 582.

## 66. PATHWAYS OF BIOSYNTHESIS: CHLOROPHYLL

Diagram summarizes present knowledge of the pathways leading to synthesis of chlorophyll in plants.



/1/ Condensation of glycine with succinyl-CoA produces  $\alpha$ -amino- $\beta$ -ketoadipate. /2/ Decarboxylates to  $\delta$ -aminolevulinate. /3/ Two molecules of  $\delta$ -aminolevulinate are condensed by a dehydrase enzyme, forming the pyrrole amine, porphobilinogen. /4/ Four molecules of porphobilinogen are condensed, forming a reduced porphyrin, uroporphyrinogen III. /5/ Autoxidation products found in congenital porphyria. /6/ Decarboxylation of uroporphyrinogen III to coproporphyrinogen III. /7/ Side products found in a *Chlorella* mutant. /8/ Oxidation of one propionic side chain to vinyl. /9/ Oxidation of one propionic side chain to vinyl, producing protoporphyrinogen. /10/ Autoxidation to protoporphyrin. /11/ Open-chain bile pigment chemically bound to protein. /12/ Magnesium protoporphyrin found in a *Chlorella* mutant. /13/ Found in a *Chlorella* mutant and etiolated barley. /14/ Three or four steps postulated, including reduction of a vinyl side chain to ethyl, oxidation of a propionic acid group and cyclization to a cyclopentanone ring. Found in a *Chlorella* mutant. /15/ Addition of two hydrogen atoms on pyrrole ring D of protochlorophyllide. /16/ Esterification of a propionic acid group with phytol, a C-20 alcohol.


**Contributor:** Granick, S.

**References:** [1] Granick, S. 1951. Ann. Rev. Plant Physiol. 2:115. [2] Shemin, D. 1955. Ciba Found. Symp. Porphyrin Biosyn. Metab., p. 4.



## 67. PATHWAYS OF PHOTOSYNTHESIS: CARBON DIOXIDE REDUCTION CYCLE

According to current evidence, photosynthetic carbon dioxide reduction follows the same general pathways in all plants. The first reaction results in formation of two molecules of phosphoglycerate from carbon dioxide and ribulose diphosphate. Phosphoglycerate is then reduced via the reverse of glycolysis (indicated by light arrows) to supply hexose phosphates for synthesis of sucrose and polysaccharides. A portion of the intermediate compounds goes through the sequence of reactions shown in the diagram, leading to regeneration of the carbon dioxide acceptor, ribulose diphosphate. It is not known whether one or both of the pathways to sedoheptulose phosphate (step 8 and steps 7 and 9) are important in carbon reduction during photosynthesis. Heavy arrows indicate directions of material transfer during steady state photosynthesis.



/1/ Ribulose diphosphate adds CO<sub>2</sub> at carbon-2 and splits hydrolytically to give two molecules of 3-phosphoglycerate.  
 /2/ The carboxyl group is reduced to an aldehyde group with the aid of ATP and NADPH. /3/ Isomerization involves transfer of two hydrogen atoms from carbon-2 to carbon-1. /4/ Aldol condensation of carbon-1 of glyceraldehyde-3-P with carbon-1 of dihydroxyacetone-3-P. /5/ Removal of the phosphate ester group from carbon-1 by hydrolysis. /6/ Two hydrogen atoms plus the glycolyl group (carbon-1, 2) of fructose transferred to glyceraldehyde-P to form xylulose-5-P, leaving erythrose-P. /7/ Aldol condensation of carbon-1 of erythrose-4-P with carbon-1 of dihydroxyacetone-3-P obtained from step 3. /8/ Transfer of triose group (carbon-1, 2, 3) from fructose-6-P to carbon-1 of erythrose-4-P, leaving glyceraldehyde-3-P from carbon-4, 5, 6 of the fructose-6-P. /9/ Hydrolysis of the phosphate ester group on carbon-1 to give inorganic phosphate. /10/ Transfer of the glycolyl group (carbon-1, 2) of sedoheptulose-7-P to carbon-1 of glyceraldehyde-3-P to give ribulose-5-P, leaving ribose-5-P. /11/ Epimerization of carbon-3 of ketopentose. Xylulose-5-P isomerizes to ribulose-5-P with phosphoketopentose epimerase. /12/ Isomerization of aldose to ketose by the transfer of two hydrogen atoms. /13/ Phosphorylation of carbon-1 by reaction with ATP.

continued


## 67. PATHWAYS OF PHOTOSYNTHESIS: CARBON DIOXIDE REDUCTION CYCLE

**Contributors:** (a) Benson, Andrew A., (b) Calvin, Melvin, and Bassham, James A.

**References:** [1] Bassham, J. A., and M. Calvin. 1957. The path of carbon in photosynthesis. Prentice-Hall, Englewood Cliffs, N. J. [2] Calvin, M., and J. A. Bassham. 1962. The photosynthesis of carbon compounds. W. A. Benjamin, New York.

## 68. PATHWAYS OF SUCROSE SYNTHESIS: INTERMEDIATES

Sucrose, common to all green plants, is the first free sugar formed by a series of steps involving phosphorylated intermediates. Photosynthesis supplies reduced pyridine nucleotides and adenosine triphosphate (ATP). Phosphoglycerate, provided by the photosynthetic carboxylation reaction, is reduced and condensed to form hexose molecules. Energy required to produce sucrose from hexose phosphates comes largely from high-energy uridine triphosphate (UTP) which becomes uridine diphosphate glucose (UDP-glucose) for condensation with fructose phosphate.



/1/ Hydrogen atom on carbon-1 shifts to carbon-2, forming the epimer, glucose-6-phosphate; furanose ring structure is changed to pyranose. /2/ Phosphate group on carbon-6 is transferred to carbon-1 through the required co-enzyme intermediate, glucose-1,6-diphosphate; Mg<sup>++</sup> is required. /3/ UTP reacts with glucose-1-phosphate to form pyrophosphate and UDP-glucose. /4/ Fructose-6-phosphate and UDP-glucose react to give UDP and an unstable sucrose phosphate. /5/ Hydrolysis occurs to give free sucrose and orthophosphate.

**Contributors:** (a) Benson, Andrew A., (b) Calvin, Melvin, and Bassham, James A.

**References:** [1] Bassham, J. A., and M. Calvin. 1957. The path of carbon in photosynthesis. Prentice-Hall, Englewood Cliffs, N. J. [2] Calvin, M., and J. A. Bassham. 1962. The photosynthesis of carbon compounds. W. A. Benjamin, New York.

## 69. PHOTOSYNTHESIS: APPARENT RATES

Photosynthesis is complicated by such factors as light intensity, temperature, CO<sub>2</sub> concentration, and certain internal conditions of the plant.

### Part I. MAXIMUM RATES: NATURAL CONDITIONS, VARIOUS LOCALES

Values are mg CO<sub>2</sub> per 100 sq cm per hour, unless otherwise indicated, and are uncorrected for respiration. Determinations were made chiefly in sunlight; those made in the shade are enclosed in parentheses.

Species	Common Name	Location	Temp. °C	CO <sub>2</sub> Fixation	Reference
(A)	(B)	(C)	(D)	(E)	(F)
Lichenes					
1 <i>Lasallia papulosa</i>	Rock tripe	Southern Sweden	28	2.0	28
2 <i>Lobaria pulmonaria</i>	Lungwort	Southern Sweden	27	(2.2)	28

*continued*

## 69. PHOTOSYNTHESIS: APPARENT RATES

### Part I. MAXIMUM RATES: NATURAL CONDITIONS, VARIOUS LOCALES

	Species	Common Name	Location	Temp. °C	CO <sub>2</sub> Fixation	Refer- ence
	(A)	(B)	(C)	(D)	(E)	(F)
Algae						
3	<i>Fucus serratus</i>	Rockweed	Helgoland	17	4.50	15
4	<i>Laminaria saccharina</i>	Kelp	Helgoland	20	1.75	15
5	<i>Porphyra laciniata</i>	Porphyra	Helgoland	17	2.09	15
6			Naples	23	1.64	15
7	<i>Ulva lactuca</i>	Sea lettuce	Naples	21-23	1.62	15
Pteridophyta						
8	<i>Dryopteris spinulosa</i>	Toothed wood fern	Southern Sweden	.....	3.96	14
9	<i>Polypodium vulgare</i>	Common polypody	Southern Sweden	.....	(5.60)	14
Gymnospermae						
10	<i>Picea abies</i>	Norway spruce	Southern Sweden	.....	1.5 (2.1) <sup>1</sup>	27
11			Germany	-2 to +3	0.096 <sup>2</sup>	31
12			Germany	6-12	0.29 <sup>2</sup>	31
13	<i>Pinus taeda</i>	Loblolly pine	North Carolina	26	14.3	19
14			Massachusetts	.....	(2.51)	18
Angiospermae						
15	<i>Acer platanoides</i>	Norway maple	Stockholm	18-20	1.22 <sup>3</sup>	27
16	<i>A. tschonoski</i>	Tschonoski maple	Japan	.....	(4.1)	13
17	<i>Allium victorialis</i>	Long-root onion	Leningrad	.....	3.66	8
18	<i>Avena</i> sp.	Oat	Denmark	20	13.0	4
19	<i>Beta vulgaris</i>	Beet	Sweden	.....	20.1	20,21
20	<i>Betula pendula</i>	European white birch	Denmark	20	6.4	5,6
21	<i>Catalpa bignonioides</i>	Southern catalpa	England	.....	4.7	7
22	<i>Chrysanthemum alpinum</i>	Alpine chrysanthemum	Alps	14	(84.1)	12
23	<i>Citrus limon</i>	Lemon	Southern USSR	25-28	15-26	9
24	<i>Cucumis sativus</i>	Cucumber	USSR	25	10.91-11.98	24,25
25	<i>Cucurbita pepo</i>	Pumpkin	Germany	.....	20 <sup>2</sup>	26
26	<i>Fagopyrum esculentum</i>	Buckwheat	USSR	.....	10.5	22
27	<i>Fagus sylvatica</i>	European beech	Denmark	20	6.6 (2.4)	3
28	<i>Fraxinus excelsior</i>	European ash	Denmark	20	9.8 (4.2)	3
29	<i>Glycine soja</i>	Soybean	USSR	25	7.93-8.78	24,25
30	<i>Gossypium hirsutum</i>	Upland cotton	USSR	.....	7.98	32
31	<i>Helianthus annuus</i>	Common sunflower	Netherlands	13-27	5.8	10
32			Java	28-36	7.8	10
33	<i>Hordeum</i> sp.	Barley	Pamirs	.....	>30	1
34	<i>Lycopersicon esculentum</i>	Tomato	Southern Sweden	20	16.8	28
35	<i>Malus pumila</i>	Common apple	Ohio	26	20 (20)	2
36	<i>Medicago sativa</i>	Alfalfa	Central Asia	.....	22.4	17
37	<i>Nicotiana tabacum</i>	Common tobacco	USSR	25	8.40	24,25
38	<i>Phaseolus vulgaris</i>	Kidney bean	Sweden	.....	17.0	30
39	<i>Phleum pratense</i>	Timothy	USSR	17-20	23	16
40	<i>Phoenix dactylifera</i>	Date palm	Algeria	.....	3.4	11
41	<i>Prunus laurocerasus</i>	Laurel cherry	Germany	-3 to 0	0.054 <sup>2</sup>	31
42			Germany	3-12	0.34 <sup>2</sup>	31
43	<i>Quercus ilex</i>	Holly oak	Italy	12-14	65.7 <sup>3</sup>	29
44	<i>Q. lyrata</i>	Overcup oak	Massachusetts	.....	(10.8)	18
45	<i>Rheum officinale</i>	Medicinal rhubarb	Germany	.....	19 <sup>3</sup>	26
46	<i>Rhododendron brachycarpum</i>	Fujiyama rhododendron	Japan	.....	(2.8)	13
47	<i>Salix glauca</i>	Gray-leaf willow	Greenland	10	4	23
48			Greenland	20	6	23
49	<i>Sambucus nigra</i>	European elder	Denmark	20	4.6 (1.7)	3
50	<i>Solanum tuberosum</i>	Potato	Southern Sweden	20	19.2	28
51	<i>Trifolium pratense</i>	Red clover	USSR	20-29	24	16
52	<i>Triticum aestivum</i>	Wheat	Central Asia	.....	36.5	17
53			Germany	-6 to 0	0.52 <sup>2</sup>	31
54			Germany	0-6	1.67 <sup>2</sup>	31
55	<i>Vicia sativa</i>	Vetch	USSR	.....	12.21	15
56	<i>Vitis vinifera</i>	European grape	Central Asia	.....	16.1	17
57	<i>Zea mays</i>	Corn	USSR	25	10.02	25

/1/ mg CO<sub>2</sub> per g fresh weight per hour. /2/ mg CO<sub>2</sub> per g dry weight per hour. /3/ Calculated from the assay of assimilates, assuming an empirical formula of CH<sub>2</sub>O.

continued

## 69. PHOTOSYNTHESIS: APPARENT RATES

### Part I. MAXIMUM RATES: NATURAL CONDITIONS, VARIOUS LOCALES

Contributors: (a) Siegel, Jack M., (b) Olson, Rodney A.

References: [1] Blagoveshchenskij, A. V. 1935. *Planta* 24:276. [2] Böhning, R. H. 1949. *Plant Physiol.* 24:222. [3] Boysen-Jensen, P. 1919. *Botan. Tidsskr.* 36:219. [4] Boysen-Jensen, P. 1932. *Stoffproduktion der Pflanzen.* G. Fischer, Jena. [5] Boysen-Jensen, P., and D. Müller. 1929. *Jahrb. Wiss. Botan.* 70:493. [6] Boysen-Jensen, P., and D. Müller. 1929. *Ibid.* 70:503. [7] Brown, H. T., and F. Escombe. 1905. *Proc. Roy. Soc. (London)*, B, 76:29. [8] Chrelashvili, M. N. 1941. *Tr. Botan. Inst. Akad. Nauk SSSR*, IV, 5:101. [9] Filippenko, I. A., E. H. Gerber, and O. K. Elpidina. 1937. *Compt. Rend. Acad. Sci. URSS* 17:323. [10] Giltay, E. 1898. *Ann. Jard. Botan. Buitenzorg* 15:43. [11] Harder, R., P. Filzer, and A. Lorenz. 1932. *Jahrb. Wiss. Botan.* 75:45. [12] Henrici, M. 1921. *Verhandl. Naturforsch. Ges. Basel* 32:107. [13] Hiramatsu, K. 1932. *Sci. Rept. Tohoku Imp. Univ., Ser. 2*, 7:239. [14] Johansson, N. 1926. *Svensk Botan. Tidskr.* 20:107. [15] Kniep, H. 1914. *Intern. Rev. Ges. Hydrobiol. Hydrog.* 7:1. [16] Kostychev, S., K. Bazyrina, and G. Vasiliev. 1927. *Biochem. Z.* 182:79. [17] Kostychev, S., and H. Kardo-Sysojeva. 1930. *Planta* 11:117. [18] Kozlowski, T. T. 1949. *Ecol. Monographs* 19:207. [19] Kramer, P. J., and W. S. Clark. 1947. *Plant Physiol.* 22:51. [20] Lundegårdh, H. 1924. *Biochem. Z.* 154:195. [21] Lundegårdh, H. 1924. *Medd. Centralanstalt. Foersoeksvaesendet Jordbruks.* 331. [22] Maximov, N. A., and T. A. Krasnoselskaja-Maximova. 1928. *Ber. Deut. Botan. Ges.* 46:383. [23] Müller, D. 1928. *Planta* 6:22. [24] Richter, A. A., K. T. Sukhorukov, and L. A. Ostapenko. 1945. *Compt. Rend. Acad. Sci. URSS* 46:40. [25] Richter, A. A., K. T. Sukhorukov, and L. A. Ostapenko. 1945. *Ibid.* 46:165. [26] Sachs, J. 1884. *Arb. Botan. Inst. Wuerzburg* 3:1. [27] Stålfelt, M. G. 1921. *Medd. Statens Skogsfoersoeksanstalt (Stockholm)* 18:221. [28] Stöcker, O. 1927. *Flora (Jena)* 121:334. [29] Von Guttenberg, H. 1927. *Planta* 4:726. [30] Yoshii, Y. 1928. *Ibid.* 6:22. [31] Zeller, O. 1951. *Ibid.* 39:500. [32] Zhdanova, L. P. 1944. *Compt. Rend. Acad. Sci. URSS* 45:353.

### Part II. MAXIMUM RATES: NEAR-OPTIMUM CONDITIONS

Values are uncorrected for respiration.

	Species	Common Name	CO <sub>2</sub> in Air %	Temp. °C	CO <sub>2</sub> Fixation, g CO <sub>2</sub> /hr			Assimilation Time (TA) <sup>2</sup> sec	Reference	
					per 100 g wet wt	per 100 g dry wt	per sq dm x 1000			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Algae										
1	<i>Chlorella pyrenoidosa</i> <sup>3</sup>	Chlorella	...	...	...	13.4	...	2.8	56	3
2	In light		...	...	...	11.5	...	4.1	36	3
3	<i>Gigartina harveyana</i> <sup>4</sup>	Seaweed	...	16	...	...	14	...	...	1
4	<i>Hormidium flaccidum</i>	Water net	...	20	...	...	...	6.8	23	4
5	<i>Ulva lactuca</i>	Sea lettuce	...	25	...	...	...	11.8	...	2
Angiospermae										
6	<i>Acer pseudoplatanus</i>	Plane-tree	5	25	0.98	3.0	16	11.8	13	5
7	Young leaves	maple	5	25	2.07	5.8	26	5.2	30	5
8	<i>Helianthus annuus</i>	Common sunflower	5	25	2.30	13.4	80	14.0	11	5
9	<i>Populus alba</i>	White poplar	5	25	1.90	6.0	40	10.0	16	5
10	<i>Sambucus nigra</i>	European elder	5	25	1.96	5.3	34	6.6	24	5
11	Green leaves		5	25	0.88	4.7	18	120	1.3	5
12	<i>Tilia cordata</i>	Little-leaf linden	5	25	1.88	5.8	28	6.6	24	5

<sup>1</sup>/<sub>1</sub>  $\sqrt{A}$ , the assimilation number, is the maximum quantity of CO<sub>2</sub> that can be reduced in unit time by unit quantity of chlorophyll. <sup>2</sup>/<sub>2</sub> The shortest time in which one molecule of chlorophyll can reduce one molecule of CO<sub>2</sub>. <sup>3</sup>/<sub>3</sub> In carbonate buffer 9. <sup>4</sup>/<sub>4</sub> In artificial seawater.

continued



## 69. PHOTOSYNTHESIS: APPARENT RATES

### Part II. MAXIMUM RATES: NEAR - OPTIMUM CONDITIONS

Contributor: Olson, Rodney A.

References: [1] Emerson, R., and L. Green. 1934. J. Gen. Physiol. 17:817. [2] Kniep, H. 1914. Intern. Rev. Ges. Hydrobiol. Hydrog. 7:1. [3] Noddack, W., and C. Kopp. 1940. Z. Physik. Chem., A, 187:79. [4] Van der Honert, T. H. 1930. Rev. Trav. Botan. Neerl. 27:149. [5] Willstätter, R., and A. Stoll. 1918. Untersuchungen über die Assimilation der Kohlensäure. J. Springer, Berlin.

### Part III. AVERAGE RATES

Values are uncorrected for respiration. Values in parentheses are maximum rates. Temp. (column C): N = under natural conditions.

	Species	Common Name	Conditions			Photosynthesis rate/hr	Reference
			Temp. °C	Light ft-c	CO <sub>2</sub> in Air		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
Algae							
1	<i>Chlorella saccharophila</i>	Chlorella	22.4	2,480	Buffer 9 <sup>1</sup>	452.3 mm <sup>3</sup> O <sub>2</sub> /100 million cells	19
2	<i>C. vulgaris viridis</i>	Chlorella	22.4	2,480	Buffer 9 <sup>1</sup>	194.7 mm <sup>3</sup> O <sub>2</sub> /100 million cells	19
3	<i>Cladophora glomerata</i>	Cladophora	17	.....	.....	0.3 moles CO <sub>2</sub> /10μ <sup>3</sup>	15
4	<i>Nostoc muscorum</i>	Nostoc	25	1,000	Buffer 9 <sup>1</sup>	13.2 ml O <sub>2</sub> /ml packed cells	2
Bryophyta							
5	<i>Hylocomium proliferum</i>	Hylocomium	....	93-186	.....	1.25-3 mg CO <sub>2</sub> /g dry wt	6
6	<i>Sphagnum girgensohnii</i>	Sphagnum moss	....	110-260	.....	2.75 mg CO <sub>2</sub> /g dry wt	6
Gymnospermae							
7	<i>Picea pungens</i>	Colorado spruce	24	2,200	Natural	0.03 mg CO <sub>2</sub> /100 leaves	4
8	<i>Pinus taeda</i>	Loblolly pine	30	2,000	Natural	2 (3.9) mg CO <sub>2</sub> /dm <sup>2</sup>	8
Angiospermae							
9	<i>Citrus limon</i>	Lemon	....	1,300	1.5%	3-5 ml O <sub>2</sub> /dm <sup>2</sup>	17
10	<i>C. sinensis</i>	Sweet orange	....	1,300	1.5%	4-6 ml O <sub>2</sub> /dm <sup>2</sup>	17
11	<i>Cornus florida</i>	Flowering dogwood	30	2,000	Natural	2 (3.06) mg CO <sub>2</sub> /dm <sup>2</sup>	8
12	<i>Cucurbita pepo</i>	Pumpkin	....	.....	.....	0.68 g/M <sup>2</sup> increase in dry wt	13
13	<i>Helianthus annuus</i>	Common sunflower	....	4,460	5%	(80) mg CO <sub>2</sub> /dm <sup>2</sup>	18
14	<i>Hordeum vulgare</i>	Barley	N	500	Natural	9-16 mg CO <sub>2</sub> /dm <sup>2</sup>	5
15	<i>Lycopersicon esculentum</i>	Tomato	....	.....	.....	16.5-18.6 mg CO <sub>2</sub> /dm <sup>2</sup>	10
16	<i>Malus sylvestris</i>	Apple	N	Natural	Natural	6.6 mg CO <sub>2</sub> /dm <sup>2</sup>	7
17	<i>Medicago sativa</i>	Alfalfa	N	Noon sun	Natural	0.75 (1.042) g/6 x 6 ft plot	14
18	<i>Oryza sativa</i>	Rice	31	1.74 cal/cm/min	Natural	9-20 mg CO <sub>2</sub> /M <sup>2</sup>	11
19	<i>Phaseolus vulgaris</i>	Kidney bean	25	1,400	Natural	5.8-16.6 mg CO <sub>2</sub> /dm <sup>2</sup>	1
20	<i>Prunus laurocerasus</i>	Laurel cherry	29.5	Noon sun	Natural	23.2 mg CO <sub>2</sub> /dm <sup>2</sup>	9
21	<i>P. persica</i>	Peach	N	Natural	Natural	0.146-0.18 g/M <sup>2</sup> increase in dry wt	12
22	<i>Quercus rubra</i>	Eastern red oak	30	2,000	Natural	5 (6.04) mg CO <sub>2</sub> /dm <sup>2</sup>	8
23	<i>Rheum rhaponticum</i>	Garden rhubarb	....	.....	.....	0.65 g/M <sup>2</sup> increase in dry wt	13
24	<i>Solanum tuberosum</i>	Potato	N	>5,000	Natural	16-20 mg CO <sub>2</sub> /dm <sup>2</sup>	3
25	<i>Zea mays</i>	Corn	N	Full sun	Natural	1.8 g CO <sub>2</sub> /M <sup>2</sup> leaf	16

/1/ Carbonate buffer 9.

Contributor: Bing, Arthur

References: [1] Bing, A. Unpublished. Cornell Univ. Ornamentals Research Laboratory, N. Y., 1953. [2] Brown, T. E. 1954. Ph.D. Thesis. Ohio State Univ., Columbus. [3] Chapman, H. W. 1951. Am. Potato J. 28(5):602. [4] Freeland, R. O. 1952. Plant Physiol. 27(4):685. [5] Gregory, F. G., and F. J. Richards. 1929. Ann. Botany (London) 43:119. [6] Harder, R. 1930. Planta 11:263. [7] Heinicke, A. J., and M. B. Hoffman. 1933. Cornell

continued

## 69. PHOTOSYNTHESIS: APPARENT RATES

### Part III. AVERAGE RATES

Univ. Agr. Expt. Sta. Bull. 577. [8] Kramer, P. J., and J. P. Decker. 1944. *Plant Physiol.* 19(2):350.  
 [9] Matthaei, G. L. C. 1905. *Phil. Trans. Roy. Soc. London, B.* 197:47. [10] Mitchell, J. W. 1936. *Botan. Gaz.* 98:87. [11] Noguti, Y. 1941. *Japan. J. Botany* 11(2):167. [12] Pickett, W. F., A. S. Fish, and W. S. Shan. 1951. *Proc. Am. Soc. Hort. Sci.* 57:111. [13] Sachs, J. 1884. *Arb. Botan. Inst. Wuerzburg* 3:1. [14] Thomas, M. D., and G. R. Hill. 1937. *Plant Physiol.* 12:285. [15] Verduin, J. 1952. *Am. J. Botany* 39(3):157. [16] Verduin, J., and W. E. Loomis. 1944. *Plant Physiol.* 19:278. [17] Wedding, R. T., L. A. Riehl, and W. H. Rhoads. 1952. *Ibid.* 27(2):269. [18] Willstätter, R., and A. Stoll. 1918. *Untersuchungen über die Assimilation der Kohlensäure.* J. Springer, Berlin. [19] Winokur, M. 1948. *Am. J. Botany* 35(5):207.

## 70. CARBON PRODUCTION AND PHOTOSYNTHETIC EFFICIENCY

### Part I. ESTIMATED ANNUAL CARBON PRODUCTION

Region	Area sq km	Carbon Fixed ton/yr		Refer- ence
		per sq km	Total	
(A)	(B)	(C)	(D)	(E)
1 Forest	44 x 10 <sup>6</sup>	250	11 x 10 <sup>9</sup>	1
2 Cultivated land	27 x 10 <sup>6</sup>	160	4.3 x 10 <sup>9</sup>	1
3 Grassland	31 x 10 <sup>6</sup>	36	1.1 x 10 <sup>9</sup>	1
4 Desert	47 x 10 <sup>6</sup>	7	0.3 x 10 <sup>9</sup>	1
5 Total land	149 x 10 <sup>6</sup>		16.7 x 10 <sup>9</sup>	1
6 Ocean	371 x 10 <sup>6</sup>	340	126 x 10 <sup>9</sup> <sup>1</sup>	2

/1/ 16 x 10<sup>9</sup> also reported [3].

Contributor: Bohning, Richard H.

References: [1] Rabinowitch, E. I. 1945. *Photosynthesis.* Interscience, New York. v. 1, p. 6.

[2] Riley, G. A. 1944. *Am. Scientist* 32:129.

[3] Steemann-Nielsen, E. 1952. *Nature* 169:956.

### Part II. ENERGY UTILIZATION IN PHOTOSYNTHESIS

Specification	Value	Refer- ence
(A)	(B)	(C)
1 Energy utilized in photosynthesis by one acre of corn plants to produce 8,732 kg glucose <sup>1</sup>	3.3 x 10 <sup>7</sup> kcal	2
2 Total solar energy available on the acre during growing season	2.043 x 10 <sup>9</sup> kcal	2
3 Photosynthetic efficiency of corn plants, i.e., percent of available energy used in photosynthesis	1.6%	2
4 Energy equivalent of earth's carbon production	(13.6±8.1) x 10 <sup>17</sup> kcal	1
5 Mean solar radiation	7.4 x 10 <sup>20</sup> kcal	1
6 Photosynthetic efficiency of the world	0.18±0.12%	1

/1/ Total sugar, as glucose, manufactured by one acre of corn plants.

Contributor: Bohning, Richard H.

References: [1] Riley, G. A. 1944. *Am. Scientist* 32:129.

[2] Transeau, E. N. 1926. *Ohio J. Sci.* 26:1.

## 71. NITROGEN FIXATION

### Part I. RHIZOBIA-INOCULATED LEGUMES

The amount of nitrogen fixed from the air by the symbiotic relationship of rhizobia with legumes is influenced by the effectiveness of the rhizobia, host species, soil and climatic conditions, and individual crop handling. Values in parentheses are ranges, estimate "e" (cf. Introduction).

Rhizobium and Host		N <sub>2</sub> Fixed kg/acre	Rhizobium and Host		N <sub>2</sub> Fixed kg/acre
(A)	(B)	(B)	(A)	(B)	(B)
1 <i>Rhizobium</i> spp.			5 <i>Rhizobium</i> spp.		
2 <i>Arachis hypogaea</i> (peanut)	19		6 <i>Stizolobium deeringianum</i> (Florida velvet bean)	30	
3 <i>Cicer arietinum</i> (garbanzo)	30		7 <i>Vigna sinensis</i> (cowpea)	41(26-53)	
4 <i>Lespedeza</i> spp. <sup>1</sup> (lespedeza)	39(15-94)		Pastures with legumes	48(5-91)	
5 <i>Pueraria thumbergiana</i> (kudzu)	49(40-57)				

/1/ *L. striata* (common lespedeza) and *L. stipulacea* (Korean lespedeza).

continued

## 71. NITROGEN FIXATION

### Part I. RHIZOBIA-INOCULATED LEGUMES

Rhizobium and Host		N <sub>2</sub> Fixed kg/acre	Rhizobium and Host		N <sub>2</sub> Fixed kg/acre
(A)		(B)	(A)		(B)
8	<i>Rhizobium japonicum</i> <i>Glycine soja</i> (soybean)	26(7-48)	15	<i>Rhizobium meliloti</i> <i>Medicago sativa</i> (alfalfa)	88(25-170)
	<i>R. leguminosarum</i>		16	<i>Melilotus alba</i> (white sweet clover)	54(30-75)
9	<i>Lens culinaris</i> (lentil)	47	17	<i>M. indica</i> (annual yellow sweet clover)	45
10	<i>Pisum sativum</i> (garden pea)	33(13-60)	18	<i>Trigonella foenum-graecum</i> (fenugreek)	37
11	<i>P. sativum arvense</i> (field pea)	23(20-28)		<i>R. phaseoli</i>	
12	<i>Vicia</i> spp. (vetch)	36(20-63)	19	<i>Phaseolus vulgaris</i> (kidney bean)	18
	<i>R. lupini</i>		20	<i>R. trifolii</i> <i>Trifolium incarnatum</i> (crimson clover)	42(32-53)
13	<i>Lupinus angustifolius</i> (lupine)	68	21	<i>T. pratense</i> (red clover)	52(19-78)
	<i>R. meliloti</i>		22	<i>T. repens</i> (white clover)	47(35-65)
14	<i>Medicago hispida denticulata</i> (toothed bur clover)	35(23-49)	23	<i>T. repens giganteum</i> (white clover)	81(72-91)

Contributor: Erdman, Lewis W.

Reference: Erdman, L. W. 1948. U. S. Dept. Agr. Farmers' Bull. 2003.

### Part II. CHARACTERISTICS OF NITROGEN-FIXING ORGANISMS

Organism	Essential Host	Oxygen	Nourishment	Limiting pH	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Bacteria						
1	<i>Azotobacter</i> <sup>1</sup>	None	Aerobic	Heterotrophic	2.9-9.2	4, 5, 8, 18, 21
2	<i>Chlorobacterium</i>	None	Anaerobic	Heterotrophic	ca. 7.0	4, 16
3	<i>Chromatium</i>	None	Anaerobic	Autotrophic	ca. 7.0	4, 16
4	<i>Clostridium</i> <sup>2</sup>	None	Anaerobic	Heterotrophic	4.0-8.0	4, 6, 15, 22
5	<i>Desulfovibrio</i>	None	Anaerobic	Heterotrophic	ca. 7.0	4, 19
6	<i>Rhizobium</i> spp. <sup>3</sup>	<i>Vigna, Lespedeza</i>	Aerobic*	Heterotrophic	3.3	1-4,7,11-14,18
7	<i>R. japonicum</i> <sup>3</sup>	<i>Glycine</i>	Aerobic	Heterotrophic	3.3	1-4,7,11-14,18
8	<i>R. leguminosarum</i> <sup>3</sup>	<i>Pisum</i>	Aerobic	Heterotrophic	4.7	1-4,7,11-14,18
9	<i>R. lupini</i>	<i>Lupinus</i>	Aerobic	Heterotrophic	3.15	1-4,7,11-14,18
10	<i>R. meliloti</i> <sup>1,3</sup>	<i>Medicago, Melilotus</i>	Aerobic	Heterotrophic	4.9	1-4,7,11-14,18
11	<i>R. phaseoli</i> <sup>3</sup>	<i>Phaseolus</i>	Aerobic	Heterotrophic	4.2	1-4,7,11-14,18
12	<i>R. trifolii</i> <sup>3</sup>	<i>Trifolium</i>	Aerobic	Heterotrophic	4.2	1-4,7,11-14,18
13	<i>Rhodomicrobium</i>	None	Anaerobic	Heterotrophic	ca. 7.0	4, 9, 17
14	<i>Rhodospseudomonas</i>	None	Anaerobic <sup>4</sup>	Heterotrophic	ca. 7.0	4,9,16,17
15	<i>Rhodospirillum</i>	None	Anaerobic <sup>4</sup>	Heterotrophic	ca. 7.0	4,9,16,17
Algae						
16	<i>Catolthrix</i> <sup>1</sup>	None	Aerobic	Autotrophic	5.7-8.5	4-6,10,20
17	<i>Nostoc</i> <sup>1</sup>	None	Aerobic	Autotrophic	5.7-8.5	4-6,10,20

/1/ Inhibited by NH<sub>4</sub> and NO<sub>3</sub>, stimulated by Mo. /2/ Inhibited by NH<sub>4</sub>, stimulated by Mo. /3/ N<sub>2</sub> fixation inhibited by 2,4-D and by seed treatments containing Cu; slightly inhibited by DDT. /4/ Fixation is best under anaerobic conditions in light, slight under aerobic conditions in dark. Organism is facultative.

Contributor: Appleman, Milo D.

References: [1] Anderson, A. J. 1946. J. Council Sci. Ind. Res. 19:1. [2] Appleman, M. D. 1941. Soil Sci. Soc. Am. Proc. 6:200. [3] Appleman, M. D. 1946. J. Am. Soc. Agron. 38:545. [4] Appleman, M. D. Unpublished. Univ. Southern California, Los Angeles, 1953. [5] Bortch, H. 1930. Arch. Mikrobiol. 1:333. [6] Bortch, H. 1936. Zentr. Bakteriell. Parasitenk., II, 95:193. [7] Bortch, H. 1937. Arch. Mikrobiol. 8:13. [8] Burk, D. 1939. Ergeb. Enzymforsch. 3:23. [9] Duchow, E., and H. C. Douglas. 1949. J. Bacteriol. 58:409. [10] Fogg, G. E. 1942. J. Exptl. Biol. 19:78. [11] Fulst, J. L., and M. G. Payne. 1947. Am. J. Botany 34:245. [12] Jensen, H. L. 1946. Proc. Linnean Soc. N. S. Wales 70:203. [13] Jensen, H. L. 1948. Ibid. 72:265. [14] Jensen, H. L., and


continued

## 71. NITROGEN FIXATION

### Part II. CHARACTERISTICS OF NITROGEN-FIXING ORGANISMS

R. C. Betty. 1943. *Ibid.* 68:1. [15] Jensen, H. L., and D. Spencer. 1947. *Ibid.* 72:73. [16] Lindstrom, E. S., et al. 1949. *J. Bacteriol.* 58:313. [17] Lindstrom, E. S., et al. 1951. *Ibid.* 61:481. [18] Milder, E. G. 1948. *Plant Soil* 1:94. [19] Sisler, F. D., and C. E. Zobell. 1951. *Science* 113:511. [20] Williams, A. E. 1951. M.S. Thesis. Univ. Wisconsin, Madison. [21] Wilson, P. W., and R. H. Burris. 1947. *Bacteriol. Rev.* 11:41. [22] Zelitch, I. 1951. *Proc. Natl. Acad. Sci. U.S.* 37:559.

## 72. NITROGEN CYCLE IN NATURE



Contributor: Thimann, Kenneth V.

Reference: Thimann, K. V. 1963. *The life of bacteria*. Ed. 2. Macmillan, New York.



## VII. RESPIRATION AND CIRCULATION

### 73. CHARACTERISTICS OF RESPIRATORY MEDIA

Water and nitrogen are the two major ecological variations in the respiratory media available to organisms. The aqueous or gaseous solvent (water or nitrogen), through which the exchange of O<sub>2</sub> and CO<sub>2</sub> occurs, is the primary substance that actively ventilates the respiratory organ. The solvent is mechanically inspired by the animal. STP = standard temperature and pressure.

Variable (A)	Media			
	Aquatic		Atmospheric	
	Ocean (B)	Fresh (C)	Sea Level (D)	6,000 Meters (E)
1 Temperature, °C	-2.0 to +30.0	2.0-32.0	0.7-15.7 <sup>1</sup>	-28.1 to -15.1
2 Pressure, total, mm Hg	760-760,000	760-20,000	760	347.5-360.2
3 Density, g/liter	1.027 <sup>2</sup> (20°C)	1.000 <sup>2</sup> (4°C)	1.223-1.290 <sup>3</sup>	0.649-0.659 <sup>3</sup>
Concentration				
4 H <sub>2</sub> O, vol %	100.00 <sup>4</sup>	100.00 <sup>4</sup>	1.00 <sup>5</sup>	1.00 <sup>5</sup>
5 N <sub>2</sub> , vol %	1.03 <sup>2</sup> (15°C)	1.33 <sup>2</sup> (15°C)	78.03 (STP)	78.03 (STP)
6 CO <sub>2</sub> , vol %	0.02 <sup>2</sup> (15°C)	0.03 <sup>2</sup> (15°C)	0.03 (STP)	0.03 (STP)
7 O <sub>2</sub> , vol %	0.58 <sup>2</sup> (15°C)	0.72 <sup>2</sup> (15°C)	20.95 (STP)	20.95 (STP)
8 Salts, ‰	34.48 <sup>2</sup>	0.18 <sup>2</sup>	.....	.....
9 pH	7.5-8.4	3.2-10.6	.....	.....
10 Inert gases, vol %	Traces	Traces	0.95 (STP)	0.95 (STP)
Partial Pressure (Tension) <sup>6</sup>				
11 H <sub>2</sub> O, mm Hg	12.79 (15°C)	6.10 (4°C)	6.40 <sup>7</sup> (15°C)	0.72 <sup>7</sup> (-15°C)
12 N <sub>2</sub> , mm Hg	593.02 (STP)	593.02 (STP)	593.02 (STP)	281.06 <sup>8</sup> (STP)
13 CO <sub>2</sub> , mm Hg	0.23 <sup>2</sup> (STP)	0.23 <sup>2</sup> (STP)	0.23 (STP)	0.11 <sup>8</sup> (STP)
14 O <sub>2</sub> , mm Hg	159.52 <sup>2</sup> (STP)	159.52 <sup>2</sup> (STP)	159.52 (STP)	75.61 <sup>8</sup> (STP)
15 Inert gases, mm Hg	7.46 (STP)	7.46 (STP)	7.46 (STP)	3.42 <sup>8</sup> (STP)
Diffusion Coefficient (ml/min/sq cm/cm distance at 760 mm Hg, 20°C)				
16 N <sub>2</sub>	.....	0.000018 <sup>9</sup> (0.53) <sup>10</sup>	.....	.....
17 CO <sub>2</sub>	.....	0.000785 <sup>9</sup> (23.1) <sup>10</sup>	.....	.....
18 O <sub>2</sub>	.....	0.000034 (1) <sup>10</sup>	11.0	.....

<sup>1/</sup> Actual range is much wider. <sup>2/</sup> Average of many determinations; varies widely with conditions of measurement. <sup>3/</sup> Density determined at temperatures given in line 1. <sup>4/</sup> Less volume of solutes. <sup>5/</sup> Varies but never absent, and always of biological significance. <sup>6/</sup> Only for water in equilibrium with the atmosphere, as at the surface of ocean or lake. <sup>7/</sup> Calculated for 50% relative humidity. <sup>8/</sup> Calculated. <sup>9/</sup> Calculated from measured value for O<sub>2</sub> (line 18) and relative coefficients (lines 16 and 17). <sup>10/</sup> Values in parentheses are relative coefficients, with O<sub>2</sub> as unity.

Contributor: McCutcheon, F. Harold

References: [1] Heilbrunn, L. V. 1952. General physiology. W. B. Saunders, Philadelphia. [2] Hodgman, C. D., ed. 1956-57. Handbook of chemistry and physics. Chemical Rubber, Cleveland. [3] Krogh, A. 1919. J. Physiol. (London) 52:391. [4] Pearse, A. S. 1939. Animal ecology. McGraw-Hill, New York. [5] Sverdrup, H. U., M. W. Johnson, and R. H. Fleming. 1946. The oceans. Prentice-Hall, New York.

## 74. LUNG VENTILATION: VERTEBRATES

Values, unless otherwise indicated, are for adult animals at rest. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Species	Common Name	Sex and Body Weight kg	Respiratory Rate breaths/min	Tidal Volume <sup>1</sup> ml	Minute Volume liters/min	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
<b>Mammalia</b>							
1	<i>Homo sapiens</i>	Man					
2		Premature Newborn	1.97(1.41-2.53) <sup>b</sup>	34.4(17.2-51.6) <sup>b</sup>	12.3(4.1-20.5) <sup>b</sup>	0.396(0.204-0.588) <sup>b</sup>	3
3		Adult	3.4(2.5-4.3) <sup>b</sup>	28.6(18.2-39.0) <sup>b</sup>	20.7(13.5-27.9) <sup>b</sup>	0.584(0.471-0.697) <sup>b</sup>	3
4		♂ 68.5	11.7(10.1-13.1)	750(575-895)	7.43(5.8-10.3)		12
5			17.1(15.7-18.2) <sup>a</sup>	1,673(1,510-1,770) <sup>a</sup>	28.6(27.3-30.9) <sup>a</sup>		12
6		♀ 54.0	21.2(18.6-23.3) <sup>a</sup>	2,030(1,900-2,110) <sup>a</sup>	42.9(39.3-45.2) <sup>a</sup>		12
7			11.7(10.4-13.0)	339(285-393)	4.5(4.0-5.1)		12
8			19 <sup>a</sup>	860(836-885) <sup>a</sup>	16.3(15.9-16.8) <sup>a</sup>		12
9			30.0(25.0-35.3) <sup>a</sup>	880(490-1,270) <sup>a</sup>	24.5(17.3-31.8) <sup>a</sup>		12
10	<i>Bos taurus</i>	Cattle	♀ (403-514)	30 <sup>a</sup>	(2,700-3,400) <sup>a</sup>	(82-104) <sup>a</sup>	5
11				(27-29) <sup>a</sup>	(3,400-4,200) <sup>a</sup>	(92-114) <sup>a</sup>	5
12	<i>Canis familiaris</i>	Dog <sup>a</sup>	(16.4-30.5)	18(11-37)	320(251-432)	5.21(3.3-7.4)	6
13	<i>Capra hircus</i>	Goat	.....	19	310	5.7	1
14	<i>Cavia porcellus</i>	Guinea pig	0.466(0.274-0.941)	90(69-104)	1.8(1.0-3.9)	0.16(0.10-0.38)	4
15	<i>Equus caballus</i>	Horse <sup>a</sup>	696	11.9(10.6-13.6)	9,060(8,520-9,680)	107	2
16	<i>Felis catus</i>	Cat	2.45	26	12.4	0.322	13
17	<i>Macaca mulatta</i>	Rhesus monkey	2.68(2.05-3.08)	40(31-52)	21.0(9.8-29.0)	0.86(0.31-1.41)	4
18	<i>Mesocricetus auratus</i>	Golden hamster	0.092(0.065-0.134)	74(33-127)	0.8(0.42-1.2)	0.06(0.033-0.083)	4
19	<i>Mus musculus</i>	House mouse	0.020(0.012-0.026)	163(84-230)	0.15(0.09-0.23)	0.024(0.011-0.036)	4
20	<i>Oryctolagus cuniculus</i>	European rabbit	.....	51(38-60)	21.0(19.3-24.6)	1.07(0.80-1.14)	4, 8, 9, 14
21	<i>Phoca vitulina</i>	Harbor seal	27.5	9(6-12) <sup>a</sup>	.....	3.97 <sup>a</sup>	7
22	<i>Rattus norvegicus</i>	Norway rat	0.113(0.063-0.152)	85.5(66-114)	0.86(0.60-1.25)	0.073(0.05-0.101)	4
23	<i>Sus scrofa</i>	Swine	♂ 225	.....	.....	37	2
<b>Aves</b>							
23	<i>Anas</i> sp.	Duck	♂	42	(35-38) <sup>a</sup>	.....	11
24			♀	110	.....	.....	11
25	<i>Anser</i> sp.	Goose	♂	20	.....	.....	11
26			♀	40	.....	.....	11
27	<i>Columba livia</i>	Street pigeon	.....	(25-30)	(4.5-5.2) <sup>10</sup>	.....	11
28	<i>Gallus domesticus</i>	Chicken	♂	(12-21)	45	.....	11
29			♀	(20-37)	.....	.....	11
30	<i>Meleagris gallopavo</i>	Turkey	♂	28	.....	.....	11
31			♀	49	.....	.....	11
32	<i>Serinus canarius</i>	Canary	.....	(96-120)	.....	.....	11
<b>Reptilia</b>							
33	<i>Malaclemys terrapin centrata</i>	Southern diamondback terrapin	(0.65-0.72)	3.7 <sup>11</sup>	14 <sup>11</sup>	0.051 <sup>12</sup>	10

<sup>1/1</sup> Air inspired or expired in one respiration. <sup>1/2</sup> Light work. <sup>1/3</sup> Heavy work. <sup>1/4</sup> Lying. <sup>1/5</sup> Standing. <sup>1/6</sup> Measurements made after 30-minute rest in hammock at 24°C; values corrected to BTPS conditions (gas at body temperature and atmospheric pressure, completely saturated with water vapor). <sup>1/7</sup> Percheron gelding. <sup>1/8</sup> Cheyne-Stokes respiration. <sup>1/9</sup> Standing; supine, 30. <sup>1/10</sup> Standing; supine, 4.7. <sup>1/11</sup> At (24-29)°C.

Contributors: (a) Stroud, Robert, and Forster, Robert E., (b) Elisberg, Edward I., (c) Hemingway, Allan

References: [1] Barcroft, J., et al. 1919. Quart. J. Med. 13:35. [2] Brody, S. 1945. Bioenergetics and growth. Reinhold, New York. [3] Cross, K. W. 1953. Ph.D. Thesis. Univ. London, England. [4] Guyton, A. C. 1947. Am. J. Physiol. 150:70. [5] Hall, W. C., and S. Brody. 1933. Missouri Univ. Agr. Expt. Sta. Res. Bull. 180:11. [6] Hemingway, A., and G. S. Nahas. Unpublished. Univ. California, Los Angeles, 1953. [7] Irving, L., et al. 1936. J. Cellular Comp. Physiol. 7:137. [8] Leegard, F. 1927. Acta Med. Scand. 67:401. [9] Leegard, F. 1930.

continued

## 74. LUNG VENTILATION: VERTEBRATES

Ibid. 74:191. [10] McCutcheon, F. H. 1943. *Physiol. Zool.* 16:255. [11] Sturkie, P. D. 1954. *Avian physiology*. Comstock, Ithaca. [12] Taylor, C. 1941. *Am. J. Physiol.* 135:27. [13] Wang, S. C., and L. F. Nims. 1948. *J. Pharmacol. Exptl. Therap.* 92:187. [14] Wright, C. I. 1934. *Ibid.* 51:327.

## 75. OXYGEN CONSUMPTION

Oxygen consumption values should be used with caution, as the figures reflect order of magnitude only.

### Part I. MAMMALS

Values are cubic millimeters oxygen per gram fresh weight per hour for adult animals.

Species	Common Name	Rate	Ref-er-ence		Species	Common Name	Rate	Ref-er-ence
(A)	(B)	(C)	(D)		(A)	(B)	(C)	(D)
1 <i>Homo sapiens</i>	Man	220 <sup>1</sup>	2	14	<i>Mus musculus</i>	House mouse	3,500 <sup>1</sup>	8
2		4,000 <sup>2</sup>	10	15			1,530 <sup>6</sup>	7
3 <i>Bos taurus</i>	Cattle, ♀	184	2	16	<i>Mustela rixosa</i>	Least weasel	5,000	13
4		390	3,14	17	<i>Myotis lucifugus</i>	Little brown bat	1,500	8
5 <i>Canis familiaris</i>	Dog	580	3	18	<i>Ornithorhynchus</i> sp.	Platypus	460	6
6 <i>Cavia</i> sp.	Guinea pig	816	2	19	<i>Oryctolagus cuniculus</i>	European rabbit	640-850	3
7 <i>Citellus undulatus parryi</i>	Parry's Arctic ground squirrel	600	13	20	<i>Ovis aries</i>	Sheep	220	2
8 <i>Dasypus</i> sp.	Armadillo	201	12	21			340	3,14
9 <i>Elephas maximus</i>	Asiatic elephant, ♀ <sup>3</sup>	155	1	22	<i>Phoca vitulina</i>	Harbor seal	540	4
10 <i>Equus caballus</i>	Horse	250	3	23	<i>Phocaena phocaena</i>	Harbor porpoise	300	11
11 <i>Felis catus</i>	Cat	710	3	24	<i>Procyon cancrivorus</i>	Crab-eating raccoon	395	13
12 <i>Mesocricetus auratus</i>	Golden hamster	2,900 <sup>4</sup>	5	25	<i>Rattus</i> sp.	Rat	2,000	6
13		70 <sup>5</sup>	5	26	<i>Sorex cinereus</i>	Gray shrew	13,700	9
				27	<i>Sus scrofa</i>	Swine	220	14

1/ Resting. 2/ Maximum work. 3/ 37 years old. 4/ Awake. 5/ Hibernating. 6/ Basal.

Contributor: Flemister, Launce J.

References: [1] Benedict, F. G. 1936. *The physiology of the elephant*. Carnegie Institution of Washington, Washington, D. C. [2] Brody, S. 1945. *Bioenergetics and growth*. Reinhold, New York. [3] Heilbrunn, L. V. 1952. *An outline of general physiology*. Ed. 3. W. B. Saunders, Philadelphia. [4] Irving, L., et al. 1935. *J. Cellular Comp. Physiol.* 7:137. [5] Lyman, C. P. 1948. *J. Exptl. Zool.* 109:55. [6] Martin, C. J. 1903. *Phil. Trans. Roy. Soc. London, B*, 195:1. [7] Morrison, P. R. 1948. *J. Cellular Comp. Physiol.* 31:281. [8] Pearson, O. P. 1947. *Ecology* 28:127. [9] Pearson, O. P. 1948. *Science* 108:44. [10] Robinson, S., A. T. Edwards, and D. B. Dill. 1937. *Ibid.* 85:409. [11] Scholander, P. F. 1940. *Hvalradets Skrifter Norske Videnskaps-Akad. Oslo* 22. [12] Scholander, P. F., et al. 1943. *J. Cellular Comp. Physiol.* 21:53. [13] Scholander, P. F., et al. 1950. *Biol. Bull.* 99:259. [14] Voit, E. 1901. *Z. Biol.* 41:113.

### Part II. VERTEBRATES OTHER THAN MAMMALS

Values are cubic millimeters oxygen per gram fresh weight per hour for adult animals.

Species	Common Name	Temp., °C	Rate	Refer-ence
(A)	(B)	(C)	(D)	(E)
Aves				
1 <i>Anas</i> sp.	Duck	.....	800	8
2 <i>Anser</i> sp.	Goose	.....	592	21

continued

*Contrails*  
75. OXYGEN CONSUMPTION

Part II. VERTEBRATES OTHER THAN MAMMALS

Species	Common Name	Temp., °C	Rate	Refer- ence	
(A)	(B)	(C)	(D)	(E)	
Aves					
3	<i>Columba</i> sp.	Pigeon	.....	710	15
4	<i>Corvus corax</i>	Raven <sup>1</sup>	.....	940	17
5	<i>Gallus domesticus</i>	Chicken	.....	497	4
6	<i>Larus hyperboreus</i>	Glaucous gull	.....	1,640	17
7	<i>Passer domesticus</i>	House sparrow	.....	2,100	3
8	<i>Selasphorus sasin</i>	Allen's hummingbird	22	13,900	14
9	<i>Serinus canarius</i>	Canary	.....	2,900	3
Reptilia					
10	<i>Alligator mississippiensis</i>	American alligator	22; 19.5	8.9; 7.5	2
11	<i>Anguis fragilis</i>	Slowworm	20	40	20
12	<i>Crotalus atrox</i>	Western diamondback rattlesnake	30; 22	35.5; 16.4	19
13	<i>Iguana tuberculata</i>	Iguana, tuberculate	30; 22	52.0; 22.2	2
14	<i>Malaclemys terrapin centrata</i>	Southern diamondback terrapin	24	35	12
15	<i>Natrix natrix</i>	European water snake	20	92-150	6,9
Amphibia					
16	<i>Rana esculenta</i>	Edible frog	20	85 <sup>2</sup>	5
17			20	437 <sup>3</sup>	5
18	<i>Triturus</i> sp.	Newt	20	110	9
Pisces					
19	<i>Anguilla anguilla</i>	European freshwater eel	25	128	13
20	<i>Carassius auratus</i>	Goldfish	20	85 <sup>4</sup>	7
21			20	160 <sup>5</sup>	7
22	<i>Cyprinus carpio</i>	Carp	19.5	100	10
23	<i>Esox lucius</i>	Northern pike	18	102	11
24	<i>Lepidosiren paradoxa</i>	South American lungfish	20	42	16
25	<i>Protopterus aethiopicus</i>	East African lungfish	20	52 <sup>6</sup>	18
26			20	10 <sup>7</sup>	18
27	<i>Salmo trutta</i>	Brown trout	15	226	11
28	<i>Scomber scombrus</i>	Atlantic mackerel	20	726	1

1/ Arctic. 2/ Winter. 3/ Summer. 4/ Resting. 5/ Active. 6/ Feeding. 7/ Fasting.

Contributor: Flemister, Launce J.

References: [1] Baldwin, F. M. 1924. Proc. Iowa Acad. Sci. 30:173. [2] Benedict, F. G. 1932. Carnegie Inst. Wash. Publ. 425. [3] Benedict, F. G., and E. L. Fox. 1933. Arch. Ges. Physiol. 232:357. [4] Benedict, F. G., W. Landauer, and E. L. Fox. 1932. Conn. Univ. Storrs Agr. Expt. Sta. Bull. 177:1. [5] Bohr, C. 1899. Skand. Arch. Physiol. 10:74. [6] Cohnheim, O. 1912. Z. Physiol. Chem. 76:298. [7] Fry, F. E., and J. S. Hart. 1948. Biol. Bull. 94:66. [8] Hari, Y., and A. Kriwusch. 1918. Biochem. Z. 88:345. [9] Hill, A. V. 1911. J. Physiol. (London) 43:379. [10] Knauthe, K. 1898. Arch. Ges. Physiol. 73:490. [11] Lindstedt, P. 1914. Z. Fischerei 14:193. [12] McCutcheon, F. H. 1943. Physiol. Zool. 16:255. [13] Montuori, A. 1913. Arch. Ital. Biol. 59:213. [14] Pearson, O. P. 1950. Condor 52:145. [15] Riddle, O. 1932. Missouri Univ. Agr. Expt. Sta. Res. Bull. 166:86. [16] Sawaya, P. 1946. Univ. Sao Paulo Fac. Filosof. Cienc. Letras, Zool., Bol. 11:255. [17] Scholander, P. F., et al. 1950. Biol. Bull. 99:259. [18] Smith, H. W. 1935. J. Cellular Comp. Physiol. 6:43. [19] Sumner, F. B., and U. N. Lanham. 1942. Biol. Bull. 82:313. [20] Vernon, H. M. 1897. J. Physiol. (London) 21:443. [21] Voit, E. 1901. Z. Biol. 41:113.

continued



## 75. OXYGEN CONSUMPTION

### Part III. INVERTEBRATES OTHER THAN PROTOZOA

Values are cubic millimeters oxygen per gram fresh weight per hour for adult animals.

Class	Species	Common Name	Temp., °C	Rate	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
Chordata						
1	Cephalochordata <sup>1</sup>	<i>Branchiostoma lanceolatum</i>	Amphioxus	20; 16	45; 35	25
2	Ascidiacea	<i>Ascidia mentula</i>	Sea squirt	25	4.8	19
Echinodermata						
3	Ophiuroidea	<i>Ophioderma longicauda</i>	Brittle star	25	8-32	7.19
4	Asteroidea	<i>Asterias rubens</i>	Starfish	15	21 <sup>a</sup>	3
5				15	24 <sup>a</sup>	
6	Holothuroidea	<i>Holothuria impatiens</i>	Sea cucumber	25	17	19
Arthropoda						
7	Crustacea	<i>Astacus fluviatilis</i>	Crayfish	15	30	16
8		<i>Carcinus maenas</i>	Shore crab	15	625	19
9		<i>Homarus americanus</i>	American lobster	15	507	5
10	Insecta	<i>Aedes aegypti</i>	Yellow-fever mosquito	26	♂2,330; ♀4,200	18
11		<i>Apis mellifera</i>	Honeybee	20	17,466 <sup>4</sup>	20
12				20	87,000 <sup>5</sup>	12
13		<i>Blatta orientalis</i>	Oriental cockroach	25; 20	450; 277	8,26
14		<i>Drosophila</i> sp.	Fruit fly	20	1,560 <sup>4</sup>	6
15				20	21,800 <sup>5</sup>	
16		<i>Formica</i> sp.	Ant	20	532	21
17		<i>Melolontha</i> sp.	June beetle	20	960	2
18		<i>Musca domestica</i>	Housefly	20	1,980	10
19		<i>Phaenicia sericata</i>	Greenbottle fly	20	95,600	9
20	Onychophora	<i>Peripatus accacioi</i>	Peripatus	30; 20; 10	226; 92; 37	17
Annelida						
21	Oligochaeta	<i>Lumbricus terrestris</i>	Earthworm	20.5	138	13
22	Polychaeta	<i>Arenicola</i> sp.	Lugworm	12	30	4
23		<i>Nereis virens</i>	Clam worm	15	26	5
Mollusca						
24	Cephalopoda	<i>Sepia officinalis</i>	Cuttlefish	15	320	19
25	Bivalvia	<i>Mytilus</i> sp.	Mussel	20	22	15
26	Gastropoda	<i>Aplysia limacina</i>	Sea hare	16	30	7
27		<i>Helix pomatia</i>	Land snail	20	94	26
28		<i>Lymnaea stagnalis</i>	Freshwater snail	20; 10	123; 36.7	28
Aschelminthes						
29	Nematoda	<i>Ascaris lumbricoides</i>	Large roundworm	37	72 <sup>b</sup> ; 156 <sup>c</sup>	14
30				37	♂112; ♀61	1
31		<i>Setaria equinum</i>	Filarial worm	38	250	23
Platyhelminthes						
32	Cestoda	<i>Diphyllobothrium latum</i>	Fish tapeworm	37	243 <sup>a</sup>	11
33	Trematoda	<i>Fasciola hepatica</i>	Liver fluke	37.5	330	24
34	Turbellaria	<i>Planaria torva</i>	Flatworm	25; 2.5	75.8; 18.9	27
Cnidaria						
35	Anthozoa	<i>Anemonia sulcata</i>	Sea anemone	18	13.4	29
36	Scyphozoa	<i>Aurelia aurita</i>	Scyphomedusa	17; 31	5.0; 3.4	22

/1/ Subphylum. /2/ Baltic Sea. /3/ North Sea. /4/ Resting. /5/ True flight. /6/ Large. /7/ Small. /8/ Proglottids.

Contributor: Flemister, Launce J.

References: [1] Adam, W. 1932. Z. Vergleich. Physiol. 16:229. [2] Battelli, F., and L. Stern. 1913. Biochem. Z. 56:50. [3] Block, K. J., and C. Schlieper. 1953. Kiel. Meeresforsch. 9:201. [4] Borden, M. A. 1931. J. Marine Biol. Assoc. U. K. 17:709. [5] Bosworth, M. W., et al. 1936. J. Cellular Comp. Physiol. 9:77. [6] Chadwick, L. E.

continued

## 75. OXYGEN CONSUMPTION

### Part III. INVERTEBRATES OTHER THAN PROTOZOA

1947. Biol. Bull. 93:229. [7] Cohnheim, O. 1912. Z. Physiol. Chem. 76:298. [8] Davis, J. G., and W. K. Slater. 1928. Biochem. J. 22:331. [9] Davis, R. A., and G. Fraenkel. 1940. J. Expt. Biol. 17:402. [10] Edwards, G. A. 1946. J. Cellular Comp. Physiol. 27:53. [11] Friedheim, E. A., and J. G. Baer. 1933. Biochem. Z. 264:329. [12] Jongbloed, J., and C. A. G. Wiersma. 1934. Z. Vergleich. Physiol. 21:519. [13] Konopacki, M. 1907. Mem. Acad. Sci. Cracovie, p. 357. [14] Kreuger, F. 1936. Zool. Jahrb. Abt. Allgem. Zool. Physiol. Tiere 57:1. [15] Krogh, A. 1941. The comparative physiology of respiratory mechanisms. Univ. Pennsylvania Press, Philadelphia. [16] Lindstedt, P. 1914. Z. Fischerei 14:193. [17] Mendes, E. G., and P. Sawaya. 1957. Ciencia Cult. (Sao Paulo) 9:120. [18] Mercado, T. I., H. L. Trembley, and T. von Brand. 1956. Physiol. Comparata Oecol. 4:200. [19] Montuori, A. 1913. Arch. Ital. Biol. 59:213. [20] Paron, M. 1909. Ann. Sci. Nat. Zool. 9:1. [21] Slowzoff, B. 1909. Biochem. Z. 19:497. [22] Thill, H. 1937. Z. Wiss. Zool. 150:51. [23] Toryu, Y. 1934. Sci. Rept. Tohoku Imp. Univ., Ser. 4, 9:61. [24] Van Grembergen, G. 1948. Enzymologica 13:241. [25] Vernon, H. M. 1896. J. Physiol. (London) 19:18. [26] Vernon, H. M. 1897. Ibid. 21:443. [27] Von Brand, T. 1936. Physiol. Zool. 9:530. [28] Von Brand, T., and B. Mehlman. 1953. Biol. Bull. 104:301. [29] Von Buddenbrock, W. 1938. Z. Vergleich. Physiol. 26:303.

### Part IV. PROTOZOA

Values are cubic millimeters oxygen per million cells per hour for mature protozoa.

	Class	Species	Temp., °C	Rate	Refer- ence
	(A)	(B)	(C)	(D)	(E)
1	Ciliata	<i>Paramecium aurelia</i> <sup>1</sup>	35	1,512	4
2			30	831	
3			25	616	
4			20	354	
5	Sporozoa	<i>Plasmodium cathemerium</i>	38	0.25	1
6	Rhizopoda	<i>Amoeba chaos chaos</i> <sup>2</sup>	35	17,749	3
7			30	13,244	
8			25	9,010	
9			20	7,050	
10			15	5,040	
11	Mastigophora	<i>Trypanosoma gambiense</i>	37	1.70	5
12			30	0.38	
13			<i>Chilomonas paramecium</i> <sup>3</sup>	25	

<sup>1/1</sup> No substrate. <sup>2/2</sup> Fed. <sup>3/3</sup> Bacteria-free.

Contributor: Flemister, Launce J.

References: [1] Maier, J., and L. T. Coggeshall. 1941. J. Infect. Diseases 69:87. [2] Mast, S. O., et al. 1936. J. Cellular Comp. Physiol. 8:125. [3] Pace, D. M., and W. H. Belda. 1944. Biol. Bull. 86:146. [4] Pace, D. M., and K. K. Kimura. 1944. J. Cellular Comp. Physiol. 24:173. [5] Von Brand, T., E. J. Tobie, and B. Mehlman. 1950. Ibid. 35:273.

## 76. RESPIRATION RATES

### Part I. BACTERIA

Rate and degree of respiration may be affected by numerous factors, such as strain characteristics, composition of growth medium, age and number of cells in an inoculum, origin of inoculum, ages of culture harvested for study, nature of solution used for washing, number of washings, and composition of the respiratory system. Data are for bacterial suspensions in the presence of glucose.

Species	Temp. °C	Culture Age hr	QO <sub>2</sub> μl/mg dry wt/hr	Refer- ence
(A)	(B)	(C)	(D)	(E)
1 <i>Aerobacter aerogenes</i>	36; 30	17; 48	47; 50	1,2
2 <i>Azotobacter chroococcum</i>	22	36	2,000-10,000	9
3 <i>Bacillus subtilis</i>	37	6-8	170	6
4 <i>B. subtilis</i> (spores)	32	98-147	10	4
5 <i>Corynebacterium</i> sp.	30	48-96	67	8
6 <i>Escherichia coli</i>	40; 32	20	200; 272	1,7
7 <i>Lactobacillus bulgaricus</i>	45; 37	8	55; 34	14
8 <i>Micrococcus luteus</i>	35	30-34	15	10
9 <i>Mycobacterium avium</i>	37	84	1	11
10 <i>M. tuberculosis</i>	38	252	4	5
11 <i>Pseudomonas fluorescens</i>	26	20	58	12
12 <i>Streptococcus pyogenes</i> , C 203 M	37.5	4	57-163	13
13 <i>S. pyogenes</i> , C 203 S	37.5	4	99-113	13
14 <i>Streptomyces coelicolor</i>	.....	72	35	3

Contributor: Silverman, Milton

References: [1] Ajl, S. J. 1950. J. Bacteriol. 59:499. [2] Ajl, S. J., and T. O. Wong. 1951. Ibid. 61:379. [3] Cochrane, V. W., and M. Gibbs. 1951. Ibid. 61:305. [4] Crook, P. G. 1952. Ibid. 63:193. [5] Edson, N. L., and G. J. Hunter. 1943. Biochem. J. 37:563. [6] Gary, N. D., and R. C. Bard. 1952. J. Bacteriol. 64:501. [7] Krebs, H. A. 1937. Biochem. J. 31:2095. [8] Levine, S., and L. O. Krampitz. 1952. J. Bacteriol. 64:645. [9] Meyerhof, O., and D. Burk. 1928. Z. Physik. Chem. (Leipzig), A, 139:117. [10] Nunheimer, T. D., and F. W. Fabian. 1942. J. Bacteriol. 44:215. [11] Oginsky, E. L., P. H. Smith, and M. Soltorovsky. 1950. Ibid. 59:29. [12] Sebek, O. K., and C. I. Randles. 1952. Ibid. 63:693. [13] Sevag, M. G., and M. Shelburne. 1942. Ibid. 43:411. [14] Stein, R. M., and W. L. Frazier. 1941. Ibid. 42:501.

### Part II. MYXOPHYTA AND FUNGI

Method (column D): Mano = manometric; Chem = chemical; Volu = volumetric. Substrate (column E): Endo = endogenous; CHO = carbohydrates; Org = organic compounds. QCO<sub>2</sub> (column H): Values *not* in parentheses are for aerobic CO<sub>2</sub> production, those in parentheses are for anaerobic CO<sub>2</sub> production.

Species	Material	Temp. °C	Method	Sub- strate	Specification	Respiration Rate μl/mg dry wt/hr <sup>1</sup>		Respiratory Quotient CO <sub>2</sub> /O <sub>2</sub>	Refer- ence
						QO <sub>2</sub>	QCO <sub>2</sub>		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Myxomycetes									
1 <i>Physarum polycephalum</i>	Plasmodium	22	Mano	Endo	50 mg/vessel	1.4 <sup>2</sup>	1.0 <sup>2</sup> (0.24 <sup>2</sup> )	0.75-0.85	2
Phycomycetes									
2 <i>Phycomyces blakesleeana</i>	Mycelia	20	Chem	CHO	1.5; 3.5; 7 da	.....	27; 13; 3	.....	14
3 <i>Rhizopus sexualis</i>	Mycelia	20	Chem	Org	52 hr	.....	25.7 <sup>3</sup>	.....	6
Ascomycetes									
4 <i>Neurospora crassa</i>	Mycelia	30	Mano	Org	Endogenous	11-38	(0-5)	.....	13
5 <i>Saccharomyces cerevisiae</i> R	Cell suspension	.....	Mano	CHO	No stored reserves	83-109	370-432 (278-299)	.....	7

<sup>1</sup>/ Unless otherwise indicated. <sup>2</sup>/ μl per mg wet weight per hour. <sup>3</sup>/ mg per g dry weight per hour.

continued

## 76. RESPIRATION RATES

### Part II. MYXOPHYTA AND FUNGI

	Species	Material	Temp. °C	Method	Sub- strate	Specification	Respiration Rate μl/mg dry wt/hr <sup>1</sup>		Respiratory Quotient CO <sub>2</sub> /O <sub>2</sub>	Ref- er- ence
							QO <sub>2</sub>	QCO <sub>2</sub>		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Ascomycetes										
6	<i>Saccharomyces cerevisiae</i> R	Cell suspen- sion	.....	Mano	CHO	Fat reserves	76	249 (322)	.....	7
7			.....	Mano	CHO	Glycogen reserves	0	63 (116)	.....	
8	<i>S. cerevisiae</i> U	Cell suspen- sion	.....	Mano	CHO	No stored reserves	10-137	160-348 (276-284)	.....	7
9			.....	Mano	CHO	Fat reserves	125	151 (261)	.....	
10			.....	Mano	CHO	Glycogen reserves	47	82 (83)	.....	
11	<i>Schizosaccharomyces octosporus</i>	Cell suspen- sion	30	Mano	CHO	Endogenous; + glu- cose	21; 90	(0.1)	.....	12
Basidiomycetes										
12	<i>Agaricus campestris</i>	Growing cul- ture	25	Volu	....	.....	1.9-2.9	2.3-4.0	0.70-0.90	4
13	<i>Puccinia graminis</i>	Urediospore	30	Mano	Endo	PO <sub>4</sub> buffer, pH 6.5, ungerminated	1.6 <sup>2</sup>	1.1 <sup>2</sup>	0.65	10
14			30	Mano	Endo	Germinated	1.4 <sup>2</sup>	1.0 <sup>2</sup>	0.70	
15	<i>Ustilago sphaerogena</i>	Sporidia <sup>4</sup>	.....	Mano	Endo	.....	75	.....	.....	1
16			.....	Mano	CHO	+ Sugars	375	.....	.....	
Fungi Imperfecti										
17	<i>Alternaria</i> sp.	Pellets	23-25	Chem	CHO	.....	.....	.....	1.26-1.31	3
18	<i>Aspergillus niger</i>	Mycelia	35; 19	Chem	CHO	+ Glucose	.....	.....	1.30; 0.98	9
19			35; 18	Chem	CHO	+ Sucrose	.....	.....	1.22; 0.91	
20			36	Chem	Org	+ Tartrate	.....	.....	1.35-2.03	
21			36	Chem	Org	+ Glycerol	.....	.....	0.82-0.86	
22	<i>Candida albicans</i>	Cell suspen- sion <sup>5</sup>	30	Mano	Endo	.....	5	.....	.....	8
23			30	Mano	CHO	+ Glucose	40	.....	.....	
24	<i>Cladosporium</i> spp.	Pellets	23-25	Chem	CHO	5 strains	.....	.....	1.10-1.28	3
25	<i>Fusarium trichothecioides</i>	Mycelia <sup>5</sup> (1 da old)	30	Mano	Endo	1; 4 hr	40; 13	31; 11	0.78; 0.84	5
26			30	Mano	CHO	1; 4 hr (+ glucose)	34; 39	64; 56	1.85; 1.55	
27		Mycelia <sup>5</sup> (3 da old)	30	Mano	Endo	1; 4 hr	14; 13	14; 12	1.01; 0.92	
28			30	Mano	CHO	1; 4 hr (+ glucose)	14; 13	19; 26	1.36; 1.97	
29	<i>Helminthosporium gramineum</i>	Pellets	23-25	Chem	CHO	.....	.....	.....	1.31	3
30	<i>Penicillium notatum</i>	Mycelia <sup>5</sup>	20-24	Mano	Endo	0; 1 da	6.5; 1.7	.....	.....	15
31		Mycelia	20-24	Mano	CHO	2; 4; 7 da	6; 16; 2	.....	.....	
32	<i>Torulopsis utilis</i>	Cell suspen- sion	30	Mano	Endo	+ Glycine	3.7 <sup>2</sup>	.....	0.86	11
33			30	Mano	Endo	+ Urea	3.5 <sup>2</sup>	.....	1.15	
34			30	Mano	Endo	+ α-Alanine	5.2 <sup>2</sup>	.....	0.89	
35			30	Mano	Endo	+ β-Alanine	4.2 <sup>2</sup>	.....	1.16	

<sup>1</sup>/ Unless otherwise indicated. <sup>2</sup>/ μl per mg wet weight per hour. <sup>4</sup>/ Washed. <sup>5</sup>/ Starved. <sup>6</sup>/ Homogenized.

Contributors: (a) Darby, Richard T., and Mandels, Gabriel R., (b) Henderson, Lavaniel L., Sr.

References: [1] Allen, P. J. 1948. Am. J. Botany 35:799. [2] Allen, P. J., and W. H. Price. 1950. Ibid. 37:393. [3] Birkinshaw, J. H., et al. 1931. Phil. Trans. Roy. Soc. London, B, 220:99. [4] Chevillard, L., A. Meyer, and L. Plantefol. 1930. Ann. Physiol. Physicochim. Biol. 6:506. [5] Gould, B. S., and A. A. Tytell. 1941. J. Gen. Physiol. 24:655. [6] Hawker, L. E., and P. M. Hepden. 1962. Ann. Botany (London) 26:619. [7] Lindegren, C. C. 1946. Arch. Biochem. Biophys. 9:353. [8] Nickerson, W. J. 1946. Am. J. Botany 33:831. [9] Porievitch, K. 1905. Ann. Sci. Nat. Bot. Biol. Vegetale, Ser. 9, 1:1. [10] Shu, P., K. G. Tanner, and G. A. Ledingham. 1954. Can. J. Botany 32:16. [11] Sperber, E. 1945. Arkiv Kemi 21A(3):1. [12] Spiegelman, S., and M. Nozawa. 1945. Arch. Biochem. Biophys. 6:303. [13] Strauss, B. S. 1952. Ibid. 36:33. [14] Wassink, E. C. 1934. Rec. Trav. Bot. Neerl. 31:583. [15] Wolf, F. T. 1947. Arch. Biochem. Biophys. 13:83.

continued



*Continued*  
76. RESPIRATION RATES

Part III. LICHENS, ALGAE, AND BRYOPHYTES

Method (column C): Mano = manometric; Cond = conductometric; Chem = chemical. Figures in parentheses are control or endogenous values.

	Division or Class, and Species	Temp. °C	Method	Respiration Rate μl/100 mg dry wt/hr <sup>1</sup>		Respiratory Quotient CO <sub>2</sub> /O <sub>2</sub>	Refer- ence
				Q <sub>O<sub>2</sub></sub>	Q <sub>CO<sub>2</sub></sub>		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
Lichenes							
Ascolichenes							
1	<i>Alectoria nigricans</i>	30; 10; 0	Mano	33; 14; 8	.....	.....	16
2	<i>Cladonia sylvatica</i>	30; 10; 0	Mano	24; 6.8; 2.9	.....	.....	16
3	<i>Parmelia nigrociliata</i>	30; 10; 0	Mano	25; 13; 4	.....	.....	16
4	<i>Peltigera aphthosa</i>	30; 10; 0	Mano	90; 33; 17	.....	.....	16
5	<i>Umbilicaria proboscidea</i>	30; 10; 0	Mano	18; 6.5; 3.5	.....	.....	16
6	<i>Usnea dasygoga</i>	.....	Cond	.....	60-90 <sup>2</sup>	.....	15
Algae							
Cyanophyta							
7	<i>Anabaena variabilis</i> <sup>a</sup>	25	Mano	170 (840)	.....	1.0 (1.1)	14
8	<i>Anacystis nidulans</i> <sup>a</sup>	39; 25	Mano	200 (500); 30 (160)	.....	1.1 (1.1); 0.9 (1.0)	14
9	<i>Nostoc muscorum</i> <sup>a</sup>	25	Mano	110 (440)	.....	.....	14
Chlorophyta							
10	<i>Chlorella pyrenoidosa</i>	20	Mano	1,700	.....	1.39	7
11		18; 3.5	Mano	890 (430); 200 (150)	.....	(0.94); (0.98)	6
12	<i>Cladophora rupestris</i>	20	Chem	33	.....	.....	8
13	<i>Spirogyra majuscula</i> <sup>4</sup>	10.4	Chem	0.5 <sup>5</sup>	.....	.....	3
14	<i>Ulothrix flacca</i>	.....	Chem	160	.....	.....	9
Phaeophyta							
15	<i>Ectocarpus siliculosus</i>	12	Chem	41 <sup>5</sup>	.....	.....	10
16	<i>Fucus vesiculosus</i>	14	Chem	.....	12.7 <sup>6</sup>	.....	2
17	<i>Laminaria digitata</i>	5	Mano	.....	0.9 <sup>7</sup>	0.67	13
Rhodophyta							
18	<i>Polysiphonia violacea</i>	11	Chem	.....	107	1.02	10
19	<i>Porphyra laciniata</i>	17	Chem	39	.....	.....	12
Bryophyta							
Musci							
20	<i>Hylocomium squarrosum</i>	30; 20; 5	Chem	.....	100; 61; 15	.....	17
21	<i>Hypnum cupressiforme</i>	18.5	Chem	2-30 <sup>9</sup>	.....	.....	5
22	<i>Mnium undulatum</i>	.....	.....	.....	7.5-97.0 <sup>2</sup>	.....	11
23	<i>Polytrichum juniperinum</i> <sup>a</sup>	18	.....	.....	1.2 <sup>5</sup> -0.7 <sup>6</sup>	1.00-0.65	1
24	<i>Sphagnum girgensohnii</i>	30; 20; 5	Chem	.....	130; 71; 20	.....	17
Hepaticae							
25	<i>Marchantia polymorpha</i>	20	Chem	.....	0.6 <sup>7</sup>	.....	4
26	<i>Riccia fluitans</i>	25	Mano	250-300	.....	.....	18

/1/ Unless otherwise indicated. /2/ Effect of moisture. /3/ After 24-hour dark starvation. /4/ Effect of pH. /5/ μl per 100 mg wet weight per hour. /6/ μg per 100 g wet weight per hour. /7/ μl per sq cm per hour. /8/ Shoots or tops only; values show change caused by growth, development, or maturation.

Contributors: (a) Mandels, Gabriel R., and Darby, Richard T., (b) Myers, Jack, (c) Henderson, Lavaniel L., Sr.

References: [1] Bastit, E. 1891. Rev. Gen. Botan. 3:255. [2] Bidwell, R. C. S. 1963. Can. J. Botany 41(1):155. [3] Bode, H. R. 1925. Jahrb. Wiss. Botan. 65:352. [4] Boysen-Jensen, P., and D. Müller. 1929. Ibid. 70:503. [5] Fraymouth, J. 1928. Ann. Botany (London) 42:75. [6] French, C. S., H. I. Kohn, and P. S. Tang. 1934. J. Gen. Physiol. 18:193. [7] Gaffron, H. 1939. Biol. Zentr. 59:288. [8] Gessner, F. 1940. Jahrb. Wiss. Botan. 89:1. [9] Harder, R. 1915. Ibid. 56:254. [10] Hoffmann, C. 1929. Ibid. 71:214. [11] Jönsson, B. 1894. Compt. Rend. 119:440. [12] Kniep, H. 1914. Intern. Rev. Ges. Hydrobiol. Hydrog. 7:1. [13] Krascheninnikoff, T. 1926. Compt. Rend. 182:939. [14] Kratz, W. A., and J. Myers. 1955. Plant Physiol. 30:275. [15] Neubauer, A. F. 1938. Beitr. Biol. Pflanz. 25:273. [16] Scholander, P. F., et al. 1952. Am. J. Botany 39:707. [17] Stålfelt, M. G. 1937. Planta 27:30. [18] Usami, S. 1937. Acta Phytochim. (Japan) 9:287.

*continued*

## 76. RESPIRATION RATES

### Part IV. TRACHEOPHYTA

Method (column E): Mano = manometric; Chem = chemical; Cond = conductometric. Figures in parentheses are control or endogenous values.

	Species	Common Name	Condition or Part	Temp. °C	Method	Respiration Rate $\mu\text{l}/100 \text{ mg wet wt}/\text{hr}^1$		Respiratory Quotient $\text{CO}_2/\text{O}_2$	Reference
						$\text{QO}_2$	$\text{QCO}_2$		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Seeds									
1	<i>Acer saccharum</i>	Sugar maple	Resting	.....	.....	.....	14	.....	42
2	<i>Avena sativa</i> <sup>2</sup>	Common oat	Coleoptile, segment	30	Mano	47-39	.....	.....	12
3	<i>Cucurbita pepo</i>	Pumpkin	Germinating	25	Chem	.....	10-117	0.94-0.62	49
4	<i>Fagopyrum esculentum</i>	Buckwheat	Germinating	25	.....	.....	41-306	0.8-1.0	48
5	<i>Glycine soja</i> <sup>3</sup>	Soybean	Germinating	.....	Mano	.....	.....	0.93-0.87	24
6	<i>Gossypium hirsutum</i> <sup>4,5</sup>	Upland cotton	Resting	26	Mano	.....	0.03-6.0	0.96-1.12	43
7	<i>Helianthus annuus</i>	Common sunflower	Resting	28	Mano	.....	.....	1.05	67
8			Germinating	25	Chem	.....	41-407	0.85-0.50	49
9	<i>Hordeum vulgare</i> <sup>4</sup>	Barley	Resting	37.8	Chem	.....	0.002 <sup>6</sup> -0.36 <sup>6</sup>	.....	7
10	<i>Juglans regia</i>	Persian walnut	Resting	28	Mano	.....	.....	0.52	67
11	<i>Juniperus virginiana</i>	Eastern red cedar	Resting	25	Mano	.....	0.05	0.76	63
12			Germinating	25	Mano	.....	6.6-25	0.84-0.97	63
13	<i>Malus pumila</i>	Common apple	Resting	19	.....	.....	2.8 <sup>5</sup>	0.86	35
14	<i>Medicago sativa</i>	Alfalfa	Resting	18	Mano	38	.....	1.08	30
15			Germinating	18	Mano	106	.....	0.86	30
16	<i>Oryza sativa</i>	Rice	Resting	.....	Mano	0.03 <sup>6</sup>	.....	1.15	25
17			Moist	.....	Mano	2.8 <sup>6</sup>	.....	1.96	25
18			Germinating	.....	Mano	4.9 <sup>6</sup>	.....	1.98	25
19			Seedling	.....	Mano	1.06 <sup>6</sup>	.....	1.00	25
20	<i>Phaseolus vulgaris</i>	Kidney bean	Germinating	.....	.....	.....	65	.....	50
21	<i>Pinus radiata</i>	Monterey pine	Resting	.....	Mano	.....	0.0013 <sup>5</sup>	.....	84
22	<i>Pisum sativum</i>	Garden pea	Resting	28	Mano	.....	.....	1.00	67
23	<i>P. sativum</i> <sup>7</sup>	Garden pea	Intact	20	Chem	.....	15 (35)	4.9 (1.1)	65
24	<i>Prunus amygdalus</i> <sup>3</sup>	Almond	Germinating	.....	Mano	.....	.....	0.7-0.86	24
25	<i>P. domestica</i>	Garden plum	Moist	25	Mano	4.7	.....	0.91	72
26	<i>P. persica</i>	Peach	Moist	25	Mano	5.8	.....	0.68	72
27	<i>Raphanus sativus</i>	Garden radish	Resting	20	Mano	7.0	.....	0.86	30
28			Germinating	20	Mano	1.03	.....	0.58	30
29	<i>Triticum aestivum</i> <sup>4</sup>	Wheat	Resting	38	Chem	.....	0.005 <sup>6</sup> -0.16 <sup>6</sup>	.....	8
30	<i>T. aestivum</i> <sup>4</sup>	Wheat	Germinating	38	Chem	.....	0.014 <sup>6</sup> -0.53 <sup>6</sup>	.....	8
31	<i>T. aestivum</i>	Wheat	Seedling	18	Chem	.....	21	.....	76
32	<i>Vicia faba</i>	Broad bean	Resting	28	Mano	.....	.....	0.99	67
33			Germinating	25	Chem	.....	.....	1.23-0.82	78
34			Seedling	20	Chem	.....	13	.....	76
35	<i>Zea mays</i> <sup>4</sup>	Corn	Resting	22	Chem	.....	0.24 <sup>6</sup> -1.2 <sup>6</sup>	.....	2
36	<i>Z. mays</i>	Corn	Germinating	25	.....	.....	10-127	0.75-1.0	48
37	<i>Z. mays</i>	Corn	Seedling	18	Chem	15	.....	.....	76
Roots									
38	<i>Allium cepa</i> <sup>3</sup>	Garden onion	Segment	25	Mano	1,390 <sup>5</sup> -1,140 <sup>5</sup>	.....	0.99-1.07	10
39	<i>Beta vulgaris</i> <sup>5,7</sup>	Beet	Intact	25	Chem	.....	0.9-0.6	0.8	17
40	<i>B. vulgaris</i> <sup>5</sup>	Beet	Segment	25	Mano	70 <sup>5</sup> -180 <sup>5</sup> -110 <sup>5</sup>	.....	1.01-0.85	77
41	<i>Chrysanthemum morifolium</i>	Florist's chrysanthemum	Intact	28	Mano	.....	.....	0.93	67
42	<i>Daucus carota</i> <sup>5</sup>	Carrot	Intact	24	Chem	.....	3.3-1.5	1.10-1.18	64
43			Intact	10	Chem	.....	1.5-0.5	1.08-1.01	64
44			Intact	0.5	Chem	.....	0.44-0.22	0.92-1.16	64
45	<i>Glycine soja</i>	Soybean	Nodule	28	Mano	60 <sup>6</sup> -430 <sup>6</sup>	.....	1.0-2.0	1
46	<i>Gossypium herbaceum</i> <sup>3</sup>	Levant cotton	Intact	38	Chem	.....	380 <sup>6</sup> -73 <sup>6</sup>	.....	38
47	<i>Hordeum vulgare</i> <sup>7</sup>	Barley	Intact	20	Cond	.....	484 <sup>6</sup> -740 <sup>6</sup>	.....	86
48	<i>Ipomoea batatas</i>	Sweet potato	Intact	35	Chem	.....	5.6-6.2	.....	41

<sup>1</sup>/ Unless otherwise indicated. <sup>2</sup>/ Effect of substrate. <sup>3</sup>/ Effect of growth, development, or maturation. <sup>4</sup>/ Effect of moisture. <sup>5</sup>/ Effect of storage or starvation. <sup>6</sup>/  $\mu\text{l}$  per 100 mg dry weight per hour. <sup>7</sup>/ Effect of oxygen.

continued

## 76. RESPIRATION RATES

### Part IV. TRACHEOPHYTA

	Species	Common Name	Condition or Part	Temp. °C	Method	Respiration Rate $\mu\text{l}/100 \text{ mg wet wt}/\text{hr}^1$		Respiratory Quotient $\text{CO}_2/\text{O}_2$	Reference
						$\text{QO}_2$	$\text{QCO}_2$		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
Roots									
49	<i>Ipomoea batatas</i>	Sweet potato	Intact	25	Chem	.....	3.2-4.0	.....	41
50			Intact	15	Chem	.....	1.4-1.9	.....	41
51			Segment	25	Mano	96	.....	1.0	81
52	<i>Lycopersicon esculentum</i>	Tomato	Excised	25	Mano	600 <sup>b</sup> -800 <sup>b</sup>	.....	1.0	37
53	<i>Malus pumila</i>	Common apple	Intact	14	Chem	.....	26 <sup>b</sup>	0.73	83
54	<i>Oryza sativa</i> <sup>3</sup>	Rice	Intact	15-18	.....	180 <sup>b</sup> -230 <sup>b</sup>	.....	.....	46
55	<i>Pastinaca sativa</i>	Parsnip	Intact	22; 1.5	Chem	.....	2.7; 1.1	.....	2
56	<i>Raphanus sativus</i>	Garden radish	Intact	28	Mano	.....	.....	0.99	67
57	<i>Triticum aestivum</i> <sup>9</sup>	Wheat	Intact	20	Chem	25 <sup>b</sup> (10 <sup>b</sup> )	.....	.....	51
58	<i>Vicia faba</i>	Broad bean	Excised	26	Mano	.....	.....	1.46	69
Stems									
59	<i>Acer rubrum</i> <sup>3</sup>	Red maple	Xylem	25	Mano	3.7-2.3	.....	.....	31
60			Cambium	25	Mano	22.4	.....	.....	31
61			Phloem	25	Mano	16.9	.....	.....	31
62	<i>Asparagus officinalis</i> <sup>5</sup>	Garden asparagus	Shoot	30	Chem	.....	915 <sup>b</sup> -254 <sup>b</sup>	.....	9
63			Intact	24	Chem	.....	35.4-13.2	1.04-0.95	64
64			Intact	10	Chem	.....	9.7-3.6	1.03-0.86	64
65			Intact	0.5	Chem	.....	3.0-2.0	0.98-0.95	64
66	<i>Elodea canadensis</i>	Canada waterweed	Shoot	20	Mano	90 <sup>b</sup>	.....	.....	29
67	<i>Equisetum telmateia</i>	Giant horsetail	Shoot or top	20	Mano	.....	6	0.78	52
68			Fruiting shoot or top	20	Mano	.....	100	0.83	52
69			Intact	Room	Mano	.....	9.6	0.80	57
70			Branchlet	Room	Mano	.....	19	0.69	57
71	<i>Fraxinus nigra</i> <sup>3</sup>	Black ash	Xylem	25	Mano	.....	31.3-1.4	.....	31
72			Phloem	25	Mano	.....	16.7	.....	31
73			Cambium	25	Mano	.....	22	.....	31
74	<i>Gladiolus</i> sp.	Gladiolus	Corm	23	Chem	.....	8.5 <sup>b</sup>	.....	21
75	<i>Gossypium herbaceum</i> <sup>3</sup>	Levant cotton	Intact	38	Chem	.....	168 <sup>b</sup> -42 <sup>b</sup>	.....	38
76	<i>Helianthus annuus</i>	Common sunflower	Shoot	25;10;5	Chem	.....	483 <sup>b</sup> ;141 <sup>b</sup> ;76 <sup>b</sup>	.....	44
77	<i>Ipomoea batatas</i> <sup>5</sup>	Sweet potato	Tuber	30	Chem	.....	1.4-7.0-2.4	.....	36
78	<i>Lycopersicon esculentum</i> <sup>3</sup>	Tomato	Segment	28	Mano	420 <sup>b</sup> -350 <sup>b</sup>	.....	0.91-0.95	44
79	<i>Malus pumila</i> <sup>9</sup>	Common apple	Intact	6	Chem	.....	1.2-4.6	.....	20
80	<i>Nicotiana glauca</i> x <i>N. langsdorffii</i> <sup>10,11</sup>	Tobacco	Callus	30	Mano	380 <sup>b</sup>	.....	1.0	56
81	<i>Phaseolus vulgaris</i> <sup>11,12</sup>	Kidney bean	Intact	30	Mano	28 <sup>b</sup> -710 <sup>b</sup>	.....	0.9-1.1	74
82	<i>P. vulgaris</i> <sup>12</sup>	Kidney bean	Shoot	24	Chem	.....	190 <sup>b</sup> (150 <sup>b</sup> )	.....	16
83	<i>Pisum sativum</i> <sup>5</sup>	Garden pea	Segment	25	Mano	334 <sup>b</sup> (532 <sup>b</sup> )	.....	0.98 (1.07)	18
84	<i>Prunus laurocerasus</i> <sup>5</sup>	Laurel cherry	Shoot	22.5	Chem	.....	14.4-2.6	.....	6
85	<i>Quercus coccifera</i> <sup>3</sup>	Kermes oak	Segment	21	Mano	.....	31-11	0.91-0.83	58
86	<i>Raphanus raphanistrum</i>	Wild radish	Intact	Room	Mano	.....	10.5	0.87	57
87	<i>Salix herbacea</i>	Pygmy willow	Shoot	20;10;0	Chem	.....	23.4;9.1;2.5	.....	80
88	<i>Solanum tuberosum</i> <sup>5</sup>	Potato	Tuber	24	Chem	.....	0.6-0.3	1.02-0.75	64
89			Tuber	10	Chem	.....	0.2-0.15	0.86-0.99	64
90			Tuber	0.5	Chem	.....	0.07-0.15	0.45-0.66	64
91	<i>Taxus baccata</i>	English yew	Shoot	28	Mano	.....	.....	0.97	67
92	<i>Triticum aestivum</i>	Wheat	Shoot	13; 8	Mano	.....	29; 19	0.98; 1.03	4

/1/ Unless otherwise indicated. /3/ Effect of growth, development, or maturation. /5/ Effect of storage or starvation. /6/  $\mu\text{l}$  per 100 mg dry weight per hour. /8/ Effect of inorganic nutrition, salts. /9/ Effect of precooling. /10/ Effect of pH. /11/ Effect of metabolic poisons. /12/ Effect of herbicides.

continued



## 76. RESPIRATION RATES

### Part IV. TRACHEOPHYTA

	Species	Common Name	Condition or Part	Temp. °C	Method	Respiration Rate $\mu\text{l}/100 \text{ mg wet wt}/\text{hr}^1$		Respiratory Quotient $\text{CO}_2/\text{O}_2$	Reference
						$\text{QO}_2$	$\text{QCO}_2$		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Stems									
93	<i>Vicia faba</i>	Broad bean	Intact	Room	Mano	.....	6.2	.....	57
94			Shoot	21	Mano	.....	62.6	0.90	57
95			Shoot	21	Mano	.....	48.8	0.87	57
96	<i>Zea mays</i> <sup>1a</sup>	Corn	Shoot	30	Mano	.....	760 <sup>b</sup>	.....	32
Leaves									
97	<i>Acer pseudoplatanus</i>	Plane-tree maple	Intact	10	Chem	.....	33	.....	66
98	<i>Allium cepa</i>	Garden onion	Bulb	22	Chem	.....	2.1	.....	2
99	<i>Antirrhinum majus</i>	Snapdragon	Intact	20	Mano	.....	16	0.88	52
100	<i>Asparagus albus</i>	White asparagus	Tendrils, phyllode, or cladode	Room	Mano	.....	22.3	0.78	57
101	<i>Beta vulgaris</i>	Beet	Intact	27	Chem	.....	23	.....	53
102	<i>Betula nana</i>	Dwarf arctic birch	Intact	20; 10	Chem	.....	66; 26	.....	80
103	<i>Catalpa bignonioides</i>	Southern catalpa	Intact	14	Chem	.....	18-25	.....	66
104	<i>Citrus limon</i>	Lemon	Intact	.....	Mano	7.7 <sup>1a</sup> -9.5 <sup>1a</sup>	.....	.....	82
105	<i>C. sinensis</i>	Sweet orange	Intact	.....	Mano	9.6 <sup>1a</sup> -12.9 <sup>1a</sup>	.....	.....	82
106	<i>Elodea canadensis</i>	Canada waterweed	Intact	.....	Mano, Chem	.....	.....	8.4	68
107	<i>Fagus sylvatica</i>	European beech	Intact	20	Chem	.....	1 <sup>1a</sup> -5 <sup>1a</sup>	.....	14
108	<i>Fragaria sp.</i> <sup>3</sup>	Strawberry	Intact	24.5	Chem	.....	10 <sup>1a</sup> -5 <sup>1a</sup>	.....	3
109	<i>Fraxinus excelsior</i> <sup>1a, 2a</sup>	European ash	Intact	20	Chem	.....	1 <sup>1a</sup> -6 <sup>1a</sup>	.....	14
110	<i>Gladiolus gandavensis</i>	Breeder's gladiolus	Intact	24	Mano	.....	18	0.64	52
111	<i>Gossypium herbaceum</i> <sup>5</sup>	Levant cotton	Intact	38	Chem	.....	224 <sup>b</sup> -94 <sup>b</sup>	.....	38
112	<i>Helianthus annuus</i> <sup>6</sup>	Common sunflower	Intact	25	Chem	.....	9 <sup>1a</sup> -3 <sup>1a</sup>	.....	75
113	<i>Hordeum vulgare</i> <sup>6</sup>	Barley	Intact	25	Chem	.....	76-15	1.2-0.8	87
114	<i>Ilex aquifolium</i>	English holly	Intact	21	Mano	.....	12	.....	76
115	<i>Ipomoea grandiflora</i>	Large moonflower	Intact	20	Mano	220 <sup>b</sup>	.....	.....	29
116	<i>Iris germanica</i> <sup>5</sup>	German iris	Intact	22.5	Chem	.....	12-13.6-5	.....	5
117	<i>Lactuca sativa</i> <sup>7</sup>	Lettuce	Intact	24	Chem	.....	3.3-2.6	1.12-0.99	63
118			Intact	10	Chem	.....	1.3-0.73	1.09-0.93	63
119			Intact	0.5	Chem	.....	0.8-0.35	0.84-0.98	63
120	<i>Lycopersicon esculentum</i> <sup>1a</sup>	Tomato	Intact	27	Mano	260 <sup>b</sup> -320 <sup>b</sup>	.....	.....	23
121	<i>L. esculentum</i> <sup>8</sup>	Tomato	Segment	30	Mano	46 (42)	.....	1.13 (1.28)	79
122	<i>L. esculentum</i> <sup>8</sup>	Tomato	Segment	28	Mano	390 <sup>b</sup> -430 <sup>b</sup>	0.96-0.91	.....	45
123	<i>Malus pumila</i> <sup>4</sup>	Common apple	Intact	33	Chem	.....	8.6 <sup>1a</sup> -43.0 <sup>1a</sup>	.....	70
124	<i>Nicotiana glauca</i> x <i>N. langsdorffii</i> <sup>3</sup>	Tobacco	Segment	25	Mano	330 <sup>b</sup> -170 <sup>b</sup>	.....	1.27-1.43	56
125	<i>Oenothera biennis</i> <sup>3</sup>	Common evening primrose	Blade	18	Mano	.....	24-12	0.83-0.70	58
126	<i>Phaseolus vulgaris</i>	Kidney bean	Intact	26	Mano	26-57	.....	.....	40
127	<i>Phleum pratense</i>	Timothy	Intact	21-26	.....	124 <sup>e</sup>	.....	.....	46
128	<i>Phoenix dactylifera</i>	Date palm	Intact	20	Chem	.....	4.5 <sup>1a</sup>	.....	28
129	<i>Pinus pinea</i>	Italian stone pine	Intact	24; 14	Mano	.....	12; 6.9	0.83; 0.82	13
130	<i>Pisum sativum</i> <sup>1a</sup>	Garden pea	Intact	27	Mano	430 <sup>b</sup> -680 <sup>b</sup>	.....	.....	23

<sup>1/</sup> Unless otherwise indicated. <sup>2a/</sup> Effect of growth, development, or maturation. <sup>3a/</sup> Effect of moisture. <sup>4a/</sup> Effect of storage or starvation. <sup>5a/</sup>  $\mu\text{l}$  per 100 mg dry weight per hour. <sup>6a/</sup> Effect of inorganic nutrition, salts. <sup>7a/</sup> Effect of light or photoperiod. <sup>8a/</sup>  $\mu\text{l}$  per sq cm per hour.

continued



## 76. RESPIRATION RATES

### Part IV. TRACHEOPHYTA

	Species	Common Name	Condition or Part	Temp. °C	Method	Respiration Rate μl/100 mg wet wt/hr <sup>1</sup>		Respiratory Quotient CO <sub>2</sub> /O <sub>2</sub>	Reference
						Q <sub>O<sub>2</sub></sub>	Q <sub>CO<sub>2</sub></sub>		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Leaves									
131	<i>Polypodium vulgare</i>	Common polypody	Frond	20	Mano	.....	10	0.92	52
132			Frond with sori	20	Mano	.....	19	1.06	52
133	<i>Populus deltoides</i> x <i>P. nigra</i>	Poplar	Intact	.....	Chem	.....	19	.....	66
134	<i>Prunus amygdalus</i>	Almond	Intact	14	Mano	.....	29	1.00	59
135	<i>P. laurocerasus</i> <sup>5</sup>	Laurel cherry	Intact	22.5	Chem	.....	20-3.4-13.6	.....	6
136	<i>Quercus coccifera</i> <sup>3</sup>	Kermes oak	Intact	21	Mano	.....	44-13	0.87-0.79	58
137	<i>Raphanus raphanistrum</i>	Wild radish	Blade	Room	Mano	.....	13.3	0.73	57
138			Petiole	Room	Mano	.....	6.2	0.86	57
139	<i>Rheum rhaponticum</i>	Garden rhubarb	Segment	30	Mano	.....	29	1.17	55
140	<i>Rhododendron fargesii</i> <sup>5</sup>	Père Farges' rhododendron	Intact	22.5	Chem	.....	13.6-5.1	.....	5
141	<i>Rosa</i> sp.	Rose	Intact	14	Mano	.....	23	0.93	59
142	<i>Salix glauca</i>	Gray-leaf willow	Intact	20;10;0	Chem	.....	78;45;13	.....	80
143	<i>Solanum tuberosum</i>	Potato	Intact	48;30	Chem	.....	137;41	.....	39
144			Intact	10	Chem	10	.....	.....	39
145	<i>Taxus baccata</i>	English yew	Intact	46;34;16	Mano	.....	55;23;6	0.89;0.80;0.86	13
146	<i>Tradescantia virginidis</i>	Wandering Jew	Intact	29	Mano	.....	.....	1.01	67
147	<i>Triticum aestivum</i>	Wheat	Intact	25	Mano	.....	40.2	0.97	57
148			Intact, etiolated	25	Mano	.....	37.5	0.98	57
149	<i>Ulmus glabra</i>	Scotch elm	Intact	16	Chem	.....	24	.....	66
150	<i>Vicia faba</i>	Broad bean	Blade	Room	Mano	.....	11.1	.....	57
151			Petiole	Room	Mano	.....	4.1	.....	57
152	<i>Vitis vinifera</i>	European grape	Blade	.....	Chem	.....	81 <sup>6</sup>	.....	54
153	<i>Yucca gloriosa</i> <sup>5</sup>	Mound lily yucca	Intact	22.5	Chem	.....	8.5-3.3	.....	5
154	<i>Zea mays</i>	Corn	Intact	26	Mano	.....	68.3	0.99	57
155			Intact, etiolated	26	Mano	.....	54.1	0.97	57
Flowers									
156	<i>Antirrhinum majus</i> <sup>3</sup>	Snapdragon	Petal	23	Mano	.....	82-70-34	1.15-1.13-1.00	52
157			Stamen	24	Mano	.....	81-106-76	.....	52
158	<i>Cucumis sativus</i> <sup>3</sup>	Cucumber	Pistil	22	Mano	.....	48-43-29	.....	52
159	<i>Gladiolus gandavensis</i>	Breeder's gladiolus	Petal	24	Mano	.....	15	0.72	52
160			Stamen	24	Mano	.....	27	0.77	52
161			Pistil	24	Mano	.....	71	0.90	52
162	<i>Helianthus annuus</i> <sup>3</sup>	Common sunflower	Inflorescence	10	Chem	.....	57 <sup>6</sup> -43 <sup>6</sup>	.....	44
163	<i>Lilium bulbiferum</i> <sup>3</sup>	Bulbil lily	Stamen	.....	.....	.....	56-21	1.14-0.98	33
164			Pistil	.....	.....	.....	58-19	1.06-1.12	33
165	<i>Pinus densiflora</i>	Japanese red pine	Pollen	25	Mano	160 <sup>6</sup>	.....	.....	60, 61
166	<i>Rosa</i> sp.	Rose	Intact	28	Mano	.....	.....	1.04	52
167	<i>Yucca gloriosa</i> <sup>3</sup>	Mound lily yucca	Pistil	16	Mano	.....	24-23-22	.....	52
168			Petal	24	Mano	.....	67-41-44	0.91-0.97-1.07	52
Fruits									
169	<i>Capsicum frutescens</i> <sup>5</sup>	Bush red pepper	Intact	24	Chem	.....	4.0-1.4	1.12-0.88	64
170			Intact	10	Chem	.....	1.2-0.58	1.27-0.88	64
171			Intact	0.5	Chem	.....	0.44-0.29	0.96	64
172	<i>Citrus limon</i>	Lemon	Intact	38;21;0	.....	.....	4.1;1.1;0.15	1.4;1.0;1.2	34

<sup>1</sup>/ Unless otherwise indicated. <sup>2</sup>/ Effect of growth, development, or maturation. <sup>3</sup>/ Effect of storage or starvation. <sup>6</sup>/ μl per 100 mg dry weight per hour.

continued

## 76. RESPIRATION RATES

### Part IV. TRACHEOPHYTA

	Species	Common Name	Condition or Part	Temp. °C	Method	Respiration Rate $\mu\text{l}/100 \text{ mg wet wt}/\text{hr}^{\text{a}}$		Respiratory Quotient $\text{CO}_2/\text{O}_2$	Reference
						$\text{QO}_2$	$\text{QCO}_2$		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Fruits									
173	<i>Citrus sinensis</i>	Sweet orange	Intact	21;10;0	.....	.....	2.0;0.8;0.15	1.1;1.1;1.2	34
174	<i>Cucumis sativus</i> <sup>5</sup>	Cucumber	Intact	24	Chem	.....	2.3-0.8	1.01-0.91	64
175			Intact	10	Chem	.....	1.0-0.4	1.01-1.10	64
176			Intact	0.5	Chem	.....	0.2-0.08	0.97-0.88	64
177	<i>Fragaria sp.</i> <sup>3</sup>	Strawberry	Intact	20	Chem	.....	3.3-5.1	0.84-0.91	62
178	<i>Helianthus annuus</i>	Common sunflower	Intact	25	Mano	.....	.....	0.96	67
179	<i>Lycopersicon esculentum</i> <sup>5</sup>	Tomato	Intact	24	Chem	.....	2.5-1.6	1.11-1.13	64
180			Intact	10	Chem	.....	0.77-0.58	1.39-1.06	64
181			Intact	0.5	Chem	.....	0.36-0.15	1.11-0.9	64
182	<i>Malus pumila</i> <sup>3</sup>	Common apple	Intact	27	Mano	.....	2.4-5.1-0.6	0.43-0.91	71
183			Intact	20	Chem	.....	1.7-0.8	.....	27
184	<i>Nicotiana tabacum</i>	Common tobacco	Intact	28	Mano	.....	.....	0.94	67
185	<i>Persea americana</i> <sup>5</sup>	American avocado	Intact	15	Chem	.....	5.8-3.6-8.1	.....	11
186	<i>Phaseolus vulgaris</i> <sup>5</sup>	Kidney bean	Intact	24	Chem	.....	16.4-6.6	1.14-1.00	64
187			Intact	10	Chem	.....	4.6-2.0	1.08-0.98	64
188			Intact	0.5	Chem	.....	0.95-0.65	0.94-0.96	64
189	<i>Pisum sativum</i> <sup>5</sup>	Garden pea	Intact	24	Chem	.....	20-12	1.32-1.06	64
190			Intact	10	Chem	.....	7.9-3.1	1.13-1.00	64
191			Intact	0.5	Chem	.....	2.2-1.4	1.07-0.96	64
192	<i>Prunus domestica</i> <sup>5</sup>	Garden plum	Intact	18	Chem	.....	1.7-3.6	.....	19
193	<i>P. domestica</i>	Garden plum	Intact	4	Chem	.....	0.5	.....	19
194	<i>P. persica</i> <sup>5</sup>	Peach	Intact	18	Chem	.....	1.4-2.0	.....	19
195			Intact	4	Chem	.....	0.4-0.3	.....	19
196	<i>Pyrus communis</i> <sup>3</sup>	Pear	Intact	18	Chem	.....	6.3-1.0-1.2	.....	26
197	<i>Quercus alba</i>	White oak	Intact	30;10;2.5	Mano	21 <sup>b</sup> ;16 <sup>b</sup> ;17 <sup>b</sup>	.....	0.71;0.30;0.16	15
198	<i>Ribes rubrum</i>	Northern red currant	Intact	28	Mano	.....	.....	1.4	67
199	<i>Rosa sp.</i>	Rose	Intact	28	Mano	.....	.....	0.86	67
200	<i>Solanum lycopersicum</i>	Nightshade	Intact	28	Mano	.....	.....	1.9	67
201	<i>Triticum aestivum</i> <sup>3</sup>	Wheat	Intact	28	Mano	340 <sup>b</sup> -8 <sup>b</sup>	.....	.....	73
202	<i>Vitis vinifera</i>	European grape	Intact	28	Mano	.....	.....	1.6	67
203	<i>Zea mays</i>	Corn	Intact	28;4.5	Chem	.....	17-11;3.5	.....	2
Whole Plants									
204	<i>Betula nana</i>	Dwarf arctic birch	Intact	16	Mano	.....	7.0 <sup>14</sup>	0.93	47
205	<i>Gossypium herbaceum</i> <sup>3</sup>	Levant cotton	Intact	38	Chem	.....	198 <sup>b</sup> -65 <sup>b</sup>	.....	38
206	<i>Helianthus annuus</i>	Common sunflower	Intact	10	Chem	.....	148 <sup>b</sup> -13 <sup>b</sup>	.....	44
207	<i>Ipomoea batatas</i> <sup>15</sup>	Sweet potato	Intact	21	Mano	.....	1-2	.....	85
208	<i>Triticum aestivum</i> <sup>5</sup>	Wheat	Intact	2	Chem	.....	38 <sup>b</sup> 13 <sup>b</sup>	.....	22

<sup>14</sup>/ Unless otherwise indicated. <sup>3</sup>/ Effect of growth, development, or maturation. <sup>5</sup>/ Effect of storage or starvation. <sup>6</sup>/  $\mu\text{l}$  per 100 mg dry weight per hour. <sup>14</sup>/  $\mu\text{l}$  per sq cm per hour. <sup>15</sup>/ Effect of wounding.

**Contributors:** (a) Mandels, Gabriel R., and Darby, Richard T., (b) Forward, Dorothy F., (c) Klein, Richard M., (d) Henderson, Lavaniel L., Sr.

**References:** [1] Allison, F. E., et al. 1940. Bot. Gaz, 101:513. [2] Appleman, C. O., and R. G. Brown. 1946. Am. J. Botany 33:170. [3] Arney, S. E. 1947. New Phytologist 46:68. [4] Aubert, E. 1892. Rev. Gen. Botan.

continued

**76. RESPIRATION RATES**

**Part IV. TRACHEOPHYTA**

- 4:421. [5] Audus, L. J. 1939. *New Phytologist* 38:284. [6] Audus, L. J. 1947. *Ann. Botany (London)*, N.S. 11:165. [7] Bailey, C. H. 1946. *Plant Physiol.* 15:257. [8] Bailey, C. H., and A. M. Gurjar. 1920. *J. Biol. Chem.* 44:17. [9] Benoy, M. P. 1929. *J. Agr. Res.* 39:75. [10] Berry, L. J. 1949. *J. Cellular Comp. Physiol.* 33:41. [11] Biale, J. B. 1946. *Am. J. Botany* 33:363. [12] Bonner, J. 1949. *Ibid.* 36:429. [13] Bonnier, G., and L. Mangin. 1884. *Ann. Sci. Nat. Zool., Ser. 6*, 19:217. [14] Boysen-Jensen, P., and D. Müller. 1929. *Jahrb. Wiss. Botan.* 70:503. [15] Brown, J. W. 1939. *Plant Physiol.* 14:621. [16] Brown, J. W. 1946. *Botan. Gaz.* 107:332. [17] Choudhury, J. K. 1939. *Proc. Roy. Soc. (London)*, B, 127:238. [18] Christiansen, G. S., and K. V. Thimann. 1950. *Arch. Biochem. Biophys.* 26:248. [19] Claypool, L. L., and F. W. Allen. 1948. *Proc. Am. Soc. Hort. Sci.* 51:103. [20] DeLong, W. A., J. H. Beaumont, and J. J. Willaman. 1930. *Plant Physiol.* 15:509. [21] Denny, F. E. 1939. *Contrib. Boyce Thompson Inst.* 10:453. [22] Dexter, S. T. 1934. *Plant Physiol.* 9:831. [23] Elliott, B. B., and A. C. Leopold. 1952. *Ibid.* 27:787. [24] Ermakov, A. I., and N. N. Ivanov. 1931. *Biochem. Z.* 231:79. [25] Erygin, P. S. 1936. *Plant Physiol.* 11:821. [26] Ezell, B. D., and F. Gerhardt. 1938. *J. Agr. Res.* 56:365. [27] Ezell, B. D., and F. Gerhardt. 1942. *Ibid.* 65:453. [28] Gabrielsen, E. K. 1931. *Planta* 14:217. [29] Genevois, L. 1927. *Biochem. Z.* 191:147. [30] Godlewski, E. 1882. *Jahrb. Wiss. Botan.* 13:491. [31] Goodwin, R. H., and D. R. Goddard. 1940. *Am. J. Botany* 27:234. [32] Groner, M. G. 1936. *Ibid.* 23:381. [33] Guilcher, J. M. 1937. *Rev. Gen. Botan.* 49:235. [34] Haller, M. H., et al. 1945. *J. Agr. Res.* 71:327. [35] Harrington, G. T. 1923. *Ibid.* 23:117. [36] Hasselbring, H., and L. A. Hawkins. 1913. *Ibid.* 5:509. [37] Henderson, J. H., and J. F. Stauffer. 1944. *Am. J. Botany* 31:528. [38] Inamdar, R. S., S. B. Singh, and T. D. Pande. 1925. *Ann. Botany (London)* 39:281. [39] Johansson, N. 1926. *Svensk Botan. Tidskr.* 20:107. [40] Johnson, C. M., and W. M. Hoskins. 1952. *Plant Physiol.* 27:507. [41] Johnstone, G. R. 1925. *Botan. Gaz.* 80:145. [42] Jones, H. A. 1920. *Ibid.* 69:127. [43] Karon, M. L., and A. M. Altschul. 1946. *Plant Physiol.* 21:506. [44] Kidd, F., C. West, and G. E. Briggs. 1921. *Proc. Roy. Soc. (London)*, B, 92:368. [45] Klein, R. M. 1951. *Arch. Biochem. Biophys.* 30:207. [46] Kostytschev, S. 1927. *Plant respiration*. Blakiston, Philadelphia. [47] Krascheninnikoff, T. 1926. *Compt. Rend.* 182:939. [48] Leach, W. 1936. *Proc. Roy. Soc. (London)*, B, 119:507. [49] Leach, W., and K. W. Dent. 1934. *Ibid.*, B, 116:150. [50] Lewin, M. 1905. *Ber. Deut. Botan. Ges.* 23:100. [51] Lundegårdh, H. 1950. *Nature* 165:513. [52] Maige, G. 1911. *Ann. Sci. Nat. Botan. Biol. Vegetale, Ser. 9*, 14:1. [53] Meyer, A., and N. T. Deleano. 1911. *Z. Botan.* 3:657. [54] Meyer, A., and N. T. Deleano. 1913. *Ibid.* 5:209. [55] Morrison, J. F. 1949. *Australian J. Exptl. Biol. Med. Sci.* 27:581. [56] Newcomb, E. H. 1950. *Am. J. Botany* 37:264. [57] Nicolas, G. 1909. *Ann. Sci. Nat. Botan. Biol. Vegetale, Ser. 9*, 10:1. [58] Nicolas, G. 1918. *Rev. Gen. Botan.* 30:209. [59] Nicolas, G. 1919. *Ibid.* 31:161. [60] Okunuki, K. 1937. *Acta Phytochim. (Japan)* 9:267. [61] Okunuki, K. 1939. *Ibid.* 11:27. [62] Overholser, E. L., M. B. Hardy, and H. D. Locklin. 1931. *Plant Physiol.* 6:549. [63] Pack, D. A. 1920. *Botan. Gaz.* 71:32. [64] Platenius, H. 1942. *Plant Physiol.* 17:179. [65] Platenius, H. 1943. *Ibid.* 18:671. [66] Plester, W. 1912. *Beitr. Biol. Pflanz.* 11:249. [67] Pringsheim, E. G. 1935. *Jahrb. Wiss. Botan.* 81:579. [68] Ronkin, R. R., and S. C. Brooks. 1942. *Science* 95:231. [69] Ruhland, W., and K. Ramshorn. 1938. *Planta* 28:471. [70] Schneider, G. W., and N. F. Childers. 1941. *Plant Physiol.* 16:565. [71] Shaw, S. T. 1942. *Ibid.* 17:80. [72] Sherman, H. 1921. *Botan. Gaz.* 72:1. [73] Shirk, H. G. 1942. *Am. J. Botany* 29:105. [74] Smith, F. G. 1948. *Plant Physiol.* 23:70. [75] Spoehr, H. A., and J. M. McGee. 1924. *Am. J. Botany* 11:493. [76] Stich, C. 1891. *Flora (Jena)* 74:1. [77] Stiles, W., and K. W. Dent. 1947. *Ann. Botany (London)*, N.S. 11:1. [78] Stiles, W., and W. Leach. 1933. *Proc. Roy. Soc. (London)*, B, 113:405. [79] Tsui, C. 1949. *Nature* 164:970. [80] Wager, H. G. 1941. *New Phytologist* 40:1. [81] Walter, E. M., and J. M. Nelson. 1945. *Arch. Biochem. Biophys.* 6:131. [82] Wedding, R. T., L. A. Riehl, and W. A. Rhoads. 1952. *Plant Physiol.* 27:269. [83] White, D. G., and N. F. Childers. 1944. *Ibid.* 19:699. [84] White, J. 1909. *Proc. Roy. Soc. (London)*, B, 81:417. [85] Whiteman, T. M., and H. A. Schomer. 1945. *Plant Physiol.* 20:171. [86] Woodford, E. K., and F. G. Gregory. 1948. *Ann. Botany (London)*, N.S. 12:335. [87] Yemm, E. W. 1935. *Proc. Roy. Soc. (London)*, B, 117:504.



## 77. HEART RATES

Heart rate varies with species, sex, age, size, environment, and temperature. Values in parentheses are ranges, estimate "c" (cf. Introduction).

### Part I. MAN

	Specification	Heart Rate beats/min	Refer- ence		Specification	Heart Rate beats/min	Refer- ence
	(A)	(B)	(C)		(A)	(B)	(C)
	Embryo		1	19	45-50 yr	72(49-100)	5
1	5th mo	156(150-160)		20	50-55 yr	72(52-94)	5
2	6th mo	154(141-155)		21	55-60 yr	75(48-108)	5
3	7th mo	149(118-156)		22	60-65 yr	73(54-100)	5
4	8th mo	142(129-152)		23	65-70 yr	75(52-96)	5
5	9th mo	146(131-173)		24	70-75 yr	75(54-104)	5
6	Premature	145(110-185)	6	25	75-80 yr	72(50-94)	5
7	Newborn	134(101-160)	5	26	>80 yr	77(63-98)	3
8	1 yr	111(84-136)	5		College students, ♂		4
9	2 yr	108(84-134)	5	27	Basal	65(45-105)	
10	4 yr	103(80-133)	5	28	Recumbent	66(40-100)	
11	5-9 yr	96(68-128)	3	29	Sitting	73(48-105)	
12	10-14 yr	87(56-120)	3	30	Standing	82(54-124)	
13	15-19 yr	79(52-112)	3		28 yr old, ♂		2
14	20-24 yr	74(41-100)	3	31	Sleeping	59.4(52.8-67.1)	
15	25-30 yr	72(52-102)	5	32	Awake	77.8(61.2-111.8)	
16	30-35 yr	70(58-104)	5		25 yr old, ♀		2
17	35-40 yr	72(56-100)	5	33	Sleeping	65.3(57.7-75.4)	
18	40-45 yr	72(50-104)	5	34	Awake	83.9(67.1-120.6)	

Contributors: (a) Johnson, Richard P., (b) Robb, Jane Sands

References: [1] Barcroft, J. 1936. *Physiol. Rev.* 16:103. [2] Boas, E. P., and E. F. Goldschmidt. 1932. The heart rate. C. C. Thomas, Springfield, Ill. p. 23. [3] Bowerman, W. G., and J. H. Brett. 1941. *Quart. Rev. Biol.* 16:90. [4] Brouha, L., and C. W. Heath. 1943. *New Engl. J. Med.* 228:473. [5] Lehmann, G. 1925. *Tabulae Biologicae* 1:140. [6] Sutliff, W. D., and E. Holt. 1925. *Arch. Internal Med.* 35:224.

### Part II. VERTEBRATES OTHER THAN MAN

Values are for adult animals, unless otherwise specified.

	Species	Common Name	Specification	Heart Rate beats/min	Refer- ence
	(A)	(B)	(C)	(D)	(E)
Mammalia					
1	<i>Bos taurus</i>	Cattle	500 kg; 38°C	46-53	2
2		Young	.....	106(100-115)	19
3		Newborn	.....	(141-160)	8
4		Embryo	.....	161	8
5	<i>Camelus bactrianus</i>	Bactrian camel	.....	(25-32)	19
6	<i>Canis familiaris</i>	Dog	5-20 kg	(72-200)	8,27,36
7		Young	1,040 g	208(145-275) <sup>1</sup>	18
8		Newborn	.....	(160-180)	8
9		Embryo	.....	(120-170)	8
10	<i>Capra hircus</i>	Goat	33 kg; 39°C	81(70-135)	2,8
11		Newborn	.....	(145-240)	1
12		Embryo	.....	(120-246)	1
13	<i>Cavia porcellus</i>	Guinea pig	300-750 g	(230-300)	8,20
14			437 g	269(225-312) <sup>1</sup>	18
15	<i>Dasyus novemcinctus</i>	Nine-banded armadillo	2.8-4.0 kg; 32-36°C	(70-100)	29
16	<i>Delphinapterus leucas</i>	Beluga whale	.....	(15-16)	34
17	<i>Didelphis marsupialis virginiana</i>	Virginia opossum	2.2-3.2 kg; 35°C	187(140-228) <sup>1</sup>	10

<sup>1</sup>/ Anesthetized.

continued



## 77. HEART RATES

### Part II. VERTEBRATES OTHER THAN MAN

	Species	Common Name	Specification	Heart Rate beats/min	Refer- ence	
	(A)	(B)	(C)	(D)	(E)	
Mammalia						
18	<i>Elephas maximus</i>	Asiatic elephant	2,000-3,000 kg; 36°C	(25-50)	2,8,36	
19	<i>Equus caballus</i>	Horse	380-450 kg	(34-55)	8,19	
20		Young	.....	63(60-71)	19	
21		Newborn	.....	(100-120)	26	
22	<i>Erinaceus europaeus</i>	European hedgehog	500-900 g; 36°C	246(234-264)	10	
23			485 g	263(200-325) <sup>1</sup>	18	
24	<i>Eutamias minimus</i>	Least chipmunk	40 g; 38.7°C	684(660-702)	10	
25	<i>Felis catus</i>	Cat	2.5 kg	(110-240) <sup>1</sup>	8,27	
26		Young	117 g	300	8	
27		Newborn	.....	300	8,21	
28	<i>Macaca irus</i>	Crab-eating macaque	.....	215	11	
29	<i>Mesocricetus auratus</i>	Golden hamster	103 g	347(276-420) <sup>1</sup>	18	
30			75-103 g	(375-425)	20	
31	<i>Mus musculus</i>	House mouse	10-20 g; 38.4°C	624(480-738) <sup>1</sup>	10	
32			17.4 g	500(450-550) <sup>1</sup>	17	
33			Young	12 g	670	4,8
34	<i>Mustela vison</i>	Mink	0.7-1.4 kg; 40.5°C	(272-414)	10	
35	<i>Myotis lucifugus</i>	Little brown bat	6 g	588 <sup>1</sup>	10	
36	<i>Ondatra zibethica</i>	Muskrat	0.8-1.3 kg; 38°C	(148-306) <sup>1</sup>	10	
37	<i>Oryctolagus cuniculus</i>	European rabbit	1,344 g	251(167-330) <sup>1</sup>	18	
38			Newborn	.....	220	13
39	<i>Ovis aries</i>	Sheep	50 kg	(70-80)	30	
40	<i>Phoca vitulina</i>	Harbor seal	20-25 kg	100	17	
41	<i>Phocaena phocaena</i>	Harbor porpoise	170 kg	(40-110)	16	
42	<i>Rattus norvegicus</i>	Norway rat	252 g	352(260-450) <sup>1</sup>	18	
43			92-210 g	305(270-350) <sup>2</sup>	15	
44			Newborn	.....	161(121-201)	21
45			Embryo	.....	(95-256)	1
46	<i>Sciurus carolinensis</i>	Gray squirrel	500-600 g; 40.1°C	390	10	
47	<i>Sorex cinereus</i>	Gray shrew	3-4 g; 38.8°C	782(588-1,320)	22	
48	<i>Sus scrofa</i>	Swine	100 kg	(60-80)	8,12,19	
49			Newborn	.....	227	21
Aves						
50	<i>Anas platyrhynchos</i>	Mallard duck	2,304 g	212(133-268)	8,31	
51	<i>Anser</i> sp.	Goose	4,000 g	80	8,23	
52			2,800 g	144	8,33	
53	<i>Archilochus colubris</i>	Ruby-throated hummingbird	4 g	615 <sup>2</sup>	24	
54	<i>Columba</i> sp.	Pigeon	240-370 g	185(141-225)	5,7,19	
55	<i>Corvus cornix</i>	Hooded crow	360 g	378(312-492)	19,31	
56	<i>Cygnus olor</i>	Mute swan	.....	257	9	
57	<i>Gallus domesticus</i>	Chicken	1,980 g	312(178-458)	19,31	
58	<i>Gyps fulvus</i>	Griffon vulture	8,310 g	199	19,31	
59	<i>Larus canus</i>	Mew gull	388 g	401(360-483)	19,31	
60	<i>Meleagris gallopavo</i>	Turkey	8,750 g	93	19,31	
61	<i>Passer domesticus</i>	House sparrow	28 g	350 <sup>2</sup> ; 902 <sup>3</sup>	24	
62			20 g	(640-910)	8,25	
63	<i>Serinus canarius</i>	Canary	16 g	514 <sup>2</sup> ; 1,000 <sup>3</sup>	24	
64	<i>Struthio camelus</i>	African ostrich	80 kg	65(60-70)	8	
65	<i>Sturnus vulgaris</i>	Starling	.....	388(375-400)	37	
66	<i>Troglodytes aedon</i>	House wren	11 g	450 <sup>2</sup> ; 950 <sup>3</sup>	24	
67	<i>Turdus migratorius</i>	American robin	.....	570(520-620)	37	
68	<i>Zenaidura macroura</i>	Mourning dove	130 g	135 <sup>2</sup> ; 570 <sup>3</sup>	24	
Reptilia and Amphibia						
69	<i>Alligator mississippiensis</i>	American alligator	.....	38	35	
70	<i>Anguis fragilis</i>	Slowworm	.....	64	19	
71	<i>Bufo</i> sp.	Toad	.....	(40-50)	19	
72	<i>Caretta</i> sp.	Loggerhead turtle	.....	11	19	
73	<i>Emys orbicularis</i>	European pond turtle	.....	(9-60)	19	

1/ Anesthetized. 2/ Basal rate. 3/ Maximum rate on nest.

continued

## 77. HEART RATES

### Part II. VERTEBRATES OTHER THAN MAN

Species	Common Name	Specification	Heart Rate beats/min	Refer- ence	
(A)	(B)	(C)	(D)	(E)	
Reptilia and Amphibia					
74	<i>Natrix natrix</i>	European water snake	169 g	(23-41)	6, 19
75	<i>Pseudemys terrapen rugosa</i>	Cuban freshwater turtle	.....	(21-44)	19
76	<i>Rana pipiens</i>	Leopard frog	.....	(37.5-60.0)	19
77	<i>Salamandra</i> sp.	Salamander	.....	(30-40)	28
Pisces and Chondrichthyes					
78	<i>Anguilla</i> sp.	Freshwater eel	.....	(39-68)	19
79	<i>Carassius auratus</i>	Goldfish	.....	(36-40)	19
80	<i>Cyprinus carpio</i>	Common carp	.....	75(72-78)	37
81	<i>Esox lucius</i>	Northern pike	.....	(38-54)	3
82	<i>Gadus morhua</i>	Atlantic cod	.....	(48-60)	3
83	<i>Ictalurus</i> sp.	Bullhead	.....	22(5-50)	14
84	<i>Melanogrammus</i> sp.	Haddock	.....	(30-40)	32
85	<i>Micropterus salmoides</i>	Largemouth black bass	.....	20(5-50)	14
86	<i>Perca fluviatilis</i>	European perch	.....	(52-66)	3
87	<i>Pleuronectes platessa</i>	European plaice	.....	(54-76)	3
88	<i>Raja</i> sp.	Skate	.....	(16-50)	19
89	<i>Salmo trutta</i>	Brown trout	.....	(30-46)	3
90	<i>Squalus acanthias</i>	Atlantic spiny dogfish	.....	(40-50)	3

*Contributors:* (a) Kruta, Vladislav, and Seliger, Václav, (b) Dawe, Albert R., (c) Robb, Jane Sands, (d) Morrison, Peter R., (e) Garb, Solomon, (f) Nardone, Roland M., (g) Lombard, Elna A., (h) Woodbury, Robert A., (i) Johnson, Richard P., (j) Walker, Sheppard M.

*References:* [1] Barcroft, J. 1936. *Physiol. Rev.* 16:103. [2] Benedict, F. G., and E. G. Retzman. 1923. *Carnegie Inst. Wash. Publ.* 324. [3] Bielig, W. 1931. *Z. Vergleich. Physiol.* 15:488. [4] Buchanan, F. 1908. *J. Physiol. (London)* 37:69. [5] Buchanan, F. 1909. *Ibid.* 38:62. [6] Buchanan, F. 1909. *Ibid.* 39:25. [7] Buchanan, F. 1910. *Sci. Progr. (London)* 5:60. [8] Clark, A. J. 1927. *Comparative physiology of the heart.* Macmillan, New York. [9] Davies, F., and E. T. B. Francis. 1950. *J. Anat.* 86:302. [10] Dawe, A. R. 1953. Ph.D. Thesis. Univ. Wisconsin, Madison. [11] DeWaart, A., and C. J. Storm. 1934. *Acta Brevia Neerl. Physiol. Pharmacol. Microbiol.* 4:130. [12] Dukes, H. H. 1955. *The physiology of domestic animals.* Ed. 7. Constock, Ithaca. [13] Hamilton, W. F., R. A. Woodbury, and E. B. Woods. 1937. *Am. J. Physiol.* 119:206. [14] Hart, I. J. 1944. *Proc. Florida Acad. Sci.* 7:221. [15] Hoskins, R. G., M. O. Lee, and E. P. Durrant. 1927. *Am. J. Physiol.* 82:621. [16] Irving, L., P. F. Scholander, and S. W. Grinnell. 1941. *J. Cellular Comp. Physiol.* 17:145. [17] Irving, L., P. F. Scholander, and S. W. Grinnell. 1941. *Ibid.* 18:283. [18] Kruta, V. 1958. *Babakova Sbirka (Praha)* 8. [19] Lehmann, G. 1925. *Tabulae Biologicae* 1:136-139. [20] Lombard, E. A. 1952. *Am. J. Physiol.* 171:189. [21] Marcuse, F. L., and A. U. Moore. 1943. *Ibid.* 139:49. [22] Morrison, P. R., A. R. Dawe, and F. A. Ryser. 1953. *Federation Proc.* 12:100. [23] Mosso, A. 1901. *Arch. Ital. Biol.* 35:21. [24] Odum, E. P. 1945. *Science* 101:153. [25] Oppenheimer, E. 1922. *Z. Ges. Exptl. Med.* 28:96. [26] Reichert, A. 1909. *Klinische Untersuchungen über die normale Pulsfrequenz unserer Haustiere.* O. Kindt, Giessen. [27] Reichert, A. 1910. *Zentr. Biochem. Biophys.* 10:170. [28] Rienmüller, J. 1932. *Arch. Ges. Physiol.* 230:782. [29] Scholander, P. F., L. Irving, and S. W. Grinnell. 1943. *J. Cellular Comp. Physiol.* 21:53. [30] Sisson, S. 1953. *The anatomy of the domestic animals.* Ed. 4. W. B. Saunders, Philadelphia. [31] Stübel, H. 1910. *Arch. Ges. Physiol.* 135:249. [32] Thesen, J. E. 1896. *Arch. Zool. Exptl. Gen.* 3(3):122. [33] Vierordt, K. 1877. *Grundriss der Physiologie des Menschen.* Ed. 5. H. Laupp, Tübingen. p. 162. [34] White, P. D., R. L. King, and J. L. Jenks. 1953. *New Engl. J. Med.* 248:69. [35] Wilber, C. G. 1960. *Comp. Biochem. Physiol.* 1:164. [36] Winterstein, H. 1910-24. *Handbuch der vergleichenden Physiologie.* G. Fischer, Jena. [37] Woodbury, R. A., and W. F. Hamilton. 1937. *Am. J. Physiol.* 119:663.

*continued*

# 77. HEART RATES

## Part III. INVERTEBRATES

Values are for adult animals, unless otherwise specified.

Class	Species	Common Name	Specification	Heart Rate beats/min	Refer- ence
(A)	(B)	(C)	(D)	(E)	(F)
Chordata					
1	Asciacea	<i>Ciona intestinalis</i>	Sea squirt	.....	(17-32) 16
2		<i>Molgula manhattensis</i>	Sea squirt	21-22°C	(30-50) 20
Arthropoda					
3	Merostomata	<i>Limulus polyphemus</i>	King crab	.....	20(18-28) 5
4	Crustacea	<i>Astacus fluviatilis</i>	Crayfish	.....	(30-60) 13,20,27
5		<i>Callinectes sapidus</i>	Blue crab	22-23°C	(25-84) 10
6		<i>Daphnia pulex</i>	Water flea	0.0008 g; 20°C	(381-418) 30
7				0.000025 g; 20°C	486
8		<i>Homarus gammarus</i>	European lobster	450 g; 20°C	50 30
9	Insecta	<i>Anopheles quadrimaculatus</i>	Malaria mosquito	Larva, stage 1; 25-27°C	131.7 15
10				Larva, stage 2; 25-27°C	134.3
11				Larva, stage 3; 25-27°C	118.6
12				Larva, stage 4; 25-27°C	106.6
13				Pupa; 25-27°C	109.1
14				Adult, ♀; 25-27°C	151.2
15		<i>Anthophora retusa</i>	Digger bee	.....	142 17
16		<i>Bombyx mori</i>	Silkworm	Resting	(40-50) 24
17				Active	(110-140) 24
18				Larva, calm	(44-66) 22
19				Larva, excited	94 22
20		<i>Calliphora</i> sp.	Blowfly	Larva; 18°C	60 <sup>1</sup> 21
21		<i>Drosophila funebris</i>	Fruit fly	28-29°C	235 26
22		<i>Dytiscus marginalis</i>	Diving beetle	Intact	(30-70) 11
23		<i>Ephestia kuehniella</i>	Mediterranean flour moth	Pupa	(6-11) 36
24		<i>Locusta migratoria</i>	Migratory locust	♂; 26°C	(25-100) <sup>1</sup> 34
25				♂; 29°C	(80-120) <sup>1</sup>
26				♀; 29°C	(80-130) <sup>1</sup>
27		<i>Melanoplus differentialis</i>	Differential grasshopper	Nymph to adult; 21-26°C	(40-70) 14
28		<i>Pediculus</i> sp.	Louse	.....	(30-48) 23
29		<i>Periplaneta americana</i>	American cockroach	18°C	72.9(22-150) <sup>1</sup> 2
30				Nymph to adult; 29°C	49(18-70) <sup>1</sup> 35
31				Nymph to adult; 26-27°C	(90-100) <sup>1</sup> 25
32				Nymph to adult; 20-26°C	117.3(99-147) 1
33				Nymph, ♂	94 8
34				Nymph, ♀	100 8
35		<i>Pieris brassicae</i>	European cabbage worm	Larva	29 32
36		<i>Prodenia eridania</i>	Southern armyworm	Larva; 29°C	58(40-90) <sup>1</sup> 35
37		<i>Tenebrio molitor</i>	Yellow mealworm	Pupa	10 28
38				Larva	(15-17) 28
39				Adult	(19-57) 3
Annelida					
40	Hirudinea	<i>Hirudo</i> sp.	Leech	.....	6 12
41	Oligochaeta	<i>Lumbricus terrestris</i>	Earthworm	.....	17(15-20) 7
42	Polychaeta	<i>Arenicola</i> sp.	Lugworm	.....	7(6-8) 6
44		<i>Nereis virens</i>	Clam worm	35°C	50 29
				7°C	4.6
Mollusca					
45	Cephalopoda	<i>Loligo</i> sp.	Squid	.....	(60-80) 5
46		<i>Sepia officinalis</i>	Cuttlefish	Perfused median ventricle, in situ	(18-30) 18,19
47	Bivalvia	<i>Mytilus edulis</i>	Mussel	0.104 g; 20°C	49.2 31
48		<i>Ostrea edulis</i>	Oyster	.....	(25-30) 33
49		<i>Pecten jacobaeus</i>	Scallop	.....	(22-50) 7,9
50	Gastropoda	<i>Aplysia limacina</i>	Sea hare	870 g; 20°C	28 31
51				347 g; 20°C	32.3

/1/ Heart exposed by dissection and examined under physiological saline.

continued



## 77. HEART RATES

### Part III. INVERTEBRATES

	Class	Species	Common Name	Specification	Heart Rate beats/min	Refer- ence
	(A)	(B)	(C)	(D)	(E)	(F)
	Mollusca					
52	Gastropoda	<i>Helix pomatia</i>	Land snail	36.7 g; 20°C	37.1	31
53				0.63 g; 20°C	47.6	
54				0.123 g; 20°C	60.3	
55		<i>Lymnaea stagnalis</i>	Freshwater snail	3.16 g; 20°C	21	31
56				0.00078 g; 20°C	55.4	
57	Polyplacophora	<i>Ischnochiton</i> sp.	Chiton	.....	(15-25)	4,5

**Contributors:** (a) Jones, Jack Colvard, (b) Kruta, Vladislav, (c) Robb, Jane Sands, (d) Nardone, Roland M., (e) Johnson, Richard P.

**References:** [1] Bellemare, E. R., and J. Belcourt. 1955. *Can. J. Zool.* 33:175. [2] Bettini, S., G. Natalizi, and M. Boccacci. 1956. *Riv. Parassitol.* 17:179. [3] Butz, A. 1957. *J. N. Y. Entomol. Soc.* 65:22. [4] Carlson, A. J. 1905. *Am. J. Physiol.* 13:396. [5] Carlson, A. J. 1906. *Ibid.* 16:47. [6] Carlson, A. J. 1908. *Ibid.* 22:353. [7] Clark, A. J. 1927. *Comparative physiology of the heart.* Macmillan, New York. [8] Coon, B. F. 1944. *J. Econ. Entomol.* 37:785. [9] Dogiel, J. 1877. *Arch. Mikroskop. Anat. Entwicklungsmech.* 14:59. [10] Dubuisson, M., and A. M. Monnier. 1931. *Arch. Intern. Physiol.* 34:180. [11] Duwez, Y., and P. Rijlant. 1936. *Compt. Rend. Soc. Biol.* 122:84. [12] Gaskell, W. H. 1914. *Phil. Trans. Roy. Soc. London, B*, 205:163. [13] Hoffman, P. 1911. *Arch. Anat. Physiol.*, p. 135. [14] Jahn, T. L., F. Crescitelli, and A. B. Taylor. 1937. *J. Cellular Comp. Physiol.* 10:439. [15] Jones, J. C. 1954. *J. Morphol.* 94:71. [16] Knoll, P. 1893. *Sitzber. Wien Akad. Wiss. Math. Naturw. Kl., I*, 102:387. [17] Kozhanchikov, I. V. 1932. *Bull. Leningrad Inst. Control Farm Forest Pests* 2:149. [18] Kruta, V. 1936. *Compt. Rend. Soc. Biol.* 122:582. [19] Kruta, V. 1937. *Casopis Lekarů Ceských* 76:1328. [20] Lehmann, G. 1925. *Tabulae Biologicae* 1:135. [21] Levy, R. 1928. *Compt. Rend. Soc. Biol.* 99:1482. [22] Masera, E. 1933. *Riv. Biol. (Perugia)* 15:225. [23] Müller, J. 1915. *Oesterr. Sanitaetsw.* 27:1. [24] Newport, G. 1837. *Phil. Trans. Roy. Soc. London, B*, 127:259. [25] Orser, W. B., and A. W. A. Brown. 1951. *Can. J. Zool.* 29:54. [26] Perttunen, V. 1955. *Ann. Entomol. Fennici* 21:78. [27] Prosser, C. L., ed. 1961. *Comparative animal physiology.* Ed. 2. W. B. Saunders, Philadelphia. p. 395. [28] Rengel, C. 1896. *Z. Wiss. Zool.* 62:1. [29] Rogers, C. G. 1911. *Am. J. Physiol.* 28:81. [30] Schwartzkopff, J. 1955. *Experientia* 11:323. [31] Schwartzkopff, J. 1956. *Verhandl. Deut. Zool. Ges.*, p. 463. [32] Tarasova, K. L. 1936. *Izv. Inst. Priklad. Zool. i Fitopatol. (Leningr.)* 6:15. [33] Von Skramlik, E. 1929. *Arch. Ges. Physiol.* 221:503. [34] Yamasaki, T., and T. Ishii. 1950. *Oyo Kontyu* 5:155. [35] Yeager, J. F., and J. B. Gahan. 1937. *J. Agr. Res.* 55:1. [36] Zeller, H. 1938. *Z. Morphol. Oekol. Tiere* 34:663.

## 78. ARTERIAL BLOOD PRESSURE

### Part I. MAN

Number of subjects: line 1, 24 infants; lines 7-18, 3,580 children; lines 19-42, 7,222 males and 7,984 females; lines 43-56, 2,998 males and 2,759 females. Values in parentheses are ranges, estimate "b" (cf. Introduction).

Age	Sex	Blood Pressure, mm Hg		Refer- ence	Age	Sex	Blood Pressure, mm Hg		Refer- ence		
		Systolic	Diastolic				Systolic	Diastolic			
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)		
1	Newborn	♂♀	80(64-96)	46(30-62)	5	4	2 yr	♂♀	99(74-124)	64(39-89) <sup>1</sup>	1
2	6 mo-1 yr	♂♀	89(60-118)	60(50-70) <sup>1</sup>	1	5	3 yr	♂♀	100(75-125)	67(44-90) <sup>1</sup>	
3	1 yr	♂♀	96(66-126)	66(41-91) <sup>1</sup>		6	4 yr	♂♀	99(79-119)	65(45-85) <sup>1</sup>	

<sup>1</sup>/ Point of muffling taken as the diastolic pressure.

continued



## 78. ARTERIAL BLOOD PRESSURE

### Part I. MAN

Age	Sex	Blood Pressure, mm Hg		Reference	Age	Sex	Blood Pressure, mm Hg		Reference		
		Systolic	Diastolic				Systolic	Diastolic			
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)		
7	5 yr	♂♀	94(80-108)	55(46-64)	2	32	35-39 yr	♀	124(97-151)	78(58-98)	3
8	6 yr	♂♀	100(85-115)	56(48-64)		33	40-44 yr	♂	129(100-159)	81(63-100)	
9	7 yr	♂♀	102(87-117)	56(48-64)		34		♀	127(94-161)	80(59-100)	
10	8 yr	♂♀	105(89-121)	57(48-66)		35	45-49 yr	♂	130(97-163)	82(61-103)	
11	9 yr	♂♀	107(91-123)	57(48-66)		36		♀	131(92-169)	82(59-104)	
12	10 yr	♂♀	109(93-125)	58(49-67)		37	50-54 yr	♂	135(97-172)	83(61-106)	
13	11 yr	♂♀	111(94-128)	59(49-69)		38		♀	137(96-179)	84(59-108)	
14	12 yr	♂♀	113(95-131)	59(49-69)		39	55-59 yr	♂	138(101-175)	84(62-106)	
15	13 yr	♂♀	115(96-134)	60(50-70)		40		♀	139(97-180)	84(61-106)	
16	14 yr	♂♀	118(99-137)	61(51-71)		41	60-64 yr	♂	142(100-183)	85(60-109)	
17	15 yr	♂♀	121(102-140)	61(51-71)		42		♀	144(100-188)	85(60-110)	
18	16 yr	♂♀	121(102-140)	61(51-71)		43	65-69 yr	♂	143(92-194)	83(64-102)	4
19	17 yr	♂♀	121(96-146)	74(56-93)	3	44		♀	154(97-211)	85(58-112)	
20		♀	116(93-139)	72(54-90)		45	70-74 yr	♂	145(93-197)	82(52-112)	
21	18 yr	♂	120(96-143)	74(55-94)		46		♀	159(108-210)	85(55-115)	
22		♀	116(94-139)	72(55-89)		47	75-79 yr	♂	146(104-188)	81(56-106)	
23	19 yr	♂	122(92-151)	75(54-95)		48		♀	158(106-210)	84(58-110)	
24		♀	115(92-138)	71(54-89)		49	80-84 yr	♂	145(95-195)	82(63-101)	
25	20-24 yr	♂	123(96-150)	76(57-96)		50		♀	157(102-212)	83(57-109)	
26		♀	116(93-139)	72(53-91)		51	85-89 yr	♂	145(98-192)	79(50-108)	
27	25-29 yr	♂	125(100-150)	78(60-95)		52		♀	154(99-209)	82(48-116)	
28		♀	117(94-139)	74(56-92)		53	90-94 yr	♂	145(99-191)	78(54-102)	
29	30-34 yr	♂	126(99-153)	79(60-98)		54		♀	150(104-196)	79(55-103)	
30		♀	120(92-147)	75(54-96)		55	95-106 yr	♂	146(92-200)	78(53-103)	
31	35-39 yr	♂	127(99-155)	80(60-101)		56		♀	149(103-195)	81(57-106)	

Contributors: (a) Master, Arthur M., (b) Van Liere, Edward J., and Lindsay, Hugh A., (c) Hartroft, W. Stanley

References: [1] Allen-Williams, G. M. 1945. Arch. Disease Childhood 20:125. [2] Graham, A. W., E. A. Hines, and R. P. Gage. 1945. Am. J. Diseases Children 69:203. [3] Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia. [4] Master, A. M., R. P. Lasser, and H. L. Jaffe. 1957. Proc. Soc. Exptl. Biol. Med. 94:463. [5] Woodbury, R. A., M. Robinow, and W. F. Hamilton. 1938. Am. J. Physiol. 122:472.

### Part II. ANIMALS OTHER THAN MAN

Values are for adult animals, unless otherwise specified. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Species	Common Name	Subjects	Anesthetic	Blood Pressure, mm Hg		Reference
					Systolic	Diastolic	
					Mean		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Mammalia							
1	<i>Bos taurus</i>	Cattle	.....	None	134(124-166)	88(80-120)	16
2		Young	4	.....	157(133-177)		15
3	<i>Canis familiaris</i>	Dog	13	None	112(95-136)	56(43-66)	27
4			22	Pentobarbital	149(108-189) 100(75-122)		21
5			67♂	Sodium barbital	134(85-190)		26
6			80♀	Sodium barbital	125(60-170)		26
7	<i>Capra hircus</i>	Goat	.....	None	120(112-126)	84(76-90)	16
8	<i>Cavia porcellus</i>	Guinea pig	8	Ether, pentobarbital, and/or procaine	77(28-140)	47(16-90)	17
9	<i>Didelphis sp.</i>	Opossum	.....	.....	(120-135)		15
10	<i>Equus caballus</i>	Horse	173♂	None	98(90-104)	64(45-86)	6

continued

## 78. ARTERIAL BLOOD PRESSURE

### Part II. ANIMALS OTHER THAN MAN

	Species	Common Name	Sub-jects	Anesthetic	Blood Pressure, mm Hg		Ref-erence
					Systolic	Diastolic	
					Mean		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
<b>Mammalia</b>							
11	<i>Equus caballus</i>	Horse	43♀	None	90(86-98)	59(43-84)	6
12		Young	5♂, 3♀	None	80	50	6
13	<i>Felis catus</i>	Cat	5	Barbital or ether	120	75	28
14			191♂	Dial-urethan	129(67-216)		22
15			208♀	Dial-urethan	121(62-200)		22
16		Newborn	.....	.....	(25-30)	.....	1
17	<i>Macaca mulatta</i>	Rhesus monkey	14	None	159(137-188)	127(112-152)	24
18	<i>Mesocricetus auratus</i>	Golden hamster	.....	Pentobarbital	(120-170)		5
19	<i>Mus musculus</i>	House mouse	9	Urethan or ether	113(95-125)	81(67-90)	29
20		Young	19	None	111(95-138)	.....	30
21	<i>Ornithorhynchus</i> sp.	Platypus	2♂, 1♀	.....	14	.....	9
22	<i>Oryctolagus cuniculus</i>	European rabbit	32	None	110(95-130)	80(60-90)	19
23		Young	.....	.....	21	.....	1
24		Newborn	1	.....	35	1	11
25	<i>Ovis aries</i>	Sheep	13	Local	114(90-140)		7
26		Newborn	.....	.....	73		1
27	<i>Phoca vitulina</i>	Harbor seal, young	1	.....	ca. 135		14
28	<i>Rattus norvegicus</i>	Norway rat	124	Pentobarbital	129(88-184)	91(58-145)	23
29			100	None	98(82-120)	.....	10
30		Young	18♂	Ether and/or amytal	124		8
31			23♀	Ether and/or amytal	116		8
32	<i>Sus scrofa</i>	Swine	.....	None	169(144-185)	108(98-120)	16
<b>Aves</b>							
33	<i>Anas platyrhynchos</i>	Mallard duck	.....	.....	162		15
34	<i>Anser</i> sp.	Goose	.....	.....	(129-176)		15
35	<i>Columba livia</i>	Street pigeon	4	None	135(120-140)	105(100-115)	29
36	<i>Corvus cornix</i>	Hooded crow	.....	.....	147		15
37	<i>Gallus domesticus</i>	Chicken	5	Barbital	130	85	28
38			13♂	.....	135	120	7
39			♀	.....	(88-171)		15
40	<i>Larus canus</i>	Mew gull	.....	.....	179		15
41	<i>Meleagris gallopavo</i>	Turkey	.....	.....	193		15
42	<i>Passer domesticus</i>	House sparrow	1	None	180	140	29
43		Fledgling	2	None	123(115-130)	.....	29
44		Pinfeatherer	3	None	108(80-135)	.....	29
45	<i>Serinus canarius</i>	Canary	4	None	220(200-250)	154(150-160)	29
46	<i>Sturnus vulgaris</i>	Starling	2	None	180(150-210)	130(100-160)	29
47	<i>Turdus migratorius</i>	American robin	2	None	118(110-125)	80	29
<b>Reptilia and Amphibia</b>							
48	<i>Bufo terrestris</i>	Southern toad	.....	.....	48		15
49	<i>Natrix natrix</i>	European water snake	.....	.....	89		15
50	<i>Pseudemys scripta elegans</i>	Red-eared turtle	5	.....	31 <sup>1</sup> ; 27 <sup>2</sup>	25 <sup>1</sup> ; 21 <sup>2</sup>	20
51	<i>Rana catesbeiana</i>	American bullfrog	6	None	43(36-56)	31(24-44)	29
52			6	None	32(28-36)	21(20-24)	2
53	<i>R. pipiens</i>	Leopard frog	6	None	31(21-33)	21(16-26)	2
<b>Pisces and Chondrichthyes</b>							
54	<i>Anguilla</i> sp.	Freshwater eel	.....	.....	(65-70)		15
55	<i>Cyprinus carpio</i>	Carp	3	None	43(40-45)	.....	29
56	<i>Esox lucius</i>	Northern pike	.....	.....	(35-84)		15
57	<i>Ictalurus punctatus</i>	Channel catfish	.....	.....	40	30	12
58			.....	.....	30	23	13
59	<i>Micropterus salmoides</i>	Largemouth black bass	.....	.....	50	40	12
60	<i>Raja punctulata</i>	Skate	.....	.....	16	7	18
61	<i>Salmo</i> sp.	Salmon	.....	.....	75(47-120)		15
62	<i>Squalus acanthias</i>	Atlantic spiny dogfish	.....	.....	32	28	3

<sup>1</sup>/ At 22°C. <sup>2</sup>/ At 16°C.

continued

**78. ARTERIAL BLOOD PRESSURE**

**Part II. ANIMALS OTHER THAN MAN**

Species	Common Name	Sub-jects	Anesthetic	Blood Pressure, mm Hg		Ref-er-ence
				Systolic	Diastolic	
				Mean		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Crustacea						
63	<i>Astacus marinus</i>	Crayfish	.....	.....	8,5	15
64	<i>Cancer irroratus</i>	Edible crab	6	.....	8	4
65	<i>Homarus americanus</i>	American lobster	14 <sup>3</sup> ; 1 <sup>4</sup>	.....	13 <sup>3</sup> ; 27 <sup>4</sup>	1 <sup>3</sup> ; 13 <sup>4</sup>
Gastropoda						
66	<i>Aplysia</i> sp.	Sea hare	.....	.....	(20-40)	25

/3/ At rest. /4/ During activity.

*Contributors:* (a) Van Liere, Edward J., and Lindsay, Hugh A., (b) Conklin, Ruth E., (c) Freed, S. Charles, (d) Freis, Edward D., (e) Heisler, Charles R., (f) Link, Roger P., (g) Rodbard, Simon, (h) Woodbury, Robert A.

*References:* [1] Barcroft, J., and D. H. Barron. 1945. *J. Exptl. Biol.* 22:63. [2] Bieter, R. N., and F. H. Scott. 1929. *Am. J. Physiol.* 91:265. [3] Burger, J. W., and S. E. Bradley. 1947. *Anat. Record* 99:670. [4] Burger, J. W., and C. M. Smythe. 1953. *J. Cellular Comp. Physiol.* 42:369. [5] Chatfield, P. O., and C. P. Lyman. 1950. *Am. J. Physiol.* 163:566. [6] Covington, N. G., and G. W. McNutt. 1931. *J. Am. Vet. Med. Assoc.* 79:603. [7] Dukes, H. H. 1955. *The physiology of domestic animals*. Ed. 7. Comstock, Ithaca. [8] Durant, R. R. 1927. *Am. J. Physiol.* 81:679. [9] Feakes, M. J., et al. 1950. *J. Exptl. Biol.* 27:50. [10] Friedman, M., and S. C. Freed. 1949. *Proc. Soc. Exptl. Biol. Med.* 70:670. [11] Hamilton, W. F., R. A. Woodbury, and E. B. Woods. 1937. *Am. J. Physiol.* 119:206. [12] Hart, I. J. 1944. *Proc. Florida Acad. Sci.* 7:221. [13] Hart, J. S. 1957. *Can. J. Zool.* 35:195. [14] Irving, L., P. F. Scholander, and S. W. Grinnell. 1942. *Am. J. Physiol.* 135:557. [15] Lehmann, G. 1925. *Tabulae Biologicae* 1:142, 143. [16] Link, R. P. Unpublished. Univ. Illinois, Urbana, 1956. [17] Marshall, L. H., and C. H. Hanna. 1956. *Proc. Soc. Exptl. Biol. Med.* 92:31. [18] Prosser, C. L., and F. A. Brown, Jr. 1961. *Comparative animal physiology*. Ed. 2. W. B. Saunders, Philadelphia. [19] Rodbard, S. 1940. *Am. J. Physiol.* 129:448. [20] Rodbard, S., and D. Feldman. 1946. *Proc. Soc. Exptl. Biol. Med.* 63:43. [21] Romagnoli, A. 1953. *Cornell Vet.* 43:161. [22] Root, M. A. 1950. *Am. J. Physiol.* 162:308. [23] Schroeder, H. A. 1942. *J. Exptl. Med.* 75:513. [24] Smith, C. C., and A. Ansevin. 1957. *Proc. Soc. Exptl. Biol. Med.* 96:428. [25] Straub, W. 1904. *Arch. Ges. Physiol.* 103:429. [26] Van Liere, E. J., J. C. Stickney, and D. F. Marsh. 1949. *Science* 109:489. [27] Wilhelmj, C. M., E. B. Waldmann, and T. F. McGuire. 1951. *Am. J. Physiol.* 166:296. [28] Woodbury, R. A., and B. E. Abreu. 1944. *Ibid.* 142:114. [29] Woodbury, R. A., and W. F. Hamilton. 1937. *Ibid.* 119:663. [30] Wu, C. H., and M. B. Visscher. 1947. *Federation Proc.* 6:231.

**79. VASCULAR AND CAPILLARY PRESSURES**

**Part I. VASCULAR PRESSURES: MAN**

Chamber or Vessel	Blood Pressure, mm Hg			Method	Refer-ence
	Systolic	Diastolic	Mean		
(A)	(B)	(C)	(D)	(E)	(F)
1 Medium veins	.....	.....	4-7	Direct measurement	3
2 Venae cavae	.....	.....	-5 to +5	Right heart catheterization	4
3 Right atrium	2-7	0-5	1-5	Right heart catheterization	2
4 Right ventricle	19-31	2-6	.....	Right heart catheterization	2
5 Pulmonary artery	16-29	5-13	10-18	Right heart catheterization	2
6 Pulmonary capillary	.....	.....	5-13	Right heart catheterization	1
7 Left atrium	6-21	1-12	2-12	Transseptal left heart catheterization	1
8 Left ventricle	90-130	5-12	.....	Transseptal left heart catheterization	1
9 Aorta	90-130	60-90	70-115	.....	4

*continued*

## 79. VASCULAR AND CAPILLARY PRESSURES

### Part I. VASCULAR PRESSURES: MAN

*Contributors:* Terry, Luther L., and Braunwald, Eugene

*References:* [1] Braunwald, E., et al. 1961. *Circulation* 24:267. [2] Fowler, N. O., et al. 1953. *Am. Heart J.* 46:264. [3] Snabel, T. G., Jr., et al. 1954. *Penna. Med. J.* 57:363. [4] Terry, L. L. Unpublished. Natl. Institutes of Health, Bethesda, Md., 1955.

### Part II. RELATIONSHIP OF PERIPHERAL ARTERIAL TO CENTRAL ARTERIAL PRESSURE: MAN

Values are expressed as percent of aorta pressure or of subclavian pressure near the aorta.

	Condition	Artery	Blood Pressure			
			Systolic	Diastolic	Mean	
	(A)	(B)	(C)	(D)	(E)	
1	Supine, at rest	Brachial	109	96	98	
2		Radial	112	93	94	
3		Femoral	110	94	96	
4	Supine, during exercise	Brachial	111	97	97	
5		Radial	113	93	93	
6		Femoral	101	95	97	
7		70° head-up tilt	Brachial	111	98	99
8		Radial	115	95	98	
9		Femoral	123	98	100	

*Contributor:* Terry, Luther L.

*Reference:* Kroeker, E. J., and E. H. Wood. 1955. *Circulation Res.* 3:623.

### Part III. VENOUS BLOOD PRESSURE: MAN

Reference level was the phlebostatic axis. Subjects were supine, breathing quietly. Forced respiration (e.g., Valsalva's maneuver) profoundly influences venous pressure. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Vein	Blood Pressure
		mm H <sub>2</sub> O
	(A)	(B)
	Median basilic vein at elbow	
1	3-5 yr	46(30-63)
2	5-10 yr	58(33-74)
3	Adult ♀	94(60-128)
4	Adult ♂	100(50-140)
5	Femoral	111(98-128)
6	Abdominal	115(70-160)
7	Dorsal metacarpal	130(70-170)
8	Great saphenous at ankle	150(110-190)
9	Dorsal pedal	175(124-210)

*Contributor:* Terry, Luther L.

*Reference:* Burch, G. E. 1950. *A primer of venous pressure.* Lea and Febiger, Philadelphia.

### Part IV. CAPILLARY BLOOD PRESSURE: VERTEBRATES

All measurements made directly by microcannulation [1]. **Capillary** (column D): A = arterial; V = venous. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Tissue	Species (Common Name)	Condition	Capillary	Pressure	Reference
					cm H <sub>2</sub> O	
	(A)	(B)	(C)	(D)	(E)	(F)
1	Eponychium (finger)	<i>Homo sapiens</i> (man)	Normal	A	43.5(28.6-65.0) <sup>1</sup>	5,7
2				V	16.5(8.0-24.5) <sup>1</sup>	
3			Hypertension	A	48.5(10.1-95.1)	2
4				V	30.8(12.8-58.0)	
5			Hyperemia	A	86.0(71.0-93.0)	7
6				V	{54.5-66.5}	
7	Mesentery	<i>Cavia porcellus</i> (guinea pig)	Decerebrate; anesthetized with veronal ether	A	38.5(31.0-49.0)	4,7
8				V	17.0(13.0-19.5)	
9		<i>Rattus</i> sp. (rat)	Decerebrate	A	30.0(22.0-34.0)	4,7
10				V	17.0(15.0-20.0)	
11		<i>Rana</i> sp. (frog)	Pithed	A	14.4(5.0-22.0)	3,7
12				V	10.1(6.7-18.0)	

<sup>1/1</sup> Varies directly with arteriolar vasodilation produced by emotion, heat, or trauma. Varies inversely with arteriolar vasoconstriction produced by emotion or cold. Varies minimally in a single capillary with time, and also from capillary to capillary. Varies directly with venous pressure as affected by hydrostatic pressure or venous obstruction.

*continued*



79. VASCULAR AND CAPILLARY PRESSURES: VERTEBRATES

Part IV. CAPILLARY BLOOD PRESSURE: VERTEBRATES

	Tissue	Species (Common Name)	Condition	Capil- lary	Pressure cm H <sub>2</sub> O	Refer- ence
	(A)	(B)	(C)	(D)	(E)	(F)
13	Muscle	<i>Rana</i> sp. (frog)	Normal; anesthetized with urethan	A	14.9(11.0-18.0)	6,7
14				V	9.5(7.0-12.7)	
15			Hyperemia	A	20.1(17.0-26.0)	7
16				V	16.0(12.0-17.5)	
17	Web	<i>Rana</i> sp. (frog)	Normal; anesthetized with urethan	A	13.9(10.0-19.0)	6,7
18				V	9.6(8.5-13.0)	
19			Normal; curarized	A	14.5(10.0-20.5)	7
20				V	10.0(8.5-15.5)	
21			Hyperemia	A	19.5(14.0-26.5)	7
22	V	16.5(15.0-17.5)				

*Contributors:* (a) Griffith, John Quintin, Jr., (b) Mendlowitz, Milton

*References:* [1] Carrier, E. B., and P. B. Rehberg. 1923. Skand. Arch. Physiol. 44:20. [2] Eichna, L. W., and J. Bordley, III. 1942. J. Clin. Invest. 21:711. [3] Landis, E. M. 1926. Am. J. Physiol. 75:548. [4] Landis, E. M. 1930. Ibid. 93:353. [5] Landis, E. M. 1930. Heart 15:209. [6] Landis, E. M. 1931. Am. J. Physiol. 98:704. [7] Landis, E. M. 1934. Physiol. Rev. 14:404.

# *Contrails*

## VIII. BLOOD

### 80. BLOOD GROUP SYSTEMS: MAN

The A-B-O, the M-N, and the Rh-Hr are the three most important blood group systems in man and are the ones most used in the study of human linkage, in tests for zygosity of twins, and in medicolegal problems of disputed parentage. Other blood group systems have been reported after an antibody giving reactions unrelated to previously described blood groups has been encountered in the serum of certain individuals (mothers of erythroblastotic babies [Kell, Kidd], patients who had received blood transfusions [Duffy, Lutheran]). The blood group systems in man depend on multiple allelic genes for their hereditary transmission. None of the blood-grouped genes is known to be sex-linked, and the genes of each blood group system appear to be located on different pairs of chromosomes, as is shown by their independent heredity. For information on other blood group systems, consult reference 1, Part I.

**Definitions:** **agglutinin** = an antibody aggregating a particular antigen; **agglutigen** = any substance that acts as an antigen and stimulates the production of agglutinin; **antisera** = a serum containing an antibody or antibodies; **antibody** = a modified serum globulin, synthesized by an animal in response to antigenic stimulus, that reacts specifically in vivo and in vitro with the homologous antigen; **antigen** = a high-molecular-weight substance, or complex, foreign to the blood stream, that, upon gaining access to the tissues of the animal, stimulates the formation of a specific antibody and reacts specifically in vivo or in vitro with the homologous antibody; **blood factors** = the specific serological properties by which an agglutigen is recognized, e.g., if blood cells are clumped by an antiserum assigned the symbol anti-**X**, they are said to have blood factor **X**; **gene** = the biologic unit of heredity, self-reproducing and located in a definite position (locus) on a particular chromosome; **allelic genes** = genes situated at corresponding loci in a pair of chromosomes; **genotype** = the fundamental hereditary constitution (or assortment of genes) of an individual; **phenotype** = the outward, visible expression of the hereditary constitution of an individual.

#### Part I. PHENOTYPES AND GENOTYPES OF THE A-B-O SYSTEM

The A-B-O blood groups are determined by two agglutinogens on the red blood cells (agglutigen A which occurs in two principal forms, A<sub>1</sub> and A<sub>2</sub>, and agglutigen B) and two corresponding, naturally occurring isoagglutinins in the serum (anti-**A** [alpha] and anti-**B** [beta]). The latter are regularly present in the serum when the corresponding agglutigen is absent from the red cells, except during the neonatal period when the antibody-producing mechanism is immature. Tests include examination of the serum for isoagglutinin content, as well as examination of the reaction of the red cells to anti-**A** and anti-**B** serums. Anti-**A** was obtained from type-B subjects, and anti-**B** from type-A subjects. Frequencies (columns B and I) are for populations of European origin. *Xx* genes (column J) segregate independently of and modify *ABO* genes.

Designation	Frequency %	Phenotype				Plasma Agglutinin	Genotype		
		A	A <sub>1</sub>	B	H		ABO Genes	Frequency %	Xx Genes
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 O	45.0	-	-	-	+	Anti- <b>A</b> , anti- <b>B</b>	OO	45.0	XX or Xx
2 A <sub>1</sub>	31.0	+	+	-	-	Anti- <b>B</b> , occasionally anti- <b>H</b>	A <sub>1</sub> A <sub>1</sub>	3.5	Xx
3							A <sub>1</sub> A <sub>2</sub>	2.6	
4							A <sub>1</sub> O	25.0	
5 A <sub>2</sub>	9.6	+	-	-	+	Anti- <b>B</b> , occasionally anti- <b>A<sub>1</sub></b>	A <sub>2</sub> A <sub>2</sub>	0.5	
6							A <sub>2</sub> O	9.2	
7 B	10.0	-	-	+		Anti- <b>A</b>	BB	0.7	
8							BO	9.3	
9 A <sub>1</sub> B	2.9	+	+	+	-	Occasionally anti- <b>H</b>	A <sub>1</sub> B	2.9	
10 A <sub>2</sub> B	1.1	+	-	+		Occasionally anti- <b>A<sub>1</sub></b>	A <sub>2</sub> B	1.1	
11 "Bombay type"	Rare	-	-	-	-	Anti- <b>A</b> , anti- <b>B</b> anti- <b>H</b>	A, B, O <sup>1</sup>	Rare	xx

/:/: Probably any combination is possible.

*Contributor:* Allen, Fred H., Jr.

*References:* [1] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Race, R. R., and R. Sanger. 1958. Blood groups

*continued*

## 80. BLOOD GROUP SYSTEMS: MAN

### Part I. PHENOTYPES AND GENOTYPES OF THE A-B-O SYSTEM

in man. Ed. 3. Blackwell, Oxford. [3] Wiener, A. S., and I. B. Wexler. 1958. Heredity of the blood groups. Grune and Stratton, New York.

### Part II. PARTIAL LIST OF ALLELIC GENES OF THE M-N SYSTEM

So far as is known, the M-N phenotype of a person is exactly what would be expected from the genotype, no suppressing effect of any one M-N gene on any other having been demonstrated. In blood typing, anti-M is obtained from humans or immunized rabbits, anti-N from immunized rabbits or from seeds of *Vicia graminea*, anti-Hu and anti-He are obtained only from immunized rabbits, and other antisera only from sensitized humans. Blood factor frequencies are for Europeans and are given in parentheses.

Gene	Gene Frequency %	Reaction with Anti-										
		M (79%)	M <sup>G</sup> (Rare)	N (71%)	S (55%)	s (89%)	U (99+%)	Hu (Rare <sup>1</sup> )	He (Rare <sup>1</sup> )	Mi <sup>A</sup> (Rare <sup>1</sup> )	Vw (Rare <sup>1</sup> )	Vr (Rare <sup>1</sup> )
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)
1 MS	25	+	-	+ <sup>2</sup>	+	-	+	-	±	±	-	-
2 Ms	28	+	-	+ <sup>2</sup>	-	+	+	-	±	-	-	±
3 MS <sup>u</sup>	Rare	+	-	-	-	-	-	-	-	-	-	-
4 NS	8	-	-	+	+	-	+	-	±	-	-	-
5 Ns	39	-	-	+	-	+	+	±	±	±	±	-
6 NS <sup>u</sup>	Rare	-	-	+	-	-	-	-	-	-	-	-
7 M <sup>G</sup>	Rare	-	+	-	-	-	-	-	-	-	-	-
8 MU	Rare	+	-	-	-	-	+	-	-	-	-	-
9 NU	Rare	-	-	+	-	-	+	-	-	-	-	-

[1] Factors Hu, He, Mi<sup>A</sup>, Vw, and Vr undoubtedly occur in other combinations than are shown here, but there is, as yet, insufficient information about these factors to prepare a complete table of the M-N genes. Other antigenic factors in the MN system are M<sub>1</sub>, M<sup>C</sup>, M<sup>O</sup>, Mu, Mt<sup>A</sup>, St<sup>A</sup>, Ri<sup>3</sup>, Cl<sup>A</sup>, and Ny<sup>A</sup>. All except M<sup>O</sup> are rare. M<sup>O</sup> is produced by M genes and by genes that produce He. [2] These genes produce a small amount of N factor or something which cross-reacts with anti-N.

Contributor: Allen, Fred H., Jr.

References: [1] Allen, F. H., Jr., P. A. Corcoran, and F. R. Ellis. 1960. Vox Sanguinis 5:224. [2] Allen, F. H., Jr., et al. 1958. Ibid. 3:81. [3] Cleghorn, T. E. 1961. M.D. Thesis. Univ. Sheffield, England. [4] Cleghorn, T. E. 1962. Nature 195:297. [5] Dunsford, I., E. W. Ikin, and A. E. Mourant. 1953. Ibid. 172:688. [6] Graydon, J. J. 1946. Med. J. Australia 2:9. [7] Ikin, E. W., and A. E. Mourant. 1951. Brit. Med. J. 1:456. [8] Jack, J. A., et al. 1960. Nature 186:642. [9] Landsteiner, K., and P. Levine. 1927. Proc. Soc. Exptl. Biol. Med. 24:600. [10] Landsteiner, K., W. R. Stratton, and M. W. Chase. 1934. J. Immunol. 27:469. [11] Levine, P., et al. 1951. Proc. Soc. Exptl. Biol. Med. 77:402. [12] Levine, P., et al. 1951. Ibid. 78:218. [13] Sanger, R., and R. R. Race. 1947. Nature 160:505. [14] Swanson, J., and G. A. Matson. 1962. Vox Sanguinis 7:585. [15] Van der Hart, M., et al. 1958. Ibid. 3:261. [16] Walsh, R. J., and C. Montgomery. 1947. Nature 160:504. [17] Wiener, A. S., and R. E. Rosenfield. 1961. J. Immunol. 87:376. [18] Wiener, A. S., L. J. Unger, and E. B. Gordon. 1953. J. Am. Med. Assoc. 153:1444.

continued



## 80. BLOOD GROUP SYSTEMS: MAN

### Part III. PHENOTYPES AND GENOTYPES OF THE Rh-Hr SYSTEM

The Rh-Hr system is the most complicated of the human blood systems. At the present time, four principal Rh blood factors ( $Rh_0$ ,  $rh^1$ ,  $rh''$ , and  $rh^w$ ) and three principal Hr factors ( $hr^1$ ,  $hr''$ , and  $hr$ ) are recognized. However, antisera for only five of these seven factors are readily available for routine clinical and medicolegal work (anti- $Rh_0$ , anti- $rh^1$ , anti- $rh''$ , anti- $hr^1$  and anti- $hr''$ ). The  $Rh_0$  factor is the most common source of clinical symptoms, as it is the most antigenic of the Rh-Hr factors. It appears to represent a special structure within the Rh-Hr agglutinin, since red cells can be coated with the  $Rh_0$ -blocking antibody without interfering with the reactions of the red cells with other antibodies such as anti- $rh^1$ , anti- $rh''$ , anti- $hr^1$ . Frequencies are for white residents of New York City. Frequencies (columns E and J) are based on estimated gene frequencies:  $r = 38\%$ ;  $r' = 0.6\%$ ;  $r'' = 0.5\%$ ;  $r^y = 0.01\%$ ;  $r^w = 0.005\%$ ;  $R^0 = 2.7\%$ ;  $R^1 = 41\%$ ;  $R^2 = 15\%$ ;  $R^z = 0.2\%$ ;  $R^{lw} = 2\%$ .

2 Rh Phenotypes			12 Rh Phenotypes					28 Rh-Hr Phenotypes					55 Genotypes (N)	
Designation	Frequency %	Reaction with Anti- $Rh_0$ (or Anti-rhesus)	Designation	Frequency %	Reaction with Anti- $rh^1$ $rh''$ $rh^w$			Designation	Frequency %	Reaction with Anti- $hr^1$ $hr''$ $hr$				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)		
1	Rh n e g a t i v e	-	rh	14.4	-	-	-	rh	14.4	+	+	+	$rr$	
2			$rh^1$	0.46	+	-	-	$rh^1rh$	0.46	+	+	+	$r^1r$	
3									$rh^1rh^1$	0.0036	-	+	-	$r^1r^1$
4			$rh^w$	0.004	+	-	+	$rh^w rh$	0.004	+	+	+	+	$r^w r$
5									$rh^w rh^1$	0.00006	-	+	-	$r^w r^1$ or $r^w r^1 w$
6			$rh''$	0.38	-	+	-	$rh'' rh$	0.38	+	+	+	+	$r'' r$
7									$rh'' rh''$	0.0025	+	-	-	$r'' r''$
8			$rh^y$	0.01	+	+	-	$rh^y rh$	0.006	+	+	-	-	$r^y r$
9									$rh_y rh$	0.008	+	+	+	$r^y r$
10									$rh_y rh^1$	0.0001	-	+	-	$r^y r^1$
11									$rh_y rh''$	0.0001	+	-	-	$r^y r''$
12									$rh_y rh_y$	0.000001	-	-	-	$r^y r^y$
13									$rh_y^w rh''$	0.00005	+	+	-	$r^w r^1$
14									$rh_y^w rh^1$	0.000001	-	+	-	$r^w r^1 w$
15	Rh p o s i t i v e	+	$Rh_0$	2.1	-	-	-	$Rh_0$	2.1	+	+	+	$R^0 R^0$ or $R^0 r$	
16			$Rh_1$	50.7	+	-	-	$Rh_1 rh$	33.4	+	+	+	$R^1 r$ , $R^1 R^0$ , or $R^0 r^1$	
17									$Rh_1 Rh_1$	17.3	-	+	-	$R^1 R^1$ or $R^1 r^1$
18			$Rh_1^w$	3.3	+	-	+	$Rh_1^w rh$	1.6	+	+	+	+	$R^1 w r$ , $R^1 w R^0$ , or $R^0 r^1 w$
19									$Rh_1^w Rh_1$	1.7	-	+	-	$R^1 w R^1$ , $R^1 r^1 w$ , $R^1 w r^1$ , $R^1 w R^1 w$ , or $R^1 w r^1 w$
20			$Rh_2$	14.6	-	+	-	$Rh_2 rh$	12.2	+	+	+	+	$R^2 r$ , $R^2 R^0$ , or $R^0 r^2$
21									$Rh_2 Rh_2$	2.4	+	-	-	$R^2 R^2$ or $R^2 r^2$
22			$Rh_z$	13.4	+	+	-	$Rh_1 Rh_2$	12.9	+	+	-	-	$R^1 R^2$ , $R^1 r^2$ , or $R^2 r^1$
23									$Rh_z rh$	0.2	+	+	+	$R^z r$ , $R^z R^0$ , or $R^0 r^z$
24									$Rh_z Rh_1$	0.2	-	+	-	$R^z R^1$ , $R^z r^1$ , or $R^1 r^z$
25									$Rh_z Rh_2$	0.07	+	-	-	$R^z R^2$ , $R^z r^2$ , or $R^2 r^z$
26									$Rh_z Rh_z$	0.0004	-	-	-	$R^z R^z$ or $R^z r^z$
27			$Rh_z^w$	0.6	+	+	+	$Rh_1^w Rh_2$	0.6	+	+	-	-	$R^1 w R^2$ , $R^1 w r^2$ , or $R^2 r^1 w$
28									$Rh_z^w Rh_1$	0.008	-	+	-	$R^1 w R^z$ , $R^1 w r^z$ , or $R^z r^1 w$

Contributor: Wiener, Alexander S.

Reference: Wiener, A. S., and I. B. Wexler. 1963. An Rh-Hr syllabus; the types and their application. Ed. 2. Grune and Stratton, New York.

continued

## 80. BLOOD GROUP SYSTEMS: MAN

### Part IV. PARTIAL LIST OF ALLELIC GENES OF THE Rh-Hr SYSTEM

Gene	Agglu- tinogen	Reaction with Anti-																		
		Rh <sub>0</sub>	rh'	rh <sup>w1</sup>	rh''	rh <sup>w2</sup>	hr'	hr''	hr	Rh <sup>A</sup>	Rh <sup>B</sup>	Rh <sup>C</sup>	Rh <sup>D</sup>	rh <sup>x</sup>	rh <sub>1</sub>	rh <sup>G</sup>	Hr	hr <sup>S</sup>	hr <sup>V</sup>	hr <sup>N</sup>
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)
1	r	rh	-	-	-	-	+	+	+	-	-	-	-	-	-	-	+	+	-	-
2	r'	rh'	-	+	-	-	-	-	+	-	-	-	-	-	+	+	+	+	-	-
3	r' <sup>w</sup>	rh' <sup>w</sup>	-	+	+	-	-	-	+	-	-	-	-	-	+	+	+	+	-	-
4	r''	rh''	-	-	-	+	-	+	-	-	-	-	-	-	-	-	+	-	-	-
5	r <sup>y</sup>	rh <sub>y</sub>	-	+	-	+	-	-	-	-	-	-	-	-	-	+	+	-	-	-
6	R <sup>0</sup>	Rh <sub>0</sub>	+	-	-	-	-	+	+	+	+	+	+	-	-	+	+	+	-	-
7	R <sup>1</sup>	Rh <sub>1</sub>	+	+	-	-	-	-	+	-	+	+	+	-	+	+	+	+	-	-
8	R <sup>1w</sup>	Rh <sub>1</sub> <sup>w</sup>	+	+	+	-	-	-	+	-	+	+	+	-	+	+	+	+	-	-
9	R <sup>2</sup>	Rh <sub>2</sub>	+	-	-	+	-	+	-	-	+	+	+	-	-	+	+	-	-	-
10	R <sup>z</sup>	Rh <sub>z</sub>	+	+	-	+	-	-	-	-	+	+	+	+	-	-	+	+	-	-
11	R <sup>0</sup>	Rh <sub>0</sub>	±	-	-	-	-	+	+	+	±	±	±	±	-	-	+	+	-	-
12	R <sup>1</sup>	Rh <sub>1</sub>	±	+	-	-	-	-	+	-	±	±	±	±	-	+	+	+	-	-
13	R <sup>2</sup>	Rh <sub>2</sub>	±	-	-	+	-	+	-	-	±	±	±	±	-	-	+	+	-	-
14	R̄ <sup>0</sup>	R̄h <sub>0</sub>	++	-	-	-	-	-	-	-	+	+	+	+	-	-	+	-	-	-
15	R̄ <sup>w</sup>	R̄h <sub>0</sub> <sup>w</sup>	++	-	+	-	-	-	-	-	+	+	+	+	-	-	+	-	-	-
16	R̄ <sup>0</sup>	R̄h <sub>0</sub>	+	-	-	-	-	+	-	+	+	+	+	-	-	+	-	-	-	-
17	R̂ <sup>0</sup>	R̂h <sub>0</sub>	+	-	-	-	-	+	+	+	+	+	+	-	-	+	-	-	-	-
18	r̄	r̄h	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
19	R <sup>1x</sup>	Rh <sub>1</sub> <sup>x</sup>	+	+	-	-	-	-	+	-	+	+	+	+	+	+	+	+	-	-
20	R <sup>2w</sup>	Rh <sub>2</sub> <sup>w</sup>	+	-	-	+	+	+	-	-	+	+	+	-	-	+	+	-	-	-
21	r <sup>v</sup>	rh <sup>v</sup>	-	-	-	-	-	+	+	+	-	-	-	-	-	-	+	+	+	+
22	R <sup>ov</sup>	Rh <sub>0</sub> <sup>v</sup>	+	-	-	-	-	+	+	+	+	+	+	-	-	+	+	+	+	+
23	r' <sup>N</sup>	rh' <sup>N</sup>	-	+	-	-	-	+	+	+	-	-	-	-	-	+	+	+	-	+
24	R <sup>1ab</sup>	Rh <sub>1</sub> <sup>ab</sup>	+	+	-	-	-	-	+	-	-	-	+	+	-	+	+	+	-	-
25	R <sup>2b</sup>	Rh <sub>2</sub> <sup>b</sup>	+	-	-	+	-	+	-	-	+	-	+	-	-	+	+	-	-	-
26	R <sup>2c</sup>	Rh <sub>2</sub> <sup>c</sup>	±	-	-	+	-	+	-	-	+	+	-	+	-	-	+	-	-	-
27	R <sup>od</sup>	Rh <sub>0</sub> <sup>d</sup>	+	-	-	-	-	+	+	+	+	+	+	-	-	-	+	+	-	-
28	r <sup>G</sup>	rh <sup>G</sup>	-	-	-	-	-	-	+	-	-	-	-	-	-	+	+	-	-	-

Contributor: Wiener, Alexander S.

Reference: Wiener, A. S., and I. B. Wexler. 1963. An Rh-Hr syllabus; the types and their applications. Ed. 2. Grune and Stratton, New York.

## 81. HEREDITY OF BLOOD GROUPS AND TYPES: MAN

Because of the possibility of coincidence, it is considered an inconclusive finding when the blood type of the child matches the blood type of the putative parent. Therefore, in cases of disputed parentage, blood tests can be used only to exclude the claim of maternity or paternity.

### Part I. A-B-O EXCLUSION

Parental Phenotype Combination		Blood Group of Child that Refutes	
Putative Mother	Putative Father	Putative Maternity	Putative Paternity
(A)	(B)	(C)	(D)
1 O	O	AB	A, B
2	A	AB	B
3	B	AB	A
4	AB	AB	O
5 A	O	None	B, AB
6	A	None	B, AB
7	B	None	None
8	AB	None	O
9 B	O	None	A, AB
10	A	None	None
11	B	None	A, AB
12	AB	None	O
13 AB	O	O	AB
14	A	O	None
15	B	O	None
16	AB	O	None

### Part II. M-N EXCLUSION

Parental Phenotype Combination		Blood Type of Child that Refutes	
Putative Mother	Putative Father	Putative Maternity	Putative Paternity
(A)	(B)	(C)	(D)
1 M	M	N	MN
2	N	N	M
3	MN	N	None
4 N	M	M	N
5	N	M	MN
6	MN	M	None
7 MN	M	None	N
8	N	None	M
9	MN	None	None

*Contributor:* Wiener, Alexander S.

*Reference:* See Part I.

*Contributor:* Wiener, Alexander S.

*Reference:* Wiener, A. S., and I. B. Wexler. 1958. Heredity of the blood groups. Grune and Stratton, New York.

### Part III. Rh-Hr EXCLUSION

This table is applicable only to matings in which at least one of the parents is Rh<sub>0</sub>-positive. Where both parents are Rh<sub>0</sub>-negative, all Rh<sub>0</sub>-positive children are necessarily excluded. Boldface figures represent phenotypes of children for whom putative *maternity* is excluded; all other figures shown represent phenotypes of children for whom putative *paternity* is excluded. Code numbers and corresponding phenotypes are given in the column headings, e.g., 1 is the code number for phenotypes rh and Rh<sub>0</sub>.

Phenotype of Putative Mother	Phenotype of Putative Father									
	1 rh Rh <sub>0</sub>	2 rh'rh Rh <sub>1</sub> rh	3 rh'rh' Rh <sub>1</sub> Rh <sub>1</sub>	4 rh''rh Rh <sub>2</sub> rh	5 rh''rh'' Rh <sub>2</sub> Rh <sub>2</sub>	6a rh'rh'' Rh <sub>1</sub> Rh <sub>2</sub>	6b rh <sub>y</sub> rh Rh <sub>z</sub> Rh <sub>0</sub>	7 rh <sub>y</sub> rh' Rh <sub>z</sub> Rh <sub>1</sub>	8 rh <sub>y</sub> rh'' Rh <sub>z</sub> Rh <sub>2</sub>	9 rh <sub>y</sub> rh <sub>y</sub> Rh <sub>z</sub> Rh <sub>z</sub>
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
1 rh Rh <sub>0</sub>	2,3,4,5, 6a,6b, 7,8,9	3,4,5,6a, 6b,7,8, 9	1,3,4,5, 6a,6b, 7,8,9	2,3,5,6a, 6b,7,8, 9	1,2,3,5, 6a,6b, 7,8,9	1,3,5,6a, 6b,7,8, 9	2,3,4,5, 6a,7,8, 9	1,3,4,5, 6a,7,8, 9	1,2,3,5, 6a,7,8, 9	1,2,3,4, 5,6a,7, 8,9
2 rh'rh Rh <sub>1</sub> rh	3,4,5, 6a,6b, 7,8,9	4,5,6a, 6b,7,8, 9	1,4,5, 6a,6b, 7,8,9	3,5,6b,7, 8,9	1,2,3,5, 6b,7,8, 9	1,5,6b,7, 8,9	3,4,5,6a, 8,9	1,4,5,6a, 8,9	1,2,3,5, 8,9	1,2,3,4, 5,6a,8, 9
3 rh'rh' Rh <sub>1</sub> Rh <sub>1</sub>	1,3,4,5, 6a,6b, 7,8,9	1,4,5,6a, 6b,7,8, 9	1,2,4,5, 6a,6b, 7,8,9	1,3,4,5, 6b,7,8, 9	1,2,3,4, 5,6b,7, 8,9	1,2,4,5, 6b,7,8, 9	1,3,4,5, 6a,6b, 8,9	1,2,4,5, 6a,6b, 8,9	1,2,3,4, 5,6b,8, 9	1,2,3,4, 5,6a, 6b,8,9
4 rh''rh Rh <sub>2</sub> rh	2,3,5, 6a,6b, 7,8,9	3,5,6b,7, 8,9	1,3,4,5, 6b,7, 8,9	2,3,6a, 6b,7,8, 9	1,2,3,6a, 6b,7,8, 9	1,3,6b,7, 8,9	2,3,5,6a, 7,9	1,3,4,5, 7,9	1,2,3,6a, 7,9	1,2,3,4, 5,6a,7, 9
5 rh''rh'' Rh <sub>2</sub> Rh <sub>2</sub>	1,2,3,5, 6a,6b, 7,8,9	1,2,3,5, 6b,7,8, 9	1,2,3,4, 5,6b, 7,8,9	1,2,3,6a, 6b,7,8, 9	1,2,3,4, 6a,6b, 7,8,9	1,2,3,4, 6b,7,8, 9	1,2,3,5, 6a,6b, 7,9	1,2,3,4, 5,6b,7, 9	1,2,3,4, 6a,6b, 7,9	1,2,3,4, 5,6a, 6b,7,9
6 6a rh'rh'' Rh <sub>1</sub> Rh <sub>2</sub>	1,3,5,6a, 6b,7,8, 9	1,5,6b,7, 8,9	1,2,4,5, 6b,7, 8,9	1,3,6b,7, 8,9	1,2,3,4, 6b,7,8, 9	1,2,4,6b, 7,8,9	1,3,5,6a, 6b,9	1,2,4,5, 6b,9	1,2,3,4, 6b,9	1,2,3,4, 5,6a, 6b,9
7 6b rh <sub>y</sub> rh Rh <sub>z</sub> Rh <sub>0</sub>	2,3,4,5, 6a,7,8, 9	3,4,5,6a, 8,9	1,3,4,5, 6a,6b, 8,9	2,3,5,6a, 7,9	1,2,3,5, 6a,6b, 7,9	1,3,5,6a, 6b,9	2,3,4,5, 6a,7,8	1,3,4,5, 6a,8	1,2,3,5, 6a,7	1,2,3,4, 5,6a, 7,8

*continued*

## 81. HEREDITY OF BLOOD GROUPS AND TYPES: MAN

### Part III. Rh-Hr EXCLUSION

	Phenotype of Putative Mother	Phenotype of Putative Father									
		1 rh Rh <sub>0</sub>	2 rh'rh Rh <sub>1</sub> rh	3 rh'rh' Rh <sub>1</sub> Rh <sub>1</sub>	4 rh''rh Rh <sub>2</sub> rh	5 rh''rh'' Rh <sub>2</sub> Rh <sub>2</sub>	6a rh'rh'' Rh <sub>1</sub> Rh <sub>2</sub>	6b rh <sub>y</sub> rh Rh <sub>2</sub> Rh <sub>0</sub>	7 rh <sub>y</sub> rh' Rh <sub>2</sub> Rh <sub>1</sub>	8 rh <sub>y</sub> rh'' Rh <sub>2</sub> Rh <sub>2</sub>	9 rh <sub>y</sub> rh <sub>y</sub> Rh <sub>2</sub> Rh <sub>2</sub>
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
8	7	rh <sub>y</sub> rh' Rh <sub>2</sub> Rh <sub>1</sub>	1,3,4,5, 6a,7,8, 9	1,4,5,6a, 8,9	1,2,4,5, 6a,6b, 8,9	1,3,4,5, 7,9	1,2,3,4, 5,6b,7, 9	1,2,4,5, 6b,9	1,2,4,5, 6a,6b, 8	1,2,3,4, 5,6b	1,2,3,4, 5,6a, 6b,8
	9	rh <sub>y</sub> rh'' Rh <sub>2</sub> Rh <sub>2</sub>	1,2,3,5, 6a,7,8, 9	1,2,3,5, 8,9	1,2,3,4, 5,6b, 8,9	1,2,3,6a, 7,9	1,2,3,4, 6a,6b, 7,9	1,2,3,4, 6b,9	1,2,3,5, 6a,7	1,2,3,4, 5,6b	1,2,3,4, 6a,6b, 7
10	9	rh <sub>y</sub> rh <sub>y</sub> Rh <sub>2</sub> Rh <sub>2</sub>	1,2,3,4, 5,6a,7, 8,9	1,2,3,4, 5,6a,8	1,2,3,4, 5,6a, 6b,8,9	1,2,3,4, 5,6a,7, 9	1,2,3,4, 5,6a, 6b,7,9	1,2,3,4, 5,6a, 6b,9	1,2,3,4, 5,6a,7, 8	1,2,3,4, 5,6a, 6b,8	1,2,3,4, 5,6a, 6b,7,8

Contributor: Wiener, Alexander S.

Reference: Wiener, A. S. 1963. J. Forensic Med. 10(1):6.

## 82. DISTRIBUTION OF BLOOD GROUPS AND TYPES IN VARIOUS POPULATIONS: MAN

### Part I. A-B-O GROUPS

	Population	Location	No. of Subjects	Frequency, %			
				Group O	Group A	Group B	Group AB
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Ainu	Shizunai	504	36.7	44.5	14.7	5.1
2	American Indian						
3	Blackfoot	Montana	115	23.5	76.5	0	0
4	Navaho	Arizona	457	72.9	26.9	0.2	0.2
5	Pueblo	New Mexico	310	78.4	20.0	1.6	0
6	Ute	Utah	104	98.1	1.9	0	0
7	Armenian	Vicinity of Marash, Turkey	330	27.3	53.9	12.7	6.1
8	Australian aborigine	Australia	805	53.1	44.7	2.1	0
9	Australian white	Sydney	220	44.6	43.6	9.1	2.7
10	Basque	San Sebastian, Spain	91	57.2	41.7	1.1	0
11	Bedouin						
12	Iraqi	Vicinity of Baghdad	338	40.8	26.6	25.8	6.8
13	Rwala	Syrian Desert	208	43.3	22.1	30.3	4.3
14	Belgian	Liège	3,500	46.7	41.9	8.3	3.1
15	Chinese	Peking	1,000	30.7	25.1	34.2	10.0
16	Danish	Copenhagen	14,304	40.6	44.0	10.9	4.5
17	Dutch	Amsterdam	23,643	44.4	43.2	8.9	3.5
18	Egyptian	Cairo	502	27.3	38.5	25.5	8.8
19	English	Southern England	106,477	45.2	43.2	8.5	3.1
20	Eskimo	Southwest Greenland	1,063	46.0	46.1	4.9	3.0
21	Estonian	Central, southern, and southeastern Estonia	1,844	32.3	36.6	22.4	8.7
22	Fijian	Fiji Islands	160	43.8	43.1	9.4	3.8
23	Filipino	Philippine Islands	382	45.0	22.0	27.0	6.0
24	Finnish	Finland	23,200	34.1	41.0	18.0	6.9
25	French	Paris	14,303	42.7	45.6	8.3	3.3
26	German	Berlin	39,174	36.5	42.5	14.5	6.5
27	Greek	Athens	21,635	43.5	38.6	13.1	4.8
28	Hindu	Calcutta	6,247	32.4	24.1	36.2	7.3
29	Hungarian	Budapest	624	36.1	41.8	15.9	6.2
30	Indonesian	Djakarta	7,129	39.2	26.8	27.3	6.7
31	Irish	Northern Ireland	10,784	52.0	34.7	10.4	2.9
32	Italian	Rome and vicinity	20,051	44.7	40.0	11.4	3.8
33	Japanese	Tokyo	33,834	31.2	38.4	21.8	8.6
34	Korean	South Korea (north of Seoul)	1,000	27.0	32.0	29.0	12.0
35	Norwegian	Oslo	8,292	37.8	50.0	8.2	4.0

continued



## 82. DISTRIBUTION OF BLOOD GROUPS AND TYPES IN VARIOUS POPULATIONS: MAN

### Part I. A-B-O GROUPS

	Population	Location	No. of Subjects	Frequency, %			
				Group O	Group A	Group B	Group AB
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
34	Pakistani, Punjabi	Vicinity of Quetta, West Pakistan	10,000	30.6	24.5	34.8	10.0
35	Papuan	Papua	753	53.7	26.8	16.3	3.2
36	Polish	Warsaw	2,886	33.1	38.9	20.1	7.9
37	Portuguese	Lisbon	7,502	41.8	47.9	7.6	2.7
38	Puerto Rican	Puerto Rico	429	48.7	38.7	9.6	3.0
39	Russian	Leningrad	1,800	28.3	39.5	22.9	9.3
40	Scottish	Glasgow	456	47.6	38.4	9.0	5.0
41	Siamese	Bangkok	6,267	37.3	21.8	33.1	7.8
42	Swedish	Kopparberg	10,732	38.5	44.7	10.9	5.8
43	Ukrainian	Kharkov	310	36.4	38.4	21.6	3.6
44	USA Negro	Iowa	6,722	49.1	26.5	20.1	4.3
45	USA white	Rochester, New York	23,787	44.4	41.8	10.1	3.8
46	Welsh	North Wales	192	47.9	32.8	16.1	3.1
47	Yugoslavian	Yugoslavia	1,527	32.8	42.7	17.9	6.6

*Contributors:* (a) Levine, Philip, (b) Levine, Victor E., (c) Wiener, Alexander S.

*References:* [1] Boyd, W. C. 1939. *Tabulae Biologicae* 17:113. [2] Boyd, W. C. 1950. *Genetics and the races of man*. Little and Brown, Boston. [3] Mourant, A. E., A. C. Kopec, and K. Domaniewska-Sobczak. 1958. *The ABO blood groups*. Blackwell, Oxford. [4] Wiener, A. S. 1943. *Blood groups and transfusion*, Ed. 3. C. C. Thomas, Springfield, Ill.

### Part II. M-N TYPES

	Population	Location	No. of Subjects	Frequency, %		
				Type M	Type N	Type MN
	(A)	(B)	(C)	(D)	(E)	(F)
1	Ainu	Shizunai	504	17.9	31.9	50.2
2	American Indian					
3	Blackfoot	Montana	95	54.7	5.3	40.0
4	Navaho	New Mexico	361	84.5	1.1	14.4
5	Pueblo	New Mexico	140	59.3	7.9	32.8
6	Ute	Utah	104	58.7	6.7	34.6
7	Armenian	Vicinity of Marash, Turkey	332	32.8	20.2	47.0
8	Australian aborigine	Australia	730	3.0	67.4	29.6
9	Basque	Spain	91	23.1	25.3	51.6
10	Bedouin					
11	Iraqi	Vicinity of Baghdad	338	38.2	13.6	48.2
12	Rwala	Syrian Desert	208	57.5	5.8	36.7
13	Belgian	Liège	3,100	28.9	20.8	50.3
14	Chinese	Hong Kong	1,029	33.2	18.2	48.6
15	Danish	Copenhagen	2,023	29.1	21.4	49.5
16	Egyptian	Cairo	613	28.3	23.1	48.6
17	English	London	1,522	30.5	21.4	48.2
18	Eskimo	East Greenland	569	83.5	0.9	15.6
19		Southwest Greenland	1,063	66.2	2.9	31.0
20	Estonian	Estonia	310	34.8	15.5	49.7
21	Fijian	Fiji Islands	200	11.0	44.5	44.5
22	Filipino	Philippine Islands	382	25.9	23.8	50.3
23	Finnish	Finland	6,926	42.3	13.7	44.0
24	French	Paris	1,400	30.1	19.8	50.1
25	German	Germany	40,255	30.2	19.7	50.0
26	Hindu	India	300	42.7	10.7	46.7
27	Hungarian	Budapest	624	33.5	18.6	47.9
28	Irish	Dublin	399	30.0	23.3	46.7
29	Italian	Modena and Sicily	736	28.9	17.1	53.9
30	Japanese	Japan	7,551	29.0	21.1	49.9
31	Korean	Korea	836	27.9	20.8	51.4

*continued*

**82. DISTRIBUTION OF BLOOD GROUPS AND TYPES IN VARIOUS POPULATIONS: MAN**

**Part II. M-N TYPES**

Population	Location	No. of Subjects	Frequency, %		
			Type M	Type N	Type MN
(A)	(B)	(C)	(D)	(E)	(F)
30 Papuan	Papua	200	7.0	69.0	24.0
31 Polish	Poland	600	28.2	22.8	49.0
32 Russian	Leningrad	763	32.2	21.2	46.5
33 Scottish	Glasgow	456	35.0	17.1	47.9
34 Swedish	Sweden	1,200	36.1	16.9	47.0
35 Ukrainian	Kharkov	310	36.1	19.6	44.3
36 USA Negro	New York City	278	28.4	21.9	49.6
37 USA white	New York City; Boston; Columbus, Ohio	6,129	29.2	21.3	49.6
38 Welsh	North Wales	192	30.7	14.0	55.3
39 Yugoslavian	Yugoslavia	1,527	30.3	17.9	51.8

Contributors: (a) Levine, Philip, (b) Wiener, Alexander S., (c) Levine, Victor E.

References: [1] Boyd, W. C. 1939. *Tabulae Biologicae* 17:113. [2] Boyd, W. C. 1950. *Genetics and the races of man*. Little and Brown, Boston. [3] Wiener, A. S. 1943. *Blood groups and transfusion*. Ed. 3. C. C. Thomas, Springfield, Ill.

**Part III. Rh-Hr TYPES**

Population	No. of Subjects	Rh Positive Frequency, %						Rh Negative Frequency, %				Reference
		Rh <sub>0</sub>	Rh <sub>1</sub> Rh <sub>1</sub>	Rh <sub>1</sub> rh	Rh <sub>2</sub>	Rh <sub>1</sub> Rh <sub>2</sub>	Rh <sub>1</sub> Rh <sub>2</sub>	rh	rh'	rh''	rh'rh''	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)
1 American Indian												
1 In Mexico	95	1.1	40.7	7.4	9.5	38.1	3.1	0	0	0	0	5
2 In Oklahoma	105	2.9	34.3	5.7	17.1	36.2	2.9	0	0.9	0	0	5
3 Ute	104	0		33.7	28.8	37.5	...	0	0	0	0	1
4 Asiatic Indian	156	1.9		70.5	5.1	12.8	...	7.1	2.6	0	0	5
5 Australian aborigine	100	4.0	39.0	14.0	21.0	15.0	6.0	0	1.0	0	0	5
6 Australian white	350	0.6		54.0	12.6	16.6	...	14.9	0.9	0.6	0	5
7 Basque	167	0.6	7.8	47.3	7.8	6.0	0	28.8	1.8	0	0	3
8 Chinese	132	0.9		60.6	3.0	34.1	...	1.5	0	0	0	5
9 Dutch	200	1.5		51.5	12.3	17.7	...	15.4	1.5	0	0	5
10 Egyptian	...	11.5	25.5	39.7	9.2	8.2	...	5.9	...	...	....	4
11 English	927	2.5	19.7	35.2	12.2	13.6	0.1	14.8	0.7	1.3	0	5
12 Eskimo	315	1.0		34.9	19.7	44.4	...	0	0	0	0	2
13 Filipino	100	0		87.0	2.0	11.0	...	0	0	0	0	5
14 German	...	2.0	19.5	35.6	13.0	13.9	0.4	14.4	0.5	0.8	....	4
15 Hindu	...	2.9	35.2	32.4	3.8	16.2	0	7.6	1.9	...	....	4
16 Indonesian	200	0.5		74.0	2.5	22.5	0	0	0	0	0.5	5
17 Italian	...	1.3	23.3	37.3	9.6	11.8	0.7	14.8	0.5	0.5	0.3	4
18 Japanese	150	0		37.4	13.3	47.3	...	1.3	0	0	0.7	5
19 Norwegian	...	1.5	15.9	35.6	13.8	14.7	...	16.2	0.7	1.2	....	4
20 Papuan	100	0	89.0	4.0	0	4.0	3.0	0	0	0	0	5
21 Puerto Rican	179	15.1		39.1	19.6	14.0	...	10.1	1.7	0.5	0	5
22 USA Negro	135	45.9	0.9	22.8	16.3	4.4	0	7.4	1.5	0.7	0	5
23	223	41.2		20.2	22.4	5.4	...	8.1	2.7	0	0	5
24 USA white	766	2.2	20.9	33.8	14.9	13.9	0.1	12.5	0.9	0.5	0	5
25	7,317	2.2		53.5	15.0	12.9	...	14.7	1.1	0.6	0.01	5

Contributors: (a) Wiener, Alexander S., (b) Levine, Philip


References: [1] Matson, G. A., and C. L. Piper. 1947. *Am. J. Phys. Anthropol.* 5:357. [2] Matson, G. A., and H. J. Roberts. 1949. *Ibid.* 7:109. [3] Mourant, A. E. 1947. *Nature* 160:505. [4] Mourant, A. E. 1954. *The distribution of the human blood groups*. C. C. Thomas, Springfield, Ill. [5] Wiener, A. S. 1946. *Am. J. Clin. Pathol.* 16:477.

## 83. BLOOD COAGULATION THEORIES

### Part I. ACCORDING TO F. C. MONKHOUSE AND W. W. COON (1963)

**Synonymous terms for clotting factors:** PLASMA THROMBOPLASTIN COMPONENT = Christmas factor, platelet cofactor-2, autoprothrombin-2; ANTIHEMOPHILIC FACTOR = antihemophilic globulin, thromboplastinogen; FACTOR V = Ac-globulin, labile factor, proaccelerin; FACTOR VII = stable factor, autoprothrombin-1, proconvertin, cothromboplastin; PLASMINOGEN = profibrinolysin; PLASMIN = fibrinolysin.

**Symbols:** —————> gives rise to; - - - - -> acts on; =====> inhibits or destroys; [ ] present in blood.



<sup>1/1</sup> Other trace proteins affect the development of thromboplastin activity in plasma. Factor XII (Hageman factor) is sensitive to surface activation; its deficiency results in prolonged clotting time but no increase in bleeding time. The exact point of action of Factor X (Stuart-Prower factor) has not yet been determined.

**Contributors:** (a) Monkhouse, Frank C., (b) Coon, William W.

**References:** [1] Astrup, T. 1956. Blood 11:781. [2] Lorand, L. 1954. Physiol. Rev. 34:742. [3] Sherry, S., W. Troll, and H. Glueck. 1954. Ibid. 34:736.


*continued*

## 83. BLOOD COAGULATION THEORIES

### Part II. ACCORDING TO P. A. OWREN (1963)

#### System I: Intrinsic Blood Coagulation

Factor VII (proconvertin) does not take part in the intrinsic blood coagulation system.



/1/ CONTACT activates INACTIVE FACTOR XII (Hageman factor) to ACTIVE FACTOR XII. /2/ ACTIVE FACTOR XII activates INACTIVE FACTOR XI (plasma thromboplastin antecedent, PTA) to ACTIVE FACTOR XI. /3/ ACTIVE FACTOR XI, FACTOR IX (antihemophilia B factor, Christmas factor), FACTOR VIII (antihemophilia A factor, antihemophilic globulin), and CALCIUM interact to form BLOOD CONVERTIN. /4/ BLOOD CONVERTIN activates INACTIVE FACTOR X (Stuart-Prower factor) to ACTIVE FACTOR X. /5/ ACTIVE FACTOR X and CEPHALIN in the presence of calcium bring about a minimal conversion of PROTHROMBIN to THROMBIN. /6/ This initially formed THROMBIN starts the accelerator system, i.e., the conversion of INACTIVE FACTOR V (proaccelerin) to ACTIVE FACTOR V (accelerin). /7/ ACTIVE FACTOR X, ACTIVE FACTOR V, and CEPHALIN interact in the presence of calcium to form PROTHROMBINASE. /8/ PROTHROMBINASE produces rapid conversion of PROTHROMBIN to THROMBIN. /9/ THROMBIN is now formed in sufficient quantities to convert FIBRINOGEN to FIBRIN.

*continued*




## 83. BLOOD COAGULATION THORIES

### Part II. ACCORDING TO P. A. OWREN (1963)

#### System II: Extrinsic (tissue-blood) Coagulation

Platelets and antihemophilic factors do not take part in the extrinsic blood coagulation system.



/1/ CONTACT (mediated through activation of factors XII and XI) and/or TISSUE THROMBOPLASTIN activates INACTIVE FACTOR VII (proconvertin) to ACTIVE FACTOR VII. /2/ TISSUE THROMBOPLASTIN (liberated by tissue injury), ACTIVE FACTOR VII, and CALCIUM interact to form TISSUE CONVERTIN. /3/ TISSUE CONVERTIN activates INACTIVE FACTOR X (Stuart-Prower factor) to ACTIVE FACTOR X. /4/ ACTIVE FACTOR X and CEPHALIN in the presence of calcium bring about a minimal conversion of PROTHROMBIN to THROMBIN. /5/ This initially formed THROMBIN starts the accelerator system, i.e., the conversion of INACTIVE FACTOR V (proaccelerin) to ACTIVE FACTOR V. /6/ ACTIVE FACTOR X, ACTIVE FACTOR V, and CEPHALIN interact in the presence of calcium to form PROTHROMBINASE. /7/ PROTHROMBINASE produces rapid conversion of PROTHROMBIN to THROMBIN. /8/ THROMBIN is now formed in sufficient quantities to convert FIBRINOGEN to FIBRIN.


*Contributor:* Owren, Paul A.

*References:* [1] Hjort, P. F. 1957. Scand. J. Clin. Lab. Invest., Suppl. 27. [2] Owren, P. A. 1947. Acta Med. Scand., Suppl. 194. [3] Waaler, B. A. 1959. Scand. J. Clin. Lab. Invest., Suppl. 37.

*continued*

## 83. BLOOD COAGULATION THEORIES

Part III. ACCORDING TO A. J. QUICK (1963)



/1/ A plasma constituent, tentatively named CONTACT FACTOR, after activation by thrombin or by contact with glass, reacts with a PLATELET FACTOR to form ERYTHROCYTIN. /2/ FACTOR IX (plasma thromboplastin component) is inactive in plasma but is activated during coagulation. The activator mechanism is not known. /3/ ERYTHROCYTIN, FACTOR VIII (thromboplastinogen), and ACTIVATED FACTOR IX (activated plasma thromboplastin component) interact to form INTRINSIC THROMBOPLASTIN. /4/ INTRINSIC THROMBOPLASTIN, CALCIUM, FACTOR V (labile factor), and possibly FACTOR X (Stuart-Prower factor) interact with PROTHROMBIN to form THROMBIN. In human blood, a large fraction of prothrombin is inactive but becomes activated during coagulation in glass. /5/ When TISSUE THROMBOPLASTIN is utilized, FACTOR VII (stable factor) is required in addition to CALCIUM and FACTOR V. /6/ THROMBIN acts enzymatically on FIBRINOGEN to convert it to FIBRIN. /7/ The prompt removal of THROMBIN by adsorption on FIBRIN holds in check the autocatalytic reaction mediated through the activation of THROMBIN on the CONTACT FACTOR.

*Contributor:* Quick, Armand J.


*References:* [1] Quick, A. J. 1943. Am. J. Physiol. 140:212. [2] Quick, A. J. 1947. Am. J. Med. Sci. 214:272. [3] Quick, A. J. 1958. Thromb. Diath. Haemorrhag. 2:226. [4] Quick, A. J. 1960. Am. J. Med. Sci. 239:51. [5] Quick, A. J. 1961. Ann. Internal Med. 55:201. [6] Quick, A. J., and J. E. Favre-Gilly. 1949. Am. J. Physiol. 158:387. [7] Quick, A. J., and C. V. Hussey. 1955. Brit. Med. J. 1:934.

*continued*

## 83. BLOOD COAGULATION THEORIES

### Part IV. ACCORDING TO W. H. SEEGERS (1963)

Prothrombin is found in the blood and may become activated in the presence of one or more procoagulants. Since prothrombin itself contains all the necessary material for the formation of thrombin, purified prothrombin can be activated to thrombin and autoprothrombin C by placing it in 25% sodium citrate solution. Consequently the activators of prothrombin are catalysts and do not enter into stoichiometric combination with prothrombin to form thrombin. Anticoagulants inhibit the activation. Ordinarily the procoagulants and anticoagulants are present in balanced proportion. This balance is readily disturbed by the procoagulants from injured tissues. Contact with foreign surfaces also promotes prothrombin activation and platelet disintegration. With certain combinations of procoagulants, prothrombin is only partially activated, and these derivatives of prothrombin themselves accelerate the conversion of prothrombin to thrombin. Thrombin functions as activator of prothrombin, and a second enzyme from prothrombin, called autoprothrombin C, functions similarly. Prothrombin activation is primarily by autocatalysis. Thrombin also functions with accelerator systems, such as plasma Ac-globulin which becomes serum Ac-globulin, and it further supports the dissolution of platelets. Plasma antithrombin eventually destroys thrombin activity. By proteolysis, thrombin splits peptides from fibrinogen and acts as a polymerase in the polymerization of the activated fibrinogen. In the presence of calcium ions and fibrin stabilizing factor, the fibrin of a normal clot forms. Vitamin K is needed for the metabolic production of prothrombin and its derivatives, whereas dicumarol may interfere with normal prothrombin metabolism.



<sup>1/1</sup> Related to prothrombin production.


*Contributor:* Seegers, Walter H.

*References:* [1] McClaughry, R. I., and J. L. Fahey. 1950. Blood 5:421. [2] Milstone, J. H. 1948. Proc. Soc. Exptl. Biol. 68:225. [3] Seegers, W. H. 1950. Circulation 1:2. [4] Seegers, W. H. In J. B. Sumner and K. Myrbäck, ed. 1951. The enzymes. Academic Press, New York. v. 1, pt. 2, p. 1006. [5] Seegers, W. H., et al. 1963. Can. J. Biochem. Physiol. 41:1047. [6] Ware, A. G., J. L. Fahey, and W. H. Seegers. 1948. Am. J. Physiol. 154:140. [7] Ware, A. G., and W. H. Seegers. 1948. Ibid. 152:567.

*continued*

83. BLOOD COAGULATION THEORIES

Part V. ACCORDING TO L. M. TOCANTINS (1960)



/1/ CONTACT of BLOOD with certain SURFACES (damaged blood vessel endothelium, glass) initiates the first changes in the inception of clotting. Blood PLATELETS adhere to the surface and to each other, swell and DISINTEGRATE. /2/ PLATELET DISINTEGRATION releases, among other substances, PLATELET FACTOR 3 (cephalin-like factor) and PLATELET FACTOR 2 (a fibrinolytic factor). /3/ Conjugation of PLATELET FACTOR 3 with FACTOR IX (platelet cofactor in plasma, plasma thromboplastin component) leads, with the aid of FACTOR X (Stuart-Prower factor), to formation of PLASMA THROMBOPLASTIN. /4/ Plasma ANTITHROMBOPLASTIN slows or blocks formation of THROMBOPLASTIN and, less effectively, offsets the action of formed THROMBOPLASTIN. Antihemophilic globulin is considered to represent various stages of development of plasma thromboplastin, or various degrees of conjugation of the platelet lipid and its plasma cofactor. /5/ Plasma ANTITHROMBOPLASTIN is probably a lipid in conjugation with FACTOR IX. /6/ THROMBOPLASTIN (from TISSUES, or generated in PLASMA) brings about, with the aid of FACTOR V (Ac-globulin) and FACTOR VII (convertin), a minimal amount of conversion of PROTHROMBIN to THROMBIN. /7/ This initial THROMBIN activates further transformation of PROTHROMBIN to THROMBIN, with the help of FACTOR V. /8/ Some of the THROMBIN may be inactivated by ANTI-THROMBIN. /9/ The THROMBIN that escapes such inactivation acts, with the aid of PLATELET FACTOR 2, to convert FIBRINOGEN to FIBRIN. Some of the THROMBIN is removed from the plasma by adsorption on FIBRIN. /10/ Adhesion of platelets to FIBRIN probably causes further PLATELET DISINTEGRATION.

*Contributor:* Tocantins, Leandro M.

*References:* [1] Silver, M. J., D. L. Turner, and L. M. Tocantins. In L. M. Tocantins, ed. 1959. Progress in hematology. Grune and Stratton, New York. v. 2. [2] Tocantins, L. M. 1943. Am. J. Physiol. 139:265. [3] Tocantins, L. M. 1944. Proc. Soc. Exptl. Biol. Med. 55:291. [4] Tocantins, L. M. 1944. Ibid. 57:211. [5] Tocantins, L. M. 1946. Blood 1:56. [6] Tocantins, L. M. 1949. Surg. Clin. North Am. 29:1835. [7] Tocantins, L. M. 1954. Blood 9:281. [8] Tocantins, L. M. 1955. The coagulation of blood: methods of study. Grune and Stratton, New York. [9] Tocantins, L. M., R. T. Carroll, and R. R. Holburn. 1951. Blood 6:720. [10] Tocantins, L. M., R. T. Carroll, and T. J. MacBride. 1948. Proc. Soc. Exptl. Biol. Med. 68:110. [11] Tocantins, L. M., R. R. Holburn, and R. T. Carroll. 1951. Ibid. 76:623.



## 84. ACID-BASE BALANCE

For additional information, consult reference 2, Part I. *Abbreviations:*  $pK'_1$  = first dissociation constant;  $f_{CO_2}$  = carbon dioxide factor.

### Part I. ACID-BASE VALUES: MAN

**Blood** (column C): C = cutaneous; A = arterial; V = venous. Values in parentheses are ranges, estimate "b" (cf. Introduction).

Subjects		Blood	pH at 37°C	Hemo- globin <sup>1</sup> mM/L	CO <sub>2</sub> Content, mM/L		CO <sub>2</sub> Pressure <sup>2</sup> mm Hg	Buffer- Base <sup>3</sup> mEq/L	Ref- er- ence	
Age	No. and Sex				Whole Blood	Plasma				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
1	4-6 da <sup>4</sup>	18♂♀	C	7.38(7.26-7.50)	10 (8.6-11.4)	15.3(11-20)	19(14.5-23.5)	31.1(23-40)	42.3(36-49)	6
2	3-15 yr	20♂♀	A	7.38	7.6	21.2	25	39.8	45.7	3,4
3	18-28 yr <sup>5</sup>	8♂	C	7.37(7.32-7.42)	9.3 <sup>6</sup>	21.9(21.1-22.7)	26.8	43.5(37.9-49.1)	48.1	5
4		7♀	C	7.39(7.36-7.42)	7.6 <sup>6</sup>	21.9(19.9-23.9)	26	40.4(37.5-43.3)	46.9	
5	24-43 yr	3♀	A	7.43	8.1 <sup>7</sup>		25.9	38.4	47	1
6	Adult	259♂	A	7.39(7.34-7.44)	8.9 (7.7-10.3)	22.2(20-24)	26.9(25-29)	41.6(35-47)	48.4(46-52)	2
7		118♂	V	7.35(7.28-7.42)	8.9 <sup>6</sup>	24.4(21-28)	29.6(26-33)	50	48.4 <sup>6</sup>	

<sup>1/1</sup> Hemoglobin concentration assumed to be 20 mM/L erythrocytes; 1 mM (single Fe-atom structure, molecular weight 16,500) combines with 22.4 ml of oxygen STP when saturated. <sup>2/2</sup> CO<sub>2</sub> pressure calculated from adjusted pH and plasma CO<sub>2</sub> content by the Henderson-Hasselbalch equation ( $pH - pK'_1 = \log \frac{CO_2 - 0.0314 CO_2 \text{ pressure}}{0.0314 CO_2 \text{ pressure}}$ ).

<sup>3/3</sup> Calculated from pH, bicarbonate, and other blood buffers, as described in reference 7. <sup>4/4</sup> Formula-fed infants; different results were obtained with breast-fed infants. <sup>5/5</sup> Observations made in the morning on fasting, seated subjects, after one-half hour rest. <sup>6/6</sup> Calculated from hematocrit value. <sup>7/7</sup> Average value for non-pregnant females [2]. <sup>8/8</sup> Assumed equal to value in arterial blood.

*Contributors:* Singer, Richard B., and Hastings, A. Baird.

*References:* [1] Alexander, J. K., et al. 1955. J. Clin. Invest. 34:511. [2] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [3] Kennedy, C., and L. Sokoloff. 1957. J. Clin. Invest. 36:1130. [4] Robinson, S. 1938. Arbeitsphysiologie 10:251. [5] Shock, N. W., and A. B. Hastings. 1934. J. Biol. Chem. 104:585. [6] Singer, R. B. Unpublished. New England Mutual Life Insurance Co., Boston, 1958. [7] Singer, R. B., and A. B. Hastings. 1948. Medicine 27:223.

### Part II. ACID-BASE VALUES: VERTEBRATES

pH adjusted to body temperature by applying correction of -0.015 per °C temperature difference. CO<sub>2</sub> pressure calculated by means of Henderson-Hasselbalch equation; value of  $pK'_1$  increases 0.005 per °C decrease in temperature, and  $f_{CO_2}$  is assumed to increase proportionately as it does in pure water. The following values for  $pK'_1$  and  $f_{CO_2}$  were used for body temperatures other than 38°C: 5°C, 6.26 and 0.0864; 10°C, 6.24 and 0.0697; 20°C, 6.19 and 0.0508; 26°C, 6.16 and 0.0434; 34°C, 6.12 and 0.0357; 40°C, 6.09 and 0.0313; 42°C, 6.08 and 0.0303. **Blood** (column B): A = arterial; M = mixed arterial and venous; V = venous; H = heart. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Species (Common Name)	Body Temp. °C (Blood)	Whole Blood pH	Erythro- cytes		Plasma						Reference	
			Hb mM/L	Vol %	CO <sub>2</sub> Content mM/L	CO <sub>2</sub> Pres- sure mm Hg	Na <sup>+</sup> mEq/L	Cl <sup>-</sup> mEq/L	H <sub>2</sub> O g/L	Pro- tein g/L		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	
Mammalia and Aves <sup>1</sup>												
1	<i>Homo sapiens</i> (man), adult♂	37 (A)	7.39 (7.33-7.45) <sup>b</sup>	9.0	45	27.0 (25-29) <sup>b</sup>	42 (36-47) <sup>b</sup>	138 (132-144) <sup>b</sup>	102 (97-108) <sup>b</sup>	940	68	B,C,F-I,2; D,E,J,K,15
2	<i>Bos taurus</i> (cat- tle), ♀	38.5 (A)	7.38 (7.27-7.49) <sup>b</sup>	7.0	40	31.0 (29-33) <sup>b</sup>	50	142 (132-152) <sup>b</sup>	104 (97-111) <sup>b</sup>	930 <sup>2</sup>	83	B,16;C,F-I, 26;D,E,K, 2;J,28

<sup>1/1</sup> Homiothermic body temperature relatively independent of environmental temperature except in hibernating animals. <sup>2/2</sup> Calculated data.

*continued*

## 84. ACID-BASE BALANCE

### Part II. ACID-BASE VALUES: VERTEBRATES

Species (Common Name)	Body Temp. °C (Blood)	Whole Blood pH	Erythro- cytes		Plasma						Reference	
			Hb mM/L	Vol %	CO <sub>2</sub> Content mM/L	CO <sub>2</sub> Pres- sure mm Hg	Na <sup>+</sup> mEq/L	Cl <sup>-</sup> mEq/L	H <sub>2</sub> O g/L	Pro- tein g/L		
												(D)
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	
Mammalia and Aves <sup>1</sup>												
3	<i>Canis familiaris</i> (dog)	38.9 (A)	7.36 (7.31-7.42)	9.0	46	21.4 (17-24)	38	147 (140-154) <sup>b</sup>	114 (108-119) <sup>b</sup>	941	67	B,14;C,F,G, 4,14,19;D, E,K,2;H-J, 27
4	<i>Cavia</i> sp. (guinea pig)	38.6 (H; A)	7.35 (7.17-7.55)	8.7	42	22.0 (16-26) <sup>3</sup>	40 (19-59)	141 (138-144)	104 (100-108)	954 <sup>2</sup>	47	B,C,F,G,21; D,E,K,2;H, I,22;J,28
5	<i>Equus caballus</i> (horse)	37.8 (V)	(7.20-7.55)	6.8	33	28.1 (24-32)	47 <sup>2</sup>	135	96	931	68	B,C,16;D,E, K,2;F,23; G,28;H-J, 30
6	<i>Felis catus</i> (cat)	38.6 (M)	7.35 (7.24-7.40)			20.4 (17-24) <sup>b</sup>	36	153 (150-156) <sup>b</sup>	120 (117-123) <sup>b</sup>	941		B,16,34; C-J,34
7	Anesthetized	38.6 (V)	7.28 (7.18-7.35)	6.8	40	21.8 (19-25)	45 (34-52)		108 (105-111)	942	76	B,16,32;C, F-J,32;D, E,K,2
8	<i>Gallus domesticus</i> (chicken)	41.7 (V)	7.54 (7.45-7.63)	6.8	32	23.0 (21-26)	26 <sup>3</sup>	154 (148-161)	117 (109-120)	960 <sup>2</sup>	36	B,16;C,24; D,E,K,2;F, 1;G,J,28; H,I,22
9	<i>Mesocricetus au- ratus</i> (golden hamster) Anesthetized	38 (H; V)	7.39 (7.37-7.44)	8.4	46	37.3 (35-39)	59 (54-61)	144 (140-151)	106 (103-108)	945 <sup>2</sup>		B,C,F,G,25; D,E,2;H,I, 22;J,28
10	Hibernating	5 (H; V)	7.44 (7.34-7.56)			42.4 (35-50)	32 (26-42)					25
11	<i>Oryctolagus cuni- culus</i> (European rabbit)	39.4 (A)	7.35 (7.21-7.57)	7.2		22.8 (13-33) <sup>3</sup>	40 (22-51)	140 (139-142)	102 (99-105)	944 <sup>2</sup>		B,16,21;C, F,G,21,35; D,H,I,35; J,28
12	<i>Ovis aries</i> (sheep)	39.1 (V)	7.44 (7.32-7.54)	7.6	32	26.2 (21-28)	38	153 (146-161)	103 (98-109)	947 <sup>2</sup>	57	B,16;C,F-I, 9;D,E,K,2; J,28
13	<i>Rattus</i> sp. (rat)	38.2 (A)	7.35 (7.26-7.44) <sup>b</sup>	9.0	46	24.0 (20-28) <sup>b</sup>	42	144 (135-155) <sup>b</sup>	104 (99-112) <sup>b</sup>	946	60	B,5,6,21;C, F-I,5,6;D, E,K,2;J,6
Reptilia, Pisces, and Chondrichthyes <sup>4</sup>												
14	<i>Alligator missis- sippiensis</i>	34 (H; M)	7.43	5.4		19.8	29		105	954 <sup>2</sup>	46	B-I,K,12;J, 28
15	(American alli- gator) <sup>5</sup>	26 (H; M)	7.48 (7.33-7.62)	4.3	22	23.5 (15-40)	38	141 (136-143)	112 (106-117)	952 <sup>2</sup>	50	B-I,K,7;J, 28
16		5 (H; M)	7.74	4.2	25	36.1	15		110	958 <sup>2</sup>	41	B-I,K,12;J, 28
17	<i>Anolis carolinensis</i> (American "cha- meleon")	26 (H; M)	7.26 (6.93-7.63)	4.2	28	15.4 (10-22)	27	157 (139-186)	127 (113-133)	958 <sup>2</sup>	41	B-I,K,10;J, 28

/1/ Homiothermic body temperature relatively independent of environmental temperature except in hibernating animals. /2/ Calculated data. /3/ Calculated from whole blood CO<sub>2</sub> content, pH, and hemoglobin, by means of nomogram of Singer and Hastings [29]. /4/ Poikilothermic body temperature dependent on environmental temperature. When temperature is decreased, pH and CO<sub>2</sub> solubility coefficient increase, and the O<sub>2</sub> dissociation curve is shifted to the left. /5/ The alligator shows a marked variation among individuals and in the same individual at different seasons, and a prolonged and extreme "alkaline tide" following meals [8].

continued

## 84. ACID-BASE BALANCE

### Part II. ACID-BASE VALUES: VERTEBRATES

Species (Common Name)	Body Temp. °C (Blood)	Whole Blood pH	Erythro- cytes		Plasma						Reference	
			Hb mM/L	Vol %	CO <sub>2</sub> Content mM/L	CO <sub>2</sub> Pres- sure mm Hg	Na <sup>+</sup> mEq/L	Cl <sup>-</sup> mEq/L	H <sub>2</sub> O g/L	Pro- tein g/L		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	
Reptilia, Pisces, and Chondrichthyes <sup>a</sup>												
18	<i>Cyprinus carpio</i> (carp)	20 <sup>b</sup> (H; V)	7.39 (7.33-7.45)	6.4	31	17.7 (14-22)	22 <sup>b</sup>	130 (126-137)	107 (96-121)	957 <sup>a</sup>	42	B,G,J,28;C, F,3;D,E,H, K,18;L,31
19		15 (V)			39	13.5 <sup>b</sup>	8.5 <sup>b</sup>		147	951		17
20	<i>Pseudemys scrip- ta elegans</i> (red- eared turtle)	26 (H)	7.65 (7.50-8.10)		25	24.4 (18-32)		125 (114-135)	92 (80-100)			33
21	<i>Raja</i> sp. (skate)	10.4 (A)	7.82	2.7	20	3.5	1.3	254 (219-289)	255 (230-285)	967 <sup>a</sup>	27	B-G,K,13;H, I,20;J,28
22	<i>Salmo gairdneri</i> (rainbow trout)	15 (V)			35	9.5 <sup>b</sup>	9.0 <sup>b</sup>		140	955		17
23	<i>Thamnophis</i> sp. (garter snake)	26 (H)	7.25 (7.12-7.50)		28	6.6 (3-16)		156 (143-169)	130 (122-143)		42	11

/s/ Calculated data. /4/ Poikilothermic body temperature dependent on environmental temperature. When temperature is decreased, pH and CO<sub>2</sub> solubility coefficient increase, and the O<sub>2</sub> dissociation curve is shifted to the left.  
/e/ Value for whole blood.


Contributors: (a) Singer, Richard B., (b) Irvin, J. Logan, (c) Hernandez, Thomas

References: [1] Ackerson, C. W., M. J. Blish, and F. E. Mussehl. 1925. *J. Biol. Chem.* 63:75. [2] Albritton, E. C., ed. 1952. *Standard values in blood*. W. B. Saunders, Philadelphia. [3] Auvergnat, R., and M. Lecondat. 1942. *Compt. Rend.* 215:92. [4] Bennett, M. A. 1926. *J. Biol. Chem.* 69:675. [5] Cooke, R. E., F. R. Coughlin, Jr., and W. E. Segar. 1952. *J. Clin. Invest.* 31:1006. [6] Cotlove, E., et al. 1951. *Am. J. Physiol.* 167:665. [7] Coulson, R. A., and T. Hernandez. Unpublished. Louisiana State Univ., New Orleans, 1963. [8] Coulson, R. A., T. Hernandez, and H. C. Dessauer. 1950. *Proc. Soc. Exptl. Biol. Med.* 74:866. [9] Denton, D. A., et al. 1951. *Acta Med. Scand., Suppl.* 261. [10] Dessauer, H. C. 1952. *Proc. Soc. Exptl. Biol. Med.* 80:742. [11] Dessauer, H. C., and W. Fox. Unpublished. Louisiana State Univ., New Orleans, 1955. [12] Dill, D. B., and H. T. Edwards. 1935. *J. Cellular Comp. Physiol.* 6:243. [13] Dill, D. B., H. T. Edwards, and M. Florkin. 1932. *Biol. Bull.* 62:23. [14] Dill, D. B., et al. 1932. *J. Biol. Chem.* 95:143. [15] Dill, D. B., et al. 1940. *Ibid.* 136:449. [16] Dukes, H. H. 1955. *The physiology of domestic animals*. Ed. 7. Comstock, Ithaca. [17] Ferguson, J. K. W., and E. C. Black. 1941. *Biol. Bull.* 80:139. [18] Field, J. B., C. A. Elvehjem, and C. Juday. 1943. *J. Biol. Chem.* 148:261. [19] Harkins, H. N., and A. B. Hastings. 1931. *Ibid.* 90:565. [20] Hartman, F. A., et al. 1941. *Physiol. Zool.* 14:476. [21] Hawkins, J. A. 1924. *J. Biol. Chem.* 61:147. [22] Hernandez, T., and R. A. Coulson. Unpublished. Louisiana State Univ., New Orleans, 1955. [23] Ichiji, N. 1922. *J. Japan. Soc. Vet. Sci.* 1:76. [24] Johnson, E. P., and W. B. Bell. 1936. *J. Infect. Diseases* 58:342. [25] Lyman, C. P., and A. B. Hastings. 1951. *Am. J. Physiol.* 167:633. [26] McSherry, B. J., and I. Griner. 1954. *Am. J. Vet. Res.* 15:509. [27] Mellors, R. C., E. Muntwyler, and F. R. Mantz. 1942. *J. Biol. Chem.* 144:773. [28] Singer, R. B. Unpublished. New England Mutual Life Insurance Co., Boston, 1953. [29] Singer, R. B., and A. B. Hastings. 1948. *Medicine* 27:223. [30] Van Slyke, D. D., et al. 1925. *J. Biol. Chem.* 65:701. [31] Vars, H. M. 1934. *Ibid.* 105:135. [32] Wallace, W. M., and A. B. Hastings. 1942. *Ibid.* 144:637. [33] Williams, J. K. Unpublished, 1959. [34] Yannet, H. 1940. *J. Biol. Chem.* 136:265. [35] Young, I. M. 1952. *Am. J. Physiol.* 170:434.

continued

### Part III. NORMAL IONIC PATTERNS, ARTERIAL BLOOD: MAN

Values are for the adult male and are based on the literature.  $X^-$  = undetermined anion residue. " $HbO_2^-$ " includes other erythrocyte buffer anions, such as organic phosphate.  $B^+$  = mEq total cation ( $Na^+$ ,  $K^+$ , etc.) in 1 liter of blood, on the basis of a hematocrit value of 45%. Buffer base = the appropriate fraction of total cations and its equivalent of total anions (labile fraction), i.e., proteinate, bicarbonate, oxyhemoglobinate, organic phosphate, and other erythrocyte buffer anions.  $CO_2$  partial pressure or tension for plasma, erythrocytes, or whole blood = 41 mm Hg.



Contributor: Singer, Richard B.

### Part IV. CLASSIFICATION OF ACID-BASE DISTURBANCES: MAN

Ranges for acid-base variables, as reported in the literature or inferred from related observations, are for adult arterial or cutaneous blood. See also normal values, Part I. Limits given are approximate. Boldface type (columns B and C) indicates the best index for existence of the particular condition.

Condition	Buffer Base <sup>1</sup> mEq/L	$CO_2$ Pressure mm Hg	Bicarbonate <sup>2</sup> mEq/L	pH at 37°C
(A)	(B)	(C)	(D)	(E)
1 Normal (arterial or cutaneous blood)	46-52	35-45	24-28	7.35-7.45
2 Metabolic acidosis (acid excess or base deficit)	<b>Always low</b> 20-46	<b>Usually low</b> 15-35	Usually low 4-24	Usually low 6.8-7.35

/1/ Buffer base for whole blood of normal hemoglobin concentration = 15 g/100 ml. A decrease in buffer base of whole blood is almost always accompanied by a decrease in plasma or extracellular  $Na^+$  relative to  $Cl^- + X^-$ , e.g., decrease in plasma  $Na^+$ , increase in plasma  $Cl^-$  or plasma  $X^-$ , or any appropriate combination. An increase in buffer base of whole blood is accompanied by an increase in  $Na^+$  relative to  $Cl^- + X^-$ , e.g., increase in plasma  $Na^+$ , decrease in plasma  $Cl^-$ , or any appropriate combination. See normal values in diagram, Part III. /2/ Comprises about 90-98% of total carbon dioxide in plasma; average, 95%.

continued



## 84. ACID-BASE BALANCE

### Part IV. CLASSIFICATION OF ACID-BASE DISTURBANCES: MAN

	Condition	Buffer Base <sup>1</sup> mEq/L	CO <sub>2</sub> Pressure mm Hg	Bicarbonate <sup>2</sup> mEq/L	pH at 37°C
	(A)	(B)	(C)	(D)	(E)
3	Respiratory acidosis (H <sub>2</sub> CO <sub>3</sub> excess)	Normal or high 46-70	Always high 45-100+	Usually high 28-45	Usually low 7.0-7.35
4	Metabolic alkalosis (base excess or acid deficit)	Always high 52-75	Normal or high 35-55	Usually high 28-50	Usually high 7.45-7.65
5	Respiratory alkalosis (H <sub>2</sub> CO <sub>3</sub> deficit)	Normal or low 40-52	Always low 10-35	Usually low 15-24	Usually high 7.45-7.70
6	Mixed acidosis (combination of lines 2 and 3)	Always low 25-45	Always high 45-100	Variable 10-35	Always low 6.8-7.35
7	Mixed alkalosis (combination of lines 4 and 5)	Always high 52-70	Always low 15-35	Variable 20-45	Always high 7.5-7.7
8	Mixed hypercapnia (combination of lines 3 and 4)	Always high 52-75	Always high 45-100	Always high 30-50	Variable 7.3-7.6
9	Mixed hypocapnia (combination of lines 2 and 5)	Always low 20-46	Always low 10-35	Always low 4-22	Variable 7.0-7.6

<sup>1/</sup> Buffer base for whole blood of normal hemoglobin concentration = 15 g/100 ml. A decrease in buffer base of whole blood is almost always accompanied by a decrease in plasma or extracellular Na<sup>+</sup> relative to Cl<sup>-</sup> + X<sup>-</sup>, e.g., decrease in plasma Na<sup>+</sup>, increase in plasma Cl<sup>-</sup> or plasma X<sup>-</sup>, or any appropriate combination. An increase in buffer base of whole blood is accompanied by an increase in Na<sup>+</sup> relative to Cl<sup>-</sup> + X<sup>-</sup>, e.g., increase in plasma Na<sup>+</sup>, decrease in plasma Cl<sup>-</sup>, or any appropriate combination. See normal values in diagram, Part III. <sup>2/</sup> Comprises about 90-98% of total carbon dioxide in plasma; average, 95%.

Contributor: Singer, Richard B.

References: [1] Peters, J. P., and D. D. Van Slyke. 1946. Quantitative clinical chemistry. Ed. 2. Williams and Wilkins, Baltimore, v. 1. [2] Singer, R. B., and A. B. Hastings. 1948. Medicine 27:223.

## 85. BLOOD VOLUMES

For additional information, consult reference 1, Part I.

### Part I. VERTEBRATES

For a summary of blood methods and interpretations, consult reference 11. Subjects were normal, adult animals. Plasma and erythrocyte volumes were obtained by various dilution methods (diluent or tagging substance is given in the pertinent method column). Venous hematocrit values were obtained by centrifuging the blood sample (3,000 rpm, 30 minutes, 18 cm radius). In most instances, whole blood volume was calculated from other values in the same study; where a tagging substance is given in column H, blood volume was determined directly by dilution of the tagged erythrocytes in whole blood, on the assumption that the erythrocyte concentration in the sampled blood represented the total body erythrocyte concentration. Method (columns C, E, H): PV = plasma volume; EV = erythrocyte volume; VH = venous hematocrit; BV = whole blood volume. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Species (Common Name)	Sub- jects	Plasma Volume		Erythrocyte Volume		Venous Hematocrit % cells	Whole Blood Volume		Ref- er- ence
			Method	ml/kg body wt	Method	ml/kg body wt		Method	ml/kg body wt	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Mammalia										
1	<i>Homo sapiens</i> (man)	25	T-1824	45.4 (32.3-54.2)	P <sup>32</sup>	30.1 (21.1-44.1)		PV + EV	75.5 (60.9-95.8)	6
2		30♂	T-1824	41.1	P <sup>32</sup>	28.0 <sup>c</sup>		PV + EV	69.1	25
3		40♂	T-1824	47.9 (39.2-62.5)	Fe <sup>55</sup> or Fe <sup>59</sup>	29.8 (21.5-36.3)		PV + EV	77.7 (63.8-97.0)	10
4		20♂	T-1824	45.7 (35.8-56.5)			44.1 (39.3-49.4)	PV 100 - VH × 100	81.6 (65.4-95.2)	30

<sup>1/</sup> Corrected for trapped plasma by factor of 0.96.

continued

# Contrails

## 85. BLOOD VOLUMES

### Part I. VERTEBRATES

Species (Common Name)	Sub- jects	Plasma Volume		Erythrocyte Volume		Venous Hematocrit % cells	Whole Blood Volume		Ref- erence	
		Method	ml/kg body wt	Method	ml/kg body wt		Method	ml/kg body wt		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
Mammalia										
5	<i>Homo sapiens</i> (man)	30♀	T-1824	40.5	P <sup>32</sup>	21.6 <sup>1</sup>		PV + EV	63.1	25
6		20♀	T-1824	44.7 (37.2-58.5)			39.8 (37.1-41.9)	$\frac{PV}{100 - VH} \times 100$	74.3 (63.0-97.5)	30
7		35♀	T-1824	48.2 (37.7-57.2)			39.2 (34.5-43.8) <sup>2</sup>	$\frac{PV}{100 - VH} \times 100$	79.5 (65.0-99.8)	28
8	<i>Bos taurus</i> (cattle)	10♀	T-1824	38.8 (36.3-40.6)			32.4 (30.3-34.9) <sup>2</sup>	$\frac{PV}{100 - VH} \times 100$	57.4 (52.4-60.6)	23
9	<i>Camelus dro- medarius</i> (Arabian camel)	19♂	T-1824	59 (47-70)			29	$\frac{PV}{100 - VH} \times 100$	83 (68-100)	3
10	<i>Canis famili- aris</i> (dog)	11	T-1824	55.2 (43.7-73.0)	P <sup>32</sup>	39.0 (28.0-55.0)	44 (35-54) <sup>1</sup>	PV + EV	94.1 (76.5-107.3)	27
11	<i>Capra hircus</i> (goat)	20	T-1824	55.9 (42.6-75.1)	Cr <sup>51a</sup>	14.7 (9.7-19.3)	24.3 (18.5-30.8) <sup>4</sup>	PV + EV	70.5 (56.8-89.4)	18
12	<i>Cavia sp.</i> (guinea pig)	13	I <sup>131</sup> rab- bit glob- ulin	39.4 (35.1-48.4) <sup>5</sup>				$\frac{PV}{100 - VH} \times 100$	75.3 (67.0-92.4)	19
13	<i>Didelphis sp.</i> (opossum)	10 <sup>b</sup>	T-1824	37.8 (29.6-52.2)	P <sup>32</sup>	19.2 (14.2-29.2)		PV + EV	57.0 (44.5-69.8)	7
14	<i>Equus cabal- lus</i> (horse)									
14	Light wt	6	BV - EV	61.9 (45.5-79.1)	P <sup>32</sup>	47.1 (39.6-57.5)	43.3(37-56)	$\frac{EV}{VH} \times 100$	109.6 (94.3-136.0)	16
15	Heavy wt	4	BV - EV	43.2 (30.6-64.1)	P <sup>32</sup>	28.5 (23.1-37.6)	40.3 (37-46)	$\frac{EV}{VH} \times 100$	71.7 (56.7-101.7)	16
16	<i>Felis catus</i> (cat)	5♂ <sup>b</sup>	T-1824	40.7 (34.6-52.0)	Cr <sup>51a</sup>	14.8 (12.2-17.7) <sup>1</sup>		PV + EV	55.5 (47.3-65.7)	9
17	<i>Macaca mu- latta</i> (rhe- sus monkey)	15♂, 3♀	T-1824	36.4 (30.0-48.4)	P <sup>32</sup>	17.7 (14.3-20.0)	39.6 (35.6-42.8) <sup>1</sup>	PV + EV	54.1 (44.3-66.6)	12
18	<i>Mus sp.</i> (mouse)	11	T-1824	48.8	P <sup>32</sup>	29.0		PV + EV	77.8	32
19	<i>Myotis lucifu- gus</i> (little brown bat)	123 <sup>b</sup>	T-1824	65			49.3 <sup>7</sup>	$\frac{PV}{100 - VH} \times 100$	130	17
20	<i>Oryctolagus cuniculus</i> (European rabbit)	29	T-1824	38.8 (27.8-51.4)	P <sup>32</sup>	16.8 (13.7-25.5)		PV + EV	55.6 (44.0-70.0)	2
21		71			P <sup>32</sup>	17.5 (13.4-22.8)	35.2 (28.6-41.0) <sup>1</sup>	$\frac{EV \times 100}{0.858(VH) - 0.2}$	57.3 (47.8-69.5)	2
22	<i>Ovis aries</i> (sheep)	5	T-1824	46.7 (43.4-52.9)	Cr <sup>51a</sup>	19.7 (16.3-23.8)		PV + EV	66.4 (59.7-73.8)	14
23	<i>Rattus norve- gicus</i> (Nor- way rat)		T-1824	40.4 (36.3-45.3) <sup>8</sup>	P <sup>32</sup>	23.7 (18.4-26.0) <sup>8</sup>	50.3 (42.3-61.5) <sup>8,9</sup>	PV + EV	64.1 (57.5-69.9) <sup>8</sup>	31
24	<i>Sus scrofa</i> (swine)									
24	45 kg	4			P <sup>32</sup>	25.9 (20.2-29.0)	39.1 (30.3-43.1) <sup>9</sup>	$\frac{EV}{VH} \times 100$	65 (61-68)	13
25	50 kg	6 <sup>e</sup>	BV - EV	41.9	BV x $\frac{VH}{100}$	27.5		P <sup>32</sup> erythro- cytes	69.4	8
Aves										
26	<i>Anas platy- rhynchos domesticus</i> (Pekin duck)	42	I <sup>131</sup> hu- man se- rum albumin	65.5				$\frac{PV}{100 - VH} \times 100$	102	21
27		2♂			Cr <sup>51a</sup>	30	38.5			24
28		2♀			Cr <sup>51a</sup>	25	43.5			24

<sup>1</sup>/ Corrected for trapped plasma by factor of 0.96. <sup>2</sup>/ Corrected for trapped plasma by factor of 0.94. <sup>3</sup>/ Cr<sup>51</sup> as sodium chromate. <sup>4</sup>/ Corrected for trapped plasma by factor of 0.81. <sup>5</sup>/ Calculated from an average hematocrit of 47.6 obtained from 10 other guinea pigs. <sup>6</sup>/ Anesthetized. <sup>7</sup>/ Cardiac blood. <sup>8</sup>/ Blood samples taken from carotid artery or tail vein. <sup>9</sup>/ Corrected for trapped plasma by factor of 0.95.

continued

## 85. BLOOD VOLUMES

### Part I. VERTEBRATES

	Species (Common Name)	Sub- jects	Plasma Volume		Erythrocyte Volume		Venous Hematocrit % cells	Whole Blood Volume		Ref- er- ence
			Method	ml/kg body wt	Method	ml/kg body wt		Method	ml/kg body wt	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Aves										
29	<i>Columba livia</i> (street pi- geon)	6	T-1824	44	Consult ref. 4	49	52	Consult ref. 4	92	5
30	<i>Gallus domes- ticus</i> (chick- en)	110♂ <sup>11</sup> , 113♀ <sup>11</sup>	T-1824					$\frac{PV}{100 - VH} \times 100$	90	20
31		3♂ <sup>12</sup>	T-1824	31	Consult ref. 4	25	45	Consult ref. 4	56	5
32		4♀ <sup>12</sup>	T-1824	44	Consult ref. 4	19	30	Consult ref. 4	63	5
33	<i>Phasianus colchicus</i>	4♂	T-1824	45	Consult ref. 4	22	33	Consult ref. 4	67	5
34	(ring-necked pheasant)	2♀	T-1824	32	Consult ref. 4	16	34	Consult ref. 4	48	5
Reptilia and Amphibia										
35	<i>Alligator mississippi- ensis</i> (Amer- ican alligator)	16♂ <sup>6</sup> , 17♀ <sup>6</sup>	T <sup>131</sup>	60.1	Cr <sup>51</sup>	12.6	22.7	PV + EV	72.7	15
36	<i>Pseudemys scripta ele- gans</i> (red-eared turtle)	26 <sup>13</sup>	T-1824 <sup>14</sup>	74.0 (58.2-90.8)			18.5 (12.0-25.4)	$\frac{PV}{100 - VH} \times 100$	90.8 (72.5-110.2)	26
37	<i>Rana cates- beiana</i> (American bullfrog)	2	T-1824	80 <sup>7</sup>			15.5 <sup>7</sup>		95	22
Pisces and Chondrichthyes										
38	<i>Ictalurus na- talis</i> (yellow bullhead)	6	T-1824	12.5 <sup>7</sup>			30.1 <sup>7</sup>		17.7	22
39	<i>Raja rhina</i> (longnose skate)	8 <sup>6</sup>	T-1824	59 (34-79)			16.8 (12.0-21.5) <sup>7,15</sup>		72 (40-95)	29
40	<i>Squalus acan- thias</i> (Atlan- tic spiny dogfish)	24 <sup>6</sup>	T-1824	55 (25-90)			18.2 (14-24) <sup>15</sup>		68 (31-109)	29

/6/ Anesthetized. /7/ Cardiac blood. /11/ New Hampshires from 6 weeks old to maturity. /12/ White Leghorns. /13/ Unfed 3-8 weeks. /14/ And high-molecular-weight dextran. /15/ 11 subjects. /16/ 25 subjects.

Contributors: (a) Reynolds, Monica, (b) Brown, Ellen

References: [1] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Armin, J., et al. 1952. J. Physiol. (London) 116:59. [3] Banerjee, S., and R. C. Bhattacharjee. 1963. Am. J. Physiol. 204:1045. [4] Bond, C. F. 1957. Ph.D. Thesis. Cornell Univ., Ithaca. [5] Bond, C. F., and P. W. Gilbert. 1958. Am. J. Physiol. 194:519. [6] Brady, L. W., et al. 1953. Surg. Gynecol. Obstet. 97:25. [7] Burke, J. D. 1954. Physiol. Zool. 27:1. [8] Bush, J. A., et al. 1955. Am. J. Physiol. 181:9. [9] Farnsworth, P. N., C. Paulino-Gonzalaez, and M. I. Gregersen. 1960. Proc. Soc. Exptl. Biol. Med. 104:729. [10] Gibson, J. G., II, et al. 1946. J. Clin. Invest. 25:838. [11] Gregersen, M. I., and R. Rawson. 1959. Physiol. Rev. 39:307. [12] Gregersen, M. I., et al. 1959. Am. J. Physiol. 196:184. [13] Hansard, S. L., H. E. Sauberlich, and C. L. Comar. 1951. Proc. Soc. Exptl. Biol. Med. 78:544. [14] Hodgetts, V. E. 1961. Australian J. Exptl. Biol. Med. Sci. 39:187. [15] Huggins, S. W. 1961. Proc. Soc. Exptl. Biol. Med. 108:231. [16] Julian, L. M., et al. 1956. J. Appl. Physiol. 8:651. [17] Kallen, F. C. 1960. Am. J. Physiol. 198:999. [18] Klement, A. W., Jr., D. E. Ayer, and E. B. Rogers. 1955. Ibid. 181:15. [19] Masouredis, S. P.,

continued

## 85. BLOOD VOLUMES

### Part I. VERTEBRATES

and L. R. Melcher. 1951. Proc. Soc. Exptl. Biol. Med. 78:264. [20] Newell, G. W., and C. S. Shaffner. 1950. Poultry Sci. 29:78. [21] Portman, O. W., K. P. McConnell, and R. H. Rigdon. 1952. Proc. Soc. Exptl. Biol. Med. 81:599. [22] Prosser, C. L., and S. J. F. Weinstein. 1950. Physiol. Zool. 23:113. [23] Reynolds, M. 1953. Am. J. Physiol. 173:421. [24] Rodnan, G. P., E. G. Ebaugh, Jr., and M. R. S. Fox. 1957. Blood 12:355. [25] Samet, P., et al. 1957. Medicine 36:211. [26] Semple, R. E. 1960. Federation Proc. 19:79. [27] Sisson, G., A. Cain, and W. S. Root. 1955. Am. J. Physiol. 180:485. [28] Steinbeck, A. W. 1954. Australian J. Exptl. Biol. Med. Sci. 32:95. [29] Thorson, T. B. 1958. Physiol. Zool. 31:16. [30] von Porat, B. 1951. Acta Med. Scand., Suppl. 256. [31] Wang, L. 1959. Am. J. Physiol. 196:188. [32] Wish, L., et al. 1950. Proc. Soc. Exptl. Biol. Med. 74:644.

### Part II. INSECTS

For additional information, consult reference 5. Hemolymph volume varies according to sex, stage of development, age, nutrition, rearing status, method of blood extraction, coagulability, and method of volume determination. Determinations were made on live or fresh-killed insects. Values in parentheses are ranges, estimate "b" or "c" as indicated (cf. Introduction).

	Species	Common Name	Stage	Method	Hemolymph Volume	Ref-er-ence
	(A)	(B)	(C)	(D)	(E)	(F)
1	<i>Aedes aegypti</i>	Yellow-fever mosquito	Larva	Exsanguination	(0.3-0.4) <sup>c</sup> cu mm/insect	17
2	<i>Apis mellifera</i>	Honeybee	Larva	Exsanguination	(25-30) <sup>c</sup> % body wt	4
3					0.04 g/insect	4
4	<i>Bombyx mori</i>	Silkworm	Larva	Exsanguination	(0.15-0.22) <sup>c</sup> ml/g body wt	3
5					0.35 ml/insect	8
6					31.2(27.6-34.8) <sup>b</sup> % body wt	15
7			Pupa	Exsanguination	(0.11-0.31) <sup>c</sup> ml/g body wt	3
8					(0.09-0.35) <sup>c</sup> ml/insect	8
9			Adult	Exsanguination	0.05 ml/insect	8
10	<i>Dytiscus</i> sp.	Diving beetle		Exsanguination	0.1 ml/insect	8
11	<i>Galleria mellonella</i>	Greater wax moth	Larva	Exsanguination	41(36.6-45.4) <sup>b</sup> % body wt (dry)	15
12	<i>Hyalophora cecropia</i>	Cecropia moth	Pupa	Exsanguination	0.25 ml/g body wt	6
13	<i>Locusta migratoria</i>	Migratory locust	Nymph	Exsanguination	<0.2 ml/insect	10
14			Adult	Exsanguination	(0-1) <sup>c</sup> cu mm/insect	10
15	<i>Periplaneta americana</i>	American cockroach	Nymph			
16			Intermolt	Dye dilution	17(11-30) <sup>c</sup> % body wt	16
17			Molting	Dye dilution	14(10-26) <sup>c</sup> % body wt	16
18			♂	Dye dilution	19.6(18.8-20.4) <sup>b</sup> % body wt	18
19				Chloride	20(16.3-23.7) <sup>b</sup> % body wt <sup>1</sup>	18
20					16.8(14.8-18.8) <sup>b</sup> % body wt <sup>a</sup>	18
21				Cell dilution	15.7(13.2-18.2) <sup>b</sup> % body wt	18
22			♀	Dye dilution	19.8(19.1-20.5) <sup>b</sup> % body wt	18
23				Chloride	18.6(14.5-22.7) <sup>b</sup> % body wt <sup>1</sup>	18
24					19.5(17.2-21.8) <sup>b</sup> % body wt <sup>a</sup>	18
25			Adult		16.8(12.3-21.3) <sup>b</sup> % body wt	18
26			Newly ecdysed	Dye dilution	21(13-35) <sup>c</sup> % body wt	16
27			24-hr old	Dye dilution	15(13-19) <sup>c</sup> % body wt	16
28			♂	Dye dilution	27.5(23.8-31.2) <sup>b</sup> % body wt	18
29				Chloride	15.3(12.9-17.7) <sup>b</sup> % body wt	18
30			♀	Dye dilution	20.9(18.8-23.0) <sup>b</sup> % body wt	18
31				Chloride	16.9(11.9-21.9) <sup>b</sup> % body wt	18
32	<i>Phormia regina</i>	Black blowfly	Larva		20 μl/insect	7
33			Adult	Dye dilution	(6.6-10.2) <sup>c</sup> μl/insect	9
34					20% body wt	9
35	<i>Popillia japonica</i>	Japanese beetle	Larva	Exsanguination	(0.9-40.8) <sup>c</sup> % body wt	2
36				Manganese	40.9(38.5-42.9) <sup>c</sup> % body wt	2
				Exsanguination	0.03 ml/insect	13

<sup>1</sup>/ Individual. <sup>2</sup>/ Pooled.

continued



## 85. BLOOD VOLUMES

### Part II. INSECTS

	Species	Common Name	Stage	Method	Hemolymph Volume	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
37	<i>Prodenia eridania</i>	Southern armyworm	Larva	Exsanguination	0.12(0.07-0.20) <sup>c</sup> ml/insect	1
38				C <sup>14</sup> inulin	0.19 ml/insect	12
39	<i>Tenebrio molitor</i>	Yellow mealworm	Larva	Dye dilution	10% body wt	11
40				Chloride	0.22 ml/g body wt	14

Contributors: (a) Jones, Jack Colvard, (b) Buck, John B.

References: [1] Babers, F. H. 1938. J. Agr. Res. 57:697. [2] Beard, R. L. 1949. J. N. Y. Entomol. Soc. 57:79. [3] Bialaszewicz, K., and C. Landau. 1938. Acta Biol. Exptl. Polish Acad. Sci. 12:307. [4] Bishop, G. H. 1923. J. Biol. Chem. 58:567. [5] Buck, J. B. 1953. In K. D. Roeder, ed. Insect physiology. J. Wiley, New York. [6] Buck, J. B., and S. Friedman. 1958. J. Insect Physiol. 2:52. [7] Evans, D. R., and V. G. Dethier. 1957-58. Ibid, 1:3. [8] Florkin, M. 1937. Acad. Roy. Belg., Classe Sci., Mem. Couron. 16:1. [9] Friedman, S. Unpublished. Natl. Institutes of Health, Bethesda, Md., 1960. [10] Hoyle, G. 1954. J. Exptl. Biol. 31:260. [11] Jones, J. C. 1957. J. Cellular Comp. Physiol. 50:423. [12] Levenbook, L. 1958. Ibid, 52:329. [13] Ludwig, D. 1951. Physiol. Zool. 24:329. [14] Munson, S. C., and J. F. Yeager. 1945. J. Econ. Entomol. 38:634. [15] Richardson, C. H., R. C. Burdette, and C. W. Eagleson. 1931. Ann. Entomol. Soc. Am. 24:503. [16] Wheeler, R. E. 1962. Federation Proc. 21(2):123. [17] Wigglesworth, V. B. 1938. J. Exptl. Biol. 15:235. [18] Yeager, J. F., and S. C. Munson. 1950. Arthropoda 1:255.

## 86. ERYTHROCYTE AND PLATELET VALUES

For information on additional species, consult reference 4, Part I.

### Part I. ERYTHROCYTE AND HEMOGLOBIN VALUES: VERTEBRATES

Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

	Species (Common Name)	Erythrocyte Count million/cu mm blood	Erythrocyte Packed Volume (Hematocrit) ml/100 ml blood	Erythrocyte Volume (Mean Cor- puscular) cu $\mu$	Hemoglobin Concentration		Erythrocyte Hemoglobin Content $\mu$ g	Erythro- cyte Dimen- sions <sup>1</sup> (Dry Film), $\mu$	Ref- er- ence
					g/100 ml blood	g/100 ml erythro- cytes			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Mammalia									
	<i>Homo sapiens</i> (man)								7-9, 12- 14, 17, 23
1	At birth <sup>2</sup>	5.7 (4.8-7.1)	56.6	106	21.5 (18.0-27.0)	38.0	38		
2	1st da	5.6 (4.7-7.0)	56.1	106	21.2 (17.7-26.5)	37.8	38		
3	1st wk	5.3 (4.5-6.4)	52.7	101	19.6 (16.2-25.5)	37.2	37		
4	2nd wk	5.1 (4.3-6.0)	49.6	96	18.0 (14.5-24.2)	36.3	35		
5	3rd wk	4.9 (4.1-6.0)	46.6	93	16.6 (13.2-23.0)	35.6	34		
6	4th wk	4.7 (3.9-5.9)	44.6	91	15.6 (12.0-21.8)	35.0	33		

<sup>1/1</sup> Dimensions for mammals are diameters. <sup>2/2</sup> When cord was clamped after placental separation rather than immediately after birth, erythrocyte count was 560,000/cu mm greater, and hemoglobin 2.6 g/100 ml greater, during first week of life. Erythrocyte and hemoglobin values were higher for heel blood (capillary) than for blood from superior sagittal sinus.

continued

## 86. ERYTHROCYTE AND PLATELET VALUES

### Part I. ERYTHROCYTE AND HEMOGLOBIN VALUES: VERTEBRATES

Species (Common Name)	Erythrocyte Count million/cu mm blood	Erythrocyte Packed Volume (Hematocrit) ml/100 ml blood	Erythrocyte Volume (Mean Cor- puscular) cu $\mu$	Hemoglobin Concentration		Erythrocyte Hemoglobin Content $\mu\mu\text{g}$	Erythro- cyte Dimen- sions <sup>1</sup> (Dry Film), $\mu$	Ref- er- ence	
				g/100 ml blood	g/100 ml erythro- cytes				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
Mammalia									
7	<i>Homo sapiens</i> (man) 2nd mo	4.5 (3.8-5.8)	38.9	85	13.3 (10.8-18.0)	34.2	30	7-9, 12- 14, 17, 23	
8	4th mo	4.5 (3.8-5.3)	36.5	79	12.4 (10.2-15.0)	34.0	27		
9	6th mo	4.6 (3.9-5.3)	36.2	78	12.3 (10.0-15.0)	34.0	27		
10	8th mo	4.6 (4.0-5.4)	35.8	77	12.1 (9.8-15.0)	33.8	26		
11	10th mo	4.6 (4.0-5.5)	35.5	77	11.9 (8.4-14.9)	33.5	26		
12	12th mo	4.6 (4.0-5.5)	35.2	77	11.6 (9.0-14.6)	33.0	25		
13	2nd yr	4.7 (3.8-5.4)	35.5	78	11.7 (9.2-15.5)	33.0	25		
14	4th yr	4.7 (3.8-5.4)	37.1	80	12.6 (9.6-15.5)	34.0	27		
15	6th yr	4.7 (3.8-5.4)	37.9	80	12.7 (10.0-15.5)	33.5	27		
16	8th yr	4.7 (3.8-5.4)	38.9	80	12.9 (10.3-15.5)	33.2	27		
17	10th yr	4.8 (3.8-5.4)	39.0	80	13.0 (10.7-15.5)	33.3	27		
18	12th yr	4.8 (3.8-5.4)	39.6	81	13.4 (11.0-16.5)	33.8	28		
19	14 yr & over Male	5.4 (4.6-6.2) <sup>b</sup>	47	(86-101)	15.8 (14.0-18.0) <sup>b</sup>	33.5	29	6,19 7.5 (7.2-7.8) <sup>b</sup>	
20	Female	4.8 (4.2-5.4) <sup>b</sup>	42	(86-101)	13.9 (11.5-16.0) <sup>b</sup>	33.5	29	7.5 (7.2-7.8) <sup>b</sup>	
21	Pregnant 6 mo	4.0 (3.5-4.8)	37 (32-42)	92	11.4 (10.2-14.0)	31	28.5	5	
22	9 mo	4.2 (3.7-5.0)	37.5 (33-43)	89	12.0 (10.8-14.4)	32	28.5		
23	Postpartum, 10 da	4.5 (4.0-5.0)	40 (35-45)	89	12.8 (11.4-14.4)	32	28.4	5	
24	<i>Bos taurus</i> (cattle)	8.1 (6.1-10.7)	40 (33-47) <sup>b</sup>	50 (47-54)	11.5 (8.7-14.5) <sup>b</sup>	29.0		5.9 1	
25	<i>Canis familiaris</i> (dog)	6.3 (4.5-8.0)	45.5 (38-53)	66 (59-68)	14.8 (11.0-18.0)	33 (30-35)	23 (21-25)	7.0 (6.2-8.0)	1
26	<i>Capra hircus</i> (goat)	16.0 (13.3-17.9)	33 (27.0-34.6)	19.3	10.5 (8.8-11.4)	34 (33-36)	6.7	4.0	22
27	<i>Cavia porcellus</i> (guinea pig)	5.6 (4.5-7.0)	42 (37-47)	77 (71-83)	14.4 (11.0-16.5)	34 (33-35)	26.0 (24.5-27.5)	7.4 (7.0-7.5)	1
28	<i>Equus caballus</i> (horse)	9.3 (8.21-10.35) <sup>b</sup>	33.4 (28-42) <sup>b</sup>		11.1 (8-14) <sup>b</sup>	33.0		5.5	1
29	<i>Felis catus</i> (cat)	8.0 (6.5-9.5)	40 (28-52)	57 (51-63)	11.2 (7.0-15.5)	28 (23-31)	14 (12-16)	6.0 (5.0-7.0)	1
30	<i>Macaca mulatta</i> (rhesus mon- key)	5.2 (3.6-6.8) <sup>b</sup>	42 (32-52) <sup>b</sup>		12.6 (10-16) <sup>b</sup>	30.0			1
31	<i>Mesocricetus</i> <i>auratus</i> (gold- en hamster)	6.96 (3.96-9.96) <sup>b</sup>	49 (39-59) <sup>b</sup>	70.0	16.0 (2.0-30.0) <sup>b</sup>	32.0	23.0	5.6 (5.4-5.8) <sup>b</sup>	11
32	<i>Mus musculus</i> (house mouse)	9.3 (7.7-12.5)	41.5	49 (48-51)	14.8 (10-19)	36 (33-39)	16 (15.5-16.5)	6.0	1

<sup>1</sup>/ Dimensions for mammals are diameters.

continued

## 86. ERYTHROCYTE AND PLATELET VALUES

### Part I. ERYTHROCYTE AND HEMOGLOBIN VALUES: VERTEBRATES

	Species (Common Name)	Erythrocyte Count million/cu mm blood	Erythrocyte Packed Volume (Hematocrit) ml/100 ml blood	Erythrocyte Volume (Mean Cor- puscular) cu $\mu$	Hemoglobin Concentration		Erythrocyte Hemoglobin Content $\mu$ g	Erythro- cyte Dimen- sions <sup>1</sup> (Dry Film), $\mu$	Ref- er- ence
					g/100 ml blood	g/100 ml erythro- cytes			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Mammalia									
33	<i>Oryctolagus cuniculus</i> (European rabbit)	5.7 (4.5-7.0)	41.5 (33-50)	61 (60-68)	11.9 (8.0-15.0)	29 (27-31)	21 (19-23)	7.5 (6.5-7.5)	1
34	<i>Ovis aries</i> (sheep)	10.3 (9.4-11.1)	31.7 (29.9-33.6)	31 (30-32)	10.9 (10.0-11.8)	34.5 (34-35)	11.0	4.8	22
35	<i>Rattus norvegicus</i> (Norway rat)	8.9 (7.2-9.6)	46 (39-53)	55 (52-58)	14.8 (12.0-17.5)	32 (30-35)	17 (15-19)	7.5 (6.0-7.5)	1,18
36	<i>Sus scrofa</i> (swine)	6.4	39.0 (38.0-40.0)	61.1 (59-63)	13.7 (13.2-14.2)	35.0	21.5 (21-22)		22
Aves									
37	<i>Anas platyrhynchos</i> (duck) <sup>3</sup>	2.8	39.5		14.8 (9-21)	38.1	52.1 (32-71)	12.8 x 6.6	21
38	<i>Anser domesticus</i> (common goose)	2.8 (2.6-3.0)	44.7 (43.1-46.2)	160 (145-174)	12.7 (11.9-13.4)	28.5 (28-29)	45.5 (40-51)	12.2 x 7.2	22
39	<i>Columba livia</i> (street pigeon)	3.2	42.3	131.0	12.8	30.0	40.0	13.2 x 6.9	22
40	<i>Gallus domesticus</i> (chicken)	2.8 (2.0-3.2)	35.6 (24.0-43.3)	127 (120-137)	10.3 (7.3-12.9)	29 (27-30)	36.6 (33-41)	11.2 x 6.8	22
41	<i>Meleagris gallopavo</i> (turkey)	2.3	38.0		11.2	23.5		15.5 x 7.5	21
Reptilia									
42	<i>Alligator mississippiensis</i> (American alligator)	0.67	30.0	450.0	8.2	27.0	123.0	23.2 x 12.1	22
43	<i>Natrix sipedon</i> (North American water snake)	0.77	35.5	465.0	10.0	28.0	131.0	19.6 x 11.0	22
44	<i>Pseudemys scripta elegans</i> (red-eared turtle)	0.69 (0.53-0.78)	17.5 (15-21)	255 (211-296)	7.3 (5.9-8.9)	41.7 (39.3-42.3)	106 (96-118)		15
45	<i>Terrapene carolina</i> (box turtle)	0.65 (0.41-0.83)	28.6 (20-38)	442 (309-587)	5.9 (5.0-8.5)	20.6 (17.4-29.7)	91 (79-131)	19 x 9	2,3
46	<i>Thamnophis sirtalis</i> (common garter snake)	1.05 (0.71-1.39)	28 (19-37)	267 (266-268)	8.5 (5.8-11.3)	31.0	82.0	18.1 x 10.3	22
Amphibia									
47	<i>Ambystoma tigrinum</i> (tiger salamander)	1.68 (1.13-1.94)	42 (27-48)	250	8.6 (5.6-10.9)	20.4	51.1		20
48	<i>Amphiuma means</i> (two-toed amphiuma)	0.03	40 (39-41)	13,857 (13,200-14,513)	9.4 (7.17-11.0)	24 (21-27)	3,287 (2,750-3,823)	62.5 x 36.3	22
49	<i>Cryptobranchus alleganiensis</i> (hellbender)	0.07	49.0	7,425	13.3	27.0	2,010	40.5 x 21.0	22
50	<i>Necturus maculosus</i> (mud puppy)	0.02	21.4	10,070	4.6	22.0	2,160	52.8 x 28.2	22

<sup>1</sup>/ Dimensions for mammals are diameters; dimensions for other vertebrates are length x width. <sup>3</sup>/ As ducks mature, hematological values progressively increase.

continued

## 86. ERYTHROCYTE AND PLATELET VALUES

### Part I. ERYTHROCYTE AND HEMOGLOBIN VALUES: VERTEBRATES

Species (Common Name)	Erythrocyte Count million/cu mm blood	Erythrocyte Packed Volume (Hematocrit) ml/100 ml blood	Erythrocyte Volume (Mean Cor- puscular) cu $\mu$	Hemoglobin Concentration		Erythrocyte Hemoglobin Content $\mu$ g	Erythro- cyte Dimen- sions <sup>1</sup> (Dry Films), $\mu$	Ref- er- ence	
				g/100 ml blood	g/100 ml erythro- cytes				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
Amphibia									
51	<i>Rana catesbeiana</i> (American bullfrog)	0.44 (0.43-0.45)	29.3 (26.6-32.0)	670 (625-716)	7.8 (7.4-8.2)	27 (26-28)	179 (174-184)	24.8 x 15.3	22
Pisces									
52	<i>Anguilla rostrata</i> (American freshwater eel)	2.48	37.9 (36.0-39.8)	156 (141-170)	9.0 (8.0-10.0)	23.5 (22-25)	36.5 (35-38)	13.0 x 8.0	22
53	<i>Cyprinus carpio</i> (carp)	0.84 (0.65-1.13)	31.3 (21-40)	311 (278-340)	10.5 (9.4-12.4)	33.5	72 (63-78)		10
54	<i>Gadus callarias</i> (rock cod)	1.55 (1.49-1.60)	29.1 (23.8-32.6)	186 (159-201)	5.9 (5.2-6.4)	20 (19-22)	38 (35-40)	12.2 x 9.0	22
55	<i>Ictalurus catus</i> (white catfish)	2.65	15.4	123	9.2	28	35	10.4 x 8.7	22
56	<i>Limanda ferruginea</i> (yellowtail flounder)	1.23 (0.78-1.61)	14.6 (8.4-18.2)	117.7 (107-138)	3.2 (2.1-4.2)	22.7 (19-25)	26.7 (26-28)	10.3 x 7.7	22
57	<i>Salvelinus fontinalis</i> (eastern brook trout)	1.01 (0.74-1.50)	27.2 (22-36)	314 (284-348)	8.5 (6.2-11.5)	31.2	75 (61-82)		10
58	<i>Scomber scombrus</i> (Atlantic mackerel)	3.94 (3.68-4.20)	57.5 (56-59)	146 (140-152)	14.9 (14.5-15.2)	26.0	37.5 (36-39)	12.5 x 8.3	22
Chondrichthyes and Agnatha									
59	<i>Dasyatis centroura</i> (rough-tail stingray)	0.30	19.0	612	3.0			20.6 x 14.3	16
60	<i>Myxine glutinosa</i> (Atlantic hagfish)	0.15 (0.12-0.19)	22.2 (19.3-27.6)	1,530 (1,470-1,560)	4.6 (4.0-5.7)	21.0	318.3 (303-330)	26.4 x 18.3	22
61	<i>Petromyzon marinus</i> (sea lamprey)	0.33	23.5	710.0	5.8			14.3 x 14.3	16
62	<i>Raja erinacea</i> (little skate)	0.09 (0.07-0.11)	7.2 (4.7-9.6)	778 (646-910)	1.4 (0.9-1.8)	19.5 (19-20)	148.5 (125-172)	24.3 x 13.9	22
63	<i>Sphyrna zygaena</i> (hammerhead shark)	0.44	23.1	526	5.4			15.2 x 11.2	16
64	<i>Squalus acanthias</i> (Atlantic spiny dogfish)	0.24	18.9	820.0	3.8			22.7 x 15.2	16

<sup>1</sup>/ Dimensions for vertebrates other than mammals are length x width.

**Contributors:** (a) Altland, Paul D., (b) Bonnycastle, Desmond D., (c) Brecher, George, (d) Cronkite, Eugene P., (e) DeMarsh, Q. B., (f) Ferguson, John H., (g) Glaser, Kurt, (h) Guest, George M., (i) Hart, J. Sanford, (j) Kisch, Bruno, (k) McCutcheon, F. Harold, (l) Mayerson, H. S., (m) Musacchia, X. J., (n) Osgood, Edwin E., (o) Rekers, Paul E., (p) Root, Raymond W., (q) Windle, William F., (r) Wintrobe, M. M., (s) Young, I. Maureen

**References:** [1] Albritton, E. C., ed. 1952. Standard values in blood. W. B. Saunders, Philadelphia. p. 42. [2] Altland, P. D., and M. Parker. 1955. Am. J. Physiol. 180:421. [3] Altland, P. D., and E. C. Thompson. 1958. Proc. Soc. Exptl. Biol. Med. 99:456. [4] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body

continued



## 86. ERYTHROCYTE AND PLATELET VALUES

### Part I. ERYTHROCYTE AND HEMOGLOBIN VALUES: VERTEBRATES

fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [5] Bethell, F. H., S. H. Gardiner, and F. MacKinnon. 1939. *Ann. Internal Med.* 13:91. [6] Brecher, G., et al. 1956. *Am. J. Clin. Pathol.* 26:1439. [7] DeMarsh, Q. B., H. L. Alt, and W. F. Windle. 1948. *Am. J. Diseases Children* 75:860. [8] DeMarsh, Q. B., et al. 1941. *J. Am. Med. Assoc.* 116:2568. [9] Elvehjem, C. A., W. H. Peterson, and D. R. Mendenhall. 1933. *Am. J. Diseases Children* 46:105. [10] Field, J. B., C. A. Elvehjem, and C. Juday. 1943. *J. Biol. Chem.* 148:261. [11] Fulton, G. P., et al. 1954. *Blood* 9:622. [12] Guest, G. M. 1938. In *Nutrition: The newer diagnostic methods*. Milbank Memorial Fund, New York. p. 138. [13] Guest, G. M., and E. W. Brown. 1957. *Am. J. Diseases Children* 93:486. [14] Guest, G. M., E. W. Brown, and M. Wing. 1938. *Ibid.* 56:529. [15] Hutton, K. E. 1961. *Am. J. Physiol.* 200:1004. [16] Kisch, B. 1951. *Exptl. Med. Surg.* 9:125. [17] Merritt, K. K., and L. T. Davidson. 1933. *Am. J. Diseases Children* 46:991. [18] Moores, R. R., et al. 1963. *Blood* 22:286. [19] Osgood, E. E. 1935. *Arch. Internal Med.* 56:849. [20] Roofe, P. G. 1961. *Anat. Record* 140:337. [21] Sturkie, P. D. 1954. *Avian physiology*. Comstock, Ithaca. [22] Wintrobe, M. M. 1934. *Folia Haematol.* 51:32. [23] Wintrobe, M. M. 1961. *Clinical hematology*. Ed. 5. Lea and Febiger, Philadelphia.

### Part II. BLOOD PLATELET COUNT: MAMMALS

Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Species (Common Name)	Platelets thousands/cu mm	Remarks	Refer- ence
(A)	(B)	(C)	(D)
<i>Homo sapiens</i> (man)			
Infant		Direct method of Wood, Vogel, and Famulener; cutaneous blood	7,19
1 At birth	227(140-290)	73 observations	
2 1 wk	233(160-320)	69 observations	
3 2 wk	242(170-370)	19 observations	
4 3 wk	269(160-380)	23 observations	
5 1 mo	277(200-370)	48 observations	
6 2 mo	320(200-470)	59 observations	
7 4 mo	324(180-450)	56 observations	
8 6 mo	350(200-480)	47 observations	
9 8 mo	346(220-480)	28 observations	
10 10 mo	340(200-450)	23 observations	
11 1 yr	339(250-470)	15 observations	
Adult	250(140-440)	13♂; direct method, phase microscopy; venous blood	1
	260(145-375)	♂♀, 185 observations; direct method, phase microscopy; venous and capillary blood	8
14 <i>Bos taurus</i> (cattle)	350(250-600)	.....	11
15	(550-600)	.....	2
16 <i>Canis familiaris</i> (dog)	326	Direct method, phase microscopy	15
17	300(100-600)	.....	11
18 <i>Capra hircus</i> (goat)	350(250-600)	.....	11
19 <i>Cavia porcellus</i>	783(525-900)	4 subjects, 8 determinations; direct method; blood from ear	16
(guinea pig)	773(680-865)	.....	18
21 <i>Equus caballus</i> (horse)	250(100-500)	.....	11
22 <i>Felis catus</i> (cat)	250(100-500)	.....	11
23 <i>Macaca mulatta</i>	344(250-750)	57 subjects	6
(rhesus monkey)	414	Direct method, no phase microscopy	12
25 <i>Mesocricetus auratus</i>	688(504-880)	10♂; direct method	10
(golden hamster)	742(500-870)	12♀; direct method	10
27 <i>Mus musculus</i> (house mouse)	1,520	92♂; direct method, phase microscopy	9
	1,190	22♂; direct method, no phase microscopy	14
29 <i>Oryctolagus cuniculus</i>	400	24 subjects; direct method, phase microscopy	3
(European rabbit)	(380-520)	12 subjects	13
31 <i>Rattus norvegicus</i>	1,240(1,100-1,380) <sup>b</sup>	60 subjects; direct method, phase microscopy	4
(Norway rat)	1,190(1,000-1,300)	18 subjects	5
33 <i>Sus scrofa</i> (swine)	445(383-507)	.....	17
34	350(250-600)	.....	11

continued

## 86. ERYTHROCYTE AND PLATELET VALUES

### Part II. BLOOD PLATELET COUNT: MAMMALS

Contributors: (a) Brecher, George, (b) Mayerson, H. S.

References: [1] Brecher, G., et al. 1950. J. Appl. Physiol. 3:365. [2] Brown, D. G., et al. 1961. Radiation Res. 15:675. [3] Cooney, D. P., et al. 1961. Acta Haematol. 26:317. [4] Cronkite, E. P., et al. 1960. In S. A. Johnson, ed. Blood platelets. Little and Brown, Boston, p. 595. [5] Hjort, P. F., and H. Paputchis. 1960. Blood 15:45. [6] Krise, G. M., et al. 1958. J. Appl. Physiol. 12:482. [7] Merritt, K. K., and L. T. Davidson. 1933. Am. J. Diseases Children 46:1008. [8] Miale, J. B. 1958. Laboratory medicine - hematology. C. V. Mosby, St. Louis. [9] Odell, T. T. Unpublished. Oak Ridge Natl. Laboratory, Oak Ridge, Tenn., 1963. [10] Otis, K., et al. 1952. Blood 7:948. [11] Pearman, V. Unpublished. Univ. Minnesota, College Veterinary Medicine, Minneapolis. [12] Pitcock, J. P., et al. 1962. Radiation Res. 16:692. [13] Rodriguez-Erdmann, F., et al. 1961. Thromb. Diath. Haemorrhag. 5:518. [14] Smith, L. H. Unpublished. Oak Ridge Natl. Laboratory, Oak Ridge, Tenn., 1963. [15] Sorensen, D. K., et al. 1960. Radiation Res. 13:669. [16] Tocantins, L. M. 1938. Medicine 17:202. [17] Trum, B. F., et al. 1959. Radiation Res. 11:326. [18] Upton, A. C., and T. T. O'Dowell, Jr. 1956. Arch. Pathol. 62:144. [19] Wood, F. C., K. M. Vogel, and L. W. Famulener. 1929. Laboratory technique. Ed. 3. T. Dougherty, New York. p. 26.

## 87. LEUKOCYTE COUNTS

For additional information, consult reference 1, Part I.

### Part I. MAN

Values were derived from smoothed curves plotted from data given in the references. Unless designated as percent (of total leukocytes), values are thousands/cu mm blood. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Age	Leukocytes, Total <sup>1</sup>	Neutrophils			Eosinophils	Basophils	Lymphocytes	Monocytes
		Total	Band <sup>2</sup>	Segmented				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 At birth	18.1(9.0-30.0) <sup>a</sup>	11.0(6.0-26.0) 61%	1.65 9.1%	9.4 52%	0.40(0.02-0.85) 2.2%	0.10(0-0.64) 0.6%	5.5(2.0-11.0) 31%	1.05(0.40-3.1) 5.8%
2 12 hr	22.8(13.0-38.0)	15.5(6.0-28.0) 68%	2.33 10.2%	13.2 58%	0.45(0.02-0.95) 2.0%	0.10(0-0.50) 0.4%	5.5(2.0-11.0) 24%	1.20(0.40-3.6) 5.3%
3 24 hr	18.9(9.4-34.0)	11.5(5.0-21.0) 61%	1.75 9.2%	9.8 52%	0.45(0.05-1.00) 2.4%	0.10(0-0.30) 0.5%	5.8(2.0-11.5) 31%	1.10(0.20-3.1) 5.8%
4 1 wk	12.2(5.0-21.0)	5.5(1.5-10.0) 45%	0.83 6.8%	4.7 39%	0.50(0.07-1.10) 4.1%	0.05(0-0.25) 0.4%	5.0(2.0-17.0) 41%	1.10(0.30-2.7) 9.1%
5 2 wk	11.4(5.0-20.0)	4.5(1.0-9.5) 40%	0.63 5.5%	3.9 34%	0.35(0.07-1.00) 3.1%	0.05(0-0.23) 0.4%	5.5(2.0-17.0) 48%	1.00(0.20-2.4) 8.8%
6 4 wk	10.8(5.0-19.5)	3.8(1.0-9.0) 35%	0.49 4.5%	3.3 30%	0.30(0.07-0.90) 2.8%	0.05(0-0.20) 0.5%	6.0(2.5-16.5) 56%	0.70(0.15-2.0) 6.5%
7 2 mo	11.0(5.5-18.0)	3.8(1.0-9.0) 34%	0.49 4.4%	3.3 30%	0.30(0.07-0.85) 2.7%	0.05(0-0.20) 0.5%	6.3(3.0-16.0) 57%	0.65(0.13-1.8) 5.9%
8 4 mo	11.5(6.0-17.5)	3.8(1.0-9.0) 33%	0.45 3.9%	3.3 29%	0.30(0.07-0.80) 2.6%	0.05(0-0.20) 0.4%	6.8(3.5-14.5) 59%	0.60(0.10-1.5) 5.2%
9 6 mo	12.0(6.0-17.5)	3.8(1.0-8.5) 32%	0.45 3.8%	3.3 28%	0.30(0.07-0.75) 2.5%	0.05(0-0.20) 0.4%	7.3(4.0-13.5) 61%	0.58(0.10-1.3) 4.8%
10 8 mo	12.2(6.0-17.5)	3.7(1.0-8.5) 30%	0.41 3.3%	3.3 27%	0.30(0.07-0.70) 2.5%	0.05(0-0.20) 0.4%	7.6(4.5-12.5) 62%	0.58(0.08-1.2) 4.7%
11 10 mo	12.0(6.0-17.5)	3.6(1.0-8.5) 30%	0.40 3.3%	3.2 27%	0.30(0.06-0.70) 2.5%	0.05(0-0.20) 0.4%	7.5(4.5-11.5) 63%	0.55(0.05-1.2) 4.6%
12 12 mo	11.4(6.0-17.5)	3.5(1.5-8.5) 31%	0.35 3.1%	3.2 28%	0.30(0.05-0.70) 2.6%	0.05(0-0.20) 0.4%	7.0(4.0-10.5) 61%	0.55(0.05-1.1) 4.8%
13 2 yr	10.6(6.0-17.0)	3.5(1.5-8.5) 33%	0.32 3.0%	3.2 30%	0.28(0.04-0.65) 2.6%	0.05(0-0.20) 0.5%	6.3(3.0-9.5) 59%	0.53(0.05-1.0) 5.0%

<sup>1</sup>/ Mean value is sum of means in columns C, F-I. <sup>2</sup>/ Includes a small percentage of myelocytes during first few days after birth. <sup>3</sup>/ Approximately 3 nucleated erythrocytes per 100 leukocytes have been found at birth.

*continued*

# Contrails

## 87. LEUKOCYTE COUNTS

### Part I. MAN

Age	Leukocytes, Total <sup>1</sup>	Neutrophils			Eosinophils	Basophils	Lymphocytes	Monocytes
		Total	Band <sup>2</sup>	Segmented				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
14 4 yr	9.1(5.5-15.5)	3.8(1.5-8.5) 42%	0.27 3.0%	3.5 39%	0.25(0.02-0.65) 2.8%	0.05(0-0.20) 0.6%	4.5(2.0-8.0) 50%	0.45(0-0.8) 5.0%
15 6 yr	8.5(5.0-14.5)	4.3(1.5-8.0) 51%	0.25 3.0%	4.0 48%	0.23(0-0.65) 2.7%	0.05(0-0.20) 0.6%	3.5(1.5-7.0) 42%	0.40(0-0.8) 4.7%
16 8 yr	8.3(4.5-13.5)	4.4(1.5-8.0) 53%	0.25 3.0%	4.1 50%	0.20(0-0.60) 2.4%	0.05(0-0.20) 0.6%	3.3(1.5-6.8) 39%	0.35(0-0.8) 4.2%
17 10 yr	8.1(4.5-13.5)	4.4(1.8-8.0) 54%	0.24 3.0%	4.2 51%	0.20(0-0.60) 2.4%	0.04(0-0.20) 0.5%	3.1(1.5-6.5) 38%	0.35(0-0.8) 4.3%
18 12 yr	8.0(4.5-13.5)	4.4(1.8-8.0) 55%	0.24 3.0%	4.2 52%	0.20(0-0.55) 2.5%	0.04(0-0.20) 0.5%	3.0(1.2-6.0) 38%	0.35(0-0.8) 4.4%
19 14 yr	7.9(4.5-13.0)	4.4(1.8-8.0) 56%	0.24 3.0%	4.2 53%	0.20(0-0.50) 2.5%	0.04(0-0.20) 0.5%	2.9(1.2-5.8) 37%	0.38(0-0.8) 4.7%
20 16 yr	7.8(4.5-13.0)	4.4(1.8-8.0) 57%	0.23 3.0%	4.2 54%	0.20(0-0.50) 2.6%	0.04(0-0.20) 0.5%	2.8(1.2-5.2) 35%	0.40(0-0.8) 5.1%
21 18 yr	7.7(4.5-12.5)	4.4(1.8-7.7) 57%	0.23 3.0%	4.2 54%	0.20(0-0.45) 2.6%	0.04(0-0.20) 0.5%	2.7(1.0-5.0) 35%	0.40(0-0.8) 5.2%
22 20 yr	7.5(4.5-11.5)	4.4(1.8-7.7) 59%	0.23 3.0%	4.2 56%	0.20(0-0.45) 2.7%	0.04(0-0.20) 0.5%	2.5(1.0-4.8) 33%	0.38(0-0.8) 5.0%
23 21 yr	7.4(4.5-11.0)	4.4(1.8-7.7) 59%	0.22 3.0%	4.2 56%	0.20(0-0.45) 2.7%	0.04(0-0.20) 0.5%	2.5(1.0-4.8) 34%	0.30(0-0.8) 4.0%

<sup>1</sup>/ Mean value is sum of means in columns C, F-I. <sup>2</sup>/ Includes a small percentage of myelocytes during first few days after birth.

**Contributors:** (a) Broun, G. O., (b) Diggs, L. W., (c) Glaser, Kurt, and Limarzi, Louis R., (d) Hamre, Christopher J., (e) Harrell, George T., (f) Osgood, Edwin E., (g) Smith, Clement A., (h) Wintrobe, M. M.

**References:** [1] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Broun, G. O. Unpublished. St. Louis Univ., School Medicine, St. Louis, 1950. [3] Glaser, K., L. R. Limarzi, and H. G. Poncher. 1950. Pediatrics 6:789. [4] Hamre, C. J., and K. K. L. Wong. 1940. Am. J. Diseases Children 60:22. [5] Hutaff, L. W., and G. T. Harrell. 1946. N. Carolina Med. J. 7:641. [6] Lippman, H. S. 1924. Am. J. Diseases Children 27:473. [7] Lucas, W. P. 1921. Ibid. 22:525. [8] Osgood, E. E., et al. 1939. Ibid. 58:61. [9] Osgood, E. E., et al. 1939. Ibid. 58:282. [10] Osgood, E. E., et al. 1939. Arch. Internal Med. 64:105. [11] Osgood, E. E., et al. 1939. J. Lab. Clin. Med. 24:905. [12] Smith, C. A. 1959. The physiology of the newborn infant. Ed. 3. C. C. Thomas, Springfield, Ill. [13] Sturgis, C. C., and F. H. Bethell. 1943. Physiol. Rev. 23:279. [14] Sunderman, F. W., and F. Boerner. 1949. Normal values in clinical medicine. W. B. Saunders, Philadelphia. [15] Washburn, A. H. 1935. Am. J. Diseases Children 50:413. [16] Wegelius, R. 1948. Acta Paediat., Suppl. 4. [17] Wintrobe, M. M. 1961. Clinical hematology. Ed. 5. Lea and Febiger, Philadelphia.

### Part II. VERTEBRATES OTHER THAN MAN

Unless designated as percent (of total leukocytes), values are thousands/cu mm blood. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Species (Common Name)	Leuko-cytes, Total	Neutrophils	Eosinophils	Basophils	Lymphocytes	Monocytes	Ref-er-ence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Mammalia							
1 <i>Bos taurus</i> (cattle)	9.2 (6.0-12.0)	2.9(1.9-3.7) 31.9(20-40)%	0.7(0.3-1.3) 7.7(3-15)%	0.06(0-0.09) 0.62(0-1)%	5.09(4.1-5.9) 55.4(45-65)%	0.48(0.27-1.40) 5.2(3-15)%	8

continued

## 87. LEUKOCYTE COUNTS

### Part II. VERTEBRATES OTHER THAN MAN

Species (Common Name)	Leuko- cytes, Total	Neutrophils	Eosinophils	Basophils	Lymphocytes	Monocytes	Ref- erence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Mammalia							
2 <i>Canis familiaris</i> (dog)	12.0 (8.0-18.0)	8.2(6.0-12.5) 68(62-80)%	0.6(0.2-2.0) 5.1(2-14)%	0.085(0-0.3) 0.7(0-2)%	2.5(0.9-4.5) 21(10-28)%	0.65(0.3-1.5) 5.2(3-9)%	1
3 <i>Capra hircus</i> (goat)	(5.0-14.0)	(2.10-3.35)	(0-1.1)	(0-0.6)	(2.10-11.25)	(0.05-0.60)	4
4 <i>Cavia</i> sp. (guinea pig)	10.0 (7.0-19.0)	4.2(2.0-7.0) 42(22-50)%	0.4(0.2-1.3) 4(2-12)%	0.07(0-0.3) 0.7(0-2)%	4.9(3.0-9.0) 49(37-64)%	0.43(0.25-2.00) 4.3(3-13)%	1
5 <i>Equus caballus</i> (horse)	(5.0-11.0)	(3.0-6.9)	(0.05-0.60)	(0-0.1)	(1.2-4.8)	(0.10-1.45)	4
6 <i>Felis catus</i> (cat)	16.0 (9.0-24.0)	9.5(5.5-16.5) 59.5(44-82)%	0.85(0.2-2.5) 5.4(2-11)%	0.02(0-0.1) 0.1(0-0.5)%	5.0(2.0-9.0) 31(15-44)%	0.65(0.05-1.40) 4(0.5-7.0)%	1
7 <i>Mus</i> sp. (mouse)	8.0 (4.0-12.0)	2.0(0.7-4.0) 25.5(12-44)%	0.15(0-0.5) 2(0-5)%	0.05(0-0.1) 0.5(0-1)%	5.5(3.0-8.5) 68(54-85)%	0.3(0-1.3) 4(0-15)%	1
8 <i>Oryctolagus cunicu- lus</i> (rabbit)	9.0 (6.0-13.0)	4.1(2.5-6.0) 46(36-52)%	0.18(0-0.4) 2(0.5-3.5)%	0.45(0.15-0.75) 5(2-7)%	3.5(2.0-5.6) 39(30-52)%	0.725(0.3-1.3) 8(4-12)%	1
9 <i>Ovis aries</i> (sheep)	7.8(5-10)	2.8(1.6-3.5) 35.7(20-45)%	0.19(0.08-0.5) 2.5(1-7)%	0.3(0-0.15) 0.4(0-2)%	4.4(3.9-5.5) 56.9(50-70)%	0.47(0.08-0.60) 6(1-8)%	8
10 <i>Rattus</i> sp. (rat)	14.0 (5.0-25.0)	3.1(1.1-6.0) 22(9-34)%	0.3(0-0.7) 2.2(0-6)%	0.1(0-0.2) 0.5(0-1.5)%	10.2(7.0-16.0) 73(65-84)%	0.3(0-0.65) 2.3(0-5)%	1
11 <i>Sus scrofa</i> (swine)	(7.0-20.0)	(2.4-10.0)	(0.05-2.00)	(0-0.8)	(3.2-12.0)	(0.05-2.00)	4
Aves							
12 <i>Anas</i> sp. (duck)	23.4	24.3%	2.1%	1.0(0-4.5)	45.8(13.0-73.5)	4.4(0.5-11.5)	6,9
13 <i>Gallus domesticus</i> (chicken)	32.6 (9.1-56.0)	9.1(3.0-18.2) <sup>1</sup> 27.8(9.1-56.0)%	0.05(0-0.23) <sup>2</sup> 1.5(0-7)%	0.9(0-2.6) 2.7(0-8)%	17.6(7.8-27.3) 54(24-84)%	4.4(0-9.7) 13.7(0-30)%	10
14 <i>Meleagris gallopavo</i>	19.0	44.5(35-65) <sup>3,4</sup>	7.5(1-24) <sup>3,5</sup>	6.9(3-11) <sup>3</sup>	36.3(22-46) <sup>3</sup>	7.3(2-11) <sup>3</sup>	5
15 (turkey)	(16.0-25.5)	45.4(39-52) <sup>4,6</sup>	2.3(0-5) <sup>5,6</sup>	5.1(1-9) <sup>6</sup>	40.9(35-48) <sup>6</sup>	6.5(3-10) <sup>6</sup>	
Reptilia							
16 <i>Pituophis sayi</i> (bull snake)	50.2 <sup>7</sup>	2.7 <sup>1,2</sup> 5.4%		0.01 0.2%	19.7 39.4%	3.4 6.9%	7
17 <i>Terrapene carolina</i> (box turtle)	37.5 (24.0-48.0)	0.01 <sup>7</sup> 0.3%	4.1 10.8%	3.0 8%	21.0 56.1%	3.5 9.4%	2,3

/1/ Heterophils with rod-shaped eosinophilic bodies. /2/ Heterophils with granular eosinophilic bodies. /3/ Supravital stain. /4/ Polymorphic myelocytes with eosinophilic rods. /5/ Polymorphic myelocytes with pseudoeosinophilic granules. /6/ Wright's stain. /7/ Includes thrombocytes.

Contributors: (a) Altland, Paul D., (b) Dunlap, J. S., (c) Rigdon, R. H.

- References: [1] Albritton, E. C., ed. 1951. Standard values in blood. W. B. Saunders, Philadelphia. p. 53.  
 [2] Altland, P. D., and M. Parker. 1955. Am. J. Physiol. 180:421. [3] Bernstein, R. E. 1938. S. African J. Sci. 35:327. [4] Craige, A. H., Jr. Unpublished. Univ. Pennsylvania, School Veterinary Medicine, Philadelphia, 1950.  
 [5] Dunlap, J. S. Unpublished. Washington State College, College Veterinary Medicine, Pullman, 1956.  
 [6] Hewitt, R. 1942. Am. J. Hyg. 36:6. [7] Ryerson, D. L. 1949. J. Entomol. Zool. 41(4):49. [8] Scarborough, R. A. 1931-32. Yale J. Biol. 4:69. [9] Sturkie, P. D. 1954. Avian physiology. Comstock, Ithaca.  
 [10] Twisselmann, N. M. 1939. Poultry Sci. 18:151.



## 88. BONE MARROW DIFFERENTIAL CELL COUNTS

### Part I. RIB: DOG

Values in parentheses are ranges, estimate "c" unless otherwise specified (cf. Introduction).

Specification	Mulligan	Mulligan	Values from Rekers and Coulter	Stasney and Higgins <sup>1</sup>	Van Loon and Clark
(A)	(B)	(C)	(D)	(E)	(F)
1 Number of subjects	21	35	36	35	81
2 Age of subjects	0.5-2.5 da	Adult	19-24 mo	Adult	Adult
3 Cells, % of total count					
Proerythroblasts	1.3(0.4-3.2) <sup>2</sup>	0.5(0-1.4) <sup>2</sup>	0.3(0-1.3) <sup>3</sup>	59(40-78)	0.6(0.2-2.7) <sup>2</sup>
Normoblasts					
Early	5.8(3.0-9.5) <sup>4</sup>	1.5(0.4-3.8) <sup>5</sup>	28.2(8.0-53.9) <sup>4</sup>		7.8(6.4-10.0) <sup>5</sup>
Intermediate	45.1(33.0-56.6) <sup>6</sup>	38.1(18.6-63.6) <sup>6</sup>	4.6(0-11.2) <sup>6</sup>		16.4(11-26) <sup>7</sup>
Late				17.4(9-26) <sup>6</sup>	
7 Myeloblasts		0.6(0-1.8)	1.9(0.2-3.7)	2.4(0-5.1) <sup>9</sup>	0.6(0.2-1.0)
8 Promyelocytes	0.8(0-2.2)	1.5(0.2-4.6)	0.7(0-3.3)	2.8(0-5.8)	1.6(0.7-2.8)
9 Myelocytes			2.7(0-9.5)		6.0(2.7-10.0)
10  Neutrophilic	4.3(2.0-6.6)	4.7(2.2-11.2)		8.9(2.8-15.0)	
11  Eosinophilic				1.2(0-2.4)	
12 Metamyelocytes	9.7(7.2-12.0)	10.5(5.6-20.0)	5.1(0-24.4) <sup>10</sup>	15.3(7.2-23.0)	3.4(1.1-4.6)
13 Band cells	20.6(14.8-28.4) <sup>11</sup>	31.0(16.8-53.8) <sup>11</sup>	42.4(16.5-62.9) <sup>11</sup>		11.7(6.8-17.0)
Segmented cells					
14  Neutrophilic	3.4(0.8-6.6)	3.9(0.2-8.6)	5.0(0.2-14.3)	5.1(0-12.5)	30.1(17-44)
15  Eosinophilic	2.4(0-5.2)	3.7(1.0-6.8)	4.7(0.2-19.3)	2.8(0-6.8)	2.0(0.4-3.8)
16  Basophilic			0.2(0-1.3)	0.1(0-0.3)	
17 Lymphocytes	3.3(1.6-6.0)	1.9(0-6.6)	0.7(0-8)	1.2(0.2-2.3)	0.9(0.2-2.7)
18 Monocytes					0.2(0-0.3)
19 Megakaryocytes			0.6(0-1.1)	0.1(0-0.5)	0.5(0-1.4)
20 Plasma cells			0.4(0-2.1)		
21 Reticulum cells				1.0(0-2.1)	
22 Unclassified cells	3.1(0.8-5.4)	2.1(0.8-6.1)	3.0(0-15.7)	0.2(0-0.7) <sup>12</sup>	
Reference	1	2	3	4	5

<sup>1/</sup> Ranges are estimate "b" (cf. Introduction). <sup>2/</sup> Pronormoblasts. <sup>3/</sup> Megaloblasts <sup>4/</sup> Erythrocytes.  
<sup>5/</sup> Basophilic normoblasts. <sup>6/</sup> Normoblasts. <sup>7/</sup> Polychromic normoblasts. <sup>8/</sup> Orthochromic normoblasts.  
<sup>9/</sup> Includes leukoblasts. <sup>10/</sup> Juvenile cells. <sup>11/</sup> Stab cells. <sup>12/</sup> Includes heterophils.

**Contributor:** Rekers, Paul E.

**References:** [1] Mulligan, R. M. 1941. Anat. Record 79:101. [2] Mulligan, R. M. 1945. Ibid. 91:161.  
 [3] Rekers, P. E., and M. Coulter. 1948. Am. J. Med. Sci. 216:643. [4] Stasney, J., and G. M. Higgins. 1937. Ibid. 193:462. [5] Van Loon, E. J., and B. B. Clark. 1943. Clin. Med. 28:1575.

### Part II. STERNUM: MAN

For additional information, consult reference 1. All values are for adults. Values in parentheses are ranges, estimate "c" unless otherwise specified (cf. Introduction).

Specification	Berman	Diggs	Israëls	Leitner	Values from Lucia and Hunt <sup>1</sup>	Osgood and Seaman <sup>1,2</sup>	Vaughan and Brockmyre	Whitby and Britton	Wintrobe
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Number of subjects	19♂♀			20♂♀	6♂♀	28♂, 24♀	42♂, 8♀		
2 Marrow aspirated, ml	1.5	(0.1-0.2)	0.2	(0.1-0.3)	0.5	(0.5-10.0)	3.0	0.25	(1.0-2.0)
3 Nucleated cells, total, thousands/cu mm				(60-100)		35 (10-100)	35.3 (9.4-74.0)		

<sup>1/</sup> Ranges are estimate "b" (cf. Introduction). <sup>2/</sup> Values are smoothed weighted means and calculated ranges.

*continued*

## 88. BONE MARROW DIFFERENTIAL CELL COUNTS

### Part II. STERNUM: MAN

Specification	Values from								
	Berman	Diggs	Israëls	Leitner	Lucia and Hunt <sup>1</sup>	Osgood and Seaman <sup>1,2</sup>	Vaughan and Brockmyre	Whitby and Britton	Wintrobe
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Cells, % of total count									
4 Proerythroblasts	3 <sup>3,4</sup>	(0-1) <sup>5</sup>	(0.5-4.0)	0.8 <sup>6</sup>	0.6(0-3.4) <sup>7</sup>	0.2(0-1) <sup>8</sup>		(0-4)	4(1-8) <sup>4</sup>
5 Normoblasts							9.5		18(7-32)
6 Early	10 <sup>9,10</sup>	(1-4) <sup>11</sup>	(1-5)	3.2 <sup>6,12</sup>	8(0-20.4) <sup>13</sup>	2.0(0-4) <sup>14</sup>	(1.5-24.0) <sup>9</sup>	(4-15)	
7 Intermediate	86 <sup>9,16</sup>	(10-20) <sup>15</sup>	(12-20)	24.4 <sup>6,9</sup>		6.0(4-8) <sup>17</sup>			
8 Late	13 <sup>13</sup>	(5-10) <sup>15</sup>	(6-10)		12(0-25) <sup>9</sup>	3.0(1-5) <sup>20</sup>		(7-19)	
9 Myeloblasts	3 <sup>21</sup>	(0-1)	(0.3-2.0)	1.2	1.0(0-2.8)	0.4(0-1) <sup>22</sup>	1.3(0-3)	(0-2.5)	2(0.3-5)
10 Promyelocytes	9	(1-5)	(1-8)	2.2	4(1.2-6.8)	1.4(0-3)		(0.5-5.0)	5(1-8)
11 Myelocytes							8.9(2-16)		
12 Neutrophilic	6	(2-10)	(5-20)	12.6 <sup>23</sup>	10.2 (4.0-16.4)	4.2(0-12) <sup>23</sup>		(2-8)	12(5-19)
13 Eosinophilic				1.4	0.8(0-1.8)			(0-1)	1.5(0.5-3)
14 Basophilic				0.02					0.3(0-0.5)
15 Metamyelocytes						6.5(3-10) <sup>24</sup>	8.8 (3.5-18.0) <sup>25</sup>		22(13-32)
16 Neutrophilic	9	(5-15)	(13-32) <sup>26</sup>	10.2	13.2 (7.8-18.6)			(10-25)	
17 Eosinophilic				0.8	1.3(0.5-2.1)			(0-2.5)	
18 Band cells						24(17-33) <sup>27</sup>	23.9(12-34)		
19 Neutrophilic	31	(10-40)		24	24.6(5.2-44)				
Segmented cells									
20 Neutrophilic	17	(10-30)	(7-30)	28.4	10(0-22.4) <sup>28</sup>	15(5-25) <sup>29</sup>	18.5(6-36) <sup>30</sup>	(10-40) <sup>28,29</sup>	20(7-30) <sup>28</sup>
21 Eosinophilic		(0-3)	(0.5-4.0)	1.8	0.3(0-0.9) <sup>28</sup>	2(0-4) <sup>29</sup>	1.9(0-6.5) <sup>30</sup>	(0-4) <sup>28,29</sup>	2(0.5-4) <sup>28</sup>
22 Basophilic	2	(0-1)	(0-1)	0.02		0.2 (0-0.5) <sup>29</sup>	0.2(0-1.5) <sup>30</sup>	(0-1) <sup>28</sup>	0.2(0-0.7) <sup>28</sup>
23 Lymphocytes	14	(5-15)	(3-20)	7.6	10.3 (1.5-19.1)	14(3-25)	16.2(7-35)	(5-20)	10(3-17)
24 Monocytes		(0-2)	(0.5-5.0)	1.4	0.5	2(0-4)	2.4(0-6)	(0-5)	2(0.5-5)
25 Megakaryocytes	5			0.8					0.4(0.03-3)
26 Plasma cells	1	(0-1)	(0-2)	1.2	1.5(0-4.1)		0.3(0-1.5)	(0-1)	0.4(0-2)
27 Reticulum cells	2			0.4 <sup>32</sup>			0.3(0-2.5) <sup>33</sup>		0.2(0.1-2)
28 Unclassified cells				3.5			0.02(0-0.5)		
29 Disintegrated cells						19(10-30)	7.9(0-18)		
Reference	2	3	4	5	6	7	8	9	10


<sup>1/1</sup> Ranges are estimate "b" (cf. Introduction). <sup>1/2</sup> Values are smoothed weighted means and calculated ranges.  
<sup>1/3</sup> Percent of red series. <sup>1/4</sup> Pronormoblasts. <sup>1/5</sup> Rubriblasts. <sup>1/6</sup> Per 100 leukocytes. <sup>1/7</sup> Megaloblasts.  
<sup>1/8</sup> Karyoblasts. <sup>1/9</sup> Normoblasts. <sup>1/10</sup> Basophilic normoblasts. <sup>1/11</sup> Prorubricytes. <sup>1/12</sup> Macroblasts.  
<sup>1/13</sup> Erythroblasts. <sup>1/14</sup> Prokaryocytes. <sup>1/15</sup> Polychromatophilic normoblasts. <sup>1/16</sup> Rubricytes. <sup>1/17</sup> Karyocytes.  
<sup>1/18</sup> Orthochromic normoblasts. <sup>1/19</sup> Metarubricytes. <sup>1/20</sup> Metakaryocytes. <sup>1/21</sup> Includes leukoblasts. <sup>1/22</sup> Granu-  
 loblasts. <sup>1/23</sup> Includes early neutrophilic myelocytes. <sup>1/24</sup> Metagranulocytes. <sup>1/25</sup> Young forms. <sup>1/26</sup> Includes band  
 cells. <sup>1/27</sup> Rhabdocytes. <sup>1/28</sup> Polymorphonuclear cells. <sup>1/29</sup> Lobocytes. <sup>1/30</sup> Filament cells. <sup>1/31</sup> Occasionally  
 present. <sup>1/32</sup> Endothelial cells. <sup>1/33</sup> Reticuloendothelial cells.

**Contributors:** (a) Diggs, L. W., (b) Osgood, Edwin E.

**References:** [1] Altman, P. L., and D. S. Dittmer, ed. 1961. Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. [2] Berman, L. 1949. Blood 4:511. [3] Diggs, L. W. In S. E. Miller, ed. 1960. Textbook of clinical pathology. Ed. 6. Williams and Wilkins, Baltimore. p. 60. [4] Israëls, M. C. G. 1955. An atlas of bone marrow pathology. Ed. 2. Grune and Stratton, New York. [5] Leitner, S. M. 1945. Die intravitale Knochenmarksuntersuchung. B. Schwabe, Basel. [6] Lucia, S. P., and M. L. Hunt. 1947. Am. J. Med. Sci. 213:686. [7] Osgood, E. E., and A. J. Seaman. 1944. Physiol. Rev. 24:46. [8] Vaughan, S. L., and F. Brockmyre. 1947. Blood, Spec. Issue 1:54. [9] Whitby, L. E. H., and C. J. C. Britton. 1963. Disorders of the blood. Ed. 9. Grune and Stratton, New York. [10] Wintrobe, M. M. 1961. Clinical hematology. Ed. 5. Lea and Febiger, Philadelphia.

# Contrails

## NORMAL BLOOD AND MARROW CELLS: MAN



- |  |                                      |                                  |
|--|--------------------------------------|----------------------------------|
| 1 Erythrocytes   | 3 <sup>c</sup> Normoblast in mitosis | 11 Neutrophil leukocytes         |
| 1 <sup>a</sup> Erythrocytes in rouleau                       | 4 Hemolyzed red cells (ghosts)       | 12 Basophil leukocyte            |
| 1 <sup>b</sup> Deformed cells (poikilocytes), crenated forms | 5 Platelets                          | 13 Polychromatophil erythroblast |
| 2 Reticulocytes stained with dilute solution of cresyl blue  | 6 Small lymphocyte                   | 13 <sup>a</sup> Hemocytoblast    |
| 3 Early normoblasts  | 7 Monocytes                          | 14 Megaloblast                   |
| 3 <sup>a</sup> Extruded nucleus                              | 8 Large lymphocytes                  | 15 Eosinophil myelocyte          |
| 3 <sup>b</sup> Late normoblast                               | 9 Megakaryocyte                      | 16 Neutrophil myelocyte          |
|  | 10 Eosinophil leukocyte              | 17 Neutrophil metamyelocyte      |

*Reference:* Best, C. H., and N. B. Taylor. 1961. The physiological basis of medical practice. Ed. 7. Williams and Wilkins, Baltimore.



## IX. BIOLOGICAL REGULATORS AND TOXINS

### 89. ENZYMES

#### Part I. CATALYTIC ACTION

**Catalytic Action** (column B) and **Cofactors** (column F): ADP = adenosine diphosphate; AMP = adenosine-5'-monophosphate; ATP = adenosine triphosphate; CoA = coenzyme A; NAD<sup>+</sup> = nicotinamide adenine dinucleotide; NADP<sup>+</sup> = nicotinamide adenine dinucleotide phosphate. **Method** (column G): Chem = chemical; Col = colorimetric; Enzy = enzymatic; Mano = manometric; Pol = polariscopic; Phys = physical; Thun = Thunberg; Titr = titrimetric; Turb = turbidity. Figures (column G) are wavelengths used for measuring light absorption.

Enzyme	Catalytic Action (Substrate → Product)	Conditions Suitable for Enzyme Action <sup>1</sup>					Occurrence
		pH	Substrate Concentration	Temp. °C	Cofactor	Method	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Aceto-coenzyme A-kinase	ATP + acetate + CoA → acetyl CoA + AMP + pyrophosphate	7.5	0.005 M	37	Mg <sup>++</sup> , K <sup>+</sup>	Chem	Yeast
2 Acetylcholine esterase	Acetylcholine → acetate + choline	7.4	3 mg/ml	37	.....	Mano	Liver, pancreas, brain, blood
3 Aconitase	Citric acid → <i>cis</i> -aconitic acid <sup>2</sup>	7.4	0.03 M	25	Fe <sup>++</sup>	240 mμ	Tissues, bacteria, yeasts, seeds, leaves
4 Adenase	Adenine → hypoxanthine + NH <sub>3</sub>	8.7	.....	40	.....	Chem	Muscle, milk, blood
5 Adenosinetriphosphatase	ATP → ADP + PO <sub>4</sub>	7.5	1 mg P/ml	37	Ca <sup>++</sup>	Chem	Brain, muscle, venoms, potatoes
6 Alcohol dehydrogenase	Ethanol → acetaldehyde	7.8	0.03%	20	NAD <sup>+</sup>	Mano	Liver, kidney, brain, blood, yeasts, bacteria, higher plants
7 Aldolase	Fructose-1, 6-diphosphate → triosephosphates	9	0.01 M	38	Co <sup>++</sup> , Fe <sup>++</sup> , or Zn <sup>++</sup>	Chem	Muscle, <i>Escherichia coli</i> , yeasts, higher plants
8 Amino acid carboxylase	Amino acid → amines + CO <sub>2</sub>	4-5.5	0.001 M	30	Pyridoxal phosphate	Mano	Liver, kidney, pancreas, bacteria, higher plants
9 D-Amino acid oxidase	D-Amino acids + O <sub>2</sub> → α-keto acids + H <sub>2</sub> O <sub>2</sub> + NH <sub>3</sub>	8.6	0.01 M	38	.....	Mano	Widespread (animals); fungi
10 L-Amino acid oxidase	L-Amino acids + O <sub>2</sub> → α-keto acids + H <sub>2</sub> O <sub>2</sub> + NH <sub>3</sub>	8.8	0.015 M	38	.....	Mano	Liver, kidney, venoms, fungi, bacteria
11 Aminotripeptidase	Tripeptide → dipeptide + amino acid	8.0	0.05 M	39	.....	Titr	Mucosa, muscle
12 α-Amylase (animal)	Starch or glycogen → dextrins + maltose	7	1%	37	NaCl	Chem	Liver, saliva, urine
13 α-Amylase (plant)	Starch or glycogen → dextrins + maltose	4.5-5.5	12 mg/ml	30	.....	Chem	Bacteria, yeasts, cereals
14 β-Amylase (animal, plant)	Starch → dextrins + maltose	4-5	12 mg/ml	...	.....	.....	Cereals, soybeans, sweet potatoes
15 Amylophosphorylase	Dextrin + glucose-1-phosphate → starch or glycogen + phosphate	6.8	0.001 M	30	Starch or glycogen	Chem	Widespread (animals, plants)
16 Amylosucrase	Sucrose → "glycogen" + fructose	5.6	10 mg/ml	23	.....	Chem	Bacteria
17 Apyrase	ATP → AMP + 2 phosphate	6.5	.....	30	Ca <sup>++</sup>	Chem	Liver, muscle, yeasts, tubers

<sup>1</sup>/ Conditions vary with the method used and with the source of the enzyme. <sup>2</sup>/ To isocitric acid.

*continued*



Part I. CATALYTIC ACTION

Enzyme	Catalytic Action (Substrate → Product)	Conditions Suitable for Enzyme Action <sup>1</sup>					Occurrence
		pH	Substrate Concentration	Temp. °C	Cofactor	Method	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
18 Arginase	L-Arginine → L-ornithine + urea	9.5	0.66%	38	Co <sup>++</sup> , Mn <sup>++</sup>	Mano	Liver, bacteria, fungi, seeds, spleen
19 Ascorbic acid oxidase	L-Ascorbic acid + O <sub>2</sub> → dehydroascorbate + H <sub>2</sub> O	6.0	0.01 N	20	.....	Mano	Widespread (plants)
20 Asparaginase	L-β-Asparagine → L-aspartic acid + NH <sub>3</sub>	8	0.5 M	40	.....	Chem	Liver, mucosa, bacteria, fungi, seeds
21 Aspartase	L-Aspartic acid → fumaric acid + NH <sub>3</sub>	7-7.5	0.1 M	37	.....	Chem	Bacteria, yeasts, leaves
22 Carbonic anhydrase	H <sub>2</sub> CO <sub>3</sub> → CO <sub>2</sub> + H <sub>2</sub> O	5-9	0.08 M	15	.....	Mano	Erythrocytes, gastric mucosa
23 Carboxypeptidase	Peptide (free COOH) → amino acid + peptide	8.5	6% edestin	25	.....	Titr	Pancreas (as xy-mogen)
24 Catalase	H <sub>2</sub> O <sub>2</sub> → H <sub>2</sub> O + O <sub>2</sub>	6.8	0.01 N	0	.....	Chem	Erythrocytes, liver, kidney, bacteria, higher plants
25 Chlorophyllase	Chlorophyll → chlorophyllide + phytol	5.9	1 mg/ml	25	CaCl <sub>2</sub>	Chem	Bacteria, leaves, stems
26 Cholesterol esterase	Cholesterol esters → cholesterol + acids	5.3 or 7	.....	...	.....	.....	Liver, kidney, spleen, intestinal mucosa, blood, pancreas, bacteria
27 Choline acetylase	Choline + acetyl CoA → acetylcholine	7	.....	...	CoA, ATP	Chem	Brain, muscle, bacteria
28 Chymotrypsin	Proteins → polypeptides + amino acids	7.6	5% casein	38	.....	Chem	Pancreas
29 Conjugase	Pteroylglutamate → pterine + glutamic acid	7-8	.....	37	Ca <sup>++</sup>	.....	Pancreas, tissues, yeasts, tubers
30 Cytochrome-c oxidase	Ferro-cytochrome-c + O <sub>2</sub> → ferri-cytochrome-c + H <sub>2</sub> O	7.2	0.0001 M	37	.....	Mano	Widespread (animals, plants)
31 Cytochrome-c peroxidase	Ferro-cytochrome-c + H <sub>2</sub> O <sub>2</sub> → ferri-cytochrome-c + H <sub>2</sub> O	7.4	1.5 x 10 <sup>-5</sup> M	20	.....	550 mμ	Yeasts
32 Cytochrome-c reductase	Ferri-cytochrome-c → ferro-cytochrome-c	7.3	2 x 10 <sup>-5</sup> M	25	NADP <sup>+</sup>	550 mμ	Liver, yeasts
33 Deoxyribonuclease	Thymonucleic acid → nucleotides	6-7	0.5%	37	Mg <sup>++</sup> , Mn <sup>++</sup>	Phys	Intestinal mucosa, pancreas, seeds
34 Dextranucrase	Sucrose → dextran + fructose	5.6	10 mg/ml	23	.....	Chem	Bacteria
35 Enolase	2-Phosphoglycerate → (enol) phosphopyruvate	7	0.1 mg P/ml	20	Mg <sup>++</sup> , Mn <sup>++</sup> , Zn <sup>++</sup>	Chem	Muscle, yeasts, leaves
36 Esterase, simple	Ethyl butyrate → ethanol + butyrate	8.0	Saturated	20	.....	Titr	Widespread (animals); seeds, fungi
37 Ficin	Proteins → amino acids and peptides?	5	.....	35	H <sub>2</sub> S, HCN, cysteine	.....	Fig tree sap
38 Fructose-1, 6-diphosphatase	Fructose diphosphate + H <sub>2</sub> O → fructose-6-phosphate + orthophosphate	8.8	0.02 M	25	Mg <sup>++</sup>	340 mμ	Spinach
39 Fumarase	Fumaric acid → L(-)malic acid	6.6	0.025 M	40	.....	Chem	Liver, muscle, bacteria, fungi, higher plants
40 β-Galactosidase (lactase)	Lactose → galactose + glucose	5.6	2.5%	38	.....	Chem	Bacteria, seeds

<sup>1</sup>/ Conditions vary with the method used and with the source of the enzyme.

*continued*

Part I. CATALYTIC ACTION

Enzyme	Catalytic Action (Substrate → Product)	Conditions Suitable for Enzyme Action <sup>1</sup>					Occurrence	
		pH	Substrate Concentration	Temp. °C	Cofactor	Method		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
41	Glucose dehydrogenase	D-Glucose → D-gluconic acid	7.4	0.2 M	38	NAD <sup>+</sup> or NADP <sup>+</sup>	Thun	Liver
42	Glucose oxidase (notatin)	D-Glucose + O <sub>2</sub> → gluconate + H <sub>2</sub> O <sub>2</sub>	6.0	1%	39	.....	Mano	Fungi
43	Glucose-6-phosphate dehydrogenase	Glucose-6-phosphate → phosphogluconate	7.5	0.02 M	38	NADP <sup>+</sup>	Mano	Blood, yeasts
44	α-Glucosidase (maltase)	Maltose → glucose	7.2	50 mg/ml	30	.....	Pol	Intestinal mucosa, fungi, malt
45	β-Glucosidase	β-Glucosides → glucose + aglycon	4.4-5.0	1 mg/ml	30	.....	Pol	Intestinal mucosa, liver, kidney
46	β-Glucuronidase	β-Glucuronide → glucuronate + alcohol	4.5	0.001 M	38	.....	Col	Widespread (animals); bacteria, higher plants
47	Glutamate dehydrogenase	Glutamate → α-ketoglutarate + NH <sub>3</sub>	8.2	0.0001 M	37	NAD <sup>+</sup> or NADP <sup>+</sup>	340 μ	Liver, kidney, muscle, brain
48	Glyoxalase	Methylglyoxal → lactate	7	1 mg/ml	25	Glutathione	Mano	Liver, kidney, muscle, blood, bacteria, fungi, seeds
49	Guanase	Guanine → xanthine + NH <sub>3</sub>	8.7	Saturated	40	.....	Chem	Liver, pancreas, spleen, kidney, seeds
50	Hexokinase	Hexose + ATP → hexosemonophosphate + ADP	7.5	0.001 M	30	Mg <sup>++</sup> , Mn <sup>++</sup>	Chem	Liver, muscle, kidney, brain, bacteria, yeasts, higher plants
51	Histaminase	Histamine → aldehyde + H <sub>2</sub> O <sub>2</sub> + NH <sub>3</sub>	6.8-7.6	0.01 M	37	.....	Mano	Widespread (animals); bacteria, fungi
52	Histidase	Histidine → glutamate + formate + NH <sub>3</sub>	8	0.01 M	38	.....	Chem	Liver
53	Hyaluronidase	Hyaluronate → acetylglucosamine + glycuronate	7.0	0.1%	37	.....	Phys	Spleen, testes, insects, venoms, bacteria
54	β-Hydroxybutyrate dehydrogenase	L-β-Hydroxybutyrate → acetoacetate	7	0.05 M	38	NAD <sup>+</sup>	Mano	Widespread (animals, plants)
55	Invertase (sucrase, saccharase)	Sucrose → glucose + fructose	4.5	4 g/25 ml	20	.....	Pol	Intestinal mucosa, invertebrates, fungi, bacteria, higher plants
56	Isocitratase	Isocitrate → succinate + glyoxylate	7.9	0.025 M	30	Mg <sup>++</sup>	Chem	Aerobic bacteria
57	Isocitrate dehydrogenase	D-Isocitrate → α-ketoglutarate + CO <sub>2</sub>	7.0	0.002 M	25	NAD <sup>+</sup> , NADP <sup>+</sup> , Mg <sup>++</sup> , Mn <sup>++</sup>	340 μ	Widespread (animals, plants)
58	Lactate dehydrogenase	Lactate → pyruvate	9.3	0.02 M	20	NAD <sup>+</sup>	Mano	Widespread (animals)
59	Lecithinase A	Lecithin → lysolecithin + fatty acid	7	Egg yolk	38	Ca <sup>++</sup>	Chem	Liver, muscle, pancreas, venoms, mushrooms
60	Lecithinase B	Lysolecithin → glycerylphosphorylcholine + fatty acid	4	.....	41	.....	Chem	Liver, spleen, pancreas, brain, fungi, seeds, rice bran

<sup>1/1</sup> Conditions vary with the method used and with the source of the enzyme.

*continued*

Part I. CATALYTIC ACTION

Enzyme	Catalytic Action (Substrate → Product)	Conditions Suitable for Enzyme Action <sup>1</sup>					Occurrence
		pH	Substrate Concentration	Temp. °C	Cofactor	Method	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
61 Leucylpeptidase	Leucyl peptides → leucine + other amino acids	8-9	0.05 M	40	Mg <sup>++</sup> , Mn <sup>++</sup>	.....	Intestinal mucosa, leaves, malt, bacteria, fungi
62 Levansucrase	Sucrose → levan + glucose + fructose	5.0-5.8	5%	...	.....	Chem	Bacteria
63 Lipase, pancreatic	Fats → glycerol + fatty acids	9	2.5 g/15 ml	30	CaCl <sub>2</sub>	Titr	Pancreas
64 Lipase, seed	Fats → glycerol + fatty acids	4.7-5.0	.....	...	Ca <sup>++</sup>	.....	Seeds
65 Lipoxidase	Linoleic acid, etc., → oxidized fatty acids	6.5	0.02%	25	.....	Col	Intestinal mucosa, muscle, seeds
66 Luciferase	Luciferin + O <sub>2</sub> → oxidized luciferin + light	7.2	10 <sup>-6</sup> M	23	Mg <sup>++</sup> , ATP	.....	Insects, ostracods, bacteria, fungi
67 Lysozyme	Bacterial cells → lysed bacterial cells	5.3	Suspension	38	.....	Turb	Nasal mucosa; latex of fig
68 Malate dehydrogenase	L(-)Malate → oxalacetate	7.2	0.025 M	37	NAD <sup>+</sup> or NADP <sup>+</sup>	Thun	Brain, kidney, liver, muscle; widespread (plants)
69 Nucleoside phosphorylase	Inosine → hypoxanthine + ribose-1-phosphate	7.5	0.001 M	30	.....	290 mμ	Spleen, lungs, liver, blood, intestinal mucosa; traces (plants)
70 Oxalacetic carboxylase	Oxalacetate → pyruvate + CO <sub>2</sub>	5.0	0.5 mg/L	30	Mn <sup>++</sup>	.....	Liver, bacteria, seeds, leaves
71 Papain	Proteins, proteoses, etc. → amino acids	7.5	2%	30	HCN, H <sub>2</sub> , S, cysteine	Col	Seeds, latex
72 Pectinesterase	Pectin → pectate + methanol	6.2	1%	30	.....	Titr	Leaves, fruit, bacteria
73 Pectinase	Pectin → galacturonide	4.0	0.5%	25	.....	Chem	Bacteria, fungi
74 Pepsin	Proteins → proteoses, peptones, amino acids	1.5-2.0	2%	20	.....	Col	Gastric mucosa
75 Peroxidase, plant	H <sub>2</sub> O <sub>2</sub> + pyrogallol, etc. → H <sub>2</sub> O + purpurogallin, etc.	4.5-6.5	0.25%	20	.....	Col	Widespread (plants)
76 Phosphocarboxyltransphosphorylase	1, 3-Diphosphoglycerate + ADP → 3-phosphoglycerate + ATP	7.9	1 mg/ml	25	Mg <sup>++</sup>	340 mμ	Muscle, yeasts
77 Phosphoenoltransphosphorylase	Phosphopyruvate + AMP → pyruvate + ATP	.....	1.5 mM	38	Mg <sup>++</sup> , K <sup>+</sup>	.....	Muscle, yeasts, higher plants
78 Phosphoglucomutase	Glucose-1-phosphate → glucose-6-phosphate	7.5-9.2	10 <sup>-6</sup> M	30	Co <sup>++</sup> , Mg <sup>++</sup> , Mn <sup>++</sup>	Chem	Widespread (animals, plants)
79 Phosphoglyceromutase	3-Phosphoglycerate → 2-phosphoglycerate	7	10 <sup>-5</sup> M	24	.....	Chem	Widespread (animals, plants)
80 Phosphomonoesterase I (alkaline)	β-Glycerophosphate → H <sub>3</sub> PO <sub>4</sub> + glycerol	9.2	0.02 M	37	Mg <sup>++</sup>	Chem	Widespread (animals); bacteria, fungi; none in higher plants
81 Phosphomonoesterase II (acid)	β-Glycerophosphate → H <sub>3</sub> PO <sub>4</sub> + glycerol	5-6	0.05 M	37	.....	Chem	Prostate, spleen, liver, bacteria, fungi, seeds, tubers
82 Phosphomonoesterase III	Monoesters of phosphate → H <sub>3</sub> PO <sub>4</sub> + alcohols	3.4-4.2	.....	...	.....	.....	Liver, spleen, fungi, seeds
83 Phosphomonoesterase IV	α-Glycerophosphate → H <sub>3</sub> PO <sub>4</sub> + glycerol	5.2-6.2	.....	...	Mg <sup>++</sup> , Mn <sup>++</sup>	Chem	Blood, bacteria, yeasts

<sup>1</sup>/ Conditions vary with the method used and with the source of the enzyme.

*continued*

*89. ENZYMES*  
Part I. CATALYTIC ACTION

Enzyme	Catalytic Action (Substrate → Product)	Conditions Suitable for Enzyme Action <sup>1</sup>					Occurrence
		pH	Substrate Concentration	Temp. °C	Cofactor	Method	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
84 Phytase	Phytate → inositol + phosphate	5.5-7.8	0.1%	37	Mg <sup>++</sup>	Chem	Blood, intestinal mucosa, fungi, seeds
85 Pyrophosphatase I	Pyrophosphate → phosphate	7.2-7.8	0.001 M	38	Mg <sup>++</sup> , Mn <sup>++</sup>	Chem	Widespread (animals); fungi, seeds
86 Pyruvic carboxylase	Pyruvate → acetaldehyde + CO <sub>2</sub>	6.0	0.15 M	30	Thiamine pyrophosphate, Mg <sup>++</sup>	Mano	Fungi, bacteria, seeds
87 Q-Enzyme	Amylose → amylopectin	7.0	.....	21	.....	.....	Liver, muscle, seeds, tubers
88 Rennin	Casein → paracasein	5.8	Raw milk	40	.....	Phys	Calf stomach
89 Ribonuclease	Ribonucleic acid → ribonucleotides	4-5	0.25 mg P/ml	25	.....	Chem	Liver, spleen, pancreas, lungs, bacteria, higher plants
90 Succinate dehydrogenase	Succinate → fumarate	7.4	0.01 M	37	Cytochrome-c ?	Thun	Widespread (animals, plants)
91 Sucrose phosphorylase	Sucrose + phosphate → fructose + glucose-1-phosphate	6.6	0.1 M	30	.....	Chem	Bacteria
92 Synthetase, glycogen	(Glucose) <sub>n</sub> + uridine diphosphate glucose → (glucose) <sub>n+1</sub> + uridine diphosphate	8.4	0.005 M	37	.....	Enzy	Animal tissue
93 Transaminase	Glutamate + oxalacetate → α-ketoglutarate + aspartate	7.5	0.02 M	40	.....	Chem	Widespread (animals, plants)
94 Triosephosphate dehydrogenase	D-Glyceraldehyde-3-phosphate → 1,3-diphosphoglycerate	8.6-9.0	0.0001 M	27	NAD <sup>+</sup>	340 mμ	Widespread (animals, plants)
95 Trypsin	Proteins (especially denatured) → polypeptides and amino acids	8-9	2.2%	25	.....	Col	Pancreatic juice
96 Tyrosinase	Catechol, etc. + O <sub>2</sub> → o-quinone, etc., + H <sub>2</sub> O	5.5-7	2 mg/ml	25	.....	Mano	Melanomas, skin; plants
97 Urease	Urea → CO <sub>2</sub> + NH <sub>3</sub>	7.0	1.5%	20	.....	Chem	Blood, gastric mucosa, insects, bacteria, fungi, seeds
98 Xanthine oxidase	Xanthine or aldehyde → uric or other acids	7.5	0.003 M	20	.....	Mano	Liver, milk

<sup>1</sup>/ Conditions vary with the method used and with the source of the enzyme.

**Contributors:** (a) Somers, G. Fred., (b) Perlman, D., (c) Campbell, Jack J. R.

**References:** [1] Boyer, P. D., H. Lardy, and K. Myrbäck, ed. 1959-63. The enzymes. Ed. 2. Academic Press, New York. [2] Dixon, M., and E. C. Webb. 1958. Enzymes. Academic Press, New York. [3] Sumner, J. B., and G. F. Somers, ed. 1953. Chemistry and methods of enzymes. Ed. 3. Academic Press, New York.

*continued*



Data are for enzymes in the crystalline state. Abbreviations: AMP = adenosine-5'-monophosphate; ADP = adenosine-5'-diphosphate; NADP<sup>+</sup> = nicotinamide adenine dinucleotide phosphate; NADPH = reduced NADP<sup>+</sup>; Tris = Tris(hydroxymethyl)amino-

	Common Name (Systematic Name) <sup>1</sup> [Code Number]	Source	Physical Properties				Optimum pH
			S <sub>20, w</sub> <sup>2</sup> sec x 10 <sup>13</sup>	D <sub>20, w</sub> <sup>2</sup> cm <sup>2</sup> sec <sup>-1</sup> x 10 <sup>-7</sup>	Molecular Weight	Isoelectric Point <sup>3</sup> pH	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Adenylic deaminase (AMP amino-hydroxylase) [3.5.4.6]	Rabbit muscle	12.3	3.76	320,000	5.6	6.4 <sup>6</sup>
2	Alcohol dehydrogenase (alcohol:NAD <sup>+</sup> oxidoreductase) [1.1.1.1]	Bakers' yeast	6.72	4.70	150,000	5.4	
3		Horse liver	5.11	5.96	84,000	6.8	8.0
4	Aldolase (Ketose-1-phosphate aldehydelyase) [4.1.2.7]	Cattle liver	8.87 <sup>8</sup>	5.20 <sup>8</sup>	159,000	6.6-6.7	9.1-9.5 <sup>9</sup>
5		Rabbit muscle	7.35	4.63	149,000	6.05	5-10
6	L-Amino acid oxidase (L-Amino acid:O <sub>2</sub> oxidoreductase, deaminating) [1.4.3.2]	Snake venom	6.8 <sup>10</sup>	5.1 <sup>10</sup>	130,000		7.2-7.5
7	α-Amylase (α-1,4-Glucan 4-glucanohydrolase) [3.2.1.1]	<i>Bacillus subtilis</i>	4.56 <sup>11</sup> 6.47 <sup>12a</sup>	7.98 <sup>11</sup> 5.72 <sup>12a</sup>	48,900 <sup>11</sup> 96,900 <sup>12</sup>		5.8-6.0
8	β-Amylase (α-1,4-Glucan maltohydrolase) [3.2.1.2]	Sweet potato	8.9	5.77	152,000	4.74-4.79	4-5
9	Carboxypeptidase A [3.4.2.1]	Cattle pancreatic juice	3.07	8.82	34,440	5.95 <sup>13</sup>	7.4 <sup>14</sup>
10	Catalase (H <sub>2</sub> O <sub>2</sub> :H <sub>2</sub> O <sub>2</sub> oxidoreductase) [1.11.1.6]	Cattle liver	11.3	4.5	225,000	5.7	4-8.5
			11.15	4.1	244,000		
11	Chymotrypsin [3.4.4.5]	Cattle pancreas	2.56	10.2	22,500 25,000	8.1-8.6 <sup>15</sup>	7-9
12	Creatine kinase (ATP:creatine phosphotransferase) [2.7.3.2]	Rabbit muscle	5.0	5.78	81,000	6.0-6.1	9.0 <sup>17</sup> 6-7 <sup>18</sup>
13	Crotonase (L-3-Hydroxyacyl-CoA hydrolase) [4.2.1.17]	Cattle liver	7.84		210,000 <sup>20</sup>		9.0 <sup>21</sup>
14	Enolase (D-2-Phosphoglycerate hydrolase) [4.2.1.1]	Yeast	5.9	8.08	67,000		7.8
15	Fumarase (L-Malate hydro-lyase [4.2.1.2])	Swine heart	9.09	4.05	220,000	7.35 <sup>22</sup>	6.3
16	β-Galactosidase (β-D-Galactoside galactohydrolase) [3.2.1.23]	<i>Escherichia coli</i>	16.24	2.12	750,000		7.3 <sup>24</sup>
17	Glucose oxidase (β-D-Glucose:O <sub>2</sub> oxidoreductase) [1.1.3.4]	<i>Penicillium amagasakiense</i>	7.93	5.02	154,000	4.35	5.6-5.8
18	Glucose-6-phosphate dehydrogenase (D-Glucose-6-phosphate:NADP <sup>+</sup> oxidoreductase)	Brewers' yeast	6.3 <sup>11</sup>		100,000 <sup>11, 21</sup>		8.5
			9.3 <sup>12</sup>		200,000 <sup>12, 21</sup>		
19	Glutamate dehydrogenase (L-Glutamate:NADP <sup>+</sup> oxidoreductase)	Cattle liver	26.6	2.54	1,000,000	4-5	8.3-8.5
20	Glyceraldehyde-3-phosphate dehydrogenase (D-Glyceraldehyde-3-phosphate:NAD <sup>+</sup> oxidoreductase)	Rabbit muscle	7.01	5.46	120,000	6-8 in phosphate <sup>16</sup>	8.5-9.0

<sup>1/</sup> Systematic names and code numbers recommended in the *Report of the Commission on Enzymes of the Interdiffusion* (D<sub>20, w</sub>) coefficients are for data normalized to standard conditions of water at 20°C and extrapolated to molar activity is defined as the number of molecules of substrate transformed per minute by one molecule of enzyme (Michaelis constant) =  $\frac{K_{-1} + K_{+2}}{K_{+1}}$ , where K<sub>+1</sub> = velocity constant for formation of the enzyme-substrate complex, for the breakdown of the enzyme-substrate complex into products. <sup>2/</sup> In 0.1 M succinate. <sup>3/</sup> V<sub>max</sub> (maximal <sup>4/</sup> In glycylglycine. <sup>5/</sup> Not extrapolated to zero protein concentration. <sup>6/</sup> Monomer. <sup>7/</sup> Dimer. <sup>8/</sup> Determined approximately 30,000. <sup>9/</sup> Depends on ionic strength of the buffer. <sup>10/</sup> Forward reaction. <sup>11/</sup> Reverse reaction. <sup>12/</sup> Approximately. <sup>13/</sup> In 0.1 M Tris. <sup>14/</sup> K<sub>m</sub>'s independent of pH but dependent on ionic strength. <sup>15/</sup> Buffer second order rate constant.

**ENZYMES**

**KINETIC PROPERTIES**

sine diphosphate; ATP = adenosine triphosphate; CoA = coenzyme A; NAD<sup>+</sup> = nicotinamide adenine dinucleotide; methane.

Kinetic Properties								Reference	
Molecular Activity			Michaelis Constant						
Substrate	Temp. °C	pH	Value <sup>d</sup>	Substrate	Temp. °C	pH	K <sub>m</sub> <sup>e</sup> moles/liter		
(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	
AMP (4.5 x 10 <sup>-5</sup> M, calculated V <sub>max</sub> ) <sup>7</sup>	30	6.4	18,300	AMP	30	6.4	1.43 x 10 <sup>-3</sup>	B,66;C-F,67;G-O,68	1
Ethanol	26	7.9	27,000	Ethanol	Room	8.2	1.4 x 10 <sup>-2</sup>	B,96;C-F,45;H-K,20;L-O,28	2
				Ethanol	23.5	8.0	2.5 x 10 <sup>-4</sup>	B,11;C,D,31;E,30;F,25;G,L-O,132	3
Fructose-1,6-di-phosphate	30	9.4	400	Fructose-1,6-di-phosphate	30	9.4	4 x 10 <sup>-3</sup>	B,G-O,91;C-F,92	4
Fructose-1,6-di-phosphate	30	7.5	4,200	Fructose-1,6-di-phosphate	30		5 x 10 <sup>-5</sup>	B,126;C-E,127;F,136;G,76;H-K,7;L,M,O,134	5
L-Leucine	38	7.45	2,800					B,G,144;C-E,12;H-K,144,145	6
Maltose	25	6.0	86,000/48,900 g enzyme	Glucosidic bonds	20-30		1.5 x 10 <sup>-3</sup>	B,H-K,33;C-E,34;G,77;L,M,O,12	7
Glycosidic linkages	30	4.8	250,000					B,G,5;C-F,32;H-K,12,32	8
Chloroacetyl-α-β-phenyllactic acid	25	7.5	7,240	Benzenesulfonyl glycyphenyl-alanine	25	7.5	1.4 x 10 <sup>-2</sup>	B,2;C,105;D,95,105;E,109;F,95;G,106;H-K,81;L-O,110	9
H <sub>2</sub> O <sub>2</sub>			10 <sup>715</sup>					B,118;C,99,120;D,98,119;E,99,119;F,143;G,H,K,19	10
Benzoyl-L-phenyl-alanine methyl-ester	25	7.8	3,000	Acetyl-L-tyrosine amide	25	7.8	2.7 x 10 <sup>-2</sup>	B,62;C,104;D,102;E,12,104;F,1;G,44;H-O,81	11
ATP, creatine	30	9	16,000	ATP	30	9	5 x 10 <sup>-4</sup> <sup>19</sup>	B,57;C-F,85;G-O,58	12
ADP, creatine	30	7	100,000	Creatine	30	7	1.6 x 10 <sup>-2</sup> <sup>19</sup>		
				ADP			8 x 10 <sup>-4</sup> <sup>19</sup>		
				Creatine phosphate			5 x 10 <sup>-3</sup> <sup>19</sup>		
Crotonyl CoA	25	7.5	730,000	Crotonyl CoA			1.56 x 10 <sup>-4</sup>	B,112;C,E,12;G-L,O,138	13
Phosphoglyceric acid		7.4	5,400					B,141;C,12,73;D,E,H,J,K,12;G,147	14
Fumarate	20	7.3	100,000	Fumarate	25		2 x 10 <sup>-6</sup> to 4 x 10 <sup>-5</sup> <sup>39</sup>	B,37,74;C,37;D,17;E,37,49;F,103;G,36;H-K,74;L,M,O,38	15
o-Nitrophenyl-β-D-galactoside	20	7.6	133,500 (V <sub>max</sub> ) <sup>7</sup>	o-Nitrophenyl-β-D-galactoside	20	7.6	1.61 x 10 <sup>-4</sup>	B,48,140;C-E,48;G-O,12	16
O <sub>2</sub>	30		17,000 <sup>28</sup>	Glucose	30		1.5 x 10 <sup>-2</sup>	B-E,G-I,K,64;F,12;L,M,O,63	17
Glucose-6-phosphate	30	8.0	67,600/100,000 g enzyme	Glucose-6-phosphate	25	8.0	2.0-5.8 x 10 <sup>-5</sup>	B,H-K,88;C,E,12;G,41;L-O,41,69	18
Glutamate			1,000 mole/1,000,000 g enzyme	Glutamate	25	8.0	1.8 x 10 <sup>-3</sup>	B,90,115,116;C-H,K,90;L-O,12	19
Glyceraldehyde-3-phosphate and NAD <sup>+</sup>	27	8.6	10,300/min/mole enzyme <sup>28</sup>	Glyceraldehyde-3-phosphate	27	8.6	3.9-5.1 x 10 <sup>-5</sup>	B,G,22;C-E,127;F,137;H-O,20	20

*national Union of Biochemistry*, 1961, Pergamon Press, New York. /<sup>2</sup>/ Values for sedimentation (S<sub>20,w</sub>) and zero protein concentration. /<sup>3</sup>/ Apparent values determined from electrophoretic mobility measurements. /<sup>4</sup>/ Molyzyme at optimal substrate concentration. Values pertain to the molecular weight given in column E. /<sup>5</sup>/ K<sub>m</sub> K<sub>-1</sub> = velocity constant for the dissociation of this complex into substrate and enzyme, and K<sub>+2</sub> = velocity constant velocity) = v/(1+K<sub>m</sub>/S), where v is the measured velocity at a substrate concentration S. /<sup>6</sup>/ Determined at 25°C. mined at 0.2 ionic strength. /<sup>14</sup>/ In Veronal buffer. /<sup>15</sup>/ Hypothetical value calculated from a "katalasefähigkeit" of /<sup>19</sup>/ K<sub>m</sub>'s are apparent constants for total substrates under specified conditions. /<sup>20</sup>/ Determined by light scattering. used was 0.05 M NaCl and 0.05 M Tris. /<sup>28</sup>/ Derived from QO<sub>2</sub> = 148,000 μl O<sub>2</sub> per mg per hour. /<sup>29</sup>/ Calculated as

*continued*

	Common Name (Systematic Name) <sup>1</sup> [Code Number]	Source	Physical Properties				Optimum pH
			S <sub>20, w</sub> <sup>a</sup> sec x 10 <sup>13</sup>	D <sub>20, w</sub> <sup>a</sup> cm <sup>2</sup> sec <sup>-1</sup> x 10 <sup>-7</sup>	Molecular Weight	Isoelectric Point <sup>b</sup> pH	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
21	Glycerokinase (ATP:glycerol phosphotransferase) [2.7.1.30]	<i>Candida mycoderma</i>	10.87	4.2	251,000	4.6	
22	α-Glycerophosphate dehydrogenase (L-Glycerol-3-phosphate:NAD <sup>+</sup> oxidoreductase) [1.1.1.8]	Rabbit muscle	4.9	5.1	78,000		7.5 <sup>29</sup>
23	Glycogen phosphorylase <sup>28</sup> (α-1,4-Glucan:orthophosphate glucosyltransferase) [2.4.1.1]	Rabbit muscle	13.2	2.6	495,000	<5.8	6.8
24	Hexokinase (ATP:D-hexose 6-phosphotransferase) [2.7.1.1]	Bakers' yeast	3.1	2.9	96,600	4.5-4.8	
25	Homoserine dehydratase (L-Homoserine hydro-lyase, deaminating) [4.2.1.15]	Rat liver	8.9	4.1	190,000		8.0
26	Lactate dehydrogenase (L-Lactate:NAD <sup>+</sup> oxidoreductase) [1.1.1.27]	Cattle heart	7.0	5.3	135,000	4.5-4.8	
27	Myokinase <sup>29</sup> (ATP:AMP phosphotransferase) [2.7.4.3]	Rabbit muscle	2.30 <sup>b</sup>	1.00 <sup>a</sup>	21,279	6.1	8.0
28	Old yellow enzyme (NADPH:(acceptor) oxidoreductase) [1.6.99.1]	Brewers' yeast	5.82	5.54	102,000	5.22	
29	Papain [3.4.4.10]	Papaya latex	2.42	10.27	20,700	8.75	5-10
30	Pepsin [3.4.4.1]	Cattle stomach mucosa	3.20	8.71	35,700	<1.0	2-4 <sup>30</sup>
31	Peroxidase (Donor:H <sub>2</sub> O <sub>2</sub> oxidoreductase) [1.11.1.7]	Horseradish	3.48		39,800	7.2	
32	Phosphoglucosomerase (D-Glucose-6-phosphate ketolisomerase) [5.3.1.9]	Rabbit muscle	7.5	5.3	140,000		9
33	Phosphoglucosomutase (D-Glucose-1,6-diphosphate:D-glucose-1-phosphate phosphotransferase) [2.7.5.1]	Rabbit muscle	3.69	4.83	74,000		7.5
34	Phosphoglycerate kinase (ATP:D-3-phosphoglycerate 1-phosphotransferase) [2.7.2.3]	Brewers' yeast	3.20		34,000		
35	Phosphoglyceromutase (D-2,3-Diphosphoglycerate:D-2-phosphoglycerate phosphotransferase) [2.7.5.3]	Bakers' yeast	6.30	5.29	112,000	5.0-5.5	5.9 or 7.0 <sup>33</sup>
36	Pyrophosphatase, inorganic (Pyrophosphate phosphohydrolase) [3.6.1.1]	Bakers' yeast	4.4	6.8	63,000	4.75	7.0
37	Pyruvate kinase (ATP:pyruvate phosphotransferase)	Rabbit muscle	10.04	3.96	237,000	6.6	
38	Ribonuclease (Polyribonucleotide:2-oligoribonucleotidotransferase, cyclizing) [2.7.7.16]	Cattle pancreas	1.87	11.14	13,683	9.604 <sup>34</sup>	7.2-8.2
39	Transketolase (D-Sedoheptulose-7-phosphate:D-glyceraldehyde-3-phosphate glycolaldehydetransferase) [2.2.1.1]	Bakers' yeast	7.34	5.0	140,000		7.6

/1/ Systematic names and code numbers recommended in the *Report of the Commission on Enzymes of the International Union of Pure and Applied Chemistry*. /2/ Diffusion (D<sub>20, w</sub>) coefficients are for data normalized to standard conditions of water at 20°C and extrapolated to zero substrate concentration. /3/ Molecular activity is defined as the number of molecules of substrate transformed per minute by one molecule of enzyme. /4/ Michaelis constant =  $\frac{K-1+K+2}{K+1}$ , where K+1 = velocity constant for formation of the enzyme-substrate complex, and K = velocity constant for the breakdown of the enzyme-substrate complex into products. /5/ V<sub>max</sub> (maximal velocity) =  $v(1+K_m/S)$ , where v = initial velocity, M phosphate. /6/ Composed of two molecules of phosphorylase b, each with a molecular weight of 242,000. /7/ Phosphate concentration. /8/ Valid only under conditions specified in reference 14. /9/ Depending upon type of assay.



**ENZYMES**

**KINETIC PROPERTIES**

Kinetic Properties								Reference	
Molecular Activity				Michaelis Constant					
Substrate	Temp. °C	pH	Value <sup>4</sup>	Substrate	Temp. °C	pH	K <sub>m</sub> <sup>5</sup> moles/liter		
(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	
Glycerol			100,000	Glycerol			6 x 10 <sup>-5</sup>	B-F,H,K,L,O,10	21
Dihydroxyacetone phosphate	20	7	20,670	α-Glycerophosphate	23	7.0	1.1 x 10 <sup>-4</sup>	B,H-K,6;C-E,135;G,L-O,148	22
Glucose-1-phosphate	30	6.7	40,000	Glucose-1-phosphate	30	6.8	5 x 10 <sup>-3</sup>	B,43;C-E,51,52;F,42;G,20;H-O,21	23
Glucose	30	8	55,000	D-Glucose	28	7.6	1.67 x 10 <sup>-4</sup>	B,20,61;C-F,61;H-K,20;L-O,39	24
Homoserine			6,400 (V <sub>max</sub> ) <sup>7</sup>	Homoserine			2 x 10 <sup>-2</sup>	B-E,75;G,H,K,L,O,12	25
Pyruvate	Room	7.0	37,000	Lactate	28.5	7.0	1.7 x 10 <sup>-2</sup>	B,114;C-F,79;H-K,80;L-O,146	26
AMP (formation of ATP)	25	8.0	28,000	ATP	25	8.0	3.3 x 10 <sup>-4</sup>	B,83;C,D,F,84;E,72;G-O,82	27
O <sub>2</sub>	37	7.4	155	NADP <sup>+</sup>			0.9 x 10 <sup>-5</sup>	B,131;C-E,30;F,128;H-L,O,55	28
Benzoyl-L-argininamide	39	5.5	2.7-6 x 10 <sup>-6</sup> /mole enzyme	Benzoyl-L-argininamide	38	6	3.9 x 10 <sup>-2</sup>	B,H-K,54;C-F,107;G,108;L-O,113	29
Carbobenzoxy-L-glutamyl-L-tyrosine ethyl ester	38	4.0	0.183	Carbobenzoxy-L-glutamyl-L-tyrosine ethyl ester	32	4.0	1.78 x 10 <sup>-3</sup>	B,46;C-E,29;F,93;G,89;H-K,15;L-O,12	30
				H <sub>2</sub> O <sub>2</sub>	25-30	4.7	10 <sup>-8</sup>	B,129;C,E,16;F,130;L-O,18	31
Glucose-6-phosphate	30	8.0	77,000	Glucose-6-phosphate	30	8.0	3 x 10 <sup>-5</sup>	B,H-K,86;C-E,87;G,20;L-O,50	32
Glucose-1-phosphate	30	7.5	12,400	Glucose-1-phosphate			10 <sup>-5</sup> <sup>21,21</sup>	B,G,78;C-E,53;H-L,O,94	33
ATP	25	6.9	110,000	ATP			1.1 x 10 <sup>-4</sup> <sup>20</sup>	B,H-L,O,14;C,E,65	34
D-2-Phosphoglycerate	30	7.0	98,000	2-Phosphoglycerate			<10 <sup>-4</sup>	B,97;C-L,O,20	35
Pyrophosphate			60,000					B,F-H,K,60;C-E,100	36
Pyruvate	25	7.5	57,000-66,000	Phosphoenolpyruvate	30	7.4	7 x 10 <sup>-5</sup>	B,8,133;C-F,142;H-K,12;L-O,71	37
Ribonucleic acid	25	5.0	7.5 moles phosphate liberated/mole enzyme					B,G,59;C,D,40;E,101;F,125;H-K,70	38
Xylulose-5-phosphate and ribose-5-phosphate			3,400	Xylulose-5-phosphate (with 0.005 M R-5-P) Ribose-5-phosphate (with 0.0019 M xylulose-5-P)			2.1 x 10 <sup>-4</sup> 4 x 10 <sup>-4</sup>	B,27,111;C-E,H,K,12;G,L,O,26	39

*national Union of Biochemistry*, 1961, Pergamon Press, New York. /2/ Values for sedimentation (S<sub>20,w</sub>) and zero protein concentration. /3/ Apparent values determined from electrophoretic mobility measurements. /4/ Molyzyme at optimal substrate concentration. Values pertain to the molecular weight given in column E. /5/ K<sub>m</sub> K<sub>-1</sub> = velocity constant for the dissociation of this complex into substrate and enzyme, and K<sub>+2</sub> = velocity constant v is the measured velocity at a substrate concentration S. /6/ Determined at 25°C. /2/ Approximately. /27/ In 0.2 /20/ Adenylate kinase. /30/ With protein; pH 4 with synthetic substrates. /21/ K<sub>m</sub> depends on glucose-1,6-diphosphate /24/ Isoelectric point value is that of the isoionic point in 0.001 M KCl.

*continued*



Part II. PHYSICAL AND

	Common Name (Systematic Name) <sup>1</sup> [Code Number]	Source	Physical Properties				Optimum pH
			S <sub>20, w</sub> <sup>2</sup> sec x 10 <sup>13</sup>	D <sub>20, w</sub> <sup>2</sup> cm <sup>2</sup> sec <sup>-1</sup> x 10 <sup>-7</sup>	Molecular Weight	Isoelectric Point <sup>3</sup> pH	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
40	Trypsin [3.4.4.4]	Cattle pancreas	2.50	9.40	23,800	10.5	7-8
41	Tyrosinase (o-Diphenol:O <sub>2</sub> oxidoreductase) [1.10.3.1]	<i>Neurospora crassa</i>	4.3	10.7	33,000	6-8	6-8
42	Urease (Urea amidohydrolase) [3.5.1.5]	Jack bean meal	18.6	3.46	483,000	5.0-5.1	8.0
43	Xanthine oxidase (Xanthine:O <sub>2</sub> oxidoreductase) [1.2.3.2]	Milk	11.4	3.6	290,000	5.3-5.4	8.3

<sup>1</sup> Systematic names and code numbers recommended in the *Report of the Commission on Enzymes of the International Union of Pure and Applied Chemistry*. <sup>2</sup> Diffusion (D<sub>20, w</sub>) coefficients are for data normalized to standard conditions of water at 20°C and extrapolated to standard conditions. <sup>3</sup> Molecular activity is defined as the number of molecules of substrate transformed per minute by one molecule of enzyme.  $K_{+1} = \frac{K_{-1} + K_{+2}}{K_{+1}}$ , where K<sub>+1</sub> = velocity constant for formation of the enzyme-substrate complex, for the breakdown of the enzyme-substrate complex into products.

**Contributor:** Noltmann, Ernst A.

**References:** [1] Anderson, E. A., and R. A. Alberty. 1948. *J. Phys. Colloid Chem.* 52:1345. [2] Anson, M. L. *J. Chem. Soc.*, p. 1212. [3] Balls, A. K., M. K. Walden, and R. R. Thompson. 1948. *J. Biol. Chem.* 173:9. [4] Beisenherz, G., et al. 1953. *Z. Naturforsch.* 8b:555. [5] Bergmann, F., and S. Dikstein. 1956. *J. Biol. Chem. Scand.* 4:715. [6] Boyer, P. D., H. A. Lardy, and K. Myrbäck, ed. 1959-63. *The enzymes*. Ed. 2. Academic Press, New York. [7] Casey, E. J., and K. J. Laidler. 1953. *J. Am. Chem. Soc.* 72:2159. [8] Cecil, R., and A. G. Ogston. *Biochem. Biophys.* 22:224. [9] Chance, B., and A. C. Maehly. 1961. In C. Long, ed. *Biochemists' handbook*. Academic Press, New York. [10] Cori, C. F., G. T. Cori, and A. A. Green. 1943. *J. Biol. Chem.* 151:39. [11] Cunningham, L. W., Jr., et al. 1953. *Discussions Faraday Soc.* 13:58. [12] Dalziel, K. 1958. *Acta Chem. Scand.* 11:1257. [13] Ehrenberg, A., and K. Dalziel. 1958. *Ibid.* 12:465. [14] Englard, S., and T. P. Singer. 1960. *Proc. 4th Intern. Congr. Biochem., Vienna, 1958*, 8:124. [15] Fling, M., N. H. 212:859. [16] Frieden, C., R. M. Bock, and R. A. Alberty. 1954. *J. Am. Chem. Soc.* 76:2482. [17] Frieden, C., 237:3027. [18] Ginsberg, A., P. Appel, and H. K. Schachman. 1956. *Arch. Biochem. Biophys.* 65:545. [19] Glaser, G. T. Cori. 1943. *Ibid.* 151:21. [20] Gutfreund, H., and J. M. Sturtevant. 1956. *Biochem. J.* 63:655. [21] Hayes, [22] Horowitz, N.H., and S. Shen. 1952. *J. Biol. Chem.* 197:513. [23] Hu, A. S. L., R. G. Wolfe, and R. J. Reithel. [24] Kahana, S. E., et al. 1960. *J. Biol. Chem.* 235:2178. [25] Keller, P. J. 1955. *Ibid.* 214:135. [26] Keller, *Ibid.* 20:115. [27] Kimmel, J. R., and E. L. Smith. 1954. *J. Biol. Chem.* 207:515. [28] Kistner, S. 1959. *Acta S. A., L. Noda, and H. A. Lardy. 1954. J. Biol. Chem.* 209:191. [29] Kuby, S. A., L. Noda, and H. A. Lardy. 1954. and M. R. McDonald. 1946. *Ibid.* 29:393. [30] Kunitz, M., and J. H. Northrop. 1936. *Ibid.* 19:991. [31] Kusae, K. 40:555. [32] Larsson-Raznikiewicz, M., and B. G. Malmström. 1961. *Arch. Biochem. Biophys.* 92:94. [33] Lee, [34] Lowry, O. H., et al. 1961. *Ibid.* 236:2746. [35] McDonald, M. R. 1948. *J. Gen. Physiol.* 32:39. [36] McQuate, 1962. *Ibid.* 237:1138. [37] Malmström, B. G. 1957. *Arch. Biochem. Biophys.* 70:58. [38] Massey, V. 1952. 1963. *Ibid.* 238:100. [39] Menzi, R., E. A. Stein, and E. H. Fischer. 1957. *Helv. Chim. Acta* 40:534. [40] Najjar, 208:225. [41] Neurath, H., and G. W. Schwert. 1950. *Chem. Rev.* 46:49. [42] Noda, L. 1958. *J. Biol. Chem.* [43] Noda, L., S. A. Kuby, and H. A. Lardy. 1954. *Ibid.* 209:203. [44] Noltmann, E. A. 1963. *Federation Proc.* and S. A. Kuby. 1961. *J. Biol. Chem.* 236:1225. [45] Northrop, J. H. 1922. *J. Gen. Physiol.* 5:263. [46] Olson, 233:365. [47] Peanasky, R. J., and H. A. Lardy. 1958. *Ibid.* 233:371. [48] Perlmann, G. E. 1955. *Advan. Protein H. Neurath. 1946. J. Biol. Chem.* 166:603. [49] Racker, E. 1950. *Ibid.* 184:313. [50] Rodwell, V. W., J. C. 46:155. [51] Samejima, T., and K. Shibata. 1961. *Arch. Biochem. Biophys.* 93:407. [52] Schachman, H. K.

**ENZYMES**

**KINETIC PROPERTIES**

Kinetic Properties								Reference
Molecular Activity			Michaelis Constant					
Substrate	Temp. °C	pH	Value <sup>4</sup>	Substrate	Temp. °C	pH	K <sub>m</sub> <sup>5</sup> moles/liter	
(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)
Benzoyl-L-arginine ester	25	7.7	1,600	Benzoyl-L-arginine ester	25	8.0	8 x 10 <sup>-5</sup>	B,62;C,24;D-F,23;G,20;H-O,81
3,4-Dihydroxy-L-phenylalanine	30	6.0	19,600	Tyrosine	30	6.0	8 x 10 <sup>-4</sup>	B-F,H-K,35;G,124;L-O,47
Urea	20	7.0	460,000	Urea	21	8.0	4 x 10 <sup>-3</sup>	B,117;C-E,121;F,122;G,L-O,139;H-K,123
				Xanthine			ca. 10 <sup>-6</sup>	B,3,56;C-F,4;G,9;L, P,13

*national Union of Biochemistry*, 1961. Pergamon Press, New York. /s/ Values for sedimentation (S<sub>20,w</sub>) and zero protein concentration. /s/ Apparent values determined from electrophoretic mobility measurements. /4/ Mzyme at optimal substrate concentration. Values pertain to the molecular weight given in column E. /s/ K<sub>m</sub> K<sub>-1</sub> = velocity constant for the dissociation of this complex into substrate and enzyme, and K<sub>+2</sub> = velocity constant

1937. *J. Gen. Physiol.* 20:663. [3] Avis, P. G., et al. 1954. *Nature* 173:1230. [4] Avis, P. G., et al. 1956. [6] Baranowski, T. 1949. *Ibid.* 180:535. [7] Baranowski, T., and T. R. Niederland. 1949. *Ibid.* 180:543. 223:765. [10] Bergmeyer, H. U., et al. 1961. *Biochem. Z.* 333:471. [11] Bonnichsen, R. K. 1950. *Acta Chem. Press, New York.* [13] Bray, R. C. 1959. *Biochem. J.* 73:690. [14] Bücher, T. 1947. *Biochim. Biophys. Acta* 1951. *Biochem. J.* 49:105. [17] Cecil, R., and A. G. Ogston. 1952. *Ibid.* 51:494. [18] Chance, B. 1949. *Arch. Van Nostrand, Princeton.* p. 383. [20] Colowick, S. P., and N. O. Kaplan, ed. 1955-62. *Methods in enzymology.* [22] Cori, G. T., M. W. Slein, and C. F. Cori. 1948. *Ibid.* 173:605. [23] Cunningham, L. W., Jr. 1954. *Ibid.* *Chem. Scand.* 12:459. [26] Datta, A. G., and E. Racker. 1961. *J. Biol. Chem.* 236:617. [27] De la Haba, G., I. G. *Biophys.* 69:555. [29] Edelhoch, H. 1957. *J. Am. Chem. Soc.* 79:6100. [30] Ehrenberg, A. 1957. *Acta Chem.* 1950. *J. Biol. Chem.* 187:213. [33] Fellig, J., E. A. Stein, and E. H. Fischer. 1957. *Helv. Chim. Acta* 40:529. Horowitz, and S. F. Heinemann. 1963. *J. Biol. Chem.* 238:2045. [36] Frieden, C., and R. A. Alberty. 1955. *Ibid.* R. G. Wolfe, Jr., and R. A. Alberty. 1957. *Ibid.* 79:1523. [39] Fromm, H. J., and V. Zewe. 1962. *J. Biol. Chem.* L., and D. H. Brown. 1955. *J. Biol. Chem.* 216:67. [42] Green, A. A. 1945. *Ibid.* 158:315. [43] Green, A. A., and J. E., Jr., and S. F. Velick. 1954. *J. Biol. Chem.* 207:225. [46] Herriott, R. M. 1938. *J. Gen. Physiol.* 21:501. 1959. *Arch. Biochem. Biophys.* 81:500. [49] Johnson, P., and V. Massey. 1957. *Biochim. Biophys. Acta* 23:544. P. J., and G. T. Cori. 1953. *Biochim. Biophys. Acta* 12:235. [53] Keller, P. J., C. Lowry, and J. F. Taylor. 1956. *Chem. Scand.* 13:1149. [56] Klenow, H., and R. Emberland. 1955. *Arch. Biochem. Biophys.* 58:276. [57] Kuby, *Ibid.* 210:65. [59] Kunitz, M. 1940. *J. Gen. Physiol.* 24:15. [60] Kunitz, M. 1952. *Ibid.* 35:423. [61] Kunitz, M., 1960. *Ann. Rept. Sci. Works Fac. Sci. Osaka Univ.* 8:43. [64] Kusai, K., et al. 1960. *Biochim. Biophys. Acta* Y.-P. 1957. *J. Biol. Chem.* 227:987. [67] Lee, Y.-P. 1957. *Ibid.* 227:993. [68] Lee, Y.-P. 1957. *Ibid.* 227:999. J. R., and M. F. Utter. 1959. *J. Biol. Chem.* 234:2151. [72] Mahowald, T. A., E. A. Noltmann, and S. A. Kuby. *Biochem. J.* 51:490. [75] Matsuo, Y., and D. M. Greenberg. 1958. *J. Biol. Chem.* 230:545. [76] Mehler, A. H. V. A. 1948. *J. Biol. Chem.* 175:281. [79] Neilands, J. B. 1952. *Ibid.* 199:373. [80] Neilands, J. B. 1954. *Ibid.* 232:237. [83] Noda, L., and S. A. Kuby. 1957. *Ibid.* 226:541. [84] Noda, L., and S. A. Kuby. 1957. *Ibid.* 226:551. 22:411. [87] Noltmann, E. A. Unpublished, Univ. California, Riverside, 1963. [88] Noltmann, E. A., C. J. Gubler, J. A., and C. B. Anfinsen. 1952. *J. Biol. Chem.* 197:67. [91] Peanasky, R. J., and H. A. Lardy. 1958. *Ibid.* *Chem.* 10:23. [94] Posternak, T., and J. P. Rossetlet. 1954. *Helv. Chim. Acta* 37:246. [95] Putnam, F. W., and Towne, and S. Grisolia. 1956. *Biochim. Biophys. Acta* 20:394. [98] Samejima, T. 1959. *J. Biochem. (Tokyo)* 1952. *J. Gen. Physiol.* 35:451. [101] Scheraga, H. A., and J. A. Rupley. 1962. *Advan. Enzymol.* 24:161.

*continued*

Part II. PHYSICAL AND

[102] Schwert, G. W., and S. Kaufman. 1951. J. Biol. Chem. 190:807. [103] Shavit, N., R. G. Wolfe, and R. A. D. M. Brown, and H. T. Hanson. 1949. Ibid. 180:33. [106] Smith, E. L., and H. T. Hanson. 1949. Ibid. 176:997. 1958. Ibid. 233:1387. [109] Smith, E. L., and A. Stockell. 1954. Ibid. 207:501. [110] Snoko, J. E., and H. Neurath. Del Campillo, and I. Raw. 1956. J. Biol. Chem. 218:971. [113] Stockell, A., and E. L. Smith. 1957. Ibid. 227:1. [116] Strecker, H. J. 1953. Ibid. 46:128. [117] Sumner, J. B. 1926. J. Biol. Chem. 69:435. [118] Sumner, J. B., 136:343. [120] Sumner, J. B., and N. Gralén. 1938. Ibid. 125:33. [121] Sumner, J. B., N. Gralén, and I.-B. [123] Sumner, J. B., and G. F. Somers. 1953. Chemistry and methods of enzymes. Ed. 3. Academic Press, Hauenstein. 1956. J. Am. Chem. Soc. 78:5287. [126] Taylor, J. F., A. A. Green, and G. T. Cori. 1948. J. Biol. 1935. Biochem. Z. 278:263. [129] Theorell, H. 1942. Enzymologia 10:250. [130] Theorell, H., and Å. Åkeson. 65:439. [132] Theorell, H., A. P. Nygaard, and R. Bonnichsen. 1955. Acta Chem. Scand. 9:1148. [133] Tietz, A., 14:488. [135] Van Eys, J., B. J. Nuenke, and M. K. Patterson, Jr. 1959. J. Biol. Chem. 234:2308. [136] Velick, [138] Wakil, S. J., and H. R. Mahler. 1954. Ibid. 207:125. [139] Wall, M. C., and K. J. Laidler. 1953. Arch. Christian, 1941-42. Biochem. Z. 310:384. [142] Warner, R. C. 1958. Arch. Biochem. Biophys. 78:494. 1960. J. Biol. Chem. 235:2013. [145] Wellner, D., and A. Meister. 1961. Ibid. 236:2357. [146] Winer, A. D., H. L., and N. Pace. 1958. Arch. Biochem. Biophys. 75:125.

Part III. CHEMICAL

Data are for crystalline or electrophoretically homogeneous enzymes.

	Enzyme	Source	Elements						
			Carbon	Hydrogen	Nitrogen	Sulfur	Phosphorus	Other	Amino N
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
1	Alcohol dehydrogenase	Yeast	52.8	6.96	16.53	1.21	.....	Zn, 0.218	....
2	Aldolase	Rabbit muscle	.....	.....	16.8	.....	.....	.....	.....
3	α-Amylase	Swine pancreas	49.46	7.18	15.52	1.33	0.05	.....	.....
4	β-Amylase	Sweet potato	.....	.....	15.1	.....	.....	.....	0.83
5	Carbonic anhydrase	Mammalian red cells	.....	.....	16.1	0.34	.....	Zn, 0.21	.....
6	Carboxypeptidase	Cattle pancreas	52.6	7.2	14.4	0.47	0	Zn, 0.182	.....
7	Catalase	Horse blood	.....	.....	16.8	.....	.....	Fe, 0.093	.....
8		Horse liver	.....	.....	16.8	.....	.....	Fe, 0.093	.....
9	α-Chymotrypsin	.....	50.0	7.06	15.5	1.85	0	Cl, 0.16	1.22
10	β-Chymotrypsin	.....	.....	.....	16.24	1.56	.....	.....	1.31
11	γ-Chymotrypsin	.....	.....	.....	16.00	1.59	.....	.....	1.34
12	Chymotrypsinogen	Cattle pancreas	50.6	7.0	15.8	1.9	0	Cl, 0.17	0.97
13	Chymotrypsinogen A	.....	.....	.....	.....	.....	.....	.....	.....
14	Cytochrome-c <sup>4</sup>	Cattle heart	52.52	7.76	15.36	1.47	0	Fe, 0.43	.....
15	Deoxyribonuclease	Cattle pancreas	50.16	6.91	14.88	1.09	0	.....	.....
16	Enolase	.....	53.62	7.55	17.34	0.38	0	.....	.....
17	D-Glyceraldehyde phosphate dehydrogenase <sup>5</sup>	Rabbit muscle	.....	.....	.....	.....	.....	.....	.....
18		Yeast	52.54	7.51	16.41	1.08	Trace	.....	.....
19	Hexokinase	Yeast	52.16	7.08	15.62	0.91	0.11	.....	.....
20	Lecithinase	.....	50.77	6.41	15.88	4.0	.....	.....	.....
21	Lipoxidase	.....	.....	.....	.....	.....	.....	.....	.....
22	Lysozyme <sup>5</sup>	.....	.....	.....	18.6	2.53	.....	.....	0.74
23	Old yellow enzyme	Yeast	51.4	7.07	16.27	0.48	0.043	.....	.....
24	Papain <sup>5</sup>	.....	.....	.....	16.1	1.2	0	.....	.....
25	Pepsin <sup>5</sup>	Cattle	51.7	6.86	14.6	0.94	0.09	.....	0.162

<sup>1/1</sup> Determined after oxidation to cysteic acid; one molecule of cysteic acid is formed from one-half of the sym- between cysteine and cystine, unless otherwise indicated. <sup>2/2</sup> Cysteine plus cystine. <sup>3/3</sup> Cystine. <sup>4/4</sup> Values for given as number of amino acid residues per molecule of enzyme.

**ENZYMES**

**KINETIC PROPERTIES**

Alberty. 1958. Ibid. 233:1382. [104] Smith, E. L., and D. M. Brown. 1952. Ibid. 195:525. [105] Smith, E. L., [107] Smith, E. L., J. R. Kimmel, and D. M. Brown. 1954. Ibid. 207:533. [108] Smith, E. L., and M. J. Parker. 1949. Ibid. 181:789. [111] Srere, P., et al. 1958. Arch. Biochem. Biophys. 74:295. [112] Stern, J. R., A. [114] Straub, F. B. 1940. Biochem. J. 34:483. [115] Strecker, H. J. 1951. Arch. Biochem. Biophys. 32:448. and A. L. Dounce: 1937. Ibid. 121:417. [119] Sumner, J. B., A. L. Dounce, and V. L. Frampton. 1940. Ibid. Eriksson-Quensel. 1938. Ibid. 125:37. [122] Sumner, J. B., and D. B. Hand. 1929. J. Am. Chem. Soc. 51:1255. New York. p. 161. [124] Sussman, A. S. 1961. Arch. Biochem. Biophys. 95:407. [125] Tanford, C., and J. D. Chem. 173:591. [127] Taylor, J. F., and C. Lowry. 1956. Biochim. Biophys. Acta 20:109. [128] Theorell, H. 1943. Arkiv Kemi Mineral. Geol. 17B(7). [131] Theorell, H., and Å. Åkeson. 1956. Arch. Biochem. Biophys. and S. Ochoa. 1958. Arch. Biochem. Biophys. 78:477. [134] Tung, T.-C., et al. 1954. Biochim. Biophys. Acta. S. F. 1949. J. Phys. Colloid Chem. 53:135. [137] Velick, S. F., and J. E. Hayes, Jr. 1953. J. Biol. Chem. 203:545. Biochem. Biophys. 43:299. [140] Wallenfels, K., et al. 1959. Biochem. Z. 331:459. [141] Warburg, O., and W. [143] Weibull, C., and A. Tiselius. 1945. Arkiv Kemi Mineral. Geol. 19A:19. [144] Wellner, D., and A. Meister. and G. W. Schwert. 1958. Ibid. 231:1065. [147] Wold, F., and C. E. Ballou. 1957. Ibid. 227:313. [148] Young,

**COMPOSITION**

Values are grams per 100 grams enzyme, unless otherwise indicated.

Amino Acids																			Refer- ence		
Alanine	Arginine	Aspartic Acid	Cysteine	Cystine <sup>1</sup> / <sub>2</sub>	Glutamic Acid	Glycine	Histidine	Isoleucine	Leucine	Lysine	Methionine	Phenylalanine	Proline	Serine	Threonine	Tryptophan	Tyrosine	Valine			
(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	(W)	(X)	(Y)	(Z)	(A')	(B')	(C')		
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	5	1	
8.56	6.33	9.7	...	1.12	11.4	5.61	4.21	7.87	11.5	9.54	1.17	3.06	5.71	6.57	7.1	2.31	5.31	7.40	5,31	2	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	6	3
...	6.0	...	...	0.79 <sup>a</sup>	...	...	...	...	...	...	4.32	...	...	...	...	...	7.0	...	2	4	
...	...	...	...	1.3	...	...	...	...	...	...	...	...	...	...	...	...	4.1	...	19,24	5	
5.16	5.06	11.7	...	1.40	10.7	5.06	3.47	7.65	9.41	7.81	0.44	7.16	3.66	10.1	9.21	3.62	10.3	5.58	1,26,30	6	
...	8.75	16.5	...	1.65 <sup>a</sup>	10.9	...	4.17	...	...	7.50	...	...	...	...	...	...	6.0	...	3	7	
...	8.90	16.5	...	1.85 <sup>a</sup>	10.3	...	3.86	...	...	6.91	...	...	...	...	...	...	5.8	...	3	8	
...	...	...	1.22	3.66	...	...	1.26	...	9.1	...	1.25	...	...	...	11.2	5.81	2.83	...	21	9	
...	...	...	1.29	3.51	...	...	1.22	...	9.4	...	1.29	...	...	...	10.6	6.40	2.87	...	21	10	
...	...	...	1.27	3.59	...	...	1.26	...	8.5	...	1.28	...	...	...	10.7	6.27	3.09	...	21	11	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	21	12	
22	4	8	...	10	3	23	2	10	19	13	2	6-7	9	30	23	7	4	22	4	13	
...	5.6	...	...	1.4	...	...	6.3	...	...	30.8	...	...	...	...	...	1.4	3.5	...	28	14	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	15	15	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	33	16	
8.5	5.2	13.9	1.1	...	8.2	7.0	4.5	6.4	6.9	10.3	3.7	6.3	4.1	6.1	7.6	2.1	4.6	10.8	5	17	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	22	18	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	17	19	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	25	20	
...	4.7	6.2	0	0	10.4	6.3	3.6	8.1	11.4	7.8	1.8	4.9	5.1	...	8.9	0.41	6.2	7.8	12	21	
12	11	21	...	8	5	12	1	6	8	6	2	3	2	10	7	6	3	6	7,9	22	
...	8.25	...	...	0.34	...	7.1	2.75	...	...	13.7	...	5.75	...	...	...	4.86	7.75	...	14	23	
13	10	17	...	6	17	23	2	10	10	9	...	4	9	11	7	5	17	15	4	24	
18	2	44	...	6	27	38	1	27	28	1	5	14	15	44	28	6	18	21	4,21	25	

metrical cystine molecule or from one molecule of cysteine. The values in this column, therefore, do not distinguish amino acids (columns J-B') are given as percent of total nitrogen. /5/ Values for amino acids (columns J-B') are

*continued*



Enzyme	Source	Elements						
		Carbon	Hydrogen	Nitrogen	Sulfur	Phosphorus	Other	Amino N
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
26 Pepsin <sup>b</sup>	Salmon	51.9	6.48	15.62	1.58	0.03	.....	....
27 Pepsinogen	.....	52.8	6.88	15.9	0.09	.....	.....	....
28 Peroxidase	Horseradish	47.0	7.35	13.2	0.43	.....	Fe, 0.13	....
29 Phospho-enoltransphosphorylase	Man	53.35	7.30	17.40	1.60	0.06	.....	....
30 Phosphorylase	Rabbit muscle	.....	.....	16.5	.....	0.027	.....	....
31 Pyrophosphatase	Yeast	54.46	7.36	16.25	0.14	0	.....	0.86
32 Rennin	.....	51.4	7.19	14.51	1.46	0.04	Cu, 0.0035	1.11
33 Ribonuclease <sup>c</sup>	.....	48.2	6.2	16.1	1.1	Trace	.....	....
34 Trypsin	.....	50.2	6.6	16.13	1.1	0	Cl, 2.85	....
35 Trypsinogen	.....	50.1	6.9	15.3	1.1	.....	.....	....
36 Urease	Jack bean	51.6	7.1	16.0	1.2	.....	Cu, 0.001	....

/1/ Determined after oxidation to cysteic acid; one molecule of cysteic acid is formed from one-half of the sym- between cysteine and cystine, unless otherwise indicated. /s/ Values for amino acids (columns J-B') are given as

Contributors: (a) Tupper, Ronald, and Watts, R. W. E., (b) Dianzani, Mario U., (c) Corley, Ralph C.

References: [1] Anson, M. L. 1937. J. Gen. Physiol. 20:663. [2] Balls, A. K., M. K. Walden, and R. R. Thompson. K. Myrbäck, ed. 1960. The enzymes. Ed. 2. Academic Press, New York. v. 4. [5] Boyer, P. D., H. Lardy, and R. E. 1963. J. Biol. Chem. 238:2691. [8] Cori, C. F., G. T. Cori, and A. A. Green. 1943. Ibid. 151:39. [9] Fevold, 1952. J. Am. Chem. Soc. 74:3181. [12] Holman, R. T., et al. 1950. Arch. Biochem. 26:199. [13] Kubowitz, F., [15] Kunitz, M. 1950. J. Gen. Physiol. 33:349. [16] Kunitz, M. 1952. Ibid. 35:423. [17] Kunitz, M., and M. R. Biochim. Biophys. Acta 39:218. [20] Norris, E. R., and D. W. Elam. 1940. J. Biol. Chem. 134:443. [21] Northrop, [22] Rafter, G. W., and G. E. Krebs. 1950. Arch. Biochem. 29:233. [23] Scheraga, H. A., and J. A. Rupley. K. H., and H. L. Fraenkel-Conrat. 1938. Ber. Deut. Chem. Ges. 71B:1076. [26] Smith, E. L., and A. Stockell. and Å. Åkeson. 1941. J. Am. Chem. Soc. 63:1804. [29] Theorell, H., and Å. Åkeson. 1943. Arkiv Kemi Mineral. E. Ronzoni. 1948. J. Biol. Chem. 173:620. [32] Velick, S. F., and L. F. Wicks. 1951. Ibid. 190:741. [33] Warburg,

90. HORMONES:

Data are for man, unless otherwise specified. **Properties** (column B): MW = molecular weight; MP = melting point; insoluble; s. = soluble; sl. = slightly; v. = very. **Symbols** (columns D, G-I): † = increased; ‡ = decreased.

Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
Adenohypophysis				
1 Adrenocorticotropin; adrenotropin, adreno- corticotropic hor- mone, ACTH, cortico- tropin	MW = 4,520 (cattle); amino acid no. = 39; one peptide chain, unbranched; IEP = 4.7-4.8; S <sub>20,w</sub> = 2.08; s. water, 60-70% alcohol	Adenohypophysis; synthetic trico- sapeptide has full activity	Adrenal repair, wt mainte- nance (hypophysectomized rat); ‡ adrenal ascorbic acid (hypophysectomized rat); urinary excretion of corti- coids (guinea pig)	Not defi- nitely known; probably similar to other polypep- tides; half-life of ACTH in blood is ‡ 3-5 min

**ENZYMES**

**COMPOSITION**

Amino Acids																			Refer- ence	
Alanine	Arginine	Aspartic Acid	Cysteine	Cystine $\frac{1}{2}$	Glutamic Acid	Glycine	Histidine	Isoleucine	Leucine	Lysine	Methionine	Phenylalanine	Proline	Serine	Threonine	Tryptophan	Tyrosine	Valine		
(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	(W)	(X)	(Y)	(Z)	(A')	(B')	(C')	
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	20	26
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	21	27
...	6.91	...	...	...	...	...	0.71	...	...	4.06	...	...	...	...	...	...	...	...	29	28
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	13	29
4.79	11.6	9.3	...	0.45	13.4	3.8	3.3	6.5	10.5	7.2	2.7	6.2	4.7	3.05	4.24	2.0	5.9	7.3	8,32	30
4.9	3.3	12.1	...	...	9.7	2.9	2.2	8.9	6.0	10.9	1.3	6.2	6.4	3.1	4.8	3.6	6.0	4.1	11,16	31
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	10	32
12	4	5	...	8	5	3	4	3	2	10	4	3	4	15	10	...	6	9	21,23	33
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	3.65	7.8	...	18,21	34
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	18	35
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	27	36

metrical cystine molecule or from one molecule of cysteine. The values in this column, therefore, do not distinguish number of amino acid residues per molecule of enzyme.

1948. J. Biol. Chem. 173:9. [3] Bonnichsen, R. K. 1947. Arch. Biochem. 12:83. [4] Boyer, P. D., H. Lardy, and K. Myrbäck, ed. 1963. Ibid. v. 7. [6] Caldwell, M. L., et al. 1952. J. Am. Chem. Soc. 74:4033. [7] Canfield, H. L. 1951. Advan. Protein Chem. 6:230. [10] Hankinson, C. L. 1943. J. Dairy Sci. 26:53. [11] Hausmann, W. and P. Ott. 1944. Biochem. Z. 317:193. [14] Kuhn, R., and P. Desnuelle. 1937. Ber. Deut. Chem. Ges. 70B:1907. McDonald. 1946. Ibid. 29:393. [18] Kunitz, M., and J. H. Northrop. 1936. Ibid. 19:991. [19] Linskog, S. 1960. J. H., M. Kunitz, and R. M. Herriott. 1948. Crystalline enzymes. Columbia Univ. Press, New York. p. 26. 1961. Advan. Enzymol. 24:175. [24] Scott, D. A., and A. M. Fisher. 1942. J. Biol. Chem. 144:371. [25] Slotta, 1954. J. Biol. Chem. 207:501. [27] Sumner, J. B. 1930. Ber. Deut. Chem. Ges. 63B:582. [28] Theorell, H., and Geol. 16A(8). [30] Vallee, B. L., and H. Neurath. 1954. J. Am. Chem. Soc. 76:5006. [31] Velick, S. F., and O., and W. Christian. 1942. Biochem. Z. 310:384.

**VERTEBRATES**

IEP = isoelectric point;  $S_{20,w}$  = sedimentation coefficient;  $D_{20,w}$  = diffusion coefficient;  $[a]_D$  = refractive index; i =

Targets	Principal Effects	Effect of Deficiency (-) Excess (+)	Secretion Inhibited by (I) Stimulated by (S)	Refer- ence
(F)	(G)	(H)	(I)	(J)
Adenohypophysis				
Adrenal cortex; mitochondria and microsomes in most tissues; melanocytes	lipid and ascorbic acid content of adrenal cortex; ↓ secretion of adrenal cortical hormones; ↑ oxidative phosphorylation, protein synthesis, glycolysis, synthesis of steroid hormones; ↑ fat transport and oxidation, muscle glycogen, amino acid metabolism	(-): adrenal cortex atrophy and hypofunction; ↓ response to stress (+): adrenal cortex hyperfunction	(I) blood level of cortisol; servo or feedback mechanism (S) stress, acting thru hypothalamus, plus corticotropin-releasing factors from adenohypophysis	3,7,21, 22,24, 26,27

*continued*

Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
<b>Adenohypophysis</b>				
2 Follicle-stimulating hormone; FSH	MW = 29,000-30,000; IEP = 5.1-5.2; stable pH 7-8, 30 min at 75°C; s. water, 50% acetone, 70% alcohol, 50% dioxane, $\frac{1}{2}$ -saturated $(\text{NH}_4)_2\text{SO}_4$	Adenohypophysis (especially sheep, swine); urine (castrates); serum (pregnant mare)	Follicular growth (hypophysectomized rat); $\uparrow$ wt of ovary and uterus (hypophysectomized, immature $\text{♀}$ rat); $\uparrow$ uterine wt or hyperemia (immature mouse)	FSH-like protein
3 Growth hormone; GH, somatotropin	MW = 27,000 <sup>1</sup> ; IEP = 4.9 <sup>1</sup> ; $D_{20,w} = 7.15 \times 10^{-7}$ ; viscosity coefficient = 7.64; s. salt solution; sl. s. water; $\text{HNO}_2$ and acetylation destroy	Adenohypophysis (various species); plasma in children, 85-570 mg GH per ml	$\uparrow$ tail length, tibial epiphyseal cartilage (hypophysectomized rat); body growth (hypophysectomized rat); inhibition of hemagglutination of erythrocytes sensitized to tannic acid and titrated with human GH; radioactivity of $^{131}\text{I}$ -labeled GH unbound to anti-GH immune serum	Unknown; probably similar to other polypeptides
4 Luteinizing hormone; LH, interstitial-cell-stimulating hormone, ICSH	Sheep: MW = 30,000, IEP = 4.6, $S_{20,w} = 3.6 \times 10^{-13}$ ; swine: IEP = 7.45, $S_{20,w} = 6.8 \times 10^{-13}$ ; s. water, 40% alcohol, dilute salt solution	Adenohypophysis (especially sheep, swine)	Interstitial cell repair (hypophysectomized mouse, rat); regeneration of breast feathers (weaver finch); $\uparrow$ wt of ventral prostate (hypophysectomized rat); $\uparrow$ testes wt (pigeon); $\downarrow$ ascorbic acid in ovaries (rat)	
5 Lactogenic hormone; LTH, lutetropin, galactin, prolactin	MW = 24,100-26,000; IEP = 5.5-5.73; $D_{20,w} = 9 \times 10^{-7}$ ; viscosity coefficient = 6.65; $[\alpha]_D^{25} = -40.5^\circ$ ; s. acid alcohol, methyl alcohol; sl. s. water, dilute salt solution; one peptide chain	Adenohypophysis (especially cattle, sheep); urine (in small amounts)	$\uparrow$ crop sac wt; crop gland proliferation (pigeon); $\uparrow$ milk secretion (rabbit)	
6 Thyrotropic hormone <sup>2</sup> ; TSH, thyrotropin	MW = 24,100-26,000; inactivated by boiling, cysteine, ketene, trypsin, pepsin, and chymotrypsin; nondiffusible; one peptide chain	Probably adenohypophysis (basophilic cells)	Uptake of $^{131}\text{I}$ by thyroid; $\downarrow$ thyroid iodine (chick); $\uparrow$ thyroid cell ht (chick); $\uparrow$ thyroid gland wt (guinea pig); $\uparrow$ swelling of thyroid slices	
<b>Neurohypophysis</b>				
7 Oxytocin; oxytocic hormone, pitocin, postlobin-O, posterior-lobe principle	IEP = 7.7; not adsorbed on charcoal; destroyed by acid, trypsin, tyrosinase; s. water, concentrated acetic acid, methyl alcohol. Amino acid comp. (cattle, swine): Cys, Tyr, Isoleu, Glu(NH <sub>2</sub> ), Asp(NH <sub>2</sub> ), Cys, Pro, Leu, Gly(NH <sub>2</sub> ).	Neurohypophysis	Contraction, isolated uterus (guinea pig); $\uparrow$ milk ejection (rabbit); $\downarrow$ blood pressure (chicken)	

<sup>1</sup>/ Values for man. Other values: simian, MW = 25,400, IEP = 5.5; cattle, MW = 45,000, IEP = 6.85; sheep, MW = lengthen. <sup>2</sup>/ Evidence for two fractions--one to stimulate hypertrophy of acinar cells and colloid secretion from factor" that can be separated from TSH.

## VERTEBRATES

Targets (F)	Principal Effects (G)	Effect of Deficiency (-) Excess (+) (H)	Secretion Inhibited by (I) Stimulated by (S) (I)	Refer- ence (J)
<b>Adenohypophysis</b>				
Ovarian follicles; seminiferous tu- bules	↑ spermatogenesis and growth of seminiferous tubules, follicle development (no ova production or estrogen secre- tion unless LH present)	(-): obesity, atrophy or im- maturity of gonads; no matu- ration of ova or sperm; ↓ li- bido and potency, hair; ↑ H <sub>2</sub> O metabolism (+): growth of follicles (may be numerous); estrogen secre- tion (FSH + LH), enlargement of tubules	(I) circulating es- trogens and possi- bly androgens (S) castration; menopause; low blood levels of es- trogens or possi- bly androgens; hy- pothalamo-pitui- tary apparatus	3,7,21, 22,24- 28,31 2
Bones, especially epiphyseal cartilage; fibroblasts, most tissues	↑ skeletal and soft-tissue growth, protein anabolism, fibroblastic activity; ↓ pan- creatic insulin (↑ in rats), maintenance of muscle growth	(-): dwarfism and/or infanti- lism; delayed closure of epi- physes (+): gigantism and/or acro- megaly; hypertrophy of vis- cera <sup>2</sup>	(I) estrogens or an- drogens (large doses)	3,7,21, 24-26, 28,31 3
Maturing graafian fol- licles and intersti- tial cells (ovaries); interstitial cells of Leydig (testes)	↑ follicle maturation (produc- tion but not maintenance of corpus luteum), estrogen secretion, androgen secre- tion from testes synergis- tic with FSH in stimulating estrogen and ovulation	(-): lack of ovulation and lutei- nization; little or no estro- gen or androgen secretion; atrophy of interstitial tissue in ovary and testis (+): ovulation and luteiniza- tion of prepared follicles; hyper- trophy of Leydig tissue; ↑ es- trogen secretion (with FSH), androgen secretion	(I) high levels of ovarian and testic- ular hormones (S) intermediate levels of ovarian hormones; hypo- thalamo-pitui- tary apparatus	3,7,21, 22,24, 26-28, 31 4
Mammary glands; corpus luteum; crop sac (pigeon)	↑ milk secretion, progester- one secretion by developed corpus luteum, uterine ni- dation and decidua; ↑ crop gland growth and secretion (pigeon)	(-): lactation failure; deficient secretion of progesterone by corpus luteum (+): lactation initiation; main- tenance of corpus luteum and progesterone secretion; re- lease of estrogen and pro- gesterone by luteal tissue; crop gland development (pigeon)	(I) or (S) effects of androgens, FSH, LH, and progester- one on target organs (S) by oxytocin, in turn (S) by suck- ling	3,7,21, 22,24, 26-28, 31 5
Glands of orbit (?); thyroid gland	↑ thyroid hormone synthesis and secretion, cell height of thyroid epithelium, thyroid size, serum protein-bound iodine; ↓ iodine and colloid content of thyroid; stimu- lates proliferation of con- nective tissue?	(-): myxedema; cretinism (some forms); ↓ basal meta- bolic rate (+): goiter; hyperthyroidism; exophthalmos <sup>4</sup> ; ↑ basal meta- bolic rate	(I) ↑ circulating thy- roxine (I) or (S) nervous system (S) ↓ circulating thyroxine	3,7,19, 21,22, 24,26, 28,31 6
<b>Neurohypophysis</b>				
Uterine and other smooth muscles; mammary glands	↑ uterine muscle contraction (and other smooth muscles to a lesser degree); facili- tates movement of sperm up fallopian tubes; initiates labor; may stimulate re- lease of prolactin	(-): delayed uterine contrac- tion (pre- or post-partum); ↓ milk flow (+): uterine contraction (es- pecially if prepared by es- trogen); ↑ milk flow	(S) suckling	3,7,22, 24,27, 28,31 7

48,000, IEP = 6.8; whale, MW = 39,900, IEP = 6.2. /2/ More characteristic of acromegaly, since long bones do not gland, and the other to accelerate uptake from blood and hormone synthesis. /4/ May be caused by "exophthalmos

*continued*



Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
<b>Neurohypophysis</b>				
8 Vasopressin; vaso-pressor principle, postlobin-V, vasopressor-antidiuretic principle, ADH	IEP = 10.8; adsorbed on charcoal; inactivated by trypsin, not pepsin; solubility same as for oxytocin. Amino acid comp. (human): Cys, Tyr, Phe, Glu(NH <sub>2</sub> ), Asp(NH <sub>2</sub> ), Cys, Pro, Arg, Gly(NH <sub>2</sub> ).	Neurohypophysis	↑ blood pressure (dog, rat); antidiuretic activity (dog, rat, excreting 5 ml urine/min)	ADH-like material
<b>Pars Intermedia</b>				
9 Melanocyte-stimulating hormones; MSH, melanotropins, intermedin, melanophore hormone, chromatophorotropic hormone	IEP = 4.1; dialyzable; moderately heat stable; destroyed by trypsin; stable acid, alkali; s. water; i. ether, acetone	Intermediate lobe (many species, including mammals); anterior lobe (birds porpoises); chemical synthesis	In vitro: Intensity of darkening of isolated section of skin at end of 60 min ( <i>Rana pipiens</i> ). Changes in reflectance correlate with amount of MSH in solution.	
α-MSH <sup>b</sup>	Polypeptide consisting of 13 amino acid residues; IEP = 10.5-11; same residues and sequence occur in ACTH			
β-MSH <sup>a</sup>	18 amino acid residues; 6 are same in kind and sequence as in α-MSH; acidic; IEP = approx. 4.1. Some interspecies differences. Both α- and β-MSH dialyzable, moderately heat stable, destroyed by trypsin; heating 10 min at 100° with 0.1 N NaOH racemizes MSH, resulting in potentiation and prolongation of darkening effect, dilution of acid; s. water; i. ether, acetone.			
<b>Pineal Body</b>				
10 Adrenoglomerulotropin; GTH; (1-methyl-6-methoxy-1, 2, 3, 4-tetrahydro-2-carboline ?)	s. alcohol, CHCl <sub>3</sub> , hexane; lipid; nonpolar; pale tan with Le Rosen reagent	Pineal body or adjacent tissues	Stimulation of aldosterone secretion (in pinealectomized, partially decerebrate dog)	Unknown
11 Melatonin; C <sub>13</sub> H <sub>16</sub> O <sub>2</sub> N <sub>2</sub> ; (N-acetyl-5-methoxytryptamine)	MW = 222; max. absorption 2,780°; s. aqueous, organic solvents; blue color with Ehrlich reagent	Pineal body (mammals); isolated from pineals (man, cattle, monkey); smaller amounts in central and peripheral nerves; in vivo precursor is serotonin	Degree of lightening of isolated sections of skin of amphibians ( <i>Rana pipiens</i> )	5-Methoxytryptamine and 5-methoxyindoleacetic acid isolated from pineal tissue; 6-hydroxymelatonin isolated from urine (man, rat)
<b>Thyroid</b>				
12 Thyroglobulin	MW = 680,000; amino acid comp.: His, Lys, <i>p</i> -Aminotryp, Tryp, Tyr, Diiodoty, Thyrox, Cys, Meth, Ala, Glyc, Leu, Val, Ser, Monoiodoty, Iodothyrox	Thyroid gland (follicles)	Limb bud growth (amphibian larva); growth (thyroidectomized rat); ↑ basal metabolic rate (thyroid-deficient subjects); ↑ O <sub>2</sub> consumption; oxidation of succinate; prevention of goiter; survival in anoxia (rat)	Same as thyroxine and proteins; released from gland by proteases and hyaluronidase

<sup>b</sup>/ Amino acid composition: Ac-Ser, Tyr, Ser, Met, Glu, His, Phe, Arg, Try, Gly, Lys, Pro, Val (NH<sub>2</sub>). <sup>a</sup>/ Amino Lys, Asp.

# Contrails

## VERTEBRATES

Targets	Principal Effects	Effect of Deficiency (-) Excess (+)	Secretion Inhibited by (I) Stimulated by (S)	Reference
(F)	(G)	(H)	(I)	(J)
Neurohypophysis				
Capillaries; arterioles; coronary vessels; kidney vascular bed and tubules; smooth muscles	kidney excretion of H <sub>2</sub> O; ↑ NaCl and urea excretion, blood pressure (constriction of capillaries); promotes renal tubular water reabsorption	(-): diabetes insipidus, diuresis; ↓ NaCl and urea excretion (+): smooth muscle contraction; ↑ H <sub>2</sub> O retention, blood pressure	(I) ↓ plasma osmotic concentration; ↑ extracellular volume (S) ↑ plasma osmotic concentration; ↓ extracellular volume	3,7,22, 24,27, 28
Pars Intermedia				
Chromatophore cells in skin of lower vertebrates (reptiles, amphibians, fishes); adipokinetic effect may involve liver	Rapidly (within a few min) expands chromatophores, causing pigment granules to disperse and color skin (either in vivo or in vitro). Not species-specific; MSH from animals expands melanocytes in human skin; increases during most of pregnancy. In anuran larvae, MSH causes expansion of chromatophores and contraction of guanophores. Also has adipokinetic effect.	(-): chromatophore contraction and guanophore expansion; complete lack causes permanent blanching of skin (+): hyperglycemia; darkening of skin; potentiated or supplemented by ACTH	(I) corticoids (?); may be a feedback mechanism (S) central nervous system; ACTH ?	3,17,18
Pineal Body				
Zona glomerulosa of adrenal cortex	Stimulates secretion of aldosterone, possibly from ACTH-dependent precursors along a branch in synthetic chain	(-): ↓ aldosterone secretion, resulting in loss of Na <sup>+</sup> from body (+): ↑ aldosterone secretion	Unknown	3,8,9
Melanocytes in skin; gonads	Blanches skin of amphibians and fishes (very potent); antagonistic to MSH; ↓ ovulation, estrus (rat); ↓ pineal wt and uptake of H <sup>3</sup> -melatonin on exposure to light (rat); ↑ estrus	(-): darkening of skin of animals with expandable melanocytes; ↑ gametogenesis, estrus, ovarian wt; precocious puberty (+): blanching of skin; delay or decrease in estrus, gametogenesis; ↓ ovarian wt; erythema or punched-out ulcer (man)	(I) light; level of circulating melatonin? (S) dark (?); ↓ concentration of circulating melatonin	3,16,32
Thyroid				
None as such	Storage form of hormone in thyroid follicles	(-): delayed maturation, epiphyseal closure, and growth; incomplete differentiation or metamorphosis; cretinism; dwarfism; goiter (usually nontoxic or hyperiodic); slows reflexes; ↓ mentality, circulation, cardiac output, respiration, internal absorption (+): acceleration of growth, metabolic rate, maturation or metamorphosis; toxic goiter; uncouples oxidation from phosphorylation; mitochondrial swelling; thyrotoxicosis	(I) antithyroid drugs (thiouracil, thiourea, thiocyanate); thyroxine; high intake of I <sub>2</sub> or I <sup>-</sup> ; severe stress (first 24-48 hr) (S) thyrotropic hormone; chronic stress	4,23, 28,31

acid composition: Asp, Glu (swine); Ser (ox); Gly, Pro, Tyr, Lys, Met, Glu, His, Phe, Arg, Try, Gly, Ser, Pro, Pro,

*continued*

Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabolites
(A)	(B)	(C)	(D)	(E)
Thyroid				
13 Thyroxine; "T <sub>4</sub> "; C <sub>15</sub> H <sub>11</sub> O <sub>4</sub> NI <sub>4</sub> ; (β-[(3,5-diiodo-4-OH-phenoxy)-3,5-diiodophenyl]-alanine)	MW = 777; needles; MP = 232-233; [α] <sub>D</sub> = -3.2° (NaOH); s. alkali water, alkali or acid alcohol; i. water, alcohol, volatile solvents	Thyroid gland; plasma; synthesized from <i>p</i> -methoxyphenol and 3,4,5-triiodonitrobenzene	Limb bud growth (amphibian larva); growth (thyroidectomized rat); ↑ basal metabolic rate (thyroid-deficient subjects); ↑ O <sub>2</sub> consumption; oxidation of succinate; prevention of goiter; survival in anoxia (rat)	Deaminated and decarboxylated to tetraiodothyroacetic acid; as glucuronides or sulfates in urine, bile, or feces; deiodinated to T <sub>3</sub> or T <sub>2</sub>
14 3,5,3'-Triiodothyronine; "T <sub>3</sub> "; C <sub>15</sub> H <sub>12</sub> O <sub>4</sub> NI <sub>3</sub>	MW = 651; MP = 233; [α] <sub>D</sub> <sup>29.5°</sup> = +21.5° in 4.75% solution; s. HCl, alkaline water	Thyroid gland; plasma; deiodination of thyroxine	Methods similar to those for thyroxine; more active in most tests; same total calorogenic effect	
Parathyroid				
15 Parathyroid hormone; parathormone, PTH	Max. MW = 8,500; 83 amino acid residues; stable dilute acid; s. water, saline, aqueous alcohol, 94% acetic acid, concentrated phenol, 50% glycerol; i. volatile organic solvents; inactivated by proteases, alkali; reversibly inactivated by mild oxidation	Parathyroids	Rate and degree of rise in serum Ca <sup>++</sup> and decrease in serum phosphate (dog); Ca-mobilizing activity 2,000-3,000 USP units/mg; prevention of fall in plasma Ca in parathyroidectomized rat	
Adrenal Cortex				
16 Aldosterone <sup>7</sup> ; C <sub>21</sub> H <sub>28</sub> O <sub>5</sub> ; (11β,21-dihydroxy-3,20-dioxopregn-4-en-18-al)	MW = 360; MP = 164; [α] <sub>D</sub> = +160° (chloroform); s. organic solvents; sl. s. water. Urine concentration of aldosterone by double isotope derivative.	Zona glomerulosa of cortex (man, cattle, dog, frog, monkey, swine); precursors probably corticosterone, progesterone	Urinary Na/K ratio (adrenalectomized rat); muscle fatigue recovery (Everse and De Fremery test in adrenalectomized rat); muscular work performance (Ingle's test in adrenalectomized rat); deposition of liver glycogen (mouse, rat); wt maintenance (adrenalectomized dog); survival and growth (Kuizenga test in young, adrenalectomized rat); convulsion prevention (anti-insulin test in mouse); protection against cold (adrenalectomized rat); ↓ circulating eosinophils, circulating lymphocytes, thymus wt; chemical methods (formaldehyde, reducing properties, reaction with phenylhydrazine)	Synthesized in vivo from corticosterone and progesterone; approx. 400 μg/da "turned over" in liver; mean 24 hr secretion (adult man) = 100 μg reduced in liver to tetrahydro derivative (inactive); mean excretion (man) = 10 μg/da, 30-40% as glucuronide, and 4-8% as free steroid; balance as other conjugates
17 Deoxycorticosterone <sup>7</sup> ; DOC, DOCA; C <sub>21</sub> H <sub>30</sub> O <sub>3</sub> ; (Δ <sup>4</sup> -pregnen-21-ol-3,20-dione)	MW = 330; MP = 141-142; [α] <sub>D</sub> = +178° (alcohol); s. acetone, benzene, chloroform, volatile solvents, vegetable oils; i. water	Adrenal cortex; synthesized commercially from cholesterol, diosgenin	protection against cold (adrenalectomized rat); ↓ circulating eosinophils, circulating lymphocytes, thymus wt; chemical methods (formaldehyde, reducing properties, reaction with phenylhydrazine)	Corticosterone; aldosterone; dehydrocorticosterone; pregnanediol; 17-ketosteroids
18 Corticosterone <sup>8</sup> ; C <sub>21</sub> H <sub>30</sub> O <sub>3</sub> ; (11β,21-dihydroxy-4-pregnen-3,20-dione)	MW = 346; MP = 180-182; [α] <sub>D</sub> = +262° (alcohol); s. organic solvents; sl. s. vegetable oils; v. sl. s. water	Adrenal cortex; blood; urine; synthesized in vitro from deoxycholic acid, pregnenolone, progesterone, cholesterol		Aldosterone; 11-dehydrocorticosterone; 17-ketosteroids (11-OH-androsterone and 11-keto-etio-cholane-3-[α]-ol-17-one)
19 Dehydrocorticosterone <sup>8</sup> ; cortisone; C <sub>21</sub> H <sub>28</sub> O <sub>4</sub> ; (Δ <sup>4</sup> -pregnen-21-ol-3,11,20-trione)	MW = 344; MP = 178-180; [α] <sub>D</sub> = +258° (alcohol); s. organic solvents; i. water	Adrenal cortex; synthesized from deoxycholic acid		Similar to those of corticosterone; 11-ketopregnanediol main metabolite

/-/ Mineralocorticoid. /s/ Glucocorticoid.

# Contrails

## VERTEBRATES

Targets (F)	Principal Effects (G)	Effect of Deficiency (-) Excess (+) (H)	Secretion Inhibited by (I) Stimulated by (S) (I)	Reference (J)
<b>Thyroid</b>				
All body cells; adeno- hypophysis	Regulation of rate of CHO, fat, protein, H <sub>2</sub> O, mineral metabolism; stimulates growth, maturation, neuromuscular function, skin development, hematopoiesis, spermatogenesis, lactation, absorption through intestinal wall; regulation of TSH secretion; ↑ O <sub>2</sub> uptake of all organs and tissues	(-): delayed maturation, epiphyseal closure, and growth; incomplete differentiation or metamorphosis; cretinism; dwarfism; goiter (usually non-toxic or hyperiodic); slows reflexes; ↓ mentality, circulation, cardiac output, respiration, internal absorption (+): acceleration of growth, metabolic rate, maturation or metamorphosis; toxic goiter; uncouples oxidation from phosphorylation; mitochondrial swelling; thyrotoxicosis	(I) antithyroid drugs (thiouracil, thiourea, thiocyanate); thyroxine; high intake of I <sub>2</sub> or I <sup>-</sup> ; severe stress (first 24-48 hr) (S) thyrotropic hormone; chronic stress	4,7,15, 13 19,22-24,26-28,31, 34  4,23, 14 27,28, 31
	Same as thyroxine; more potent in goiter prevention, calorigenic activity, regulation of growth and metabolism			
<b>Parathyroid</b>				
Bones; kidneys, other soft tissues ?	↑ renal tubular absorption of Ca, distal renal tubular excretion of PO <sub>4</sub> . Controls Ca and PO <sub>4</sub> level in blood via mineral exchange between blood and bones, and PO <sub>4</sub> excreted by kidneys.	(-): hypocalcemia; irritability of nervous system; convulsive seizures; tetany; phosphate retention (+): hypercalcemia, renal calculi; ↑ calcium and phosphate excretion, osteoclastic activity; ↓ growth	(I) high serum (Ca <sup>++</sup> ) (S) low serum (Ca <sup>++</sup> ); high serum (PO <sub>4</sub> )	5,7,21, 15 22,24, 26-28
<b>Adrenal Cortex</b>				
Distal renal tubule	Promotes renal excretion of K, and retention of Na and Cl; indirectly promotes reabsorption of H <sub>2</sub> O in distal tubules; 25-100 times more potent than deoxycorticosterone	(-): ↓ Na <sup>+</sup> , H <sub>2</sub> O in blood and muscles; hemoconcentration; loss of Na through kidneys (+): reverse of above; hypertension; congestive heart failure	(I) ↓ K <sup>+</sup> concentration in blood, secretion of adrenoglomerulotropin; ↑ Na <sup>+</sup> , hemodilution (S) glomerulotropin; ACTH; ↑ K <sup>+</sup> , blood pressure in carotid arteries; ↓ Na <sup>+</sup> , dehydration of blood	6,10 16 21,22, 24,27
Distal renal tubule	Similar to aldosterone; chiefly of historical interest; normally secreted only in trace amounts	(+): acute Na <sup>+</sup> retention and K <sup>+</sup> loss; hypertension; polyuria; ↑ plasma volume and edema	Unknown	6,7,10, 17 13,15, 21,22, 24,26-28
Muscles, liver, capillaries, kidneys, pancreas (?), integument, lymphoid organs and bone marrow, circulating blood cells	Effects of glucocorticoids are qualitatively comparable, but differ quantitatively for glycogen deposition in liver, muscle work performance, hypersensitivity reactions, thymus involution, gluconeogenesis, maintenance of renal function, K/Na ratio and H <sub>2</sub> O balance, muscular fatigue recovery, cold protection, CHO metabolism,	(-): asthenia; hemoconcentration; skin pigmentation; ↓ wt, blood glucose, liver glycogen, stress resistance, blood pressure, secondary sex characteristics, growth in young; ↑ K/Na ratio in serum. Slowing of electrical discharges in nerves reversed by hydrocortisone.	(S) ACTH (α-corticotropin)	6,7,10, 18 15,20-22,24, 26-28, 33
		(+): reverse of above; Cushing's syndrome; hirsutism; negative		6,7,10, 19 13,21, 22,24, 26-28, 33

*continued*



## 90. HORMONES:

Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
<b>Adrenal Cortex</b>				
20 Hydrocortisone <sup>s</sup> ; 17-hydroxycorti- costerone, corti- sol; C <sub>21</sub> H <sub>30</sub> O <sub>5</sub> ; ( $\Delta^4$ -pregnene- 11 $\beta$ , 17 $\alpha$ , 21-triol- 3, 20-dione)	MW = 362; MP = 217- 220; [a] <sub>D</sub> = +167° (al- cohol); s. chloroform, ether, vegetable oils; sl. s. water	Adrenal cortex; syn- thesized from de- oxycholic acid	Same as for aldoster- one; Porter-Silber reaction; <i>m</i> -dinitro- benzene reaction	Tetrahydrocortisol; cortisone
21 Cortisone <sup>s</sup> ; 17-OH- 11-dehydrocorti- costerone; C <sub>21</sub> H <sub>28</sub> O <sub>5</sub> ; ( $\Delta^4$ - pregnene-17 $\alpha$ , 21- diol-3, 11, 20- trione)	MW = 360; MP = 220- 224; [a] <sub>D</sub> = +209° (al- cohol); s. acetone, chloroform, benzene, ether, vegetable oils; sl. s. water	Adrenal cortex; pla- centa; synthesized in vitro from squa- lene, cholesterol, pregnenolone, pro- gesterone	Same as for aldoster- one; Porter-Silber reaction	Tetrahydrocortisone; $\beta$ -cortolone
22 Fluorocortisone; C <sub>21</sub> H <sub>29</sub> O <sub>5</sub> F; (9 $\alpha$ - fluoro-17 $\beta$ -[1- keto-2-hydroxy- ethyl]- $\Delta^4$ -androstene-3,- 11-dione-17 $\alpha$ -ol)	MW = 380; MP = 233- 234 (acetate); [a] <sub>D</sub> <sup>23°</sup> = +123° (acetate)	Synthetic	Same as for aldoster- one	
<b>Adrenal Medulla</b>				
23 Epinephrine; ad- renaline, supra- renine, adrenine	MW = 183; MP (L) = 207-211; [a] <sub>D</sub> = -50° to -53.5°; s. alkali, acid; sl. s. water; v. sl. s. alcohol; i. ether, chloroform	Adrenal medulla; synthesized (com- mercially) from catechol, (in vivo) tyrosine; methyl- ation of norepineph- rine	Isolated heart stimula- tion (frog); uterus relaxation (cat); pupil dilation (cat); $\downarrow$ peri- stalsis (rabbit); $\uparrow$ blood pressure (cat)	Catechol derivatives; 50% to metanephrine (5-methoxyepineph- rine), 30% to 3- methoxy-4-hydroxy- mandelic acid (VMA) in urine
24 Norepinephrine; arterenol, norad- renaline	MW = 169; MP (L-bi- tartrate) = 163-164, (L-HCl) = 146-147; solubility same as for epinephrine	Adrenal medulla; various nerves, especially splenic; spleen; heart; blood vessels	Same as for epineph- rine in all tests ex- cept pressor and con- traction of gravid uterus	Methylated to epineph- rine by adrenals in presence of ATP; conjugated normeta- nephrine and VMA in urine
<b>Ovaries</b>				
25 Equilenin; C <sub>18</sub> H <sub>18</sub> O <sub>2</sub> ; ( $\Delta^1$ , 3, 5:10, 6, 8-estra- pentaen-3-ol-17- one)	MW = 266; MP = 258- 259; [a] <sub>D</sub> = +89° (diox- ane); s. volatile sol- vents; sl. s. alcohol, vegetable oils; i. water	Synthesized; 4 ster- eoisomers; natural estrogen from urine (pregnant mare)	Phenolsulfonic acid (Kober); ZnCl <sub>2</sub> /ben- zoyl Cl, sulfanilic acid/NaNO <sub>2</sub> (Pincus); spectrophotometric;	One inactive stereo- isomer is produced by catalytic dehy- drogenation of es- trone
26 Equilin; C <sub>18</sub> H <sub>20</sub> O <sub>2</sub> ; ( $\Delta^1$ , 3, 5:10, 7-es- tratetraen-3-ol- 17-one)	MW = 268; MP = 238- 240; [a] <sub>D</sub> = +308° (di- oxane); s. volatile solvents; sl. s. vege- table oils; i. water	Urine (pregnant mare)	estrus changes in va- gina (immature mouse, rat) (Allen- Doisy)	Easily dehydrogenated to equilenin
27 Estradiol-17 $\beta$ ; di- hydrotheelin, di- hydrofolliculin, dihydroestrone, di-OH-estrin; C <sub>18</sub> H <sub>24</sub> O <sub>2</sub> ; (3,- 17 $\beta$ -di-OH-1, 3,- 5:10 estratriene)	MW = 272; MP = 178; [a] <sub>D</sub> = +82° (dioxane); s. alkali, volatile sol- vents; sl. s. vegetable oils; i. water	Urine (pregnant woman, mare, rab- bit); ovary (swine); testes (stallion); human placenta; synthesized from cholesterol. Es- tradiol-17 $\alpha$ from urine (pregnant cattle, goat)	See equilenin. Estima- tion of dehydrogen- ases in placenta.	Estradiol-17 $\alpha$ con- verted to estrone, but not estradiol- 17 $\beta$ , in calves

/s/ Glucocorticoid.

## VERTEBRATES

Targets	Principal Effects	Effect of Deficiency (-) Excess (+)	Secretion Inhibited by (I) Stimulated by (S)	Reference	
(F)	(G)	(H)	(I)	(J)	
<b>Adrenal Cortex</b>					
Chiefly muscle, liver, connective tissue; also capillaries, kidneys, pancreas (?), integument, lymphoid organs and bone marrow, circulating blood cells	collagen maintenance, normal capillary permeability, protection against stress and shock. Corticosterone is of minor importance, but has weak antianabolic and catabolic effects. Hydrocortisone and cortisone principal glucocorticoids (man); stimulate gluconeogenesis and redistribution of fat; lymphocytopenia.	N balance; hyperglycemia; osteoporosis; obesity; muscle wasting; diabetes; alkalosis; inhibition of inflammatory responses and wound healing; diuresis; suppression of antibody formation; hyperglycemia (hydrocortisone); ↑ gastric acidity (hydrocortisone)	(S) ACTH (α-corticotropin)	1,6,7, 10,13, 21,22, 24,26-28,33	20
				1,6,7, 10,13, 21,22, 24,26-28,33	21
Joints affected with rheumatoid arthritis	(See preceding page)	(See preceding page)		2,11, 12,29	22
<b>Adrenal Medulla</b>					
Sympathetic nervous system; heart muscle; smooth and skeletal muscle; liver phosphorylase; adenohypophysis; lipase	Relaxation of nongravid uterus, peripheral arterioles, bronchial muscles; ↑ contraction of heart muscle (↑ output and rate), gravid uterus, iris muscle (radial), capillaries; ↑ spleen size, glucogenolysis,	(-): no clinical syndrome (+): over-secretion rare; may cause paroxysmic hypertension and, in some instances, sustained hypertension. Norepinephrine: pheochromocytoma.	(S) sympathetic nervous system, via splanchnic nerve; stress	6,7,15, 20,22, 24,26-28,30	23
Similar to epinephrine but causes arteriolar constriction; tends to decrease cardiac output; pressor effect is not reversed by ergotoxine	blood coagulation, ACTH secretion, salivary and sweat gland secretions; ↓ peristalsis; moderate lipolysis. Epinephrine: slight pressor effect; moderately increased cardiac output; great excitation of central nervous system; eosinopenia; hyperglycemia; greatly ↑ basal metabolic rate. Norepinephrine: great pressor effect; slight or no ↑ cardiac output; no excitation of central nervous system; slight eosinopenia and hyperglycemia; moderately ↑ basal metabolic rate.			6,7,22, 24,27	24
<b>Ovaries</b>					
All female sex organs; mammary glands; mucous membranes; adenohypophysis	Estrogenic. Endometrial proliferation; development and maintenance of vaginal mucosa, cornification of superficial layer; antagonizes androgen effects; ↑ mammary gland duct development, uterine motility, growth of axillary and pubic hair (♀, human), growth of down (♀, bird), growth of all female secondary sex organs. β-Estradiol stimulates <i>trans</i> dehydrogenases in human placenta, slows growth of skeleton, promotes closure of epiphysis; moderately stimulates protein anabolism and calcification of bones.	Unknown		3,10, 15,21, 24,26, 27	25
		Unknown		3,15, 21,24, 26,27	26
All female sex organs; mammary glands; mucous membranes; adenohypophysis; osteoblasts		(-): immaturity or atrophy of accessory sex organs; lack of secondary sex characteristics and female behavior patterns; ↓ mammary gland development; delayed epiphyseal closure; long bones continue to grow; osteoporosis (+): precocious maturity; hypertrophy of accessory sex organs and mammary glands; estrus changes; cystic	(I) high blood levels of estrogens ? (S) LH, combined FSH and LH	3,7,10, 15,21, 22,24, 26-28, 31,33	27

continued

Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
Ovaries				
28 Estriol; theelol, tri-OH-estrin; $C_{18}H_{24}O_3$ ; ( $\Delta^{1,3,5:10}$ -estratriene-3,16 $\alpha$ ,17 $\beta$ -triol)	MW = 288; MP = 282; $[\alpha]_D = +53$ to $+63^\circ$ (dioxane); s. volatile solvents, alkali; sl. s. vegetable oils; i. water	Urine (pregnant woman); human placenta; synthesized from estrone, $\beta$ -estradiol	See equilenin	$\beta$ -Estradiol; 4-pregnene-20 $\beta$ -ol-3-one ↓ estrone; 4-pregnene-20 $\alpha$ -ol-3-one ↓ $\alpha$ -estradiol ↓ estriol (principal metabolite in urine; excreted as water-soluble inactive glucuronide)
29 Estrone; theelin, folliculin, keto-OH-estrin; $C_{18}H_{24}O_2$ ; ( $\Delta^{1,3,5:10}$ -estratrien-3-ol-17-one)	MW = 270; MP = 255; $[\alpha]_D = +170^\circ$ (dioxane); s. volatile solvents, alkali; v. sl. s. water; sl. s. vegetable oils	Adrenal cortex; urine (man, bull, steer); synthesized from cholesterol, diosgenin	See equilenin	
30 Progesterone; progestin, luteosterone; $C_{21}H_{30}O_2$ ; ( $\Delta^4$ -pregnene-3,20-dione)	MW = 314; MP ( $\alpha$ ) = 128, ( $\beta$ ) = 121; $[\alpha]_D = +172$ to $+182^\circ$ (dioxane); s. volatile solvents; sl. s. vegetable oils; i. water	Corpus luteum; placenta; adrenal cortex; synthesized from cholesterol or stigmasterol	See equilenin	Pregnanediol; 17 $\alpha$ -hydroxyprogesterone
31 Relaxin; releasin	Polypeptide; IEP = 5.5; s. water and 95% alcohol	Serum (pregnant woman, cat, dog, mare, rabbit, sow); placenta; corpus luteum; ovaries; endometrium	Degree of relaxation of pelvic ligaments (guinea pig)	Unknown
Placenta <sup>3</sup>				
32 Chorionic gonadotropin; HCG, prolan	MW = 30,000; IEP = 2.95; $[\alpha]_D^{20} = -68^\circ$ ; s. water, 50% acetone, 60% alcohol; 12,000 I.U./mg	Placenta, blood, urine (pregnant female); peak in 60-75 days after first day of last menses	Corpus luteum formation (mouse); ovarian wt (rat); ovulation (rabbit); ovarian cell repair (immature, hypophysectomized rat)	Human chorionic gonadotropin excretion; basis of Aschheim-Zondek pregnancy test
Testes				
33 Testosterone; $C_{19}H_{28}O_2$ ; ( $\Delta^4$ -androsen-17 $\beta$ -ol-3-one)	MW = 288; MP = 155-156; $[\alpha]_D = +109^\circ$ (alcohol); absorption maximum = 238 m $\mu$ ; s. alcohol, ether, volatile solvents; sl. s. vegetable oils; i. water	Testes (man, bear, bull, rat, stallion); synthesized from cholesterol, acetate, and dehydroandrosterone	Alkali <i>m</i> -dinitrobenzene; SbCl <sub>3</sub> -acetic anhydride (colorimetric); chromatography; comb growth (capon); blackening of bill (sparrow); † wt of seminal vesicle, prostate (castrated rat). Testosterone: Girard's reagent T, followed by polarography.	Androsterone; etiocholanolone; epianandrosterone; estradiol; estrone (urine)
34 Dehydroepiandrosterone; dehydroisoandrosterone; $C_{19}H_{28}O_2$ ; ( $\Delta^5$ -androsen-3 $\beta$ -ol-17-one)	MW = 288; MP (leaflets) = 152-153; $[\alpha]_D = +10.9^\circ$ (alcohol); s. alcohol, ether, benzene; precipitated by digitonin; i. water	Urine (man, bull, pregnant cow); testes (bull); synthesized from cholesterol		May be intermediate in synthesis of androgens from acetate or pregnalone; metabolized to androstenedione
35 Androsterone; $C_{19}H_{30}O_2$ ; (androstane-3-[ $\alpha$ ]-ol-17-one)	MW = 290; MP = 185.5; $[\alpha]_D = +87.8^\circ$ (dioxane); s. volatile solvents; sl. s. vegetable oils; not precipitated by digitonin; i. water	Urine (man, bull, pregnant cow); synthesized from cholesterol or sitosterol		Dehydroepiandrosterone, $\Delta^4$ -androstene-3,17-dione, 17-hydroxyprogesterone. Found only in urine (probably a metabolite of testosterone).

<sup>3</sup>/o Placenta also produces estrogens and progesterone.

# Contrails

## VERTEBRATES

Targets (F)	Principal Effects (G)	Effect of Deficiency (-) Excess (+) (H)	Secretion Inhibited by (I) Stimulated by (S) (I)	Refer- ence (J)	
Ovaries					
All female sex organs; mammary glands; mucous membranes; adenohypophysis; osteoblasts	See entry 25	hyperplasia of endometrium; blocking of ovulation; skeletal growth deceleration (premature closing of epiphyses); excessive calcification of tissues, if parathyroids are normal  (See preceding page)	(I) high blood levels of estrogens ? (S) LH, combined FSH and LH	3,7,10, 15,21, 22,24, 26-28, 31,33  3,7,10, 15,21, 22,24, 26-28, 31,33	28  29
Endometrium of uterus; mammary gland lobules and alveoli; kidney tubules; adenohypophysis	Luteinizing. Preparation of endometrium for implantation of zygote; ↓ uterine contractions; ↑ mammary gland development, metabolism and excretion of estrogens; stimulation of protein catabolism and galactose oxidation	(-): lack of normal cyclic changes and of endometrial development for implantation and gestation (+): pregestational changes; pregnancy prolongation; inhibition of uterine growth, especially of endometrium; ↑ Na and K excretion, catabolism	(S) LTH	3,7,10, 15,21, 22,24, 26-28, 31,33, 34	30
Pubic ligaments	Relaxation of pubic ligament by replacing bone with connective tissue; synergistically with other hormones promotes mammary development and softening of cervix	Not definitely known	(S) estrogens ?	3,7,22, 24,31, 33	31
Placenta <sup>a</sup>					
Ovaries, especially corpus luteum	Corpus luteum maintenance in pregnancy; in nonpregnant women, stimulates ovulation. Human chorionic gonadotropin → luteinizing corpus luteum; pregnant mare serum → luteinization, follicle development.	(-): abortion (+): toxemias of pregnancy ?	(I) sex steroids	7,20, 22,26-28,31	32
Testes					
All male sex organs; adenohypophysis; muscle; hair follicles; epiphyses of long bones; vocal cords	Androgenic. ↑ development of male secondary sex organs and sex characteristics, libido, folliculoid and luteoid activity (immature female), basal metabolic rate and protein anabolism; positive balances of N, Ca, K, P; ↓ creatinuria and amino acid catabolism.	(-): immaturity or atrophy of accessory sex organs; lack of secondary sex characteristics and male behavior patterns; poor muscle development and function; delayed closure of epiphyses; ↓ excretion of 17-ketosteroids in urine (+): precocious sex development; hypertrophy of accessory sex organs; ↑ skeletal growth until closure of epiphyses, muscle mass, hirsutism, excretion of 17-ketosteroids in urine; ↓ scalp hair ?	(I) androgens (S) FSH + ICSH	3,10, 15,21, 22,24, 26-28, 31,33  3,10, 15,21, 22,24, 26,27, 31,33  3,7,10, 15,21, 22,24, 26-28, 31,33	33  34  35

continued



Name; Synonyms; Chemical Formula; (Systemic Name)	Properties	Sources	Assay Methods	Metabo- lites
(A)	(B)	(C)	(D)	(E)
<b>Pancreas</b>				
36 Insulin protein (polypeptide) <sup>10</sup>	MW = 36,000; monomer composed of two polypeptide chains; destroyed by oxidizing or reducing agents. MP = 223; IEP = 5.3; s. water, alcohol; precipitated by protein precipitants; stable acid; destroyed by alkali.	$\beta$ -cells of islets of Langerhans (various species)	Convulsions (mouse); hypoglycemia (starved rabbit); glucose uptake by diaphragm (rat, mouse); metabolism of glucose-1-C <sup>14</sup> by fat pad (rat); rate of displacement of beef insulin-I <sup>131</sup> by human insulin (immunized guinea pig); $\downarrow$ blood glucose (adrenodemedullated, hypophysectomized, diabetic rat)	
37 Glucagon (polypeptide) <sup>10</sup>	MW = 3,485; 29 amino acid residues; two polypeptide chains of different lengths; IEP = 7.5-8.5; s. dilute alkali and acid; i. water; stable alkali	$\alpha$ -cells of islets of Langerhans	Rise of blood glucose level at intervals of 20, 30, 45, 60, 90, and 120 min after intravenous injection; rate of glycogenolysis in liver	Not known
<b>Gastrointestinal Tract</b>				
38 Cholecystokinin; CCK	MW = 5,000-10,000 (?); IEP = 5.0-5.5; dialyzable; s. water	Upper intestinal mucosa	Contraction of gallbladder (dog)	Metabolized as protein ?
39 Enterocrinin	s. acid, water, alcohol; i. ether, acetone; salted out by NaCl	Intestinal mucosa (man, cat, cow, dog, hog)	$\uparrow$ flow of intestinal juice in jejunum (dog)	Metabolized as protein ?
40 Enterogastrone	s. water; i. organic solvents; dialyzable; destroyed by pepsin	Duodenal mucosa	$\downarrow$ stimulating effect of exogenous histamine on HCl secretion	Urogastrone ?
41 Gastrin	IEP = 8; heat stable; destroyed by ultraviolet, alkali, pepsin; dialyzable	Gastric mucosa, especially pylorus	No standard assay method	Unknown
42 Pancreozymin	Salted out by NaCl; s. absolute alcohol, water	Upper intestinal mucosa	Increase in enzymes in pancreatic juice	Unknown
43 Secretin	MP = 234; salted out by NaCl, CCl <sub>3</sub> COOH; s. dilute acidic water	Upper intestinal mucosa	Volume of pancreatic juice (dog)	Unknown
44 Urogastrone	Same as for enterogastrone	Urine	Same as for enterogastrone	Unknown
<b>Nervous System</b>				
45 Acetylcholine; ACh	MW = 163; MP (Br) = 143; quaternary salt unstable; s. water, alcohol, ether, oils; i. ether; salts v. s. water	Ganglia of parasympathetic nervous system; brain; synthesized commercially	None	

<sup>10/</sup> Data are for cattle.

**Contributors:** (a) Pritham, Gordon H., (b) Russell, Jane A.

**References:** [1] Borman, A., and F. M. Singer. 1954. Federation Proc. 13:185. [2] Borman, A., F. M. Singer, Williams and Wilkins, Baltimore. v. 1. [4] Danowski, T. S. 1962. Ibid. v. 2. [5] Danowski, T. S. 1962. Ibid.

# Contrails

Targets (F)	Principal Effects (G)	Effect of Deficiency (-) Excess (+) (H)	Secretion Inhibited by (I) Stimulated by (S) (I)	Refer- ence (J)
<b>Pancreas</b>				
All tissues involved in carbohydrate metabolism; hexokinases in mitochondria	Regulates CHO and fatty acid metabolism; stimulates hexokinases, phosphorylases; promotes uptake of glucose by cells; oxidation of CHO; stimulates lipogenesis, transport and oxidation of lipids; stimulates amino acid transport into cells and protein synthesis; stimulates synthesis of mucopolysaccharides. ↓ gluconeogenesis, ketogenesis.	(-): diabetes mellitus (hyperglycemia, glycosuria, ketonuria; ↓ wt and blood volume, negative nitrogen balance); delayed wound healing; gangrene; dwarfism; ↓ response to growth hormone; ↓ HPO <sub>4</sub> uptake by cells; ↓ ATP; ↓ glycogenesis, lipogenesis, proteogenesis (+): hyperinsulinism (hypoglycemia, convulsions, nausea, muscular weakness, anxiety, confusion); ↑ food intake, fat and protein deposition	(I) low blood glucose; antagonized by growth hormone, glucosteroids (corticoids), epinephrine, thyroxine, glucagon (S) vagus stimulation; ↑ blood glucose; ↑ growth hormone; ↑ ACTH, thyroxine, estrogens	7,15, 21,22, 24,26, 28 36
Similar to insulin	Antagonistic to insulin; raises blood glucose; promotes glycogenesis in liver but not muscle; inhibits synthesis of fatty acids from precursors; ↑ O <sub>2</sub> consumption by tissues, excretion of ions by kidney	(-): low blood glucose (+): high blood glucose; opposes action of insulin	(I) high blood glucose? (S) low blood glucose?	27,28, 30,31 37
<b>Gastrointestinal Tract</b>				
Gallbladder	↑ contraction and emptying of gallbladder	?	(S) fat, protein, acid in duodenum	14,21, 28 38
Secretory cells of ileum, jejunum	↑ secretion of succus entericus, volume rate and enzyme concentration	?	(S) food in intestine	14,21 39
Stomach	↓ motor activity and acid secretion of stomach	?	(S) sugar, fat in intestine	14,21, 28 40
Parietal (HCl-producing) cells, stomach	↑ HCl secretion (but not pepsin) by gastric mucosa	?	(S) mechanical distention; protein degradation products	14,21, 28 41
Enzyme-secreting cells of pancreas	↑ enzyme secretion by pancreas (no effect on volume rate)	?	(S) peptones, amino acids, soaps, fats in duodenum	14,21 42
Pancreas (acinar or exocrine), liver	↑ volume rate of pancreatic enzymes (no effect on concentration), bile secretion	(-): hyposecretion of pancreatic enzymes and bile (+): excess doubtful	(S) HCl, protein degradation products, digested fat or bile in small intestine	14,21, 28 43
Gastric mucosa and muscularis	↓ HCl secretion, stomach muscle contractions	?		14,21, 28 44
<b>Nervous System</b>				
Muscles, especially involuntary (ACh released at neuromuscular junctions, synapses)	Conduction of electrical impulses along nerve fibers; arteriolar dilation; effects (cholinergic) generally opposite to those of epinephrine; ↓ heart rate	?		15,22, 24,27, 28,30 45

and P. Numerof. 1954. Proc. Soc. Exptl. Biol. Med. 86:570. [3] Danowski, T. S. 1962. Clinical endocrinology. v. 3. [6] Danowski, T. S. 1962. Ibid. v. 4. [7] Emmons, C. W., ed. 1950. Hormone assay. Academic Press,

*continued*

*Continued*  
90. HORMONES: VERTEBRATES

New York. [8] Farrell, G. L. 1958. *Physiol. Rev.* 38:709. [9] Farrell, G. L. 1961. *Arch. Biochem. Biophys.* 94(3):543. [10] Fieser, L. F., and M. Fieser. 1949. *Natural products related to phenanthrene*. Ed. 3. Reinhold, New York. [11] Fried, J., and E. Sabo. 1953. *J. Am. Chem. Soc.* 75:2273. [12] Fried, J., and E. Sabo. 1954. *Ibid.* 76:1455. [13] Gordon, E. S., ed. 1950. *Steroid hormones. Symposium*. Univ. Wisconsin Press, Madison. [14] Grossman, M. I. 1950. *Physiol. Rev.* 30(1):33. [15] Lange, N. A., ed. 1956. *Handbook of chemistry*. Ed. 9. Handbook Publications, Sandusky, Ohio. [16] Lerner, A. B., and J. D. Case. 1960. *Federation Proc.* 19:590. [17] Lerner, A. B., and T. H. Lee. 1962. *Vitamins Hormones* 20:337. [18] Li, C. H. 1961. *Ibid.* 19:321. [19] Means, J. H., et al. 1963. *The thyroid and its diseases*. Ed. 3. McGraw-Hill, New York. [20] Moulton, F. R., ed. 1944. *The chemistry and physiology of hormones*. American Association for the Advancement of Science, Washington, D. C. [21] Pincus, G., and K. V. Thimann, ed. 1948. *The hormones*. Academic Press, New York. v. 1. [22] Pincus, G., and K. V. Thimann, ed. 1950. *Ibid.* v. 2. [23] Pitt-Rivers, R. V., and J. R. Tata. 1959. *The thyroid hormones*. Pergamon Press, New York. [24] Selye, H. 1949. *Textbook of endocrinology*. Ed. 2. *Acta Endocrinologica*, Montreal. [25] Smith, R. W., Jr., O. H. Gaebler, and C. N. H. Long. 1955. *Hypophyseal growth hormone, nature and actions*. McGraw-Hill, New York. [26] Soffer, L. J. 1956. *Diseases of the endocrine glands*. Ed. 2. Lea and Febiger, Philadelphia. [27] Stecher, P. G., et al., ed. 1960. *The Merck index*. Ed. 7. Merck, Rahway, N. J. [28] Turner, C. D. 1960. *General endocrinology*. Ed. 3. W. B. Saunders, Philadelphia. [29] Ward, L. E., et al. 1954. *Proc. Staff Meetings Mayo Clinic* 29:649. [30] West, E. S., and W. R. Todd. 1961. *Textbook of biochemistry*. Ed. 3. Macmillan, New York. [31] Williams, R. H. 1962. *Textbook of endocrinology*. Ed. 3. W. B. Saunders, Philadelphia. [32] Wurtman, R. J., J. Axelrod, and E. W. Chu. 1963. *Science* 141:277. [33] Young, W. C., ed. 1961. *Sex and internal secretions*. Ed. 3. Williams and Wilkins, Baltimore. [34] Zondek, H. 1944. *Diseases of the endocrine glands*. Ed. 4. Williams and Wilkins, Baltimore.

91. ENDOCRINE ORGANS AND HORMONES: INVERTEBRATES

	Phylum and Class	Possible Endocrine Organs or Areas <sup>1</sup>	Possible Endocrine Factors [Chemistry]	Effects
	(A)	(B)	(C)	(D)
1	Chordata <sup>2</sup>	Some neurosecretion in brain		
2		Endostyle	Minimal thyroxinogenesis	?
3	Echinodermata	Some neurosecretion in circum-oral nerve-ring and radial nerves	[Polypeptide?]	Stimulation of spawning Water balance
4				
5	Arthropoda Arachnida	Brain-and-neurohemal-organ neurosecretory system		
6	Crustacea	X-organ-and-sinus-gland neurosecretory system (in eyestalk)	Somatic chromatophorotropins [Protein-polypeptide?]: A substance (=PLH of F. Brown) UDH ( <i>Uca</i> -darkening hormone)	Somatic pigmentation: Concentration of red pigments Dispersion of melanins
7			Retinal chromatophorotropin(s)	Retinal pigment movements in light and dark adaptation
8			Molt-influencing hormones (MIH and MAH)	Probable inhibition or acceleration of Y-organ secretion
9			Gonad-inhibiting hormones (in all ♀ crustaceans and ♂ crabs)	
10			Metabolic factors?	O <sub>2</sub> consumption, Ca <sup>++</sup> metabolism, water metabolism (probably related to molting); blood sugar regulation (diabetogenic)

<sup>1</sup>/ Neurosecretion, as used in this column, refers to the presence of special secretory-appearing neurons, regardless of evidence for actual neurohormone production. <sup>2</sup>/ Cephalochordata and Urochordata only.

*continued*

*Continued*

**91. ENDOCRINE ORGANS AND HORMONES: INVERTEBRATES**

Phylum and Class	Possible Endocrine Organs or Areas <sup>1</sup>	Possible Endocrine Factors [Chemistry]	Effects
(A)	(B)	(C)	(D)
11 Arthropoda Crustacea	Brain(?) - and - postcommissure-organ neurosecretory system	Somatic chromatophorotropins [Protein-polypeptide?]; A substance  A' substance  B substance (=CDH of F. Brown)	Somatic pigmentation:  Concentration of red pigments Concentration of white and some red and black pigments Dispersion of red pigments
12	Thoracic-ganglion-and-pericardial-organ neurosecretory system	[5-Hydroxytryptamine?; 5,6-dihydroxytryptamine?; polypeptide?]	Frequency and amplitude of heart beat
13	Thoracic-ganglion-and-anterior-ramifications neurosecretory system		Rate of gas transport?
14	Neurosecretion in brain and ventral ganglia unrelated to known neurohemal areas		
15	Y-organ	Molting hormone (insect ecdyson?)	Rate of molting in adult; under control of sinus gland MIH and MAH
16	Androgenic gland	Androgenic hormone	Male gonad, gonoduct, and secondary sex character development. Under control of sinus gland testis-inhibitory factor in crabs?
17	Ovary?		Female secondary sex characters
18 Insecta	Brain-and-corpus-cardiacum neurosecretory system	"Brain hormone" = ecdysiotropin [Protein? or steroid related to cholesterol?]	Stimulates secretion by ecdysial (molting) gland
19	Tritocerebral neurosecretion	Chromatophorotropin in phasmids and <i>Corethra</i> larva	
20	Neurosecretion in subesophageal ganglion		Activity regulation in cockroaches; release of chromatophorotropin in phasmids; egg diapause in <i>Bombyx</i>
21	Neurosecretion in ventral ganglia		
22	Corpus cardiacum (intrinsic function)	Myotropic factors	Motility of oviduct, gut, Malpighian tubules
23		Cardioaccelerator factor (produced by pericardial cells under corpus cardiacum stimulation?) [Orthodiphenol?]	Heart beat
24		Diabetogenic factor	Blood sugar regulation
25			Depression of spontaneous activity in nerve cords
26	Corpus allatum	Neotenin = "juvenile" hormone [Fat-soluble; related to farnesol (C <sub>15</sub> H <sub>25</sub> OH)?]	Larval and nymphal development; gonadotropic in some ? insects (not Lepidoptera or phasmids)
27	Corpora cardiaca and allata	Metabolic factors?	Water metabolism?
28	Ecdysial (molting) gland: ventral, prothoracic, thoracic, peritrateal, etc.	Ecdyson = growth and differentiation (GD) hormone = molting hormone [C <sub>27</sub> H <sub>44</sub> O <sub>6</sub> (polyhydroxysteroid)]	Larval and pupal molting (protein synthesis, quinone tanning in integument, tyrosine metabolism); differentiation of adult structures; diapause termination
29	Gonads	Sex hormones?	
30 Symphyla <sup>2</sup>	Brain-and-cerebral-gland neurosecretory system		Molt inhibition

/:/ Neurosecretion, as used in this column, refers to the presence of special secretory-appearing neurons, regardless of evidence for actual neurohormone production. /s/ Also Chilopoda, Diplopoda, and Pauropoda.

*continued*



**91. ENDOCRINE ORGANS AND HORMONES: INVERTEBRATES**

	Phylum and Class	Possible Endocrine Organs or Areas <sup>1</sup>	Possible Endocrine Factors [Chemistry]	Effects
	(A)	(B)	(C)	(D)
31	Arthropoda Onychophora	Neurosecretion in brain, ventral cord, pedal nerves; neurohemal infracerebral organs?		
32	Annelida Hirudinea	Neurosecretion in brain and nerve cord; neurohemal organ at base or posterior surface of brain		Reproduction?
33	Oligochaeta	Same as in Hirudinea		Secondary sex characteristics; gamete maturation; regeneration; pigmentation
34	Polychaeta	Same as in Hirudinea	"Juvenile hormone"	Gonad inhibition; inhibition of heteronereid transformation
35				Regeneration
36	Echiuroidea	Neurosecretion in ventral cord		
37	Sipunculoidea	Neurosecretion in brain; anterior neurohemal area ("finger organs")		Reproduction? Myotropic factor?
38	Mollusca Cephalopoda	Some neurosecretion in brain		
39		Optic glands	Inhibited by light through optic tracts	Stimulation of sexual maturation and gonad development
40	Bivalvia	Neurosecretion in most ganglia		Inhibition of spawning
41	Scaphopoda	Neurosecretion in various ganglia		
42	Gastropoda	Neurosecretion in all ganglia		Gonad development? Water balance.
43		Gonads		Reproductive tract development
44	Aschelminthes <sup>2</sup>	Neurosecretion in cephalic nerve ring		
45	Nemertina	Neurosecretion in brain		Gonad inhibition in female
46		Cerebral organ		Spawning?
47	Platyhelminthes	Neurosecretion in brain	[Water-soluble]	Regeneration
48	Cnidaria	Neurosecretion (diffuse?)		Growth stimulation and inhibition

<sup>1/2</sup> Neurosecretion, as used in this column, refers to the presence of special secretory-appearing neurons, regardless of evidence for actual neurohormone production. <sup>1/4</sup> Nematoda only.

*Contributor:* Bern, Howard A.

*References:* [1] Bern, H. A., and I. R. Hagadorn. 1964. In T. H. Bullock and G. A. Horridge. Structure and function in the nervous system of invertebrates. W. H. Freeman, San Francisco. p. 358. [2] Carlisle, D. B., and F. G. Knowles. 1959. Cambridge Monographs Exptl. Biol. 10. [3] Gorbman, A., and H. A. Bern. 1962. A textbook of comparative endocrinology. J. Wiley, New York. p. 377. [4] Heller, H., and R. B. Clark, ed. 1962. Mem. Soc. Endocrinol. 12. [5] Heller, H., and U. S. von Euler, ed. 1963. Comparative endocrinology. Academic Press, New York. v. 2. [6] Jenkin, P. M. 1962. Animal hormones. Pergamon Press, London. [7] Ortman, R. 1960. In J. Field, ed. Handbook of physiology. American Physiological Society, Washington, D. C. sect. 1, v. 2, p. 1039. [8] Scharrer, B. 1955. In G. Pincus and K. V. Thimann, ed. The hormones. Academic Press, New York. v. 3, p. 57. [9] Scharrer, E., and B. Scharrer. 1963. Neuroendocrinology. Columbia Univ. Press, New York. [10] Scheer, B. T. 1960. Vitamins Hormones 18:141. [11] Takewaki, K., ed. 1962. Gen. Comp. Endocrinol., Suppl. 1. [12] Welsh, J. H. 1959. Comp. Endocrinol., Proc. Columbia Univ. Symp. Cold Spring Harbor, N. Y., 1958, p. 121. [13] Wigglesworth, V. B. 1959. The control of growth and form. Cornell Univ. Press, Ithaca.

**92. RELATIVE ACTIVITY OF GROWTH REGULATORS: PLANTS**

**Part I. CELL ELONGATION OF OAT COLEOPTILES**

Elongation effect was determined by floating 15 apical sections (3 mm in length) of decapitated *Avena* coleoptiles, 90-92 hours old, on the surface of 25 ml of solution in a covered Petri dish, at 24°C for 24 hours. Where concentrations greater than  $10^{-5} M$  were required for an elongation of 0.15 mm, the pH of the solutions was adjusted to 5.6 with NaOH. Activity Index =  $\frac{\text{molar concentration of indole-3-acetic acid inducing an elongation of 0.15 mm}}{\text{molar concentration of growth regulator inducing an elongation of 0.15 mm}} \times 100$ .

Compound		Activity Index	Compound		Activity Index
(A)		(B)	(A)		(B)
<b>Indole acids</b>					
1	3-Indoleacetic acid ( $5 \times 10^{-8} M$ )	100	59	Phenoxy acids	
2	Indole-3-acetonitrile	250	60	2,4-Dimethylphenoxyacetic acid	0.5
3	4-Chloroindole-3-acetic acid	140	61	2,5-Dimethylphenoxyacetic acid	0.2
4	4,7-Dichloro-2-methylindole-3-acetic acid	0.1	62	3,5-Dimethylphenoxyacetic acid	0
5	5,7-Dichloro-2-methylindole-3-acetic acid	1.5	63	2,4,6-Trimethylphenoxyacetic acid	0
6	5-Fluoroindole-3-acetic acid	50	64	2-Methyl-4-chlorophenoxyacetic acid	25
7	6-Fluoroindole-3-acetic acid	100	65	3-Methylsulfonylphenoxyacetic acid	0
8	5-Hydroxyindole-3-acetic acid	0.5	66	2-Nitrophenoxyacetic acid	0
9	2-Methylindole-3-acetic acid	1.5	67	3-Nitrophenoxyacetic acid	0.2
10	Indole-3-butyric acid	15	68	4-Nitrophenoxyacetic acid	0.1
11	Indole-3-propionic acid	1.5	69	2,4-Dinitrophenoxyacetic acid	0
<b>Phenoxy acids</b>					
12	Phenoxyacetic acid	0.03	70	2-Phenylphenoxyacetic acid	0
13	2-Acetylphenoxyacetic acid	0	71	4-Phenylphenoxyacetic acid	0
14	3-Acetylphenoxyacetic acid	0.02	72	$\alpha$ -Methyl- $\gamma$ -phenoxybutyric acid	0
15	4-Acetylphenoxyacetic acid	0	73	$\gamma$ -Phenoxybutyronitrile	0
16	3-Aminophenoxyacetic acid	0.005		$\alpha$ -Phenoxypropionic acid	0.5
17	4-Aminophenoxyacetic acid	0.02	74	Phenyl compounds	
18	2-Bromophenoxyacetic acid	0.1	75	Phenylacetic acid	1
19	3-Bromophenoxyacetic acid	2.5	76	$\alpha$ -Aminophenylacetic acid	0
20	4-Bromophenoxyacetic acid	1.5	77	4-Aminophenylacetic acid	0.05
21	2,4-Dibromophenoxyacetic acid	12.5	78	4-Chlorophenylacetic acid	1
22	2,6-Dibromophenoxyacetic acid	0	79	3-Fluorophenylacetic acid	1.5
23	2,4,6-Tribromophenoxyacetic acid	0	80	4-Fluorophenylacetic acid	1.5
24	3-Carboxyphenoxyacetic acid	0	81	2,5-Dihydroxyphenylacetic acid	0.02
25	4-Carboxyphenoxyacetic acid	0	82	3-Iodophenylacetic acid	10
26	2-Chlorophenoxyacetic acid	0.06	83	4-Iodophenylacetic acid	0
27	3-Chlorophenoxyacetic acid	2	84	2,4-Dimethylphenylacetic acid	0.5
28	4-Chlorophenoxyacetic acid	5	85	3,5-Dimethylphenylacetic acid	0.5
29	2,4-Dichlorophenoxyacetic acid	50	86	2,4,6-Trimethylphenylacetic acid	0
30	2,6-Dichlorophenoxyacetic acid	0	87	4-Nitrophenylacetic acid	0
31	3,5-Dichlorophenoxyacetic acid	0	88	4-Phenylphenylacetic acid	0
32	2,4,5-Trichlorophenoxyacetic acid	25	89	Diphenylacetic acid	2
33	2,4,6-Trichlorophenoxyacetic acid	0	90	Phenylacetoneitrile	1.5
34	2,4-Dichloro-6-methylphenoxyacetic acid	0	91	$\gamma$ -Phenylbutyric acid	0.05
35	2,4-Dichloro-5-nitrophenoxyacetic acid	0.2	92	<i>N</i> -Phenylglycine	1
36	3-Cyanophenoxyacetic acid	0.02	93	4-Chlorophenylglycine	0
37	4-Cyanophenoxyacetic acid	0	94	Phenylpropionic acid	0
38	2-Ethyl-4-chlorophenoxyacetic acid	0	95	2-Chlorophenylpropionic acid	0
39	2-Fluorophenoxyacetic acid	0	96	3-Chlorophenylpropionic acid	0
40	3-Fluorophenoxyacetic acid	0.02	97	4-Chlorophenylpropionic acid	0
41	4-Fluorophenoxyacetic acid	5	98	$\beta$ -Phenylpropionic acid	0
42	2,4-Difluorophenoxyacetic acid	2	99	5-Phenylthioglycolic acid	0.07
43	2-Trifluoromethylphenoxyacetic acid	0		4-Chlorophenylthioglycolic acid	0.5
44	3-Trifluoromethylphenoxyacetic acid	7	100	Benzoic acids	
45	3-Pentafluorosulfurphenoxyacetic acid	1	101	Benzoic acid	0
46	3-Hydroxyphenoxyacetic acid	0.07	102	2-Acetoxybenzoic acid	0
47	4-Hydroxyphenoxyacetic acid	0.01	103	2-Aminobenzoic acid	0
48	2-Iodophenoxyacetic acid	0.1	104	2-Amino-3,5-diiodobenzoic acid	0
49	3-Iodophenoxyacetic acid	7	105	2-Bromobenzoic acid	0.1
50	4-Iodophenoxyacetic acid	0	106	3-Bromobenzoic acid	0
51	2,4-Diiodophenoxyacetic acid	0	107	4-Bromobenzoic acid	0
52	2-Isopropylphenoxyacetic acid	0	108	2-Chlorobenzoic acid	0.05
53	2-Methoxyphenoxyacetic acid	0	109	3-Chlorobenzoic acid	0
54	3-Methoxyphenoxyacetic acid	0.1	110	4-Chlorobenzoic acid	0
55	4-Methoxyphenoxyacetic acid	0.03	111	2,4-Dichlorobenzoic acid	0
56	2-Methylphenoxyacetic acid	0.2	112	2,5-Dichlorobenzoic acid	1
57	3-Methylphenoxyacetic acid	0.07	113	2,6-Dichlorobenzoic acid	0.1
58	4-Methylphenoxyacetic acid	0.05	114	Pentachlorobenzoic acid	0
			115	2-Chloro-4-fluorobenzoic acid	0
				2-Chloro-6-fluorobenzoic acid	0.1

*continued*

**92. RELATIVE ACTIVITY OF GROWTH REGULATORS: PLANTS**

**Part I. CELL ELONGATION OF OAT COLEOPTILES**

Compound		Activity Index	Compound		Activity Index
(A)		(B)	(A)		(B)
Benzoic acids			Miscellaneous compounds		
116	2-Chloro-5-nitrobenzoic acid	0	139	Carboxymethyl dimethyldithiocarbamate	0.5
117	4-Ethyl-3-mercaptobenzoic acid	0	140	Ethoxycarbonylmethyl dibutyldithiocarbamate	0
118	2-Fluorobenzoic acid	0	141	Ethoxycarbonylmethyl diethyldithiocarbamate	0.1
119	2,5-Difluorobenzoic acid	0	142	Ethoxycarbonylmethyl dimethyldithiocarbamate	1
120	2-Fluoro-5-aminobenzoic acid	0	143	1-Cyclohexenylacetic acid	0.1
121	2-Fluoro-5-chlorobenzoic acid	0	144	$\Delta$ -1-Cyclopentenylacetic acid	0
122	2-Fluoro-3,5-dichlorobenzoic acid	0	145	Ferroceneacetic acid	0
123	2-Trifluoromethylbenzoic acid	0	146	Ferrocenediacetic acid	0
124	2-Iodobenzoic acid	0	147	Ferrocenepropionic acid	0
125	2,3,5-Triiodobenzoic acid	50	148	Gibberellic acid	100
126	3,4,5-Triiodobenzoic acid	0	149	5-Indanyloxyacetic acid	0
127	2,6-Dimethylbenzoic acid	0.05	150	1-Naphthaleneacetic acid	50
128	2,6-Dimethyl-3-bromobenzoic acid	3	151	1-Naphthaleneacetoneitrile	100
129	2,6-Dimethyl-3-chlorobenzoic acid	2	152	1-Naphthoic acid	0
130	2,6-Dimethyl-3-iodobenzoic acid	2.5	153	2-Naphthoxyacetic acid	0.7
131	2,6-Dimethyl-3-nitrobenzoic acid	0.1	154	2-Phenanthreneacetic acid	0
132	2,4,6-Trimethylbenzoic acid	0	155	3-Pyridoxyacetic acid	0
133	2-Nitrobenzoic acid	0.1	156	3-Pyridylacetic acid	0.01
Miscellaneous compounds			157	Quinoline-5-oxyacetic acid	0
134	Adamantine-1-acetic acid	0	158	Quinoline-6-oxyacetic acid	0
135	Azulene-1-acetic acid	1	159	5-Chloroquinoline-6-oxyacetic acid	0
136	Azulene-1-acetonitrile	4	160	5-Chloroquinoline-8-oxyacetic acid	0
137	Azulene-1-carboxylic acid	0	161	8-Chloroquinoline-5-oxyacetic acid	0
138	Benzothiazyl-2-oxyacetic acid	0.5			

Contributor: Muir, Robert M.

References: [1] Muir, R. M., and C. Hansch. 1953. Plant Physiol. 28:218. [2] Muir, R. M., and C. Hansch. 1955. Ann. Rev. Plant Physiol. 6:157. [3] Muir, R. M., and C. Hansch. 1961. Plant Growth Regulation, Intern. Conf., 4th, Yonkers, N. Y., 1959, p. 249. [4] Muir, R. M., and C. Hansch. 1961. Nature 190:741. [5] Muir, R. M., C. Hansch, and J. Gally. 1961. Plant Physiol. 36:222.

**Part II. STEM CURVATURE OF SLIT PEA AND LEAF EXPANSION OF BEAN**

Compound		Slit Pea Stem Curvature <sup>1</sup>	Bean Leaf Expansion <sup>2</sup>	Compound		Slit Pea Stem Curvature <sup>1</sup>	Bean Leaf Expansion <sup>2</sup>
(A)		(B)	(C)	(A)		(B)	(C)
Indole acids				Phenoxy compounds			
1	3-Indoleacetic acid	100	<18	15	3,5-Dichlorophenoxyacetic acid	<0.05	<44
2	$\beta$ -(3-Indole)-propionic acid	.....	<19	16	2,4,5-Trichlorophenoxyacetic acid	500	<4,740
3	$\gamma$ -(3-Indole)-butyric acid	190	<40	17	2,4,6-Trichlorophenoxyacetic acid	0.4	294
Phenoxy compounds				18	2,3,4,6-Tetrachlorophenoxyacetic acid	1	.....
4	Phenoxyacetic acid	0	<11	19	2,4-Difluorophenoxyacetic acid	12	5,360
5	2-Bromophenoxyacetic acid	.....	<23	20	2,4-Diodophenoxyacetic acid	.....	344
6	4-Bromophenoxyacetic acid	.....	6,160	21	4-Chloro-2-methylphenoxyacetic acid	500	513
7	2,4-Dibromophenoxyacetic acid	.....	11,500	22	2,4,6-Trimethylphenoxyacetic acid	0	.....
8	2,4,6-Tribromophenoxyacetic acid	0.1	.....	23	DL- $\alpha$ -(2,4-Dichlorophenoxy)-propionic acid	600	16,800
9	2-Chlorophenoxyacetic acid	4	<19				
10	3-Chlorophenoxyacetic acid	.....	<37				
11	4-Chlorophenoxyacetic acid	200	18,700				
12	2,4-Dichlorophenoxyacetic acid	200-1,200	23,500				
13	2,5-Dichlorophenoxyacetic acid	15	<69				
14	2,6-Dichlorophenoxyacetic acid	3-4	137				

<sup>1/1</sup> Expressed as percent of activity of 3-indoleacetic acid [4]. <sup>2/2</sup> Activity expressed as reciprocal of dose (micromoles) causing 50% repression of leaf expansion [2].

*continued*



**92. RELATIVE ACTIVITY OF GROWTH REGULATORS: PLANTS**

**Part II. STEM CURVATURE OF SLIT PEA AND LEAF EXPANSION OF BEAN**

Compound		Slit Pea Stem Curvature <sup>1</sup>	Bean Leaf Expansion <sup>2</sup>	Compound		Slit Pea Stem Curvature <sup>1</sup>	Bean Leaf Expansion <sup>2</sup>		
(A)	(B)	(C)	(A)	(B)	(C)	(B)	(C)		
Phenoxy compounds			Phenyl acids						
24	<i>n</i> -a-(2,4-Dichlorophenoxy)-propionic acid	1,200	.....	40	2,4-Dinitrophenylacetic acid	0.1	<23		
25	$\beta$ -(2,4-Dichlorophenoxy)-propionic acid	.....	<47	41	Phenylthioacetic acid	0	.....		
26	$\gamma$ -(2,4-Dichlorophenoxy)-butyric acid	.....	18,500	42	4-Chloro-2-methylphenylthioacetic acid	200	.....		
27	<i>n</i> -Butyl 2,4-dichlorophenoxyacetate	.....	23,100	Benzoic acids					
28	2,4-Dichlorophenoxyacetylchloride	.....	19,900	43	2-Chlorobenzoic acid	.....	<5		
29	2,4-Dichlorophenoxyacetamide	.....	7,760	44	3-Chlorobenzoic acid	.....	<15		
30	2,4-Dichlorophenoxyacetanilide	.....	30,800	45	4-Chlorobenzoic acid	.....	<8		
31	2,4-Dichlorophenoxyethanol	.....	22	46	2,4-Dichlorobenzoic acid	0	<19		
32	2,4-Dichlorophenoxyethylamine	.....	296	47	2,5-Dichlorobenzoic acid	.....	204		
33	2,4-Dichlorophenoxythioacetic acid	.....	20,300	48	3,4-Dichlorobenzoic acid	0	<19		
Phenyl acids			Naphthalene compounds						
34	Phenylacetic acid	3-6; 10	<3	49	2,3,5-Trichlorobenzoic acid	.....	2,130		
35	$\gamma$ -Phenylbutyric acid	2	.....	50	2,3,6-Trichlorobenzoic acid	200	.....		
36	4-Bromophenylbutyric acid	15	.....	51	3,4,5-Trichlorobenzoic acid	.....	<45		
37	2,4-Dichlorophenylacetic acid	15	.....	52	1-Naphthaleneacetic acid	250; 370	<100		
38	<i>N</i> -(2,4-Dichlorophenyl)-glycine	.....	2.04	53	2-Naphthaleneacetic acid	100	<19		
39	<i>S</i> -(2,4-Dichlorophenyl)-thiocolic acid	.....	<47	54	1-Naphthaleneacetamide	10	.....		
			Naphthoxy compounds						
			55 1-Naphthoxyacetic acid					.....	<40
			56 2-Naphthoxyacetic acid					.....	319
			57 1-Naphthoxyacetamide					.....	25
			Reference					1, 3-5	2, 6

/1/ Expressed as percent of activity of 3-indoleacetic acid [4]. /2/ Activity expressed as reciprocal of dose (micro-moles) causing 50% repression of leaf expansion [2].

*Contributors:* Brown, James W., and Weintraub, Robert L.

*References:* [1] Bonner, J. 1950. Plant biochemistry. Academic Press, New York. [2] Brown, J. W., and R. L. Weintraub. 1950. Botan. Gaz. 111:448. [3] Thimann, K. V. 1951. In F. Skoog, ed. Plant growth substances. Univ. Wisconsin Press, Madison. p. 32. [4] Thimann, K. V. 1952. Plant Physiol. 27:392. [5] Thimann, K. V. Unpublished. Harvard Univ., Cambridge, 1953. [6] Weintraub, R. L., J. W. Brown, and J. A. Throne. Unpublished. Fort Detrick, Maryland, 1953.

**93. ANTIMETABOLITES**

Metabolite	Antimetabolite <sup>1</sup>	Structural Alteration	Alteration Affects
(A)	(B)	(C)	(D)
1 Acetic acid	Fluoroacetic acid	F for H	.....
2 Adenine	Benzimidazole & derivatives	2 C for 2 N; side-chain alterations	Microorganisms, animals
3	Triazolopyrimidines	N for C	Microorganisms, animals
4	Diaminopurine	NH <sub>2</sub> for H	Bacteria, animals
5 $\alpha$ -Alanine	Glycine	H for CH <sub>3</sub>	Bacteria
6 $\beta$ -Alanine	$\beta$ -Aminobutyric acid	CH <sub>3</sub> for H	Yeast
7	Propionic acid	H for NH <sub>2</sub>	Bacteria
8	Asparagine	COOH for H; CONH <sub>2</sub> for COOH	Yeast
9 <i>p</i> -Aminobenzoic acid	Sulfanilamide & derivatives	SO <sub>2</sub> NH <sub>2</sub> or derivatives for COOH	Microorganisms
10	<i>p</i> -Aminobenzamide	CONH <sub>2</sub> for COOH	Bacteria
11	Carbarsonic & related arsenicals	Arsenic for C in a COOH group; derivatives	Microorganisms, animals
12	Phosphanilic acid	PO <sub>3</sub> H <sub>2</sub> for COOH	Microorganisms
13	Heterocyclic acids <sup>2</sup>	N or S for C	Bacteria
14	Ring-substituted PAB	Halogen or alkyl for H	Bacteria

/1/ Structural analog. /2/ E.g., 6-aminonicotinic acid.

*continued*



*Continails*  
93. ANTIMETABOLITES

Metabolite	Antimetabolite <sup>1</sup>	Structural Alteration	Alteration Affects	
(A)	(B)	(C)	(D)	
15	<i>p</i> -Aminobenzoic acid	<i>p</i> -Aminoacetophenone & derivatives	COR for COOH	Bacteria
16		<i>p</i> -Nitrobenzoic acid	NO <sub>2</sub> for NH <sub>2</sub>	Bacteria
17	Arginine	Canavanine	O for CH <sub>2</sub>	Bacteria
18	Ascorbic acid	Glucoscorbic acid	Addition of CHO <sub>H</sub> & optical inversion	Animals, liver enzymes
19	Aspartic acid	Hydroxyaspartic acid	OH for H	Bacteria
20		Aspartophenone	C <sub>6</sub> H <sub>5</sub> for OH	Bacteria
21	Biotin	Desthiobiotin & derivatives	2 H for S	Microorganisms
22		Biotin sulfone	SO <sub>2</sub> for S	Microorganisms
23		Ureylene-cyclohexyl aliphatic acids	2 C for S; derivatives with shorter side chains	Microorganisms
24		Desthioisobiotin	Loss of S, geometric isomerism	Insects
25		Ureylene-tetrahydrofuryl aliphatic sulfonic acids	O for S, SO <sub>3</sub> H for COOH	Microorganisms
26		Homobiotin	Addition of -CH <sub>2</sub> -	Microorganisms
27	Choline	Triethyl choline	3 ethyls for 3 methyls	Frog muscle, mice
28	Coccarboxylase	Thiaminethiazole pyrophosphate	Loss of pyrimidine portion	Carboxylase
29	Cytidine	Adenosine	OH for H, loss of imidazole ring	<i>Neurospora</i> mutant
30	Desthiobiotin	2-Oxyimidazole aliphatic acids	H for CH <sub>3</sub>	Microorganisms
31	Glutamic acid	Methionine sulfoxide	SOCH <sub>3</sub> for COOH	Bacteria
32		Hydroxyglutamic acid	OH for H	Bacteria
33		<i>N</i> -Alkylglutamines	<i>N</i> -Alkyl for OH	Bacteria
34	Guanine	Triazolopyrimidines	N for C	Bacteria
35		Benzimidazole	2 C for 2 N	Microorganisms
36	Histamine	Imidazole & derivatives	Elimination or substitution of part of molecule	Smooth muscle, histamine shock in animals
37		Diphenhydramine	Opening of ring, O for N, alkylation of N, C	Smooth muscle, histamine shock in animals
38		Tripelennamine	Opening of ring, alkylation of N	Smooth muscle, histamine shock in animals
39	Hypoxanthine	Hydroxytriazolopyrimidine	N for C	Bacteria
40	Indoleacetic acid	Phenyl butyric acid	Elimination of N and shift of 1 C	Plants
41		Skatyl sulfonic acid	SO <sub>3</sub> H for COOH	Plants
42	Inositol	Hexachlorocyclohexane	6 Cl for 6 OH	Fungi, plants, pancreatic amylase
43	Isoleucine	Leucine	Position isomerism of 1 CH <sub>3</sub>	Bacteria
44	Leucine	<i>o</i> -Leucine	Optical inversion	Bacteria
45	Lysine	Arginine	Guanidino for amino, elimination of CH <sub>2</sub>	<i>Neurospora</i> mutant
46	Methionine	Methoxinine	O for S	Bacteria
47		Ethionine	CH <sub>3</sub> for H	Bacteria and animals
48		Norleucine	CH <sub>2</sub> for S	Bacteria
49	Nicotinic acid (or amide)	Pyridine-3 sulfonic acid or amide	SO <sub>3</sub> H for COOH	Microorganisms, animals <sup>3</sup>
50		3-Acetylpyridine	COCH <sub>3</sub> for COOH	Animals; not microorganisms
51		5-Thiazole carboxamide	S for CH=CH	Certain bacteria
52	Pantothenic acid	Thiopianic acid (pantoyl-taurine) & derivatives	SO <sub>3</sub> H and derivatives for COOH	Microorganisms, pantothenate-utilizing enzymes; not animals
53		Pantothenyl alcohol	CH <sub>2</sub> OH for COOH	Microorganisms; not animals
54		<i>α</i> - or <i>β</i> -Methyl pantothenic acid	CH <sub>3</sub> for H	Microorganisms
55		Other substituted pantoamides	Alkyl or OH- and NH <sub>2</sub> -alkyl for CH <sub>2</sub> CH <sub>2</sub> COOH	Microorganisms
56		Phenyl pantothenone	COC <sub>6</sub> H <sub>5</sub> for COOH	Microorganisms
57		Salicylyl <i>β</i> -alanine	<i>o</i> -Hydroxy-benzoyl for pantoyl	Microorganisms
58		<i>γ</i> '-Methyl pantothenic acid	CH <sub>3</sub> for H	Bacteria
59	Phenylalanine	<i>β</i> -Hydroxyphenylalanine	OH for H	Bacteria
60		Thienylalanine	S for CH=CH	Microorganisms, animals
61		Furylalanine	O for CH=CH	Microorganisms
62		Halogenated phenylalanines	Halogen for H	Microorganisms
63	Pimelic acid	2,4-Dichlorosulfanilidocaproic acid	Dichlorosulfanilide for COOH	Biotin-independent microorganisms

<sup>1</sup>/ Structural analog. <sup>3</sup>/ Animal alcohol and lactic dehydrogenases, not animals in vivo.

*continued*

# Contrails

## 93. ANTIMETABOLITES

Metabolite	Antimetabolite <sup>1</sup>	Structural Alteration	Alteration Affects
(A)	(B)	(C)	(D)
64	Porphyrins <sup>4</sup>	Porphyrins lacking vinyl groups	Bacteria
65	Pteroylglutamic acid	Addition of two glutamic acids	Transplanted tumors
66	Xanthopterin	Loss of <i>p</i> -aminobenzoyl glutamic acid	Transplanted tumors
67	Pyridoxine	H for OH	Microorganisms, animals
68	2-Ethyl-3-amino-4-ethoxymethyl-5-amino-methyl pyridine	CH <sub>3</sub> for H, NH <sub>2</sub> for OH, Et for H	Microorganisms
69	Riboflavin	2 Cl for 2 CH <sub>3</sub>	Microorganisms
70	Isoriboflavin	Shift in position of CH <sub>3</sub>	Animals; not bacteria
71	Corresponding phenazine	2 C for 2 N, 2 NH <sub>2</sub> for 2 OH	Microorganisms, animals
72	Galactoflavin	Dulcetyl for ribityl	Animals
73	Lumiflavin	CH <sub>3</sub> for ribityl	Bacteria
74	Araboflavin	Inversion of position of OH	Animals
75	Succinic acid	Loss of CH <sub>2</sub>	Succinic oxidase
76	Malonic acid	SO <sub>3</sub> H for H	Succinic oxidase
77	Testosterone	Estradiol	Animals
78	Thiamine	Benzene ring for cyclohexane ring, loss of CH <sub>3</sub>	Microorganisms, animals
79	Pyrithiamine	CH=CH for S	Microorganisms, animals
80	Oxythiamine	OH for NH <sub>2</sub>	Animals, fish thiaminase
81	Butylthiamine	Butyl for CH <sub>3</sub>	Animals
81	Aminobenzyl-methylthiazolium chloride	2 C for 2 N, loss of side chains	Fish thiaminase
82	Thymine	5-Substituted dioxypyrimidines	Bacteria
83	2,4-Diamino- or dithiothymine	NO <sub>2</sub> or Br or NH <sub>2</sub> or OH for CH <sub>3</sub>	Bacteria
84	Thyroxine	NH <sub>2</sub> or SH for OH	Bacteria
84	Ethers of diiodotyrosine	<i>p</i> -Nitrobenzyl or <i>p</i> -nitrophenylethyl or benzyl for <i>p</i> -hydroxydiiodophenyl	Tadpoles
85	$\alpha$ -Tocopherol	Opening of ring by addition of O	Animals
86	Tryptophan	Indole of acrylic acid	Bacteria
87	Naphthylacrylic acid	Loss of NH <sub>3</sub> , C=C for N	Bacteria
88	Styrylacetic acid	Loss of NH <sub>2</sub> , substitution of aliphatic unsaturated side chain for pyrrole ring <sup>1</sup>	Bacteria
89	Methyltryptophans	CH <sub>3</sub> for H	Bacteria
90	Benzothienylalanine	S for N	Bacteria
91	Indole	Loss of side chain	Bacteriophage plus bacteria
92	Tyrosine	3-Fluorotyrosine	Rats
93	Uracil	OH for H	Bacteria
94	Thiouracil	S for O	Bacteria, plant seed germination
95	Vitamin K	Dicumarol & derivatives	Animals
96	Iodinol	O for C, side-chain alterations	Bacteria
97	$\alpha$ -Tocopherol quinone	2 N for 2 C, side-chain alterations	Animals
98	2,3-Dichloronaphthoquinone	2 CH <sub>3</sub> for benzene ring	Animals
98		2 Cl for alkyl side chains	Microorganisms
99	2-Substituted-3-hydroxynaphthoquinones	OH for H, change in alkyl substituent	Animals; not bacteria
100	Methoxynaphthoquinone	OCH <sub>3</sub> for CH <sub>3</sub>	Microorganisms

<sup>1/1</sup> Structural analog. <sup>4/4</sup> E.g., hematin and protoporphyrin.

Contributor: Woolley, D. W.

Reference: Woolley, D. W. 1952. A study of antimetabolites. J. Wiley, New York. pp. 33-36.

## Part I. PHYSICAL AND

Abbreviations: d. = decomposes; s. = soluble; i. = insoluble; sl. = slightly; l. = less;

Antibiotic (Synonym)	Source	Molecular Formula and Weight	Nature	Crystal Form and Color
(A)	(B)	(C)	(D)	(E)
1 Actinomycins	<i>Streptomyces</i> spp.	Varies with the amino acid content	Chromopeptides differing in amino acid portions of the molecule; usually occur naturally as mixtures of two or more components	Red or yellow-red plates, prisms, or needles
2 Amphotericin B <sup>1</sup>	<i>Streptomyces nodosus</i>	C <sub>46</sub> H <sub>73</sub> NO <sub>20</sub> ; 959 (neut. equiv.) <sup>2</sup>	Amphoteric conjugated heptaene	Deep yellow prisms or needles from dimethyl formamide
3 Bacitracins (Ayfivin)	<i>Bacillus subtilis</i> , <i>B. licheniformis</i>	Bacitracins A: C <sub>66</sub> H <sub>103</sub> O <sub>16</sub> N <sub>17</sub> S; 1,470±10% (actual) Bacitracins B: C <sub>71</sub> H <sub>112</sub> O <sub>17</sub> N <sub>18</sub> S	Weakly basic polypeptides	White, highly hygroscopic, amorphous powder
4 Carbomycin <sup>4</sup> (Antibiotic M 4209)	<i>Streptomyces</i> spp.	C <sub>42</sub> H <sub>67</sub> NO <sub>16</sub> ; 842 (calc.); 830-860 (neut. equiv.)	Monobasic macrolide	Slender, white, blunt-ended, needle-shaped crystals; colorless laths from ethanol; rectangular plates from ethanol-water
5 Chloramphenicol (Antibiotic 8-44; Levomycetin; Sintomycetin; Synthomycin)	<i>Streptomyces venezuelae</i> ; by synthesis	C <sub>11</sub> H <sub>12</sub> N <sub>5</sub> O <sub>2</sub> Cl <sub>2</sub> ; 323 (calc.); 310 (Rast)	Neutral	Colorless, elongated plates or fine needles
6 Colistin <sup>5</sup>	<i>Bacillus colistinus</i>	C <sub>45</sub> H <sub>85</sub> N <sub>13</sub> O <sub>10</sub> ; 1200±50; about 969 (actual)	Heteromeric decapeptide	Colorless
7 Erythromycin <sup>6</sup> (Erythromycin A)	<i>Streptomyces erythraeus</i>	C <sub>37</sub> H <sub>67</sub> NO <sub>13</sub> ; 733.9 (calc.)	Basic macrolide	White needles
8 Griseofulvin	<i>Penicillium</i> spp.; by synthesis	C <sub>17</sub> H <sub>17</sub> O <sub>6</sub> Cl; 309-398 (actual); 352.5 (calc.)	Neutral	Massive colorless rhombic or octahedral crystals from benzene
9 Kanamycins <sup>8</sup>	<i>Streptomyces kanamyceticus</i>	Kanamycin A: C <sub>18</sub> H <sub>36</sub> N <sub>4</sub> O <sub>11</sub> ; 427-490 (actual) Kanamycin B: C <sub>16</sub> H <sub>32</sub> N <sub>4</sub> O <sub>10</sub> ; 385-560 (Barger); 463.3 (calc.) Kanamycin C <sup>9</sup> : C <sub>18</sub> H <sub>36</sub> N <sub>4</sub> O <sub>11</sub> ; 415-625 (Barger)	Basic	Base: fine colorless needles Sulfate: small, irregular, white, prismatic crystals after repeated crystallization
10 Neomycins <sup>10</sup>	<i>Streptomyces</i> spp.	Complex: C <sub>23</sub> H <sub>46</sub> N <sub>6</sub> O <sub>13</sub> ; 507-669 (ebull.)	Basic	Sulfate, HCl, and free base: colorless amorphous solids
11 Novobiocin	<i>Streptomyces griseoflavus</i> , <i>S. niveus</i> , <i>S. spheroides</i>	C <sub>31</sub> H <sub>36</sub> N <sub>2</sub> O <sub>11</sub> ; 592±25 (ebull.); 610 (Rast); 618 (X ray)	Dibasic acid	White to cream-colored amorphous solid or pale yellow crystals (two crystalline forms, I and II)
12 Nystatin (Fungicidin)	<i>Streptomyces noursei</i> ; other <i>Streptomyces</i> species	C <sub>46</sub> H <sub>77</sub> O <sub>19</sub> ; 933 (calc.)	Amphoteric conjugated tetraene	Pale yellow microcrystals or needles

<sup>1</sup>/ The same *Streptomyces* culture produces amphotericin A, a conjugated tetraene antifungal agent [46, 47].  
<sup>2</sup>/ Carbomycin B occurs in beers of *Streptomyces halstedii* [68]. <sup>3</sup>/ Identical with polymyxin E [132]. <sup>4</sup>/ Eryth- partially resolidifies and then melts at 190-193°C [54]. <sup>5</sup>/ A complex of kanamycins A, B, and C [90, 105]. <sup>6</sup>/ Iso- years, under refrigeration [21].

## ANTIBIOTICS

### CHEMICAL CHARACTERISTICS

v. = very; calc. = calculated; neut. equiv. = neutral equivalent; ebull. = ebullient.

Melting Point °C	Solubility	Stability	Reference	
(F)	(G)	(H)	(I)	
215-255 d.	s. benzene, ethanol acetone; sl.s. water, ether; i. aqueous dilute mineral acids and alkalis, petroleum ether	Thermostable at pH 6-7; relatively stable at acid pH; unstable at alkaline pH	126,127	1
Gradual decomposition above 170	s. dimethyl sulfoxide, acetic <i>N,N</i> -dimethylformamide; sl.s. propylene glycol, glacial acetic acid, <i>N,N</i> -dimethylformamide; i. water, chloroform, methanol, ether, pyridine, alcoholic KOH <sup>1</sup>	Dry solids stable for long periods at moderate temperatures; undergoes decomposition in solution at pH 4-10	B,46,47;C,E-G,45;D,H,120	2
	s. methanol, ethanol, water; sl.s. acetone, cyclohexanone, propanols, butanols, pyridine; i. ether, chloroform, benzene, acetone, ethyl acetate, petroleum ether	Relatively thermostable, especially at pH 4 and 5; unstable above pH 9; aqueous solutions unstable after storage at room temperature <sup>3</sup>	B,73;C,29,30;D,9,28;E,9;G,H,6,15,73	3
210-218 d.	Acid salts: s. water, most organic solvents Base: i. water, hexane	Most stable at pH 5-7 (11 days at 25°C); stable for several months in dark at room temperature; unstable below pH 3 and above pH 9	B,51,72,115;C,44,51,72;D,44,51,115,133;E,51,115,121;F,H,44,51,121;G,115,121	4
149.7-150.7	s. propylene glycol, methanol, ethanol, butanol, ethyl acetate, acetone, diethyl ether; sl.s. chloroform, water, alkali; i. acid, benzene, petroleum ether, vegetable oils	Thermostable; alkali labile	B,27,48,80;C,11,101;D,10,48,60,101;E,10,48,57;F,10,11,48,57,101;G,10,48,57,60,101;H,10,11,48,57	5
Base: 245-249 d. Phosphate: 145	s. water, lower alcohols	Stable at 40°C for at least 60 days; salt solutions stable at pH 2-6, less stable above pH 6	B,E,F,79;C,G,79,106;D,33	6
Complex: 82-83.5 Base: 136-140 <sup>7</sup> HCl: 170-173	s. alcohols, acetone, chloroform, acetonitrile, ethyl acetate; l.s. ether, ethylene dichloride, amyl acetate, water HCl: v.s. water, lower alcohols	Stable at -25° to +4°C; stable 4 days at 37°C; unstable at 60° or 100°C	B,84;C,130;D,84,130;E,F,54;G,54,84;H,62	7
220-221	s. <i>N,N</i> -dimethylformamide 12-14% at room temperature, acetic acid, dioxane, benzene, ether, ethanol; sl.s. chloroform, ethyl acetate, toluene, acetone, ligroin; i. water, petroleum ether	Thermostable	B,20,35,116,118;C,59,93;D,E,93;F,59;G,8,93;H,18	8
Decompose over a wide range	Base: s. water; sl.s. lower alcohols; i. nonpolar solvents HCl: v.s. water; s. methanol; sl.s. ethanol; i. acetone, ethyl acetate, butyl acetate, ether, benzene	Thermostable, especially at neutral pH	B,119;C,32,85,89,122;D,32;E,32,89,122;F,32,90,105,122;G,32,105,122;H,31,58	9
	s. water; i. organic solvents	Thermostable; crude neomycin stable at pH 2-9; highly purified preparations stable to alkali only	124,125	10
I: 152-156 d. II: 174-178 d.	s. methanol, ethanol, butanol, acetic acid, dioxane, water above pH 7.5; i. water below pH 7.5, ether, benzene, carbon tetrachloride, chloroform Mono- and di-sodium salts: v.s. water	Dry material stable at 24°C in absence of light; dilute aqueous solutions stable at pH 2 at 24°C; half-life, 60 days at pH 7-10	B,111,128;C,67,69,74;D,F,69,74;E,69,74,88;G,88;H,70	11
Gradual decomposition above 160 without melting at 250	s. <i>N,N</i> -dimethylformamide, <i>N,N</i> -dimethylacetamide, propylene glycol; l.s. methanol, aqueous <i>n</i> -propanol or isopropanol, water-saturated butanol; i. water, most organic solvents	Unstable; most stable as a dry powder <sup>11</sup> ; aqueous suspensions less stable than alcoholic solutions; aqueous suspensions stable at least eight months at -25°C	B,65;C,G,21;D,F,107;E,43	12

<sup>12</sup>/ Perchloric acid in acetic acid [45]. <sup>13</sup>/ Bacitracin F, a transformation product, is formed above pH 7 [30, 64]. romycins B and C also occur in same fermentation liquors [131]. <sup>17</sup>/ If slow rate of heating continues, compound meric with kanamycin A [89]. <sup>10</sup>/ A complex of neomycins B and C which are isomeric [125]. <sup>11</sup>/ At least 4½

*continued*



## Part I. PHYSICAL AND

Antibiotic (Synonym)	Source	Molecular Formula and Weight	Nature	Crystal Form and Color
(A)	(B)	(C)	(D)	(E)
13 Oleandomycin (Antibiotic PA 105)	<i>Streptomyces antibioticus</i>	C <sub>35</sub> H <sub>61</sub> NO <sub>12</sub> ; 715 (ebull.); 687.84 (calc.)	Basic mac- rolide	Long white needles; colorless prisms
14 Paromomycin (Aminosidin; Catenulin; Hydroxymycin)	<i>Streptomyces</i> spp.	C <sub>23</sub> H <sub>45</sub> N <sub>5</sub> O <sub>14</sub>	Basic	Amorphous; white
15 Penicillins			Strong mono- basic car- boxylic acid	
16 Ampicillin	Semisynthetic	C <sub>16</sub> H <sub>19</sub> N <sub>3</sub> O <sub>4</sub> S		Fine needles
17 Benzylpenicil- lin (Penicil- lin G)	<i>Penicillium</i> spp.; <i>Asper- gillus</i> spp.	C <sub>16</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub> S; 331 (actual)		Colorless prisms
18 Cephalosporin N (Penicillin N; Synnematin B)	<i>Cephalosporium</i> spp.; <i>Strepto- myces</i> sp.	C <sub>14</sub> H <sub>21</sub> N <sub>3</sub> O <sub>6</sub> S	A hydrophilic penicillin	Ba salt; white powder
19 Methicillin	Semisynthetic	Sodium salt: C <sub>17</sub> H <sub>19</sub> N <sub>2</sub> O <sub>6</sub> SNa		White crystalline solid
20 Oxacillin	Semisynthetic	Sodium salt: C <sub>19</sub> H <sub>18</sub> N <sub>3</sub> O <sub>5</sub> SNa		
21 Phenethicillin	Semisynthetic	Potassium salt: C <sub>17</sub> H <sub>19</sub> N <sub>2</sub> O <sub>5</sub> SK; 402.51		Colorless crystals
22 Phenoxymeth- ylpenicillin (Penicillin V)	<i>Penicillium notatum</i>	Potassium salt: C <sub>16</sub> H <sub>17</sub> N <sub>2</sub> O <sub>5</sub> SK		
23 Polymyxins	<i>Bacillus poly- myxa</i>	Polymyxin B <sub>1</sub> ·5HCl: C <sub>56</sub> H <sub>104</sub> N <sub>16</sub> O <sub>14</sub> Cl <sub>5</sub> ; 1,150±10% (actual) Polymyxin D·4HCl: C <sub>50</sub> H <sub>97</sub> N <sub>15</sub> O <sub>15</sub> Cl <sub>4</sub> ; 1,150 (actu- al); 1,144 (calc.) Average for A, B, D, E: 1,250 (actual)	Basic poly- peptides	Birefringent; no definite structure
24 Ristocetins <sup>12</sup>	<i>Nocardia lurida</i>	Ristocetin A: 2,500 (freezing point depression); 5,000 (ultra- centrifuge)	Amphoteric	A sulfate: hexagonal prismatic rods B sulfate: needles
25 Spiramycins <sup>13</sup> (Antibiotic R.P. 5337; Foromacidins; Sequamycin)	<i>Streptomyces ambofaciens</i>	I: C <sub>45</sub> H <sub>78</sub> N <sub>2</sub> O <sub>15</sub> ; 886 (calc.) II: C <sub>47</sub> H <sub>80</sub> N <sub>2</sub> O <sub>16</sub> ; 928 (calc.) III: C <sub>48</sub> H <sub>82</sub> N <sub>2</sub> O <sub>16</sub> ; 942 (calc.)	Basic mac- rolides	Complex; amorphous white powder
26 Streptomycin	<i>Streptomyces</i> spp.	C <sub>21</sub> H <sub>39</sub> N <sub>7</sub> O <sub>12</sub> ; 581.58 (calc.)	Strongly basic	Reineckate: thin plates Helianthate: dark red crystals HCl: white amorphous powder Tri-HCl: monoclinic prisms
27 Streptovaricin <sup>14</sup>	<i>Streptomyces spectabilis</i>			Partially crystalline; bright orange-yellow

<sup>12</sup>/ Two components distinguishable by paper chromatography; B is three-to-four times as active as A [98]. <sup>13</sup>/ The five components [110].

**ANTIBIOTICS**

**CHEMICAL CHARACTERISTICS**

Melting Point °C	Solubility	Stability	Reference	
(F)	(G)	(H)	(I)	
Base: 110 d. Chloroform sol- vate: 120-121	Base: s. most organic solvents, acidic media; sl.s. water, ligroin HCl: v.s. water	Stable at room temperature pH 2-9 in aqueous solution; unstable when heated at acid pH	B,112;C,49,72, 113;D,49,72;E, 49,113;F,G,49; H,112,113	13
	v.s. water; l.s. methanol; sl.s. absolute ethanol; i. other organic solvents	Stable in aqueous solutions at pH 1.5-10.0	B,7,55,61;C,D,7, 104;E,55;G,7,55; H,34	14
	Acids: s. alcohols, ketones, ethers, esters, aromatic hydrocarbons; sl.s. water; i. aliphatic hydrocarbons Na salts: s. water, methanol, ethanol; sl.s. dry butanol, ketones, ethyl ace- tate		24,53	15
202 d.	Free acid: sl.s. water; more s. in al- kaline solutions	Acid stable; unstable to penicillin- ase	B,C,E,F,39;G,H, 77	16
Na salt: 215 d.		Labile to acids, heat, penicillinase	24,53	17
	i. most organic solvents Ba salt: s. water; sl.s. methanol; i. ethanol	Unstable at room temperature be- low pH 4 and above 9; unstable at pH 7 in presence of heavy metal ions; unstable to penicillinase	B,D,1,3;C,1,91;E, 2;G,2,3;H,3,91	18
		Unstable to acid; stable to penicil- linase	B,H,37,76,77;C, 37;E,37,103	19
Hydrated Na salt: 188 d.		Acid stable; stable to penicillinase	38	20
K salt: 230-231 d.		Acid stable; unstable to penicillin- ase	B,C,78,97;E,F,97; H,78	21
K salt: 263 d.		Acid stable; unstable to penicillin- ase	B,53,95;C,109;F, 108;H,56,95	22
228-235 d.	Base: sl.s. water; i. alcohol HCl: s. water, methanol; l.s. higher al- cohols; i. ether, esters, hydrocarbons, chlorinated solvents	Thermo- and acid-stable; alkali unstable	B,4;C,14,50,63;D, 22;E,100;F,G, 14,23;H,23	23
	s. acidic aqueous solution; l.s. at neu- tral pH; i. organic solvents Bases are less water-soluble than salts	Stable at acid pH; unstable at alka- line pH	98	24
I: 134-137 II: 130-133 III: 128-131	Base: s. most organic solvents; sl.s. water Sulfate: s. water, lower aliphatic alco- hols	Biological activity lost on acid hy- drolysis	B,E,G,99;C,72,96; D,72,99;F,H,96	25
Reineckate: 164- 165 d. Helianthate: 220- 226 d. Tri-HCl: gradual decomposition without melting	s. water; sl.s. lower alcohols; i. other organic solvents	Stable at pH 3-7; less stable to heat, acid, alkali	123	26
144-147 d.	v.s. N,N-dimethylformamide, 95% ethanol; s. methanol, butanol, lower ketones, methylene chloride, chloro- form, lower acetates; sl.s. water; i. hexane, ether, carbon tetrachloride, benzene	Biological activity destroyed in al- kaline solution	B,110;E-H,129	27

three components I, II, and III are identical with formacidins A, B, and C [96]. /14/ A complex containing at least

*continued*

Part I. PHYSICAL AND

Antibiotic (Synonym)	Source	Molecular Formula and Weight	Nature	Crystal Form and Color
(A)	(B)	(C)	(D)	(E)
28 Tetracyclines Chlortetracycline	<i>Streptomyces aureofaciens</i>	C <sub>22</sub> H <sub>23</sub> N <sub>2</sub> O <sub>8</sub> Cl; 478.88 (calc.)	Amphoteric	Base: yellow, acicular to bladed crystals HCl: rhomboid; clear vitreous lemon-yellow
29 Demethylchlortetracycline	<i>Streptomyces aureofaciens</i>	C <sub>21</sub> H <sub>21</sub> N <sub>2</sub> O <sub>8</sub> Cl; 464.6 (calc.)	Amphoteric	Yellow crystals
30 Oxytetracycline	<i>Streptomyces rimosus</i> , <i>S. hygroscopicus</i>	C <sub>22</sub> H <sub>24</sub> N <sub>2</sub> O <sub>9</sub> ; 460.43 (calc.)	Amphoteric	Anhydrous base: pale yellow substance Dihydrate: thick hexagonal plates or thick needles HCl: bright yellow needles from methanol or platelets from water
31 Tetracycline	<i>Streptomyces aureofaciens</i> ; catalytic dehalogenation	C <sub>22</sub> H <sub>24</sub> N <sub>2</sub> O <sub>8</sub> ; 444.43 (calc.)	Amphoteric	Trihydrate: yellow orthorhombic or equant to tabular crystals
32 Tyrothricin <sup>15</sup> Gramicidin	Component of tyrothricin	C <sub>148</sub> H <sub>210</sub> N <sub>30</sub> O <sub>26</sub> ; 2826 (calc.)	Mixture of neutral polypeptides	Colorless platelets with pointed or rectangular ends
33 Tyrocidine	Component of tyrothricin	Tyrocidine A: C <sub>66</sub> H <sub>86</sub> N <sub>13</sub> O <sub>13</sub> ; 1270 (actual) Tyrocidine B: C <sub>68</sub> H <sub>88</sub> N <sub>14</sub> O <sub>13</sub> ; 1346 (calc.)	Mixture of basic polypeptides	HCl: fine colorless needles or rods
34 Vancomycin	<i>Streptomyces orientalis</i>	HCl: 3,300 (estimated from sedimentation data) Base: 785 (minimum molecular wt) Sulfate: 2,013-2,238 (minimum molecular wt)	Amphoteric	HCl: white solid Base: crystalline rosettes

<sup>15</sup>/ Composed of approximately 20% gramicidin and 80% tyrocidine, which in turn are mixtures of polypeptides [71];

**Contributor:** Porter, John N.

**References:** [1] Abraham, E. P. 1962. *Pharmacol. Rev.* 14:473. [2] Abraham, E. P., G. G. Newton, and C. W. Brown, and G. Brownlee. 1947. *Ibid.* 160:263. [5] American Cyanamid Company. 1957. *Brit. Patent* 775,115. [8] Ashton, G. C., and A. Rhodes. 1955. *Chem. Ind. (London)*, p. 1183. [9] Barry, G. T., J. D. Gregory, and Clin. Invest. 28:1051. [12] Battersby, A. R., and L. C. Craig. 1952. *J. Am. Chem. Soc.* 74:4019. [13] Battersby, [15] Bennett, R. E., J. F. Dudley, and M. W. Shepard. 1951. *Ind. Eng. Chem.* 43:1488. [16] Bohonos, N., et al. P. J. Curtis, and H. G. Hemming. 1946. *Brit. Mycol. Soc. Trans.* 29:173. [19] Broschard, R. W., et al. 1949. *Trans. N. Y. Acad. Sci.*, II, 19:447. [22] Brownlee, G. 1949. *Ann. N. Y. Acad. Sci.* 51:875. [23] Catch, J. R., *The chemistry of penicillin.* Princeton Univ. Press, Princeton. [25] Conover, L. H. 1955. *U.S. Patent* 2,699,054. Crooks, Jr. 1949. *Ibid.* 71:2463. [28] Craig, L. C., J. D. Gregory, and G. T. Barry. 1949. *J. Clin. Invest.* 28:1014. Konigsberg. 1957. *J. Org. Chem.* 22:1345. [31] Cron, M. J., et al. 1958. *Ann. N. Y. Acad. Sci.* 76:27. [32] Cron, 43:495. [34] Davisson, J. W., I. A. Solomons, and T. M. Lees. 1952. *Antibiot. Chemotherapy* 2:460. [35] Day, *Ann. N. Y. Acad. Sci.* 51:218. [37] Doyle, F. P., J. H. C. Nayler, and G. N. Rolinson. 1960. *U.S. Patent* 2,951,839.

## ANTIBIOTICS

### CHEMICAL CHARACTERISTICS

Melting Point °C	Solubility	Stability	Reference	
(F)	(G)	(H)	(I)	
Base: 168-169 d. HCl: decomposes above 210	Base: v.s. aqueous solutions above pH 8.5, dioxane, pyridine, cellosolves, carbitol; l.s. methanol, ethanol, butanol, acetone, ethyl acetate, benzene; i. ether, petroleum ether HCl: s. water, methanol; sl.s. ethanol	Most stable at pH 2.5; less stable at neutral and alkaline pH	B,41;C,114;D,19,25,42,102;E,42;F,G,19,42;H,36,52,102	28
Base: 170-175 d. Sesquihydrate: 174-178 d.	More water-soluble than other tetracyclines	More stable than other tetracyclines at acid pH, and oxytetracycline and chlortetracycline at alkaline pH	B,F,81,82;C,E,G,H,82;D,19,25,42,102	29
Anhydrous base: 184.5-185.5 d. Dihydrate: 181-182 HCl: 190-194	Anhydrous base: s. methanol, ethanol, acetone, propylene glycol, dioxane; sl.s. water, butanol, 90% aqueous acetone, 95% methanol; i. ether, petroleum ether HCl: s. acetone, methanol, ethanol, other polar organic solvents; i. ether, petroleum ether	Most stable at pH 2.5; less stable at neutral and alkaline pH	B,52,117;C,102;D,19,25,42,102;E-G,52,102;H,36,52,102	30
Anhydrous base: 160-168 Trihydrate: 170-175 d. HCl: 214 d.	Trihydrate: v.s. methanol; s. ethanol, butanol, ethyl acetate, chloroform; sl.s. water, benzene, ether; i. petroleum ether HCl: s. water; l.s. methanol, ethanol; i. ether, hydrocarbons	More stable than chlortetracycline and oxytetracycline; most stable at acid pH	B,25,86,114;C,25,26;D,19,25,42,102;E,5,87;F,17,25,26;G,16;H,86	31
228-248 (depends on last solvent present during drying)	s. chloroform, benzene, ethanol, acetone, hot ethyl acetate, 10% HCl; sl.s. water, ether; i. petroleum ether, dilute mineral acids, dilute alkali	Thermostable	B-E,G,H,71;F,92	32
240 d.	s. methanol, ethanol, acetic acid, pyridine (especially in presence of water); sl.s. water, dry acetone, dioxane; i. ether, hydrocarbons, chloroform, electrolytes	Thermostable	B,F-H,71;C,13,75,94;D,E,12;F-H,71	33
	HCl: s. water; l.s. aqueous methanol; i. higher alcohols, acetone, ether	Most stable at pH 3-5; unstable at 37°C in glycine buffer at other than pH 3-5; 10% loss in six months at 5°C and pH 3-5	B,D,G,83;C,H,66;E,66,83	34

*Bacillus brevis* is source of tyrothricin [40].

Hale. 1954. *Biochem. J.* 58:94. [3] Abraham, E. P., et al. 1953. *Nature* 171:343. [4] Ainsworth, G. C., A. M. [6] Anker, H. S., et al. 1948. *J. Bacteriol.* 55:249. [7] Arcamone, F., et al. 1959. *Giorn. Microbiol.* 7:251. L. C. Craig. 1948. *J. Biol. Chem.* 175:485. [10] Bartz, Q. R. 1948. *Ibid.* 172:445. [11] Bartz, Q. R. 1949. *J. A. R.*, and L. C. Craig. 1952. *Ibid.* 74:4023. [14] Bell, P. H., et al. 1949. *Ann. N. Y. Acad. Sci.* 51:897. 1953-54. *Antibiot. Ann.*, p. 49. [17] Boothe, J. H., et al. 1953. *J. Am. Chem. Soc.* 75:4621. [18] Brian, P. W., *Science* 109:199. [20] Brossi, A., et al. 1960. *Helv. Chim. Acta* 43:2071. [21] Brown, R., and E. L. Hazen. 1957. T. S. G. Jones, and S. Wilkinson. 1949. *Ibid.* 51:917. [24] Clarke, H. T., J. R. Johnson, and R. Robinson. 1949. [26] Conover, L. H., et al. 1953. *J. Am. Chem. Soc.* 75:4622. [27] Controulis, J., M. C. Rebstock, and H. M. [29] Craig, L. C., W. Hausmann, and J. R. Weisiger. 1953. *J. Biol. Chem.* 200:765. [30] Craig, L. C., and W. M. J., et al. 1958. *J. Am. Chem. Soc.* 80:752. [33] Dautrevaux, M., and G. Biserte. 1961. *Bull. Soc. Chim. Biol.* A. C., J. Nabney, and A. I. Scott. 1960. *Proc. Chem. Soc.*, p. 284. [36] Dornbush, A. C., and E. J. Pelcak. 1948. [38] Doyle, F. P., et al. 1961. *Nature* 192:1183. [39] Doyle, F. P., et al. 1962. *J. Chem. Soc.* p. 1440.

*continued*



94. ANTIBIOTICS

Part I. PHYSICAL AND CHEMICAL CHARACTERISTICS

- [40] Dubos, R. J. 1939. *J. Exptl. Med.* 70:1. [41] Duggar, B. M. 1948. *Ann. N. Y. Acad. Sci.* 51:177. [42] Duggar, B. M. 1949. U.S. Patent 2,482,055. [43] Dutcher, J. D., G. Boyack, and S. Fox. 1953-54. *Antibiot. Ann.*, p. 191. [44] Dutcher, J. D., et al. 1953. *Antibiot. Chemotherapy* 3:910. [45] Dutcher, J. D., et al. 1956-57. *Antibiot. Ann.*, p. 866. [46] Dutcher, J. D., et al. 1959. U.S. Patent 2,908,611. [47] Dutcher, J. D., et al. 1959. U.S. Patent 2,908,612. [48] Ehrlich, J., et al. 1947. *Science* 106:417. [49] Els, H., W. D. Celmer, and K. Murai. 1958. *J. Am. Chem. Soc.* 80:3777. [50] Few, A. V., and J. H. Schulman. 1953. *Biochem. J.* 54:171. [51] Finlay, A. C., and P. P. Regna. 1953. *Intern. Congr. Microbiol.*, 6th, Rome, Symp. 4:58. [52] Finlay, A. C., et al. 1950. *Science* 111:85. [53] Florey, H. W., et al. 1949. *Antibiotics*. Oxford Univ. Press, London. [54] Flynn, E. H., et al. 1954. *J. Am. Chem. Soc.* 76:3121. [55] Frohardt, R. P., et al. 1959. U.S. Patent 2,916,485. [56] Goodey, R., K. N. Reed, and J. Stephens. 1955. *J. Pharm. Pharmacol.* 7:692. [57] Gottlieb, D., et al. 1948. *J. Bacteriol.* 55:409. [58] Granatek, A. P., S. Duda, and F. H. Buckwalter. 1960. *Antibiot. Chemotherapy* 10:148. [59] Grove, J. F., et al. 1952. *J. Chem. Soc.*, p. 3949. [60] Gruhzt, O. M., et al. 1949. *J. Clin. Invest.* 28:943. [61] Hagemann, G., et al. 1962. U.S. Patent 3,052,605. [62] Haight, T. H., and M. Finland. 1952. *New Engl. J. Med.* 247:227. [63] Hausmann, W., and L. C. Craig. 1954. *J. Am. Chem. Soc.* 76:4892. [64] Hausmann, W., J. R. Weisiger, and L. C. Craig. 1955. *Ibid.* 77:721. [65] Hazen, E. L., and R. Brown. 1951. *Proc. Soc. Exptl. Biol. Med.* 76:93. [66] Higgins, H. M., et al. 1957-58. *Antibiot. Ann.*, p. 906. [67] Hinman, J. W., et al. 1956. *J. Am. Chem. Soc.* 78:1072. [68] Hochstein, F. A., and K. Murai. 1954. *Ibid.* 76:5080. [69] Hoeksema, H., J. L. Johnson, and J. W. Hinman. 1955. *Ibid.* 77:6711. [70] Hoeksema, H., et al. 1956. *Antibiot. Chemotherapy* 6:143. [71] Hotchkiss, R. D. 1944. *Advan. Enzymol.* 4:153. [72] Hütter, R., W. Keller-Schierlein, and H. Zähler. 1961. *Arch. Mikrobiol.* 39:158. [73] Johnson, B. A., H. Anker, and F. L. Meleney. 1945. *Science* 102:376. [74] Kaczka, E. A., et al. 1955. *J. Am. Chem. Soc.* 77:6404. [75] King, T. P., and L. C. Craig. 1956. *Ibid.* 77:6624. [76] Knox, R. 1961. *Nature* 192:492. [77] Knudsen, E. T. 1963. *Antibiot. Chemotherapia* 11:118. [78] Knudsen, E. T., and G. N. Rolinson. 1959. *Lancet* 2:1105. [79] Koyama, Y. 1957. *Giorn. Ital. Chemioterap.* 4:279. [80] Long, L. M., and H. D. Troutman. 1949. *J. Am. Chem. Soc.* 71:2469. [81] McCormick, J. R. D., et al. 1957. *Ibid.* 79:4561. [82] McCormick, J. R. D., et al. 1959. U.S. Patent 2,878,289. [83] McCormick, M. H., et al. 1955-56. *Antibiot. Ann.*, p. 606. [84] McGuire, J. M., et al. 1952. *Antibiot. Chemotherapy* 2:281. [85] Maeda, K., et al. 1957. *J. Antibiotics (Tokyo)*, A, 10:228. [86] Minieri, P. P., et al. 1953-54. *Antibiot. Ann.*, p. 81. [87] Minieri, P. P., et al. 1956. U.S. Patent 2,734,018. [88] Mullins, J. D., and T. J. Macek. 1960. *J. Am. Pharm. Assoc. Sci. Ed.* 49:245. [89] Murase, M. 1961. *J. Antibiotics (Tokyo)*, A, 14:367. [90] Murase, M., et al. 1961. *Ibid.*, A, 14:156. [91] Newton, G. G., and E. P. Abraham. 1954. *Biochem. J.* 58:103. [92] Olesen, P. E., and L. Szabo. 1959. *Nature* 183:749. [93] Oxford, A. E., H. Raistrick, and P. Simonart. 1939. *Biochem. J.* 33:240. [94] Paladini, A., and L. C. Craig. 1954. *J. Am. Chem. Soc.* 76:688. [95] Parker, G., R. J. Cox, and D. Richards. 1955. *J. Pharm. Pharmacol.* 7:683. [96] Paul, R., and S. Tchelitcheff. 1957. *Bull. Soc. Chim. France*, Ser. 5, p. 443. [97] Perron, Y. G., et al. 1959-60. *Antibiot. Ann.*, p. 107. [98] Philip, J. E., J. R. Schenck, and M. P. Hargie. 1956-57. *Ibid.*, p. 699. [99] Pinnert-Sindico, S., et al. 1954-55. *Ibid.*, p. 724. [100] Porter, J. N., et al. 1949. *Ann. N. Y. Acad. Sci.* 51:857. [101] Rebstock, M. C., et al. 1949. *J. Am. Chem. Soc.* 71:2458. [102] Regna, P. P., et al. 1951. *Ibid.* 73:4211. [103] Rolinson, G. N., et al. 1960. *Lancet* 2:564. [104] Schillings, R. T., and C. P. Schaffner. 1961. *Intersci. Conf. Antimicrobial Agents Chemotherapy, Proc.*, p. 274. [105] Schmitz, H., et al. 1958. *J. Am. Chem. Soc.* 80:2911. [106] Schwartz, B. S., et al. 1959-60. *Antibiot. Ann.*, p. 41. [107] Seneca, H. 1955-56. *Ibid.*, p. 697. [108] Sheehan, J. C., and K. R. Henery-Logan. 1957. *J. Am. Chem. Soc.* 79:1262. [109] Sheehan, J. C., and K. R. Henery-Logan. 1959. *Ibid.* 81:3089. [110] Siminoff, P., et al. 1957. *Am. Rev. Tuberc. Pulmonary Diseases* 75:576. [111] Smith, C. G., et al. 1956. *Antibiot. Chemotherapy* 6:135. [112] Sobin, B. A., A. R. English, and W. D. Celmer. 1954-55. *Antibiot. Ann.*, p. 827. [113] Sobin, B. A., J. B. Routien, and T. M. Lees. 1956. U.S. Patent 2,757,123. [114] Stephens, C. R., et al. 1952. *J. Am. Chem. Soc.* 74:4976. [115] Tanner, F. W., Jr., et al. 1952. *Antibiot. Chemotherapy* 2:441. [116] Taub, D., C. H. Kuo.

*continued*

## 94. ANTIBIOTICS

### Part I. PHYSICAL AND CHEMICAL CHARACTERISTICS

and N. L. Wendler. 1962. Chem. Ind. (London), p. 1617. [117] Tresner, H. D., and E. J. Backus. 1956. Appl. Microbiol. 4:243. [118] Udagawa, I., and S. Abe. 1961. J. Antibiotics (Tokyo), A, 14:215. [119] Umezawa, H., et al. 1957. Ibid., A, 10:181. [120] Vandeputte, J., J. L. Wachtel, and E. T. Stiller. 1955-56. Antibiot. Ann., p. 587. [121] Wagner, R. L., et al. 1953. J. Am. Chem. Soc. 75:4684. [122] Wakazawa, T., et al. 1961. J. Antibiotics (Tokyo), A, 14:180. [123] Waksman, S. A. 1949. Streptomycin; nature and practical applications. Williams and Wilkins, Baltimore. [124] Waksman, S. A. 1953. Neomycin: nature, formation, isolation, and practical application. Rutgers Univ. Press, New Brunswick. [125] Waksman, S. A., ed. 1958. Neomycin; its nature and practical application. Williams and Wilkins, Baltimore. [126] Waksman, S. A., E. Katz, and L. C. Vining. 1958. Proc. Natl. Acad. Sci. U.S. 44:602. [127] Waksman, S. A., and H. A. Lechevalier. 1962. The actinomycetes. Williams and Wilkins, Baltimore. v. 3. [128] Wallick, H., et al. 1955-56. Antibiot. Ann., p. 909. [129] Whitfield, G. B., et al. 1957. Am. Rev. Tuberc. 75:584. [130] Wiley, P. F., et al. 1957. J. Am. Chem. Soc. 79:6062. [131] Wiley, P. F., et al. 1957. Ibid. 79:6074. [132] Wilkinson, S. 1963. Lancet 1:922. [133] Woodward, R. B. 1957. Angew. Chem. 69:50.

### Part II. BIOLOGICAL ACTIVITY

Antibiotic (A)	Biological Activity (B)	Clinical Results (C)
1 Actinomy- cins	<b>In vitro:</b> Inhibitory to gram-positive bacteria; less active against mycobacteria; virtually inactive against gram-negative bacteria and fungi. <b>In vivo:</b> Effectiveness in animals is limited by high toxicity, but there are several reports of activity on tumors. Many actinomycins cause splenic atrophy in animals after multiple doses. [54]	Short remissions have been obtained in Hodgkin's disease and chronic lymphatic leukemia with actinomycin C. Diarrhea, mucosal inflammation, alopecia, and liver damage have been noted. [3] Actinomycin D has produced a number of temporary remissions in carcinoma of the breast, malignant melanoma, and lymphosarcoma. Toxicity is similar to that produced with actinomycin C. [51]
2 Amphoteri- cin B	<b>In vitro:</b> An antifungal agent which is most active against fungi possessing a yeast phase (apparently acts against oxidative metabolic processes). Activity, which is markedly influenced by pH, is reduced above pH 7.5. Inhibits <i>Blas-tomyces dermatitidis</i> , <i>Candida albicans</i> , <i>Clado-sporium trichoides</i> , <i>Coccidioides immitis</i> , <i>Cryptococcus neoformans</i> , <i>Histoplasma capsu-latum</i> , <i>Paracoccidioides brasiliensis</i> and the yeast phase only of <i>Sporotrichum schenckii</i> . <b>In vivo:</b> Biological cures have been limited mostly to cases in which drug administration occurred with, or shortly after, inoculation. [34,43,47]	Useful in treating such systemic mycoses as histoplasmosis, blastomycosis, coccidioidomycosis, and cryptococcosis. Usefulness in moniliasis has not been adequately established. Preferred route of administration is intravenous, but intrathecal or local instillations may be employed. Poor absorption from the gastrointestinal tract precludes oral administration. Mild reversible azotemia, anorexia, headache, chills and fever are common toxic manifestations. Other effects include anemia, intestinal disturbance, and rash. [34,43,47]
3 Bacitracins	<b>In vitro:</b> In general, active against gram-positive bacteria (especially micrococci); little or no activity against gram-negative organisms, <i>Bacillus subtilis</i> , or fungi. Little evidence of cross-resistance with other antibiotics. Bactericidal. <b>In vivo:</b> Controlled staphylococcal meningitis in dogs, <i>Treponema pallidum</i> in rabbits, clostridial infections in guinea pigs (if promptly administered), and pinworms in mice. [13,31]	Used in the topical therapy of superficial infections of the skin, mucous membranes, eye, ear, etc. Frequently combined with polymyxin or neomycin for specific drug-resistant staphylococcal sepsis. Amoebic colitis, dysentery due to various organisms, and pinworms treated successfully by oral administration. Nephrotoxicity and pain at site generally result from parenteral treatments. [13,23]
4 Carbomy- cin	<b>In vitro:</b> Bacteriostatic. Active mainly against gram-positive and a few gram-negative bacteria, large viruses, rickettsiae, and certain protozoa. Cross-resistance with erythromycin and other macrolides. <span style="float: right;"><i>continued</i></span>	Useful in infections caused by gram-positive organisms resistant to penicillin, but seems to have no advantage over erythromycin in common pyogenic infections. Usually administered orally. Low toxicity, similar to that produced by erythromycin. [4,13]

*continued*

## 94. ANTIBIOTICS

### Part II. BIOLOGICAL ACTIVITY

Antibiotic (A)	Biological Activity (B)	Clinical Results (C)
Carbomycin (See preceding page)	<b>In vivo:</b> Good protection against <i>Diplococcus pneumoniae</i> , <i>Staphylococcus aureus</i> , <i>Pasteurella multocida</i> , <i>Clostridium tetani</i> , many rickettsiae, the viruses of psittacosis, ornithosis, lymphogranuloma venereum, human and feline pneumonitis, and sporadic encephalomyelitis. Inactive against <i>Bacillus anthracis</i> , <i>Mycobacterium tuberculosis</i> . [13]	
5 Chloramphenicol	<b>In vitro:</b> Inhibits protein synthesis and is primarily bacteriostatic. Active against gram-positive and gram-negative bacteria, rickettsiae, and large viruses; no activity against fungi. Active against <i>Borrelia</i> spp., <i>Entamoeba histolytica</i> , <i>Tritrichomonas foetus</i> . <b>In vivo:</b> Active in a wide variety of infections caused by the organisms specified under "in vitro." [21,56]	Agent of choice in typhoid fever and some types of meningitis, notably <i>Haemophilus influenzae</i> meningitis. Commonly used in systemic infections caused by staphylococci, particularly those resistant to other drugs. Useful in treating rickettsial diseases, infections caused by resistant gram-negative enteric bacteria, and severe dysenteries. Administered orally or parenterally. Toxic manifestations have included gastrointestinal upset, hypersensitivity, and blood dyscrasias. [21,56]
6 Colistin	<b>In vitro:</b> Primarily bactericidal. Highly inhibitory for most strains of the coli-aerogenes, <i>Pseudomonas</i> , <i>Salmonella</i> , and <i>Shigella</i> groups, being more active than polymyxin B. Inhibits some species of <i>Candida</i> and certain other fungi. Most strains of <i>Proteus</i> , <i>Neisseria</i> , and gram-positive organisms are resistant. <b>In vivo:</b> Colistinmethanesulfonate is less toxic in mice, but not in rats, than the sulfate. Both compounds are highly effective orally or parenterally against <i>Escherichia coli</i> or <i>Klebsiella pneumoniae</i> infections in mice. [42,57]	Colistin sulfate, administered orally, is particularly effective in infectious infant enteritis. Intramuscular administration of colistinmethanesulfonate has given encouraging results in pertussis, influenza, meningitis, urinary tract infections, gram-negative septicemias, and endocarditis when caused by sensitive organisms. The most potentially serious toxic effect is renal damage. [35,41]
7 Erythromycin	<b>In vitro:</b> Bacteriostatic. Active primarily against gram-positive bacteria, but a few gram-negative organisms, certain rickettsiae, and large viruses are also susceptible. Active against <i>Entamoeba histolytica</i> and <i>Trichomonas vaginalis</i> . <b>In vivo:</b> Good protection in infections produced by gram-positive organisms, large viruses, rickettsiae, oxyurids, and <i>E. histolytica</i> . Rapid increase in resistance noted. Cross-resistance to a varying degree with other macrolides. [13,18]	Effective in streptococcal, pneumococcal, and staphylococcal infections, especially those caused by strains resistant to penicillin. Useful in patients allergic to penicillin. Occasional use in <i>Haemophilus</i> infections, venereal diseases, pertussis, diphtheria, and amoebiasis. Usually administered by the oral route. Has a low order of toxicity, but gastrointestinal side effects limit high oral dosage. Jaundice has occasionally been observed. [13,18]
8 Griseofulvin	<b>In vitro:</b> An antifungal agent with no activity against bacteria. The majority of fungi are sensitive, including most human pathogens, notable exceptions being <i>Candida albicans</i> , <i>Saccharomyces cerevisiae</i> , <i>Torulopsis utilis</i> , and oomycetes. Mycostatic. <b>In vivo:</b> When administered orally, active in dogs and guinea pigs infected with <i>Trichophyton rubrum</i> and <i>Microsporum canis</i> . Presence of griseofulvin in hair shafts was demonstrated. [8,28]	Oral administration useful in the treatment of superficial infections of the hair, nails, and skin. Localizes and concentrates in these keratinized tissues. Cures may require long, continued dosage. Ineffective in deep mycoses. Very low toxicity, but occasional mild gastrointestinal upset, diarrhea, headache, and drug rashes may occur. [8,28,40]
9 Kanamycins	<b>In vitro:</b> Bactericidal. Active against most strains of staphylococci, <i>Vibrio</i> , <i>Salmonella</i> , <i>Shigella</i> , mycobacteria, coliforms, <i>Proteus</i> , and some strains of <i>Pseudomonas</i> . Inactive against streptococci, pneumococci, anaerobes, yeasts, and fungi. Some cross-resistance with paromomycin and neomycin, less cross-resistance with streptomycin. <b>In vivo:</b> Effective in protecting mice against <i>Klebsiella pneumoniae</i> , <i>Diplococcus pneumoniae</i> , <i>Proteus vulgaris</i> , and <i>Staphylococcus aureus</i> ; no protection against virulent strains of <i>Streptococcus pyogenes</i> . Active against tuberculosis in guinea pigs. [17,22,58-60]	Useful in infections caused by staphylococci, <i>Escherichia coli</i> , <i>Proteus</i> , <i>Klebsiella</i> , and some strains of <i>Pseudomonas</i> . Therapy usually correlates well with sensitivity tests. Used orally to obtain preoperative sterilization of the bowel. Commonest parenteral administration is by the intramuscular route. Usefulness in tuberculosis is limited by rapid development of resistance and by ototoxicity. Other side effects include nephrotoxicity, eosinophilia, pain at site of injection, and rashes. The kanamycin in present use is primarily A, which is less toxic than B. [59,60]

continued



## 94. ANTIBIOTICS

### Part II. BIOLOGICAL ACTIVITY

Antibiotic	Biological Activity	Clinical Results
(A)	(B)	(C)
10	<p><b>Neomycins</b></p> <p><b>In vitro:</b> Bactericidal. Active primarily against gram-positive cocci, gram-negative rods, acid-fast bacilli, and actinomycetes. More active against staphylococci than streptococci. Little or no activity against yeasts, filamentous fungi, viruses, or protozoa.</p> <p><b>In vivo:</b> Protected mice and chick embryos from lethal doses of <i>Staphylococcus aureus</i> and <i>Salmonella typhosa</i>; effective in tuberculosis in guinea pigs, and in <i>Klebsiella pneumoniae</i>, <i>Proteus vulgaris</i>, <i>Pseudomonas aeruginosa</i>, and cholera infections in mice. [13,53,60]</p>	<p>Employed orally for preoperative bowel preparation, management of liver failure, and management of gastroenteritis due to <i>Escherichia coli</i>. A topical solution or ointment is used for superficial infections. Not used parenterally. Intraperitoneal or intrapleural injections may cause respiratory arrest or insufficiency. Ototoxic and nephrotoxic. Oral treatment may cause gastrointestinal distress. [13,53,60]</p>
11	<p><b>Novobiocin</b></p> <p><b>In vitro:</b> Bactericidal. Staphylococci, including strains resistant to other antibiotics, are uniformly susceptible. Streptococci, while generally sensitive, are more variable. Gram-negative bacteria, except for strains of <i>Haemophilus</i>, are generally insensitive. Activity reduced in presence of serum. Rapid induction of resistance.</p> <p><b>In vivo:</b> Protected mice in acute and chronic staphylococcal infections, but less active than erythromycin in streptococcal and pneumococcal infections. Also active in mice against <i>Pasteurella multocida</i>, <i>Proteus vulgaris</i>, and <i>Haemophilus influenzae</i>. No activity in tuberculosis, or in fungal, rickettsial, and viral infections. [13]</p>	<p>Primary use in staphylococcal pyoderms insensitive to other antibiotics; some usefulness in pneumococcal pneumonia, undulant fever, and in genitourinary infections due to <i>Proteus</i>, <i>Staphylococcus aureus</i>, or <i>Streptococcus faecalis</i>. Calcium and sodium salts absorbed via the oral route (the route commonly used). Much patient-to-patient variation in serum binding; the dosage therefore is determined on an individual basis. Tendency for resistant strains to appear in clinical practice. Skin rash, urticaria, drug fever, and occasional leucopenia may be noted. Very high doses may induce liver damage. [13,37]</p>
12	<p><b>Nystatin</b></p> <p><b>In vitro:</b> Fungistatic and fungicidal. Active against most fungi, including <i>Candida albicans</i> and species of <i>Blastomyces</i>, <i>Coccidioides</i>, <i>Cryptococcus</i>, <i>Epidermophyton</i>, <i>Histoplasma</i>, <i>Microsporium</i>, and <i>Trichophyton</i>. Affects production of certain enzymes by sensitive organisms; also affects cell permeability. Has laboratory use in controlling yeast and mold contaminations in biological samples, culture media, tissue cultures, etc.</p> <p><b>In vivo:</b> Protected mice infected with strains of <i>Candida</i> or <i>Histoplasma</i>. [13,20]</p>	<p>Effective against various forms of moniliasis when administered by the topical or oral routes, by inhalation, or by local injection. Frequently used prophylactically with broad spectrum antibacterial antibiotics. Oral use curtailed by poor absorption. Temporary gastric upset may occur with oral treatment, and severe toxicity with injection. [13,20]</p>
13	<p><b>Oleandomycin</b></p> <p><b>In vitro:</b> Bacteriostatic. Mostly active against gram-positive bacteria. Gram-negative activity limited to <i>Haemophilus</i>, <i>Brucella</i>, and <i>Neisseria</i>. Cross-resistant with erythromycin and other macrolides.</p> <p><b>In vivo:</b> Good therapeutic activity in mice against <i>Streptococcus pyogenes</i> and <i>Staphylococcus aureus</i>. [14,48]</p>	<p>Employed in infections caused by gram-positive cocci, especially in cases of penicillin sensitivity. Administered orally or intravenously. Triacetyloleandomycin has similar use and is pharmacologically superior, but is not suitable for parenteral injection because of low solubility. Toxicity is mostly confined to gastrointestinal distress. [14]</p>
14	<p><b>Paromomycin</b></p> <p><b>In vitro:</b> Bactericidal and bacteriostatic. Broad spectrum of activity against gram-positive and gram-negative organisms and mycobacteria. Moderate activity against <i>Vibrio comma</i>, non-mammalian mycobacteria, and <i>Shigella</i> spp. Inactive against <i>Pseudomonas aeruginosa</i>.</p> <p><b>In vivo:</b> Especially effective against staphylococci and gram-negative bacilli in experimental mouse infections. Moderately tuberculostatic when given parenterally to mice and guinea pigs. Effective in amoebic infections in dogs and rats. [11,44]</p>	<p>Oral form is used for chronic and acute intestinal amoebiasis, enteric bacterial infections, preoperative bowel treatment, and management of hepatic coma. May exert a mild laxative side effect. Nephrotoxic when administered parenterally. [44,60]</p>
15	<p><b>Penicillins</b></p> <p><b>Ampicillin</b></p> <p><b>In vitro:</b> Slightly less active than penicillin G against pyogenic cocci. Active against many gram-negative bacteria, including <i>Escherichia coli</i>, <i>Haemophilus influenzae</i>, <i>Salmonella</i> spp., <i>Shigella</i> spp.; <i>Proteus</i> activity varies with the</p>	<p>Efficacious in acute and chronic respiratory infections, in many urinary infections, and in peritonitis and wound infections. May be useful in cholecystitis, meningitis, and endocarditis, especially when caused by streptococci of</p>

continued



## 94. ANTIBIOTICS

### Part II. BIOLOGICAL ACTIVITY

Antibiotic	Biological Activity	Clinical Results
(A)	(B)	(C)
16	<p>Ampicillin (See preceding page)</p> <p>strain. Inactive against <i>Aerobacter aerogenes</i>, <i>Pseudomonas pyocyanea</i>, and staphylococci resistant to penicillin G.</p> <p><b>In vivo:</b> Stable in acid medium and well-absorbed orally. Active in animals against infections produced by sensitive organisms. [26,50]</p>	<p>group D. Use in infections from <i>Proteus</i> and coliforms should be governed by sensitivity tests. Can be given orally or intramuscularly. Nontoxic but is cross-allergenic with other penicillins. [50]</p>
16	<p>Benzylpenicillin (Penicillin G)</p> <p><b>In vitro:</b> Bactericidal. In general, highly active against gram-positive bacteria. Most strains of the following are sensitive to low concentrations: <i>Bacillus</i>, <i>Clostridium</i>, <i>Corynebacterium</i>, <i>Diplococcus</i>, <i>Micrococcus</i>, <i>Streptococcus</i>, <i>Actinomyces</i>, <i>Borrelia</i>, <i>Leptospira</i>, and <i>Treponema</i>. With the exception of <i>Haemophilus</i> and <i>Neisseria</i>, most gram-negative bacteria are not sensitive. Inactive against <i>Mycobacterium</i> spp., pleuropneumonia-like organisms, yeasts, fungi, viruses, rickettsiae, protozoa. Resistance develops in a slow, stepwise manner. Destroyed by penicillinase (organisms producing penicillinase are resistant).</p> <p><b>In vivo:</b> Activity in experimental infections in general follows the "in vitro" antimicrobial spectrum. [5,12]</p>	<p>Drug of choice in infections caused by all strains and species of <i>Streptococcus</i>; pneumococci; nonpenicillinase-producing strains of staphylococci; gonococci; spirochetes, <i>Borrelia</i>, and spiral organisms of the mouth; clostridia; <i>Corynebacterium</i>; and anthrax. Aqueous, crystalline penicillin G is used for rapid effect or high serum concentrations. For longer-lasting activity (e.g., in prophylaxis of rheumatic fever or glomerulonephritis), procaine (or benzathine) penicillin G is administered intramuscularly. Usefulness limited by allergic reactions, sensitivity to penicillinase, and acid instability which precludes oral dosage. [5,30]</p>
17	<p>Cephalosporin N</p> <p><b>In vitro:</b> Bactericidal. Less active than other important penicillins against gram-positive bacteria, but shows good activity against many gram-negative bacteria, its activity against <i>Salmonella</i> being more than tenfold that of penicillin G.</p> <p><b>In vivo:</b> Protected mice from infection with <i>Salmonella typhimurium</i>, <i>S. typhosa</i>, <i>Escherichia coli</i>, <i>Proteus vulgaris</i>, and <i>P. mirabilis</i>, and chicks with <i>S. pullorum</i>. Activity in vivo sometimes greater than is indicated by "in vitro" tests. [1]</p>	<p>Effective in limited trials in the treatment and control of typhoid fever in man. Has also been used for the treatment of gonorrhea and syphilis in patients sensitive to penicillin G. [1]</p>
18	<p>Methicillin</p> <p><b>In vitro:</b> Bactericidal. As active as penicillin G against penicillin-sensitive, or resistant, staphylococci. Less active against streptococci.</p> <p><b>In vivo:</b> Activity corresponds to that indicated in "in vitro." [26]</p>	<p>Particularly useful in infections caused by penicillin-resistant staphylococci. Must be administered parenterally. Low toxicity, but shows complete cross-allergenicity with other penicillins. [26]</p>
19	<p>Oxacillin</p> <p><b>In vitro:</b> Activity resembles that of methicillin, but oxacillin is five-to-eight times more active against resistant staphylococci.</p> <p><b>In vivo:</b> Resembles penicillin V in stability, absorption, and serum level, but oxacillin binds more readily to serum than any other available penicillin. [2,25]</p>	<p>As useful as methicillin in same types of infections, but oxacillin is resistant to acid and therefore is an effective oral antibiotic against resistant staphylococci. [2,25]</p>
20	<p>Phenethicillin</p> <p><b>In vitro:</b> Has spectrum similar to that of penicillin V.</p> <p><b>In vivo:</b> As active as penicillin V against <i>Diplococcus pneumoniae</i> and <i>Streptococcus pyogenes</i> infections, and more active against <i>Staphylococcus aureus</i> infections. Not active against strains resistant to penicillin G. [16]</p>	<p>A penicillin capable of being administered orally. Its use parallels that of penicillin V, but phenethicillin is reported to give higher and more constant blood-level concentrations than does potassium penicillin V. [32]</p>
21	<p>Phenoxy-methylpenicillin (Penicillin V)</p> <p><b>In vitro:</b> Spectrum of activity is the same as for penicillin G, but V is more stable at pH of less than 6.5.</p> <p><b>In vivo:</b> Protection in pneumococcal and other infections equivalent to that produced by penicillin G. Acid stability permits good protection via the oral route. [7,9]</p>	<p>Range of activity in treating various infections is comparable to that of penicillin G. Has the advantage of oral administration. [7,9]</p>
22	<p>Polymyxins</p> <p><b>In vitro:</b> Bactericidal. Active mainly against gram-negative bacteria, including <i>Pseudomonas</i>. <i>Proteus</i> strains are often resistant. Little or no activity against fungi.</p> <p><b>In vivo:</b> Good protection in mice infected with</p>	<p>Used mainly in serious infections due to <i>Pseudomonas aeruginosa</i>, such as urinary tract infections and meningitis. Effective in gram-negative infections of body surfaces (eye, ear, skin wounds) and of body spaces (joint, pleural,</p>

continued

## 94. ANTIBIOTICS

### Part II. BIOLOGICAL ACTIVITY

Antibiotic	Biological Activity	Clinical Results
(A)	(B)	(C)
Polymyxins (See preceding page)	<i>Salmonella typhosa</i> , <i>Bordetella pertussis</i> , and <i>Klebsiella pneumoniae</i> , and in chicks infected with <i>Pasteurella multocida</i> and <i>S. gallinarum</i> . No protection against streptococci and staphylococci. [13,23]	dural). Used topically or by mouth, as in <i>Salmonella enteritis</i> . Often combined with other drugs. Parenteral use may cause neuro- and nephro-toxicity, pain at site of injection, fever, and rashes. [13,23]
23 Ristocetins	<b>In vitro:</b> Bactericidal and bacteriostatic. Specifically active against gram-positive bacteria and mycobacteria; inactive against gram-negative bacteria (including <i>Haemophilus influenzae</i> and <i>Neisseria</i> spp.), yeasts, filamentous fungi, and protozoa. No cross-resistance with other antibiotics. <b>In vivo:</b> Controlled infections of mice caused by <i>Streptococcus pyogenes</i> , <i>Staphylococcus aureus</i> , and <i>Diplococcus pneumoniae</i> . Not effective against tuberculosis in mice. [19,39]	Most useful in severe staphylococcal and enterococcal infections resistant to other antimicrobials. Useful in short-term therapy of enterococcal endocarditis. Only administered intravenously. Toxicity is directly related to size of dosage and may include disturbances of the hematopoietic system, phlebitis, fever, and rash. [39]
24 Spiramycins	<b>In vitro:</b> Bacteriostatic. Especially active on gram-positive bacteria and to a lesser extent on mycobacteria and some gram-negative bacteria. Active on rickettsiae. Cross-resistance with other macrolides. <b>In vivo:</b> Protected mice infected with hemolytic streptococci, pneumococci, and staphylococci. Activity in vivo is greater than is indicated by "in vitro" tests. [13,36]	Active in various staphylococcal infections; particularly useful against organisms resistant to other antimicrobials. Promising results in gonococcal infections. Orally administered. Toxicities are very low or nonexistent. [13]
25 Streptomycin	<b>In vitro:</b> Bactericidal. Active against gram-positive and gram-negative bacteria, including mycobacteria. No activity against fungi, rickettsiae, or viruses. Rapid development of resistance. <b>In vivo:</b> Excellent protection in a wide variety of experimental infections, including <i>Bacillus anthracis</i> , <i>Brucella abortus</i> , <i>Diplococcus pneumoniae</i> , <i>Mycobacterium tuberculosis</i> , <i>Neisseria meningitidis</i> , <i>Pasteurella pestis</i> , <i>P. tularensis</i> , <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i> , and species of <i>Haemophilus</i> , <i>Klebsiella</i> , <i>Pseudomonas</i> , <i>Salmonella</i> , <i>Shigella</i> . [45,52]	Often combined with bacteriostatic agents, such as one of the tetracyclines or sulfonamides, chloramphenicol, etc. Is used alone or in combination against tuberculosis, <i>Haemophilus influenzae</i> meningitis, subacute bacterial endocarditis, tularemia, plague, <i>Klebsiella pneumoniae</i> , or <i>Escherichia coli</i> infections. Parenteral or topical application. Use in tuberculosis is limited by rapid development of resistant strains. May cause ototoxicity, but is less likely to do so than dihydrostreptomycin or a combination of the two. [45,52]
26 Streptovaricin	<b>In vitro:</b> The streptovaricin complex and components A, B, and C are very active against gram-positive bacteria and mycobacteria, and show activity against some gram-negative bacteria and fungi. Components D and E have only low activity against gram-positive bacteria. <b>In vivo:</b> Highly effective in the tuberculous mouse, component C being the most active. Effectiveness increased when combined with isoniazid. [29,38,46]	Only slight improvement in patients with advanced pulmonary tuberculosis when administered orally (50 mg/kg daily dosage). Some gastrointestinal upset observed. Results when combined with isoniazid were not superior to those with isoniazid alone, and were inferior to isoniazid-pyrazinamide treatment. [6,29]
27 Tetracyclines <sup>1</sup>	<b>In vitro:</b> Bacteriostatic. Broad spectrum of activity against gram-positive and gram-negative bacteria. Also active against coccidia, amoebae, and balantidiasis. Superior to penicillin against <i>Bacillus anthracis</i> . Effective against rickettsiae and larger viruses. No activity against fungi. Resistance does not develop readily, but there is almost complete cross-resistance among the four major tetracyclines. <b>In vivo:</b> Good protection in laboratory animals against sensitive microorganisms in the groups listed under "in vitro." Little or no protection against <i>Mycobacterium tuberculosis</i> . No activity against small viruses or fungi. [8,10,27,33,49]	The commercially available tetracyclines are useful in infections caused by a wide range of microorganisms, including group A streptococci, staphylococci, pneumococci, meningococci, <i>Bacillus anthracis</i> , <i>Haemophilus influenzae</i> , <i>Bordetella pertussis</i> , <i>Entamoeba histolytica</i> , rickettsiae, and viruses of the lymphogranuloma-psittacosis group. Toxicity is very low and primarily confined to gastrointestinal upset, although a phototoxic reaction has been reported for demethylchlortetracycline. Dosage is usually via the oral route, although parenteral injection or topical application may be used as required. [8,10,27,33,49]

<sup>1</sup>/ Chlortetracycline, demethylchlortetracycline, oxytetracycline, and tetracycline.

continued



## 94. ANTIBIOTICS

### Part II. BIOLOGICAL ACTIVITY

Antibiotic	Biological Activity	Clinical Results
(A)	(B)	(C)
28 Tyrothricin <sup>a</sup>	<p><b>In vitro:</b> Tyrothricin is a mixture of approximately 20% gramicidin (the active component) and 80% tyrocidine. It is bacteriostatic at low concentrations and bactericidal at high (1.0 µg/ml, or more). Active against corynebacteria, diplococci, staphylococci, streptococci, lactobacilli, and some fungi.</p> <p><b>In vivo:</b> Tyrocidine loses most of its antibacterial activity in the presence of animal tissues, although it shows some effect against gram-negative cocci. Tyrothricin gives protection against pneumococci, streptococci, and staphylococci, but produces hemolysis and acute lethal toxicity upon injection. [55]</p>	Tyrothricin is employed as a solution, spray, or in troches against sensitive gram-positive bacteria. Sometimes combined with bacitracin. The toxicity of the antibiotic precludes parenteral use. Since it is effective only when in direct contact with microorganisms, it is of little or no value in deep-seated infections. [55]
29 Vancomycin	<p><b>In vitro:</b> Bactericidal. Uniformly active against pathogenic staphylococci. Also inhibits hemolytic streptococci, pneumococci, enterococci, gonococci, corynebacteria, and clostridia. Not active against most gram-negative bacteria, mycobacteria, fungi, and yeasts. Unaffected by serum and by pH of the test medium. Resistance develops slowly, and there is no cross-resistance with other antibiotics.</p> <p><b>In vivo:</b> Mice received complete protection from staphylococcal infections when vancomycin was injected subcutaneously. No protection against <i>Toxoplasma gondii</i>. [15,24,39]</p>	Used primarily in severe staphylococcal infections: pneumonia and empyema, infections of skin and soft tissues, septicemia and endocarditis, osteomyelitis, and enterocolitis. Intravenous administration. Occasional phlebitis, chills, fever, renal irritation, urticaria, macular rashes, and hearing loss, but these side effects are less common with more highly purified preparations. Rare instances have been noted of cross-allergy with other antibiotics. [15,24,39]

<sup>a</sup>/ Gramicidin and tyrocidine.

**Contributor:** Porter, John N.

- References:** [1] Abraham, E. P. 1962. *Pharmacol. Rev.* 14:473. [2] Barber, M., and P. Waterworth. 1962. *New Engl. J. Med.* 266:246. [3] Begemann, H. 1960. *Ann. N. Y. Acad. Sci.* 89:454. [4] British Medical Association. 1963. *Brit. Med. J.* 1:1213. [5] Bunn, P. A. 1961. *Pediat. Clin. N. Am.* 8:981. [6] Des Prez, R., et al. 1959. *Am. Rev. Respirat. Diseases* 80:431. [7] Diding, N. A., and A. R. Frisk. 1955-56. *Antibiot. Ann.*, p. 529. [8] Dowling, H. F. 1955. *Antibiot. Monographs* 3. [9] Elias, W. F., and H. J. Merrion. 1955-56. *Antibiot. Ann.*, p. 511. [10] Finland, M., and L. P. Garrod. 1960. *Brit. Med. J.* 2:959. [11] Fisher, M. W., et al. 1959-60. *Antibiot. Ann.*, p. 293. [12] Florey, H. W., et al. 1949. *Antibiotics*. Oxford Univ. Press, London. [13] Florey, M. E. 1960. *The clinical application of antibiotics*. Oxford Univ. Press, London. v. 4. [14] Foltz, E. L. 1961. *Pediat. Clin. N. Am.* 8:1133. [15] Geraci, J. E., et al. 1956-57. *Antibiot. Ann.*, p. 90. [16] Gourevitch, A., G. A. Hunt, and J. Lein. 1959-60. *Ibid.*, p. 111. [17] Gourevitch, A., et al. 1958. *Ann. N. Y. Acad. Sci.* 76:31. [18] Griffith, R. S., and H. R. Black. 1961. *Pediat. Clin. N. Am.* 8:1115. [19] Grundy, W. E., et al. 1956-57. *Antibiot. Ann.*, p. 687. [20] Hazen, E. L., and R. Brown. 1960. *Ann. N. Y. Acad. Sci.* 89:258. [21] Hodgman, J. E. 1961. *Pediat. Clin. N. Am.* 8:1027. [22] Hunt, G. A., and A. J. Moses. 1958. *Ann. N. Y. Acad. Sci.* 76:81. [23] Jawetz, E. 1961. *Pediat. Clin. N. Am.* 8:1057. [24] Kirby, W. M. M. 1963. *Antibiot. Chemotherapia* 11:84. [25] Kirby, W. M. M., L. S. Rosenfeld, and J. Brodie. 1962. *J. Am. Med. Assoc.* 181:739. [26] Knudsen, E. T. 1963. *Antibiot. Chemotherapia* 11:118. [27] Lepper, M. H. 1956. *Antibiot. Monographs* 7. [28] Lofferer, O., and G. Riehl. 1962. *Antibiot. Chemotherapia* 10:335. [29] McCune, R. M., et al. 1957. *Am. Rev. Tuberc. Pulmonary Diseases* 75:659. [30] Martin, W. J. 1961. *Med. Ann. District Columbia* 30:516. [31] Meloney, F. L., and B. A. Johnson. 1949. *Am. J. Med.* 7:794. [32] Morigi, E. M. E., W. B. Wheatley, and H. Albright. 1959-60. *Antibiot. Ann.*, p. 127. [33] Musselman, M. M. 1956. *Antibiot. Monographs* 6. [34] Newcomer, V. D., et al. 1959. *J. Chronic Diseases* 9:353. [35] Petersdorf, R. G., and J. J. Florde. 1963. *J. Am. Med. Assoc.* 183:123. [36] Pinnert-Sindico, S., et al. 1954-55. *Antibiot. Ann.*, p. 724. [37] Pratt, R. 1962. *J. Pharm. Sci.* 51:1.

*continued*

Part II. BIOLOGICAL ACTIVITY

- [38] Rhuland, L. E., K. F. Stern, and H. R. Reames. 1957. *Am. Rev. Tuberc. Pulmonary Diseases* 75:588. [39] Riley, H. D., Jr. 1961. *Pediat. Clin. N. Am.* 8:1073. [40] Roth, F. J., Jr. 1960. *Ann. N. Y. Acad. Sci.* 89:247. [41] Schöenberg, H. 1963. *Antibiot. Chemotherapia* 11:136. [42] Schwartz, B. S., et al. 1959-60. *Antibiot. Ann.*, p. 41. [43] Seabury, J. H., and H. E. Dascomb. 1960. *Ann. N. Y. Acad. Sci.* 89:202. [44] Shafei, A. Z. 1959. *Antibiot. Med. Clin. Therapy* 6:275. [45] Shaw, E. B., and H. B. Bruyn. 1961. *Pediat. Clin. N. Am.* 8:1013. [46] Siminoff, P., et al. 1957. *Am. Rev. Tuberc. Pulmonary Diseases* 75:576. [47] Smith, D. E. 1961. *Pediat. Clin. N. Am.* 8:1099. [48] Sobin, B. A., A. R. English, and W. D. Celmer. 1954-55. *Antibiot. Ann.*, p. 827. [49] Spitzzy, K. H. 1962. *Antibiot. Chemotherapia* 10:193. [50] Stewart, G. T. 1963. *Pharmakotherapia* 1:197. [51] Tan, C. T. C., et al. 1960. *Ann. N. Y. Acad. Sci.* 89:426. [52] Waksman, S. A. 1949. *Streptomycin: nature and practical applications*. Williams and Wilkins, Baltimore. [53] Waksman, S. A. 1953. *Neomycin: nature, formation, isolation, and practical application*. Rutgers Univ. Press, New Brunswick. [54] Waksman, S. A., and H. A. Lechevalier. 1962. *The actinomycetes*. Williams and Wilkins, Baltimore. v. 3. [55] Welch, H., and C. N. Lewis. 1953. *Antibiotic therapy*. Medical Encyclopedia, New York. [56] Woodward, T. E., and C. L. Wisseman, Jr. 1958. *Antibiot. Monographs* 8. [57] Wright, W. W., and H. Welch. 1959-60. *Antibiot. Ann.*, p. 61. [58] Yanagisawa, K., et al. 1958. *Ann. N. Y. Acad. Sci.* 76:88. [59] Yow, M. D., and H. Abu-Nassar. 1963. *Antibiot. Chemotherapia* 11:148. [60] Yow, M. D., and E. M. Yow. 1961. *Pediat. Clin. N. Am.* 8:1043.

95. ANTICOAGULANTS

Anticoagulant dosage is usually determined in vitro from the clotting time for heparin and heparinoid compounds, or the prothrombin time for indirect anticoagulants [5,20,21,27]. In vivo, the dosage required to prevent coagulation may be many times that indicated by the test in vitro [30].

**Direct Anticoagulants:** Heparin and heparinoid compounds injected intravenously give a prolonged coagulation time. For some purposes, the high peak blood levels resulting from intravenous administration are essential [26]. Heparin is inactive orally [13], and intramuscular and subcutaneous injections are not as generally effective as those given intravenously [32]. The subcutaneous route, however, is the commonly accepted method of choice in doses of 75-100 mg, given each 8-12 hours (depending on the clotting time) [32]. It is desirable that the clotting time be approximately twice that of the normal control before the next dose is administered [32]. The effect on clotting time is increased and prolonged by caronamide [29] and phosphorylated hesperidin [10]. Value for maximum clotting time can be estimated from the effect of clotting time in vitro [24]. International unit of heparin = 1/130 mg of the international standard; commercial heparin = 90-120 units/mg.

**Indirect Anticoagulants:** These drugs usually are given orally. Dosage is dependent on the technique used to detect the change in prothrombin time and on individual susceptibility to the drug. A number of individuals in various species are refractory to one or more of these drugs, as judged by the effect on prothrombin time [25]. Significantly different results are obtained with tests for coagulation factors. The action of these drugs is cumulative and can be greatly enhanced by following the initial dose with a series of smaller doses. Such administration also avoids toxic side effects. Dicumarol must be dissolved with a small amount of 5N sodium hydroxide for intravenous use. It can be given intraperitoneally in suspension in propylene glycol (10-100 mg/ml). Warfarin and tromexan are more soluble than dicumarol and can be given intraperitoneally in neutral solution.

**Route** (column B): iv = intravenous; im = intramuscular; ip = intraperitoneal; sc = subcutaneous; rec = rectal; po = oral. Values in parentheses are ranges, estimate "b" (cf. Introduction).

	Species (Common Name)	Dose and Route	Maximum Clotting Time or Prothrombin Time		Achieve- ment of Thera- peutic Effect	Re- covery Time	Remarks	Refer- ence
			Control	Experimental				
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Heparin <sup>1,2</sup>								
1	<i>Homo sapi- ens</i> (man)	1,000 units (10 mg); iv	.....	<4.5 min	.....	.....	28 hyporeactors	9
2			.....	4.5-7 min	.....	.....	50 normal reactors	
3			.....	>7 min	.....	.....	7 hyperreactors	

<sup>1/</sup> Intra-arterial injection of 1 mg heparin/100 ml of blood, plus 1.2 mg protamine/100 ml of blood into venous outflow of organ (limb, kidney), gives satisfactory local heparinization (without general heparinization) in man and dog [15]. <sup>2/</sup> To neutralize heparin in man and dog, slowly inject protamine in the amount of 1.2-2.5 x weight of heparin in blood, as determined by in vitro titration; excess of protamine will be anticagulant [7,15,22].

*continued*



# Contrails

## 95. ANTICOAGULANTS

Species (Common Name)	Dose and Route	Maximum Clotting Time or Prothrombin Time		Achieve- ment of Thera- peutic Effect	Re- covery Time	Remarks	Ref- er- ence
		Control	Experimental				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Heparin <sup>1,2</sup>							
4	<i>Homo sapi- ens</i> (man)	5,000 units; iv	15 min	115 min	.....	.....	9
5		150 mg; iv	.....	>80 min	.....	4 hr	26
6		100 mg; iv	.....	>80 min	.....	4 hr	Incoagulable for 1 hr
7		75 mg; iv	.....	80 min	.....	4 hr	Incoagulable for 1/2 hr
8		50 mg; iv	.....	55 min	.....	2 hr	.....
9		150 mg; im	12 min	30 min	.....	10 hr	.....
10		100 mg; im	12 min	27 min	.....	6 hr	.....
11		25, 50, 75 mg; im	12 min	18 min	1 hr	3-4 hr	.....
12	<i>Canis fami- liaris</i>	30 units/kg; iv	.....	34 min	.....	32 min	24
13		100 units/kg; iv	.....	>2 hr	.....	.....	.....
14	(dog)	450 units/kg; sc	.....	0.25-3.5 hr	.....	2-13 hr	20
15		2-3 units/kg/min for 2 hr; iv	20 sec	5 min	.....	4 hr	31
16		65-290 units/kg; iv	20 sec	5 min	.....	200 min	.....
17	<i>Rattus nor- vegicus</i>	1,950 units/kg; sc	.....	>24 hr	.....	6 hr	22
18	(Norway rat)	5,500 units/kg; sc	.....	>24 hr	.....	12 hr	.....
19	<i>Oryctolagus cuniculus</i>	10.5 mg/kg <sup>3</sup> (50 units/kg)	.....	.....	.....	100 min	16
20	(European rabbit)	2.5 mg/kg	.....	>60 min	.....	2 hr	3
21		1.0 mg/kg	.....	40 min	.....	1 hr	.....
Dextran Sulfate							
22	<i>Oryctolagus cuniculus</i>	21 mg/kg	.....	>80 min	.....	3 hr	3
23	(European rabbit)	17 mg/kg	.....	>80 min	.....	100 hr	.....
24		7 mg/kg	.....	>80 min	.....	50 min	.....
Thrombocid (Xylan Polysulfuric Acid)							
25	<i>Oryctolagus cuniculus</i>	30 mg/kg	.....	>60 min	.....	2.5 hr	3
26	(European rabbit)	15 mg/kg	.....	65 min	.....	1.5 hr	.....
Dicumarol (3, 3'-Methylene-bis-4-hydroxycoumarin)							
27	<i>Homo sapi- ens</i> (man)	300 mg	.....	.....	.....	.....	1
28		250 mg	.....	30 sec	.....	.....	19
29		500 mg	.....	63 sec	3-4 da	Varies	.....
30		750 mg	.....	106 sec	.....	.....	.....
31	<i>Canis fami- liaris</i> (dog)	5 mg/kg	10.7(6.7-14.7) sec	37.8(30.4-45.2) sec	.....	6 da	25
32	<i>Mesocricet- us auratus</i> (golden hamster)	5 mg/kg	.....	No effect	.....	.....	25
33	<i>Mus muscu- lus</i> (house mouse)	15 mg/kg	10.6(7.8-13.4) sec	14.0(12.4-15.6) sec	.....	24 hr	25
34	<i>Rattus nor- vegicus</i> (Norway rat)	5 mg/kg	14.0(11.4-16.6) sec	19.9(18.7-21.1) sec	.....	24 hr	25
35	<i>Oryctolagus cuniculus</i>	5 mg/kg	8.0(7.6-8.4) sec	25.4(8.4->60) sec	.....	10 da	25
36	(European rabbit)	2.5 mg	8.0(7.6-8.4) sec	10.6(6.2-15.0) sec	.....	5 da	.....
37		10 mg/kg	15-16 sec	27 sec	1 da	4 da	19
38		25 mg/kg	15-16 sec	80 sec	1 da	5 da	.....
39		50 mg/kg	15-16 sec	6.5 min	2.5 da	8 da	.....
40		100 mg/kg	15-16 sec	20 min	3 da	8 da	.....
41		0.37 mg	28 sec	32 sec	1 da	3 da	28
42		0.75 mg	28 sec	40 sec	1 2/3 da	4 2/3 da	.....
43		1.5 mg	28 sec	46 sec	2 da	8 da	.....

/1/ Intra-arterial injection of 1 mg heparin/100 ml of blood, plus 1.2 mg protamine/100 ml of blood into venous out-flow of organ (limb, kidney), gives satisfactory local heparinization (without general heparinization) in man and dog [15]. /2/ To neutralize heparin in man and dog, slowly inject protamine in the amount of 1.2-2.5 x weight of heparin in blood, as determined by in vitro titration; excess of protamine will be anticoagulant [7,15,22]. /3/ Crude heparin.

continued

## 95. ANTICOAGULANTS

Species (Common Name)	Dose and Route	Maximum Clotting Time or Prothrombin Time		Achievement of Thera- peutic Effect	Re- covery Time	Remarks	Ref- er- ence	
		Control	Experimental					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Dicumarol (3,3'-Methylene-bis-4-hydroxycoumarin)								
44	<i>Oryctolagus cuniculus</i> (European rabbit);	3 mg	28 sec	67 sec	2.5 da	9 da	Susceptible animals only; 12.5% plasma prothrombin time	28
45	6 mg	28 sec	85 sec	3 da	10 da			
46	<i>Gallus domesticus</i> (chicken)	100 mg/kg	.....	.....	.....	.....	Minimum effective dose	27
47	300 mg/kg	.....	.....	.....	.....	.....	Effective dose	
Tromexan (3,3'-Carboxymethylene-bis-4-hydroxycoumarin)								
48	<i>Homo sapiens</i> (man)	1,200 mg	14 sec	25 sec	30 hr	56 hr	.....	6
49	1,200 mg	14 sec	32 sec	30 hr	60 hr	.....		
50	<i>Canis familiaris</i> (dog)	10 mg/kg	15 sec	40 sec	2 da	5 da	.....	17
51	40 mg/kg	15 sec	55 sec	3 da	5 da	.....		
52	<i>Oryctolagus cuniculus</i> (European rabbit)	400 mg	12 sec	30 sec	30 hr	48 hr	.....	6
53	<i>Gallus domesticus</i> (chicken);	100-500 mg/kg	.....	.....	.....	.....	Minimum effective dose	27
Warfarin (3-[ $\alpha$ -Acetylbenzyl]-4-hydroxycoumarin)								
54	<i>Homo sapiens</i> (man)	45-50 mg initial	12 sec	28 sec	.....	.....	5-10 mg/da for maintenance	8, 32
55	100 mg; rec	.....	40 sec	18-24 hr	6 da	.....	.....	14
56	<i>Rattus norvegicus</i> (Norway rat)	0.03 mg/kg/da	25-36 sec	No effect	.....	.....	.....	2
57	0.05 mg/kg/da	25-36 sec	55 sec	.....	.....	.....		
58	0.10 mg/kg/da	25-36 sec	2 min	.....	.....	.....		
59	<i>Oryctolagus cuniculus</i> (European rabbit);	40 mg/kg; ip	9.6 sec	36.7 sec	48 hr	.....	.....	18
60	5 mg/kg; ip	7.9 sec	35.2 sec	48 hr	.....	.....		
Liquamar (3-[1'-Phenylpropyl]-4-hydroxycoumarin)								
61	<i>Homo sapiens</i> (man)	18 mg	20.9 sec	40 sec	96 hr	144 hr	.....	4
62	21 mg	20.9 sec	36 sec	96 hr	168 hr	Minimum effective dose		
63	<i>Oryctolagus cuniculus</i> (European rabbit);	2.5 mg	14 sec	22 sec	48 hr	96 hr	.....	4
64	4 mg	14 sec	22 sec	72 hr	160 hr	.....		
65	<i>Gallus domesticus</i> (chicken);	50 mg/kg	.....	.....	.....	.....	Minimum effective dose	27
EDC (Ethylidene-bis-4-hydroxycoumarin)								
66	<i>Homo sapiens</i> (man)	0.5 g	.....	Great individual variation	.....	.....	0.2 g maintenance dose	11
67	<i>Oryctolagus cuniculus</i> (European rabbit);	20 mg; po	7 sec	10 sec	.....	.....	.....	11
68	30 mg; po	7 sec	27 sec	.....	.....	.....		
Phenindione (2-Phenyl-1,3-indanedione)								
69	<i>Canis familiaris</i> (dog)	50 mg/kg	9-13 sec	30 sec	26 hr	36 hr	.....	23
70	<i>Oryctolagus cuniculus</i> (European rabbit)	50 mg/kg	10-12 sec	22 sec	25 hr	40 hr	.....	23
Dipaxin (2-Diphenylacetyl-1,3-indanedione)								
71	<i>Homo sapiens</i> (man)	20 mg	.....	.....	1-2 da	6-10 da	2-4 mg maintenance dose	12

Contributors: (a) Jaques, Louis B., (b) Wright, Irving S.

References: [1] Allen, E. V. 1947. J. Am. Med. Assoc. 134:323. [2] Ashwin, J., and L. B. Jaques. Unpublished.

continued

Univ. Saskatchewan, Saskatoon, Can., 1957. [3] Astrup, T., H. I. K. Flyger, and J. Gormsen. 1955. Scand. J. 1952. Glasgow Med. J. 33:225. [6] Burke, G. E., and I. S. Wright. 1951. Circulation 3:164. [7] Chargaff, E., 97:753. [9] De Takats, G. 1943. Surg. Gynecol. Obstet. 77:31. [10] Evans, J. M., I. Hsu, and T. H. Korthalis. et al. 1952. Proc. Soc. Exptl. Biol. Med. 81:678. [13] Fisher, A., and T. Astrup. 1939. Ibid. 42:81. [14] Freeman, J. Med. 255:1026. [16] Gross, P. 1929. Proc. Soc. Exptl. Biol. Med. 26:383. [17] Gruber, C. M., et al. 1951. 4:144. [19] Jansen, K. F. 1944. Dikumarin. E. Munksgaard, Copenhagen. [20] Jaques, L. B. 1950. New Engl. Best. 1938. Acta Med. Scand., Suppl. 90:190. [23] Jaques, L. B., E. Lepp, and E. Gordon. 1950. Blood Clotting L. B., et al. 1957. Arch. Intern. Pharmacodyn. 111:478. [26] Jorpes, J. E. 1950. Acta Chir. Scand. 99:476. [28] Link, K. P. 1943. Harvey Lectures 39:162. [29] Sirak, H. D., R. S. McCleery, and C. P. Artz. 1948. Surgery Med. 42:968. [32] Wright, I. S. Unpublished. New York Hospital, New York, 1959.

## 96. ANIMAL

### Part I.

For information on standardization of methods of extraction, preparation, and purification of venoms, consult alinase; E = cholinesterase; F = deoxyribonuclease; G = diaminoxidase; H = diastase; I = dipeptidase; J = endopep- P = lipase; Q = 5-nucleotidase; R = ophio-adenosine triphosphatase; S = phosphatase; T = phospholipase; U = phos- enzyme activity known not to be present.

Species (Common Name)	Distribution	Adult Length ft	Fibrinogen <sup>1</sup>		Pro- thrombin <sup>1</sup>		Enzyme Activity <sup>2</sup>	Mouse Toxicity <sup>3</sup> mg/kg
			Coag- ulate	De- stroy	Acti- vate	De- stroy		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Crotalidae <sup>4</sup>								
1 <i>Ancistrodon contortrix</i> (southern U.S. copper- head)	Eastern and southern United States	2-3½					E, N, U, N, X	LD <sub>50</sub> 25.6 LD <sub>100</sub> 53.0
2 <i>A. piscivorus</i> (eastern cottonmouth)	Southeastern United States to central Texas	3-5	-	+	-	+	D, E, K, L, N, O, Q, T, U, X, X	LD <sub>50</sub> 25.8 LD <sub>100</sub> 45.0
3 <i>Bothrops atrox</i> (fer-de- lance)	Central Mexico to eastern Argentina, Martinique, Tobago, Trinidad, & St. Lucia	4½-6½	+	-	-	+	E, F, N, R, X	LD <sub>50</sub> 62.8- 99.2
4 <i>B. jararaca</i> (jararaca)	Brazil to northern Argentina & Paraguay	3½-4½	+	-	+	-	A, E, J, L, N, R, X	LD <sub>50</sub> 2.0-22.6
5 <i>Crotalus adamanteus</i> (eastern diamondback rattlesnake)	Southeastern United States, in lowlands	4½-6½	+	+	-	-	D, E, L, N, Q, U, X, X	LD <sub>50</sub> 14.5
6 <i>C. atrox</i> (western dia- mondback rattlesnake)	Southwestern United States, northern Mexico	3½-5½	-	+	-	-	A, E, L, N, R, X	LD <sub>50</sub> 7.5 LD <sub>100</sub> 19.0
7 <i>C. durissus terrificus</i> (cascabel)	Southern Mexico to Uruguay & Argentina, mostly in highlands	4-5½	+	-	±	-	E, F, L, N, O, R, X	LD <sub>50</sub> 0.6 LD <sub>50</sub> 0.1 (in- travenous)
8 <i>C. viridis</i> (prairie rattle- snake)	Western United States, southwestern Canada, northern Mexico	2½-5					E, N	LD <sub>50</sub> 7.2
9 <i>Lachesis muta</i> (bush- master)	Costa Rica to northern S. America	8-11					X	LD <sub>100</sub> 57.0
10 <i>Sistrurus catenatus</i> (east- ern massasauga)	Southern Ontario, central to southwestern United States	2-3					E, N, R	LD <sub>50</sub> 5.2 LD <sub>100</sub> 9.0

<sup>1/1</sup> Information from reference 23, unless otherwise specified in column M. <sup>1/2</sup> Presence in venom determined by  
unless otherwise stated. <sup>1/4</sup> Fangs front, movable, hollow; pit between eye and nostril; more than 80 species.

## ANTICOAGULANTS

Clin. Lab. Invest. 7:204. [4] Bourgain, R., et al. 1954. Circulation 10:680. [5] Brown, A., and A. S. Douglas. and K. B. Olson. 1937. J. Biol. Chem. 122:153. [8] Clatanoff, D. V., and O. O. Meyer. 1956. Arch. Internal Med. 1955. Am. Surgeon 21:745. [11] Fantl, P., and M.H. Nance. 1947. Med. J. Australia 2:133. [12] Field, J. B., D. J., and O. O. Meyer. 1956. Ibid. 92:52. [15] Gordon, L. A., V. Richards, and H. A. Perkins. 1956. New Engl. Arch. Intern. Pharmacodyn. 87:402. [18] Heisey, S. R., J. P. Saunders, and K. C. Olson. 1956. J. Agr. Food Chem. J. Med. 243:395. [21] Jaques, L. B. 1955. Rev. Hematol. 10:379. [22] Jaques, L. B., A. F. Charles, and C. H. Allied Probl., Trans. Conf., 3rd, p. 11. [24] Jaques, L. B., and A. G. Ricker. 1948. Blood 3:1197. [25] Jaques, [27] Koller, T., and W. R. Merz, ed. 1955. Intern. Conf. Thrombosis Embolism, 1st, Basel, 1954. Proc. 24:811. [30] Solandt, D. Y., and C. H. Best. 1940. Lancet 1:1042. [31] Willis, P. W., et al. 1953. J. Lab. Clin.

## TOXINS

### REPTILES

reference 17. **Enzyme Activity** (column H): *A* = bradykininogen; *B* = carboxypolypeptidase; *C* = catalase; *D* = cephalinase; *K* = flavin adenine dinucleotide; *L* = hyaluronidase; *M* = invertase; *N* = L-amino-acid oxidase; *O* = lecithinase; phosphodiesterase; *V* = phosphomonoesterase; *W* = polypeptidase; *X* = protease. Slash mark (/) through letter indicates

Symptoms of Envenomation in Man	Mortality %	Available Antiserum	Reference	
(J)	(K)	(L)	(M)	
<i>Crotalidae</i> <sup>4</sup>				
Local pain, swelling and necrosis; lymphangitis and lymphadenitis; sweating, nausea, vomiting. Severe cases: shock, petechiae, bloody stools.	<1	Yes	H,40,43,74,114;I,68;J,12,15,84	1
Similar to poisoning by <i>A. contortrix</i> , but more severe; local necrosis more marked.	2-10	Yes	H,19,23,26,40,41,43,74,91,92,114;I,68;J,54,84,99	2
Local pain, edema and lymphadenopathy; bleeding from fang punctures, gums, nose and other body orifices; low prothrombin, prolonged clotting time; moderate to high leukocytosis; hematuria. Severe cases: shock, failure of pupils to react to light, respiration irregular. Autopsy: hemorrhagic necrosis at site of bite; hemorrhages into muscles, bowel, central nervous system; blood incoagulable.	10-20	Yes	H,7,23,43,101,114,116,118;I,86;J,49	3
Similar to poisoning by <i>B. atrox</i> . Autopsy: generalized visceral hemorrhages, cerebral hemorrhage, hemoglobinuric nephrosis.	5-15	Yes	H,7,23,27,42,43,67,100,109,114,116;I,87,88;J,68	4
Local pain, edema and ecchymoses; dryness of mouth, vomiting, shock, hemolytic anemia. Severe cases: muscular twitching, paresthesia, cyanosis, afibrogenemia, anemia, proteinuria, blood in feces, speech difficulty, sensation of yellow vision, unconsciousness.	5-15	Yes	H,21-23,40,41,43-45,93,114,117;I,55,68;J,2,55,61,108	5
Similar to poisoning by <i>C. adamanteus</i> , but neurotic symptoms less marked. Severe cases: profound shock.	5-15	Yes	H,23,24,64,109,114,116,117;I,55,68,84;J,25,55	6
Similar to poisoning by <i>Naja naja</i> , except that no edema occurs.	>40	Yes	H,17,23,27,36,63,101,114,116,117;I,68,86;J,16,68	7
Usual local symptoms of <i>Crotalus</i> poisoning; also thirst, abdominal pain, vomiting, diarrhea, dyspnea. Severe cases: excitement, hypertonicity of muscles, paresthesia, convulsions, cyanosis, respiratory failure, clouding of consciousness, weakness, and sweating.	1-2	Yes	H,114;I,68;J,8,20,25,55,81	8
Inadequate information. Rapid death preceded by severe, shocklike state.	Usually 100	Yes	H,7;I,84;J,68	9
Pain, edema, ecchymoses; weakness, sweating, vomiting. Severe cases: hemolytic anemia.	1-5	Yes	H,114,116;I,55,68,84;J,55,59	10

characteristic activity, rather than by specific isolation, of enzyme. /s/ Dry venom administered subcutaneously,

*continued*



Species (Common Name)	Distribution	Adult Length ft	Fibrinogen <sup>1</sup>		Pro- thrombin <sup>1</sup>		Enzyme Activity <sup>2</sup>	Mouse Toxicity <sup>3</sup> mg/kg
			Coag- ulate	De- stroy	Acti- vate	De- stroy		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Crotalidae <sup>4</sup>								
11	<i>Trimeresurus mucrosquamatus</i> (Taiwan habu)	Southeastern China, Formosa	3-4					MLD 7.6
Viperidae <sup>5</sup>								
12	<i>Bitis arietans</i> (puff adder)	Africa, southern Arabia	2½-5	-	+	-	-	A, E, N, R, U, V, X MLD 7.5
13	<i>Echis carinatus</i> (saw-scaled viper)	India, Iraq, Arabia, Africa (north of equator)	1½-2½	+				B, E, I, J, L, N, R, W, X LD <sub>50</sub> 3.3
14	<i>Vipera berus</i> (European viper)	British Isles, across northern Europe and Asia to Japan	1½-2½					A, N Mean LD 6.5
15	<i>V. russelli</i> (Russell's viper)	India, Burma, southern China, Formosa, Java	4-5½	+	-	-	+	B, E, H, I, J, K, L, M, N, O, P, Q, R, T, U, X, W, X MLD 1.0 LD <sub>50</sub> 5.0
Hydrophidae <sup>6</sup>								
16	<i>Enhydrina schistosa</i> (beaked sea snake)	Persian Gulf to S. China Sea, and south to Ceylon & northern Australia; mostly coastal waters	3-3½	-	-	-	-	O MLD 0.05-0.13
Elapidae <sup>7</sup>								
17	<i>Acanthophis antarcticus</i> (death adder)	Most of Australia (excluding Tasmania), New Guinea, nearby islands	1¾-3	+?		+?		E, L, N, O, R, T LD <sub>100</sub> 0.5
18	<i>Bungarus candidus caeruleus</i> (Indian krait)	India, Burma, Malay Peninsula, Java, Sumatra, Celebes	3½-4½					E, K, N, X LD <sub>50</sub> 1.0 LD <sub>100</sub> 3.0
19	<i>Demansia textilis</i> (brown snake)	Most of Australia (excluding Tasmania)	4½-7					E, N, R, T LD <sub>100</sub> 0.25
20	<i>Dendroaspis angusticeps</i> (eastern green mamba)	Eastern Africa, Ethiopia to Natal	6-13					E, N, R LD <sub>50</sub> 3.5
21	<i>Denisonia superba</i> (Australian copperhead)	Southeastern Australia	3½-5	+?		+?		E, L, N, O, R, T LD <sub>100</sub> 1.2
22	<i>Hemachatus haemachates</i> (ringhals)	S. Africa	3-4	+	-	±	-	E, N, O, Q, R, U, V, X Mean LD 1.5
23	<i>Micrurus corallinus</i> (coral snake)	Subtropical S. America	2-4					E, L, N, R
24	<i>M. fulvius</i> (eastern coral snake)	Southern United States to northeastern Mexico	2-3½					LD (MLD?) 1.0
25	<i>Naja naja</i> (Indian cobra)	Southern Asia to Indonesia, Formosa, Philippines	4-6	-	-	-	+	B, E, H, I, J, L, M, N, O, P, R, U, V, W, X MLD 0.75 LD <sub>50</sub> 0.20

<sup>1/1</sup> Information from reference 23, unless otherwise specified in column M. <sup>2/2</sup> Presence in venom determined by unless otherwise stated. <sup>4/4</sup> Fangs front, movable, hollow; pit between eye and nostril; more than 80 species. <sup>6/6</sup> Fangs front, grooved though virtually fused for most of length; more than

## TOXINS

### REPTILES

Symptoms of Envenomation in Man	Mor- tality %	Avail- able Anti- serum	Reference	
(J)	(K)	(L)	(M)	
<b>Crotalidae<sup>4</sup></b>				
Local pain, ecchymoses, blistering; little systemic reaction.	2-10	Yes	I,57;J,95,111	11
<b>Viperidae<sup>5</sup></b>				
Severe local edema, necrosis and sloughing; restlessness, weak pulse, dyspnea, gastrointestinal hemorrhages.	11-40	Yes	H,23,40,109,114,116; I,38;J,39,90	12
Local pain and edema; ecchymoses and hemorrhages from mucous membranes and into viscera; profound anemia, abdominal pain, impaired liver function; prothrombin time greatly prolonged, thrombocytopenia. Autopsy: intestinal and retroperitoneal hemorrhage.	11-40	Yes	D,58;H,27,33,34,36,56,114,115;L,68;J,69,82;L,39	13
Local pain and edema of bitten extremity, sometimes extending into trunk; hemorrhages along lymphatics. Little systemic reaction; sometimes vomiting, sweating, abdominal pain, faintness, cyanosis, shock. Ptosis common after bite by <i>V. berus bosniensis</i> .	1-5	Yes	H,109,119;I,85;J,75,83,98,102,106	14
Rapidly spreading edema with extravasation of blood, epistaxis and petechiae, abdominal pain, vomiting, paralytic ileus, collapse, shock, albuminuria, prolonged clotting time. Terminally: loss of consciousness, failure of pupils to react to light, circulatory failure. Autopsy: subcutaneous hemorrhages near site of bite, meningeal congestion, blood in lungs.	11-40	Yes	H,19,23,33-36,40,41,44,45,60,74,79,80,91,114,116;I,57,77;J,13,112	15
<b>Hydrophidae<sup>6</sup></b>				
No local reaction; latent period minutes to few hours. Giddiness, muscular aching followed by muscle weakness, ptosis, trismus, hypertension. Death from respiratory failure, cardiac arrest, or acute renal failure. Autopsy: marked, widespread myonecrosis; renal congestion with distal tubular necrosis.	12-25	Yes	D-H,10;I,9,68;J,K,66,78	16
<b>Elapidae<sup>7</sup></b>				
Similar to poisoning by <i>Notechis scutatus</i> except that peripheral circulatory failure is more common and hemorrhagic phenomena occur.	11-40	Yes	D,F,104;H,19,28,111,114,116;I,71;J,53	17
Little pain or local reaction. Latent period may extend to 12 hours, followed by abdominal pain, staggering gait, dysphagia, dyspnea, ptosis, stiffness of jaws, coma, respiratory paralysis, cardiac failure.	77	Yes	H,48,74,91,114;I,48,77;J,1,31	18
Latent period to 12 hours, followed by abdominal pain, vomiting, headache, dizziness, weakness, rapid pulse and subnormal temperature, respiratory and circulatory collapse, hemoglobinuria, and peripheral thromboses.	11-40	Yes	H,19,114,116;I,52;J,53	19
Local pain and swelling, salivation, paralysis of vocal cords, sweating, vomiting, restlessness, drowsiness or collapse followed by coma; dyspnea and respiratory failure.	>40	Yes	H,114,116;I,14;J,76	20
Similar to poisoning by <i>Notechis scutatus</i> . Rapid loss of muscle tone and consciousness, peripheral circulatory failure.	11-40	Yes	D,F,104;H,11,19,28,103,114,116;I,50;J,53	21
Pain; dyspnea; weak, thready pulse; cyanosis; collapse. Venom sprayed at eyes; effects resemble those produced by <i>Naja nigricollis</i> .	2-10	Yes	H,5,23,40,114,116;I,85;J,29,90	22
Numbness without pain at bite. Early symptoms: headache, swelling of face and lips; hyperesthesia, sore throat, ptosis, photophobia, normal pupillary reflex, vomiting, cramps, dyspnea, loss of muscle tone, tachycardia. Later symptoms: backache, irritability, salivation, bradycardia, dysuria, albuminuria.	11-40	Yes	H,114-117;J,113	23
Cyclic pains radiating from site of bite; somnolence, dyspnea, dysphagia, sweating; soreness of face, throat, and eyes.	5-20	No	I,62;J,110	24
Pain radiating from site of bite; edema, numbness, tremors, ptosis, drooping of head, salivation, speech difficulty, giddiness, muscular incoordination and weakness, blindness, progressive depression of respiration, convulsions, incontinence of urine and feces. Pupils react to light, and heart continues to beat, after respiration has ceased.	11-40	Yes	H,3,6,23,30,32-34,36,40,46,65,72-74,79,80,94,114,117;I,77;J,1,13,31	25

characteristic activity, rather than by specific isolation, of enzyme. /s/ Dry venom administered subcutaneously, /s/ Fangs front, movable, hollow; approximately 50 species. /s/ Fangs short, front, permanently erect; approximately 150 species.

*continued*

Species (Common Name)	Distribution	Adult Length ft	Fibrinogen <sup>1</sup>		Pro- thrombin <sup>1</sup>		Enzyme Activity <sup>2</sup>	Mouse Toxicity <sup>3</sup> mg/kg	
			Coag- ulate	De- stroy	Acti- vate	De- stroy			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	
<b>Elapidae<sup>7</sup></b>									
26	<i>Naja nigricollis</i> (black-necked cobra)	Africa (south of Sahara in savanna area)	5-6				+	E,N,R,T, X	MLD 2.5
27	<i>Notechis scutatus</i> (tiger snake)	Most of Australia except northern part	3½-6	-	-	+	-	E,L,N,R, S,T,U, V,X	LD <sub>50</sub> (guinea pig) 6.5 µg/kg LD <sub>100</sub> 0.3
28	<i>Ophiophagus hannah</i> (king cobra)	Western India, Burma, Philippines, Indonesia, southern China, Thailand	12-16					E,N,R	
29	<i>Oxyuranus scutellatus</i> (taipan)	Northern Australia, New Guinea	6-11	+?		+?		O,T	LD <sub>100</sub> 0.17
30	<i>Pseudochis porphyriacus</i> (Australian black snake)	Eastern & southern Australia	4-6½	+?		+?		N,O,T	LD <sub>100</sub> 3.5
<b>Colubridae<sup>8</sup></b>									
31	<i>Dispholidus typus</i> (boom-slang)	Africa, forested portions (south of Sahara)	4½-5½				+?	X	MLD 10.0
<b>Helodermatidae</b>									
32	<i>Heloderma suspectum</i> (Gila monster)	Southwestern United States (chiefly Arizona), northwestern Mexico (chiefly Sonora)	1½-2						LD <sub>50</sub> (rat), 20.18 (lyophilized venom)

/1/ Information from reference 23, unless otherwise indicated in column M. /2/ Presence in venom determined by unless otherwise stated. /7/ Fangs front, grooved though virtually fused for most of length; more than 150 species. /8/ Antiserum for crotalid envenomation believed to be effective.

**Contributors:** (a) Minton, Sherman A., Jr., (b) Schöttler, Werner H. A., (c) Slotta, Karl H., (d) Graydon, John J.,

**References:** [1] Ahuja, M. L., and G. Singh. 1954. Indian J. Med. Res. 42:661. [2] Andrews, E. H., and C. B. K. M. 1951. Med. J. Australia 38(1):147. [5] Björk, W. 1961. Biochim. Biophys. Acta 49:195. [6] Bovet, F., and 12:442. [8] Bulger, J. J., and A. K. Northrop. 1951. J. Am. Med. Assoc. 147:1134. [9] Carey, J. E., and E. A. [11] Chain, E., and E. S. Duthie. 1940. Brit. J. Exptl. Pathol. 21:324. [12] Chotkowski, L. A. 1949. New Engl. lished. South African Institute for Medical Research, Johannesburg, 1963. [15] Corkill, N. L. 1932. Indian J. St. Louis. p. 1252. [17] do Amaral, A. 1960. Haffkine Inst. (Bombay) Symp., 1959, Proc., p. 128. [18] Doery, 1928. Calif. Western Med. 29:237. [21] Dunn, E. E. 1934. J. Pharmacol. Exptl. Therap. 50:386. [22] Dunn, E. E. 1941. Biochem. J. 35:872. [25] Ehrlich, P. 1928. Bull. Antivenin Inst. Am. 2:65. [26] Fairbairn, D. 1945. J. 1938. J. Physiol. (London) 94:232. [29] Fitzsimons, F. W. 1921. Snakes of South Africa. T. M. Miller, Capetown. Med. Gaz. 67:81. [32] Ghosh, B. N., 1936. J. Indian Chem. Soc. 13:450. [33] Ghosh, B. N. 1940. Oesterr. B. N., and S. S. De. 1936. Ibid. 13:627. [36] Ghosh, B. N., P. K. Dutt, and D. K. Chowdhury. 1939. Ibid. 16:75. A. W. Schaafsma. 1935. Trans. Roy. Soc. Trop. Med. Hyg. 28:601. [39] Gray, H. H. 1962. Ibid. 56:390. Ibid. 32:597. [42] Henriques, O. B., et al. 1960. Ciencia Cult. (Sao Paulo) 12(3-4):175. [43] Houssay, B. A., and 1951. J. Biol. Chem. 193:91. [45] Hurst, R. O., J. A. Little, and G. C. Butler. 1951. Ibid. 188:705. [46] Iyengar, Inst. Pasteur 84:959. [48] Jaques, R. Unpublished. Ciba, Basel, Switzerland, 1955. [49] Jutzy, D., et al. 1953. Ibid. 17(2):33. [52] Kellaway, C. H. 1931. Ibid. 18(2):747. [53] Kellaway, C. H. 1942. Ibid. 29(2):171. [54] Kelly, [56] Kundu, M. L., S. S. De, and B. N. Ghosh. [57] Kuwajima, Y. 1953. Japan. J. Exptl. Med. 23:457. [58] Lefrou, Indiana Acad. Sci. 45:253. [60] McClean, D., and C. W. Hale. 1941. Biochem. J. 35:159. [61] McCreary, T., and

## TOXINS

### REPTILES

Symptoms of Envenomation in Man	Mortality %	Available Antiserum	Reference	
(J)	(K)	(L)	(M)	
Elapidae <sup>7</sup>				
Similar to poisoning by <i>N. naja</i> . Venom frequently sprayed at eyes; contact produces acute, intense ophthalmia. Systemic poisoning does not occur from such contact, and permanent damage to vision is rare.	11-40	Yes	G,47;H,19,74,114,116;I,38;J,68	26
Latent period 15-60 minutes, followed by nausea, vomiting, faintness, drowsiness, sweating. Later symptoms: dullness of sensation, staggering, dysphagia, slurred speech, ptosis, dilation of pupils and failure to react to light; rapid, weak pulse and respiration; progressive dyspnea and death from respiratory failure.	>40	Yes	H,11,18,19,23,40,114,116;I,71,105;J,53	27
Similar to poisoning by <i>Naja</i> species. Symptoms develop rapidly; death often occurs in 30-60 minutes.	>40	Yes	H,114,116;J,107	28
Similar to poisoning by <i>Notechis scutatus</i> . Flaccid paralysis of limbs, intercostal and bulbar paralysis; often rapidly fatal.	>40	Yes	D,F,104;H,19,28,103;I,70;J,4	29
Local pain and swelling, vomiting, hemorrhages from nose and mouth, prostration, hematuria.	<1	Yes	D,F,104;H,19,28,103,114;I,51;J,53	30
Colubridae <sup>8</sup>				
Local pain, swelling and hemorrhage; ecchymoses, defibrination syndrome, bleeding from nose and mouth, sometimes from all mucous membranes and skin; headache, vomiting, collapse; temperature normal or subnormal.	High	Yes	F,H,I,37;J,37,96	31
Helodermatidae				
Local pain, swelling, hyperemia, weakness, hyperpnea, tinnitus, nausea, vomiting. Death from respiratory paralysis and cardiac failure.	<1->40	No <sup>8</sup>	I,97;J,89	32

characteristic activity, rather than by specific isolation, of enzyme. /s/ Dry venom administered subcutaneously, /e/ Numerous species, mostly harmless; fangs rear, immovable, grooved in venomous species; dangerous if handled.

and Morgan, F. G., (e) Christensen, P. Agerholm, (f) do Amaral, Afranio

Pollard. 1953. J. Florida Med. Assoc. 40:388. [3] Augustinsson, K.-B. 1949. Arch. Biochem. 23:111. [4] Benn, D. Bovet. 1943. Ann. Inst. Pasteur 69:309. [7] Brazil, V., and N. R. Pestana. 1909. Rev. Med. Cir. Sao Paulo Wright. 1960. Trans. Roy. Soc. Trop. Med. Hyg. 54:50. [10] Carey, J. E., and E. A. Wright. 1961. Ibid. 55:153. J. Med. 241:600. [13] Chowhan, J. S. 1938. Antiseptic (Madras, India) 35:544. [14] Christensen, P. A. Unpub. Med. Res. 20:599. [16] do Amaral, A. 1951. In R. B. H. Gradwohl, ed. Clinical tropical medicine. C. V. Mosby, H. M. 1958. Biochem. J. 70:535. [19] Doery, H. M., and J. E. Pearson. 1961. Ibid. 78:820. [20] Doughty, J. F. 1934. Ibid. 50:393. [23] Eagle, H. 1937. J. Exptl. Med. 65:613. [24] East, M. E., J. Madinaveitia, and A. R. Todd. Biol. Chem. 157:633. [27] Favilli, G. 1940. Nature 145:866. [28] Feldberg, W., H. F. Holden, and C. H. Kellaway. [30] Ganguly, S. N., and M. T. Malkana. 1936. Indian J. Med. Res. 24:281. [31] Gharpurey, K. G. 1932. Indian Chemiker-Ztg. 43:158. [34] Ghosh, B. N., and D. K. Chowdhury. 1938. J. Indian Chem. Soc. 15:566. [35] Ghosh, [37] Grasset, E., and A. W. Schaafsma. 1940. S. African Med. J. 14:236. [38] Grasset, E., A. Zoutendyk, and [40] Gulland, J. M., and E. M. Jackson. 1938. Biochem. J. 32:590. [41] Gulland, J. M., and E. M. Jackson. 1938. J. Negrete. 1918. Rev. Inst. Bacteriol. Dept. Nacl. Hig. (Buenos Aires) 1:341. [44] Hurst, R. O., and G. C. Butler. N. K., K. B. Sehra, and B. Mukerji. 1938. Indian J. Med. Res. 26:487. [47] Iizard, Y., and P. Boquet. 1953. Ann. Am. J. Trop. Med. Hyg. 2:129. [50] Kellaway, C. H. 1929. Med. J. Australia 16(1):358. [51] Kellaway, C. H. 1930. H. A. 1922. Therap. Gaz. 38:846. [55] Klauber, L. M. 1956. Rattlesnakes. Univ. California Press, Berkeley, v.2. G., and J. Martignoles. 1954. Ann. Inst. Pasteur 86:446. [59] Lyon, M. W., Jr., and C. A. Bishop. 1936. Proc. H. Wurzel. 1959. J. Am. Med. Assoc. 170:268. [62] Macht, D. I. 1947. Copeia (4):269. [63] Madinaveitia, J. 1939.

continued



Biochem. J. 33:1470. [64] Madinaveitia, J. 1941. Ibid. 35:447. [65] Manwaring, W. H. 1910. Z. Immunitaets-Azevedo. 1949. Mem. Inst. Butantan 22:47. [68] Minton, S. A., Jr. Unpublished. Indiana Univ. Medical Center, 1954. Papers Intern. Conf. Animal Venoms, Meeting Am. Assoc. Advan. Sci., 12th, Berkeley, p. 359. [71] Morgan, Melbourne, 1955. [72] Mounter, L. A. 1951. Biochem. J. 49:xlv. [73] Mounter, L. A. 1951. Ibid. 50:122. [76] Pitman, C. R. S. 1938. A guide to the snakes of Uganda. Uganda Society, Kampala. [77] Rao, S. S., and A. C. 1938. Indian J. Med. Res. 26:249. [80] Roy, A. C., and R. N. Chopra. 1938. Ibid. 26:241. [81] Russell, Z. Hyg. Infektionskrankh. 124:141. [84] Schöttler, W. H. A. 1951. Am. J. Trop. Med. 31:489. [85] Schöttler, W. H. A. 1955. Ibid. 12:877. [88] Schöttler, W. H. A. 1958. Ibid. 19:341. [89] Shannon, F. 1953. Herpetologica Biochem. 27:348. [92] Singer, T. P., and E. B. Kearney. 1950. Ibid. 29:190. [93] Sinsheimer, R. L., and J. F. [95] Sonneborn, D. G. 1946. U.S. Naval Med. Bull. 46:105. [96] Spies, S. K., L. F. Malherbe, and W. J. Pepler. [98] Stanley-Jones, D., and C. E. S. Harris. 1942. Brit. Med. J. 2:395. [99] Swarzwelder, J. 1950. Am. J. Trop. et al. 1952. J. Biol. Chem. 195:207. [102] Tallqvist, H., and K. Österlund. 1962. Nord. Med. 68:1073. Univ. Melbourne, Australia, 1956. [105] Trethewie, E. R., and A. J. Day. 1948. Australian J. Exptl. Biol. Med. India. Bombay Natural History Society, Bombay. [108] Watt, H. F., and C. B. Pollard. 1954. J. Florida Med. D. M. Darling. 1950. Herpetologica 6:197. [111] Wu, Y. K., and Y. H. Tsui. 1945. Chinese Med. J. 63A:148. 1948. Advan. Enzymol. 8:459. [115] Zeller, E. A. 1948. Ibid. 8:475. [116] Zeller, E. A. 1950. Helv. Chim. [118] Zeller, E. A., and A. Maritz. 1945. Helv. Chim. Acta 28:365. [119] Zeller, E. A., A. Maritz, and B. Iselin.

Part II.

Toad	Distribution	Bufagins <sup>1</sup>			Bufotoxins <sup>2</sup>
		Name (Proposed Formula)	Action or Effect <sup>4</sup>	Toxicity <sup>5</sup> mg/kg	Name (Proposed Formula)
(A)	(B)	(C)	(D)	(E)	(F)
1 <i>Bufo alvarius</i> (Colorado River toad)	Southwestern United States				Alvarobufotoxin
2 <i>B. americanus</i> (American toad)	Eastern United States	Americobufagin	Digitalis-like action		Americobufotoxin
3 <i>B. arenarum</i> (sand toad)	Argentina	Arenobufagin (C <sub>25</sub> H <sub>34</sub> O <sub>6</sub> )	Digitalis-like action; emesis; systolic standstill	0.092 ±0.005	Arenobufotoxin (C <sub>39</sub> H <sub>60</sub> O <sub>11</sub> N <sub>4</sub> )
4 <i>B. bufo</i> (European toad)	Europe				Vulgarobufotoxin (C <sub>38</sub> H <sub>60</sub> O <sub>11</sub> N <sub>4</sub> )
5 <i>B. formosus</i> (Japanese toad)	Japan	Gamabufagin (C <sub>27</sub> H <sub>38</sub> O <sub>6</sub> )	Digitalis-like action; emesis; ventricular fibrillation	0.101 ±0.005	Gamabufotoxin (C <sub>41</sub> H <sub>60</sub> O <sub>11</sub> N <sub>4</sub> )
6 <i>B. woodhousii fowleri</i> (Fowler's toad)	Southeastern United States	Fowlerobufagin (C <sub>23</sub> H <sub>33</sub> O <sub>6</sub> )	Digitalis-like action; emesis; ventricular fibrillation	0.218 ±0.012	Fowlerobufotoxin
7 <i>B. gargarizans</i> (Cantor's toad)	China	Cinobufagin (C <sub>25</sub> H <sub>31</sub> O <sub>5</sub> ) <sup>9</sup>	Digitalis-like action on vagus, vagus center, myocardium; emesis; clonic or tonic convulsions after paralysis	0.219 ±0.011	Cinobufotoxin (C <sub>43</sub> H <sub>64</sub> O <sub>12</sub> N <sub>4</sub> ) or (C <sub>39</sub> H <sub>58</sub> O <sub>11</sub> N <sub>4</sub> )
		Telecinobufagin (C <sub>24</sub> H <sub>34</sub> O <sub>5</sub> )		0.102 ±0.007	
8 <i>B. marinus</i> (marine toad)	Circumtropical	Marinobufagin <sup>10</sup> (C <sub>24</sub> H <sub>32</sub> O <sub>5</sub> )	Digitalis-like action; emesis; ventricular fibrillation	0.555 ±0.028	Marinobufotoxin (C <sub>38</sub> H <sub>58</sub> O <sub>10</sub> N <sub>4</sub> ) or (C <sub>42</sub> H <sub>62</sub> O <sub>11</sub> N <sub>4</sub> ) <sup>11</sup>
9 <i>B. quercicus</i> (oak toad)	Southeastern United States	Quercicobufagin (C <sub>23</sub> H <sub>34</sub> O <sub>5</sub> )	Digitalis-like action; emesis; ventricular fibrillation	0.097 ±0.004	Quercicobufotoxin

<sup>1/2</sup> Bufagins are steroid-type compounds [1, 8, 20, 32]. <sup>2/2</sup> Bufotoxins are the conjugation product of the specific molecule [13]. <sup>4/4</sup> On cat, guinea pig, rabbit, pigeon, frog. <sup>5/5</sup> Average fatal dose for cat (intravenous) [2]. <sup>6/6</sup> On [12]. <sup>10/10</sup> Also reported as occurring in *B. Paracnemis* (Argentina) [18, 34, 35]. <sup>11/11</sup> Also reported as C<sub>42</sub>H<sub>64</sub>O<sub>11</sub>N<sub>4</sub>

**TOXINS**

**REPTILES**

forsch. 6:513. [66] Marsden, A. T. H., and H. A. Reid. 1961. Brit. Med. J. 1:1290. [67] Martirani, L., and M. P. Indianapolis. 1963. [69] Mole, R. H., and A. Everard. 1947. Quart. J. Med., N.S.16:291. [70] Morgan, F. G. F. G., and J. J. Graydon. Unpublished. Commonwealth of Australia, Dept. of Health, Serum Laboratories, [74] Noc, F. 1904. Ann. Inst. Pasteur 18:387. [75] Petitpierre, M. 1934. Schweiz. Med. Wochschr. 64:372. M. E. Kulkarni. 1952. Haffkine Inst. (Bombay) Rept., p. 49. [78] Reid, H. A. 1961. Brit. Med. J. 1:1284. [79] Roy, F. E. 1960. Am. J. Med. Sci. 239:1. [82] Salou, G. 1951. Med. Trop. 11:655. [83] Schöttler, W. H. A. 1942. W. H. A. 1951. Ibid. 31:836. [86] Schöttler, W. H. A. 1952. Bull. World Health Organ. 5:293. [87] Schöttler, 9:125. [90] Shircore, J. O. 1947. E. African Med. J. 24:200. [91] Singer, T. P., and E. B. Kearney. 1950. Arch. Koerner. 1952. J. Biol. Chem. 198:293. [94] Slotta, K. 1955. Fortschr. Chem. Org. Naturstoffe 12:406. 1962. S. African Med. J. 36:834. [97] Stahnke, H. L. Unpublished. Arizona State College, Tempe, 1956. Med. 30:575. [100] Taborda, A. R., and L. C. Taborda. 1940. Mem. Inst. Butantan 14:183. [101] Taborda, A. R., [103] Trethewie, E. R. 1939. Australian J. Exptl. Biol. Med. Sci. 17:145. [104] Trethewie, E. R. Unpublished. Sci. 26:37. [106] Walker, C. W. 1945. Brit. Med. J. 2:13. [107] Wall, F. 1928. Poisonous terrestrial snakes of Assoc. 41:367. [109] Werle, E., R. Kehl, and K. Koebke. 1950. Biochem. Z. 320:372. [110] Werler, J. E., and [112] Wynon, P. H. 1945. Brit. Med. J. 2:919. [113] Yered, D. 1942. Gaz. Clin. 40:261. [114] Zeller, E. A. Acta 33:821. [117] Zeller, E. A., B. Iselin, and A. Maritz. 1946. Helv. Physiol. Pharmacol. Acta 4:233. 1945. Ibid. 28:1615.

**TOADS**

Bufotoxins <sup>2</sup>		Bufotenines <sup>3</sup>			Other Compounds Isolated	Reference
Action or Effect <sup>6</sup>	Toxicity <sup>5</sup> mg/kg	Name (Proposed Formula)	Action or Effect <sup>7</sup>	Cardiac Arrest <sup>8</sup>		
(G)	(H)	(I)	(J)	(K)	(L)	(M)
Digitalis-like action; emesis; systolic standstill	0.756 ±0.075	Alvarobufotenine (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; slight pressor action; diastolic standstill	1:5,000 dilution	Cholesterol, ergosterol	6,8,13
Digitalis-like action		Americobufotenine (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; marked pressor action			6,9,13
Digitalis-like action; emesis; ventricular fibrillation	0.406 ±0.012	Arenobufotenine A (C <sub>12</sub> H <sub>20</sub> O <sub>3</sub> N <sub>2</sub> ) Arenobufotenine B (C <sub>14</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; slight pressor action	1:5,000 dilution	Cholesterol, epinephrine	8,13,15
Emesis; ventricular fibrillation	0.292 ±0.017	Vulgarobufotenine (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; marked pressor action		Cholesterol, ergosterol	7,8,13, 14
Persistent action; slight pressor action; emesis; ventricular fibrillation	0.374 ±0.027	Gamabufotenine (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; marked pressor action	1:5,000 dilution	Cholesterol, epinephrine, bufotenidine	8,13,16, 17,22, 33
Emesis	0.792 ±0.054	Fowlerobufotenine (C <sub>13</sub> H <sub>20</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; marked pressor action; epinephrine-like action	1:5,000 dilution	Bufotenedine	6,8,13, 22,33
Emesis; vasopressor effect, followed by cardiac collapse, death in systole; prolongation of P-R interval	0.359 ±0.024	Cinobufotenine (C <sub>12</sub> H <sub>16</sub> ON <sub>2</sub> )	Oxytotic; miotic; intense, short vasopressor action; contraction of smooth muscle not inhibited by atropine	1:10,000 dilution	Cholesterol, epinephrine, norepinephrine, bufotenedine	3,6,7, 11-13, 22-31
More emetic than bufagin	0.417 <sup>12</sup> ±0.022	Marinobufotenine (C <sub>12</sub> H <sub>14</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; no pressor action	1:5,000 dilution	Epinephrine, cholesterol, ergosterol, 5-hydroxytryptamine	1,5-7, 10,13, 19
Ventricular systolic standstill		Quercicobufotenine (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytotic; slight pressor action		Cholesterol?	6,13

bufagin with one molecule of suberyl-arginine [8]. /s/ Bufotenines are organic bases having an indole ring in the cat, pigeon, frog. /7/ On cat, guinea pig, frog. /s/ Frog, heart perfusion method. /s/ Also reported as C<sub>29</sub>H<sub>38</sub>O<sub>7</sub> [21]. /12/ Also reported as 0.43 [5] and 0.49 [10].

*continued*

Toad	Distribution	Bufagins <sup>1</sup>			Bufotoxins <sup>2</sup>
		Name (Proposed Formula)	Action or Effect <sup>4</sup>	Toxicity <sup>5</sup> mg/kg	Name (Proposed Formula)
(A)	(B)	(C)	(D)	(E)	(F)
10 <i>Bufo regularis</i> (leopard toad)	South Africa	Regularobufagin (C <sub>23</sub> H <sub>34</sub> O <sub>5</sub> )	Digitalis-like action; emesis; ventricular fibrillation	0.153 ±0.006	Regularobufotoxin (C <sub>37</sub> H <sub>60</sub> O <sub>10</sub> N <sub>4</sub> )
11 <i>B. valliceps</i> (Mexican toad)	Eastern Mexico, Texas & Louisiana	Vallicepobufagin (C <sub>23</sub> H <sub>34</sub> O <sub>5</sub> )	Nausea; emesis; A-V block, and ventricular standstill	0.201 ±0.017	Vallicepobufotoxin
12 <i>B. viridis</i> (green toad)	Europe	Viridobufagin (C <sub>23</sub> H <sub>34</sub> O <sub>5</sub> )	Nausea; emesis; increased intestinal tone; ventricular fibrillation	0.111 ±0.008	Viridobufotoxin (C <sub>37</sub> H <sub>60</sub> O <sub>10</sub> N <sub>4</sub> )

/1/ Bufagins are steroid-type compounds [1, 8, 20, 32]. /2/ Bufotoxins are the conjugation product of the specific molecule [13]. /4/ On cat, guinea pig, rabbit, pigeon, frog. /5/ Average fatal dose for cat (intravenous) [2]. /6/ On 10.7 mg per animal.

Contributors: (a) Chen, K. K., and Herrmann, Roy G., (b) Shannon, F. A.

References: [1] Abel, J. J., and D. I. Macht. 1912. J. Pharmacol. Exptl. Therap. 3:319. [2] Chen, K. K. 1945. Med. 76:372. [4] Chen, K. K., and A. L. Chen. 1933. J. Pharmacol. Exptl. Therap. 49:503. [5] Chen, K. K., and Chen. 1933. Ibid. 49:548. [8] Chen, K. K., and A. L. Chen. 1933. Ibid. 49:561. [9] Chen, K. K., and A. L. Chen. 97:511. [11] Chen, K. K., H. Jensen, and A. L. Chen. 1931. Ibid. 97:512. [12] Chen, K. K., H. Jensen, and A. L. Exptl. Biol. Med. 29:905. [14] Chen, K. K., H. Jensen, and A. L. Chen. 1933. J. Pharmacol. Exptl. Therap. 47:307. Ibid. 49:14. [17] Chen, K. K., H. Jensen, and A. L. Chen. 1933. Ibid. 49:26. [18] Deulofeu, V., and J. R. Mendive. 57:2733. [21] Jensen, H., and K. K. Chen. 1930. J. Biol. Chem. 87:31. [22] Jensen, H., and K. K. Chen. 1936. H. M., and K. K. Chen. 1951. J. Pharmacol. Exptl. Therap. 102:286. [25] Meyer, K. 1948. Experientia 4:385. Ibid. 32:1599. [29] Meyer, K. 1949. Ibid. 32:1993. [30] Meyer, K. 1949. Pharm. Acta Helv. 24:222. [31] Shimizu, H., W. Konz, and H. Mittasch. 1934. Ann. Chem. 513:1. [34] Xavier, A. A., J. Vellard, and M. Miguelote-Vianna. 108:1085.

Part III. MARINE ORGANISMS

Chordata	
1	<p>CATFISH STING, caused by contact with spine located in front of the soft-rayed portion of the dorsal and pectoral fins.</p> <p><b>Species &amp; Distribution:</b> <i>Bagre marina</i> (sea catfish); Cape Cod to Brazil  <i>Clarias batrachus</i> (catfish); India to Indonesia, Philippines  <i>Galeichthys felis</i> (sea catfish); Cape Cod to Gulf of Mexico  <i>Heteropneustes fossilis</i> (catfish); India, Ceylon, Vietnam  <i>Plotosus lineatus</i> (sea catfish); vicinity of river mouths in Indo-Pacific area</p> <p><b>Symptoms:</b> Instant stinging pain, usually localized. Primary shock. Pallor about wound, occasionally some edema. Rarely serious. [12, 20, 57, 64, 106-108]</p>
2	<p>CIGUATERA POISONING, caused by ingestion of any one of a number of species of fishes, the flesh of which becomes toxic under certain conditions.</p> <p><b>Species &amp; Distribution:</b> <i>Acanthurus glaucopareius</i> (surgeonfish); tropical Pacific, Indonesia, Philippines  <i>A. triostegus</i> (Indo-Pacific convict fish); Hawaiian &amp; Johnson Islands  <i>Albula vulpes</i> (ladyfish); warm seas  <i>Alutera scripta</i> (longtail filefish); warm seas  <i>Aprion virescens</i> (blue-gray snapper); tropical Indo-Pacific  <i>Balistoides niger</i> (triggerfish); tropical Pacific, China, Japan  <i>Caranx hippos</i> (jack); tropical Atlantic  <i>Cephalopholis argus</i> (blue-spotted argus); tropical Indo-Pacific  <i>Clupanodon thrissa</i> (gizzard shad); tropical Pacific, Japan, China, Formosa, Korea, Indonesia, India  <i>Engraulis japonica</i> (anchovy); China, Japan, Korea, Formosa  <i>Epibolus insidiator</i> (Indo-Pacific long-jawed wrasse); tropical Indo-Pacific  <i>Epinephelus fuscoguttatus</i> (mottled grouper); Indo-Pacific</p>

continued



**TOXINS**

**TOADS**

Bufotoxins <sup>2</sup>		Bufotenines <sup>3</sup>			Other Compounds Isolated	Reference
Action or Effect <sup>4</sup>	Toxicity <sup>5</sup> mg/kg	Name (Proposed Formula)	Action or Effect <sup>7</sup>	Cardiac Arrest <sup>8</sup>		
(G)	(H)	(I)	(J)	(K)	(L)	(M)
Emesis; ventricular fibrillation	0.477 ±0.026	Regularobufotenine	Oxytocic; marked pressor action		Epinephrine <sup>13</sup>	4,8,13
		Vallicepobufotenine (C <sub>11</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytocic; slight pressor action; decreased amplitude and arrest of heart contractions	1:1,000 dilution		6,13
Action similar to, but weaker than, that of viridobufagin	0.270 ±0.012	Viridobufotenine A (C <sub>12</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> )	Oxytocic; marked pressor action	1:5,000 dilution	Cholesterol, ergosterol	8,13,16
		Viridobufotenine B (C <sub>12</sub> H <sub>20</sub> O <sub>3</sub> N <sub>2</sub> )	Oxytocic; slight pressor action			

bufagin with one molecule of suberyl-arginine [8]. <sup>3</sup>/ Bufotenines are organic bases having an indole ring in the cat, pigeon, frog. <sup>7</sup>/ On cat, guinea pig, frog. <sup>8</sup>/ Frog, heart perfusion method. <sup>13</sup>/ 4.3-5.0% of whole venom,

Ann. Rev. Physiol. 7:682. [3] Chen, K. K., R. C. Anderson, and F. G. Henderson. 1951. Proc. Soc. Exptl. Biol. A. L. Chen. 1933. Ibid. 49:514. [6] Chen, K. K., and A. L. Chen. 1933. Ibid. 49:526. [7] Chen, K. K., and A. L. Chen. 1934. Arch. Intern. Pharmacodyn. 47:297. [10] Chen, K. K., H. Jensen, and A. L. Chen. 1931. Am. J. Physiol. Chen. 1931. J. Pharmacol. Exptl. Therap. 43:13. [13] Chen, K. K., H. Jensen, and A. L. Chen. 1932. Proc. Soc. [15] Chen, K. K., H. Jensen, and A. L. Chen. 1933. Ibid. 49:1. [16] Chen, K. K., H. Jensen, and A. L. Chen. 1933. 1938. Ann. Chem. 534:288. [19] Jensen, H. 1932. Science 75:53. [20] Jensen, H. 1935. J. Am. Chem. Soc. Ibid. 116:87. [23] Kotake, M., and K. Kuwada. 1939. Sci. Papers Inst. Phys. Chem. Res. (Tokyo) 35:419. [24] Lee, [26] Meyer, K. 1949. Helv. Chim. Acta 32:1238. [27] Meyer, K. 1949. Ibid. 32:1593. [28] Meyer, K. 1949. S. 1916. J. Pharmacol. Exptl. Therap. 8:347. [32] Shoppee, C. W. 1942. Ann. Rev. Biochem. 11:137. [33] Wieland, 1931. Compt. Rend. Soc. Biol. 108:1082. [35] Xavier, A. A., J. Vellard, and M. Miguelote-Vianna. 1931. Ibid.

**Part III. MARINE ORGANISMS**

Chordata
<i>Gnathodentex aureolineatus</i> (snapper); Tuamotu Archipelago west to eastern Africa
<i>Katsuwonus pelamis</i> (skipjack); circumtropical
<i>Lactophrys trigonus</i> (trunkfish); Atlantic coast of tropical America north to Cape Cod
<i>Lactoria cornutus</i> (trunkfish); tropical Pacific
<i>Lethrinus miniatus</i> (snapper-like fish); Polynesia west to eastern Africa
<i>Lutjanus bohar</i> (twinspot snapper); tropical Pacific to eastern Africa, Red Sea
<i>L. gibbus</i> (red snapper); tropical Indo-Pacific
<i>L. monostigma</i> (one-spot snapper); Polynesia west to Red Sea, China
<i>L. vaigiensis</i> (red snapper); Polynesia west to eastern Africa, Japan
<i>Mycteroperca venenosa</i> (sea bass); western tropical Atlantic
<i>Pagellus erythrinus</i> (porgy); Mediterranean & Black Seas, eastern Atlantic from British Isles & Scandinavia to Azores, Canary Islands, Fernando Po
<i>Pagrus pagrus</i> (porgy); E. Atlantic, Mediterranean
<i>Paradicichthys venenatus</i> (Chinaman fish); Australia
<i>Parupeneus chryseerydros</i> (surmullet); Polynesia west to eastern Africa
<i>Plectropomus oligacanthus</i> (sea bass); Indonesia, Philippines, Caroline & Marshall Islands
<i>Scarus caeruleus</i> (blue parrot fish); Florida, W. Indies
<i>S. microrhinos</i> (parrot fish); Indo-Pacific
<i>Sphyrna barracuda</i> (great barracuda); Indo-Pacific, Hawaii to Red Sea, western Atlantic from Brazil to W. Indies, Florida, Bermuda
<i>Tetragomurus cuvieri</i> (squaretail); temperate waters
<i>Upeneus arge</i> (surmullet); Polynesia, Micronesia

*continued*



Chordata

- CIGUATERA POISONING (See preceding page)
- Symptoms:** Onset within 36 hours; tingling about the lips, tongue, and throat, followed by numbness of some parts. Dryness of mouth, nausea, vomiting, and abdominal cramps common. Headache, dizziness, pallor, restlessness, weakness, blurring of vision, itching, ataxia, and convulsions may occur. Deaths reported.
- Chemistry:** Heat-stable; soluble in some organ solutions, but reported insoluble in water; dialyzable. [3, 6, 57, 59, 69, 79, 105, 110, 130]
- 3 GYMNOTHORAX POISONING, caused by ingestion of any of several species of fishes, the flesh of which becomes toxic under certain conditions.
- Species & Distribution:** *Gymnothorax flavimarginatus* (moray eel); Hawaiian Islands west to eastern Africa  
*G. javanicus* (moray eel); Hawaiian Islands west to eastern Africa  
*G. meleagris* (white-spotted moray eel); Hawaiian Islands west to eastern Africa, Japan south to Australia  
*G. pictus* (speckled moray eel); Polynesia to eastern Africa  
*G. undulatus* (brown moray eel); Hawaiian Islands to Red Sea & eastern Africa
- Symptoms:** Tingling and numbness about lips, tongue, throat, hands, and feet; feeling of heaviness in the legs. Nausea, vomiting, abdominal cramps, joint pains, difficulty in swallowing and breathing, weakness, ataxia, and convulsions may occur. Deaths reported.
- Chemistry & Toxicology:** Small molecular substance. Death in mice produced by 1 ml (ip). [57, 66, 104]
- 4 SCORPION-FISH STING, caused by contact with dorsal, pelvic, or anal spines. Venoms produced by venom glands differ in three genera: *Pterois*, *Scorpaena*, and *Synanceja*.
- Species & Distribution:** *Apistus carinatus* (scorpion fish); coasts of India, Indonesia, Philippines, China, Japan, Australia  
*Centropogon australis* (waspfish); coasts of New South Wales, Queensland  
*Choridactylus multibarbis* (stonefish); coasts of India, Philippines, Polynesia  
*Inimicus didactylus* (lumpfish); Philippines, Malaya  
*I. japonicus* (stonefish); coasts of Japan  
*Minous monodactylus* (stonefish); coasts of Japan, China, S. Pacific Islands  
*Notesthes robusta* (scorpion fish); coasts of New South Wales, Queensland  
*Pterois antennata* (thread-finned zebra fish); tropical & temperate seas  
*P. lumulata* (tiger fish); coasts of Japan, Banka Islands  
*P. volitans* (turkey fish); Red Sea, Indian Ocean, Melanesia, Micronesia, Polynesia, coasts of China, Japan, Australia  
*Scorpaena guttata* (California scorpion fish); central California coast to Gulf of California  
*S. plumieri* (sculpin); Atlantic coast of N. America to Brazil  
*S. porcus* (hogfish); Atlantic coast of Europe from English Channel to Canary Islands, Mediterranean & Black Seas  
*Scorpaenopsis diabolus* (scorpion fish); coasts of Indonesia, Australia, Melanesia, Polynesia  
*Synanceja horrida* (poison stonefish); coasts of India, E. Indies, China, Philippines, Australia  
*S. verrucosa* (poison stonefish); Red Sea, coasts of Africa, India, & Australia, tropical Indo-Pacific
- Symptoms:** Immediate, severe, localized pain; pallor about wound, and sometimes symptoms of primary shock. Later manifestations include severe pain and weakness of involved extremity, dyspnea, headache, nausea, and vomiting. Coma and deaths reported.
- Chemistry & Toxicology:** Several proteins with lethal effect in one band, as separated on gel electrophoresis. LD<sub>50</sub> = 200 µg protein/kg. Produces hypotension similar to that caused by stingray and weever-fish venoms. No adrenergic-blocking action. [8, 13, 29, 56, 57, 60-62, 97, 98, 120, 124, 128, 138]
- 5 STINGRAY STING, caused by contact with bilaterally serrated, dentinal caudal spine or sting. Venom contained within ventrolateral grooves of sting, enveloped in an integumentary sheath.
- Species & Distribution:** *Aetobatus narinari* (spotted duck-billed ray); tropical & warm-temperate areas of Atlantic, Pacific, & Indo-Pacific oceans, Red Sea  
*Dasyatis dipterurus* (diamond stingray); coast of British Columbia to Central America  
*D. pastinaca* (stingray); northeastern Atlantic & Indian Oceans, Mediterranean & Red Seas  
*Gymnura marmorata* (butterfly stingray); coast of California to Mexico  
*Myliobatis californicus* (bat stingray); coast of Oregon to Lower California  
*Potamotrygon motoro* (freshwater stingray); rivers of Paraguay, Amazon south to Rio de Janeiro  
*Urolophus halleri* (round stingray); southern Pacific coast of N. America
- Symptoms:** Immediate, intense, localized pain, with increased skin temperature and discoloration or pallor about wound. Often symptoms of primary shock. Lacerated-type wound with ragged edges, some edema. Nausea, vomiting, headache, weakness, sweating, cramps, and diarrhea reported. Death extremely rare.
- Chemistry & Toxicology:** Several soluble protein fractions of low-to-average molecular weight, extremely labile and rapidly inactivated on heating. LD<sub>50</sub> of crude extract = 15 mg/kg. Small doses produce vaso-dilation; large doses, vasoconstriction. Direct effect on the heart, but not on the central nervous system or neuromuscular junction. [7, 52, 58, 111-113, 115-119]
- 6 TETRAODON POISONING, caused by ingestion of any of a number of species of puffers, the flesh of which may become toxic under certain conditions. Toxicity thought to be related to food-chain habits. Liver, gonads, intestines, and skin most toxic.
- Species & Distribution:** *Arothron hispidus* (puffer); tropical Pacific to Japan & Red Sea  
*A. meleagris* (white-spotted puffer); west coast of Central America to Indonesia  
*A. nigropunctatus* (black-spotted puffer); Polynesia, Indo-Pacific, Japan to eastern Africa & Red Sea

continued

## 96. ANIMAL TOXINS

### Part III. MARINE ORGANISMS

	Chordata
	<p><i>Canthigaster margaritatus</i> (sharp-nosed puffer); Red Sea, eastern Africa, Indonesia, China  <i>C. rivulatus</i> (sharp-nosed puffer); Japan, Hawaiian Islands  <i>Chilomycterus spinosus</i> (spiny boxfish); W. Indies, Brazil, S. Africa  <i>Colomesus psittacus</i> (freshwater puffer); rivers of Guiana, northern Brazil, W. Indies  <i>Diodon holacanthus</i> (balloonfish); tropical Atlantic, Pacific, &amp; Indian Oceans  <i>Fugu basilevskianus</i> (puffer); northern China, northwestern Korea  <i>F. chrysops</i> (puffer); Pacific coast of central Japan  <i>F. niphobles</i> (puffer); Japan  <i>F. ocellatus</i> (puffer); China, Japan, Philippines  <i>F. pardalis</i> (puffer); China, Japan  <i>F. pseudomus</i> (puffer); E. China &amp; Yellow Seas  <i>F. rubripes</i> (puffer); China to Korea, Sea of Japan, Pacific  <i>F. stictonotus</i> (puffer); southern Korea, E. China Sea, Japan  <i>F. vermicularis</i> (puffer); E. China Sea, Japan  <i>F. xanthopterus</i> (puffer); China, Korea, southern Japan  <i>Lagocephalus laevigatus inermis</i> (smooth puffer); eastern Africa, tropical Indian Ocean, Australia, E. China Sea, southern Japan  <i>L. lunaris</i> (smooth puffer); Red Sea, southern &amp; eastern Africa, India to Australia, China, Japan  <i>L. sceleratus</i> (smooth puffer); east coast of Africa to Philippines, southern Japan, Australia, Tahiti  <i>Mola mola</i> (common ocean sunfish); temperate &amp; tropical seas  <i>Sphaeroides annulatus</i> (bull's-eye puffer); Baja California to Peru, Galapagos Islands  <i>S. maculatus</i> (northern puffer); Atlantic coast of N. America to Guiana  <i>S. spengleri</i> (bandtail puffer); coasts of Texas, Florida, W. Indies, Brazil, Canary Islands, west coast of Africa  <i>Tetraodon lineatus</i> (puffer); rivers of Africa  <i>Torquigener hamiltoni</i> (puffer); Australia, Melanesia, Polynesia</p>
	<p><b>Symptoms:</b> Onset within 10-50 minutes, with tingling and numbness about mouth, lips and tongue, excessive salivation, weakness, nausea, vomiting, and difficulty in swallowing. Paresthesia and paralysis may occur over different parts of the body; convulsions and coma reported. Mortality rate about 50%.</p> <p><b>Chemistry &amp; Toxicology:</b> Suggested formula, C<sub>16</sub>H<sub>31</sub>NO<sub>16</sub>; soluble; LD<sub>50</sub> = approximately 10 µg/kg. Direct effect on neuromuscular transmission and on nerve and muscle without depolarization. Affects heart contractile force. Causes ascending-type paralysis in laboratory animals. [10, 36, 45, 46, 57, 65, 73, 77, 90, 131, 139-142]</p>
7	<p><b>TOADFISH STING</b>, caused by contact with opercular or dorsal spines. Venom produced in gland at base of spines.</p> <p><b>Species &amp; Distribution:</b> <i>Batrachus cirrhosus</i> (toadfish); Red Sea  <i>B. didactylus</i> (paddefisk); Mediterranean Sea &amp; nearby Atlantic coasts  <i>B. grunniens</i> (toadfish); coasts of Malaya, Burma, India, Ceylon  <i>Opsanus tau</i> (oyster toadfish); Atlantic coast of United States  <i>Thalassophryne reticulata</i> (venomous toadfish); Pacific coast of S. America</p> <p><b>Symptoms:</b> Onset immediate, with local pain, tenderness, increased skin temperature, and sometimes primary shock. Nausea and occasionally vomiting. Pain spreads throughout affected part; some swelling and redness about the wound. Few systemic effects; rarely serious. [9, 35, 42, 43, 49, 53, 57, 135]</p>
8	<p><b>TURTLE POISONING</b>, caused by ingestion of any one of several species of marine turtles, the flesh of which becomes toxic under certain conditions. Rare.</p> <p><b>Species &amp; Distribution:</b> <i>Chelonia mydas</i> (green sea turtle); tropical &amp; subtropical seas  <i>Dermochelys coriacea</i> (leatherback sea turtle); circumtropical, occasionally found in tropical waters  <i>Eretmochelys imbricata</i> (hawksbill sea turtle); tropical &amp; subtropical seas</p> <p><b>Symptoms:</b> Onset within 1-24 hours, with nausea, vomiting, severe abdominal cramps, increased salivation, difficulty in swallowing, and subsequent paresthesia about mouth, lips, tongue, and throat; diarrhea and tendency toward somnolence. Coma and deaths reported. [18, 57, 129]</p>
9	<p><b>WEEVER-FISH STING</b>, caused by contact with opercular or dorsal spines or stings. Spines covered by integumentary sheath; venom-producing cells within grooves of spine.</p> <p><b>Species &amp; Distribution:</b> <i>Trachinus araneus</i> (weever); Mediterranean Sea  <i>T. draco</i> (greater weever); Norway to N. Africa, Mediterranean &amp; Adriatic Seas  <i>T. radiatus</i> (weever); Mediterranean Sea  <i>T. vipera</i> (lesser weever); southern North Sea, English Channel &amp; Mediterranean Sea</p> <p><b>Symptoms:</b> Immediate, intense, localized pain, increasing in severity and spreading within an hour to entire extremity. Occasionally symptoms of primary shock. Increased skin temperature about wound, some edema, and localized discoloration or pallor. Puncture-type wound; necrosis reported in some cases. Weakness, headache, nausea, vomiting, and muscle fasciculations also reported.</p> <p><b>Chemistry &amp; Toxicology:</b> Toxin is a "muco" substance of a combined polysaccharide-protein nature. Also contains noradrenaline, adrenaline, histamine, cholinesterase, and a trace of 5-hydroxytryptamine. Free of sulfur, lecithinase, and phosphodiesterase. Direct effect on the heart, but no effect on the central nervous system or neuromuscular junction. Little hemotoxic or anticoagulant activity. [2, 13, 14, 32, 33, 50, 54, 102, 114]</p>

continued



## 96. ANIMAL TOXINS

### Part III. MARINE ORGANISMS

	Echinodermata
10	<p>SEA-CUCUMBER POISONING, caused by contact.</p> <p><b>Species &amp; Distribution:</b> <i>Euapta lappa</i> (sea cucumber); Bahamas  <i>Holothuria argus</i> (sea cucumber); Torres Strait, Pacific, Palau Islands  <i>H. atra</i> (sea cucumber); Guam, Palau Islands, Pacific  <i>H. tubulosa</i> (sea cucumber); Mediterranean  <i>Leptosynapta ooplax</i> (sea cucumber); Japan  <i>Paracaudina chilensis</i> (sea cucumber); Japan, colder seas off coasts of N. &amp; S. America  <i>Stichopus variegatus</i> (curry fish); Palau Islands, Pacific  <i>Thelenota ananas</i> (prickly red fish); Palau Islands, Japan</p> <p><b>Symptoms:</b> Localized pain, inflammation, tenderness</p> <p><b>Chemistry &amp; Toxicology:</b> Holothurin A (C<sub>50</sub>-52H<sub>81</sub>-85O<sub>5</sub>-6SNa), a steroid saponin, is comparable as a blocking agent to procaine and physostigmine on desheathed sciatic nerve of frogs, except that the action is irreversible. Produces irreversible block of direct and indirect twitch response, along with contracture, in mammalian nerve-muscle preparation. [15, 17, 23, 40, 41, 91-93, 96, 121]</p>
11	<p>SEA-URCHIN STING, caused by contact with spines or globiferous pedicellariae of any of several species of sea urchins.</p> <p><b>Species &amp; Distribution:</b> <i>Asthenosoma ijimai</i> (sea urchin); southern Japan to Molucca Sea  <i>Diadema setosum</i> (reef urchin); Indo-Pacific from eastern Africa to Polynesia, China, Japan. Related species in W. Indies, Hawaiian Islands.  <i>Toxopneustes elegans</i> (sea urchin); Japan  <i>T. pileolus</i> (sea urchin); Indo-Pacific from eastern Africa to Melanesia, Japan</p> <p><b>Symptoms:</b> Immediate, localized pain; some redness about wound and aching in injured part. In more severe cases, primary shock, muscular paralysis, and respiratory distress may occur. Death rare.</p> <p><b>Chemistry &amp; Toxicology:</b> Heat-stable; curare-like activity. [13, 21, 22, 28, 44, 89, 100, 127, 132]</p>
Annelida	
12	<p>WORM BITE, caused by bite or stinging setae.</p> <p><b>Species &amp; Distribution:</b> <i>Eurythoe complanata</i> (bristle worm); Gulf of Mexico, tropical Pacific  <i>Glycera dibranchiata</i> (bloodworm); Canadian coast to N. Carolina  <i>Lunbriconereis heteropoda</i> (marine worm); Japan</p> <p><b>Symptoms:</b> Pain similar to that caused by bee sting; some localized edema, increased skin temperature, redness, and itching.</p> <p><b>Chemistry &amp; Toxicology:</b> A tertiary amine; suggested formula, C<sub>5</sub>H<sub>11</sub>NS<sub>2</sub>. Stimulates parasympathetic activity. Direct effect on the heart and nervous system. [57, 68, 75, 94, 95]</p>
Mollusca	
13	<p>CONE STING, caused by contact with venom apparatus consisting of a bulb, duct, radular sheath, and radular teeth.</p> <p><b>Species &amp; Distribution:</b> <i>Conus aulicus</i> (court cone); Polynesia to Indian Ocean  <i>C. geographus</i> (geography cone); Indo-Pacific from eastern Africa to Polynesia  <i>C. striatus</i> (striated cone); Indo-Pacific from eastern Africa to Australia  <i>C. textile</i> (textile cone); Polynesia to Red Sea  <i>C. tulipa</i> (tulip cone); Polynesia to Red Sea</p> <p><b>Symptoms:</b> Immediate, localized pain; ischemia, and sometimes numbness about puncture wound. Tingling or paresthesia about injured part; may spread. Muscular incoordination, paresis or paralysis, and visual disturbances reported. Deaths reported within 3-5 hours.</p> <p><b>Chemistry &amp; Toxicology:</b> Toxin contains a protein, quaternary ammonium compounds, and possibly amines. Produces hyperexcitability, ataxia, dyspnea, respiratory distress in mice; paralysis in fishes and mammals. [1, 19, 24, 25, 30, 37, 72, 76, 126, 137]</p>
14	<p>OCTOPUS BITE. Venom apparatus consists of anterior and posterior salivary glands, salivary ducts, buccal mass, and mandibles or beak.</p> <p><b>Species &amp; Distribution:</b> <i>Octopus apollyon</i> (octopus); Pacific coast of N. America  <i>O. macropus</i> (octopus); Europe, Mediterranean and Red Seas, Indian Ocean, Malaysia, coasts of China, Japan, Australia  <i>O. vulgaris</i> (octopus); warm seas</p> <p><b>Symptoms:</b> Burning or tingling about wound punctures. Pain spreads to involve entire extremity. Wound bleeds freely. Some increased skin temperature; local redness and edema reported.</p> <p><b>Chemistry &amp; Toxicology:</b> Composed of octopamine, serotonin, serotonin decomposition product, histamine, dopamine, and other substances. Anticoagulant and cardiotoxic activity; no cholinesterase or aminoxidase activity. [4, 5, 11, 47, 55, 57, 67, 82]</p>
15	<p>PARALYTIC SHELLFISH POISONING, caused by ingestion of mollusks which have fed upon toxic dinoflagellates.</p> <p><b>Species &amp; Distribution:</b> <i>Cardium edule</i> (cockle); European seas  <i>Donax serra</i> (white mussel); South Africa  <i>Ensis directus</i> (razor clam); New England coast to Florida  <i>Modiolus modiolus</i> (horse mussel); Pacific coast of N. America from Arctic to Baja California; circum-boreal.</p>

continued

## Part III. MARINE ORGANISMS

	Mollusca
	<p><i>Mya arenaria</i> (soft shell clam); coasts of Britain, Scandinavia, Greenland, Japan, Atlantic &amp; Pacific coasts of N. America</p> <p><i>Mytilus californianus</i> (ocean mussel); eastern Alaska south to Socorro Island</p> <p><i>M. edulis</i> (mussel); Arctic to S. Carolina, Alaska to Baja California; found in most temperate waters</p> <p><i>Saxidomus giganteus</i> (butter clam); Alaska to central California</p> <p><i>S. nuttalli</i> (butter clam); northern California to Baja California</p> <p><i>Schizothaerus nuttalli</i> (gaper clam); Alaska to Baja California; northern coast of Japan</p> <p><i>Spisula solidissima</i> (surf clam); Labrador to N. Carolina</p> <p><b>Symptoms:</b> See entry 21.</p> <p><b>Chemistry &amp; Toxicology:</b> See entry 21. [16, 31, 39, 48, 51, 57, 83, 85-87, 90, 122, 123, 125]</p>
	Cnidaria
16	<p>HYDROID STING, caused by contact with nematocysts.</p> <p><b>Species &amp; Distribution:</b> <i>Millepora alcicornis</i> (stinging coral); tropical Pacific, Indian Ocean, Caribbean &amp; Red Seas</p> <p><b>Symptoms:</b> Burning sensation, itching, localized redness, occasional pustules and desquamation.</p> <p><b>Chemistry &amp; Toxicology:</b> See entry 18. [15, 71, 103, 134, 136]</p>
17	<p>JELLYFISH STING, caused by contact with nematocysts located for the most part on the tentacles.</p> <p><b>Species &amp; Distribution:</b> <i>Carybdea alata</i> (sea wasp); tropical Pacific, Atlantic, &amp; Indian Oceans</p> <p><i>Chiropsalmus quadrigatus</i> (sea wasp); northern Australia, Philippines, Indian Ocean</p> <p><i>Cyanea capillata</i> (giant jellyfish); N. Atlantic &amp; N. Pacific Oceans, New England to Arctic Ocean, France to northern Russia, Baltic Sea; Alaska to Puget Sound, Japan and China</p> <p><i>Dactylometra quinquecirrha</i> (pink-fringed jellyfish); Azores &amp; New England to the tropics, W. Africa, Indian Ocean, western Pacific Ocean, Malaya to Japan &amp; Philippines</p> <p><b>Symptoms:</b> Burning sensation with itching, localized edema and redness.</p> <p><b>Chemistry &amp; Toxicology:</b> (See also entry 18.) Meduscongectin present in some animals. Slow-reacting, histamine-liberating substance. Toxin hypersensitizes temperature perception organ. [34, 38, 57, 70, 74, 81, 101, 133]</p>
18	<p>PHYSALIA STING, caused by contact with nematocysts.</p> <p><b>Species &amp; Distribution:</b> <i>Physalia physalis</i> (Portuguese man-o'-war); tropical Atlantic. Related species in Indo-Pacific, Hawaii, southern Japan.</p> <p><b>Symptoms:</b> Burning, localized pain; itching, edema, redness; erythematous wheals and paresthesia in some cases. Gastrointestinal symptoms, muscular weakness, and contractures in more severe cases. Respiratory distress and secondary shock may develop. Fatal cases reported.</p> <p><b>Chemistry &amp; Toxicology:</b> Toxin contains several quaternary ammonium compounds, the most toxic being tetramine. Serotonin and histamine present; also histamine releasers and possibly several peptides; LD<sub>99</sub> = approximately 2.2 mg/kg. [38, 71, 78, 80, 84, 100, 101, 103]</p>
19	<p>SEA-ANEMONE STING, caused by contact with nematocysts.</p> <p><b>Species &amp; Distribution:</b> <i>Actinia equina</i> (sea anemone); E. Atlantic from Arctic to Gulf of Guinea, Mediterranean &amp; Black Seas, Sea of Azov</p> <p><i>Adamsia palliata</i> (cloak anemone); Norway to Spain; Mediterranean Sea</p> <p><i>Anemonia sulcata</i> (sea anemone); E. Atlantic from Norway &amp; Scotland to the Canary Islands; Mediterranean Sea</p> <p><i>Sagartia elegans</i> (sea anemone); Iceland to Atlantic coast of France, Mediterranean Sea, coast of Africa</p> <p><b>Symptoms:</b> Stinging sensation on contact; may be followed by itching, localized edema, and redness. Papules develop in severe cases. Death very rare. "Sponge fisherman's disease" attributed to <i>Sagartia</i>.</p> <p><b>Chemistry &amp; Toxicology:</b> Toxin composed of several substances, including congestin and thalassin. Congestins are water-soluble, heat-resistant, and produce vomiting, diarrhea, and visceral congestion. Thalassin is a water-soluble, alcohol-precipitated crystal, antagonistic to congestin. Scratching and sneezing in dogs caused by 100 µg. Tetramine also present. [15, 23, 74, 99, 100, 109, 136, 143]</p>
	Porifera
20	<p>SPONGE STING, caused by contact.</p> <p><b>Species &amp; Distribution:</b> <i>Fibulia nolitangere</i> (brown sponge); W. Indies</p> <p><i>Tedania ignis</i> (fire sponge); W. Indies</p> <p><i>T. toxicalis</i> (fire sponge); California coast</p> <p><b>Symptoms:</b> Burning sensation, itching, urticaria, and occasionally localized edema.</p> <p><b>Toxicology:</b> Intraperitoneal injections of crude extracts lethal to mice [27, 63]. Kills aquarium animals [26].</p>
	Protozoa
21	<p>PARALYTIC SHELLFISH POISONING, caused by ingestion of mollusks which have fed on toxic dinoflagellates. (See also entry 15.)</p> <p><b>Species &amp; Distribution:</b> <i>Gonyaulax catenella</i> (dinoflagellate); Pacific coast of United States</p> <p><i>G. polygramma</i> (dinoflagellate); Japan, S. Africa</p> <p><i>G. tamarensis</i> (dinoflagellate); Atlantic coast of N. America</p> <p><i>Gymnodinium brevis</i> (dinoflagellate); Florida coast</p> <p><i>Pyrodinium phoneus</i> (dinoflagellate); North Sea</p>

continued



96. ANIMAL TOXINS  
Part III. MARINE ORGANISMS

Protozoa
<p><b>PARALYTIC SHELLFISH POISONING</b> (See preceding page)</p> <p><b>Symptoms:</b> Onset within 10 minutes to 4 hours, with weakness, thirst, and numbness about lips, mouth, tongue, and fingertips; followed by muscular incoordination, progressive paralysis (ascending type), and respiratory failure. Death may occur within 2-24 hours.</p> <p><b>Chemistry &amp; Toxicology:</b> Basic substance forms salt with mineral acids. Optical rotation of 130°, with no absorption in ultraviolet. Formula, C<sub>10</sub>H<sub>17</sub>N<sub>7</sub>O<sub>4</sub>·2HCl; molecular weight, 372. Direct effect on heart and myoneural junction. One of most potent toxins known, approximately 3 mg fatal to humans. [51, 83, 86, 88, 90, 122, 123, 125]</p>

*Contributors:* (a) Halstead, Bruce W., and Carscallen, Leona J., (b) Russell, Findlay E.

- References:* [1] Allan, J. 1935. Med. J. Australia 22(2):554. [2] Allman, G. J. 1840. Ann. Mag. Nat. Hist., Ser. 1, 6:161. [3] Arcisz, W. 1950. U.S. Fish Wildlife Serv. Spec. Sci. Rept. Fisheries 27. [4] Baglioni, S. 1909. Arch. Ital. Biol. 51:349. [5] Baldwin, E. 1948. Dynamic aspects of biochemistry. Macmillan, New York. p. 285. [6] Banner, A. H., et al. 1960. Ann. N. Y. Acad. Sci. 90:770. [7] Bassler, H. 1942. Science 96:274. [8] Bayley, H. H. 1940. Trans. Roy. Soc. Trop. Med. Hyg. 34:227. [9] Bean, B. A., and A. C. Weed. 1910. Proc. U.S. Natl. Museum 38:511. [10] Bensen, J. 1956. J. Forensic Sci. 1:119. [11] Berry, S. S., and B. W. Halstead. 1954. Leaflets Malacol. 1:59. [12] Bhimachar, B. S. 1944. Proc. Indian Acad. Sci., B, 19:65. [13] Calmette, A. 1908. Venoms, venomous animals, and antivenomous serum-therapeutics. J. Bale and Danielsson, London. [14] Carlisle, D. B. 1962. J. Marine Biol. Assoc. U. K. 42:155. [15] Castellani, A., and A. J. Chalmers. 1919. Venomous animals, manual of tropical medicine. Ed. 3. W. Wood, New York. [16] Chambers, J. S., and H. W. Magnusson. 1950. U.S. Fish Wildlife Serv. Spec. Sci. Rept. Fisheries 53. [17] Chanley, J. D., et al. 1960. Ann. N. Y. Acad. Sci. 90:902. [18] Chevallier, A., and E. A. Duchesne. 1851. Ann. Hyg. (Paris) 46:108, 125. [19] Cilento, R. 1944. Some poisonous plants, sea and land animals of Australia and New Guinea. Smith and Paterson, Brisbane. [20] Citterio, V. 1925. Atti Soc. Ital. Sci. Nat. 64:1. [21] Clark, A. H. 1950. Bull. Raffles Museum 22:56. [22] Clark, A. H. Unpublished, 1953. [23] Cleland, J. B. 1912. Australasian Med. Gaz. 32:297. [24] Clench, W. J. 1946. Harvard Univ. Occasional Papers Mollusks 1:49. [25] Cox, J. C. 1885. Proc. Linnean Soc. N. S. Wales 9:44. [26] De Laubenfels, M. W. 1932. Proc. U.S. Natl. Museum 81:85. [27] De Laubenfels, M. W. 1953. A guide to the sponges of eastern North America. Univ. Miami Marine Laboratory, Coral Gables. p. 19. [28] Earle, K. V. 1940. Trans. Roy. Soc. Trop. Med. Hyg. 33:447. [29] Edean, R. 1961. Australian J. Marine Freshwater Res. 12:177. [30] Edean, R., and C. Rudkin. 1963. Toxicon 1:49. [31] Engelsen, H. 1922. Norsk Tidsskr. Militaermed. 26:192. [32] Evans, H. M. 1906. Brit. Med. J. 2:23. [33] Evans, H. M. 1923. Phil. Trans. Roy. Soc. London, B, 212(1):19, 27. [34] Evans, H. M. 1943. Sting-fish and seafarer. Faber and Faber, London. [35] Fish, C. J., and M. C. Cobb. 1949. Noxious marine animals of the central and western North Pacific. Woods Hole Oceanographic Institute, Woods Hole, Mass. p. 50. [36] Fish, C. J., and M. C. Cobb. 1954. U.S. Fish Wildlife Serv. Res. Rept. 36. [37] Flecker, H. 1936. Med. J. Australia 23(1):464. [38] Flecker, H. 1952. Ibid. 39(1):35. [39] Fowler, L. H. 1943. Nat. Hist. 51:228. [40] Frey, D. G. 1951. Copeia (2):175. [41] Friess, S. L., et al. 1960. Ann. N. Y. Acad. Sci. 90:893. [42] Froes, H. P. 1932. Rev. Sud-Am. Med. Chir. 3:871. [43] Froes, H. P. 1933. J. Trop. Med. Hyg. 36:134. [44] Fujiwara, T. 1935. Annotationes Zool. Japon. 15:62. [45] Fukuda, T. 1951. Clin. Studies 29(2). [46] Furukawa, T., T. Sasaoka, and Y. Hosoyo. 1959. Japan. J. Physiol. 9:143. [47] Ghiretti, F. 1960. Ann. N. Y. Acad. Sci. 90:726. [48] Gibbard, J., F. C. Collier, and E. F. Whyte. 1939. Can. Public Health J. 30:193. [49] Gill, T. 1907. Smithsonian Inst. Misc. Collections 48:391. [50] Gressin, L. 1884. These 289. Faculté de Médecine, Paris. [51] Grindley, J. R., and F. J. R. Taylor. 1962. Nature 195:1324. [52] Gudger, E. W. 1943. Bull. Hist. Med. 14:467. [53] Günther, A. 1864. Proc. Zool. Soc. London 1:155. [54] Haavaldsen, R., and F. Fonnum. 1963. Nature 199:286. [55] Halstead, B. W. 1949. Leaflets Malacol. 1:17. [56] Halstead, B. W. 1951. Calif. Med. 74:395. [57] Halstead, B. W. 1959. Dangerous marine animals. Cornell Maritime Press, Cambridge, Md. p. 146. [58] Halstead, B. W., and N. C. Bunker. 1953. Am. J. Trop. Med. Hyg. 2:115. [59] Halstead, B. W., and N. C. Bunker. 1954. Zoologica 39:61.

*continued*

Part III. MARINE ORGANISMS

- [60] Halstead, B. W., M. J. Chitwood, and F. R. Modglin. 1955. *Anat. Record* 122(3):317. [61] Halstead, B. W., M. J. Chitwood, and F. R. Modglin. 1955. *Trans. Am. Microscop. Soc.* 74(2):145. [62] Halstead, B. W., M. J. Chitwood, and F. R. Modglin. 1956. *Ibid.* 75(4):381. [63] Halstead, B. W., and R. C. Habekost. Unpublished. College Medical Evangelists, Los Angeles, Calif., 1954. [64] Halstead, B. W., L. S. Kuninobu, and H. B. Hebard. 1953. *Trans. Am. Microscop. Soc.* 72:297. [65] Halstead, B. W., and W. M. Lively. 1954. *U.S. Armed Forces Med. J.* 5:157. [66] Halstead, B. W., and R. J. Ralls. 1954. *Science* 119:160. [67] Hartman, W. J., et al. 1960. *Ann. N. Y. Acad. Sci.* 90:637. [68] Hashimoto, Y., and T. Okaichi. 1960. *Ibid.* 90:667. [69] Hessel, D. W., B. W. Halstead, and N. H. Peckham. 1960. *Ibid.* 90:788. [70] Hogberg, B. G., G. Tufvesson, and B. Uvnas. 1960. *Acta Physiol. Scand.* 38:135. [71] Hyman, L. H. 1940. *The invertebrates.* McGraw-Hill, New York. v. 1., p. 390. [72] Iredale, T. 1935. *Nautilus* 49:41. [73] Ishihara, F. 1918. *Mitt. Med. Fak. Univ. Tokyo* 20:373. [74] Junk, W., ed. 1937. *Tabulae Biologicae* 13:5. [75] Klawe, W. L., and L. M. Dickie. 1957. *Bull. Fisheries Res. Board Can.* 115:1. [76] Kohn, A. J., P. R. Saunders, and S. Wiener. 1960. *Ann. N. Y. Acad. Sci.* 90:706. [77] Lalone, R. C., E. D. De Villey, and E. Larson. 1963. *Toxicon* 1(4). [78] Lane, C. E. 1960. *Ann. N. Y. Acad. Sci.* 90:742. [79] Lee, R. K. C., and H. Q. Pang. 1945. *Am. J. Trop. Med.* 25:281. [80] Lenhoff, H. M., and W. F. Loomis, ed. 1961. *Biology of hydra and some other coelenterates.* Univ. Miami Press, Coral Gables. [81] Light, S. F. 1914. *Philippine J. Sci., D.* 9:198. [82] Livon, C., and A. Briot. 1905. *Compt. Rend. Soc. Biol.* 58:878. [83] McFarren, E. F., et al. 1957. *Proc. Natl. Shellfish Assoc.* 47:114. [84] McNeil, F. A., and E. C. Pope. 1943. *Australian Museum Mag.* 8:127. [85] Medcof, J. C., et al. 1947. *Bull. Fisheries Res. Board Can.* 75:1. [86] Meyer, K. F. 1931. *Am. J. Public Health* 21:762. [87] Meyer, K. F. 1953. *New Engl. J. Med.* 249:843. [88] Mold, J. D. 1947. Thesis. Northwestern Univ., Evanston, Ill. [89] Mortensen, T. 1943. A monograph of the echinoidea. C. A. Reitzel, Copenhagen. v. 3, pts. 2, 3. [90] Murtha, E. F. 1960. *Ann. N. Y. Acad. Sci.* 90:820. [91] Nigrelli, R. F. 1952. *Zoologica* 37:89. [92] Nigrelli, R. F., and S. Jakowska. 1960. *Ann. N. Y. Acad. Sci.* 90:884. [93] Nigrelli, R. F., and P. Zahl. 1952. *Proc. Soc. Exptl. Biol. Med.* 81:379. [94] Nitta, S. 1934. *Yakugaku Zasshi* 54:648. [95] Nitta, S. 1941. *Tokyo Igaku Zasshi* 55:285. [96] Paradise, W. E. J. 1924. *Med. J. Australia* 11(2):650. [97] Pawlowsky, E. N. 1909. *Anat. Anz.* 34:314. [98] Pawlowsky, E. N. 1914. *Zool. Jahrb.* 38:427. [99] Pawlowsky, E. N., and A. K. Stein. 1929. *Arch. Dermatol. Syphilis* 157:647. [100] Phillips, C., and W. H. Brady. 1953. *Sea pests.* Univ. Miami Press, Coral Gables. [101] Phisalix, M. 1922. *Animaux venimeux et venins.* G. Masson, Paris. [102] Pohl, J. 1893. *Prager Med. Wochschr.* 18:31. [103] Pope, E. C. 1953. *Australian Museum Mag.* 11:16. [104] Ralls, R. J., and B. W. Halstead. 1955. *Am. J. Trop. Med. Hyg.* 4(1):136. [105] Randall, J. E. 1958. *Bull. Marine Gulf Caribbean* 8:236. [106] Reed, H. D. 1900. *Proc. Am. Assoc. Advan. Sci.* 49:232. [107] Reed, H. D. 1907. *Am. Naturalist* 41:553. [108] Reed, H. D. 1924. *J. Morphol.* 38:431. [109] Richet, C. 1903. *Compt. Rend. Soc. Biol.* 55:246. [110] Ross, S. G. 1947. *Med. J. Australia* 34(2):617. [111] Russell, F. E. 1953. *Am. J. Med. Sci.* 226:611. [112] Russell, F. E. 1959. *Public Health Rept.* 74:855. [113] Russell, F. E., W. C. Barritt, and M. D. Fairchild. 1957. *Proc. Soc. Exptl. Biol. Med.* 96:634. [114] Russell, F. E., and J. A. Emery. 1960. *Ann. N. Y. Acad. Sci.* 90:805. [115] Russell, F. E., M. D. Fairchild, and J. Michaelson. 1958. *Med. Arts Sci.* 12:78. [116] Russell, F. E., and R. D. Lewis. 1956. *Publ. Am. Assoc. Advan. Sci.* 44:43. [117] Russell, F. E., and T. E. Long. 1961. In H. R. Viets, ed. *Myasthenia gravis.* C. C. Thomas, Springfield, Ill. p. 101. [118] Russell, F. E., and A. van Harreveld. 1954. *Arch. Intern. Physiol.* 62(3):322. [119] Russell, F. E., et al. 1958. *Am. J. Med. Sci.* 235(5):566. [120] Saunders, P. 1960. *Ann. N. Y. Acad. Sci.* 90:798. [121] Saville-Kent, W. 1900. *The great barrier reef of Australia.* W. H. Allen, London. p. 239. [122] Schantz, E. J. 1960. *Ann. N. Y. Acad. Sci.* 90:843. [123] Schantz, E. J., et al. 1958. *J. Assoc. Offic. Agr. Chemists* 41:160. [124] Smith, J. L. 1958. *Ichthyol. Bull.* 12:167. [125] Sommer, H., and K. F. Meyer. 1941. *Calif. Dept. Public Health Weekly Bull.* 20:53. [126] Sugitani, F. 1930. *Venus* 2:151. [127] Taft, C. H. 1945. *Texas Rept. Biol. Med.* 3:341. [128] Tange, Y. 1953. *Yokohama Med. Bull.* 4:178. [129] Taylor, E. H. 1921. *Amphibians and turtles of the Philippine Islands.* Bureau of Printing, Manila. p. 162. [130] Tennent, J. E. 1861. *Natural history of Ceylon.* Longmans, Green; London. p. 324. [131] Tsuda, K., M. Kawamura, and

*continued*

## 96. ANIMAL TOXINS

### Part III. MARINE ORGANISMS

R. Hayotsu. 1958. Chem. Pharm. Bull. (Tokyo) 6:225. [132] Tweedie, M. W. F. 1945. Poisonous animals of Malaya. Malaya, Singapore. [133] Uvnas, B. 1960. Ann. N. Y. Acad. Sci. 90:751. [134] Von Lendenfeld, R. 1887. Quart. J. Microscop. Sci., N.S. 27:393. [135] Wallace, L. B. 1893. J. Morphol. 8:563. [136] White, R. P. 1934. The dermatergoses or occupational affections of the skin. H. K. Lewis, London. p. 404. [137] Whyte, J. M., and R. Endean. 1962. Toxicon 1:25. [138] Wiener, S. 1958. Med. J. Australia 45(2):219. [139] Yokoo, A. 1952. Proc. Japan Acad. 28:200. [140] Yokoo, K. 1948. Hiroshima Igaku 2:52. [141] Yokoo, K. 1948. Riken Iho (Tokyo) 24:136. [142] Yudkin, W. H. 1945. J. Cellular Comp. Physiol. 25:85. [143] Zervos, S. G. 1934. Paris Med. 93:89.

## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
1 <i>Abrus precatorius</i> (rosary pea); seeds, possibly root; southern Florida & tropics	Abrin (N-methyl- tryptophane) and abric acid [49]	Onset may be delayed several hours to two days; vomiting, diarrhea, acute gastroenteritis, chills, sometimes con- vulsions [34]. Severe cases: death from heart failure [49].	Two ounces of seed fatal to horse; one seed chewed may be fatal to child [49]. Boiled seeds may be eaten, but in quantity cause headache [10].
2 <i>Aconitum napellus</i> (aconite monks- hood); roots, leaves, flowers, seeds; northeastern United States, Canada, Europe, Asia [23]	Principally aconitine, aconine, napelline; eight other alka- loids reported [51]	Tingling, burning sensation in tongue, throat, skin; great restlessness, dysp- nea, slow pulse, muscular weakness, incoordination, cold and livid skin, pu- pillary constriction, followed by dilation [44]; vomiting, diarrhea, convulsions, possibly death in 1-8 hours by respira- tory or cardiac paralysis [17].	Young leaves mistakenly eaten for parsley, and the roots for horseradish. Considered most danger- ous of all British plants [17].
3 <i>Agrostemma githago</i> (corn cockle); seeds; United States, Canada, Europe [34]	Githagin, agrostem- mic acid (saponins) [34,49]	Man: vertigo, diarrhea, depressed breathing [34]; irritation of digestive tract, vomiting, headache, sharp pains in spine, difficult locomotion, some- times coma and death [8]. Horses, cattle: colic, diarrhea, muscular tremors, rigidity, coma, death [9].	Milled seeds sometimes in wheat flour [17]; frequent ingestion of small amounts results in chronic githagism [34]; 0.25-1 lb/100 lb live wt fatal to stock [34].
4 <i>Amanita phalloides</i> (death cup); entire fungus; N. America, Europe, E. Indies, Australia	Phalloidin [17]; <i>Amanita</i> hemolysin possibly a factor in poisoning when mushroom is eaten raw [16]	After 6-15 hours, abdominal pain [17]; vomiting, diarrhea, intense thirst, re- current drowsiness [23,28]; respiratory and circulatory depression, delirium, and sometimes convulsions [40]; jaun- dice, hepatitis and renal disturbances, coma, death from heart failure.	One of the most deadly fungi, mortality about 50%; cause of majority of "mushroom deaths" in U.S. [17,40,44]. Genus contains other equally poisonous, and some edi- ble, species [21].
5 <i>Antiaris toxicaria</i> (upas tree); milky sap; southern Asia, E. Indies [40]	Antiarin ( $\alpha$ -, $\beta$ -) [7]	Skin irritation, blistering, swelling [37]; vomiting, convulsions, death [7].	One of the principal arrow poisons; more potent than digitalis [7,40]. Cloth made from bark causes severe itching if sap not completely re- moved [7].
6 <i>Astragalus</i> spp. (lo- cowed); fresh plant; northern hemisphere [34]	Selenium; locoine in some species [34]	Horses, cattle: dullness, weakness, ir- regularity in behavior, impaired vision, edema of eyelids, loss of muscular con- trol, depraved appetite, emaciation, starvation, death. Sheep: same symptoms as above and pos- sibly blindness [34].	Hazard to livestock indus- try in U.S.; toxicity var- ies with locality [8]; some species harmless.

continued



# Contrails

## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
7 <i>Atropa belladonna</i> (belladonna); entire plant, especially seeds, roots, leaves; eastern United States, Europe, Asia [34]	Chiefly hyoscyamine; atropine and hyoscyamine in small amounts [17]; six other alkaloids reported [51]	Man, acute: dryness of skin, mouth, throat [10]; difficulty in swallowing, flushing of face, cyanosis, mydriasis, nausea, vomiting, constipation, slurred speech, giddiness, stupor, coma, rapid and weak pulse, fever, death from asphyxia and heart failure [23]. Chronic: erythema, urticaria, vesicular eruptions, slurred speech, mydriasis, glaucoma, muscular tremors or twitchings; sudden withdrawal causes nausea, salivation, perspiration [6]. Cattle: mydriasis, constipation, rapid pulse, labored breathing, frenzy, paralysis [23].	Cultivated as source of the drug belladonna [35]. Children and grazing animals often poisoned by eating fruit of plant. Flesh of rabbits which have fed on plant is toxic to humans [17].
8 <i>Cannabis sativa</i> (hemp); upper leaves, flower bracts of female plants; United States, Mexico, tropical America, Europe, temperate Asia [34]	Cannabinol and cannabidiol, the latter isomerized to highly active tetrahydrocannabinol [47]	Man: exaltation, inebriety, confusion, followed by central nervous system depression. Prolonged addiction may produce dullness or mania [20,45]; large quantities may result in death from cardiac depression [10].	Dried leaves and bracts smoked by drug addicts; seeds harmless [34,49]. Used in "bird seed"; yields hemp seed oil for paints, etc.; plant stem yields hemp fiber for cordage, carpets, etc.
9 <i>Cicuta</i> spp. (water hemlock); primarily roots; leaves, stems less toxic; northern temperate regions [17,34]	Cicutoxin [34]	Man, other animals: abdominal pain, nausea, vomiting, diarrhea, mydriasis, labored breathing, foaming at mouth, weak and rapid pulse, epileptoid convulsions, death from respiratory failure [17,34].	Genus includes the most poisonous plants in U.S. Fatalities have resulted from mistaking roots for parsnips [17,34].
10 <i>Claviceps purpurea</i> (ergot claviceps); sclerotium; N. America, Europe, Asia, Australia [35]	Ergotoxine, ergotamine, ergonovine, ergometrine [18]	Man, acute: vomiting, diarrhea, respiratory difficulties, visual and motor disturbances followed by convulsions, lowered blood pressure, shallow respiration, unconsciousness. In pregnancy, possibly uterine hemorrhage, abortion, peripheral gangrene. Chronic, convulsive type: vomiting, itching, paresthesia, analgesia of extremities, anorexia or uncontrollable hunger, diarrhea, muscle contracture, delirium, sometimes a tabes-like complex. Gangrenous type: pustules may form, limbs swell and become hot, and gangrene may follow. [44] Cattle: gastrointestinal irritation, gangrene of extremities, uterine contractions, nervous disturbances [45].	Occurs on rye, wheat, oats, barley, and other grasses; cause of many cases of poisoning (ergotism) in man and livestock [4,45]. Ergot preparations valued medically, chiefly for effect on muscles of uterus [17]; ergometrine causes abortion [18].
11 <i>Colchicum autumnale</i> (autumn crocus); entire plant, principally mature corms [38], seeds [32]; United States, Europe, N. Africa [17]	Colchicine [34]	Man: burning in throat; 6-8 hours later a feeling of suffocation, oppression in chest, difficulty in swallowing, vomiting, diarrhea, colic, tenesmus, giddiness, weakness in legs, arthralgia, cyanosis, labored breathing, convulsions [31,45]. Death from respiratory exhaustion in 7-36 hours; consciousness preserved to end. Other animals: nausea, vomiting, colic, diarrhea, hematuria, depression, unconsciousness, paralysis, mydriasis, profuse perspiration, death in 1-3 days [45].	Widely grown in flower gardens. Colchicine arrests mitosis [32]; employed in treatment of gout [43]; also used in plant breeding.
12 <i>Conium maculatum</i> (poison hemlock); fruit, especially unripe; stems, leaves, roots; N. America, temperate S. America, Europe, N. Africa, Asia [17,34]	Coniine, conhydrine, N-methylconiine, coniceine [9,34]; 3 other alkaloids reported [51]	Man: mydriasis [17]; paralysis of extremities, muscular weakness, often blindness, death from respiratory paralysis [23,34]. Cattle: mydriasis [17]; inappetence, salivation, bloating, muscular weakness [34]; coma [23]. Horses: mydriasis [17]; nausea, grinding of teeth, rapid and labored respiration, paralysis, death from respiratory failure [23].	Leaves most toxic when plant is flowering; root less toxic in spring [34]. Resemblance of fruit to anise, leaves to parsley, and root to parsnips responsible for many human fatalities [17]. Plant commonly fatal to livestock [8].

continued



# Contrails

## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
13 <i>Croton tiglium</i> (purging croton); roots, leaves, bark, seeds; southern Asia, E. Indies, Pacific Islands, Africa [40]	Croton, croton resin [37]; ricinine [51]	Vomiting, drastic purging, possibly col- lapse and death [40]. Croton oil is a skin irritant, causing reddening, swell- ing, pustules [7].	Croton oil formerly a hu- man and veterinary pur- gative; abandoned as too violent. Now used to lu- bricate machinery. Smoke from burning wood inflames eyes [7].
14 <i>Datura stramonium</i> (jimsonweed datu- ra), <i>D. metel</i> (Hin- du datura), and other species; en- tire plant, espe- cially seeds; tem- perate, tropical and subtropical regions	Hyoscyamine, also hyos- cyamine (optically active forms), and the racemic mix- ture, atropine [3]	Man: headache, nausea, vertigo, thirst, dry and burning sensation in skin, loss of muscular control, mydriasis. Acute poisoning results in mania, convulsions, death [34]; nonfatal poisoning usually causes loss of memory, mental confu- sion [40,45]. Cattle: mydriasis, suspension of secre- tions or diarrhea, rapid heart action, paralysis, death from asphyxia. Swine: convulsive twitching [23].	Daturas widely grown for ornament. Children poisoned by eating seeds or sucking flower. In Asia and Africa, adults poisoned by ingesting seeds for intoxicating ef- fect. Accidental mixing of seed with grain, or gath- ering of young plants with other greens, also respon- sible for poisoning. [50] Products used medicinally, also criminally, for narcotic effect [45].
15 <i>Delphinium</i> spp. (larkspur); seeds, leaves; to a lesser degree, roots; north temperate regions, especially western United States	Delphinine, delphi- noidine, delphisine, staphisagrine [34]; 30 other alkaloids reported [51]	Man: burning and inflammation of mouth and pharynx, lowered blood pressure, nausea, abdominal pain, labored respi- ration, itching, cyanosis. Other animals: uneasiness, stiffness, staggering, constipation, frothing at mouth, nausea, bloating [34]; spasms, respiratory failure [49].	Second to <i>Astragalus</i> in causing fatalities among livestock in U.S. [34]; leaves and seeds may cause dermatitis. Horti- cultural varieties com- mon in flower gardens. Seeds long used in inse- cticide [49].
16 <i>Digitalis purpurea</i> (common foxglove), and <i>D. lanata</i> (Gre- cian foxglove); en- tire plant, espe- cially seeds, leaves; western Europe, western United States, other areas [23]	<i>D. purpurea</i> : diac- tyldigilanid A and B; <i>D. lanata</i> : digi- lanid A, B, and C	Anorexia, nausea, vomiting, slow and pronounced pulse in early stages [34]; cardiac arrhythmias, diarrhea, abdomi- nal pain, headache, fatigue, malaise, drowsiness, convulsions, rapid irregu- lar pulse, death in severe cases [19].	Common in flower gardens [34]. Dry plants in hay have poisoned horses and cattle; fresh leaves fatal to turkeys [17]. Digitalis and derivatives used in cardiovascular therapy [37].
17 <i>Dioscorea hispida</i> (wild yam); entire plant, tubers; southern Asia, E. Indies, Pacific Islands [7]	Dioscorine [40,50]	Discomfort, then burning, in throat; gid- diness, vomiting of blood, suffocation, drowsiness, exhaustion [7]; paralysis of nervous system [40,50].	Raw tubers a frequent cause of death in Philippines [40]. Edible after grating, boiling, repeated washing and soaking [7].
18 <i>Erythroxylon coca</i> (Huanuco cocaine tree); leaves; northern S. America, tropics of both hemi- spheres [40]	Cocaine; 13 other alkaloids reported [51]	Acute: general central nervous system stimulation, followed by depression, numbness of tongue, paralysis of respi- ratory centers, cyanosis, shallow and irregular breathing, often sudden death from asphyxia [44].	Leaves commonly chewed as a stimulant by Indians of Peru and Bolivia. Co- caine used as a local anesthetic; misused by drug addicts.
19 <i>Euphorbia</i> spp. (euphorbia); milky sap; worldwide [34]	A complex substance, euphorbiosteroid [17]; euphorbon, euphorboresene, and an acrid sub- stance reported [7, 9,34,52]; also cyan- ogenetic glucosides in some species [18]	Man: dermatitis, eye irritation, tempo- rary blindness [23]; swelling around mouth and eyes, burning in mouth and throat, sneezing, vomiting, diarrhea, hemorrhagic gastroenteritis [49]; faint- ing, death [7,34]. Other animals: blistering of skin, loss of hair, weakness, collapse, death [34].	Various species grown for ornament. Euphorbium derived from <i>E. resinif- era</i> (gum euphorbia), formerly used medici- nally, now in paint as protectant [7]; <i>Euphorbia</i> sap mixed with arrow poison as cohesive irri- tant [49].
20 <i>Helleborus niger</i> (black Christmas rose); rootstock, leaves; Europe, United States [34]	Helleborin, hellebo- rein, hellebrin [17, 34,37]	Man: severe dermatitis in some individ- uals [34]; violent inflammation of mu- cous membranes of stomach and intes- tines [36]; vomiting, dizziness, convul- sions, sometimes death; effect on heart similar to digitalis [37].	Cultivated in flower gar- dens; rootstock former- ly source of an official drug [37].

continued

## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
21 <i>Hippomane mancinella</i> (manchineel); milky sap, fruit; Florida, Central America, northern S. America, W. Indies [27]	Physostigmine or similar alkaloid, plus a sapogenin [27]	Man: severe burning of skin, swelling and possibly hemorrhage in eyes; temporary blindness from sap. [1,27] Fruit causes gastroenteritis which may be fatal; ulceration of intestinal tract proceeds slowly [27,34].	Apparently more toxic in summer than in winter [27]. Sap used as arrow poison; smoke from burning wood toxic [1].
22 <i>Hyoscyamus niger</i> (black henbane); entire plant, especially leaves; northwestern United States, Canada [34], Mediterranean region, Asia, Oceania	Hyoscyamine, hyoscyne, atropine, and other alkaloids [22, 34,36]	<i>See Atropa belladonna</i>	Usually avoided by animals because of unpleasant taste. Children poisoned by eating seeds and pods [34]. Roots mistaken for parsnips [17]. Leaves and flowering tops used in medicine [35].
23 <i>Jatropha curcas</i> (Barbados nut); seeds, milky sap; tropics [49], southern Florida	Curcin [50]	Man: burning in throat, bloating, dizziness, vomiting, diarrhea, drowsiness, possibly dysuria and mydriasis [45]; severe leg cramps, deafness [24]; violent purgative action often fatal to children [1]. Other animals: hemorrhagic enteritis, staggering, dull vision, mydriasis, bloating, paralysis, somnolence, convulsions, fever, shivering, coma, death in 1-3 days in acute cases [45].	Sap used as fish poison [7]; seeds yield "hell oil," formerly given as purgative, now used for soap in Cape Verde Islands and Ecuador. Seeds of <i>J. multifida</i> , a tropical ornamental, often cause poisoning in children.
24 <i>Kalmia latifolia</i> (mountain laurel), and <i>K. angustifolia</i> (lambkill kalmia); all parts except wood; northeastern United States, Canada, Pacific coast [8]	Andromedotoxin [34]	Man: similar to symptoms produced in other animals, plus pain in head, sweating, tingling of skin [10]. Other animals (usually sheep): salivation, flow of tears, secretions from nose, frothing at mouth, impaired vision or blindness, dizziness, irregular respiration, vomiting, convulsions, paralysis of limbs, coma, death [8,34].	Children have been poisoned by eating leaves mistaken for wintergreen [8]. Frequent cause of fatal poisoning of livestock, especially sheep.
25 <i>Lathyrus sativus</i> (grass peavine); seeds, mature plant; southern United States, southern Europe, N. Africa, Asia [34,45]	$\beta$ -Aminopropionitrile [47]	Man: pain in back [49]; sudden weakness in legs; further ingestion may cause leg paralysis. Other animals: similar to symptoms produced in man, plus asphyxia. Cattle also develop constipation, weak pulse, numbness of skin [45]; general weakening of tendons and ligaments, tissue fragility; connective tissue malformations, such as exostoses, hernias, and aneurysms [47].	<i>L. sativus</i> , <i>L. cicera</i> , and <i>L. clymenum</i> used as food and fodder [49], but cause many cases of lathyrism in man and livestock [45]. <i>L. odoratus</i> , the sweet pea of flower gardens, has fatally poisoned children.
26 <i>Leucaena glauca</i> (white popinac lead tree); leaves (especially immature), bark, roots; southern United States, tropics [40,46]	Mimosine (leucenol) [46]	Horses, mules, donkeys: alopecia of manes and tails, possibly deformation or loss of hoofs [23,50]; in severe cases, lameness, debility, death from hunger and thirst [18]. Swine: total alopecia, impaired vision, emaciation, various degrees of paralysis, respiratory failure [23,50].	A fodder plant for cattle, sheep, goats (immune to toxicity) [23]. Ripe seeds used as coffee substitute [39]. Young leaves and unripe seeds, cooked as vegetables, occasionally cause loss of hair in humans [30].
27 <i>Lupinus</i> spp. (lupine); seeds most toxic, pods less, leaves least; temperate regions [45]	Lupanine and 30 other alkaloids reported [51]	Liver damage due to poisoning by ingestion. European lupinosis, chronic: anemia, cachexia. Acute: fever, general jaundice, coma, paralysis, constipation, then hemorrhagic diarrhea, swelling of ears, eyelids, lips, nose. American lupinosis: frothing at mouth, dyspnea, frenzied actions, nausea, bloating, coma, possibly death [17,18, 45].	Toxicity varies with season and location; some species harmless [34]. Many livestock deaths, especially of sheep, from ingestion of lupine seeds in quantity [18].

continued

## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
28 <i>Manihot esculenta</i> (cassava); roots (especially skin and juice), mature leaves, stems, fruit; tropics [7,40]	Cyanogenetic gluco- sides [42]	Man, livestock: nausea [50]; rapid and labored breathing; rapid, irregular, weak pulse; twitching, staggering, spasms of neck and legs, convulsions, mydriasis, coma, death from respira- tory paralysis [45].	Bitter cultivars, high in CN <sup>-</sup> , yield starch for tapioca; sweet cultivars, with little or no CN <sup>-</sup> , widely cooked as starchy vegetables [42].
29 <i>Melia azedarach</i> (chinaberry); fruit pulp, bark, flowers; southern United States, tropical America, S. Africa, southwestern Asia [45]	Azadarin (margosine) affects central nervous system [34, 45,49]	Man, leaf poisoning: stomatitis, decrease in urine formation, violent and bloody vomiting [45]. Fruit poisoning: nausea, vomiting, labored breathing, palpitation, paralysis [34]. Other animals (especially swine): vomit- ing, colic, diarrhea, labored breathing, cyanosis, convulsions or paralysis, death by asphyxia [49].	Common shade tree. Roots, bark, leaves, flowers, fruit used for stupefying fish [1]; vari- ous parts of tree used in folk medicine [49]; seeds yield medicinal oil [49].
30 <i>Metopium toxiferum</i> (poisonwood); en- tire plant, espe- cially sap; southern Florida, W. Indies [11], Bahamas	Probably similar to poison ivy	Dermatitis similar to that caused by poi- son ivy [11,34]. Blistering may continue for weeks, readily spreading from one area to another; may be accompanied by intense itching, burning. Severe cases require hospitalization.	Smoke from burning wood highly irritating [12]. Clear, sticky sap turns black on exposure to air.
31 <i>Nerium oleander</i> (oleander); leaves, bark, roots, flow- ers; southern United States, tropics, subtropics [40]	Principally neriin, oleandrin, and foli- nerin, resembling digitalis in action; rosagenin (in bark) similar to strych- nine [49]	Man: vomiting, slow and irregular pulse, bloody diarrhea, death from cardiac or respiratory paralysis. Other animals: similar to symptoms pro- duced in man, plus sweating, gnashing of teeth, groaning, sometimes polyuria [45].	Popular ornamental shrub. Often poisons grazing animals; 15-20 g fresh plant fatal to horse. Smoke from burning plant toxic; meat roasted on skewers of oleander wood, or food stirred with oleander sticks, fa- tally toxic [1,23,45].
32 <i>Papaver somniferum</i> (opium poppy); milky exudate from incised unripe seed pod, which is dried as "opium" [34]; southern Europe, Asia, tropics, subtropics	Morphine (chiefly); also codeine, the- baine, papaverine, and narcotine [45]	Acute: central nervous system depres- sion, symmetrical pinpoint pupils, de- pressed respiration, cyanosis, coma, death from depression of respiration and circulation. Chronic: varies with individual case [48].	Cultivated in flower gar- dens as well as for com- mercial drug production. Animals infrequently poi- soned by eating seed pods [34]. Seeds harmless, commonly used in bakery products.
33 <i>Phytolacca ameri- cana</i> (pokeberry); roots, seeds, ma- ture (red) stems; eastern & southern United States [34], Europe, S. Africa [49]	Phytolaccine [51]	Burning and bitterness in mouth, vomit- ing, purging, spasms, sometimes con- vulsions, death from respiratory paral- ysis [34].	Young shoots edible if well-cooked [34]; fruit juice harmless, used as food coloring [19]. Root has been accidentally gathered with shoots, or mistaken for parsnips or horseradish [34,49].
34 <i>Prunus</i> spp. (choke- cherry, other wild cherries); leaves (especially wilted), bark, seeds; north- ern hemisphere, Orient [34]	Hydrocyanic acid formed by action of enzymes upon amygdalin (?) or prunasin [34]	Animals: uneasiness, staggering, falling, convulsions, labored breathing, bloat- ing, death [34].	Frequent cause of fatal poisoning of livestock [34].
35 <i>Rhus toxicodendron</i> (poison ivy), and <i>R. vernix</i> (poison su- mac); entire plant; N. America [34]	Urushiol [14]	Man: skin irritation, swelling, blisters, extreme discomfort, itching; sometimes fatal to children [34].	Smoke from burning plant toxic; estimated 1,000,000 Americans get ivy poisoning each year [14].
36 <i>Ricinus communis</i> (castor bean); seeds; southern United States, trop- ics, subtropics [34]	Ricin (a toxalbumin), and the less toxic alkaloid, ricinine [18]; also a castor bean allergen [41]	Man: burning in mouth, throat, stomach; vomiting; diarrhea; thirst; rapid, then faint pulse; cramps of abdomen, legs; convulsions; shallow respiration. Other animals: hemorrhagic enteritis,	Some varieties cultivated as ornamentals; 2-3 seeds may be fatal to child, 6 may kill horse [49]. Seeds yield castor

continued



## 97. PLANT TOXINS

Plant (Common Name); Toxic Portion; Distribution	Toxic Principle	Signs and Symptoms Produced	Remarks
(A)	(B)	(C)	(D)
<i>Ricinus communis</i> (See preceding page)		staggering, dulled vision, heart weakness, bloating, paralysis, convulsions, fever, shivering, coma, death in 1-3 days [45].	oil; processors subject to conjunctivitis, dermatitis, and respiratory allergy; efforts being made to detoxify protein-rich presscake for feed [41].
37 <i>Senecio</i> spp. (groundsel); entire plant, especially seeds; worldwide	Of numerous toxic alkaloids isolated, senecifolidine, isatidine, pterophine, retrorsine, sclerartine, and senecionine are responsible for "bread poisoning" and liver cirrhosis [49].	Man: abdominal pain, vomiting, ascites, enlarged liver, emaciation, bloody diarrhea. Generally fatal if not treated early; when not immediately fatal, liver damage may bring about subsequent death [45]. Grazing animals: yawning, inappetence, emaciation, staggering, colic [34]; unconsciousness, death from liver cirrhosis [49].	Seeds of various species in harvested grain considered responsible for "bread poisoning" [13]; senecio poisoning common in livestock [34].
38 <i>Strophanthus</i> spp. (strophanthus); seeds, roots, bark; southern Florida, tropical America, tropical Africa [40]	An alkaloid, trigonelline [51], and 64 cardiac glucosides and aglycons reported [49]	Vomiting, slow and irregular pulse, blurred vision, delirium, death from circulatory failure of cardiac origin [13,44].	Arrow poisons from several species [40]. <i>S. sarmentosus</i> is source of sarmentogenin, which is chemically converted to cortisone [33,49].
39 <i>Strychnos nuxvomica</i> (nuxvomica poison nut); seeds, leaves, bark, wood, flowers; India, Hawaii [2,23,40]	Strychnine, brucine [40]; six other alkaloids reported [51]	Action on spinal cord causes excessive reflex irritability, followed by rapid tonic convulsions with intermissions of exhaustion and sweating; extreme muscular rigidity, asphyxia, death. Mind not affected [40].	Strychnine formerly used as stimulant; tonic in minute amounts [40]. Children poisoned by pills. Poisoned grain used as gopher bait.
40 <i>S. toxifera</i> (curare poison nut); bark, roots; Central America, northern S. America [1]	Principally toxiferines I-XII [29], caracurines I-IX; 12 other alkaloids reported [51]	Haziness of vision, relaxation of facial muscles, inability to raise head, loss of muscular contraction in arms and legs, depressant effects on muscles of respiration, muscle nerve end-plate paralysis [15].	Often main ingredient in certain kinds of curare famed as Indian blowgun poison [15,29]. Tubocurarine chloride (U.S.P.) now used as skeletal muscle relaxant in shock therapy and as diagnostic aid [25].
41 <i>Thevetia peruviana</i> (thevetia); seeds, leaves, bark, roots, milky sap; tropics [7,40]	Thevetin, thevetoxin, neriifolin [49]	Man: vomiting, diarrhea, high blood pressure, erratic heart beat, death from asphyxiation and sudden heart paralysis [2,40]. Contact with sap may inflame and blister skin [1].	Seeds used as fish poison, and for suicide and homicide.
42 <i>Veratrum viride</i> (American false hellebore); entire plant; United States, Canada [8, 26]	Protoveratrine A and B, germerine, jervine; 15 other alkaloids reported [51]	Man: vomiting, abdominal pain, muscular weakness, spasms, possibly convulsions, rapid pulse, shallow breathing, semiconsciousness, death from asphyxia [34].	This and related species yield veratrum, a therapeutic agent for hypertension [26].
43 <i>Zigadenus</i> spp. (death camas); leaves, stems, flowers, bulb; northern hemisphere, especially United States [34,38]	Zygadenine, similar to veratrine and cevadine in action [34]	Animals: salivation, vomiting, lowered temperature, staggering or collapse, labored breathing, paralysis, possibly coma and death [5,34].	Frequent cause of fatal poisoning of livestock. Children occasionally poisoned by eating bulb [34].

Contributors: Larson, Edward, and Morton, Julia F.

References: [1] Allen, P. H. 1943. Am. J. Trop. Med. 5:23. [2] Arnold, H. L. 1944. Poisonous plants of Hawaii. Tongg, Honolulu. [3] Avery, A. G., S. Satina, and J. Rietsema. 1959. Blakeslee: the genus *Datura*. Ronald Press, New York. [4] Barger, G. 1931. Ergot and ergotism. Gurney and Jackson, London. [5] Beath, O. A., et al. 1939. Univ. Wyoming Agr. Expt. Sta. Bull. 31. [6] Brookes, V. J. 1958. Poisons: their properties, chemical identification, symptoms, and emergency treatments. Ed. 2. Van Nostrand, New York. [7] Burkill, I. H.

continued



1935. A dictionary of the economic products of the Malay Peninsula. Crown Agents for the Colonies, London.
- v. 1, 2. [8] Chesnut, V. K. 1898. U.S. Dept. Agr. Div. Botany Bull. 20. [9] Connor, H. E., and N. M. Adams. 1951. New Zealand Dept. Sci. Ind. Res. Bull. 99. [10] Council of Scientific and Industrial Research. 1948. The wealth of India. New Delhi. [11] Crooks, D. M., and L. W. Kephart. 1946. U.S. Dept. Agr. Farmers' Bull. 1972. [12] Dahlgren, B. E., and P. C. Standley. 1944. U.S. Bur. Med. Surg. Navy Med. 127. [13] Dalziel, J. M. 1948. Useful plants of west tropical Africa. Crown Agents for the Colonies, London. [14] Dawson, C. R. 1956. Trans. N. Y. Acad. Sci., II, 18(5):427. [15] Fanshawe, D. B. 1950. Brit. Guiana Forest Dept. Bull. 2. [16] Ford, W. W., and E. D. Clark. 1914. Mycologia 6:167. [17] Forsyth, A. A. 1954. Min. Agr. Fisheries (London) Bull. 161. [18] Gardner, C. A., and H. W. Bennetts. 1956. The toxic plants of western Australia. West Australia Newspaper, Perth. [19] Goodman, L. S., and A. Gilman. 1955. The pharmacological basis of therapeutics. Ed. 2. Macmillan, New York. [20] Grollman, A. 1958. Pharmacology and therapeutics. Ed. 3. Lea and Febiger, Philadelphia. [21] Grossman, C. M., and B. Malbin. 1954. Ann. Internal Med. 40:249. [22] Henry, T. A. 1949. The plant alkaloids. Ed. 4. Blakiston, Philadelphia. [23] Hurst, E. 1942. Poison plants of New South Wales. Snelling, Sydney, Australia. [24] Kirtikar, K. R. 1903. Poisonous plants of Bombay. Bombay. [25] Krantz, J. C., and C. J. Carr. 1961. The pharmacologic principles of medical practice. Ed. 5. Williams and Wilkins, Baltimore. [26] Kraye, O., and G. H. Acheson. 1946. Physiol. Rev. 26:383. [27] Lauter, W. M., and P. A. Foote. 1955. J. Am. Pharm. Assoc. Sci. Ed. 44:361. [28] Lewes, D. 1948. Brit. Med. J. 2:383. [29] Lilly Research Laboratories. 1951. Res. Today 7:2. [30] McKee, H. S. 1958. South Pacific Comm. Quart. Bull. 8(3):62. [31] Macleod, I. G., and L. Phillips. 1947. Ann. Rheumatic Diseases 6:224. [32] Manske, R. H. F., and H. L. Holmes, ed. 1952. The alkaloids. Academic Press, New York. v. 2. [33] Monachino, J. 1950. J. N. Y. Botan. Garden 51(602):25. [34] Muenscher, W. C. 1951. Poisonous plants of the United States. Rev. ed. Macmillan, New York. [35] Nayar, S. L., and I. C. Chopra. 1951. Distribution of British pharmacopoeial drug plants and their substitutes growing in India. Council of Scientific and Industrial Research, New Delhi. [36] Nelson, A. 1951. Medical botany. E. and S. Livingstone, London. [37] Osol, A., and G. E. Farrar. 1955. The dispensatory of the United States of America. Ed. 25. J. B. Lippincott, Philadelphia. [38] Pammel, L. H. 1910-11. A manual of poisonous plants. Torch Press, Cedar Rapids, Ia. [39] Pratt, R., and H. W. Youngken, Jr. 1956. Pharmacognosy. Ed. 2. J. B. Lippincott, Philadelphia. [40] Quisumbing, E. 1951. Philippines Dept. Agr. Nat. Resources Tech. Bull. 16. [41] Raymond, W. D. 1961. Trop. Sci. 3(1):19. [42] Rogers, D. J. 1963. Bull. Torrey Botan. Club 90(1):43. [43] Salter, W. T. 1952. A textbook of pharmacology. W. B. Saunders, Philadelphia. [44] Sollmann, T. H. 1957. A manual of pharmacology. Ed. 8. W. B. Saunders, Philadelphia. [45] Steyn, D. G. 1934. Toxicology of plants in South Africa. Central News Agency, South Africa. [46] Takahashi, M., and J. C. Ripperton. 1949. Hawaii Agr. Expt. Sta. Bull. 100. [47] U.S. Department of Health, Education, and Welfare. 1963. Natl. Inst. Health Record 15(15):5. [48] Von Oettingen, W. F. 1952. Poisoning. P. B. Hoeber, New York. [49] Watt, J. M., and M. G. Breyer-Brandwijk. 1962. Medicinal and poisonous plants of southern and eastern Africa. Ed. 2. E. and S. Livingstone, London. [50] Webb, L. J. 1948. Australia Council Sci. Ind. Res. Bull. 232. [51] Willaman, J. J., and B. C. Schubert. 1961. U.S. Dept. Agr. Tech. Bull. 1234. [52] Youngken, H. W. 1948. Textbook of pharmacognosy. Ed. 6. Blakiston, Philadelphia.

## X. BIOPHYSICAL AND BIOCHEMICAL CHARACTERISTICS

### 98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS

All data are for crystalline substances, unless otherwise specified. In Parts I-V, the selection of substances was restricted to natural carbohydrates found free (or in chemical combination and released on hydrolysis) and to biological oxidation products of the natural carbohydrates. In Part VI, the selection of oligosaccharides was restricted to those substances found free. The nomenclature conforms with that of the British-American report as published in the *Journal of Organic Chemistry*, 28:281 (1963). Substances have been arranged alphabetically under the name of the parent sugar within groups formulated according to increasing carbon content (excluding carbon in substituents), with synonymous common names in parentheses. **Melting Point:** b.p. = boiling point; d. = decomposes; s. = sinters. **Specific Rotation** was determined in water at concentrations of 1-5 g per 100 ml of solution and at 20°-25°C, unless otherwise specified; other temperatures or wavelengths are shown in brackets; c = grams solute per 100 ml of solution. The literature has been covered by means of *Chemical Abstracts* through 1962.

#### Part I. NATURAL MONOSACCHARIDES: ALDOSES AND KETOSES

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
	(A)	(B)	(C)	(D)	(E)
Aldoses					
1	D-Glyceraldehyde	C <sub>3</sub> H <sub>6</sub> O <sub>3</sub>	.....	+13.5±0.5 (syrup)	122
2	D-Glyceraldehyde, 3-deoxy-3,3-C-bis(hydroxymethyl)- (Cordycepose)	C <sub>5</sub> H <sub>10</sub> O <sub>4</sub>	.....	-26 (c 0.6, C <sub>2</sub> H <sub>5</sub> OH)	9,10
3	D-Glyceraldehyde, 3,3-bis(C-hydroxymethyl)- (Apiose)	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	.....	+5.6 (c 10) [150°] (syrup)	101,114
4	β-D-Arabinose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	155	-175 → -103	38,46,97, 99
5	D-Arabinose, 2-O-methyl-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	Syrup	-102	38,73
6	α-L-Arabinose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	158 amorphous	+55.4 → +105	112
7	β-L-Arabinose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	160	+190.6 → +104.5	81
8	D-L-Arabinose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	163.5-164.5	None	98,121
9	α-L-Lyxose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	105	+5.8 → +13.5	1
10	L-Lyxose, 5-deoxy-3-C-formyl- (Streptose)	C <sub>6</sub> H <sub>10</sub> O <sub>5</sub>	.....	.....	61
11	L-Lyxose, 3-C-formyl- (Hydroxystreptose)	C <sub>6</sub> H <sub>10</sub> O <sub>6</sub>	.....	.....	108
12	Pentose, 4,5-anhydro-5-deoxy-D-erythro-	C <sub>5</sub> H <sub>8</sub> O <sub>3</sub>	.....	.....	47
13	Pentose, 2-deoxy-D-erythro-	C <sub>5</sub> H <sub>10</sub> O <sub>4</sub>	96-98	-91 → -58	24
14	D-Ribose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	87	-23.1 → -23.7	87
15	D-Ribose, 2-C-hydroxymethyl- (Hamamelose)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	.....	-7.1 [λ 578]	33
16	α-D-Xylose	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	145	+93.6 → +18.8	50,52
17	D-Xylose, 5-deoxy-	C <sub>5</sub> H <sub>10</sub> O <sub>4</sub>	.....	+16	36
18	β-D-Xylose, 2-O-methyl-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	137-138	-21 → +34	64,65,73
19	α-D-Xylose, 3-O-methyl-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	95	+45 → +19	5,64,65
20	D-Allose, 6-deoxy-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	140-143 146-148	+1.6 [18°] (c 0.6) -4.7 → 0	57 25
21	D-Allose, 6-deoxy-2,3-di-O-methyl- (Mycinose)	C <sub>8</sub> H <sub>16</sub> O <sub>5</sub>	102-106	-46 → -29	25
22	Amicetose (a trideoxy hexose)	C <sub>6</sub> H <sub>12</sub> O <sub>3</sub>	Oil, b.p. 65-70	+28.6 (CHCl <sub>3</sub> )	106
23	Antiarose	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	.....	Levo	60
24	α-D-Galactose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	167	+150.7 → +80.2	96
25	β-D-Galactose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	143-145	+52.8 → +80.2	96,124
26	D-Galactose, 3,6-anhydro-	C <sub>6</sub> H <sub>10</sub> O <sub>5</sub>	.....	+21.3 [100°]	4,85
27	α-D-Galactose, 6-deoxy- (D-Fucose; Rhodeose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	140-145	+127 → +76.3 (c 10)	116
28	D-Galactose, 6-deoxy-3-O-methyl- (Digitalose)	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	106 <sup>1</sup> , 119 <sup>2</sup>	+106	66
29	D-Galactose, 6-deoxy-4-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	131-132	+82	34
30	D-Galactose, 6-deoxy-2,3-di-O-methyl-	C <sub>8</sub> H <sub>16</sub> O <sub>5</sub>	.....	+73	58,102
31	α-D-Galactose, 3-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	144-147	+150.6 → +108.6	89
32	α-D-Galactose, 6-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	122-123	+117 → +77.3	40,83
33	L-Galactose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	.....	See D-Galactose	
34	α-L-Galactose, 3,6-anhydro-	C <sub>6</sub> H <sub>10</sub> O <sub>5</sub>	.....	-39.4 → -25.2	3
35	α-L-Galactose, 6-deoxy- (L-Fucose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	145	-124.1 → -76.4	77
36	L-Galactose, 6-deoxy-2-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	149-150	-75±4 (c 0.5)	2
37	L-Galactose, 6-sulfate	C <sub>6</sub> H <sub>12</sub> O <sub>9</sub> S	.....	-47 (c 0.2) (Na salt)	111
38	D-L-Galactose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	143-144, 163	None (racemic)	82,115
39	α-D-Glucose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	146, 83 (H <sub>2</sub> O)	+112 → +52.7	7
40	β-D-Glucose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	148-150	+18.7 → +52.7	7
41	D-Glucose, 6-acetate	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	135	+48	26
42	D-Glucose, 2,3-di-O-methyl-	C <sub>8</sub> H <sub>16</sub> O <sub>6</sub>	85-86, 121	+50	22,58,119

/1/ Original melting point. /2/ Melting point after four-months' storage.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part I. NATURAL MONOSACCHARIDES: ALDOSES AND KETOSES**

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
	(A)	(B)	(C)	(D)	(E)
<b>Aldoses</b>					
43	D-Glucose, 6-O-benzoyl- (Vaccinin)	C <sub>13</sub> H <sub>16</sub> O <sub>7</sub>	Amorphous	+48 (C <sub>2</sub> H <sub>5</sub> OH)	84
44	α-D-Glucose, 6-deoxy- (Chinovose; Epirhamnose; Glucomethyllose; Isorhamnose; Isorhodoose; Quinovose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	139-140	+73.3 → +29.7 (c 8)	88
45	α-D-Glucose, 6-deoxy-3-O-methyl- (D-Thevetose)	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	116	+84 → +33	32
46	D-Glucose, 6-sulfonic acid, 6-deoxy- (6-Sulfoquinovose)	C <sub>6</sub> H <sub>12</sub> O <sub>8</sub> S	173-174	+87 <sup>a</sup>	80
47	D-Glucose, 3-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	162-167	+98 → +59.5	19
48	α-L-Glucose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	141-143	-95.5 → -51.4	27
49	L-Glucose, 6-deoxy-3-O-methyl- (L-Thevetose)	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	126-129	-36.9±2	15
50	D-Gulose, 6-deoxy-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	.....	.....	30,31,70
51	Hexose, 2,6-dideoxy-D-arabino- <sup>4</sup>	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	148	+46.6 [18°]	9,10
52	Hexose, 2,6-dideoxy-3-O-methyl-D-arabino- (D-Oleandrose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	.....	-11	110
53	Hexose, 3,6-dideoxy-D-arabino- (Tyvelose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	.....	+24±2	30
54	Hexose, 2,6-dideoxy-3-O-methyl-L-arabino- (L-Oleandrose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	62-63	+11.9±2.5	16
55	Hexose, 3,6-dideoxy-L-arabino- (Ascarylose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	.....	-24±2	30
56	Hexose, 2,6-dideoxy-3-O-methyl-D-lyxo- (Diginose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	90-92	+56±4	105,109
57	Hexose, 2,6-dideoxy-L-lyxo- (L-Fucose, 2-deoxy-)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	103-106	-61.6	53,125
58	Hexose, 2,6-dideoxy-3-O-methyl-L-lyxo-	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	78-85	-65	93
59	Hexose, 2,6-dideoxy-D-ribo- (Digitoxose; D-Altrose, 2,6-dideoxy-)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	110	+46.4	59,76
60	Hexose, 2,6-dideoxy-3-O-methyl-D-ribo- (Cymarose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	93	+52	54
61	Hexose, 3,6-dideoxy-D-ribo- (Paratose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	.....	+10±2(c 0.9)	23,31
62	Hexose, 4,6-dideoxy-3-O-methyl-D-ribo- (D-Gulose, 4,6-dideoxy-3-O-methyl-; Chalcose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	96-99	+120 → +76	25
63	Hexose, 2,6-dideoxy-D-xylo- (Boivinose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	96-98	-3.9 → +3.9	17
64	Hexose, 2,6-dideoxy-3-O-methyl-D-xylo- (Sarmen-tose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	78-79	+12 → +15.8	55
65	Hexose, 3,6-dideoxy-D-xylo- (Abequose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	.....	-3.2±0.6	118
66	Hexose, 2,6-dideoxy-3-C-methyl-L-xylo- (Mycarose)	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub>	128-129	-31.1	90
67	Hexose, 2,6-dideoxy-3-C-methyl-3-O-methyl-L-xylo- (Cladinose)	C <sub>8</sub> H <sub>16</sub> O <sub>4</sub>	oil, b.p. 120-132 (0.25 mm)	-23.1	29
68	Hexose, 3,6-dideoxy-L-xylo- (Colitose)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	.....	+4 (H <sub>2</sub> O); -51±2 (CH <sub>3</sub> OH)	72
69	D-Idose <sup>5</sup>	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	.....	.....	41
70	L-Idose, 1,6-anhydro-	C <sub>6</sub> H <sub>10</sub> O <sub>5</sub>	.....	.....	6
71	α-D-Mannose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	133	+29.3 → +14.5	68,69
72	β-D-Mannose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	132	-16.3 → +14.5	95
73	D-Mannose, 6-deoxy- (D-Rhamnose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	86-90	-7.0	75
74	α-L-Mannose, 6-deoxy-monohydrate (L-Rhamnose)	C <sub>6</sub> H <sub>14</sub> O <sub>6</sub>	93-94	-8.6 → +8.2	8,52
75	β-L-Mannose, 6-deoxy-	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	123-125	+38.4 → +8.9	78
76	L-Mannose, 6-deoxy-2-O-methyl-	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	.....	.....	24
77	L-Mannose, 6-deoxy-3-O-methyl- (L-Acofriose)	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	114-115	+30 [18°]	45
78	L-Mannose, 6-deoxy-2,4-di-O-methyl-	C <sub>8</sub> H <sub>16</sub> O <sub>5</sub>	82	-19 [16°]	20,74
79	L-Mannose, 6-deoxy-5-C-methyl-4-O-methyl- (Noviose)	C <sub>8</sub> H <sub>16</sub> O <sub>5</sub>	128-130	+19.9 (50% C <sub>2</sub> H <sub>5</sub> OH)	44
80	Rhodinose (a 2,3,6-trideoxyhexose)	C <sub>6</sub> H <sub>12</sub> O <sub>3</sub>	.....	-11±1.6	18
81	D-Talose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	128-132	+16.9	120
82	D-Talose, 6-deoxy- (D-Talomethyllose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	129-131	+20.6	75
83	L-Talose, 6-deoxy- (L-Talomethyllose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	116-118	-19.5±2 [18°]	103
84	L-Talose, 6-deoxy-2-O-methyl- (L-Acovenose)	C <sub>7</sub> H <sub>14</sub> O <sub>5</sub>	.....	-19.4	113
85	Heptose, D-glycero-D-galacto-	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	139-140	+47 → +64 (c 0.5)	104
86	Heptose, D-glycero-D-manno-	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	.....	.....	86,94
87	Heptose, D-glycero-L-manno-	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	.....	.....	78,79
<b>Ketoses</b>					
88	Dihydroxyacetone	C <sub>3</sub> H <sub>6</sub> O <sub>3</sub>	80 (dimer)	None	42

/a/ As a methyl glycoside cyclo-hexylamine salt. /4/ Included because of speculations concerning it in biological processes. /5/ Either D-idose or L-altrose is in the polysaccharide varianose.

*continued*



**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part I. NATURAL MONOSACCHARIDES: ALDOSES AND KETOSES**

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
	(A)	(B)	(C)	(D)	(E)
<b>Ketoses</b>					
89	Tetralose, <i>L-glycero</i> - <sup>6</sup> ( <i>L</i> -Erythrulose; Ketoerythritol; <i>L</i> -Threulose)	C <sub>4</sub> H <sub>8</sub> O <sub>4</sub>	Syrup	+12	11,12
90	Pentulose, <i>D-erythro</i> - (Adonose; <i>D</i> -Ribulose)	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	Syrup	+16.6 [27°]	71
91	Pentulose, <i>L-erythro</i> - ( <i>L</i> -Ribulose)	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	.....	-16.6	92
92	Pentulose, <i>D-threo</i> - ( <i>D</i> -Xylulose)	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	.....	-33	43
93	Pentulose, 5-deoxy- <i>D-threo</i> -	C <sub>5</sub> H <sub>10</sub> O <sub>4</sub>	.....	-5±1 (CH <sub>3</sub> OH)	36
94	Pentulose, <i>L-threo</i> - ( <i>L</i> -Xylulose; <i>L</i> -Lyxulose; Xyloketose)	C <sub>5</sub> H <sub>10</sub> O <sub>5</sub>	Syrup	+33.1	67
95	Hexulose, $\beta$ - <i>D-arabino</i> - ( $\beta$ - <i>D</i> -Fructose; Levulose)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	102-104 <sup>7</sup>	-133.5 → -92	49,51,126, 127
96	Hexulose, 6-deoxy- <i>D-arabino</i> - ( <i>D</i> -Rhamnulose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	.....	-13±2	48
97	Hexulose, <i>D-lyxo</i> - ( <i>D</i> -Tagatose)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	131-132	+2.7 → -4, -5	91
98	5-Hexulose, <i>D-lyxo</i> -	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	158	-86.6	117
99	Hexulose, 6-deoxy- <i>L-lyxo</i> - ( <i>L</i> -Fuculose)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	.....	.....	37
100	Hexulose, <i>D-ribo</i> - ( <i>D</i> -Psicose)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	Amorphous	+4.7	14,123
101	Hexulose, <i>L-xylo</i> - ( <i>L</i> -Sorbose)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	159-161	-43.1	100
102	Hexulose, 6-deoxy- <i>L-xylo</i> -	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	88	-25±2 ( <i>c</i> 0.7)	48
103	Heptulose, <i>D-altro</i> - (Sedoheptulose; Sedoheptose)	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	Amorphous	+2.5 ( <i>c</i> 10)	63
104	Heptulose-hemihydrate, <i>L-galacto</i> - (Perseulose)	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub> · 1/2 H <sub>2</sub> O	110-115	-90 → -80	13,39
105	Heptulose, <i>L-gulo</i> -	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	.....	-28	107
106	Heptulose, <i>D-ido</i> -	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	172	-34±8 ( <i>c</i> 0.3)	35
107	Heptulose, <i>D-manno</i> - (Mannoketoheptose; <i>D</i> -Mannotagatoheptose)	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	152	+29.4	62
108	Heptulose, <i>D-talo</i> -	C <sub>7</sub> H <sub>14</sub> O <sub>7</sub>	.....	.....	21
109	Octulose, <i>D-glycero-L-galacto</i> -	C <sub>8</sub> H <sub>16</sub> O <sub>8</sub>	.....	-57, -43.4 → -13.4	56,104
110	Octulose, <i>D-glycero-D-manno</i> -	C <sub>8</sub> H <sub>16</sub> O <sub>8</sub>	.....	+20 (CH <sub>3</sub> OH)	21

/6/ Early literature refers to this as *D-erythrose*. /7/ The ·1/2 H<sub>2</sub>O and ·2H<sub>2</sub>O forms also exist.

*Contributors:* Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

*References:* [1] Alberda van Ekenstein, W., and J. J. Blanksma. 1914. Chem. Weekblad 11:189. [2] Anderson, J. D., P. Andrews, and L. Hough. 1957. Chem. Ind. (London), p. 1453. [3] Araki, C., and S. Hirase. 1953. Bull. Chem. Soc. Japan 26:463. [4] Araki, C., and S. Hirase. 1956. Ibid. 25:770. [5] Aspinall, G. O., and J. E. McKay. 1958. J. Chem. Soc., p. 1059. [6] Baggett, N., P. J. Stoffyn, and R. W. Jeanloz. 1963. J. Org. Chem. 28:1041. [7] Bates, F. J., et al. 1942. Natl. Bur. Std. (U.S.) Circ. 728. [8] Berend, L. 1878. Ber. Deut. Chem. Ges. 11:1353. [9] Bergmann, M., et al. 1921. Ibid. 54:454. [10] Bergmann, M., et al. 1922. Ibid. 55:158. [11] Bertrand, G. 1900. Compt. Rend. 130:1330. [12] Bertrand, G. 1904. Bull. Soc. Chim. France, Ser. 3, 23:681. [13] Bertrand, G. 1909. Ibid., Ser. 4, 5:629. [14] Binkley, W. W., and M. L. Wolfrom. 1948. J. Am. Chem. Soc. 70:3940. [15] Blindenbacher, F., and T. Reichstein. 1948. Helv. Chim. Acta 31:1669. [16] Blindenbacher, F., and T. Reichstein. 1948. Ibid. 31:2061. [17] Bolliger, H. R., and T. Reichstein. 1953. Ibid. 36:302. [18] Brockmann, H., et al. 1963. Naturwissenschaften 50:43. [19] Chanley, J. D., et al. 1959. J. Am. Chem. Soc. 81:5180. [20] Charalambous, G., and E. E. Percival. 1954. J. Chem. Soc., p. 2443. [21] Charlson, A. J., and N. K. Richtmyer. 1960. J. Am. Chem. Soc. 82:3428. [22] Christensen, G. M., and F. Smith. 1957. Ibid. 79:4492. [23] Davies, D. A. L., et al. 1958. Nature 181:822. [24] Deriaz, R. E., et al. 1949. J. Chem. Soc., p. 1879. [25] Dion, H. W., P. W. K. Woo, and Q. R. Bartz. 1962. J. Am. Chem. Soc. 84:380, 1512. [26] Duff, R. B. 1957. J. Chem. Soc., p. 4730. [27] Fischer, E. 1890. Ber. Deut. Chem. Ges. 23:2618. [28] Fischer, E. 1896. Ibid. 29:324. [29] Flynn, E. H., et al. 1954. J. Am. Chem. Soc. 76:3121. [30] Fouquey, C., et al. 1958. Compt. Rend. 246:2417. [31] Fouquey, C., et al. 1958. Nature 182:944. [32] Frerejacque, M. 1950. Compt. Rend. 230:127. [33] Freudenberg, K., and F. Blümmel. 1924. Ann. Chem. 440:45. [34] Galmarini, O., and V. Deulofeu. 1961. Tetrahedron 15:76. [35] Gorin, P. A. J., et al. 1953. J. Chem. Soc., p. 1537.

*continued*



**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part I. NATURAL MONOSACCHARIDES: ALDOSES AND KETOSES**

- [36] Gorin, P. A. J., et al. 1953. *Ibid.*, p. 2140. [37] Green, M., and S. S. Cohen. 1956. *J. Biol. Chem.* 219:557. [38] Halliburton, G. J., and R. J. McIlroy. 1949. *J. Chem. Soc.*, p. 299. [39] Hann, R. M., and C. S. Hudson. 1939. *J. Am. Chem. Soc.* 61:336. [40] Hassid, W. Z., and J. Su. 1962. *Biochemistry* 1:468. [41] Haworth, W. N., H. Raistrick, and M. Stacey. 1935. *Biochem. J.* 29:2668. [42] Heilbron, J. M., and H. M. Bunbury. 1943. *Dictionary of organic compounds.* Oxford Univ. Press, New York. v. 1, p. 813. [43] Hickman, J., and G. Ashwell. 1956. *J. Am. Chem. Soc.* 78:6209. [44] Hinman, J. W., E. L. Caron, and H. Hocksema. 1957. *Ibid.* 79:3789. [45] Hirst, E. L., E. E. Percival, and R. S. Williams. 1958. *J. Chem. Soc.*, p. 1942. [46] Hockett, R., and C. S. Hudson. 1934. *J. Am. Chem. Soc.* 56:1632. [47] Hogenkamp, H. P. C., and H. A. Barker. 1961. *J. Biol. Chem.* 236:3097. [48] Hough, L., and J. K. N. Jones. 1952. *J. Chem. Soc.*, p. 4052. [49] Hudson, C. S., et al. 1916. *J. Am. Chem. Soc.* 38:1216. [50] Hudson, C. S., et al. 1917. *Ibid.* 39:1013. [51] Hudson, C. S., et al. 1917. *Ibid.* 39:1025. [52] Isbell, H. S., and W. W. Pigman. 1937. *Natl. Bur. Std. (U.S.) J. Res.* 18:141. [53] Iselin, B., and T. Reichstein. 1944. *Helv. Chim. Acta* 27:1200. [54] Jacobs, W. A., et al. 1930. *J. Biol. Chem.* 88:519. [55] Jacobs, W. A., et al. 1932. *Ibid.* 96:355. [56] Jones, J. K. N., et al. 1960. *Can. J. Chem.* 38:753. [57] Keller, M., and T. Reichstein. 1949. *Helv. Chim. Acta* 32:1607. [58] Khare, M. P., O. Schindler, and T. Reichstein. 1962. *Ibid.* 45:1534. [59] Kiliani, H. 1896. *Arch. Pharm.* 234:486. [60] Kiliani, H. 1913. *Ber. Deut. Chem. Ges.* 46:667. [61] Kuehl, F. A., Jr., et al. 1946. *J. Am. Chem. Soc.* 68:2679. [62] LaForge, F. B., et al. 1917. *J. Biol. Chem.* 28:511. [63] LaForge, F. B., et al. 1917. *Ibid.* 30:61. [64] Laidlaw, R. A., et al. 1950. *J. Chem. Soc.*, p. 528. [65] Laidlaw, R. A., et al. 1954. *Ibid.*, p. 752. [66] Lamb, I. D., and S. Smith. 1936. *Ibid.*, p. 442. [67] Levene, P. A., et al. 1914. *J. Biol. Chem.* 18:319. [68] Levene, P. A., et al. 1923. *Ibid.* 57:329. [69] Levene, P. A., et al. 1924. *Ibid.* 59:129. [70] Levene, P. A., et al. 1935. *Ibid.* 111:335. [71] Levene, P. A., et al. 1936. *Ibid.* 115:731. [72] Luderitz, O., et al. 1958. *Biochem. Z.* 330:193. [73] Lynch, D. L., H. O. Olney, and L. M. Wright. 1958. *J. Sci. Food Agr.* 9:56. [74] MacLennan, A. P., et al. 1960. *Biochem. J.* 74:3p. [75] Markovitz, A. 1962. *J. Biol. Chem.* 237:1767. [76] Micheel, F. 1930. *Ber. Deut. Chem. Ges.* 63:347. [77] Minsaa, J. 1933. *Rec. Trav. Chim.* 50:424. [78] Missale, G., A. Colajacomo, and I. Bologna. 1960. *Bull. Soc. Ital. Biol. Sper.* 36:1885. [79] Missale, G., A. Colajacomo, and I. Bologna. 1961. *Chem. Abstr.* 55:24869. [80] Miyano, M., and A. A. Benson. 1962. *J. Am. Chem. Soc.* 84:59. [81] Montgomery, E. M., and C. S. Hudson. 1934. *Ibid.* 56:2074. [82] Neuberg, C., and J. Wohlgenuth. 1902. *Z. Physiol. Chem.* 36:224. [83] Nunn, J. R., and M. M. von Holdt. 1957. *J. Chem. Soc.*, p. 1094. [84] Ohle, H. 1922. *Biochem. Z.* 131:611. [85] O'Neill, A. N. 1955. *J. Am. Chem. Soc.* 77:2837. [86] Palleroni, N. J., and M. Doudoroff. 1956. *J. Biol. Chem.* 218:535. [87] Phelps, F. P., H. S. Isbell, and W. W. Pigman. 1934. *J. Am. Chem. Soc.* 56:747. [88] Pigman, W. W., and R. M. Goepf, Jr. 1948. *Chemistry of the carbohydrates.* Academic Press, New York. p. 106. [89] Reber, F., and T. Reichstein. 1945. *Helv. Chim. Acta* 28:1164. [90] Regna, P. P., et al. 1953. *J. Am. Chem. Soc.* 75:4625. [91] Reichstein, T., and W. Bossard. 1934. *Helv. Chim. Acta* 17:753. [92] Reichstein, T., and W. Bossard. 1934. *Ibid.* 17:996. [93] Renkonen, O., O. Schindler, and T. Reichstein. 1959. *Ibid.* 42:182. [94] Richtmyer, N. K., and A. J. Charlson. 1960. *J. Am. Chem. Soc.* 82:3428. [95] Rüber, C. N., et al. 1927. *Ber. Deut. Chem. Ges.* 60:2402. [96] Rüber, C. N., et al. 1929. *J. Chem. Soc.*, p. 2173. [97] Ruff, O. 1898. *Ber. Deut. Chem. Ges.* 31:1576. [98] Ruff, O. 1899. *Ibid.* 32:550. [99] Ruff, O. 1899. *Ibid.* 32:554. [100] Schlubach, H. H., and J. Vorwerk. 1933. *Ibid.* 66:1251. [101] Schmidt, O. T. 1930. *Ann. Chem.* 483:115. [102] Schmidt, O. T. 1944. *Ibid.* 556:179. [103] Schmitz, J. 1948. *Helv. Chim. Acta* 31:1719. [104] Sephton, H. H., and N. K. Richtmyer. 1963. *J. Org. Chem.* 28:1691. [105] Shoppe, C. W., and T. Reichstein. 1942. *Helv. Chim. Acta* 25:1611. [106] Stevens, C. L., et al. 1962. *J. Org. Chem.* 27:2991. [107] Stewart, L. C., N. K. Richtmyer, and C. S. Hudson. 1952. *J. Am. Chem. Soc.* 74:2206. [108] Stodola, F. H. 1951. *Ibid.* 73:5912. [109] Tamm, C., and T. Reichstein. 1948. *Helv. Chim. Acta* 31:1630. [110] Tsechesche, R., and G. Buschauer. 1957. *Ann. Chem.* 603:59. [111] Turvey, J. R., and D. A. Rees. 1961. *Nature* 189:831. [112] Vogel, H. 1928. *Helv. Chim. Acta* 11:1210. [113] Von Euw, J., and T. Reichstein. 1950. *Ibid.* 33:485. [114] Vongerichten, E.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part I. NATURAL MONOSACCHARIDES: ALDOSES AND KETOSES**

1902. Ann. Chem. 321:71. [115] Von Lippmann, E. O. 1922. Ber. Deut. Chem. Ges. 55:3038. [116] Votocek, E., and V. Valentin. 1930. Chem. Zentr. 2:2543. [117] Weidenhagen, R., and G. Bernsee. 1960. Angew. Chem. 72:109. [118] Westphal, O., et al. 1959. Ann. Chem. 620:8. [119] White, E. V., and P. S. Rao. 1953. J. Am. Chem. Soc. 75:2617. [120] Wiley, P. F., and M. V. Sigal, Jr. 1958. Ibid. 80:1010. [121] Wohl, A., and F. Momber. 1893. Ber. Deut. Chem. Ges. 26:742. [122] Wohl, A., and F. Momber. 1917. Ibid. 50:456. [123] Wolfrom, M. L., et al. 1945. J. Am. Chem. Soc. 67:1793. [124] Wolfrom, M. L., et al. 1954. Ibid. 76:1198. [125] Wyss, E., H. Jäger, and O. Schindler. 1960. Helv. Chim. Acta 43:664. [126] Young, F. E., F. T. Jones, and O. R. Black. 1952. J. Am. Chem. Soc. 74:5798. [127] Young, F. E., F. T. Jones, and H. J. Lewis. 1952. J. Phys. Chem. 56:1093.

**Part II. NATURAL MONOSACCHARIDES: AMINO SUGARS**

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Reference
	(A)	(B)	(C)	(D)	(E)
<b>Aldosamines</b>					
1	D-Ribose, 3-amino-3-deoxy-	C <sub>5</sub> H <sub>11</sub> NO <sub>4</sub>	158-158.5 d.	-24.6 (hydrochloride)	24
2	D-Galactose, 2-amino-2-deoxy- (Galactosamine; Chondrosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	185	+121 → +80 (hydrochloride)	14
3	α-L-Galactose, 2-amino-2,6-dideoxy- (L-Fucosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>4</sub>	192-193 d.	-119 → -92 [27°] (hydrochloride)	1,13
4	α-D-Glucose, 2-amino-2-deoxy- (Glucosamine; Chitosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	88	+100 → +47.5	26
5	β-D-Glucose, 2-amino-2-deoxy-	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	110-111	+28 → +47.5	26
6	D-Glucose, 3-amino-3-deoxy- (Kanosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	128 d.	+19 [14°]	4,16,17
7	D-Glucose, 6-amino-6-deoxy-	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	161-162 d.	+23 → +50.1 (hydrochloride)	4,16,17
8	D-Glucose, 2,6-diamino-2,6-dideoxy- (Neosamine C)	C <sub>6</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	>230	+61.5 (dihydrochloride)	18,19
9	D-Glucose, 3,6-dideoxy-3-dimethylamino- (Myca- minose)	C <sub>8</sub> H <sub>17</sub> NO <sub>4</sub>	115-116	+31 (hydrochloride)	7
10	D-Glucose, 4,6-dideoxy-4-dimethylamino-	C <sub>8</sub> H <sub>17</sub> NO <sub>4</sub>	192-193	+45.5 (hydrochloride)	22
11	L-Glucose, 2-deoxy-2-methylamino-	C <sub>7</sub> H <sub>15</sub> NO <sub>5</sub>	130-132	-64	27
12	D-Gulose, 2-amino-1,6-anhydro-2-deoxy-	C <sub>6</sub> H <sub>11</sub> NO <sub>4</sub>	250-260 d.	+41±2 (hydrochloride)	8
13	D-Gulose, 2-amino-2-deoxy-	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	152-162 d.	+5.6 → -18.7 (hydrochloride)	23
14	Hexose, 3,4,6-trideoxy-3-dimethylamino-D-xylo- (Desosamine; Picrocine)	C <sub>8</sub> H <sub>17</sub> NO <sub>3</sub>	189-191 d.	+49.5 (c 10) (hydrochloride)	2
15	Hexose, a 4-acetamido-2-amino-2,4,6-trideoxy-	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	216-219	+115 → +94 [26°] (c 0.05)	21
16	Hexose, an amino-deoxy-3-O-carboxyethyl-	C <sub>9</sub> H <sub>17</sub> NO <sub>7</sub>	.....	.....	20
17	Hexose, a 2,6-diamino-2,6-dideoxy- (Neosamine B; Paramose)	C <sub>6</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	135-150 d.	+17.5 (c 0.9) (hydrochloride)	18,19
18	Hexose, a 3-dimethylamino-2,3,6-trideoxy- (Rhodosamine)	C <sub>8</sub> H <sub>17</sub> NO <sub>3</sub>	.....	.....	3
19	D-Mannose, 2-amino-2-deoxy- (Mannosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	142 d.	-4.3 (c 9) (hydrochloride)	12
20	D-Mannose, 3-amino-3,6-dideoxy- (Mycosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>4</sub>	162	-11.5 (hydrochloride)	25
21	D-Talose, 2-amino-2-deoxy- (Talosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	151-153	+3.4 → -5.7 (c 0.9) (hydrochloride)	5,11
22	L-Talose, 2-amino-2,6-dideoxy- (Pneumosamine)	C <sub>6</sub> H <sub>13</sub> NO <sub>4</sub>	162-163	+6.9 → +10.4 (hydrochloride)	1
<b>Ketosamines</b>					
23	Pentulose, 1-(o-carboxyanilino)-1-deoxy-D-erythro-	C <sub>12</sub> H <sub>14</sub> NO <sub>6</sub>	.....	.....	6,15
24	Hexulose, 1-(o-carboxyanilino)-1-deoxy-D-arabino-	C <sub>13</sub> H <sub>16</sub> NO <sub>7</sub>	.....	.....	15
25	Hexulose, 5-amino-5-deoxy-L-xylo-	C <sub>6</sub> H <sub>13</sub> NO <sub>5</sub>	174-176	-62	9
26	Hexulose, 6-deoxy-6-(N-methylacetamido)-L-xylo-	C <sub>9</sub> H <sub>17</sub> NO <sub>6</sub>	.....	.....	10

*continued*

## 98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS

### Part II. NATURAL MONOSACCHARIDES: AMINO SUGARS

*Contributors:* Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

*References:* [1] Barker, S. A., et al. 1961. *Nature* 189:303. [2] Bentley, H. R., K. G. Cunningham, and F. A. Spring. 1951. *J. Chem. Soc.*, p. 2301. [3] Brockmann, H., and T. Waehneltd. 1961. *Naturwissenschaften* 48:717. [4] Cron, M. J., et al. 1958. *J. Am. Chem. Soc.* 80:2342. [5] Crumpton, M. J. 1957. *Nature* 180:605. [6] Day, C. H., and F. W. E. Gibson. 1959. *Biochem. J.* 72:580. [7] Hochstein, F. A., and P. Regna. 1955. *J. Am. Chem. Soc.* 77:3353. [8] Jeanloz, R. W. 1959. *Ibid.* 81:1956. [9] Jones, J. K. N., et al. 1961. *Can. J. Chem.* 39:965. [10] Jones, J. K. N., et al. 1961. *Ibid.* 39:2400. [11] Kuhn, R., et al. 1957. *Ann. Chem.* 612:65. [12] Kuhn, R., et al. 1959. *Ibid.* 628:172. [13] Kuhn, R., et al. 1959. *Ibid.* 628:186. [14] Levene, P. A. 1916. *J. Biol. Chem.* 26:147. [15] Lingens, F., and H. Hellmann. 1960. *Ann. Chem.* 630:84. [16] Maeda, K., et al. 1958. *J. Antibiotics (Tokyo)*, A, 11:73. [17] Maeda, K., et al. 1960. *Chem. Abstr.* 53:20526. [18] Rinehart, K. L., et al. 1960. *J. Am. Chem. Soc.* 82:3938. [19] Rinehart, K. L., et al. 1961. *Ibid.* 83:643, 2964. [20] Salton, M. R. J. 1957. *Nature* 180:338. [21] Sharon, C. W., and R. W. Jeanloz. 1960. *J. Biol. Chem.* 235:1. [22] Stevens, C. L., et al. 1963. *J. Am. Chem. Soc.* 85:1552. [23] Van Tamelen, E. E., et al. 1956. *Ibid.* 78:4817. [24] Waller, C. W., et al. 1953. *Ibid.* 75:2025. [25] Walters, D. R., J. D. Dutcher, and O. Wintersteiner. 1963. *J. Org. Chem.* 28:995. [26] Westphal, O., and H. Holzmann. 1942. *Ber. Deut. Chem. Ges.* 75:1274. [27] Wolfrom, M. L., and A. Thompson. 1947. *J. Am. Chem. Soc.* 69:1847.

### Part III. NATURAL ALDITOLS AND INOSITOLS (with Inososes and Inosamines)

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
	(A)	(B)	(C)	(D)	(E)
Alditols					
1	Glycerol	C <sub>3</sub> H <sub>8</sub> O <sub>3</sub>	20	None	24
2	Glycerol, 1-deoxy- (1,2-Propane-diol) <sup>1</sup>	C <sub>3</sub> H <sub>8</sub> O <sub>2</sub>	Oil, b.p. 188-189	None (racemic)	25
3	Erythritol	C <sub>4</sub> H <sub>10</sub> O <sub>4</sub>	118-120	None (meso)	9
4	Erythritol, 1,4-dideoxy- (2,3-Butyleneglycol)	C <sub>4</sub> H <sub>10</sub> O <sub>2</sub>	25, 34	None (meso)	57,59
5	D-Threitol, 1,4-dideoxy-	C <sub>4</sub> H <sub>10</sub> O <sub>2</sub>	19	-13.0	57
6	L-Threitol, 1,4-dideoxy-	C <sub>4</sub> H <sub>10</sub> O <sub>2</sub>	.....	+10.2	21
7	D,L-Threitol, 1,4-dideoxy-	C <sub>4</sub> H <sub>10</sub> O <sub>2</sub>	7.6	None (racemic)	59
8	D-Arabinitol	C <sub>5</sub> H <sub>12</sub> O <sub>5</sub>	103	+7.82 (c 8, borax solution)	6
9	L-Arabinitol	C <sub>5</sub> H <sub>12</sub> O <sub>5</sub>	101-102	-32 (c 0.4, 5% molybdate)	7,47,54
10	Ribitol (Adomitol)	C <sub>5</sub> H <sub>12</sub> O <sub>5</sub>	102	None (meso)	57
11	Galactitol (Dulcitol)	C <sub>6</sub> H <sub>14</sub> O <sub>6</sub>	186-188	None (meso)	48
12	D-Glucitol (Sorbitol)	C <sub>6</sub> H <sub>14</sub> O <sub>6</sub>	112	-1.8 [15°]	56
13	D-Glucitol, 1,5-anhydro- (Polygalitol)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	140-141	+42.4	46
14	L-Iditol	C <sub>6</sub> H <sub>14</sub> O <sub>6</sub>	73.5	-3.5 (c 10)	10
15	D-Mannitol	C <sub>6</sub> H <sub>14</sub> O <sub>6</sub>	166	-0.21	13
16	D-Mannitol, 1,5-anhydro- (Styracitol)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	157	-49.9	60
17	Heptitol, D-glycero-D-galacto- (Heptitol, L-glycero-D-manno-; Perseitol)	C <sub>7</sub> H <sub>16</sub> O <sub>7</sub>	183-185, 188	-1.1	28,36
18	Heptitol, D-glycero-D-gluco- (Heptitol, L-glycero-D-talo-; β-Sedoheptitol)	C <sub>7</sub> H <sub>16</sub> O <sub>7</sub>	131-132	+46 (5% NH <sub>4</sub> molybdate)	14
19	Heptitol, D-glycero-D-manno- (Heptitol, D-glycero-D-talo-; Volemitol)	C <sub>7</sub> H <sub>16</sub> O <sub>7</sub>	153	+2.65	12
20	Octitol, D-erythro-D-galacto-	C <sub>8</sub> H <sub>18</sub> O <sub>8</sub> ·H <sub>2</sub> O	169-170	-11 (5% NH <sub>4</sub> molybdate)	14
Inositols					
21	Betitol (a dideoxy inositol)	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub>	224	.....	55
22	Bioinosose (scyllo-Inosose; myo-Inosose-2; a deoxy keto inositol)	C <sub>6</sub> H <sub>10</sub> O <sub>6</sub>	198-200	None (meso)	29,43

<sup>1</sup>/ The 1-phosphate ester of this diol is said to occur in brain tissue and sea-urchin eggs [33].

*continued*



**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part III. NATURAL ALDITOLS AND INOSITOLS (with Inososes and Inosamines)**

Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
(A)	(B)	(C)	(D)	(E)
<b>Inositols</b>				
23 <i>d</i> -Bornesitol (a <i>myo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	200	+31.6	20,22
24 <i>l</i> -Bornesitol (a <i>myo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	205-206	-32.1	11
25 Conduritol (a 2,3-dehydro-2,3-dideoxyinositol)	C <sub>6</sub> H <sub>10</sub> O <sub>4</sub>	142-143	None (meso)	30
26 Cordycepic acid (a tetrahydroxycyclohexanecarboxylic acid) <sup>2</sup>	C <sub>7</sub> H <sub>12</sub> O <sub>6</sub>	.....	.....	15
27 Dambonitol (a <i>myo</i> -inositol dimethyl ether)	C <sub>8</sub> H <sub>16</sub> O <sub>6</sub>	206	None (meso)	17
28 DL-Inositol	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	253	None (racemic)	53
29 <i>d</i> -Inositol	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	.....	+60	8
30 <i>l</i> -Inositol	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	240	-65	51
31 Laminitol (a <i>C</i> -methyl <i>myo</i> -inositol)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	266-269	-3	32
32 Liriodendritol (a <i>myo</i> -inositol dimethyl ether)	C <sub>8</sub> H <sub>16</sub> O <sub>6</sub>	224	-25	40
33 <i>muco</i> -Inositol monomethyl ether	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	322-325	.....	3
34 <i>myo</i> -Inositol ( <i>meso</i> -Inositol)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	217-218	None (meso)	35
35 <i>d</i> - <i>myo</i> -Inosose-1 (a deoxy keto inositol)	C <sub>6</sub> H <sub>10</sub> O <sub>6</sub>	138-139	+19.6	34
36 Mytilitol (a <i>C</i> -methyl <i>scyllo</i> -inositol)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	259	None (meso)	1
37 <i>neo</i> -Inosamine-2 (a deoxy amino inositol)	C <sub>6</sub> H <sub>13</sub> O <sub>5</sub> N	239-241 d.	None (meso)	4
38 <i>d</i> -Ononitol (a <i>myo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	172	+6.6	41
39 <i>d</i> -Pinitol (a <i>dextro</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	186	+65.5	37
40 <i>l</i> -Pinitol (a <i>levo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	186	-65	5,42
41 <i>l</i> -Quebrachitol (a <i>levo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	190-191	-80.2 [28°]	2,16
42 <i>d</i> -Quercitol (a deoxy <i>dextro</i> -inositol)	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	235	+24.2	45
43 <i>d</i> -Quinic acid (a trideoxy carboxy <i>dextro</i> -inositol)	C <sub>7</sub> H <sub>12</sub> O <sub>6</sub>	164	+44 (c 10)	55
44 <i>l</i> -Quinic acid (a trideoxy carboxy <i>levo</i> -inositol)	C <sub>7</sub> H <sub>12</sub> O <sub>6</sub>	162	-42.1	23
45 Quinic acid, 5-dehydro-	C <sub>7</sub> H <sub>10</sub> O <sub>6</sub>	140-142 (138 s.)	-82.4 [28°]	58
46 Scyllitol ( <i>scyllo</i> -Inositol; Cocositol)	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	352-353	None (meso)	38,44
47 Sequoyitol (a <i>myo</i> -inositol monomethyl ether)	C <sub>7</sub> H <sub>14</sub> O <sub>6</sub>	234-235	None (meso)	50
48 Shikimic acid (a 3,4-anhydro-quinic acid)	C <sub>7</sub> H <sub>10</sub> O <sub>5</sub>	183-184	-200 [16°]	19
49 Shikimic acid, 5-dehydro-	C <sub>7</sub> H <sub>8</sub> O <sub>5</sub>	150-152	-57.5 [28°] (EtOH)	49
50 Streptamine (2,4-diaminodideoxyscyllitol)	C <sub>6</sub> H <sub>14</sub> O <sub>4</sub> N <sub>2</sub>	88, 210-250 d.	None (meso)	27
51 Streptamine, 2-deoxy-	C <sub>6</sub> H <sub>14</sub> O <sub>3</sub> N <sub>2</sub>	.....	None (meso)	18,31
52 Streptadine (1,3-Dideoxy-1,3-diguandino-scyllitol)	C <sub>8</sub> H <sub>18</sub> N <sub>6</sub> O <sub>4</sub>	.....	None (meso)	39
53 Viburnitol (a deoxy <i>levo</i> -inositol) <sup>3</sup>	C <sub>6</sub> H <sub>12</sub> O <sub>5</sub>	174	-73.9	26

/2/ Strong evidence that cordycepic acid is really D-mannitol [52]. /3/ Not an enantiomorph of *d*-quercitol; other isomeric relationship is involved.

**Contributors:** Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

**References:** [1] Ackermann, D. 1921. Ber. Deut. Chem. Ges. 54:1938. [2] Adams, R., D. C. Pease, and J. H. Clark. 1940. J. Am. Chem. Soc. 62:2194. [3] Adhikari, S. K., R. A. Bell, and W. E. Harvey. 1962. J. Chem. Soc., p. 2829. [4] Allen, G. R., Jr. 1956. J. Am. Chem. Soc. 78:5691. [5] Anderson, L., et al. 1958. Arch. Biochem. Biophys. 78:518. [6] Asahina, Y., and M. Yanagita. 1934. Ber. Deut. Chem. Ges. 67:799. [7] Ashida, K. 1944. J. Agr. Chem. Soc. Japan 20:621. [8] Ballou, C. E., and A. B. Anderson. 1953. J. Am. Chem. Soc. 75:648. [9] Bamberger, M., and A. Landsiedl. 1940. Monatsh. Chem. 21:571. [10] Bertrand, G. 1905. Bull. Soc. Chim. France, Ser. 3, 33:166. [11] Bien, S., and D. Ginsburg. 1958. J. Chem. Soc., p. 3189. [12] Bougault, J., and G. Allard. 1902. Compt. Rend. 135:796. [13] Braham, J. M. 1919. J. Am. Chem. Soc. 41:1707. [14] Charlson, A. J., and N. K. Richtmyer. 1960. Ibid. 82:3428. [15] Chatterjee, R., K. A. Srinivasan, and P. C. Maiti. 1957. J. Am. Pharm. Assoc. Sci. Ed. 46:114. [16] DeJong, A. W. K. 1906. Rec. Trav. Chim. 25:48. [17] DeJong, A. W. K. 1908. Ibid. 27:257. [18] Dutcher, J. D., and M. N. Donin. 1957. Ibid. 74:3420. [19] Eijkman, J. F. 1885. Ibid. 4:32 [20] Flint, E. R., and B. Tollens. 1892. Ann. Chem. 272:288. [21] Fulmer, E. I., L. A. Underkofler, and A. C. Bantz. 1943. J. Am. Chem. Soc. 65:1425. [22] Girard, A. 1871. Compt. Rend. 73:426. [23] Gorter, K. 1908. Ann. Chem. 359:221. [24] Heilbron, I. M., and H. M. Bunbury. 1943. Dictionary of organic compounds. Oxford Univ. Press, New York, v.2. [25] Heilbron, I. M., and H. M. Bunbury. 1943. Ibid. v.3. [26] Herissey, H., and G. Poirot. 1936. Compt. Rend. 203:466. [27] Holland, G. F., et al. 1958. J. Am.

*continued*



**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part III. NATURAL ALDITOLS AND INOSITOLS (with Inososes and Inosamines)**

Chem. Soc. 80:6031. [28] Jones, J. K. N., and R. A. Wall. 1961. Nature 189:746. [29] Kluyver, A. J., and A. G. J. Boezaardt. 1939. Rec. Trav. Chim. 58:958. [30] Kubler, K. 1909. Chem. Abstr. 3:1150. [31] Kuehl, F. A., Jr., M. N. Bishop, and K. Folkers. 1951. J. Am. Chem. Soc. 73:881. [32] Lindberg, B. 1946-47. Arkiv. Kemi Mineral. Geol. 23A(2). [33] Lindberg, B., and B. Wickberg. 1959. Arkiv.Kemi 13:447. [34] Magasanik, B., and E. Chargaff. 1948. J. Biol. Chem. 175:929. [35] Maquenne, M. 1887. Ann. Chim. Phys. (Paris), Ser. 6, 12:80. [36] Maquenne, M. 1890. Ibid., Ser. 6, 19:5. [37] Maquenne, M. 1891. Ibid., Ser. 6, 22:264. [38] Müller, H. 1907. J. Chem. Soc. 91:1767. [39] Peck, R. L., et al. 1946. J. Am. Chem. Soc. 68:776. [40] Plouvier, V. 1955. Compt. Rend. 241:765. [41] Plouvier, V. 1955. Ibid. 241:983. [42] Plouvier, V. 1956. Ibid. 243:1913. [43] Posternak, T. 1936. Helv. Chim. Acta 19:1333. [44] Posternak, T. 1942. Ibid. 25:746. [45] Prunier, L. 1878. Ann. Chim. Phys. (Paris), Ser. 5, 15:5. [46] Richtmyer, N. K., C. J. Carr, and C. S. Hudson. 1943. J. Am. Chem. Soc. 65:1477. [47] Richtmyer, N. K., and C. S. Hudson. 1951. Ibid. 73:2249. [48] Rogerson, H. 1912. J. Chem. Soc. 101:1040. [49] Salamon, I. I., and B. D. Davis. 1953. J. Am. Chem. Soc. 75:5567. [50] Sherrard, E. C., and E. F. Kurth. 1929. Ibid. 51:3139. [51] Smith, R. H. 1954. Biochem. J. 57:140. [52] Sprecher, M., and D. B. Sprinson. 1963. J. Org. Chem. 28:2490. [53] Tanret, C. 1907. Compt. Rend. 145:1196. [54] Touster, O., and S. O. Harwell. 1958. J. Biol. Chem. 230:1031. [55] Von Lippmann, E. O. 1901. Ber. Deut. Chem. Ges. 34:1159. [56] Von Lippmann, E. O. 1927. Ibid. 60:161. [57] Ward, G. E., et al. 1944. J. Am. Chem. Soc. 66:541. [58] Weiss, U., B. D. Davis, and E. S. Mingioli. 1953. Ibid. 75:5572. [59] Wilson, C. E., and H. J. Lucas. 1936. Ibid. 58:2396. [60] Zervas, L. 1930. Ber. Deut. Chem. Ges. 63:1689.

**Part IV. NATURAL ALDONIC, URONIC, AND ALDARIC ACIDS**

A number of "keto" acids, reported in the literature as stemming from biological systems, have not been included since they are still grossly undefined [4,27].

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Reference
	(A)	(B)	(C)	(D)	(E)
<b>Aldonic Acids</b>					
1	D-Glyceric acid	C <sub>3</sub> H <sub>6</sub> O <sub>4</sub>	Gum	Dextro	15
2	L-Glyceric acid	C <sub>3</sub> H <sub>6</sub> O <sub>4</sub>	Gum	Levo	15
3	D-Arabinonic acid	C <sub>5</sub> H <sub>10</sub> O <sub>6</sub>	114-116	+10.5 (c 6)	35
4	L-Arabinonic acid	C <sub>5</sub> H <sub>10</sub> O <sub>6</sub>	118-119	-9.6 → -41.7 <sup>1</sup>	32
5	L-Arabinonic-1, 4-lactone	C <sub>5</sub> H <sub>8</sub> O <sub>5</sub>	97-99	-72	5
6	D-Ribonic acid	C <sub>5</sub> H <sub>10</sub> O <sub>6</sub>	112-113	-17.0	22
7	D-Xylonic acid	C <sub>5</sub> H <sub>10</sub> O <sub>6</sub>	.....	-2.9 → +20.1 <sup>1</sup>	33
8	L-Xylonic acid	C <sub>5</sub> H <sub>10</sub> O <sub>6</sub>	.....	-91.8 <sup>1</sup>	20
9	D-Altronic acid	C <sub>6</sub> H <sub>12</sub> O <sub>7</sub>	.....	+11.5 → +24.8 <sup>1</sup> (Ca salt, N HCl)	19,34
10	D-Galactonic acid	C <sub>6</sub> H <sub>12</sub> O <sub>7</sub>	122	-11.2 → +57.6 <sup>1</sup>	21,30
11	D-Gluconic acid	C <sub>6</sub> H <sub>12</sub> O <sub>7</sub>	130-132 (110-112 s.)	-6.7 → +11.9 <sup>1</sup>	31
12	L-Gulonic acid	C <sub>6</sub> H <sub>12</sub> O <sub>7</sub>	Exists only in soln.	[ca. 0°]	8
13	Hexosonic acid, 2-deoxy-D-arabino-	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	93-95	+68 (lactone)	14,40
14	2-Hexulosonic acid, D-arabino-	C <sub>6</sub> H <sub>10</sub> O <sub>7</sub>	.....	-81.7 (Na salt)	26
15	2-Hexulosonic acid, 3-deoxy-D-erythro-	C <sub>6</sub> H <sub>10</sub> O <sub>6</sub>	.....	-29.2 (c 6, Ca salt)	25
16	2-Hexulosonic acid, D-lyxo-	C <sub>6</sub> H <sub>10</sub> O <sub>7</sub>	169	-5	12
17	5-Hexulosonic acid, D-arabino-	C <sub>6</sub> H <sub>10</sub> O <sub>7</sub>	108-109	.....	3
18	5-Hexulosonic acid, D-xylono-	C <sub>6</sub> H <sub>10</sub> O <sub>7</sub>	.....	-14.5	7
19	D-Mannonic acid	C <sub>6</sub> H <sub>12</sub> O <sub>7</sub>	.....	-15.6	23
20	D-Gluconic acid, O-β-D-galactopyranosyl-(1 → 4)- (Lactobionic acid)	C <sub>12</sub> H <sub>22</sub> O <sub>12</sub>	.....	+25.1 (Ca salt)	37
<b>Uronic Acids</b>					
21	L-Lyxuronic acid	C <sub>5</sub> H <sub>8</sub> O <sub>6</sub>	.....	.....	1,2

<sup>1</sup>/ Equilibrates with the lactone.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part IV. NATURAL ALDONIC, URONIC, AND ALDARIC ACIDS**

Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
(A)	(B)	(C)	(D)	(E)
Uronic Acids				
22 $\beta$ -D-Galacturonic acid	$C_6H_{10}O_7$	160	+27 $\rightarrow$ +55.6	11
23 $\alpha$ -D-Galacturonic acid·monohydrate	$C_6H_{12}O_8$	159-160 (110-115 s.)	+97.9 $\rightarrow$ +50.9	11
24 D-Galacturonic acid, 2-amino-2-deoxy-	$C_6H_{11}O_6N$	160 d.	+84.5 (pH 2 HCl)	17,18
25 $\beta$ -D-Glucuronic acid	$C_6H_{10}O_7$	156	+11.7 $\rightarrow$ +36.3	39
26 D-Glucuronic acid, 2-amino-2-deoxy-	$C_6H_{11}O_6N$	120-172 d.	+55	16,41
27 D-Glucuronic acid, 3-O-methyl-	$C_7H_{12}O_7$	Syrup	+6	10,24
28 L-Guluronic acid	$C_6H_{10}O_7$	.....	.....	6,13
29 L-Iduronic acid	$C_6H_{10}O_7$	.....	+30	9
30 $\beta$ -D-Mannuronic acid	$C_6H_{10}O_7$	165-167	-47.9 $\rightarrow$ -23.9	36
31 $\alpha$ -D-Mannuronic acid·monohydrate	$C_6H_{12}O_8$	110 s., 120-130 d.	+16 $\rightarrow$ -6.1 (c 6.8)	36
Aldaric Acids				
32 D-Tartaric acid	$C_4H_6O_6$	170	-15	28
33 L-Tartaric acid	$C_4H_6O_6$	170	+15 [15°]	38
34 L-Malic acid	$C_4H_6O_5$	100	-2.3 (c 8.4)	29

*Contributors:* Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

*References:* [1] Ameyama, M., and K. Kondo. 1958. Bull. Agr. Chem. Soc. Japan 22:271. [2] Ameyama, M., and K. Kondo. 1958. Chem. Abstr. 52:20408. [3] Ashwell, G., A. J. Wahba, and J. Hickman. 1960. J. Biol. Chem. 235:1559. [4] Ashwell, G., et al. 1963. Ibid. 238:1577. [5] Assarsson, A., B. Lindberg, and H. Borbrueygen. 1959. Acta Chem. Scand. 13:1395. [6] Bernhauer, K., and I. Irrgang. 1935. Biochem. Z. 280:360. [7] Boutroux, L. 1890. Ann. Chim. Phys. (Paris), Ser. 6, 21:565. [8] Burns, J. J. 1957. J. Am. Chem. Soc. 79:1257. [9] Cifonelli, J. A., J. Ludowieg, and A. Dorfman. 1958. J. Biol. Chem. 233:541. [10] Das Gupta, P. C., and P. B. Sarkar. 1954. Textile Res. J. 24:705, 1071. [11] Ehrlich, F., and F. Schubert. 1929. Ber. Deut. Chem. Ges. 62:1987. [12] Ettel, V., J. Liebster, and M. Tadra. 1952. Chem. Abstr. 46:7526. [13] Fischer, F. G., and H. Dörfel. 1955. Z. Physiol. Chem. 302:186. [14] Fischer, H. O. L., and G. Dangschat. 1937. Helv. Chim. Acta 20:705. [15] Frankland, P., and J. McGregor. 1893. J. Chem. Soc. 63:513. [16] Heyns, K., et al. 1955. Chem. Ber. 88:188. [17] Heyns, K., et al. 1957. Ibid. 90:2443. [18] Heyns, K., et al. 1959. Ibid. 92:2435. [19] Hickman, J., and G. Ashwell. 1960. J. Biol. Chem. 235:1566. [20] Kanfer, J., G. Ashwell, and J. J. Burns. 1960. Ibid. 235:2518. [21] Kiliani, H. 1922. Ber. Deut. Chem. Ges. 55:75. [22] Ladenburg, K., et al. 1944. J. Am. Chem. Soc. 66:1217. [23] Levene, P. A. 1924. J. Biol. Chem. 59:123. [24] Marsh, C. A. 1952. J. Chem. Soc., p. 1578. [25] Merrick, J. M., and S. Roseman. 1960. J. Biol. Chem. 235:1274. [26] Ohle, H., and R. Wolter. 1930. Ber. Deut. Chem. Ges. 63:843. [27] Palleroni, N. J., et al. 1956. J. Biol. Chem. 223:499. [28] Pasteur, L. 1850. Ann. Chim. Phys. (Paris), Ser. 3, 28:71. [29] Pasteur, L. 1852. Ann. Chem. 82:331. [30] Pryde, J. 1923. J. Chem. Soc. 123:1808. [31] Rehorst, K. 1928. Ber. Deut. Chem. Ges. 61:163. [32] Rehorst, K. 1930. Ibid. 63:2280. [33] Rehorst, K. 1933. Ann. Chem. 503:154. [34] Richtmyer, N. K., et al. 1939. J. Am. Chem. Soc. 61:343. [35] Robbins, G. B., and F. W. Upson. 1940. Ibid. 62:1074. [36] Schoeffel, E., and K. P. Link. 1933. J. Biol. Chem. 100:397. [37] Stodola, F. H., and L. B. Lockwood. 1947. Ibid. 171:213. [38] Walden, P. 1896. Ber. Deut. Chem. Ges. 29:1701. [39] Weinmann, F. 1929. Ibid. 62:1637. [40] Williams, A. K., and R. G. Eagon. 1959. J. Bacteriol. 77:167. [41] Williamson, A. R., and A. Zamenhof. 1963. J. Biol. Chem. 238:2255.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part V. NATURAL CARBOHYDRATE PHOSPHATE ESTERS**

Wavelength (column F): D = the sodium D line, 5896 Ångstrom units.

Substance (Synonym)	Hydrolysis Constant <sup>1</sup> k x 10 <sup>3</sup>	Temp. °C	Medium	Specific Rotation [α] <sub>D</sub>	Wave-length	Compound	Concentration	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Dihydroxyacetone phosphate	33.7	100	N HCl	.....	.....	.....	.....	34
2 D-Glyceraldehyde 3-phosphate	37.5	100	N HCl	+12	D	Free acid	.....	19,34, 54
3 D-Glyceric acid 2-phosphate	.....	....	.....	+24.3, +13 -68, +5	D D	Free acid Free acid	(N HCl) (NH <sub>4</sub> molybdate)	6,35 6,55
4 D-Glyceric acid 3-phosphate	1.8 <sup>2</sup>	125	N HCl	-14.5, +14 -725	D D	Ba salt Ba salt	(N HCl) (Molybdate ion)	6,35 56
5 Glyceric acid phosphate, 2-deoxy-	15 <sup>2</sup>	90	N HCl	.....	.....	.....	.....	8
6 D-Glyceric acid 1,3-diphosphate	26	38	Water	Very small	.....	.....	.....	58
7 D-Glyceric acid 2,3-diphosphate	.....	....	.....	-2, -4 -4, -5, +4.6	D D	Ba salt Na salt	6-17 (N HNO <sub>3</sub> ) 6-28	4,23 4,23, 78
8 L-Glycerol 1-phosphate (α-L-Glycerophosphate; L-Glycerin 1-phosphate)	0.15	80	Water pH 6.3	+1.0	D	Ag salt	6.5	5,36, 87
9 D-Erythritol 4-phosphate	.....	....	.....	-2.6	D	Free acid	.....	73
10 D-Erythrose 4-phosphate	.....	....	.....	0	D	Free acid	.....	7
11 L-Erythrulose 1-phosphate	10 <sup>2</sup>	100	N HCl	.....	.....	.....	.....	15
12 D-Arabinose 5-phosphate	<3 <sup>2</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	86
13 L-Arabinofuranose 1-phosphate	.....	....	.....	+16.9	D	Free acid	.....	93
14 L-Arabinopyranose 1-phosphate	.....	....	.....	+48.2	D	Ba salt	.....	93
15 Pentose 1-phosphate, α-D-erythro-	.....	....	.....	+34.5	D	Cyclohexylamine salt	.....	48
16 Pentose 1-phosphate, β-D-erythro-	13-17 <sup>2</sup>	....	Acetate buffer pH 4.5	-15.8	D	Cyclohexylamine salt	.....	20,48, 88
17 Pentose 5-phosphate, β-D-erythro-	50 <sup>2</sup>	100	N HCl	+19	D	Free acid	0.47	48,65
18 Pentose 1,5-diphosphate, D-erythro-	>3 <5 <sup>3</sup>	100 100	pH 4 N HCl	.....	.....	.....	.....	83 83
19 D-Ribitol 5-phosphate (L-Ribitol 1-phosphate)	<5 <sup>2</sup>	100	N HCl	.....	.....	.....	.....	3
20 α-D-Ribofuranose 1-phosphate	1.25 <sup>2</sup>	20	0.01 N HCl	+40.3	D	Dicyclohexylamine salt	(Water)	84,92, 93
21 β-D-Ribofuranose 1-phosphate	0.63 <sup>2</sup>	20	0.01 N HCl	-9.3 -13.6	D D	Ba salt Dicyclohexylamine salt	..... (Ethanol)	92,93 84
22 D-Ribopyranose 1-phosphate	1200 1.25 <sup>2</sup>	25 20	0.5 N HCl 0.1 N HCl	-12.9 -47.1	D D	Free acid Ba salt	..... (5% acetic acid)	32,41 92,93
23 D-Ribose 2-phosphate	.....	....	.....	-6.8	D	Ba salt	.....	33
24 D-Ribose 3-phosphate	.....	....	.....	-6.8	D	Ba salt	.....	33
25 D-Ribose 5-phosphate	4.5 0.5	100 100	0.25 N HCl 0.25 N HCl	-9.7 +6.0 +18±2	D D D	Na salt Ba salt Free acid	..... ..... (0.2-1 N HCl)	1,44 1,46 26,57
26 α-D-Ribose 1,5-diphosphate	1.6	70	0.01 N HCl	+20.8	D	Tetracyclohexylamine salt	0.43	85
27 D-Ribose 5-phosphate 1-pyrophosphate	30 <sup>4</sup>	65	pH 4	.....	.....	.....	.....	37

<sup>1/1</sup> For the first ester group that lies farthest in the sugar carbon-chain structure from the primary hydroxyl carbon (or asymmetric center) which determines the D or L configuration of the parent sugar. <sup>1/2</sup> Calculated by the contributors from data of the original investigator, using k = 0.30/time in minutes for 50% hydrolysis. <sup>1/3</sup> For the second ester group. <sup>1/4</sup> For the pyrophosphate group.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part V. NATURAL CARBOHYDRATE PHOSPHATE ESTERS**

Substance (Synonym)	Hydrolysis Constant <sup>1</sup> k x 10 <sup>3</sup>	Temp. °C	Medium	Specific Rotation [α] <sub>D</sub>	Wave-length	Compound	Concentration	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
28 D-Ribulose 5-phosphate	5	100	N HCl	-29	D	Free acid	(0.2 N HCl)	27,30
29 L-Ribulose 5-phosphate	<0.5	90	N H <sub>2</sub> SO <sub>4</sub>	+28	D	Free acid	(0.2 N HBr)	75,76
30 D-Ribulose 1, 5-diphosphate	15 <sup>b</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	29
31 D-Xylose 5-phosphate	4	100	N HCl	+3.2	D	Na salt	.....	45
				+5	D	Ba salt	.....	45
32 D-Xylulose phosphate	86	100	N HCl	.....	.....	.....	.....	22,77
33 D-Fructose 1-phosphate	70	100	N HCl	-64.2	5461	Free acid	11.3	82
				-39	5461	Ba salt	6.1	82
34 D-Fructose 6-phosphate	4.4	100	N HCl	+3.6	D	Ba salt	10	59,69
35 D-Fructose 1, 6-diphosphate	52	100	N HCl	+4.1	D	Free acid	13.6	50,59
36 L-Fuculose 1-phosphate	60	100	N HCl	-2.3	D	Ba salt	5.1	24
37 α-D-Galactose 1-phosphate	5.9	37	0.25 N HCl	+108	D	K salt	2.6	38,39
				+148	D	Free acid	1.7 (0.2 N HCl)	38,39
				+92; +113	D; 5461	Ba salt	.....	38,39
				+78.5	D	Dicyclohexylamine salt	(pH 7.8)	64
38 β-D-Galactose 1-phosphate	5.6	37	0.25 N HCl	+31.3	D	Ba salt·3H <sub>2</sub> O	1.2	66
				+21	D	Dicyclohexylamine salt	(pH 7.8)	64
39 D-Galactose 1-phosphate, 2-amino-2-deoxy-	.....	.....	.....	178	D	N-Acetyl derivative	.....	13,14
40 D-Galactose 6-phosphate	.....	.....	.....	+25.2	D	Ba salt	[16°]	31,80, 81
				-11.9	D	CH <sub>3</sub> β-D-galactoside dicyclohexylamine salt	0.5	79
41 D-Galactose 6-phosphate, 2-amino-2-deoxy-	8 <sup>d</sup>	110	6 N HCl	48.4	D	N-Acetyl derivative	(0.05 M Na acetate)	18,47
42 D-Gluconic acid 6-phosphate	0.21	100	N HCl	+0.2	5461	Free acid	.....	61,68
				+18	5461	Free acid lactone	.....	68
43 D-Glucose 1-phosphate, 2-amino-2-deoxy-	60 <sup>d</sup>	100	N HClO <sub>4</sub>	+79	D	N-Acetyl derivative, α-form	.....	11,51
				-1.6	D	N-Acetyl derivative, β-form	.....	9
44 D-Glucose 6-phosphate, 2-amino-2-deoxy-	0.06 <sup>d</sup>	100	N HCl	+54	D	Free acid	.....	2
				+29.5	D	N-Acetyl derivative	8 (0.5 M Na acetate)	18
45 α-D-Glucose 1-phosphate	1.3	37	0.25 N HCl	+118	D	Free acid	.....	16
	5 <sup>c</sup>	33	N HCl	+0.5	D	Dibrucine salt·8H <sub>2</sub> O	.....	91
				+75.5	D	Ba salt	.....	16
				+78; +90	D; 5461	K salt·2H <sub>2</sub> O	.....	90
				+64	D	Dicyclohexylamine salt	(pH 7.8)	64
46 β-D-Glucose 1-phosphate	15 <sup>b</sup>	33	N HCl	-20	D	Dibrucine salt·10H <sub>2</sub> O	.....	91
				+7.3	D	Dicyclohexylamine salt	.....	64
47 D-Glucose 6-phosphate	0.23	100	N HCl	+35.7;	D; 5461	Free acid	.....	68,69
				+41.4	D; 5461	Ba salt	8.4	68,69
				+18; +21.2	D; 5461	K salt	.....	40
				+21.2	D	K salt	.....	40

<sup>1</sup>/ For the first ester group that lies farthest in the sugar carbon-chain structure from the primary hydroxyl carbon (or asymmetric center) which determines the D or L configuration of the parent sugar. <sup>2</sup>/ Calculated by the contributors from data of the original investigator, using k = 0.30/time in minutes for 50% hydrolysis. <sup>3</sup>/ Both groups hydrolyze equally. <sup>4</sup>/ Constants determined on the brucine salt.

*continued*



**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part V. NATURAL CARBOHYDRATE PHOSPHATE ESTERS**

Substance (Synonym)	Hydrolysis Constant <sup>1</sup> k x 10 <sup>3</sup>	Temp. °C	Medium	Specific Rotation [α] <sub>D</sub>	Wave-length	Compound	Concentration	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
D-Glucose 6-phosphate (See preceding page)	0.23	100	N HCl	+61	D	CH <sub>3</sub> α-D-glucoside dicyclohexylamine salt	.....	79
48 D-Glucose phosphate, 2-amino-3-O-(2-carboxyethyl)-2-deoxy- (Muramic acid phosphate)	0.8 <sup>2</sup>	100	6 N HCl	.....	.....	.....	.....	47
49 α-D-Glucose 1,6-diphosphate	0.78	30	N H <sub>2</sub> SO <sub>4</sub>	+83±4	D	Free acid	0.2	63
50 β-D-Glucose 1,6-diphosphate	3.15	30	N H <sub>2</sub> SO <sub>4</sub>	-19±2	D	Free acid	0.2	63
51 D-Glucuronic acid 1-phosphate	.....	.....	.....	+53.6	D	Tri K salt form	[19°]	10
52 2-Hexulosonic acid 6-phosphate, 3-deoxy-D-erythro- <sup>7</sup>	5-6 <sup>2</sup>	100	N HCl	.....	.....	.....	.....	49
53 myo-Inositol 1-phosphate (myo-Inositol 3-phosphate)	0.99	100	pH 2.0	+3.4	D	Dicyclohexylamine salt	(pH 9)	62
				-9.8	D	Dicyclohexylamine salt	(pH 2)	62
54 myo-Inositol 2-phosphate	.....	.....	.....	None (meso)	.....	.....	.....	62
55 D-Mannitol 1-phosphate	<0.5 <sup>2</sup>	100	N HCl	.....	.....	.....	.....	89
56 D-Mannose 6-phosphate	0.29	100	N HCl	+15.1	5461	Free acid	1.7	69
				+3.5	5461	Ba salt	0.7	41
57 Shikimic acid 5-phosphate	.....	.....	.....	-107.6	D	K salt·H <sub>2</sub> O	[29°]	88
58 L-Sorbose 1-phosphate	60 <sup>2</sup>	100	N HCl	-16.5	D	Mono K salt	.....	25,52
				-7.2	D	Ba salt·2H <sub>2</sub> O	(0.1 N HCl)	52
59 2-Heptulose 7-phosphate, D-altro- (Sedoheptulose 7-phosphate)	0.28 <sup>2</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	28
60 2-Heptulose 1,7-diphosphate, D-altro- (Sedoheptulose 1,7-diphosphate)	20 <sup>4</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	28
	0.28 <sup>3</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	28
61 Unidentified ketoheptose monophosphate <sup>5</sup>	4	100	N HCl	+8	5461	Ba salt	.....	70
62 α, α'-Diglycerophosphate	150 <sup>2</sup>	100	N HNO <sub>3</sub>	.....	.....	.....	.....	53
63 1-Glycerophosphoryl-myo-inositol	15 <sup>2</sup>	100	N HCl	-14	D	Cyclohexylamine salt	6 (pH 3.5)	43
64 α-Lactose 1-phosphate	2 <sup>2</sup>	37	N HCl	73.3	D	Ba salt	.....	21,71, 72
65 β-Lactose 1-phosphate	6 <sup>2</sup>	37	N HCl	24.8	D	Ba salt·5H <sub>2</sub> O	.....	21,71, 72
66 Sucrose 1-phosphate	5.9 <sup>2</sup>	100	N H <sub>2</sub> SO <sub>4</sub>	.....	.....	.....	.....	42
67 Trehalose phosphate	0.16	.....	.....	+185	D	.....	(0.1 N HCl)	12,67
				+132	5461	Ba salt	.....	12,67

<sup>1</sup>/ For the first ester group that lies farthest in the sugar carbon-chain structure from the primary hydroxyl carbon (or asymmetric center) which determines the D or L configuration of the parent sugar. <sup>2</sup>/ Calculated by the contributors from data of the original investigator, using k = 0.30/time in minutes for 50% hydrolysis. <sup>3</sup>/ For the second ester group. <sup>4</sup>/ For the pyrophosphate group. <sup>5</sup>/ D-arabino-Hexulosonic acid is also found in natural systems as a phosphate, but characterization constants are unknown [17]. <sup>6</sup>/ A D-arabino-3-hexulose phosphate and a manno-heptulose phosphate are known in nature, but their characterization constants are unknown [60,74].

**Contributors:** Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

**References:** [1] Albaum, H. C., and W. W. Umbreit. 1947. J. Biol. Chem. 167:369. [2] Anderson, J. M., and E. Percival. 1956. J. Chem. Soc., p. 814. [3] Baddiley, J., et al. 1956. Ibid., p. 4583. [4] Baer, E., et al. 1948. J. Am. Chem. Soc. 70:1394. [5] Baer, E., et al. 1950. J. Biol. Chem. 185:763. [6] Ballou, C. E., et al. 1954. J. Am. Chem. Soc. 76:3188. [7] Ballou, C. E., et al. 1955. Ibid. 77:5967. [8] Ballou, C. E., et al. 1956. Ibid. 78:3718. [9] Baluja, G., et al. 1960. J. Chem. Soc., p. 4678. [10] Barker, S. A., et al. 1958. Ibid., p. 4128.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part V. NATURAL CARBOHYDRATE PHOSPHATE ESTERS**

- [11] Brown, D. H. 1953. *J. Biol. Chem.* 204:877. [12] Cabib, E., and L. F. Leloir. 1958. *Ibid.* 231:259.  
 [13] Cardini, C. E., and L. F. Leloir. 1953. *Arch Biochem. Biophys.* 45:55. [14] Cardini, C. E., and L. F. Leloir. 1957. *J. Biol. Chem.* 225:317. [15] Charalampous, F. C., and G. C. Mueller. 1953. *Ibid.* 201:161.  
 [16] Cori, C. F., S. P. Colowick, and G. T. Cori. 1937. *Ibid.* 121:465. [17] De Ley, J., and S. Verhofstede. 1955. *Naturwissenschaften* 42:584. [18] Distler, J. J., J. M. Merrick, and S. Roseman. 1958. *J. Biol. Chem.* 230:497. [19] Fischer, H. O. L., and E. Baer. 1932. *Ber. Deut. Chem. Ges.* 65:337, 1040. [20] Friedkin, M. 1950. *J. Biol. Chem.* 184:449. [21] Gander, J. E., W. E. Petersen, and P. D. Boyer. 1957. *Arch Biochem. Biophys.* 69:85. [22] Glock, G. 1952. *Biochem. J.* 52:575. [23] Greenwald, I. 1925. *J. Biol. Chem.* 63:339.  
 [24] Heath, E. C., and M. A. Ghalambor. 1962. *Ibid.* 237:2423. [25] Hers, H. G. 1952. *Biochim. Biophys. Acta* 8:416. [26] Horecker, B. L., et al. 1950. *Arch. Biochem.* 29:232. [27] Horecker, B. L., et al. 1951. *J. Biol. Chem.* 193:383. [28] Horecker, B. L., et al. 1955. *Ibid.* 212:827. [29] Horecker, B. L., et al. 1956. *Ibid.* 218:785. [30] Hurwitz, J., et al. 1956. *Ibid.* 218:769. [31] Inouye, T., M. Tannenbaum, and D. Y. Hsia. 1962. *Nature* 193:67. [32] Kalckar, H. M. 1947. *J. Biol. Chem.* 167:477. [33] Khym, J. X., D. G. Doherty, and W. E. Cohn. 1954. *J. Am. Chem. Soc.* 76:5523. [34] Kiessling, W., et al. 1934. *Ber. Deut. Chem. Ges.* 67:869. [35] Kiessling, W., et al. 1935. *Ibid.* 68:243. [36] Kiessling, W., et al. 1938. *Ibid.* 71:123. [37] Kornberg, A., I. Lieberman, and E. S. Simms. 1955. *J. Biol. Chem.* 215:389. [38] Kosterlitz, H. W. 1939. *Biochem. J.* 33:1087. [39] Kosterlitz, H. W. 1943. *Ibid.* 37:318. [40] Lardy, H. A., and H. O. L. Fischer. 1946. *J. Biol. Chem.* 164:513. [41] Leloir, L. F., et al. 1951. *Fortschr. Chem. Org. Naturstoffe* 8:47. [42] Leloir, L. F., et al. 1955. *J. Biol. Chem.* 214:157. [43] Lepage, M., R. Mumma, and A. A. Benson. 1960. *J. Am. Chem. Soc.* 82:3713. [44] Levene, P. A., et al. 1933. *J. Biol. Chem.* 101:419. [45] Levene, P. A., et al. 1933. *Ibid.* 102:307. [46] Levene, P. A., et al. 1934. *Ibid.* 104:299. [47] Liu, T., and E. C. Gotschlich. 1963. *Ibid.* 238:1928. [48] MacDonald, D. L., and H. G. Fletcher, Jr. 1962. *J. Am. Chem. Soc.* 84:1262. [49] MacGee, J., and M. Doudoroff. 1954. *J. Biol. Chem.* 210:617. [50] MacLeod, M., and R. Robison. 1933. *Biochem. J.* 27:286. [51] Maley, F., G. F. Maley, and H. A. Lardy. 1956. *J. Am. Chem. Soc.* 78:5303. [52] Mann, K. M., and H. A. Lardy. 1950. *J. Biol. Chem.* 187:339. [53] Maruo, B., and A. A. Benson. 1957. *J. Am. Chem. Soc.* 79:4564. [54] Meyerhof, O., et al. 1943. *J. Biol. Chem.* 149:71. [55] Meyerhof, O., et al. 1949. *Ibid.* 179:1371. [56] Meyerhof, O., et al. 1949. *Ibid.* 179:1381. [57] Michelson, A. M., and A. R. Todd. 1949. *J. Chem. Soc.*, p. 2476. [58] Negelein, E., and H. Brömel. 1939-40. *Biochem. Z.* 303:132. [59] Neuberg, C., H. Lustig, and M. Rothenberg. 1944. *Arch. Biochem.* 3:33. [60] Nordahl, A., and A. A. Benson. 1954. *J. Am. Chem. Soc.* 76:5054. [61] Patwardhan, V. R. 1934. *Biochem. J.* 28:1854. [62] Pizer, F. L., and C. E. Ballou. 1959. *J. Am. Chem. Soc.* 81:915, 4745. [63] Posternak, T. 1949. *J. Biol. Chem.* 180:1269. [64] Putman, E. W., and W. Z. Hassid. 1957. *J. Am. Chem. Soc.* 79:5057. [65] Racker, E. 1952. *J. Biol. Chem.* 196:347. [66] Reithel, F. J. 1945. *J. Am. Chem. Soc.* 67:1056. [67] Robison, R., et al. 1928. *Biochem. J.* 22:1277. [68] Robison, R., et al. 1931. *Ibid.* 25:323. [69] Robison, R., et al. 1932. *Ibid.* 26:2191. [70] Robison, R., et al. 1938. *Nature* 142:114. [71] Sasaki, R., and K. Taniguchi. 1959. *Nippon Nogei Kagaku Kaishi* 33:183. [72] Sasaki, R., and K. Taniguchi. 1960. *Chem. Abstr.* 54:308. [73] Shetter, J. K. 1956. *J. Am. Chem. Soc.* 78:3722. [74] Sie, H. G., V. N. Nigam, and W. H. Fishman. 1959. *Ibid.* 81:6083. [75] Simpson, F. J., and W. A. Wood. 1956. *Ibid.* 78:5452. [76] Simpson, F. J., and W. A. Wood. 1958. *J. Biol. Chem.* 230:473. [77] Stumpf, P. K., and B. L. Horecker. 1956. *Ibid.* 218:753. [78] Sutherland, E. W., T. Posternak, and C. F. Cori. 1949. *Ibid.* 181:153. [79] Szabo, P., and L. Szabo. 1960. *J. Chem. Soc.*, p. 3762. [80] Tanaka, T. 1961. *Yakugaku Zasshi* 81:797. [81] Tanaka, T. 1961. *Chem. Abstr.* 55:27064. [82] Tanko, B., and R. Robison. 1935. *Biochem. J.* 29:961. [83] Tarr, H. L. A. 1957. *Chem. Ind. (London)*, p. 562. [84] Tener, G. M., et al. 1957. *J. Am. Chem. Soc.* 79:441. [85] Tener, G. M., et al. 1958. *Ibid.* 80:1999. [86] Volk, W. A. 1959. *J. Biol. Chem.* 234:1931. [87] Weil-Malherbe, H., and R. H. Green. 1951. *Biochem. J.* 49:286. [88] Weiss, U., and E. S. Mingioli. 1956. *J. Am. Chem. Soc.* 78:2894. [89] Wolff, J. B., and N. O. Kaplan. 1957. *J. Biol. Chem.* 218:849. [90] Wolfrom, M. L., et al. 1941. *J. Am.*

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part V. NATURAL CARBOHYDRATE PHOSPHATE ESTERS**

Chem. Soc. 63:1051. [91] Wolfrom, M. L., et al. 1942. Ibid. 64:23. [92] Wright, R. S., and H. G. Khorana. 1956. Ibid. 78:811. [93] Wright, R. S., and H. G. Khorana. 1958. Ibid. 80:1994.

**Part VI. NATURAL OLIGOSACCHARIDES**

Substances have been arranged in groups (disaccharides, trisaccharides, etc.) according to increasing carbon atom content in the parent component monosaccharide units. Within groups, substances are arranged alphabetically according to the name of the initial glycosyl monosaccharide unit in the oligosaccharide. **Substance** (column A): Gal = galactose; Man = mannose; Xyl = xylose; G = glucose; Fru = fructose; Fuc = fucose; *p* = pyranose; *f* = furanose.

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Reference
	(A)	(B)	(C)	(D)	(E)
1	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 1)- <i>D</i> -glycerol	C <sub>9</sub> H <sub>18</sub> O <sub>8</sub>	150-152	+155	111
2	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 1)- <i>D</i> -glycerol	C <sub>9</sub> H <sub>18</sub> O <sub>8</sub>	139-140	+3.8	13
3	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 2)-glycerol (Floridoside)	C <sub>9</sub> H <sub>18</sub> O <sub>8</sub>	86-87, 128	+151, +165	14, 70, 84
4	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 1)- <i>L</i> -glycerol	C <sub>9</sub> H <sub>18</sub> O <sub>8</sub>	131.5-133	+159	111
5	<i>O</i> -α- <i>D</i> -Man $p$ -(1 → ?)- <i>L</i> -glyceric acid	C <sub>9</sub> H <sub>16</sub> O <sub>9</sub>	88-89	+105 [15°]	15
6	<i>O</i> - <i>D</i> -Xyl $p$ -(1 → 1)- <i>D</i> -G $p$ <sup>1</sup>	C <sub>11</sub> H <sub>20</sub> O <sub>10</sub>	Amorphous	-36.5	68
7	<i>O</i> - <i>D</i> -Xyl $p$ -(1 → 6)- <i>D</i> -G $p$	C <sub>11</sub> H <sub>20</sub> O <sub>10</sub>	208	+24.1 → -3.3	26, 94
8	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 3)- <i>D</i> -arabinitol	C <sub>11</sub> H <sub>22</sub> O <sub>10</sub>	138-139	-81	41
9	<i>O</i> - <i>D</i> -Fru $f$ -(2 → 2)- <i>D</i> -Fru $f$ (Alliuminoside)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	92-93	-23.8	78, 79
10	<i>O</i> - <i>D</i> -Fru $f$ -(2 → 1)- <i>D</i> -Fru $f$ (Inulobiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	-26.4	81, 82
11	<i>O</i> - <i>D</i> -Fru $f$ -(2 → ?)- <i>D</i> -Fru $f$ (Sogdianose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	156-158	-16.4	78, 79
12	<i>O</i> - <i>D</i> -Fru $f$ -(2 → ?)- <i>D</i> -G $p$	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	-21	94
13	<i>O</i> -β- <i>D</i> -Fru $f$ -(2 → 1)-α- <i>D</i> -G $p$ (Sucrose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	188, 170 <sup>2</sup>	+66.5 ( <i>c</i> 26)	9
14	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 6)- <i>D</i> -Gal $p$ (Swietenose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	+149	30, 31, 91
15	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 4)-α- <i>D</i> -G $p$ (α-Lactose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub> ·H <sub>2</sub> O	202	+83.5 → +52.6	21, 90
16	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 4)-β- <i>D</i> -G $p$ (β-Lactose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	252	+34.2 → +53.6	21, 89, 90
17	<i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 6)- <i>D</i> -G $p$ (Allolactose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	165	+25	66, 67
18	<i>O</i> - <i>D</i> -Gal-(? → ?)- <i>D</i> -G (Gynolactose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	205	-27	66, 67
19	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 6)-β- <i>D</i> -G $p$ (β-Melibiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub> ·2H <sub>2</sub> O	82-85	+111.7 → +129.5	9
20	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 6)-β- <i>D</i> -glucitol (Melibiotol)	C <sub>12</sub> H <sub>24</sub> O <sub>11</sub>	173-175	+111	4
21	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 1)- <i>myo</i> -inositol	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub> ·2H <sub>2</sub> O	220-222	+135.6	12
22	<i>O</i> - <i>D</i> -Gal $p$ -(1 → ?)-mannitol	C <sub>12</sub> H <sub>24</sub> O <sub>11</sub>	162	-55.5	69
23	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 3)- <i>D</i> -Fru (Turanose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	157	+22 → +75.3	29
24	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 4)- <i>D</i> -Fru $f$ (Maltulose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	113-115 d. (monohydrate)	58 → 64	28, 103
25	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 6)- <i>D</i> -Fru $f$ (Palatinose; Iso-maltulose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	+97.2	4, 99
26	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 1)-α- <i>D</i> -G $p$ (Trehalose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub> ·2H <sub>2</sub> O	97, 203	+178.3 ( <i>c</i> 7)	9, 92
27	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 2)- <i>D</i> -G $p$ (Kojibiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	175	+135	86, 87, 95, 96
28	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 3)- <i>D</i> -G $p$ (Nigerose; Sakebiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	+145	23, 95, 96
29	<i>O</i> -β- <i>D</i> -G $p$ -(1 → 3)-α- <i>D</i> -G $p$ (α-Laminaribiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	160-163, 202-205	+23.4 → +19	8, 100
30	<i>O</i> -β- <i>D</i> -G $p$ -(1 → 3)-β- <i>D</i> -G $p$ (β-Laminaribiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	188-192	+7.5 → +20.8	4, 16
31	<i>O</i> -β- <i>D</i> -G $p$ -(1 → 4)-β- <i>D</i> -G $p$ (β-Cellobiose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	225	+14.2 → +34.6 ( <i>c</i> 8)	65
32	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 4)-α- <i>D</i> -G $p$ (α-Maltose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	108	+173	22
33	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 4)-β- <i>D</i> -G $p$ (β-Maltose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub> ·H <sub>2</sub> O	102-103	+112.5 → +130	22
34	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 6)- <i>L</i> -G $p$ (Isomaltose)	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	.....	+103.2	95, 96, 109
35	<i>O</i> -α- <i>D</i> -G $p$ -(1 → 1)-α- <i>D</i> -GN (Trehalosamine)	C <sub>12</sub> H <sub>23</sub> NO <sub>10</sub>	.....	+176 ( <i>c</i> 0.02, dil. HCl)	2, 3
36	<i>O</i> -β- <i>D</i> -G $p$ -(1 → 1)-mannitol	C <sub>12</sub> H <sub>21</sub> O <sub>11</sub>	140-141	-18.0	42
37	<i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 6)- <i>O</i> -β- <i>D</i> -Gal $p$ -(1 → 1)-glycerol	C <sub>15</sub> H <sub>28</sub> O <sub>13</sub>	196-198	+90	104
38	<i>O</i> -α- <i>D</i> -Man $p$ -(1 → 3)- <i>O</i> -α- <i>D</i> -Gal $p$ -(1 → 2)-glycerol	C <sub>15</sub> H <sub>28</sub> O <sub>13</sub>	.....	.....	43
39	<i>O</i> - <i>D</i> -Fru $f$ -(2 → ?)- <i>O</i> - <i>D</i> -Fru $f$ -(2 → 2)- <i>D</i> -Fru $f$ (Polygontin)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	207-208	-52.9	78, 79

<sup>1/2</sup> The free sugar does not exist in nature, but its dibenzoyl derivatives do. <sup>2/2</sup> Compound crystallizes in one of two forms, depending on the solvent used.

*continued*



98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS

Part VI. NATURAL OLIGOSACCHARIDES

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation $[\alpha]_D$	Reference
	(A)	(B)	(C)	(D)	(E)
40	<i>O</i> - <i>D</i> -Fruf-(2 → ?)- <i>O</i> - <i>D</i> -Fruf-(2 → 2)- <i>D</i> -Fruf (Trifructan)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	-22.3	81,82
41	<i>O</i> - <i>D</i> -Fruf-(2 → ?)- <i>O</i> - <i>D</i> -Fruf-(2 → 1)- <i>D</i> -Gp	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	+22	5,24
42	<i>O</i> - <i>β</i> - <i>D</i> -Fruf-(2 → 6)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Neokestose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	+15	24
43	<i>O</i> - <i>L</i> -Fuc-(1 → 2)- <i>O</i> - <i>D</i> -Galp-(1 → 4)- <i>β</i> - <i>D</i> -Gp	C <sub>18</sub> H <sub>32</sub> O <sub>15</sub>	230-231 d.	-57.5(c 0.2)	37, 49
44	<i>O</i> - <i>D</i> -Gal-(? → ?)- <i>O</i> - <i>D</i> -Fru-(? → ?)- <i>D</i> -Fru (Labiase)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub> ·3H <sub>2</sub> O	126 s., b.p. 205	+136.7	76,77
45	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(1 → 6)- <i>α</i> - <i>D</i> -Gp (Planteose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub> ·2H <sub>2</sub> O	123-124	+125.2	19,97,98
46	[ <i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)-] <sub>2</sub> <i>α</i> - <i>D</i> -Gp (Manninotriose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	150 amorphous	+167	88
47	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 2)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Umbelliferose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	+125	105
48	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 3)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	.....	45
49	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Raffinose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub> ·5H <sub>2</sub> O	80, 118-120	+105, +123.1	44,71
50	<i>O</i> - <i>D</i> -Galp-(1 → 4)- <i>O</i> - <i>D</i> -Gp-(1 → ?)- <i>L</i> -Fuc	C <sub>18</sub> H <sub>32</sub> O <sub>15</sub>	.....	.....	49
51	<i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Isokestose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	148, 189-190	+29.3	6,7,73
52	<i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(6 → 2)- <i>β</i> - <i>D</i> -Fruf (Kestose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	145	+28	1,18
53	<i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 3)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(2 → 1)- <i>α</i> - <i>D</i> -Gp (Melezitose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub> ·2H <sub>2</sub> O	153-154	+88.2	34,93
54	<i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 4)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Erllose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	.....	+121.8	102,108
55	<i>O</i> - <i>β</i> - <i>D</i> -Gp-(1 → 6)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Gentianose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	210	+33.4	10,47
56	[ <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 4)-] <sub>2</sub> <i>α</i> - <i>D</i> -Gp (Maltotriose)	C <sub>18</sub> H <sub>32</sub> O <sub>16</sub>	Amorphous	+160	85,110
57	<i>O</i> - <i>D</i> -Gp-(1 → 1)- <i>O</i> - <i>D</i> -Gp-(1 → 6)- <i>D</i> -mannitol	C <sub>18</sub> H <sub>34</sub> O <sub>16</sub>	.....	-14	34
58	<i>N</i> -Acetylneuraminic acid-(1 → ?)- <i>O</i> - <i>β</i> - <i>D</i> -Galp-(1 → 4)- <i>β</i> - <i>D</i> -Gp (Neuraminlactose)	.....	.....	10-26	75
59	Fructo-tetraose (Veronicin; Campanulin)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	170, 188	-23, -29.4 [30°]	57,59,61,62
60	<i>O</i> - <i>D</i> -Fruf-(2 → 1)- <i>O</i> - <i>D</i> -Fruf-(2 → 6)- <i>O</i> - <i>D</i> -Gp-(2 → 2)- <i>D</i> -Fruf (Neobifurcose)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	+14.4	74
61	<i>O</i> - <i>L</i> -Fuc-(1 → ?)- <i>O</i> - <i>L</i> -Fuc-(1 → 3)- <i>O</i> - <i>β</i> - <i>D</i> -Galp-(1 → 4)- <i>D</i> -Gp (Lactodifucotetraose)	C <sub>24</sub> H <sub>42</sub> O <sub>19</sub>	.....	-106	38
62	<i>O</i> - <i>L</i> -Fuc-(1 → ?)- <i>O</i> - <i>β</i> - <i>L</i> -Fuc-(1 → ?)- <i>O</i> - <i>β</i> - <i>D</i> -Galp- <i>D</i> -Gp	C <sub>24</sub> H <sub>42</sub> O <sub>19</sub>	.....	-17.1	48,50,51
63	<i>O</i> - <i>D</i> -Galp-(1 → ?)- <i>O</i> - <i>D</i> -Galp-(1 → ?)- <i>O</i> - <i>D</i> -Fruf-(2 → 1)- <i>D</i> -Gp (Sesamose) <sup>3</sup>	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	.....	25
64	[ <i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)-] <sub>2</sub> <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Stachyose, Manneotetraose)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	170 (140 s.)	+146.3	20,64,88
65	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(1 → 1)- <i>α</i> - <i>D</i> -Galp (Lychnose)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	+153 to +154	106
66	<i>O</i> - <i>α</i> - <i>D</i> -Galp-(1 → 6)- <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(3 → 1)- <i>α</i> - <i>D</i> -Galp (Isolychnose)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	.....	107
67	<i>O</i> - <i>β</i> - <i>D</i> -Galp-(1 → 3)- <i>O</i> - <i>β</i> - <i>D</i> -GpNAc-(1 → 3)- <i>O</i> - <i>β</i> - <i>D</i> -Galp-(1 → 4)- <i>D</i> -Gp (Lacto- <i>N</i> -tetraose)	C <sub>26</sub> H <sub>45</sub> NO <sub>21</sub>	200-205 d.	+38	35
68	<i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)-[ <i>O</i> - <i>β</i> - <i>D</i> -Fruf-(6 → 2)-] <sub>2</sub> <i>β</i> - <i>D</i> -Fruf-	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	-7	72
69	<i>O</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>D</i> -Fruf-(2 → 6)- <i>O</i> - <i>D</i> -Fruf-(1 → 2)- <i>D</i> -Fruf	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	156	+8	73
70	<i>O</i> - <i>D</i> -Gp-(1 → ?)-[ <i>D</i> -Fruf-] <sub>2</sub> <i>D</i> -Fruf	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	+15.6	80,83
71	[ <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 4)-] <sub>2</sub> <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 2)- <i>β</i> - <i>D</i> -Fruf (Maltosylsucrose)	C <sub>27</sub> H <sub>42</sub> O <sub>21</sub>	.....	.....	108
72	[ <i>O</i> - <i>α</i> - <i>D</i> -Gp-(1 → 4)-] <sub>3</sub> <i>D</i> -Gp (Maltotetraose)	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	.....	+176.4	101
73	Scorodose	C <sub>24</sub> H <sub>42</sub> O <sub>21</sub>	200 amorphous	-41.5	32,33
74	Fructo-pentaose	C <sub>30</sub> H <sub>52</sub> O <sub>26</sub>	.....	+8	72
75	<i>O</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>D</i> -Fruf-(2 → 6)-[ <i>O</i> - <i>D</i> -Fruf-(1 → 2)-] <sub>2</sub> <i>D</i> -Fruf	C <sub>30</sub> H <sub>52</sub> O <sub>26</sub>	.....	.....	73

<sup>3</sup>/ The evidence for this compound is based on paper chromatographic and methylation studies.

*continued*



98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS

Part VI. NATURAL OLIGOSACCHARIDES

	Substance (Synonym)	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Reference
	(A)	(B)	(C)	(D)	(E)
76	<i>O</i> -α- <i>L</i> -Fucp-(1 → 2)- <i>O</i> -β- <i>D</i> -Galp-(1 → 3)- <i>O</i> -β- <i>D</i> -GpNAc-(1 → 3)- <i>O</i> -β- <i>D</i> -Galp-(1 → 4)- <i>D</i> -Gp (Lacto- <i>N</i> -fucopentaose I)	C <sub>32</sub> H <sub>55</sub> NO <sub>25</sub>	216	-11 → -16.3	36
77	<i>O</i> -α- <i>L</i> -Fucp-(1 → 4)- <i>O</i> -β- <i>D</i> -Galp-(1 → 3)- <i>O</i> -β- <i>D</i> -GpNAc-(1 → 3)- <i>O</i> -β- <i>D</i> -Galp-(1 → 4)- <i>D</i> -Gp (Lacto- <i>N</i> -fucopentaose II)	C <sub>32</sub> H <sub>55</sub> NO <sub>25</sub>	213-215	-28 → +30.4	39
78	[ <i>O</i> -α- <i>D</i> -Galp-(1 → 6)-] <sub>3</sub> <i>O</i> -α- <i>D</i> -Gp-(1 → 2)-β- <i>D</i> -FruF (Verbascose)	C <sub>30</sub> H <sub>52</sub> O <sub>26</sub>	219-220, 253	+169.9	11,54,55
79	<i>O</i> - <i>D</i> -Gp-(1 → 2)- <i>O</i> - <i>D</i> -FruF-(2 → 6)-[ <i>O</i> - <i>D</i> -FruF-(1 → 2)-] <sub>2</sub> <i>D</i> -FruF	C <sub>30</sub> H <sub>52</sub> O <sub>26</sub>	.....	-3.5	73
80	<i>O</i> -α- <i>D</i> -Gp-(1 → 2)-[ <i>O</i> -β- <i>D</i> -FruF-(6 → 2)-] <sub>3</sub> β- <i>D</i> -FruF	C <sub>30</sub> H <sub>52</sub> O <sub>26</sub>	.....	-11.2	72
81	<i>O</i> -α- <i>L</i> -Fucp-(1 → 4)- <i>O</i> -β- <i>D</i> -Galp-(1 → 3)- <i>O</i> -β- <i>D</i> -GpNAc-(1 → 3)- <i>O</i> -β- <i>D</i> -Galp-(1 → 4)- <i>O</i> -α- <i>L</i> -Fucp-(1 → 3)- <i>D</i> -Gp (Lacto- <i>N</i> -difucohexaose II)	C <sub>38</sub> H <sub>65</sub> NO <sub>29</sub>	218-220 d.	-68.8	40
82	[ <i>O</i> - <i>D</i> -Galp-] <sub>4</sub> <i>O</i> - <i>D</i> -Galp-(1 → 2)- <i>D</i> -FruF (Lycopose)	C <sub>36</sub> H <sub>62</sub> O <sub>31</sub>	270	+187	56,63
83	Fructo-hexaose	C <sub>36</sub> H <sub>62</sub> O <sub>31</sub>	178	-41	58,60
84	[ <i>O</i> -α- <i>D</i> -Galp-(1 → 6)-] <sub>4</sub> <i>O</i> -α- <i>D</i> -Gp-(1 → 2)-β- <i>D</i> -FruF (Ajugose)	C <sub>36</sub> H <sub>62</sub> O <sub>31</sub> ·6H <sub>2</sub> O	204-205	+163	27
85	<i>O</i> -α- <i>D</i> -Gp-(1 → 2)-[ <i>O</i> -β- <i>D</i> -FruF-(6 → 2)-] <sub>4</sub> -β- <i>D</i> -FruF	C <sub>36</sub> H <sub>62</sub> O <sub>31</sub>	.....	-19	72
86	<i>O</i> - <i>D</i> -Gp-(1 → ?)-[ <i>O</i> - <i>D</i> -FruF-] <sub>4</sub> <i>D</i> -FruF	C <sub>35</sub> H <sub>62</sub> O <sub>31</sub>	.....	-5.3	80,83
87	Fructo-heptaose	C <sub>42</sub> H <sub>72</sub> O <sub>36</sub>	.....	-35.7	52,53
88	[ <i>O</i> -α- <i>D</i> -Galp-(1 → 6)-] <sub>5</sub> <i>O</i> -α- <i>D</i> -Gp-(1 → ?)-β- <i>D</i> -FruF	C <sub>42</sub> H <sub>72</sub> O <sub>36</sub>	246-248	+168	27
89	[ <i>O</i> -α- <i>D</i> -Galp-(1 → 6)-] <sub>4</sub> <i>O</i> -α- <i>D</i> -Gp-(1 → 2)- <i>O</i> -β- <i>D</i> -FruF-(3 → 1)- <i>O</i> -α- <i>D</i> -Galp	C <sub>42</sub> H <sub>72</sub> O <sub>36</sub>	.....	.....	17
90	Dilacto- <i>N</i> -tetraose	C <sub>52</sub> H <sub>88</sub> N <sub>2</sub> O <sub>41</sub>	.....	.....	46
91	[ <i>O</i> -α- <i>D</i> -Galp-(1 → 6)-] <sub>6</sub> <i>O</i> -α- <i>D</i> -Gp-(1 → ?)-β- <i>D</i> -FruF	C <sub>48</sub> H <sub>82</sub> O <sub>41</sub>	267-268	+168	27
92	Difucolacto- <i>N</i> -tetraose	C <sub>64</sub> H <sub>108</sub> N <sub>2</sub> O <sub>49</sub>	.....	.....	46

Contributors: Wolfrom, Melville L.; Maher, George G.; and Pagnucco, Rinaldo G.

References: [1] Albon, N., et al. 1953. *J. Chem. Soc.*, p. 24. [2] Arcamone, F., and F. Bizioli. 1957. *Gazz. Chim. Ital.* 87:896. [3] Arcamone, F., and F. Bizioli. 1959. *Chem. Abstr.* 52:4503. [4] Assarsson, A., and O. Theander. 1958. *Acta Chem. Scand.* 12:1319. [5] Bacon, J. S. D. 1959. *Nature* 184 (Suppl. 25): 1957. [6] Ballio, A., and S. Russi. 1956. *Gazz. Chim. Ital.* 86:476. [7] Ballio, A., and S. Russi. 1958. *Chem. Abstr.* 52:3922. [8] Barry, V. C. 1941. *Proc. Roy. Dublin Soc.* 24:423. [9] Bates, F. J., et al. 1942. *Natl. Bur. Std. (U.S.) Circ.* 440. [10] Binaghi, R., and P. Falqui. 1926. *Chem. Zentr.* 2:44. [11] Bourquelot, E., and M. Bridel. 1910. *Comp. Rend.* 151:760. [12] Brown, R. J., and R. F. Serro. 1953. *J. Am. Chem. Soc.* 75:1040. [13] Carter, H. E., R. H. McCluer, and E. D. Slifer. 1956. *Ibid.* 78:3735. [14] Colin, H., et al. 1937. *Bull. Soc. Chim. France, Ser. 5*, 4:277. [15] Colin, H., et al. 1939. *Compt. Rend.* 208:1450. [16] Connell, J. J., E. L. Hirst, and E. G. V. Percival. 1950. *J. Chem. Soc.*, p. 3494. [17] Courtois, J. E., P. Le Dizet, and A. Wickström. 1955. *Bull. Soc. Chim. Biol.* 40:1059. [18] Dedonder, R. 1951. *Compt. Rend.* 232:1134. [19] French, D., et al. 1953. *J. Am. Chem. Soc.* 75:709. [20] French, D., et al. 1953. *Ibid.* 75:3664. [21] Gillis, J. 1920. *Rec. Trav. Chim.* 39:88, 677. [22] Gillis, J. 1931. *Chem. Zentr.* 1:256. [23] Haq, S., et al. 1958. *J. Chem. Soc.*, p. 1342. [24] Haq, S., et al. 1961. *Can. J. Chem.* 39:1165. [25] Hatanka, S. 1959. *Arch. Biochem. Biophys.* 82:188. [26] Helferich, B., and H. Rauch. 1927. *Ann. Chem.* 455:168. [27] Herissey, H., et al. 1954. *Bull. Soc. Chim. Biol.* 36:1507. [28] Hough, L., J. K. N. Jones, and E. L. Richards. 1953. *J. Chem. Soc.*, p. 2005. [29] Hudson, C. S., and E. Pacsu. 1930. *J. Am. Chem. Soc.* 52:2519. [30] Ingle, T. R., and B. V. Bhide. 1958. *J. Indian Chem. Soc.* 35:516. [31] Ingle, T. R., and B. V. Bhide. 1959. *Chem. Abstr.* 53:11243. [32] Kihara, Y. 1936. *J. Agr. Chem. Soc. Japan* 13:1044. [33] Kihara, Y. 1937. *Chem. Abstr.* 31:3013. [34] Kuhn, R., et al.

*continued*

**98. CARBOHYDRATES: PHYSICAL AND CHEMICAL CHARACTERISTICS**

**Part VI. NATURAL OLIGOSACCHARIDES**

1926. Ber. Deut. Chem. Ges. 59:1655. [35] Kuhn, R., et al. 1953. Chem. Ber. 86:827. [36] Kuhn, R., et al. 1956. Chem. Ber. 89:504, 2514. [37] Kuhn, R., et al. 1956. Ibid. 89:2513. [38] Kuhn, R., et al. 1958. Ann. Chem. 611:249. [39] Kuhn, R., et al. 1958. Chem. Ber. 91:364. [40] Kuhn, R., et al. 1960. Ibid. 93:647. [41] Lindberg, B., et al. 1952. Acta Chem. Scand. 6:1052. [42] Lindberg, B., et al. 1953. Ibid. 7:1119. [43] Lindberg, B., et al. 1955. Ibid. 9:1093, 1097. [44] Loiseau, D. 1876. Compt. Rend. 82:1058. [45] MacLeod, A. M., and H. McCorquodale. 1958. Nature 182:815. [46] Malpress, F. H., and F. E. Hytten. 1958. Biochem. J. 68:708. [47] Meyer, A. 1882. Z. Physiol. Chem. 6:135. [48] Montreuil, J. 1955. St. Jans Hosp., Brugge (Belg.), 3e Colloq., p. 209. [49] Montreuil, J. 1956. Compt. Rend. 242:192. [50] Montreuil, J. 1956. Ibid. 242:828. [51] Montreuil, J. 1957. Chem. Abstr. 51:14950. [52] Murakami, S. 1937. Acta Phytochim. (Japan) 10:43. [53] Murakami, S. 1937. Chem. Abstr. 31:8570. [54] Murakami, S. 1940. Ibid. 34:3694. [55] Murakami, S. 1940. Proc. Imp. Acad. Tokyo 16:14. [56] Murakami, S. 1942. Acta Phytochim. (Japan) 13:37. [57] Murakami, S. 1944. Ibid. 14:101. [58] Murakami, S. 1949. Ibid. 15:105. [59] Murakami, S. 1949. Ibid. 15:109. [60] Murakami, S. 1949. Chem. Abstr. 43:8451c. [61] Murakami, S. 1949. Ibid. 43:8451e. [62] Murakami, S. 1951. Ibid. 45:8599. [63] Murakami, S. 1956. Ibid. 45:3465. [64] Onuki, M. 1933. Chem. Zentr. 2:367. [65] Peterson, F. C., and C. O. Spencer. 1927. J. Am. Chem. Soc. 49:2882. [66] Polonovski, M., and A. Lespagnol. 1931. Compt. Rend. 192:1319. [67] Polonovski, M., and A. Lespagnol. 1932. Ibid. 195:465. [68] Power, F. B., and A. H. Salway. 1914. J. Chem. Soc. 105:1062. [69] Pueyo, G. 1959. Compt. Rend. 248:2788. [70] Putman, E. W., and W. Z. Hassid. 1954. J. Am. Chem. Soc. 76:2221. [71] Scheibler, C. 1886. Ber. Deut. Chem. Ges. 19:2868. [72] Schlubach, H. H., et al. 1957. Ann. Chem. 606:130. [73] Schlubach, H. H., et al. 1958. Ibid. 614:126. [74] Schlubach, H. H., et al. 1961. Ibid. 647:41. [75] Schneir, M., R. J. Winzler, and M. E. Rafelson. 1962. Biochem. Prepn. 9:1. [76] Strepkov, S. M. 1939. J. Gen. Chem. (USSR) 9:1489. [77] Strepkov, S. M. 1940. Chem. Abstr. 34:2798. [78] Strepkov, S. M. 1958. Zh. Obshch. Khim. 28:3143. [79] Strepkov, S. M. 1959. Chem. Abstr. 53:10053. [80] Strepkov, S. M. 1959. Ibid. 53:20302. [81] Strepkov, S. M. 1959. Ibid. 53:21686. [82] Strepkov, S. M. 1959. Dokl. Akad. Nauk SSSR 124:1344. [83] Strepkov, S. M. 1959. Ibid. 125:216. [84] Su, J., and W. Z. Hassid. 1962. Biochemistry 1:468. [85] Sugihara, J. M., and M. L. Wolfrom. 1949. J. Am. Chem. Soc. 71:3357. [86] Takiura, K., and K. Koizuma. 1962. Yakugaku Zasshi 82:852. [87] Takiura, K., and K. Koizuma. 1963. Chem. Abstr. 58:6911. [88] Tanret, C. 1902. Bull. Soc. Chim. France, Ser. 3, 27:947. [89] Tanret, C. 1905. Z. Physik. Chem. (Leipzig) 53:692. [90] Trey, H. 1903. Ibid. 46:620. [91] Turton, C. N., et al. 1955. J. Am. Chem. Soc. 77:2565. [92] Von Lippmann, E. O. 1921. Ber. Deut. Chem. Ges. 45:3431. [93] Von Lippmann, E. O. 1927. Ibid. 60:161. [94] Wallenfels, K., and J. Lehrmann. 1957. Chem. Ber. 90:1000. [95] Watanabe, T., and K. Aso. 1960. Chem. Abstr. 54:23111. [96] Watanabe, T., and K. Aso. 1960. Tohoku J. Agr. Res. 11:109. [97] Wattiez, N., and M. Hans. 1943. Bull. Acad. Roy. Med. Belg. 8:386. [98] Wattiez, N., and M. Hans. 1945. Chem. Abstr. 39:4849. [99] Weidenhagen, R., and S. Lorenz. 1957. Angew. Chem. 69:641. [100] Weissmann, B., and K. Meyer. 1954. J. Am. Chem. Soc. 76:1753. [101] Whistler, R., and J. H. Duffy. 1955. Ibid. 77:1017. [102] White, J., et al. 1953. Ibid. 75:1259. [103] White, J., et al. 1959. Arch. Biochem. Biophys. 80:386. [104] Wickberg, B. 1959. Acta Chem. Scand. 12:1183, 1187. [105] Wickström, A., et al. 1956. Ibid. 10:1199. [106] Wickström, A., et al. 1958. Bull. Soc. Chim. France, Ser. 5, p. 1410. [107] Wickström, A., et al. 1959. Ibid., Ser. 5, p. 871. [108] Wolf, J. P. and W. H. Ewart. 1955. Arch. Biochem. Biophys. 58:365. [109] Wolfrom, M. L., et al. 1949. J. Am. Chem. Soc. 71:125. [110] Wolfrom, M. L., et al. 1949. Ibid. 71:2873. [111] Zemplén, G., and A. Gerecs. 1935. Ber. Deut. Chem. Ges. 68:2054.

## 99. GLYCOSIDES: CHARACTERISTICS,

**Melting Point** (columns C and K): d. = decomposes. **Solubility** (columns E-G): abs. = absolute; acet. = acetone; bz. = sl. = slightly; v. = very.

Glycoside	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Solubility			Occurrence
				H <sub>2</sub> O	Alcohol	Other	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Absinthin	C <sub>30</sub> H <sub>40</sub> O <sub>8</sub>	68	.....	sl. s.	s.	s. bz., chl., eth., NaOH	Wormwood
2 Aloin	Mixture	.....	.....	s.	s.	sl. s. chl., eth.	<i>Aloe</i> spp.
3 Amygdalin	C <sub>20</sub> H <sub>27</sub> O <sub>11</sub> N	220 <sup>1</sup>	-42	8.3 g/100 ml (h.)	sl. s.	i. eth.	Almonds
4 Apiin	C <sub>26</sub> H <sub>42</sub> O <sub>10</sub>	228 <sup>1</sup>	.....	s. h.	sl. s. h.	i. eth.	Celery; parsley
5 Arbutin	C <sub>12</sub> H <sub>16</sub> O <sub>7</sub>	195-200	-64	12.5 g/100 ml	7.7 g/100 ml	i. chl., CS <sub>2</sub> , eth.	Leaves of cranberry, pear tree
6 Barbaloin	C <sub>20</sub> H <sub>18</sub> O <sub>8</sub>	148	.....	s.	s.	sl. s. eth.	<i>Aloe</i> spp.
7 Bryonin	C <sub>48</sub> H <sub>66</sub> O <sub>18</sub>	208	.....	sl. s.	s.	i. chl., eth.	<i>Bryonia alba</i>
8 Carminic acid	C <sub>22</sub> H <sub>20</sub> O <sub>13</sub>	136 d.	.....	s.	s.	s. eth., NaOH; i. chl.	Cochineal
9 Coniferin	C <sub>16</sub> H <sub>22</sub> O <sub>8</sub>	185	-68	0.5 g/100 ml	sl. s.	i. eth.	Conifers; sugar beet
10 Convallatoxin	C <sub>29</sub> H <sub>42</sub> O <sub>10</sub>	247	0	0.05 g/100 ml	s.	sl. s. chl., eth.	Lily of the valley
11 Convolvulin	C <sub>54</sub> H <sub>96</sub> O <sub>27</sub>	155-168	.....	sl. s.	s.	s. acet.; i. eth.	Jalap resin, Canadian hemp
12 Crocin	C <sub>44</sub> H <sub>64</sub> O <sub>26</sub>	186 d. <sup>1</sup>	.....	sl. s.	sl. s.	i. chl., eth.	Saffron, crocus, gardenia
13 Cymarín	C <sub>30</sub> H <sub>44</sub> O <sub>9</sub>	139	+35	s.	.....	s. chl., me. al.	<i>Apocynum</i> & <i>Strophanthus</i> spp.
14 Daphnin	C <sub>15</sub> H <sub>16</sub> O <sub>9</sub>	125 d. <sup>1</sup>	-115	sl. s.	s.	s. NaOH; i. eth.	<i>Daphne</i> spp.
15 Diginin	C <sub>28</sub> H <sub>40</sub> O <sub>7</sub>	155-183	-176	i.	.....	s. CCl <sub>4</sub> , chl.; sl. s. eth.	Leaves of <i>Digitalis purpurea</i>
16 Digitonin	C <sub>55</sub> H <sub>90</sub> O <sub>29</sub>	235 d. <sup>1</sup>	-54	s.	1.8 g/100 ml (abs.)	i. chl., eth.	Seeds of <i>Digitalis purpurea</i>
17 Digitoxin	C <sub>41</sub> H <sub>64</sub> O <sub>13</sub>	256	+4.8	0.001 g/100 ml	1.7 g/100 ml	i. eth.	<i>Digitalis lanata</i> & <i>D. purpurea</i>
18 Digoxin	C <sub>41</sub> H <sub>64</sub> O <sub>14</sub>	265 d.	+13.3	0.001 g/100 ml	0.45 g/100 ml	i. chl., eth.	<i>Digitalis lanata</i>
19 Esculin	C <sub>15</sub> H <sub>16</sub> O <sub>9</sub>	205 d. <sup>1</sup>	-38	0.175 g/100 ml	5 g/100 ml	s. h. chl., NaOH; sl. s. eth.	Leaves and bark of horse-chestnut tree
20 Gaultherin	C <sub>19</sub> H <sub>26</sub> O <sub>12</sub>	179-180	-58	s.	s.	s. acet.; i. eth.	Wintergreen plant
21 Gitonin	C <sub>50</sub> H <sub>82</sub> O <sub>23</sub>	272	-51	sl. s.	0.98 g/100 ml	s. acet.; i. eth.	<i>Digitalis purpurea</i>
22 Gitoxin	C <sub>41</sub> H <sub>64</sub> O <sub>14</sub>	285	+3.5	sl. s.	.....	s. eth.; i. acet.	<i>Digitalis lanata</i> & <i>D. purpurea</i>
23 Gratiolin	C <sub>43</sub> H <sub>70</sub> O <sub>15</sub>	235-237	.....	sl. s.	s.	s. chl.; sl. s. eth.	<i>Gratiola</i> sp.
24 Hesperidin	C <sub>28</sub> H <sub>34</sub> O <sub>15</sub>	260-262	-76	v. sl. s.	sl. s. me. al.	s. NaOH; i. chl., eth.	Citrus plants
25 Indican	C <sub>14</sub> H <sub>17</sub> O <sub>6</sub> N	176-178	-66	s.	s.	sl. s. chl., eth.	<i>Indigofera</i> spp.
26 Iridin	C <sub>24</sub> H <sub>26</sub> O <sub>13</sub> N	208	.....	v. sl. s.	s. h.	i. chl., eth.	Rhizome of <i>Iris</i> spp.
27 Jalapin	C <sub>34</sub> H <sub>56</sub> O <sub>16</sub>	131-150	.....	sl. s.	s.	s. chl., eth.	Scammony resin
28 Khellinin	C <sub>19</sub> H <sub>20</sub> O <sub>10</sub>	175 <sup>1</sup>	0	sl. s.	s. me. al.	i. eth.	Seeds of toothpick ammi
29 Ouabain	C <sub>29</sub> H <sub>44</sub> O <sub>12</sub>	185 d.	-32.5	1.2 g/100 ml	1 g/100 ml	sl. s. chl., eth.	Seeds of <i>Strophanthus gratus</i>
30 Phlorizin	C <sub>21</sub> H <sub>24</sub> O <sub>10</sub>	110 <sup>1</sup>	-52	0.1 g/100 ml	25 g/100 ml	i. chl., eth.	Bark of fruit trees
31 Picrocrocin	C <sub>16</sub> H <sub>26</sub> O <sub>7</sub>	154-156	-50	s.	s.	sl. s. chl., eth.	Crocus
32 Quercitrin	C <sub>21</sub> H <sub>20</sub> O <sub>11</sub>	182-185 <sup>1</sup>	.....	sl. s. h.; i. c.	s.	s. NaOH; i. eth.	Bark of quercitron
33 Rutin	C <sub>27</sub> H <sub>30</sub> O <sub>16</sub>	215 d.	.....	sl. s. c.	sl. s.	i. chl., eth.	Buckwheat plant
34 Salicin	C <sub>13</sub> H <sub>18</sub> O <sub>7</sub>	199-201	-67	4 g/100 ml	1 g/100 ml	i. chl., eth.	Bark of poplar, willow
35 Sarsasaponin	C <sub>45</sub> H <sub>74</sub> O <sub>17</sub>	240	-66	s.	s. h.	sl. s. eth.	Radix of sarsaparilla

/: / Hydrated salt.

# Contrails

## OCCURRENCE, AND USES

benzene; c. = cold; chl. = chloroform; eth. = ether; h. = hot; i. = insoluble; me. al. = methyl alcohol; s. = soluble;

Uses	Aglycone	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Sugar	Reference	
(I)	(J)	(K)	(L)	(M)	(N)	
In alcoholic beverages	.....	.....	.....	Glucose	6,30,50	1
Cathartic; amenorrhea	.....	.....	.....	Pentoses	17	2
No medical use	p-Mandelonitrile	.....	.....	Glucose	18,21,60	3
No medical use	Apigenin	350	.....	Glucose; apiose	26,62,70	4
No medical use	Hydroquinone	170	.....	Glucose	19,32,57	5
Cathartic	Aloe-emodin	224	.....	Glucose	9,48	6
Homeopathic therapy	Bryogenin	.....	.....	Glucose	1,36	7
Indicator; pigment in color photography, paints, bacteriology	Carmine red	.....	.....	.....	17	8
Preparation of vanillin	Coniferyl alcohol	73-74	.....	Glucose	39,40,55	9
Cardiotonic	Strophanthidin	235	+43.1	Rhamnose	13,34,43	10
No medical use	Methylethylacetic acid; tiglic acid	.....	.....	Glucose; rhodose	33	11
Process of algae	Crocetin	285	.....	Gentiobiose	23,27,28	12
Cardiotonic	Strophanthidin	235	+43.1	Cymarose	61	13
Inflammation and vesication of skin; epispastic	7,8-Dihydroxycoumarin	253 d.	.....	Glucose	15,63	14
Cardiotonic	Diginigenin	115	-226	Giginose	44,51	15
Test for cholesterol and some other sterols	Digitogenin	250	-81	Glucose; galactose	16	16
Cardiotonic	Digitoxigenin	253	+19.1	Digitose	64	17
Cardiotonic	Digoxigenin	222	+27	Digitose	53	18
Sunburn protective	Esculetin	270 d.	.....	Glucose	37,58	19
Source of methyl salicylate	Methyl salicylate	-8.6	+1.2	Glucose; xylose	7,46	20
Similar to digitonin	Gitogenin	272	-61	Galactose; xylose	24,25	21
Cardiotonic	Gitoxigenin	235	+38.5	Digitoxose	10,54,65	22
No medical use	Gratiogenin	198	.....	Glucose	17	23
Decrease capillary fragility	3',5,7-Trihydroxy flavanone	390 d.	.....	Glucose	20,52,66	24
.....	Indoxyl	390	.....	Glucose	41,45,47	25
No medical use	Irigenin	186	.....	Glucose	3,4	26
No medical use	Jalapinolic	67-69	.....	Various	42	27
Smooth muscle relaxant	2-Hydroxymethyl-5-methoxy-furanochrome	155	.....	Glucose	56	28
Cardiotonic	Ouabagenin	255	+11.3	Rhamnose	22,44	29
Additive to lubricating oils; induce experimental glycosuria	Phloretin	271	.....	Glucose	8,38,67	30
Coloring; flavoring	Safranin	.....	.....	Glucose	28,31	31
Textile dye	Quercetin	314 d.	.....	Glucose	14,71	32
Decrease capillary fragility	Quercetin	313-314	.....	Glucose; rhamnose	2,11,68,69	33
Analgesic	Saligenin	87	.....	Glucose	12,29	34
Manufacture of pregnane compounds	Sarsapogenin	200	-75	Glucose; rhamnose	35,59	35

*continued*



99. GLYCOSIDES: CHARACTERISTICS,

	Glycoside	Chemical Formula	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Solubility			Occurrence
					H <sub>2</sub> O	Alcohol	Other	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
36	Solanine	C <sub>45</sub> H <sub>73</sub> O <sub>15</sub> N	285 d.	-60	i.	s. h.	i. chl., eth.	<i>Solanum</i> spp.
37	Streptomycin hydrochloride	C <sub>21</sub> H <sub>39</sub> O <sub>12</sub> N <sub>7</sub> ·3HCl	.....	-84	s.	v. sl. s.	i. chl., eth.	Cultures of <i>Streptomyces griseus</i>
38	Tannic acid	C <sub>76</sub> H <sub>52</sub> O <sub>46</sub>	210-215 d.	.....	v. s.	sl. s.	v. sl. s. chl., eth.	Bark of oak, sumac

Contributors: (a) Hamerslag, Frank E., (b) Calesnick, Benjamin

References: [1] Angeletti, A., and D. Ponte. 1934. Gazz. Chim. Ital. 64:569. [2] Attree, G. F., and A. G. Perkin. p. 152. [5] Bell, R. C., and L. H. Briggs. 1942. Ibid., p. 1. [6] Bourcet, P. 1898. Bull. Soc. Chim. France, 1933. Bull. Soc. Chim. Biol. 15:531. [9] Cahn, R. S., and J. L. Simonsen. 1932. J. Chem. Soc., p. 2573. [10] Cloetta, Regl. Res. Lab. AIC-114. [12] Evans, W. E., Jr., H. K. Iwamoto, and J. C. Krantz, Jr. 1945. J. Am. Pharm. [14] Freudenberg, K., and E. Vollbrecht. 1922. Ann. Chem. 429:303. [15] Gandini, A. 1941. Chem. Abstr. 35:1394. Laboratories Research Division, Radnor, Penna., 1954. [18] Haworth, W. N., and B. Wylam. 1923. J. Chem. Pharm. Assoc. Sci. Ed. 30:629. [21] Hudson, C. S. 1924. J. Am. Chem. Soc. 46:483. [22] Jacobs, W. A., and [24] Kiliani, H. 1916. Ber. Deut. Chem. Ges. 49:701. [25] Kiliani, H. 1918. Ibid. 51:1613. [26] Klein, G. 1932. [28] Kuhn, R., and A. Winterstein. 1934. Ber. Deut. Chem. Ges. 67B:344. [29] Kunz, A. 1926. J. Am. Chem. [32] Mannich, C. 1912. Arch. Pharm. 250:547. [33] Mannich, C., and P. Schumann. 1938. Ibid. 276:211. 1940. J. Am. Chem. Soc. 62:3349. [36] Masson, A. 1893. Bull. Soc. Chim. France, Ser. 3, 9:1054. [37] Merz, Soc., p. 1170. [39] Patterson, R. F., and H. Hibbert. 1943. J. Am. Chem. Soc. 65:1862. [40] Pauly, H., and K. 91:1715. [42] Power, F. B., and H. Rogerson. 1912. Ibid. 101:398. [43] Reichstein, T., and A. Katz. 1943. A. 1927. J. Chem. Soc., p. 1937. [46] Robertson, A., and R. B. Waters. 1931. Ibid., p. 1881. [47] Robertson, E. Bugie, and S. A. Waksman. 1944. Proc. Soc. Exptl. Biol. Med. 55:66. [50] Schmutz, J., and T. Reichstein. [52] Sieburg, E. 1913. Arch. Pharm. 251:154. [53] Smith, S. 1930. J. Chem. Soc., p. 508. [54] Smith, S. 1931. Chem. Ges. 74B:1549. [57] Tschitschibabin, A. E., A. W. Kirssanov, and M. G. Rudenko. 1930. Ann. Chem. 48:726. [60] Viehoveer, A., and H. Mack. 1935. Am. J. Pharm. 107:397. [61] Von Euw, J., and T. Reichstein. K. Sturm. 1930. Ber. Deut. Chem. Ges. 63B:1299. [64] Windaus, A., and C. Freese. 1925. Ibid. 58B:2503. [67] Zemplen, G., R. Bognar, and I. Szekely. 1943. Ibid. 76B:386. [68] Zemplen, G., and A. Gerecs. 1935. Ibid. 76B:776. [71] Zemplen, G., et al. 1928. Ibid. 61B:2486.

100. FATTY ACIDS: PHYSICAL AND

Boiling Point (column F): d. = decomposes. Solubility (column K): a. = acid; acet. = acetone; ac. = acetic; hex. = hexane; h. = hot; me. = methyl; pent. = pentane; pet. = petroleum; pyr. = pyridine; s. = soluble; sl. = slightly;

Acid		Chemical Formula	Molecular Weight	Melting Point °C	Boiling Point °C/mm <sup>1</sup>	Specific Gravity <sup>2</sup>	
Systematic Name	Common Name						
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Saturated Fatty Acids							
1	Methanoic	Formic	HCOOH	46.0	8.4	100.5	1.220 <sup>20°</sup>
2	Ethanoic	Acetic	CH <sub>3</sub> COOH	60.1	16.7	118.2	1.049 <sup>20°</sup>
3	Propanoic	Propionic	C <sub>2</sub> H <sub>5</sub> COOH	74.1	-22.0	141.1	0.992 <sup>20°</sup>

<sup>1</sup>/ 760 mm of mercury (atmospheric pressure), unless otherwise specified. <sup>2</sup>/ At temperature indicated in superscript. <sup>3</sup>/ Milligrams KOH required to neutralize one gram of acid. <sup>4</sup>/ Grams of iodine absorbed by 100

**OCCURRENCE, AND USES**

Uses	Aglycone	Melting Point °C	Specific Rotation [α] <sub>D</sub>	Sugar	Reference	
(I)	(J)	(K)	(L)	(M)	(N)	
No medical use	Solanidine	219	-29	Glucose; galactose; rhamnose	5,44	36
Tuberculosis; susceptible gram-negative bacteria; leprosy; granuloma inguinale	Streptidine	.....	.....	L-Streptose	49	37
Tanning; mordant in dyeing and printing; topically as astringent and styptic	Gallic acid	235 d.	.....	Glucose	17	38

1927. J. Chem. Soc., p. 234. [3] Baker, W. 1928. Ibid., p. 1022. [4] Baker, W., and R. Robinson. 1929. Ibid., Ser. 3, 19:537. [7] Bridel, M. M., and S. Grillon. 1928. Compt. Rend. 187:609. [8] Bridel, M. M., and A. Kramer. M. 1926. Arch. Exptl. Pathol. Pharmacol. 112:261. [11] Eskew, R. K. 1946. U.S. Bur. Agr. Ind. Chem. Eastern Assoc., Sci. Ed., 34:207. [13] Fieser, L. F., and R. P. Jacobsen. 1937. J. Am. Chem. Soc. 59:2335. [16] Gisvold, O. 1934. J. Am. Pharm. Assoc., Sci. Ed., 23:664. [17] Hamerslag, F. E. Unpublished. Wyeth Soc. 123:3120. [19] Helferich, B., and W. Reischel. 1938. Ann. Chem. 533:278. [20] Higby, R. H. 1941. J. Am. N. M. Bigelow. 1932. J. Biol. Chem. 96:647. [23] Karrer, P., and A. Helfenstein. 1930. Helv. Chim. Acta 13:392. Handbuch der Pflanzenanalyse. J. Springer, Wien. Bd. 3(2), S. 858. [27] Kühn, A. 1940. Angew. Chem. 53:1. Soc. 48:262. [30] Ludwig, H. 1861. Arch. Pharm. 158:129. [31] Lutz, H. E. W. 1930. Biochem. Z. 226:97. [34] Mannich, C., and G. Siewert. 1942. Ber. Deut. Chem. Ges. 75B:737. [35] Marker, R. E., and J. Krueger. K. W., and W. Hagemann. 1941. Naturwissenschaften 29:650. [38] Müller, A., and A. Robertson. 1933. J. Chem. Feuerstein. 1927. Ber. Deut. Chem. Ges. 60B:1031. [41] Perkin, A. G., and W. P. Bloxam. 1907. J. Chem. Soc. Pharm. Acta Helv. 18:521. [44] Reichstein, T., and H. Reich. 1946. Ann. Rev. Biochem. 15:155. [45] Robertson, A., and R. B. Waters. 1933. Ibid., p. 30. [48] Rosenthaler, L. 1932. Arch. Pharm. 270:214. [49] Schatz, A., 1947. Pharm. Acta Helv. 22:167. [51] Shoppee, C. W., and T. Reichstein. 1940. Helv. Chim. Acta 23:975. Ibid., p. 23. [55] Solntsev, A. A. 1944. Chem. Abstr. 38:3780. [56] Späth, E., and W. Gruber. 1941. Ber. Deut. 479:303. [58] Tunmann, O. 1916. Chem. Zentr. 87(I):1277. [59] Van der Haar, A. W. 1929. Rec. Trav. Chim. 1948. Helv. Chim. Acta 31:883. [62] Vongerichten, E. 1901. Ann. Chem. 318:121. [63] Wessely, F., and [65] Windaus, A., and G. Schwarte. 1925. Ibid. 58B:1515. [66] Zemplen, G., and R. Bogнар. 1942. Ibid. 75B:1043. Ibid. 68B:1318. [69] Zemplen, G., and A. Gerecs. 1938. Ibid. 71B:2520. [70] Zemplen, G., and L. Mester. 1943.

**CHEMICAL CHARACTERISTICS**

al. = alcohol; bz. = benzene; chl. = chloroform; cyc. = cyclohexane; eth. = ether; glac. = glacial; hept. = heptane; tol. = toluene; v. = very; w. = water.

Refractive Index <sup>3</sup> $n_{\frac{OC}{D}}$	Neutralization Value <sup>4</sup>	Iodine Value (Calculated) <sup>5</sup>	Solubility	Source	Reference	
(H)	(I)	(J)	(K)	(L)	(M)	
Saturated Fatty Acids						
1.3714 <sup>20°</sup>	1,219	.....	s.w.	Red ant	13,28,29	1
1.3718 <sup>20°</sup>	934.2	.....	s.w.	Vinegar	15,28,29	2
1.3874 <sup>20°</sup>	757.3	.....	s.al., chl., eth., w.	Milk and milk products	28,29	3

superscript, referred to water at 4°C. /s/ Refractive index (n) is given for the sodium D-line at temperature shown grams of acid.

*continued*

100. FATTY ACIDS: PHYSICAL AND

Acid		Chemical Formula	Molecular Weight	Melting Point °C	Boiling Point °C/mm <sup>1</sup>	Specific Gravity <sup>2</sup>	
Systematic Name	Common Name						
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
<b>Saturated Fatty Acids</b>							
4	Butanoic	Butyric	C <sub>3</sub> H <sub>7</sub> COOH	88.1	-7.9	163.5	0.9587 <sup>200</sup>
5	Pentanoic	Valeric	C <sub>4</sub> H <sub>9</sub> COOH	102.1	-34.5	187	0.9422 <sup>00</sup>
6	Hexanoic	Caproic	C <sub>5</sub> H <sub>11</sub> COOH	116.2	-3.4	205.8	0.9292 <sup>00</sup>
7	Heptanoic	Heptylic <sup>a</sup>	C <sub>6</sub> H <sub>13</sub> COOH	130.2	-10.5	223.0	0.92215 <sup>200</sup>
8	Octanoic	Caprylic	C <sub>7</sub> H <sub>15</sub> COOH	144.2	16.7	239.7	0.9102 <sup>00</sup>
9	Nonanoic	Pelargonic	C <sub>8</sub> H <sub>17</sub> COOH	158.2	12.5	255.6	0.9072 <sup>00</sup>
10	Decanoic	Capric	C <sub>9</sub> H <sub>19</sub> COOH	172.3	31.6	270	0.8858 <sup>400</sup>
11	Undecanoic <sup>7</sup>	Undecylic	C <sub>10</sub> H <sub>21</sub> COOH	186.3	29.3	284	0.9905 <sup>250</sup>
12	Dodecanoic	Lauric	C <sub>11</sub> H <sub>23</sub> COOH	200.3	44.2	225/100	0.8690 <sup>500</sup>
13	Tridecanoic	Tridecylic	C <sub>12</sub> H <sub>25</sub> COOH	214.3	41.5	236/100	0.8458 <sup>800</sup>
14	Tetradecanoic	Myristic	C <sub>13</sub> H <sub>27</sub> COOH	228.4	53.9	250/100	0.8622 <sup>540</sup>
15	Pentadecanoic	Pentadecylic	C <sub>14</sub> H <sub>29</sub> COOH	242.2	52.3	202.5/10	0.8423 <sup>800</sup>
16	Hexadecanoic	Palmitic	C <sub>15</sub> H <sub>31</sub> COOH	256.4	63.1	268/100	0.8487 <sup>700</sup>
17	Heptadecanoic	Margaric	C <sub>16</sub> H <sub>33</sub> COOH	270.4	61.3	220/10	0.8536 <sup>00</sup>
18	Octadecanoic	Stearic	C <sub>17</sub> H <sub>35</sub> COOH	284.5	69.6	213/5	0.8390 <sup>800</sup>
19	Nonadecanoic	Nonadecylic	C <sub>18</sub> H <sub>37</sub> COOH	298.5	68.6	299/10	0.8771 <sup>240</sup>
20	Eicosanoic	Arachidic	C <sub>19</sub> H <sub>39</sub> COOH	312.5	76.5	204/1	0.8240 <sup>1000</sup>
21	Docosanoic	Behenic	C <sub>21</sub> H <sub>43</sub> COOH	340.6	81.5	306/60	0.8221 <sup>1000</sup>
22	Tetradecanoic	Lignoceric	C <sub>23</sub> H <sub>47</sub> COOH	368.6	86.0	272/10	0.8207 <sup>1000</sup>
23	Hexacosanoic	Cerotic	C <sub>25</sub> H <sub>51</sub> COOH	396.7	88.5	.....	0.8198 <sup>1000</sup>
24	Octacosanoic	Montanic	C <sub>27</sub> H <sub>55</sub> COOH	424.7	90.9	.....	0.8191 <sup>1000</sup>
25	Triacantanoic	Melissic	C <sub>29</sub> H <sub>59</sub> COOH	452.8	93.6	.....	.....
26	Dotriacontanoic	Lacceroic	C <sub>31</sub> H <sub>63</sub> COOH	480.0	96.2	.....	.....
27	Tetracontanoic	Gheddic	C <sub>33</sub> H <sub>67</sub> COOH	508.9	98.4	.....	.....
28	Pentatriacontanoic	Ceroplastic	C <sub>34</sub> H <sub>69</sub> COOH	522.9	98.4	.....	.....
<b>Unsaturated Fatty Acids (Monoethenoic)</b>							
29	<i>trans</i> -2-Butenoic	Crotonic	C <sub>4</sub> H <sub>6</sub> O <sub>2</sub>	86.1	72	189.0	0.9648 <sup>00</sup>
30	<i>cis</i> -2-Butenoic	Isocrotonic	C <sub>4</sub> H <sub>6</sub> O <sub>2</sub>	86.1	15.5	169.3	1.0312 <sup>150</sup>
31	2-Hexenoic	Isohydroisobic	C <sub>6</sub> H <sub>10</sub> O <sub>2</sub>	114.1	32	217	0.9652 <sup>00</sup>
32	4-Decenoic	Obtusilic	C <sub>10</sub> H <sub>18</sub> O <sub>2</sub>	170.2	.....	149/13	0.9197 <sup>200</sup>
33	9-Decenoic	Caproleic	C <sub>10</sub> H <sub>18</sub> O <sub>2</sub>	170.2	.....	142/4	0.9238 <sup>150</sup>
34	4-Dodecenoic	Linderic	C <sub>12</sub> H <sub>22</sub> O <sub>2</sub>	198.3	1.0-1.3	171/13	0.9081 <sup>200</sup>
35	5-Dodecenoic	Denticetic	C <sub>12</sub> H <sub>22</sub> O <sub>2</sub>	198.3	.....	.....	0.9130 <sup>150</sup>
36	9-Dodecenoic	Lauroleic	C <sub>12</sub> H <sub>22</sub> O <sub>2</sub>	198.3	.....	142/4	.....
37	4-Tetradecenoic	Tsuzuic	C <sub>14</sub> H <sub>26</sub> O <sub>2</sub>	226.4	18.0-18.5	185-188/13	0.9024 <sup>200</sup>
38	5-Tetradecenoic	Physeteric	C <sub>14</sub> H <sub>26</sub> O <sub>2</sub>	226.4	.....	190-195/15	0.9046 <sup>200</sup>
39	9-Tetradecenoic	Myristoleic	C <sub>14</sub> H <sub>26</sub> O <sub>2</sub>	226.4	-4	.....	0.9018 <sup>200</sup>
40	9-Hexadecenoic	Palmitoleic	C <sub>16</sub> H <sub>30</sub> O <sub>2</sub>	254.4	-0.5 to +0.5	131/0.06	.....
41	6-Octadecenoic	Petroselinic	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.5	32-33	237.5/18	0.8824 <sup>350</sup>
42	<i>cis</i> -9-Octadecenoic	Oleic	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.5	13.4(α), 16.3(β)	234/15	0.8905 <sup>200</sup>
43	<i>trans</i> -9-Octadecenoic	Elaidic	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.5	44.5	288/100	0.8517 <sup>90</sup>
44	<i>trans</i> -11-Octadecenoic	Vaccenic	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.5	44	.....	0.8563 <sup>700</sup>
45	9-Eicosenoic	Gadoleic	C <sub>20</sub> H <sub>38</sub> O <sub>2</sub>	310.5	24-24.5	220/6	0.8882 <sup>250</sup>
46	11-Eicosenoic	Gondoic	C <sub>20</sub> H <sub>38</sub> O <sub>2</sub>	310.5	23.5-24	267/15	.....
47	11-Docosenoic	Cetoleic	C <sub>22</sub> H <sub>42</sub> O <sub>2</sub>	338.6	32.5-33	.....	.....
48	13-Docosenoic	Erucic	C <sub>22</sub> H <sub>42</sub> O <sub>2</sub>	338.6	34.7	242/5	0.85321 <sup>700</sup>
49	15-Tetracosenoic	Nervonic <sup>a</sup>	C <sub>24</sub> H <sub>46</sub> O <sub>2</sub>	366.6	42.5-43.0	.....	.....
50	17-Hexacosenoic	Ximenic	C <sub>26</sub> H <sub>50</sub> O <sub>2</sub>	394.7	45-45.5	.....	.....
51	21-Triacontenoic	Lumequeic	C <sub>30</sub> H <sub>58</sub> O <sub>2</sub>	450.8	.....	.....	.....
<b>Unsaturated Fatty Acids (Dienuic)</b>							
52	2,4-Pentadienoic	β-Vinylacrylic	C <sub>5</sub> H <sub>6</sub> O <sub>2</sub>	98.1	80	110 d.	.....
53	2,4-Hexadienoic	Sorbic	C <sub>6</sub> H <sub>8</sub> O <sub>2</sub>	112.1	134.5	228 d.	.....
54	2,4-Decadienoic	Stillingic	C <sub>10</sub> H <sub>16</sub> O <sub>2</sub>	168.2	.....	.....	.....
55	2,4-Dodecadienoic	.....	C <sub>12</sub> H <sub>20</sub> O <sub>2</sub>	196.3	.....	.....	.....
56	9,12-Hexadecadienoic	.....	C <sub>16</sub> H <sub>28</sub> O <sub>2</sub>	252.4	.....	.....	.....
57	<i>cis</i> -9, <i>cis</i> -12-Octadecadienoic	α-Linoleic	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	280.5	-5.2 to -5.0	202/1.4	0.9038 <sup>180</sup>

<sup>1</sup>/ 760 mm of mercury (atmospheric pressure), unless otherwise specified. <sup>2</sup>/ At temperature indicated in superscript. <sup>3</sup>/ Milligrams KOH required to neutralize one gram of acid. <sup>4</sup>/ Grams of iodine absorbed by 100



## CHEMICAL CHARACTERISTICS

Refractive Index <sup>a</sup> $n_{D}^{20^{\circ}C}$	Neutralization Value <sup>d</sup>	Iodine Value (Calculated) <sup>e</sup>	Solubility	Source	Reference	
(H)	(I)	(J)	(K)	(L)	(M)	
<b>Saturated Fatty Acids</b>						
1.33906 <sup>200</sup>	636.8	.....	s.al., eth., w.	Butterfat	13,29,36	4
1.4086 <sup>200</sup>	549.3	.....	s.al., eth.; sl.s.w.	Essential oils	29,36	5
1.41635 <sup>200</sup>	483.0	.....	s.al., eth.; sl.s.w.	Butterfat, palm oils	29,36	6
1.42302 <sup>00</sup>	431.0	.....	s.al., eth.; v.sl.s.w.	Violet-leaf oil	29,36	7
1.42852 <sup>00</sup>	389.1	.....	s.al., bz., eth.; v.sl.s.w.	Butterfat, palm-kernel oils	13,29,36	8
1.4322 <sup>200</sup>	354.6	.....	s.al., chl., eth.; v.sl.s.w.	Butterfat, hair fat	29,36	9
1.42855 <sup>400</sup>	325.7	.....	s.al., eth., pet.eth.; v.sl.s.w.	Butterfat, palm-kernel oils	13,29,36	10
1.42027 <sup>00</sup>	301.2	.....	s.al., chl., eth., pet.eth.	Hair fat (human)	29,36	11
1.42616 <sup>00</sup>	280.1	.....	s.acet., al., eth., pet.eth.	Lauraceae oils	13,29,36	12
1.42866 <sup>00</sup>	261.8	.....	s.acet., al., eth., pet.eth.	Hair fat (human)	14,29,31,36	13
1.42737 <sup>00</sup>	245.7	.....	s.acet., al., eth., pet.eth.	Myristicaceae fats	13,29,36	14
1.42927 <sup>00</sup>	231.5	.....	s.acet., al., eth., pet.eth.	Mutton, hair, & milk fats	29,36	15
1.43097 <sup>00</sup>	218.8	.....	s.acet., h.al., eth., pet.eth.	Palm-pulp oils	29,36	16
1.43247 <sup>00</sup>	207.5	.....	s.acet., h.al., eth., pet.eth.	Hair & mutton fats	29,36	17
1.43377 <sup>00</sup>	197.2	.....	s.acet., h.al., eth., pet.eth.	Animal fats generally	13,29,36	18
1.45122 <sup>50</sup>	188.0	.....	s.acet., h.al., eth., pet.eth.	Ox fat	29	19
1.42501 <sup>000</sup>	179.5	.....	s.bz., chl., eth., pet.eth.	Peanut oil, rambutan fat	13,29,36	20
1.42701 <sup>000</sup>	164.7	.....	sl.s.al., eth.	Moringa (ben) oils	13,29,36	21
1.42871 <sup>000</sup>	152.2	.....	s.ac.a., bz., CS <sub>2</sub> , eth.	Beech-tar paraffin	28,29,36,39	22
1.43011 <sup>000</sup>	141.4	.....	s.h.acet., h.chl., h.me.al.	Insect, wool, & leaf waxes	13,29,36	23
1.43131 <sup>000</sup>	132.1	.....	s.h.ac.a., h.bz., h.me.al.	Insect, leaf, & montan waxes	13,29,36	24
1.43231 <sup>000</sup>	123.9	.....	s.chl., CS <sub>2</sub> , h.me.al.	Insect & mineral waxes	13,29,36	25
.....	116.7	.....	s.h.acet., h.bz., chl.	Stick-lac wax	13,17,29,36	26
.....	110.2	.....	s.h.acet., h.bz., chl.	Ghedda wax	13,29,32,36	27
.....	107.3	.....	s.h.acet., h.bz., chl.	<i>Ceroplastes rubens</i>	13,29,32,36	28
<b>Unsaturated Fatty Acids (Monoethenoic)</b>						
1.42288 <sup>000</sup>	651.7	294.9	s.acet., al., tol., w.	Croton oil	29	29
1.44572 <sup>000</sup>	651.7	294.9	s.al., pet.eth., w.	Croton oil	29	30
1.44604 <sup>000</sup>	491.5	222.5	s.CS <sub>2</sub> , eth.	Japanese mint oils	29,36	31
1.44972 <sup>000</sup>	329.6	149.1	s.bz., eth.	<i>Lindera</i> seed oils	2,13,29,36	32
1.45071 <sup>500</sup>	329.6	149.1	s.al., eth.	Milk fats, whale oil	13,29,36,37	33
1.45292 <sup>000</sup>	282.9	128.0	s.bz., chl., eth.	<i>Lindera</i> seed oils	13,29,36	34
1.45351 <sup>500</sup>	282.9	128.0	s.bz., chl., eth.	Herring & whale oils	13,29,36,39	35
.....	282.9	128.0	s.bz., chl., eth.	Milk fats, cochineal wax	13,29,36	36
1.45572 <sup>000</sup>	247.9	112.2	s.bz., pet.eth.	<i>Lindera</i> seed oils	2,13,29,36	37
1.45522 <sup>000</sup>	247.9	112.2	s.bz., eth., pet.eth.	Whale & fish oils	13,29,36,39	38
1.45192 <sup>000</sup>	247.9	112.2	s.bz., eth., pet.eth.	Milk fats, whale oil	2,13,29,36	39
.....	220.5	99.8	s.bz., eth., pet.eth.	Marine oils, milk fats	2,13,18,29	40
1.45334 <sup>000</sup>	198.6	89.9	s.al., eth., pet.eth.	Parsley-seed oil	13,21,22,29, 36	41
1.45823 <sup>200</sup>	198.6	89.9	s.acet., eth., me.al.	Olive oil, pork fat	13,29,36	42
1.44685 <sup>000</sup>	198.6	89.9	s.al., chl., eth., pet.eth.	Beef & sheep fats	29	43
1.44067 <sup>000</sup>	198.6	89.9	s.acet., me.al.	Milk, beef, & sheep fats	13,29,36	44
1.45972 <sup>500</sup>	180.7	81.8	s.acet., me.al., pet.eth.	Sperm & fish-liver oils	13,16,21,29	45
.....	180.7	81.8	s.al., me.al.	Crucifer, jojoba, & fish oils	13,16,29	46
.....	165.7	75.0	s.al.	Marine oils	13,21,29,39	47
1.44447 <sup>000</sup>	165.7	75.0	v.s.eth., me.al.	Crucifer oils	13,29,30,36	48
.....	153.0	69.2	s.acet., al., eth.	Shark-liver oil, cerebrosides	13,21,29,36	49
.....	142.2	64.3	s.bz., chl., eth., pet.eth.	<i>Ximenia</i> oils	13,29	50
.....	124.5	56.3	s.bz., chl., eth., pet.eth.	<i>Ximenia</i> oils	13,29	51
<b>Unsaturated Fatty Acids (Dienoic)</b>						
.....	572.0	517.5	v.s.al., eth.; s.h.w.	Synthetic	29,36	52
.....	500.4	452.7	s.al., eth.; sl.s.w.	Mountain-ash berry	29,36	53
.....	333.5	301.7	s.acet., eth., hex.	<i>Stillingia</i> oils	29	54
.....	285.8	258.6	s.acet., eth., pet.eth.	<i>Sebastiania fruticosa</i> seed oil	29	55
.....	222.3	201.1	s.acet., eth., pet.eth.	<i>Asclepias syriaca</i> seed oil	7	56
1.46992 <sup>000</sup>	200.1	181.0	s.acet., al., eth., pet.eth.	Numerous seed oils	13,29,36	57

superscript, referred to water at 4°C. /a/ Refractive index (n) is given for the sodium D-line at temperature shown grams of acid. /s/ Also called enanthic acid. /r/ Also called hendecanoic acid. /e/ Also called selacholeic acid.

continued



Acid		Chemical Formula	Molecular Weight	Melting Point °C	Boiling Point °C/mm <sup>1</sup>	Specific Gravity <sup>2</sup>
Systematic Name	Common Name					
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Unsaturated Fatty Acids (Dienoic)						
58	<i>trans</i> -9, <i>trans</i> -12-Octadecadienoic	Linolelaidic	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	280.5	28-29	.....
59	<i>trans</i> -10, <i>trans</i> -12-Octadecadienoic	.....	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	280.5	55.5-56	.....
60	11,14-Eicosadienoic	.....	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	308.4	.....	.....
61	13,16-Docosadienoic	.....	C <sub>22</sub> H <sub>40</sub> O <sub>2</sub>	336.6	.....	.....
62	17,20-Hexacosadienoic	.....	C <sub>26</sub> H <sub>48</sub> O <sub>2</sub>	392.7	61	.....
Unsaturated Fatty Acids (Trienoic)						
63	6,10,14-Hexadecatrienoic	Hiragonic	C <sub>16</sub> H <sub>26</sub> O <sub>2</sub>	250.4	.....	180-190/15 0.9296 <sup>200</sup>
64	7,10,13-Hexadecatrienoic	.....	C <sub>16</sub> H <sub>26</sub> O <sub>2</sub>	250.4	.....	.....
65	<i>cis</i> -6, <i>cis</i> -9, <i>cis</i> -12-Octadecatrienoic	γ-Linolenic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	.....	.....
66	<i>trans</i> -8, <i>trans</i> -10, <i>cis</i> -12-Octadecatrienoic	α-Calendic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	40-40.5	.....
67	<i>trans</i> -8, <i>trans</i> -10, <i>trans</i> -12-Octadecatrienoic	β-Calendic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	77-78	.....
68	<i>cis</i> -8, <i>trans</i> -10, <i>cis</i> -12-Octadecatrienoic	.....	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	.....	.....
69	<i>cis</i> -9, <i>cis</i> -12, <i>cis</i> -15-Octadecatrienoic	α-Linolenic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	-10 to -11.3	157/0.001 0.914 <sup>200</sup>
70	<i>trans</i> -9, <i>trans</i> -12, <i>trans</i> -15-Octadecatrienoic	Linolenelaidic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	29-30	.....
71	<i>cis</i> -9, <i>trans</i> -11, <i>trans</i> -13-Octadecatrienoic	α-Eleostearic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	48-49	235/15
72	<i>trans</i> -9, <i>trans</i> -11, <i>trans</i> -13-Octadecatrienoic	β-Eleostearic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	71.5	.....
73	<i>cis</i> -9, <i>trans</i> -11, <i>cis</i> -13-Octadecatrienoic	Punicic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	43.5-44	..... 0.9027 <sup>500</sup>
74	<i>trans</i> -9, <i>trans</i> -11, <i>trans</i> -13-Octadecatrienoic	.....	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.4	.....	.....
75	5,8,11-Eicosatrienoic	.....	C <sub>20</sub> H <sub>34</sub> O <sub>2</sub>	306.5	.....	.....
76	8,11,14-Eicosatrienoic	.....	C <sub>20</sub> H <sub>34</sub> O <sub>2</sub>	306.5	.....	.....
Unsaturated Fatty Acids (Tetraenoic)						
77	4,8,11,14-Hexadecatetraenoic	.....	C <sub>16</sub> H <sub>24</sub> O <sub>2</sub>	248.4	.....	.....
78	6,9,12,15-Hexadecatetraenoic	.....	C <sub>16</sub> H <sub>24</sub> O <sub>2</sub>	248.4	.....	.....
79	4,8,12,15-Octadecatetraenoic	Moroctic	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276.4	.....	208-213/15 0.9297 <sup>200</sup>
80	6,9,12,15-Octadecatetraenoic	.....	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276.4	-57.4 to -56.6	.....
81	9,11,13,15-Octadecatetraenoic	α-Parinaric	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276.4	85-86	.....
82	9,11,13,15-Octadecatetraenoic	β-Parinaric	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276.4	95-96	.....
83	9,12,15,18-Octadecatetraenoic	.....	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276.4	.....	.....
84	4,8,12,16-Eicosatetraenoic	.....	C <sub>20</sub> H <sub>32</sub> O <sub>2</sub>	304.5	.....	217-220/10 0.9263 <sup>200</sup>
85	5,8,11,14-Eicosatetraenoic	Arachidonic	C <sub>20</sub> H <sub>32</sub> O <sub>2</sub>	304.5	-49.5	163/1 0.9082 <sup>200</sup>
86	6,10,14,18-Eicosatetraenoic ?	.....	C <sub>20</sub> H <sub>32</sub> O <sub>2</sub>	304.5	.....	..... 0.9263 <sup>200</sup>
87	4,7,10,13-Docosatetraenoic	.....	C <sub>22</sub> H <sub>36</sub> O <sub>2</sub>	332.5	.....	.....
88	7,10,13,16-Docosatetraenoic	.....	C <sub>22</sub> H <sub>36</sub> O <sub>2</sub>	332.5	.....	.....
89	8,12,16,19-Docosatetraenoic	.....	C <sub>22</sub> H <sub>36</sub> O <sub>2</sub>	332.5	.....	.....
Unsaturated Fatty Acids (Penta- and Hexa-enoic)						
90	4,8,12,15,18-Eicosapentaenoic	Timnodonic ?	C <sub>20</sub> H <sub>30</sub> O <sub>2</sub>	302.5	.....	..... 0.9399 <sup>150</sup>
91	5,8,11,14,17-Eicosapentaenoic	.....	C <sub>20</sub> H <sub>30</sub> O <sub>2</sub>	302.5	-54.4 to -53.8	.....
92	4,7,10,13,16-Docosapentaenoic	.....	C <sub>22</sub> H <sub>34</sub> O <sub>2</sub>	330.5	.....	.....
93	4,8,12,15,19-Docosapentaenoic	Clupanodonic	C <sub>22</sub> H <sub>34</sub> O <sub>2</sub>	330.5	.....	207-212/2 0.9356 <sup>200</sup>
94	7,10,13,16,19-Docosapentaenoic	.....	C <sub>22</sub> H <sub>34</sub> O <sub>2</sub>	330.5	.....	.....
95	4,7,10,13,16,19-Docosahexaenoic	.....	C <sub>22</sub> H <sub>32</sub> O <sub>2</sub>	328.5	-44.5 to -44.1	.....
96	4,8,12,15,18,21-Tetracosahexaenoic	Nisinic	C <sub>24</sub> H <sub>36</sub> O <sub>2</sub>	356.6	.....	..... 0.9452 <sup>200</sup>

<sup>1</sup>/ 760 mm of mercury (atmospheric pressure), unless otherwise specified. <sup>2</sup>/ At temperature indicated in superscript. <sup>3</sup>/ Milligrams KOH required to neutralize one gram of acid. <sup>5</sup>/ Grams of iodine absorbed by 100

## CHEMICAL CHARACTERISTICS

Refractive Index <sup>3</sup> $n_D^{20}$	Neutralization Value <sup>4</sup>	Iodine Value (Calculated) <sup>5</sup>	Solubility	Source	Reference	
(H)	(I)	(J)	(K)	(L)	(M)	
<b>Unsaturated Fatty Acids (Dienoic)</b>						
.....	200.1	181.0	s.al., eth., me.al., pet.eth.	Isomerized $\alpha$ -acid	29,36	58
.....	200.1	181.0	s.acet., cyc., eth.	<i>Chilopsis linearis</i> seed oil	23	59
.....	181.9	164.5	s.acet., eth., pet.eth.	Shark-liver oil	29	60
.....	166.7	150.8	s.acet., eth.	<i>Brassica</i> seed oils	29	61
.....	142.9	129.3	s.eth., pet.eth.	<i>Sphaciospongia</i> sponge	29	62
<b>Unsaturated Fatty Acids (Trienoic)</b>						
1.4850 <sup>50°</sup>	224.1	304.1	s.al., eth.	Sardine oil	13,29,36,38	63
.....	224.1	304.1	s.al., eth.	<i>Brassica napus</i> leaves	29	64
.....	201.5	273.5	s.acet., eth., me.al.	<i>Oenothera biennis</i> seed oil	29,36	65
.....	201.5	273.5	s.acet., pent.	<i>Calendula officinalis</i> seed oil	8,29,36	66
.....	201.5	273.5	s.me.al., pet.eth.	Elaidinized $\alpha$ -acid	8	67
.....	201.5	273.5	v.s.acet., al., pent., pet.eth.	<i>Jacaranda</i> oils	10,29	68
1.4678 <sup>50°</sup>	201.5	273.5	s.acet., al., eth., pet.eth.	Linseed, perilla, & hemp oils	13,29,36	69
.....	201.5	273.5	s.me.al., pet.eth.	Elaidinized $\alpha$ -acid	29,36	70
1.5112 <sup>50°</sup>	201.5	273.5	s.al., cyc., eth., pet.eth.	Tung, po-yak, & neou oils	13,28,29,36	71
1.5002 <sup>75°</sup>	201.5	273.5	s.al., eth., me.al., pet.eth.	Elaidinized $\alpha$ -acid	13,29,36	72
1.5114 <sup>50°</sup>	201.5	273.5	s.al., pent., pet.eth.	<i>Trichosanthes</i> oils	13,29,36	73
.....	201.5	273.5	s.acet., al., CS <sub>2</sub> , pent.	<i>Catalpa ovata</i> seed oil	9,29	74
.....	183.1	248.3	s.CS <sub>2</sub> , hept., me.al.	Bovine-liver phosphatides	29	75
.....	183.1	248.3	s.CS <sub>2</sub> , hept., me.al.	Bovine-liver phosphatides	29	76
<b>Unsaturated Fatty Acids (Tetraenoic)</b>						
.....	225.9	408.8	s.acet., al., eth., pet.eth.	Sardine oil	29	77
1.4870 <sup>29°</sup>	225.9	408.8	s.acet., al., CS <sub>2</sub> , eth., pent.	Pilchard & herring oils	29	78
1.4911 <sup>20°</sup>	203.0	367.3	s.acet., al., eth., pet.eth.	Sardine oil	13,29,36,38	79
1.4888 <sup>16°</sup>	203.0	367.3	s.CS <sub>2</sub> , me.al.	Pilchard & herring oils	29	80
.....	203.0	367.3	s.acet., al., eth., pet.eth.	<i>Parinarium &amp; Impatiens</i> seed oils	13,29,36	81
.....	203.0	367.3	s.eth., pet.eth.	Elaidinized $\alpha$ -acid	13,29,36	82
.....	203.0	367.3	s.CS <sub>2</sub> , me.al.	Herring oil	29	83
1.4915 <sup>20°</sup>	184.3	333.4	s.acet., eth.	Fish & whale oils	13,29,36	84
1.4824 <sup>20°</sup>	184.3	333.4	s.acet., eth., me.al., pet.eth.	Brain, liver, egg, & glandular lipids	13,29,36	85
1.4935 <sup>20°</sup>	184.3	333.4	s.acet., me.al., pet.eth.	Sardine oil	29	86
.....	168.7	305.4	s.acet., me.al., pet.eth.	Brain phosphatides	24,29	87
.....	168.7	305.4	s.CS <sub>2</sub> , hept., me.al.	Brain & bovine-liver phosphatides	24,29	88
.....	168.7	305.4	s.acet., me.al., pet.eth.	Shark-liver oil	1,29	89
<b>Unsaturated Fatty Acids (Penta- and Hexa-enoic)</b>						
1.5109 <sup>15°</sup>	185.5	419.6	s.bz., chl., eth., pet.eth.	Sardine & bonito oils	13,29,36	90
1.4977 <sup>23°</sup>	185.5	419.6	s.hept., me.al.	Bovine-liver lipids, pilchard oil	25	91
.....	169.8	384.0	s.chl., hept., me.al.	Brain lipids	24,29	92
1.5014 <sup>20°</sup>	169.8	384.0	s.acet., eth., pet.eth.	Marine oils	13,29,36	93
.....	169.8	384.0	s.bz., chl., me.al., pet.eth.	Bovine-liver lipids, herring oil	26,29	94
1.5017 <sup>26°</sup>	170.8	463.6	s.bz., chl., me.al., pet.eth.	Bovine-liver & hog-brain lipids, pilchard oil	19,24,29	95
1.5122 <sup>20°</sup>	157.4	427.1	s.bz., chl., eth., pet.eth.	Whale & fish-liver oils	13,29,36	96

superscript, referred to water at 4°C. /s/ Refractive index ( $n$ ) is given for the sodium D-line at temperature shown grams of acid.

*continued*

## 100. FATTY ACIDS: PHYSICAL AND

Acid		Chemical Formula	Molecular Weight	Melting Point °C	Boiling Point °C/mm <sup>1</sup>	Specific Gravity <sup>2</sup>
Systematic Name	Common Name					
(A)	(B)	(C)	(D)	(E)	(F)	(G)
<b>Hydroxyalkanoic Acids</b>						
97	2-Hydroxydodecanoic	2-Hydroxylauric	C <sub>12</sub> H <sub>24</sub> O <sub>3</sub>	216.3	73-74	.....
98	12-Hydroxydodecanoic	Sabincic	C <sub>12</sub> H <sub>24</sub> O <sub>3</sub>	216.3	84	.....
99	2-Hydroxytetradecanoic	2-Hydroxymyristic	C <sub>14</sub> H <sub>28</sub> O <sub>3</sub>	244.4	81.5-82	.....
100	11-Hydroxypentadecanoic	Convolvulinolic	C <sub>15</sub> H <sub>30</sub> O <sub>3</sub>	258.4	63.5-64	.....
101	2-Hydroxyhexadecanoic	2-Hydroxypalmitic	C <sub>16</sub> H <sub>32</sub> O <sub>3</sub>	272.4	86-87	.....
102	11-Hydroxyhexadecanoic	Jalapinolic	C <sub>16</sub> H <sub>32</sub> O <sub>3</sub>	272.4	68-69	.....
103	16-Hydroxyhexadecanoic	Juniperic	C <sub>16</sub> H <sub>32</sub> O <sub>3</sub>	272.4	95	.....
104	2-Hydroxyoctadecanoic	2-Hydroxystearic	C <sub>18</sub> H <sub>36</sub> O <sub>3</sub>	300.5	91	.....
105	23-Hydroxydocosanoic	Phellonic	C <sub>22</sub> H <sub>44</sub> O <sub>3</sub>	356.6	95-96	.....
106	2-Hydroxytetracosanoic	Cerebronic	C <sub>24</sub> H <sub>48</sub> O <sub>3</sub>	384.6	99.5-100.5	.....
107	3,11-Dihydroxytetradecanoic	Ipurolic	C <sub>14</sub> H <sub>28</sub> O <sub>4</sub>	260.4	100-101	.....
108	2,15-Dihydroxypentadecanoic	Dihydroxypentadecyclic	C <sub>15</sub> H <sub>30</sub> O <sub>4</sub>	274.4	102-103	.....
109	15,16-Dihydroxyhexadecanoic	Ustilic A	C <sub>16</sub> H <sub>32</sub> O <sub>4</sub>	288.4	112-113	.....
110	9,10-Dihydroxyoctadecanoic	9,10-Dihydroxystearic	C <sub>18</sub> H <sub>36</sub> O <sub>4</sub>	316.5	141 <sup>3</sup>	.....
111	9,10-Dihydroxyoctadecanoic	9,10-Dihydroxystearic	C <sub>18</sub> H <sub>36</sub> O <sub>4</sub>	316.5	90 <sup>10</sup>	.....
112	11,12-Dihydroxyeicosanoic	11,12-Dihydroxyarachidic	C <sub>20</sub> H <sub>40</sub> O <sub>4</sub>	344.5	130 <sup>9</sup>	.....
113	2,15,16-Trihydroxyhexadecanoic	Ustilic	C <sub>16</sub> H <sub>32</sub> O <sub>5</sub>	304.4	140	.....
114	9,10,16-Trihydroxyhexadecanoic	Aleuritic	C <sub>16</sub> H <sub>32</sub> O <sub>5</sub>	304.4	100	.....
<b>Keto, Epoxy, and Cyclo Fatty Acids</b>						
115	4-Ketopentanoic	Levulinic	C <sub>5</sub> H <sub>8</sub> O <sub>3</sub>	116.1	37.2	154/15
116	6-Ketooctadecanoic	Lactarinic	C <sub>18</sub> H <sub>34</sub> O <sub>3</sub>	298.5	87	.....
117	4-Keto-9,11,13-octadecatrienoic	α-Licanic	C <sub>18</sub> H <sub>28</sub> O <sub>3</sub>	292.4	74-75	.....
118	4-Keto- <i>trans</i> -9, <i>trans</i> -11, <i>trans</i> -13-octadecatrienoic	β-Licanic	C <sub>18</sub> H <sub>28</sub> O <sub>3</sub>	292.4	99.5	.....
119	<i>cis</i> -12,13-Epoxy- <i>cis</i> -9-octadecenoic	Vernolic	C <sub>18</sub> H <sub>32</sub> O <sub>3</sub>	296.5	31-32	.....
120	<i>cis</i> -3,10-Epoxyoctadecanoic	Epoxystearic	C <sub>18</sub> H <sub>34</sub> O <sub>3</sub>	298.5	57.5-58	.....
121	ω-(2- <i>n</i> -Octylcycloprop-1-enyl)-octanoic	Sterculic	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	294.5	18	.....
122	ω-(2- <i>n</i> -Octylcyclopropyl)-octanoic	Lactobacillic	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	296.5	28-29	.....
123	13-(2-Cyclopentenyl)-tridecanoic	Chaulmoogric	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	280.2	68.5	247.5/20
124	11-(2-Cyclopentenyl)-hendecanoic	Hydnocarpic	C <sub>16</sub> H <sub>28</sub> O <sub>2</sub>	252.2	60.5	.....
125	9-(2-Cyclopentenyl)-nonanoic	Alepric	C <sub>14</sub> H <sub>24</sub> O <sub>2</sub>	224.2	48.0	.....
126	7-(2-Cyclopentenyl)-heptanoic	Aleprylic	C <sub>12</sub> H <sub>20</sub> O <sub>2</sub>	196.2	32.0	.....
127	5-(2-Cyclopentenyl)-pentanoic	Aleprestic	C <sub>10</sub> H <sub>16</sub> O <sub>2</sub>	168.1	Liquid	.....
128	2-Cyclopentenyl-1-oic	Aleprolic	C <sub>6</sub> H <sub>8</sub> O <sub>2</sub>	112.1	Liquid	.....
129	13-(2-Cyclopentenyl)-6-tridecenoic	Gorlic	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	278.2	6.0	232.5
<b>Hydroxy Unsaturated Acids</b>						
130	16-Hydroxy-7-hexadecenoic	Ambrettolic	C <sub>16</sub> H <sub>30</sub> O <sub>3</sub>	270.5	25	.....
131	9-Hydroxy-12-octadecenoic	.....	C <sub>18</sub> H <sub>34</sub> O <sub>3</sub>	298.5	.....	.....
132	<i>d</i> -12-Hydroxy- <i>cis</i> -9-octadecenoic	Ricinoleic	C <sub>18</sub> H <sub>34</sub> O <sub>3</sub>	298.5	5, 7.7, & 16	225/10
133	<i>d</i> -12-Hydroxy- <i>trans</i> -9-octadecenoic	Ricinelaidic	C <sub>18</sub> H <sub>34</sub> O <sub>3</sub>	298.5	52-53	.....
134	2-Hydroxy-15-tetracosenoic	Hydroxynervonic	C <sub>24</sub> H <sub>46</sub> O <sub>3</sub>	382.6	65	.....
135	9-Hydroxy-10,12-octadecadienoic	.....	C <sub>18</sub> H <sub>32</sub> O <sub>3</sub>	296.5	.....	.....

1/ 760 mm of mercury (atmospheric pressure), unless otherwise specified. 2/ At temperature indicated in superscript. 4/ Milligrams KOH required to neutralize one gram of acid. 5/ Grams of iodine absorbed by 100

## CHEMICAL CHARACTERISTICS

Refractive Index <sup>a</sup> $n_D^{20}$	Neutralization Value <sup>4</sup>	Iodine Value (Calculated) <sup>5</sup>	Solubility	Source	Reference
(H)	(I)	(J)	(K)	(L)	(M)
<b>Hydroxyalkanoic Acids</b>					
.....	259.4	.....	s.al., me.al.	Wool wax	29 97
.....	259.4	.....	s.al., h.bz.	Juniper wax	3,13,29 98
.....	229.1	.....	s.al., chl., eth.	Beeswax, wool wax	29 99
.....	217.1	.....	s.al., chl., eth.	Convolvulin resin	13,29,34 100
.....	206.0	.....	s.al., me.al.	Wool wax	29,40 101
.....	206.0	.....	s.al., eth.	Jalap-root resin	13,29,33 102
.....	206.0	.....	s.al., bz., eth.	Conifer waxes	13,21,29,36 103
.....	186.7	.....	s.al., me.al.	Wool wax	29,40 104
.....	157.3	.....	s.acet., chl., eth., glac.ac.a, pyr.	Cork	5,13,29,36 105
.....	145.9	.....	s.acet., h.al., eth., pyr.	Cerebrosides	13,29,36 106
.....	215.5	.....	s.chl., eth.	<i>Ipomoea purpurea</i>	13,29,33 107
.....	204.5	.....	s.me.al.	Ustilic acid B	29 108
.....	194.5	.....	s.me.al.	Fermentation of <i>Ustilago zeae</i>	29 109
.....	177.3	.....	s.h.al.; sl.s.eth.	Castor oil	13,29 110
.....	177.3	.....	s.al., eth., h.w.	Soils and straw	29 111
.....	162.9	.....	s.acet., eth.	Rabbit's-ear & mustard-seed oils	13,29 112
.....	184.3	.....	s.me.al.	Fermentation of <i>Ustilago zeae</i>	29 113
.....	184.3	.....	s.me.al.	Shellac wax	29 114
<b>Keto, Epoxy, and Cyclo Fatty Acids</b>					
1.442 <sup>15.8</sup> <sup>o</sup>	483.2	.....	v.s.al., eth., w.	Hexoses + HCl	29 115
.....	188.0	.....	s.h.al., chl., eth.	<i>Lactarius</i> mushrooms	13,21,29,36 116
.....	191.9	260.4	s.h.pet.eth.	Oiticica oil	4,15,29,36 117
.....	191.9	260.4	s.h.pet.eth.	Elaidinized $\alpha$ -acid	4,15,29,36 118
.....	189.3	85.6	s.acet., al., hex.	<i>Vernonia anthelmintica</i> seed oil	27,29 119
.....	188.0	.....	s.acet., al., hex.	<i>Tragopogon porrifolius</i> seed oil	6 120
.....	190.5	86.2	s.eth.	<i>Sterculia foetida</i> seed oil	29 121
.....	189.2	.....	s.acet., eth., pet.eth.	<i>Lactobacillus plantarum</i> lipids	29 122
.....	200.1	90.5	s.acet., chl., eth.	<i>Hydnocarpus</i> seed oils	12,13,21,29,36 123
.....	222.3	100.6	s.al., chl., pet.eth.	<i>Hydnocarpus</i> seed oils	21,29,36 124
.....	250.1	113.1	s.al., eth., pet.eth.	<i>Hydnocarpus</i> seed oils	29 125
.....	285.8	129.3	s.acet., eth., pet.eth.	<i>Hydnocarpus</i> seed oils	29 126
.....	333.5	150.8	s.acet., eth., pet.eth.	<i>Hydnocarpus</i> seed oils	29 127
.....	500.4	226.4	s.acet., eth., pet.eth.	<i>Hydnocarpus</i> seed oils	29 128
1.4782 <sup>25</sup> <sup>o</sup>	201.5	182.5	s.h.al.	Gorli oil	11,13,29,36 129
<b>Hydroxy Unsaturated Acids</b>					
.....	207.5	93.9	s.al., eth.	Musk-seed oil	29,36 130
.....	188.0	85.0	s.acet., al., eth.	<i>Strophanthus</i> seed oils	29 131
1.4716 <sup>20</sup> <sup>o</sup>	188.0	85.0	s.acet., al., eth.	Castor & ergot oils	13,29,36 132
.....	188.0	85.0	s.acet., al., eth.	Elaidinized ricinoleic acid	13,29,36 133
.....	146.6	66.3	s.acet., al., chl., eth., pyr.; sl.s. pet.eth.	Cerebrosides	13,29,36 134
.....	189.2	171.2	s.acet., al., pent.	<i>Tragopogon porrifolius</i> seed oil	8 135

superscript, referred to water at 4°C. /<sub>3</sub>/ Refractive index ( $n$ ) is given for the sodium D-line at temperature shown grams of acid. /<sub>8</sub>/ *Erythro*. /<sub>10</sub>/ *Threo*.

continued



100. FATTY ACIDS: PHYSICAL AND

Acid		Chemical Formula	Molecular Weight	Melting Point °C	Boiling Point °C/mm <sup>1</sup>	Specific Gravity <sup>3</sup>
Systematic Name	Common Name					
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Hydroxy Unsaturated Acids						
136	13-Hydroxy-9,11-octadecadienoic	.....	C <sub>18</sub> H <sub>32</sub> O <sub>3</sub>	296.5	.....	.....
137	18-Hydroxy- <i>cis</i> -9, <i>trans</i> -11, <i>trans</i> -13-octadecatrienoic	α-Kamlolenic	C <sub>18</sub> H <sub>30</sub> O <sub>3</sub>	294.4	77-78	.....
138	18-Hydroxy- <i>trans</i> -9, <i>trans</i> -11, <i>trans</i> -13-octadecatrienoic	β-Kamlolenic	C <sub>18</sub> H <sub>30</sub> O <sub>3</sub>	294.4	88-89	.....
Branched-Chain Fatty Acids						
139	3-Methylbutanoic	Isovaleric	C <sub>5</sub> H <sub>10</sub> O <sub>2</sub>	102.1	-37.6	176.7
140	<i>d</i> -6-Methyloctanoic	.....	C <sub>9</sub> H <sub>18</sub> O <sub>2</sub>	158.2	.....	0.937 <sup>150</sup>
141	8-Methyldecanoic	.....	C <sub>11</sub> H <sub>22</sub> O <sub>2</sub>	186.3	-18.5	.....
142	10-Methylhendecanoic	Isolauric	C <sub>12</sub> H <sub>24</sub> O <sub>2</sub>	200.3	41.2	.....
143	<i>d</i> -10-Methyldodecanoic	.....	C <sub>13</sub> H <sub>26</sub> O <sub>2</sub>	214.3	6.2-6.5	.....
144	11-Methyldodecanoic	Isoundecylic	C <sub>13</sub> H <sub>26</sub> O <sub>2</sub>	214.3	39.4-40	.....
145	12-Methyltridecanoic	Isomyristic	C <sub>14</sub> H <sub>28</sub> O <sub>2</sub>	228.4	53.6	.....
146	<i>d</i> -12-Methyltetradecanoic	.....	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	242.4	25.8	.....
147	13-Methyltetradecanoic	Isopentadecylic	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	242.4	52.2	.....
148	14-Methylpentadecanoic	Isopalmitic	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256.4	62.4	.....
149	<i>d</i> -14-Methylhexadecanoic	.....	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	270.4	38.0	.....
150	15-Methylhexadecanoic	.....	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	270.4	60.5	.....
151	10-Methylheptadecanoic	.....	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	284.5	33.5	.....
152	16-Methylheptadecanoic	Isostearic	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	284.5	69.5	.....
153	<i>l</i> - <i>D</i> -10-Methyloctadecanoic	Tuberculostearic	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	298.5	13.2	175-178/0.7
154	<i>d</i> -16-Methyloctadecanoic	.....	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	298.5	49.9-50.7	0.887 <sup>250</sup>
155	18-Methylnonadecanoic	Isoarachidic	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	312.5	75.3	.....
156	<i>d</i> -18-Methyleicosanoic	.....	C <sub>21</sub> H <sub>42</sub> O <sub>2</sub>	326.6	55.6	.....
157	20-Methylheneicosanoic	Isobehehic	C <sub>22</sub> H <sub>44</sub> O <sub>2</sub>	340.6	79.5	.....
158	<i>d</i> -20-Methyldocosanoic	.....	C <sub>23</sub> H <sub>46</sub> O <sub>2</sub>	354.6	62.1	.....
159	22-Methyltricosanoic	Isolignoceric	C <sub>24</sub> H <sub>48</sub> O <sub>2</sub>	368.6	83.1	.....
160	<i>d</i> -22-Methyltetracosanoic	.....	C <sub>25</sub> H <sub>50</sub> O <sub>2</sub>	382.7	67.8	.....
161	24-Methylpentacosanoic	Isocerotic	C <sub>26</sub> H <sub>52</sub> O <sub>2</sub>	396.7	86.9	.....
162	<i>d</i> -24-Methylhexacosanoic	.....	C <sub>27</sub> H <sub>54</sub> O <sub>2</sub>	410.7	72.9	.....
163	26-Methylheptacosanoic	Isomontanic	C <sub>28</sub> H <sub>56</sub> O <sub>2</sub>	424.7	89.3	.....
164	<i>d</i> -28-Methyltriacontanoic	.....	C <sub>31</sub> H <sub>62</sub> O <sub>2</sub>	466.8	80.7	.....
165	2,4,6-( <i>D</i> )-Trimethyloctacosanoic	Mycoceranic(mycocerosic)	C <sub>31</sub> H <sub>62</sub> O <sub>2</sub>	466.8	27-28	.....
166	2-Methyl- <i>cis</i> -2-butenic	Angelic	C <sub>5</sub> H <sub>8</sub> O <sub>2</sub>	100.1	45	185
167	2-Methyl- <i>trans</i> -2-butenic	Tiglic	C <sub>5</sub> H <sub>8</sub> O <sub>2</sub>	100.1	65.5	198.5
168	4-Methyl-3-pentenoic	Pyroterebic	C <sub>6</sub> H <sub>10</sub> O <sub>2</sub>	114.1	.....	207
169	<i>d</i> -2,4( <i>L</i> ),6( <i>L</i> )-Trimethyl- <i>trans</i> -2-tetracosenoic	C <sub>27</sub> -Phthienic (mycolipenic)	C <sub>27</sub> H <sub>52</sub> O <sub>2</sub>	408.7	39.5-41	.....

<sup>1</sup>/ 760 mm of mercury (atmospheric pressure), unless otherwise specified. <sup>2</sup>/ At temperature indicated in superscript. <sup>4</sup>/ Milligrams KOH required to neutralize one gram of acid. <sup>5</sup>/ Grams of iodine absorbed by 100

Contributor: Markley, Klare S.

References: [1] Baudert, P. 1942. Bull. Soc. Chim. France, Ser. 5, 9:922. [2] Bosworth, A. W., and F. B. Brown. W. B., and E. H. Farmer. 1935. Biochem. J. 29:631. [5] Chibnall, A. C., S. H. Piper, and E. F. Williams. 1936. and C. Y. Hopkins. 1960. Can. J. Chem. 38:805. [8] Chisholm, M. J., and C. Y. Hopkins. 1960. Ibid. 38:2500. 1962. J. Org. Chem. 27:3137. [11] Cole, H. I., and H. T. Cardoss. 1938. J. Am. Chem. Soc. 60:612. [12] Cole, biochemistry. Interscience, New York. [14] Dorinson, A., M. R. McCorkle, and A. W. Ralston. 1942. J. Am. [16] Foreman, H. D., and J. B. Brown. 1944. Oil Soap (Chicago) 21:183. [17] Francis, F., and S. H. Piper. 1939. 19:519. [19] Hammond, E. G., and W. W. Lundberg. 1953. J. Am. Oil Chemists' Soc. 30:438. [20] Hansen, R. P.,

## CHEMICAL CHARACTERISTICS

Refractive Index <sup>3</sup> $n_{\text{D}}^{20}$	Neutralization Value <sup>4</sup>	Iodine Value (Calculated) <sup>5</sup>	Solubility	Source	Reference
(H)	(I)	(J)	(K)	(L)	(M)
<b>Hydroxy Unsaturated Acids</b>					
.....	189.2	171.2	s.acet., al., pent.	<i>Tragopogon porrifolius</i> seed oil	8 136
.....	190.5	258.6	.....	Kamala-seed oil	29 137
.....	190.5	258.6	.....	Elaidinized $\alpha$ -acid	29 138
<b>Branched-Chain Fatty Acids</b>					
1.40178 <sup>22,40</sup>	549.3	.....	s.al., chl., eth.; sl.s.w.	Dolphin & porpoise head oils	13,28,29,36 139
.....	354.6	.....	s.acet., eth., me.al., pet.eth.	Wool grease	13,29,36,40 140
.....	301.2	.....	s.acet., eth., me.al., pet.eth.	Wool grease	13,29,36,40 141
.....	280.1	.....	s.acet., eth., me.al., pet.eth.	Wool grease	13,29,36,40 142
1.4424 <sup>250</sup>	261.8	.....	s.bz., chl., me.al., pet.eth.	Wool grease, butter, mutton tallow	13,29,36,40 143
1.4293 <sup>600</sup>	261.8	.....	s.acet., al., me.al., pet.eth.	Butterfat	29 144
.....	245.7	.....	s.acet., me.al., pet.eth.	Wool grease, butterfat	13,29,36,40 145
1.4327 <sup>590</sup>	231.5	.....	s.chl., eth., me.al., pet.eth.	Wool grease, butter, mutton tallow	13,29,36,40 146
1.4312 <sup>590</sup>	231.5	.....	s.me.al., pet.eth.	Butterfat	29 147
1.4293 <sup>700</sup>	218.8	.....	s.acet., eth., me.al., pet.eth.	Wool grease, butterfat	13,29,36,40 148
.....	207.5	.....	s.acet., eth., me.al., pet.eth.	Wool grease, mutton tallow	13,29,36,40 149
1.4315 <sup>700</sup>	207.5	.....	s.acet., eth., pet.eth.	Wool grease, beef tallow	13,29,36,40 150
.....	197.2	.....	s.acet., glac.ac.a.	Butterfat	20 151
.....	197.2	.....	s.acet., eth., pet.eth.	Wool grease	13,29,36,40 152
1.4512 <sup>250</sup>	188.0	.....	s.acet., al., me.al., pent.	Human tubercle bacilli lipids	13,29,36 153
.....	188.0	.....	s.acet., me.al., pet.eth.	Wool grease	13,29,35,36,40 154
.....	179.5	.....	s.al., eth., pet.eth.	Wool grease	13,29,36,40 155
.....	171.8	.....	s.acet., chl., pet.eth.	Wool grease	13,29,36,40 156
.....	164.7	.....	s.chl., eth., me.al., pet.eth.	Wool grease	13,29,36,40 157
.....	158.2	.....	s.acet., chl., eth., pet.eth.	Wool grease	13,29,36,40 158
.....	152.2	.....	s.acet., chl., pet.eth.	Wool grease	13,29,36,40 159
.....	146.6	.....	s.al., bz., chl., pet.eth.	Wool grease	13,29,36,40 160
.....	141.4	.....	s.acet., chl., glac.ac.a.	Wool grease	13,29,36,40 161
.....	136.6	.....	s.bz., chl., glac.ac.a., pet.eth.	Wool grease	13,29,36,40 162
.....	132.1	.....	s.bz., chl., glac.ac.a., pet.eth.	Wool grease	13,29,36,40 163
.....	120.2	.....	s.bz., chl., glac.ac.a., pet.eth.	Wool grease	13,29,36,40 164
.....	120.2	.....	s.chl., pet.eth.	Tubercle bacilli lipids	13,29 165
1.4434 <sup>470</sup>	560.4	253.6	v.s.eth.; s.al.; sl.s.w.	<i>Angelica</i> root & Roman camomile oils	29 166
1.4342 <sup>810</sup>	560.4	253.6	v.s.h.w.; s.al., eth.	<i>Croton tiglium</i> root & Roman camomile oils	29 167
.....	491.6	222.4	s.al., chl., eth.	<i>Calotropis procera</i> sap	29 168
1.4598 <sup>250</sup>	137.3	62.1	s.acet., me.al., pet.eth.	Tubercle bacilli lipids	29 169

superscript, referred to water at 4°C. /s/ Refractive index ( $n$ ) is given for the sodium D-line at temperature shown grams of acid.

1933. J. Biol. Chem. 103:115. [3] Bougault, J., and L. Bourdier. 1909. J. Pharm. Chim., Ser. 6, 30:10. [4] Brown, Ibid. 30:100. [6] Chisholm, M. J., and C. Y. Hopkins. 1959. Chem. Ind. (London), p. 1154. [7] Chisholm, M. J., [9] Chisholm, M. J., and C. Y. Hopkins. 1962. J. Chem. Soc., p. 573. [10] Chisholm, M. J., and C. Y. Hopkins. H. I., and H. T. Cardoss. 1939. Ibid. 61:2349. [13] Deuel, H. J., Jr. 1951-57. The lipids; their chemistry and Chem. Soc. 64:2739. [15] Dyson, M. G. 1950. A manual of organic chemistry. Longmans, Green; London. v.1. J. Am. Chem. Soc. 61:577. [18] Gupta, R. S., A. Grollman, and S. C. Niyogy. 1953. Proc. Natl. Inst. Sci. India F. B. Shorland, and N. J. Cooke. 1951. Chem. Ind. (London), p. 839. [21] Heilbron, I. M. 1934. Dictionary of

continued

## 100. FATTY ACIDS: PHYSICAL AND

organic compounds. Eyre and Spottiswoode, London. [22] Hilditch, T. P., and E. E. Jones. 1928. Biochem. J. Bongard. 1952. Z. Physiol. Chem. 291:104. [25] Klenk, E., and W. Montag. 1957. Ann. Chem. 604:4. [26] Klenk, Riemenschneider. 1962. J. Am. Oil Chemists' Soc. 39:334. [28] Lange, N. A. 1946. Handbook of chemistry. Ed. [30] Noller, C. R., and R. H. Talbot. 1943. Organic syntheses collection. J. Wiley, New York. v. 12. [31] Nunn, H. Rogerson. 1910. J. Am. Chem. Soc. 32:106. [34] Power, F. B., and H. Rogerson, 1912. J. Chem. Soc. 101(T):1. Fatty acids and their derivations. J. Wiley, New York. [37] Smedley, I. 1912. Biochem. J. 6:451. [38] Teresi, The chemistry and technology of waxes. Ed. 2. Reinhold, New York. [40] Weitkamp, A. W. 1945. J. Am. Chem.

## 101. FATS AND OILS: PHYSICAL

Values are typical rather than average, and frequently were derived from specific analyses for particular samples source, treatment, and age of a fat or oil. **Specific Gravity** (column D) was calculated at the specified temperature parentheses (column D), was measured at the specified temperature (degrees centigrade). **Refractive Index** (column

	Fat or Oil	Source	Constants				
			Melting (or Solidification) Point, °C	Specific Gravity (or Density)	Refractive Index $n_{D}^{40°C}$	Iodine Value	Saponification Value
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Land Animals Butterfat	<i>Bos taurus</i>	32.2	0.911 <sup>40°/15°</sup>	1.4548	36.1	227
2	Depot fat	<i>Homo sapiens</i>	(15)	0.918 <sup>15°</sup>	1.4602	67.6	196.2
3	Lard oil	<i>Sus scrofa</i>	(30.5)	0.919 <sup>15°</sup>	1.4615	58.6	194.6
4	Neat's-foot oil	<i>B. taurus</i>	.....	0.910 <sup>25°</sup>	1.464 <sup>25°</sup>	69-76	190-199
5	Tallow, beef	<i>B. taurus</i>	.....	.....	.....	49.5	197
6	Tallow, mutton	<i>Ovis aries</i>	(42.0)	0.945 <sup>15°</sup>	1.4565	40	194
Marine Animals							
7	Cod-liver oil	<i>Gadus morhua</i>	.....	0.925 <sup>25°</sup>	1.481 <sup>25°</sup>	165	186
8	Herring oil	<i>Clupea harengus</i>	.....	0.900 <sup>60°</sup>	1.4610 <sup>60°</sup>	140	192
9	Menhaden oil	<i>Brevoortia tyrannus</i>	.....	0.903 <sup>60°</sup>	1.4645 <sup>60°</sup>	170	191
10	Sardine oil	<i>Sardinops caerulea</i>	.....	0.905 <sup>60°</sup>	1.4660 <sup>60°</sup>	185	191
11	Sperm oil, body	<i>Physeter macrocephalus</i>	.....	.....	.....	76-88	122-130
12	Sperm oil, head	<i>P. macrocephalus</i>	.....	.....	.....	70	140-144
13	Whale oil	<i>Balaena mysticetus</i>	.....	0.892 <sup>60°</sup>	1.460 <sup>60°</sup>	120	195
Plants							
14	Babassu oil	<i>Attalea funifera</i>	22-26	(0.893 <sup>60°</sup> )	1.443 <sup>60°</sup>	15.5	247
15	Castor oil	<i>Ricinus communis</i>	(-18.0)	0.961 <sup>15°</sup>	1.4770	85.5	180.3
16	Cocoa butter	<i>Theobroma cacao</i>	34.1	0.964 <sup>15°</sup>	1.4568	36.5	193.8
17	Coconut oil	<i>Cocos nucifera</i>	25.1	0.924 <sup>15°</sup>	1.4493	10.4	268
18	Corn oil	<i>Zea mays</i>	(-20.0)	0.922 <sup>15°</sup>	1.4734	122.6	192.0
19	Cottonseed oil	<i>Gossypium hirsutum</i>	(-1.0)	0.917 <sup>25°</sup>	1.4735	105.7	194.3
20	Linseed oil	<i>Linum usitatissimum</i>	(-24.0)	0.938 <sup>15°</sup>	1.4782 <sup>25°</sup>	178.7	190.5
21	Mustard oil	<i>Brassica hirta</i>	.....	0.9145 <sup>15°</sup>	1.475	102	174
22	Neem oil	<i>Melia azadirachta</i>	-3	0.917 <sup>15°</sup>	1.4615	71	194.5
23	Niger-seed oil	<i>Guizotia abyssinica</i>	.....	0.925 <sup>15°</sup>	1.471	128.5	190
24	Oiticica oil	<i>Licania rigida</i>	.....	0.974 <sup>25°</sup>	.....	140-180	.....
25	Olive oil	<i>Olea europaea sativa</i>	(-6.0)	0.918 <sup>15°</sup>	1.4679	81.1	189.7
26	Palm oil	<i>Elaeis guineensis</i>	35.0	0.915 <sup>15°</sup>	1.4578	54.2	199.1
27	Palm-kernel oil	<i>E. guineensis</i>	24.1	0.923 <sup>15°</sup>	1.4569	37.0	219.9
28	Peanut oil	<i>Arachis hypogaea</i>	(3.0)	0.914 <sup>15°</sup>	1.4691	93.4	192.1
29	Perilla oil	<i>Perilla frutescens</i>	.....	(0.935 <sup>15°</sup> )	1.481 <sup>25°</sup>	195	192
30	Poppy-seed oil	<i>Papaver somniferum</i>	(-15)	0.925 <sup>15°</sup>	1.4685	135	194
31	Rapeseed oil	<i>Brassica campestris</i>	(-10)	0.915 <sup>15°</sup>	1.4706	98.6	174.7
32	Safflower oil	<i>Carthamus tinctorius</i>	.....	(0.900 <sup>60°</sup> )	1.462 <sup>60°</sup>	145	192
33	Sesame oil	<i>Sesamum indicum</i>	(-6.0)	0.919 <sup>25°</sup>	1.4646	106.6	187.9
34	Soybean oil	<i>Glycine soja</i>	(-16.0)	0.927 <sup>15°</sup>	1.4729	130.0	190.6
35	Sunflower-seed oil	<i>Helianthus annuus</i>	(-17.0)	0.923 <sup>15°</sup>	1.4694	125.5	188.7
36	Tung oil	<i>Aleurites fordii</i>	(-2.5)	0.934 <sup>15°</sup>	1.5174 <sup>25°</sup>	168.2	193.1
37	Wheat-germ oil	<i>Triticum aestivum</i>	.....	.....	.....	125	.....

/1/ Caproic. /2/ Caprylic. /3/ Capric. /4/ Butyric. /5/ Decenoic. /6/ C<sub>12</sub> monoethenoic. /7/ C<sub>14</sub> monoethenoic. polyethenoic. /13/ C<sub>22</sub> polyethenoic. /14/ Behenic. /15/ C<sub>14</sub> polyethenoic. /16/ Gadoleic. /17/ C<sub>24</sub> polyethenoic. cludes behenic. /24/ Licanic. /25/ Eleostearic.

## CHEMICAL CHARACTERISTICS

22:326. [23] Hopkins, C. Y., and M. J. Chisholm. 1962. Chem. Ind. (London), p. 2064. [24] Klenk, E., and W. E., and M. J. Tomuschat. 1957. Z. Physiol. Chem. 308:165. [27] Krewson, C. F., J. S. Ard, and R. W. 6. Handbook Publications, Sandusky, Ohio. [29] Markley, K. S. 1960-61. Fatty acids. Ed. 2. Interscience, New York. J. R. 1952. J. Chem. Soc., p. 313. [32] Piper, S. H., et al. 1934. Biochem. J. 28:2175. [33] Power, F. B., and [35] Prout, F. S., J. Cason, and A. W. Ingersoll. 1947. J. Am. Chem. Soc. 69:1233. [36] Ralston, A. W. 1948. J. D. Unpublished. U.S. Naval Radiological Defense Laboratory, San Francisco, Calif. [39] Warth, A. H. 1956. Soc. 67:447.

## AND CHEMICAL CHARACTERISTICS

(especially the constituent fatty acids). Extreme variations may occur, depending on a number of variables such as (degrees centigrade) and referred to water at the same temperature, unless otherwise specified. **Density**, shown in E) was measured at 40°C, unless otherwise specified.

Constituent Fatty Acids, g/100 g total fatty acids												
Saturated						Unsaturated						
Lauric	Myris- tic	Palmi- tic	Stearic	Ara- chidic	Other	Palmit- oleic	Oleic	Lino- leic	Lino- lenic	Other		
(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)		
2.5	11.1	29.0	9.2	2.4	2.0 <sup>1</sup> ; 0.5 <sup>2</sup> ; 2.3 <sup>3</sup>	4.6	26.7	3.6	....	3.6 <sup>4</sup> ; 0.1 <sup>5</sup> ; 0.1 <sup>6</sup> ; 0.9 <sup>7</sup> ; 1.4 <sup>8</sup> ; 1.0 <sup>9</sup> ; 1.0 <sup>10</sup> ; 0.4 <sup>11</sup>		1
....	2.7	24.0	8.4	....	.....	5	46.9	10.2	....	2.5 <sup>5</sup>		2
....	1.3	28.3	11.9	....	.....	2.7	47.5	6	....	0.2 <sup>7</sup> ; 2.1 <sup>8</sup>		3
....	.....	17-18	2-3	....	.....	....	74-76	....	....	.....		4
....	6.3	27.4	14.1	....	.....	....	49.6	2.5	....	.....		5
....	4.6	24.6	30.5	....	.....	....	36.0	4.3	....	.....		6
....	5.8	8.4	0.6	....	.....	20.0	←29.1→	....	....	25.4 <sup>12</sup> ; 9.6 <sup>13</sup>		7
....	7.3	13.0	Trace	....	.....	4.9	....	....	20.7	30.1 <sup>12</sup> ; 23.2 <sup>13</sup>		8
....	5.9	16.3	0.6	0.6	.....	15.5	....	....	29.6	19.0 <sup>12</sup> ; 11.7 <sup>13</sup> ; 0.8 <sup>14</sup>		9
....	5.1	14.6	3.2	....	.....	11.8	←17.8→	....	....	18.1 <sup>12</sup> ; 14.0 <sup>13</sup> ; trace <sup>7</sup> ; 15.4 <sup>15</sup>		10
1	5	6.5	.....	....	.....	26.5	37	19	....	1 <sup>13</sup> ; 4 <sup>7</sup> ; 19 <sup>15</sup>		11
16	14	8	2	....	3.5 <sup>3</sup>	15	17	6.5	....	4 <sup>8</sup> ; 14 <sup>7</sup> ; 6.5 <sup>15</sup>		12
0.2	9.3	15.6	2.8	....	.....	14.4	35.2	....	....	13.6 <sup>12</sup> ; 5.9 <sup>13</sup> ; 2.5 <sup>7</sup> ; 0.2 <sup>17</sup>		13
44.1	15.4	8.5	2.7	0.2	0.2 <sup>1</sup> ; 4.8 <sup>2</sup> ; 6.6 <sup>3</sup>	....	16.1	1.4	....	.....		14
←.....	.....	2.4	.....	.....	.....	....	7.4	3.1	....	87 <sup>13</sup>		15
....	.....	24.4	35.4	....	.....	....	38.1	2.1	....	.....		16
45.4	18.0	10.5	2.3	0.4 <sup>13</sup>	0.8 <sup>1</sup> ; 5.4 <sup>2</sup> ; 8.4 <sup>3</sup>	0.4	7.5	Trace	....	.....		17
....	1.4	10.2	3.0	....	.....	1.5	49.6	34.3	....	.....		18
....	1.4	23.4	1.1	1.3	.....	2.0	22.9	47.8	....	.....		19
....	.....	6.3	2.5	0.5	.....	....	19.0	24.1	47.4	0.2 <sup>14</sup>		20
....	1.3 <sup>20</sup>	.....	.....	....	.....	....	27.2 <sup>20</sup>	16.6 <sup>20</sup>	1.8 <sup>20</sup>	1.1 <sup>14</sup> ; 1.0 <sup>21</sup> ; 51.0 <sup>22</sup>		21
....	2.6 <sup>20</sup>	14.1 <sup>20</sup>	24.0 <sup>20</sup>	0.8 <sup>20</sup>	.....	....	58.5 <sup>20</sup>	....	....	.....		22
....	3.3 <sup>20</sup>	8.2 <sup>20</sup>	4.8 <sup>20</sup>	0.5 <sup>20</sup>	.....	....	30.3 <sup>20</sup>	57.3 <sup>20</sup>	....	.....		23
←.....	.....	11.3 <sup>23</sup>	.....	.....	.....	....	6.2	....	....	82.5 <sup>24</sup>		24
....	Trace	6.9	2.3	0.1	.....	....	84.4	4.6	....	.....		25
....	1.4	40.1	5.5	....	.....	....	42.7	10.3	....	.....		26
46.9	14.1	8.8	1.3	....	2.7 <sup>2</sup> ; 7.0 <sup>3</sup>	....	18.5	0.7	....	.....		27
....	.....	8.3	3.1	2.4	.....	....	56.0	26.0	....	3.1 <sup>14</sup> ; 1.1 <sup>21</sup>		28
←.....	.....	9.6 <sup>23</sup>	.....	.....	.....	....	17.8	....	17.5	.....		29
....	.....	4.8 <sup>20</sup>	2.9 <sup>20</sup>	....	.....	....	30.1 <sup>20</sup>	62.2 <sup>20</sup>	....	.....		30
....	.....	1	.....	....	.....	....	32	15	1	50 <sup>22</sup>		31
←.....	.....	6.8 <sup>23</sup>	.....	.....	.....	....	18.6	70.1	3.4	.....		32
....	.....	9.1	4.3	0.8	.....	....	45.4	40.4	....	.....		33
0.2	0.1	9.8	2.4	0.9	.....	0.4	28.9	50.7	6.5	0.1 <sup>7</sup>		34
....	.....	5.6	2.2	0.9	.....	....	25.1	66.2	....	.....		35
←.....	.....	4.6 <sup>23</sup>	.....	.....	.....	....	4.1	0.6	....	90.7 <sup>25</sup>		36
←.....	.....	16.0 <sup>23</sup>	.....	.....	.....	....	28.1	52.3	3.6	.....		37

<sup>1</sup>/ Gadoleic plus erucic. <sup>2</sup>/ C<sub>12</sub> n-pentadecanoic. <sup>10</sup>/ C<sub>17</sub> margaric. <sup>11</sup>/ 12-Methyl tetradecanoic. <sup>12</sup>/ C<sub>20</sub> <sup>13</sup>/ Ricinoleic. <sup>15</sup>/ Includes behenic and lignoceric. <sup>20</sup>/ Percent by weight. <sup>21</sup>/ Lignoceric. <sup>22</sup>/ Erucic. <sup>23</sup>/ In-

continued



## 101. FATS AND OILS: PHYSICAL AND CHEMICAL CHARACTERISTICS

Contributors: (a) Harwood, H. J., (b) Geyer, Robert P.

References: [1] Bailey, A. E. 1945. Industrial oil and fat products. Interscience, New York. [2] Deuel, H. J., Jr. 1951. The lipids; their chemistry and biochemistry. Interscience, New York. v. 1. [3] Hilditch, T. P. 1956. The chemical constitution of natural fats. Ed. 3. J. Wiley, New York.

## 102. WAXES: PHYSICAL AND CHEMICAL CHARACTERISTICS

Specific Gravity (column C) was calculated at the specified temperature, degrees centigrade, and referred to water at the same temperature. Density, shown in parentheses (column C), and Refractive Index (column D) were measured at the specified temperature, degrees centigrade.

Wax	Melting Point °C	Specific Gravity or (Density)	Refractive Index $n_D^{20}$	Iodine Value	Acid Value	Saponifi- cation Value	Ref- erence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Bamboo leaf	79-80	(0.961 <sup>250</sup> )	.....	7.8 <sup>1</sup>	14.5	43.4	9
2 Bayberry (myrtle)	46.7-48.8	(0.985 <sup>150</sup> )	1.436 <sup>800</sup>	2.9 <sup>2</sup> -3.9 <sup>3</sup>	3.5	20.5-21.7	7,10
3 Beeswax, crude	62-66	(0.927-0.970 <sup>150</sup> )	1.439-1.483 <sup>400</sup>	6.8-16.4 <sup>2</sup>	16.8-35.8	89.3-149.0	10
4 Beeswax, white, U.S.P.	61-69	(0.959-0.975 <sup>150</sup> )	1.447-1.465 <sup>650</sup>	7-11 <sup>3</sup>	17-24	90-96	10
5 Beeswax, yellow	62-65	(0.960-0.964 <sup>150</sup> )	1.443-1.449 <sup>650</sup>	6-11	18-24	90-97	10
6 Candelilla, refined	67-69	(0.982-0.986 <sup>150</sup> )	1.454-1.463 <sup>850</sup>	14.4-20.4	12.7-18.1	35-86	1
7 Cape berry <sup>a</sup>	40.5-45.0	(1.004-1.007 <sup>150</sup> )	1.450 <sup>450</sup>	0.6-2.4	2.5-3.7	211-215	10
8 Carandá	79.7-84.5	(0.990 <sup>250</sup> )	.....	8.0-8.9	5.0-9.5	64.5-78.5	2,10
9 Carnauba	83-86	0.990-1.001 <sup>150</sup>	1.467-1.472 <sup>400</sup>	7.2-13.5	2.9-9.7	78-95	10
10 Castor oil, hydrogenated	83-88	(0.980-0.990 <sup>200</sup> )	.....	2.5-8.5	1.0-5.0	177-181	6,10
11 Chinese insect	81.5-84.0	0.950-0.970 <sup>150</sup>	1.457 <sup>400</sup>	1.4	0.2-1.5	73-93	10
12 Cotton	68-71	0.959 <sup>150</sup>	.....	24.5	32	70.6	8
13 Cranberry	207-218	(0.970-0.975 <sup>150</sup> )	.....	44.2-53.2 <sup>2</sup>	42.2-59.1	131-134	10
14 Douglas-fir bark	59.0-72.8	(1.030 <sup>250</sup> )	1.468 <sup>800</sup>	25.8-62.5	58.6-80.7	112-200	10
15 Esparto	67.5-78.1	0.988 <sup>150</sup>	.....	22-23	22.7-23.9	69.8-79.3	10
16 Flax	61.5-69.8	0.908-0.985 <sup>150</sup>	.....	21.6-28.8	17.5-48.3	77.5-101.5	10
17 Ghedda, E. Indian beeswax	60.5-66.4	0.956-0.973 <sup>150</sup>	1.440 <sup>500</sup>	5.6-12.6	5.8-7.9	84.5-118.3	10
18 Indian corn	80-81	.....	.....	4.2 <sup>2</sup>	1.9	120.3	5
19 Japan wax	48-53	0.975-0.993 <sup>150</sup>	.....	4.5-12.5	6-20	206.5-237.5	10
20 Jojoba	11.2-11.8	0.864-0.899 <sup>250</sup>	1.465 <sup>250</sup>	81.7-88.4 <sup>2</sup>	0.2-0.6	92.2-95.0	10
21 Madagascar	88	.....	.....	3.2-5.3	17.7-28.0	140.0-159.6	10
22 Microcrystalline, amber	64-91	0.913-0.943 <sup>150</sup>	1.424-1.452 <sup>800</sup>	0	0	0	10
23 Microcrystalline, white	71-89	0.928-0.941 <sup>150</sup>	1.441 <sup>800</sup>	0	0	0	10
24 Montan, crude	76-86	(1.010-1.020 <sup>250</sup> )	.....	13.9-17.6	22.7-31.0	59.4-92.0	10
25 Montan, refined	77-84	(1.010-1.030 <sup>250</sup> )	.....	10-14	24-43	72-103	10
26 Orange peel	44.0-46.5	0.985 <sup>150</sup>	1.502 <sup>200</sup>	115.7 <sup>2</sup>	48.3	120.9	4
27 Ouricury, refined	79.0-83.8	1.053 <sup>150</sup>	.....	6.9-7.8 <sup>2</sup>	3.4-21.1	61.8-85.8	10
28 Ozocerite, refined	74.4-75.0	0.907-0.920 <sup>150</sup>	.....	0	0	0	10
29 Palm	74-86	(0.991-1.045 <sup>150</sup> )	.....	8.9-16.9 <sup>2</sup>	5.0-10.6	64.5-104.0	10
30 Paraffin, American	49-63	0.896-0.925 <sup>150</sup>	1.442-1.448 <sup>800</sup>	0	0	0	10
31 Peat wax, natural	73-76	0.980 <sup>150</sup>	.....	16-40	60.0-73.3	73.9-136.0	10
32 Rice bran, refined	75.3-79.9	.....	1.469 <sup>300</sup>	11.1-19.4	15-17	56.9-104.4	10
33 Shellac wax	79-82	0.971-0.980 <sup>150</sup>	.....	6.0-8.8 <sup>3</sup>	12.1-24.3	63.8-83.0	10
34 Sisal hemp	74-81	1.007-1.010 <sup>150</sup>	.....	28-29 <sup>2</sup>	16-19 <sup>2</sup>	56-58	10
35 Sorghum grain	77-82	.....	.....	15.7-20.9	10.1-16.2	16-44	10
36 Spanish moss	79-80	.....	.....	33.0	25.0	120.4	3
37 Spermaceti	42-50	0.905-0.945 <sup>150</sup>	1.440 <sup>700</sup>	4.8-5.9	2.0-5.2	108-134	10
38 Sugarcane, crude	52-67	0.988-0.998 <sup>250</sup>	.....	32-84	24-57	128-177	10
39 Sugarcane, double-refined	77-82	0.961-0.979 <sup>250</sup>	1.510 <sup>250</sup>	13-29	8-23	55-95	10
40 Wool wax, refined	36-43	0.932-0.945 <sup>150</sup>	1.478-1.482 <sup>400</sup>	15.0-46.9	5.6-22.0	80-127	10

<sup>1/</sup> Wijs test. <sup>2/</sup> Hanus test. <sup>3/</sup> Hubl test. <sup>4/</sup> *Myrica cordifolia*.

Contributor: Warth, Albin H.

References: [1] Alcocer, G., and J. M. Sanders. 1910. Anales Inst. Med. Nacl. Mexico 2:155. [2] De Medeiros Transcuso, A. 1948. Rev. Quim. Ind. (Rio de Janeiro) 17(192):21. [3] Feurt, S. D., and L. E. Fox. 1952. Science

continued

## 102. WAXES: PHYSICAL AND CHEMICAL CHARACTERISTICS

117:600. [4] Gonzales-Trigo, G. 1950. *Anales Bromatol.* (Madrid) 2:31. [5] Jamieson, G. S. 1943. *Vegetable fats and oils*. Ed. 2. Reinhold, New York. [6] McLoud, E. S. Unpublished. S. C. Johnson and Son, Inc., Racine, Wis., 1955. [7] Smith, W. R., and P. B. Wade. 1903. *J. Am. Chem. Soc.* 25:629. [8] Tomm, W. H., Jr., and E. P. Schoch. 1946. *Ind. Eng. Chem.* 38:413. [9] Tuzimoto, H. 1939. *J. Soc. Chem. Ind. Japan* 42 (Suppl. 396). [10] Warth, A. H. 1956. *The chemistry and technology of waxes*. Ed. 2. Reinhold, New York.

## 103. PHOSPHATIDES AND CEREBROSIDES: PHYSICAL AND CHEMICAL CHARACTERISTICS

Data are for synthetic lipids containing known fatty acids. Naturally occurring lipids contain a spectrum of fatty acids [11-13, 16-23, 26] and have different properties dependent on their fatty-acid composition. **Solubility** (column F): ac.a. = acetic acid; acet. = acetone; bz. = benzene; c.tet. = carbon tetrachloride; chl. = chloroform; die. = diethyl; e.acet. = ethyl acetate; e.al. = ethanol; eth. = ether; ethy. = ethylene; glac. = glacial; gly. = glycol; i. = insoluble; me.al. = methanol; monome. = monomethyl; pet. = petroleum; pyr. = pyridine; sl. = slightly; s. = soluble; v. = very; w. = water.

Lipid	Chemical Formula	Molecular Weight	Melting Point °C	Specific Rotation $[\alpha]_D$	Solubility	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Phosphatidylcholines (Lecithins)							
1	Dihexanoyl-L- $\alpha$ -phosphatidylcholine	C <sub>20</sub> H <sub>42</sub> O <sub>9</sub> NP	471.5	.....	+10.9 in chloroform	v.s.acet., chl., e.al., me.al., w.; i.die.eth.	7
2	Diocanoyl-L- $\alpha$ -phosphatidylcholine	C <sub>24</sub> H <sub>50</sub> O <sub>9</sub> NP	527.6	.....	+9.65 in chloroform-methyl alcohol (1:1)	v.s.acet., chl., e.al., me.al.; i.die.eth.	7
3	Didecanoyl-L- $\alpha$ -phosphatidylcholine	C <sub>28</sub> H <sub>58</sub> O <sub>9</sub> NP	583.7	.....	+8.75 in chloroform-methyl alcohol (1:1)	v.s.chl., e.al., me.al.; s.acet.; i.die.eth.	7
4	Dimyristoyl-L- $\alpha$ -phosphatidylcholine	C <sub>36</sub> H <sub>74</sub> O <sub>9</sub> NP	695.9	237-237.5	+7.0 in chloroform-methyl alcohol (1:1)	v.s.me.al., pyr.; s.acet.; i.w.	6
5	Dipalmitoyl-L- $\alpha$ -phosphatidylcholine	C <sub>40</sub> H <sub>78</sub> O <sub>9</sub> NP	748.1	.....	+6.6 in chloroform-methyl alcohol (1:1)	v.s. 90% acet., chl., e.al., me.al.; i.pet.eth., w.	14
6	Dipalmitoyl-L- $\alpha$ -phosphatidylcholine	C <sub>40</sub> H <sub>82</sub> O <sub>9</sub> NP	752.1	234-235	+6.6 in chloroform-methyl alcohol (1:1)	v.s.me.al., pyr.; s.acet.; i.w.	6
7	Dioley-L- $\alpha$ -phosphatidylcholine	C <sub>44</sub> H <sub>86</sub> O <sub>9</sub> NP	804.1	.....	+6.2 in chloroform-methyl alcohol (1:1)	v.s. 90% acet., die.eth., e.al., me.al.; s.pet.eth.	5
8	Distearoyl-L- $\alpha$ -phosphatidylcholine	C <sub>44</sub> H <sub>90</sub> O <sub>9</sub> NP	808.2	230.5-231.5	+6.1 in chloroform-methyl alcohol (1:1)	s.me.al., pyr.; sl.s.acet.; i.w.	6
Phosphatidylethanolamines (Cephalins)							
9	Dimyristoyl-L- $\alpha$ -phosphatidylethanolamine	C <sub>33</sub> H <sub>66</sub> O <sub>8</sub> NP	635.9	195-196	+6.7 in chloroform	v.s.chl.; s.bz., c.tet., e.al., pyr.; i.acet., die.eth., e.acet., pet.eth.	8
10	Dipalmitoyl-L- $\alpha$ -phosphatidylethanolamine	C <sub>37</sub> H <sub>74</sub> O <sub>8</sub> NP	692.0	186-187	+6.4 in chloroform	v.s.chl.; s.bz., c.tet., e.al., pyr.; i.acet., die.eth., e.acet., pet.eth.	8
11	Dioley-L- $\alpha$ -phosphatidylethanolamine	C <sub>41</sub> H <sub>78</sub> O <sub>8</sub> NP	744.1	.....	+6.0 in chloroform	v.s.acet., chl., die.eth., e.al., pet.eth.; i.w.	4
12	Distearoyl-L- $\alpha$ -phosphatidylethanolamine	C <sub>41</sub> H <sub>82</sub> O <sub>8</sub> NP	748.1	180-182	+6.0 in chloroform-acetic acid (1:1)	v.s.chl.; s.bz., c.tet., e.al., pyr.; i.acet., die.eth., e.acet., pet.eth.	8
Phosphatidylserines							
13	Distearoyl-L- $\alpha$ -phosphatidylserine	C <sub>42</sub> H <sub>82</sub> O <sub>10</sub> NP	792.1	159-161	-14.0 in chloroform	s.bz., chl.; i.w.	9
Phosphatidylglycerols							
14	(Dioley-L- $\alpha$ -glycerylphosphoryl)-L- $\alpha$ -glycerol	C <sub>42</sub> H <sub>79</sub> O <sub>10</sub> P	775.0	.....	+2.0 in chloroform	v.s.acet., bz., chl., die.eth., e.al., ethy.gly. monome.eth., me.al., pet.eth.; i.w.	3
15	(Distearoyl-L- $\alpha$ -glycerylphosphoryl)-L- $\alpha$ -glycerol	C <sub>42</sub> H <sub>83</sub> O <sub>10</sub> P	779.1	66.5-67.0	+2.0 in chloroform	v.s.acet., bz., chl., die.eth., e.al., ethy.gly. monome.eth., me.al., pet.eth.	3

*continued*

## 103. PHOSPHATIDES AND CEREBROSIDES: PHYSICAL AND CHEMICAL CHARACTERISTICS

Lipid	Chemical Formula	Molecular Weight	Melting Point °C	Specific Rotation $[\alpha]_D$	Solubility	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
<b>Phosphatidylinositols</b>						
16	Beef liver monophosphoinositide	.....	.....	+5.86 in chloroform	v.s.w.; s.chl., chl./e.al. (3/1), e.acet., glac.ac.a.; i.acet., e.al., me.al.	15
<b>Phosphatidic Acids</b>						
17	Dimyristoyl-L- $\alpha$ -glyceroylphosphoric acid	C <sub>31</sub> H <sub>61</sub> O <sub>8</sub> P	592.8	61.5-62.5	+4.4 in chloroform	v.s.acet., bz., die.eth., e.al., glac.ac.a.
18	Dipalmitoyl-L- $\alpha$ -glyceroylphosphoric acid	C <sub>35</sub> H <sub>69</sub> O <sub>8</sub> P	648.9	70-71	+4.0 in chloroform	v.s.acet., bz., die.eth., e.al., glac.ac.a.
19	Diioleoyl-L- $\alpha$ -glyceroylphosphoric acid	C <sub>39</sub> H <sub>73</sub> O <sub>8</sub> P	701.0	.....	+3.8 in chloroform	v.s.acet., bz., chl., die.eth., 99% e.al., pet.eth.
20	Distearoyl-L- $\alpha$ -glyceroylphosphoric acid	C <sub>39</sub> H <sub>77</sub> O <sub>8</sub> P	705.0	75.5-76.5	+3.8 in chloroform	v.s.die.eth., e.al., glac.ac.a.; s.acet., bz.
<b>Sphingomyelins</b>						
21	Palmitoylsphingomyelin	C <sub>39</sub> H <sub>81</sub> O <sub>7</sub> N <sub>2</sub> P	721.1	209-211		25
22	Stearoylsphingomyelin	C <sub>41</sub> H <sub>85</sub> O <sub>7</sub> N <sub>2</sub> P	749.1	209-210		
23	Lignocerylsphingomyelin	C <sub>47</sub> H <sub>97</sub> O <sub>7</sub> N <sub>2</sub> P	833.3	213-216		
<b>Ceramides</b>						
24	N-Lignoceryl-D-sphingosine	C <sub>42</sub> H <sub>83</sub> NO <sub>3</sub>	649.8	93-95	-2.0 in chloroform	24
25	N-Lignoceryl-D-dihydrosphingosine	C <sub>42</sub> H <sub>85</sub> NO <sub>3</sub>	651.9	102-103	0.0 in pyridine	
<b>Cerebrosides</b>						
26	N-Behenyl-D-sphingosyl- $\beta$ -D-glucopyranoside (Gaucher's cerebroside)	C <sub>46</sub> H <sub>89</sub> NO <sub>3</sub>	827.9	182-183	-7.6 in pyridine	24
27	N-Lignoceryl-D-sphingosyl- $\beta$ -D-galactopyranoside (cerasine)	C <sub>48</sub> H <sub>93</sub> NO <sub>8</sub>	811.9	182	-3.4 in pyridine	
28	N-Cerebronyl-D-sphingosyl- $\beta$ -D-galactopyranoside (phrenosine)	C <sub>48</sub> H <sub>93</sub> NO <sub>9</sub>	811.9	195	+4.4 in pyridine	

Contributor: O'Brien, John S.

**References:** [1] Baer, E. 1951. J. Biol. Chem. 189:235. [2] Baer, E., and D. Buchnea. 1958. Arch. Biochem. Biophys. 78:294. [3] Baer, E., and D. Buchnea. 1958. J. Biol. Chem. 232:895. [4] Baer, E., and D. Buchnea. 1959. J. Am. Chem. Soc. 81:1758. [5] Baer, E., D. Buchnea, and A. G. Newcombe. 1956. Ibid. 78:232. [6] Baer, E., and M. Katex. 1950. Ibid. 72:942. [7] Baer, E., and V. Mahadevan. 1959. Ibid. 81:249. [8] Baer, E., and J. Maurukas. 1952. Ibid. 74:152. [9] Baer, E., and J. Maurukas. 1955. J. Biol. Chem. 212:25. [10] Baer, E., and J. Maurukas. 1955. Ibid. 212:39. [11] Bowyer, D. E., et al. 1963. Biochim. Biophys. Acta 70:423. [12] Brockerhoff, H. 1961. Arch. Biochem. Biophys. 93:641. [13] Farquhar, J. W. 1962. Ibid. 60:80. [14] Hanahan, D. J., and M. E. Jayko. 1955. Biochem. Prepn. 4:12. [15] Hanahan, D. J., and J. N. Olley. 1958. J. Biol. Chem. 231:813. [16] Hanahan, D. J., R. M. Watts, and D. Pappajohn. 1960. J. Lipid Res. 1:421. [17] Kishimoto, Y., and N. S. Radin. 1959. Ibid. 1:72. [18] Marcus, A., et al. 1962. J. Clin. Invest. 41:2198. [19] O'Brien, J. S., D. L. Fillerup, and J. F. Mead. 1964. J. Lipid Res. 5:109. [20] O'Brien, J. S., D. L. Fillerup, and J. F. Mead. In press, 1964. [21] O'Brien, J. S., and G. Rouser. 1962. Federation Proc. 21:284. [22] Radin, N. S., and Y. Akahori. 1961. J. Lipid Res. 2:335. [23] Rapport, M., V. P. Skipski, and C. C. Sweeley. 1961. Ibid. 2:148. [24] Shapiro, D., and H. M. Flowers. 1961. J. Am. Chem. Soc. 83:3327. [25] Shapiro, D., H. M. Flowers, and S. Spector-Schefer. 1959. Ibid. 81:4360. [26] Trams, E. G., L. E. Guiffrida, and A. Karmen. 1962. Nature 193:680.



104. STEROLS: PHYSICAL AND CHEMICAL CHARACTERISTICS

	Substance (Systematic Name) <sup>1,2</sup>	Chemical Formula	Melt- ing Point °C	Spe- cific Rota- tion <sup>3</sup> [α] <sub>D</sub>	Source	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
1	(Δ <sup>3,5</sup> -Cholestadien-7-one)	C <sub>27</sub> H <sub>42</sub> O	112	-305	Testis, spleen (swine); sclerotic aorta	35,63,68
2	(Δ <sup>4,6</sup> -Cholestadien-3-one)	C <sub>27</sub> H <sub>42</sub> O	80	+35	Sclerotic aorta; spleen (swine)	34,63
3	(Δ <sup>5,7,22</sup> -Cholestatrien-3β-ol)	C <sub>27</sub> H <sub>42</sub> O	?	?	Shellfish	86
4	(Δ <sup>4</sup> -Cholesten-3-one)	C <sub>27</sub> H <sub>44</sub> O	81	+89	Feces; hypophysis, testis (swine)	62,64,65
5	7-Dehydrocholesterol (Δ <sup>5,7</sup> -Cholesta- dien-3β-ol)	C <sub>27</sub> H <sub>44</sub> O	150	-114	Cholesterol; skin (swine); snail	21,94
6	22-Dehydrocholesterol (Δ <sup>5,22</sup> -Choles- tadien-3β-ol)	C <sub>27</sub> H <sub>44</sub> O	135	-57	Shellfish; red algae	79,83
7	24-Dehydrocholesterol (Δ <sup>5,24(25)</sup> - Cholestadien-3β-ol)	C <sub>27</sub> H <sub>44</sub> O	117	-38	Barnacle	28
8	Zymosterol (Δ <sup>8,24</sup> -Cholestadien-3β-ol)	C <sub>27</sub> H <sub>44</sub> O	108	+47	Yeast	74,90,92
9	(Cholestane-3,6-dione)	C <sub>27</sub> H <sub>44</sub> O <sub>2</sub>	175	....	Testis (swine)	64
10	7-Ketocholesterol (Δ <sup>5</sup> -Cholesten-3β- ol-7-one)	C <sub>27</sub> H <sub>44</sub> O <sub>2</sub>	170	-104	Testis (cattle, swine)	63,64
11	Cholesterol <sup>4</sup> (Δ <sup>5</sup> -Cholesten-3β-ol)	C <sub>27</sub> H <sub>46</sub> O	149	-39	All animal cells; spinal cord; wool grease; red algae	50
12	Lathosterol <sup>4</sup> (Δ <sup>7</sup> -Cholesten-3β-ol)	C <sub>27</sub> H <sub>46</sub> O	122	+5.7	Skin; cholesterol	32,42
13	(Coprostan-3-one)	C <sub>27</sub> H <sub>46</sub> O	63	+36	Ambergris	49
14	(Δ <sup>4</sup> -Cholestene-3β,6β-diol)	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	258	+9.0	Spleen (swine)	63
15	7α-Hydroxycholesterol (Δ <sup>5</sup> -Cholestene- 3β,7α-diol)	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	184	-93	Sclerotic aorta; spleen (swine); serum (pregnant mare)	35,63,96
16	7β-Hydroxycholesterol (Δ <sup>5</sup> -Cholestene- 3β,7β-diol)	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	178	+7.2	Liver (cattle, swine); spleen (swine); serum (pregnant mare)	19,37,55,63
17	22-Hydroxycholesterol (Δ <sup>5</sup> -Cholestane- 3β,20α-diol)	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	186	-39	Lily	75,84
18	(Cholestan-3β-ol-6-one)	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	143	....	Spleen (swine)	63
19	(Cholestane-3β,5α-diol-6-one)	C <sub>27</sub> H <sub>46</sub> O <sub>3</sub>	236	....	Cholesterol; liver (swine)	71
20	Cholestanol <sup>4</sup> (Cholestan-3β-ol)	C <sub>27</sub> H <sub>48</sub> O	142	+24	Cholesterol; sclerotic aorta	54,69,70
21	Coprostanol (Coprostan-3β-ol)	C <sub>27</sub> H <sub>48</sub> O	101	+28	Feces	56
22	Epicoprostanol (Coprostan-3α-ol)	C <sub>27</sub> H <sub>48</sub> O	117	+32	Feces; ambergris	56
23	(Cholestane-3β,5,6β-triol)	C <sub>27</sub> H <sub>48</sub> O <sub>3</sub>	239	+3.2	Liver (cattle); testis (swine); sclerotic aorta	35,38,68
24	Dehydroergosterol (Δ <sup>5,7,9(11),22</sup> - Ergostatetraen-3β-ol)	C <sub>28</sub> H <sub>42</sub> O	146	+149	Yeast; ergot	67,93
25	14-Dehydroergosterol (Δ <sup>5,7,14,22</sup> - Ergostatetraen-3β-ol)	C <sub>28</sub> H <sub>42</sub> O	198	-396	Mold	8
26	24-Dehydroergosterol (Δ <sup>5,7,22,24(28)</sup> - Ergostatetraen-3β-ol)	C <sub>28</sub> H <sub>42</sub> O	118	-78	Yeast	23
27	Fungisterol (Δ <sup>7</sup> -Ergosten-3β-ol)	C <sub>28</sub> H <sub>44</sub> O	148	-0.2	Ergot	88
28	Ergosterol (Δ <sup>5,7,22</sup> -Ergostatrien- 3β-ol)	C <sub>28</sub> H <sub>44</sub> O	165	-130	Ergot; yeast; <i>Aspergillus niger</i>	20,29,39,57, 77,78
29	22-Dihydroergosterol (Δ <sup>5,7</sup> -Ergosta- dien-3β-ol)	C <sub>28</sub> H <sub>46</sub> O	153	-109	Ergot	66
30	Brassicasterol (Δ <sup>5,22</sup> -Ergostadien- 3β-ol)	C <sub>28</sub> H <sub>46</sub> O	148	-64	Rapeseed; invertebrates	16
31	24-Methylenecholesterol <sup>5</sup> (Δ <sup>5,24(28)</sup> - Ergostadien-3β-ol)	C <sub>28</sub> H <sub>46</sub> O	144	-42	Sponge; mollusks; honeybee	10,17,27,43, 44
32	5-Dihydroergosterol (Δ <sup>7,22</sup> -Ergosta- dien-3β-ol)	C <sub>28</sub> H <sub>46</sub> O	174	-20	Yeast	6,7,24
33	(Δ <sup>7,24(28)</sup> -Ergostadien-3β-ol)	C <sub>28</sub> H <sub>46</sub> O	131	+6.0	Starfish	29
34	Episterol (Δ <sup>7,24(28?)</sup> -Ergostadien- 3β-ol)	C <sub>28</sub> H <sub>46</sub> O	151	-5	Yeast	6,7,89,91
35	Ascosterol (Δ <sup>8,23(?)</sup> -Ergostadien- 3β-ol)	C <sub>28</sub> H <sub>46</sub> O	147	+45	Yeast	90-92
36	Fecosterol (Δ <sup>8,24(28)</sup> -Ergostadien- 3β-ol)	C <sub>28</sub> H <sub>46</sub> O	162	+42	Yeast	90,92
37	Cerevisterol (Δ <sup>7,22</sup> -Ergostadiene- 3β,5α,6β-triol)	C <sub>28</sub> H <sub>46</sub> O <sub>3</sub>	265	-79	Yeast; ergot	41,88
38	Haliclonasterol	C <sub>28</sub> H <sub>48</sub> O	141	-41.5	Sponge; green algae	11
39	Campesterol(Δ <sup>5</sup> -24-Isoergosten-3β-ol)	C <sub>28</sub> H <sub>48</sub> O	158	-33	Rapeseed; soybean; wheat germ	30

/1/ The numbers after the symbol "Δ" indicate the position of double bonds in the basic cyclopentano perhydrophenanthrene ring. /2/ Methyl sterols not included. /3/ Chloroform solvent for most determinations. /4/ Also isolated from invertebrates. /5/ Earlier preparations containing this substance referred to as chalinasterol and ostreasterol.

*continued*



104. STEROLS: PHYSICAL AND CHEMICAL CHARACTERISTICS

Substance (Systematic Name) <sup>1, 2</sup>	Chemical Formula	Melt- ing Point °C	Spe- cific Rota- tion <sup>3</sup> [α] <sub>D</sub>	Source	Reference
(A)	(B)	(C)	(D)	(E)	(F)
40 Neospongosterol (Δ <sup>22</sup> -24-Isoergosten-3β-ol)	C <sub>28</sub> H <sub>48</sub> O	153	+10	Sponge	14
41 Fucosterol (Δ <sup>5,24(28)</sup> -Stigmastadien-3β-ol)	C <sub>28</sub> H <sub>48</sub> O	124	-38	Brown algae	35,40
42 Aptostanol	C <sub>28</sub> H <sub>50</sub> O	135	+22	Sponge	18
43 Ergostanol (Ergostan-3β-ol)	C <sub>28</sub> H <sub>50</sub> O	143	+16	Plant	47
44 β-Sitosterol (Δ <sup>5</sup> -Stigmasten-3β-ol)	C <sub>29</sub> H <sub>50</sub> O	140	-36	Cottonseed; calycanthus seed; cinchona bark; wheat germ; rubber; invertebrates	3,5,25,51,52,87
45 Corbisterol (Δ <sup>5,7,22</sup> -Stigmastadien-3β-ol)	C <sub>29</sub> H <sub>46</sub> O	154	-114	Shellfish	58,80
46 Chondrillasterol (Δ <sup>7,22</sup> -24-Isoergostadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	164	-2	Green algae; sponge	12,15
47 Poriferasterol (Δ <sup>5,22</sup> -24-Isostigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	156	-49	Sponge; shellfish	53,69,81,85
48 Sargasterol (Δ <sup>5,24(28)</sup> -20-Isostigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	133.5	-47.5	Algae	82
49 Δ <sup>5</sup> -Avenasterol (Δ <sup>5,11(?)</sup> -Stigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	137	-37.6	Oats	46
50 Δ <sup>7</sup> -Avenasterol (Δ <sup>7,11(?)</sup> -Stigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	145	+8.8	Oats	46
51 Stigmasterol (Δ <sup>5,22</sup> -Stigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	170	-49	Calabar bean; soybean	61,95
52 α-Spinasterol (Δ <sup>7,22</sup> -Stigmastadien-3β-ol)	C <sub>29</sub> H <sub>48</sub> O	175	-2.7	Spinach; senega root; alfalfa; colocyath; starfish	26,31,34-36,73
53 Palysterol	C <sub>29</sub> H <sub>50</sub> O	140	-47	Sea anemone	13
54 Clionasterol (Δ <sup>5</sup> -24-Isostigmasten-3β-ol?)	C <sub>29</sub> H <sub>50</sub> O	138	-37	Sponge; shellfish	48,59,69,81
55 γ-Sitosterol (Δ <sup>5</sup> -24-Isostigmasten-3β-ol)	C <sub>29</sub> H <sub>50</sub> O	148	-43	Soybean; wheat germ; rye germ	5,9,22,33
56 (Δ <sup>7</sup> -Stigmasten-3β-ol)	C <sub>29</sub> H <sub>50</sub> O	145	+9	Wheat germ; starfish	45
57 Dihydrositosterol (Stigmastan-3β-ol)	C <sub>29</sub> H <sub>52</sub> O	140	+25	Grains	1-5,60
58 (Δ <sup>22</sup> -Stigmastan-3β-ol)	C <sub>29</sub> H <sub>52</sub> O	159	+3.3	Root	76
59 Dicholesteryl ether	C <sub>54</sub> H <sub>90</sub> O	196	-38	Spinal cord	72

<sup>1/1</sup> The numbers after the symbol "Δ" indicate the position of double bonds in the basic cyclopentano perhydrophenanthrene ring. <sup>2/2</sup> Methyl sterols not included. <sup>3/3</sup> Chloroform solvent for most determinations.

*Contributors:* (a) Idler, D. R., and Tamura, T., (b) Reich, Hans, (c) Kuck, Kathryn D.

*References:* [1] Anderson, R. J. 1924. J. Am. Chem. Soc. 46:1450. [2] Anderson, R. J., F. P. Nabenhauer, and R. L. Shriner. 1927. J. Biol. Chem. 71:389. [3] Anderson, R. J., and R. L. Shriner. 1926. J. Am. Chem. Soc. 48:2976. [4] Anderson, R. J., and R. L. Shriner. 1927. J. Biol. Chem. 71:401. [5] Anderson, R. J., R. L. Shriner, and G. O. Burr. 1926. J. Am. Chem. Soc. 48:2987. [6] Barton, D. H. R. 1945. J. Chem. Soc., p. 813. [7] Barton, D. H. R. 1946. Ibid., p. 512. [8] Barton, D. H. R., and T. Bruun. 1951. Ibid., p. 2728. [9] Bengtsson, B. E. 1935. Z. Physiol. Chem. 237:46. [10] Bergmann, W. 1934. J. Biol. Chem. 104:317, 553. [11] Bergmann, W., and R. J. Feeney. 1949. J. Org. Chem. 14:1078. [12] Bergmann, W., and R. J. Feeney. 1950. Ibid. 15:812. [13] Bergmann, W., R. J. Feeney, and A. N. Swift. 1951. Ibid. 16:1337. [14] Bergmann, W., D. H. Gould, and E. M. Low. 1945. Ibid. 10:570. [15] Bergmann, W., and F. H. McTigue. 1948. Ibid. 13:738. [16] Bergmann, W., and R. C. Ottke. 1949. Ibid. 14:1085. [17] Bergmann, W., H. P. Schedl, and E. M. Low. 1945. Ibid. 10:587. [18] Bergmann, W., et al. 1950. Ibid. 15:96. [19] Bergstrom, S., and O. Wintersteiner. 1941. J. Biol. Chem. 141:597. [20] Bills, C. E., O. N. Massengale, and P. S. Prickett. 1930. Ibid. 87:259. [21] Bock, F., and F. Wetter. 1938. Z. Physiol. Chem. 256:33. [22] Bonstedt, K. 1928. Ibid. 176:269. [23] Breivik, O. N., et al. 1954. J. Org. Chem. 19:1734. [24] Callow, R. K. 1931. Biochem. J. 25:87. [25] Cook,

*continued*

**104. STEROLS: PHYSICAL AND CHEMICAL CHARACTERISTICS**

- J. W., and M. F. C. Paige. 1944. *J. Chem. Soc.*, p. 336. [26] Dam, H., et al. 1939. *Helv. Chim. Acta* 22:310. [27] Fagerlund, U. H. M., and D. R. Idler. 1956. *J. Org. Chem.* 21:372. [28] Fagerlund, U. H. M., and D. R. Idler. 1957. *J. Am. Chem. Soc.* 79:6473. [29] Fagerlund, U. H. M., and D. R. Idler. 1959. *Ibid.* 81:401. [30] Fernholz, E., and H. B. MacPhillamy. 1941. *Ibid.* 63:1155. [31] Fernholz, E., and M. L. Moore. 1939. *Ibid.* 61:2467. [32] Fieser, L. F. 1951. *Ibid.* 73:5007. [33] Gloyer, S. W., and H. A. Schuette. 1939. *Ibid.* 61:1901. [34] Hamilton, B., and W. O. Kermack. 1952. *J. Chem. Soc.*, p. 5051. [35] Hardegger, E., L. Ruzicka, and E. Tagmann. 1943. *Helv. Chim. Acta* 26:2205. [36] Hart, M. C., and F. W. Heyl. 1932. *J. Biol. Chem.* 95:311. [37] Haslewood, G. A. D. 1939. *Biochem. J.* 33:709. [38] Haslewood, G. A. D. 1941. *Ibid.* 35:709. [39] Heiduschka, A., and H. Lindner. 1929. *Z. Physiol. Chem.* 181:15. [40] Heilbron, I., R. F. Phipers, and H. R. Wright. 1934. *J. Chem. Soc.*, p. 1572. [41] Honeywell, E. M., and C. E. Bills. 1932. *J. Biol. Chem.* 97:xxxix. [42] Idler, D. R., and C. A. Baumann. 1952. *Ibid.* 195:623. [43] Idler, D. R., and U. H. M. Fagerlund. 1955. *J. Am. Chem. Soc.* 77:4142. [44] Idler, D. R., and U. H. M. Fagerlund. 1957. *Chem. Ind. (London)*, p. 432. [45] Idler, D. R., A. A. Kandutsch, and C. A. Baumann. 1953. *J. Am. Chem. Soc.* 75:4325. [46] Idler, D. R., et al. 1953. *Ibid.* 75:1712. [47] Karrer, P., and W. Bürgi. 1951. *Helv. Chim. Acta* 34:832. [48] Kind, C. A., and W. Bergmann. 1942. *J. Org. Chem.* 7:341. [49] Lederer, E., et al. 1946. *Helv. Chim. Acta* 29:1354. [50] Lettré, H., and H. H. Inhoffen. 1936. *Über Sterine, Gallensäuren und verwandte Naturstoffe*. F. Enke, Stuttgart. p. 98. [51] Liebermann, C. 1884. *Ber. Deut. Chem. Ges.* 17:868. [52] Liebermann, C. 1885. *Ibid.* 18:1803. [53] Lyon, A. M., and W. Bergmann. 1942. *J. Org. Chem.* 7:428. [54] McArthur, C. S. 1942. *Biochem. J.* 36:559. [55] MacPhillamy, H. B. 1940. *J. Am. Chem. Soc.* 62:3518. [56] Marker, R. E., et al. 1942. *Ibid.* 64:818. [57] Massengale, O. N., C. E. Bills, and P. S. Prickett. 1931. *J. Biol. Chem.* 94:213. [58] Matsumoto, T., and T. Tamura. 1956. *Nippon Kagaku Zasshi* 77:1596. [59] Mazur, A. 1941. *J. Am. Chem. Soc.* 63:883, 2442. [60] Nabenhauer, F. P., and R. J. Anderson. 1926. *Ibid.* 48:2972. [61] Ott, A. C., and C. D. Ball. 1944. *Ibid.* 66:489. [62] Prelog, V., and H. C. Beyerman. 1945. *Experientia* 1:64. [63] Prelog, V., L. Ruzicka, and P. Stein. 1943. *Helv. Chim. Acta* 26:2222. [64] Prelog, V., et al. 1947. *Ibid.* 30:1080. [65] Rosenheim, O., and T. A. Webster. 1943. *Biochem. J.* 37:513. [66] Ruis, A. S. 1943. *Anales Real Acad. Farm.* 3:201. [67] Ruppel, E. 1942. *Bull. Soc. Chim. Biol.* 24:324. [68] Ruzicka, L., and V. Prelog. 1943. *Helv. Chim. Acta* 26:975. [69] Schönheimer, R., H. von Behring, and R. Hummel. 1930. *Z. Physiol. Chem.* 192:93. [70] Schönheimer, R., et al. 1930. *Ibid.* 192:73, 93. [71] Schwenk, E., N. T. Werthessen, and H. Rosenkrantz. 1952. *Arch. Biochem. Biophys.* 37:247. [72] Silberman, H., and S. Silberman-Martyncewa. 1945. *J. Biol. Chem.* 159:603. [73] Simpson, J. C. E. 1937. *J. Chem. Soc.*, p. 730. [74] Smedley-MacLean, I. 1928. *Biochem. J.* 22:22. [75] Stabursvik, A. 1953. *Acta Chem. Scand.* 7:1220. [76] Takeda, K. 1958. *Chem. Pharm. Bull. (Tokyo)* 6:536. [77] Tanret, C. 1908. *Ann. Chim. Phys. (Paris)* 8:15, 313. [78] Tanret, C. 1908. *Compt. Rend.* 147:75. [79] Tarr, H. L. A. 1958. *Ann. Rev. Biochem.* 27:236. [80] Toyama, Y., M. Kita, and T. Tanaka. 1952. *Bull. Chem. Soc. Japan* 25:355. [81] Toyama, Y., T. Takagi, and T. Tanaka. 1953. *Ibid.* 26:154. [82] Tsuda, K., et al. 1958. *J. Am. Chem. Soc.* 80:921. [83] Tsuda, K., et al. 1959. *Chem. Pharm. Bull. (Tokyo)* 7:747. [84] Tsuda, K., et al. 1959. *J. Am. Chem. Soc.* 81:5987. [85] Valentine, F. R., Jr., and W. Bergmann. 1941. *J. Org. Chem.* 6:452. [86] Van der Vliet, J. 1945. *Rec. Trav. Chim.* 67:265. [87] Wallis, E. S., and P. N. Chakravorty. 1938. *Ibid.* 2:335. [88] Wieland, H., and G. Coutelle. 1941. *Ann. Chem.* 548:270. [89] Wieland, H., and G. A. C. Gough. 1930. *Ibid.* 482:36. [90] Wieland, H., and Y. Kanaoka. 1937. *Ibid.* 530:146. [91] Wieland, H., F. Rath, and H. Hesse. 1941. *Ibid.* 548:34. [92] Wieland, H., et al. 1929. *Ibid.* 473:300. [93] Windaus, A. 1928. *Ibid.* 465:148. [94] Windaus, A., and F. Bock. 1937. *Z. Physiol. Chem.* 245:168. [95] Windaus, A., and A. Hauth. 1906. *Ber. Deut. Chem. Ges.* 39:4378. [96] Wintersteiner, O., and J. R. Ritzmann. 1940. *J. Biol. Chem.* 136:697.

## 105. PROTEINS: PHYSICAL AND CHEMICAL CHARACTERISTICS

**Method** (column C): l = light scattering; s = sedimentation velocity; d = diffusion constant; e = sedimentation equilibrium; o = osmotic pressure; x = X-ray diffraction; v = intrinsic viscosity; a = chemical analysis or combining ratio.

Protein	Source	Molecular Weight		Sedimentation Coefficient <sup>1</sup>	Partial Specific Volume <sup>2</sup>	Isoelectric Point pH <sup>3</sup> (Ionic Strength)	Reference
		Method	Value				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 G-Actin	Rabbit muscle	l	80,000	3.7	.....	4.5	89,108
2 Adrenocorticotrophic hormone	Sheep hypophysis	sd	20,000	2.0	0.75 <sup>4</sup>	4.6-4.7(0.1)	53,54
3 Aldolase	Rabbit muscle	sd?	147,000	7.9	0.740	6.05(0.2)-7.7(0.02)	6,119,120,128
4 Amandin	Almond	e	288,000	11.4	0.746	.....	116
5 α-Amylase	Barley malt	o	60,000	.....	.....	6	99
6 β-Amylase	Sweet potato	sd	150,000	8.9	0.749 <sup>4</sup>	4.74-4.79(0.1)	24
7 Bence-Jones	Human urine <sup>5</sup>	e; sd	35,000-37,000	3.55	0.749	5.18	115
8 Botulinum toxin, type A	<i>Clostridium botulinum</i>	sd	900,000 <sup>6</sup>	17	0.75	.....	90
9 Botulinum toxin, type B	<i>C. botulinum</i>	sd	500,000	.....	.....	.....	131
10 Bushy stunt virus	Tomato <sup>5</sup>	x	10,800,000	132	0.739	4.11(0.02)	7,51,65,73
11 Carbonic anhydrase	Cattle blood	sd	30,000	2.8	0.749 <sup>4</sup>	5.3(0.1)	85
12 Carboxypeptidase	Cattle pancreas	sd; vd	32,000-34,000	3.07	0.75 <sup>4</sup>	6.0(0.2)	91,106
13 Cardiotoxin	Cobra venom	d	46,000	.....	0.75 <sup>4</sup>	.....	94
14 Casein (caseinogen)	Cow milk	o	33,600	10.4	0.728	4.6(0.0025-0.01)	14,66,70,82
15 Catalase	Human blood	s; a	220,000	11.2-11.3	0.73 <sup>4</sup>	.....	17,37
16 Chorionic gonadotropin	Human urine <sup>7</sup>	sd	100,000	4.3	0.76	.....	63
17 α-Chymotrypsin	Cattle pancreas	sd; sv	22,500	2.4-2.7	0.73	8.1(0.1)-8.6(0.01)	5,96,98
18 α-Chymotrypsinogen	Cattle pancreas	sd; sv	22,500	2.5	0.72	9.5(0.01)	5,97
19 Colostrum globulin, immune	Cow colostrum	d	160,000-190,000	7	.....	5.85(0.1)	103,105
20 Conalbumin	Chicken egg white	a	70,000-77,000	.....	.....	6.8(0.1)	133
21 Concanavalin A	Jack bean	sd	96,000	6.0	0.73	.....	111
22 Crotoxin	Rattlesnake venom	e; sd	30,000	3.1	0.704	4.7(0.1)	32,56
23 Cytochrome-c	Cattle or horse heart	sd	16,000	1.9	0.707	10.65 <sup>6</sup> (0.1)	114,121-123
24 Diphtheria antitoxin	Horse plasma	sd	180,000	7.2	0.745 <sup>4</sup>	6.0	80,86
25 Diphtheria toxin	<i>Corynebacterium diphtheriae</i>	sd	74,000	4.6	0.736	4.1(0.005) <sup>9</sup>	79,86
26 Edestin	Hemp seed	sd	310,000	12.8	0.745	5.5 <sup>10</sup>	2,113,117
27 Enolase	Yeast	e	67,000	5.8	.....	.....	21
28 Excelsin	Brazil nut	o; e	210,000	11.8	0.743	.....	13,116
29 Fetuin	Fetal calf or sheep blood	sd	50,000	3.1-3.4	0.70	3.5(0.2)	84
30 Fibrinogen	Human blood	o; sv	400,000-580,000	9	.....	5.4(0.1)	76,109
31 Fumarase	Swine heart	e	220,000	9.1	.....	.....	21
32 Gelatin	Collagenous tissues	o	5,000-400,000	.....	.....	4.9(0.01)	1,5
33 Gliadin	Wheat	e	27,000	2.1	0.71	6.5 <sup>10</sup>	46,118
34 Globin	Horse blood	a	31,000 <sup>11</sup>	2.5	0.749	7.5(0.1) <sup>12</sup>	31,72
35 α-Globulin	Barley	sd	26,000	2.5	0.72	5.0(0.1)	92
36 γ-Globulin	Barley	sd	170,000	8.3	.....	5.7(0.1)	92
37 Gluten	Wheat	e	39,000-4,600,000	2.5	0.700	7	64,118
38 α-Glycerolphosphate dehydrogenase	Rabbit muscle	e	78,000-87,000	4.9	.....	.....	21
39 Glycerol-3-phosphate dehydrogenase	Rabbit muscle	e	140,000	7.7	.....	.....	21

<sup>1</sup>/ Specific sedimentation velocity in units of 10<sup>-13</sup>, under standard conditions of water at 20°C. <sup>2</sup>/ Cubic centimeters increase in volume of solution per gram of protein dissolved. <sup>3</sup>/ pI at which protein does not move in an electric field. <sup>4</sup>/ Assumed value. <sup>5</sup>/ Pathological. <sup>6</sup>/ At pH 7.5, dissociation product has molecular weight of 70,000 [130,132]. <sup>7</sup>/ During pregnancy. <sup>8</sup>/ At 0°C for the oxidized (ferri-) form. <sup>9</sup>/ By cataphoresis. <sup>10</sup>/ Based on solubility minimum. <sup>11</sup>/ Composed of two approximately equal subunits. <sup>12</sup>/ For human globin.

continued



105. PROTEINS: PHYSICAL AND CHEMICAL CHARACTERISTICS

Protein	Source	Molecular Weight		Sedimentation Coefficient <sup>1</sup>	Partial Specific Volume <sup>2</sup>	Isoelectric Point pH <sup>3</sup> (Ionic Strength)	Reference	
		Method	Value					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
40	Growth hormone	Beef hypophysis	a; sd	39,000-49,000	3.1-3.7	0.76	6.85(0.1)	52,55,59,107
41	Hemocyanin	<i>Helix pomatia</i>	sd	8,900,000	103	0.738	5.05(0.02)	11,125
42	Hemoglobin A <sup>13</sup>	Human blood	a	64,450	4.5	0.749 <sup>14</sup>	6.87(0.1)	10,31,38,81,95,114
43	Hexokinase	Baker's yeast	sd	97,000	3.1	0.740 <sup>4</sup>	4.5-4.8(0.02 M) <sup>9</sup>	47
44	Insulin <sup>15</sup>	Beef pancreas	o; sd	36,000	3.5	0.707	5.2(0.033)	27,33,39,77,102,136
45	$\alpha$ -Lactalbumin	Cow milk	sd	15,000	1.75	0.735	.....	30
46	Lactate dehydrogenase	Rabbit muscle	e	135,000	6.9	.....	.....	21
47	Lactogenic hormone	Cattle or sheep hypophysis	o	26,500	.....	.....	5.7(0.05)	57,58
48	$\beta$ -Lactoglobulin A <sup>16</sup>	Cow milk	x	17,500	2.25	0.751	5.09	12,45,83,101,126,127
49	Lactoglobulin, immune	Cow milk	sd	180,000	7	.....	5.8(0.1)	4,104
50	$\alpha$ -Lipovitellin	Egg yolk	sd	400,000	10.9	0.777	.....	8
51	$\beta$ -Lipovitellin	Egg yolk	sd	400,000	10.4	0.777	.....	8
52	$\alpha$ -Livetin	Egg yolk	s	67,000	4.4	.....	.....	69
53	$\beta$ -Livetin	Egg yolk	s	45,000	3.0	.....	.....	69
54	$\gamma$ -Livetin	Egg yolk	sd	151,000	7.6	0.726	.....	68
55	Lysozyme chloride	Chicken egg white	x	13,900	2.1	0.722	11.35(0.1)	5,78,134
56	Metakentrin	Sheep hypophysis	o; a	40,000	3.6	.....	4.6(0.1)	60
57	Myosin	Rabbit muscle	sd; o	840,000-858,000	7.1	.....	5.4(0.1-0.5)	25,88,89
58	Old yellow enzyme	Brewer's yeast	sd; e	105,000	5.8	0.753	5.22(0.02)	44,124
59	Ovalbumin	Chicken egg	sd	44,000	3.55	0.749	4.58(0.1)	49,61,114
60	Pepsin	Swine gastric mucosa	x	33,000-37,000	3.3	.....	2.75-3.0	74,75,87
61	Phosvitin	Egg yolk	sd	30,900	3.14	0.545	.....	41
62	Relaxin	Pregnant sow ovary	a; s	9,000	.....	.....	.....	29
63	Ribonuclease	Cattle pancreas	x	13,400	1.85	0.709	7.8(0.055)	4,15,93
64	Ricin	Castor bean	sd	77,000-85,000	4.8-5.3	0.75 <sup>4</sup>	5.2-5.5	43
65	Salmine	Salmon testes	a	7,000	<1	.....	12	52,71,129
66	Serum albumin	Human blood	x	65,600	4.6	0.733	4.9	18,62,76
67	$\alpha_1$ -Serum globulin	Human blood	sv	200,000	5.0	0.841	.....	76
68	$\beta_1$ -Serum globulin	Human blood	sv	90,000	5.5	0.725	.....	76
69	$\gamma$ -Serum globulin	Human blood	sv	160,000	7.2	0.739	5.7(0.1) <sup>17</sup>	3,19,76
70	$\gamma_1$ -Serum globulin antipneumococcus	Human blood	sd	190,000	7.4	0.745 <sup>4</sup>	5.6(0.1) <sup>16</sup>	22,42
71	Tetanus toxin	<i>Clostridium tetani</i>	a; s	67,000	4.5	.....	.....	23
72	Thymus nucleohistone	Calf thymus	e; sd	2,000,000	31.0	0.658	.....	16
73	Thyroglobulin	Swine thyroid	sd	700,000	19.2	0.72	4.58(0.02)	36
74	Tobacco mosaic virus	Tobacco leaves <sup>5</sup>	sd; sv; vd	33,000,000	185	0.72	3.49(0.02)	26,50
75	Trypsin	Cattle pancreas	a	20,700	2.5 <sup>13</sup>	.....	10.8(0.03 M)	9,20,40
76	Trypsin inhibitor	Cattle pancreas	o	6,000	.....	.....	.....	48
77	Tuberculin protein	<i>Mycobacterium tuberculosis</i>	sd	32,000	1.9-4.9	.....	4.3(0.03)	100
78	Tyrosinase	<i>Pseudomonas campestris</i>	sd	100,000	6.4	0.75 <sup>4</sup>	<5	67
79	Urease	Jack bean	sd	480,000	18.6	0.73	5.0(0.012) <sup>1c</sup>	5,110,112
80	Zein	Corn	sd	40,000	1.9	0.73 <sup>4</sup>	.....	135

/1/ Specific sedimentation velocity in units of  $10^{-13}$ , under standard conditions of water at 20°C. /2/ Cubic centimeters increase in volume of solution per gram of protein dissolved. /3/ pH at which protein does not move in an electric field. /4/ Assumed value. /5/ Pathological. /6/ By cataphoresis. /7/ Based on solubility minimum. /8/ Several well-defined protein subunits, of which the commonest are the  $\alpha$  chain (molecular weight = 15,128) and the  $\beta$  chain (molecular weight = 15,870). /9/ For horse blood. /10/ The A component molecular weight = 5,733, and sedimentation constant = 1.2 [28, 34, 35]. /11/ Two types, A and B, having same molecular weight and sedimentation constants. Isoelectric point for  $\beta$ -lactoglobulin B = 5.23. /12/ Depends on fraction employed. /13/ Hyperimmune horse blood. /14/ Value for diisopropylphosphate derivative.

continued



**105. PROTEINS: PHYSICAL AND CHEMICAL CHARACTERISTICS**

*Contributors:* (a) Ward, Wilfred H., (b) Deutsch, Marshall E., (c) Evans, Robert John, (d) McMeekin, T. L.

- References:* [1] Abribat, M., J. Pouradier, and A. M. Venet. 1949. *J. Polymer Sci.* 4:523. [2] Adair, G. S., and M. E. Adair. 1936. *Proc. Roy. Soc. (London), B*, 120:422. [3] Alberty, R. A. 1949. *J. Phys. Colloid Chem.* 53:114. [4] Alberty, R. A., E. A. Anderson, and J. W. Williams. 1948. *Ibid.* 52:217. [5] Anderson, E. A., and R. A. Alberty. 1948. *Ibid.* 52:1345. [6] Baranowski, T., and T. R. Niederland. 1949. *J. Biol. Chem.* 180:543. [7] Bernal, J. D., and I. Fankuchen. 1941. *J. Gen. Physiol.* 25:111. [8] Bernardi, G., and W. H. Cook. 1960. *Biochim. Biophys. Acta* 44:96. [9] Bier, M., and F. F. Nord. 1951. *Arch. Biochem. Biophys.* 33:320. [10] Braunitzer, G., et al. 1961. *Z. Physiol. Chem.* 325:283. [11] Brohult, S. 1947. *J. Phys. Colloid Chem.* 51:206. [12] Bull, H. B. 1946. *J. Am. Chem. Soc.* 68:745. [13] Burk, N. F. 1937. *J. Biol. Chem.* 120:63. [14] Burk, N. F., and D. M. Greenberg. 1930. *Ibid.* 87:197. [15] Carlisle, C. H., and H. Scouloudi. 1951. *Proc. Roy. Soc. (London), A*, 207:496. [16] Carter, R. O. 1941. *J. Am. Chem. Soc.* 63:1960. [17] Cecil, R., and A. G. Ogston. 1948. *Biochem. J.* 43:205. [18] Cohn, E. J., W. L. Hughes, Jr., and J. H. Weare. 1947. *J. Am. Chem. Soc.* 69:1753. [19] Cohn, M., H. F. Deutsch, and L. R. Wetter. 1950. *J. Immunol.* 64:381. [20] Cunningham, L. W., Jr., et al. 1953. *Discussions Faraday Soc.* 13:58. [21] Deal, W. C., et al. 1963. *Biochem. Biophys. Res. Commun.* 10:49. [22] Deutsch, H. F., and J. C. Nichol. 1948. *J. Biol. Chem.* 176:797. [23] Dunn, M. S., M. N. Camien, and L. Pillemer. 1949. *Arch. Biochem.* 22:374. [24] Englard, S., and T. P. Singer. 1950. *J. Biol. Chem.* 187:213. [25] Erdős, T., and O. Snellman. 1948. *Biochim. Biophys. Acta* 2:642. [26] Eriksson-Quensel, I.-B., and T. Svedberg. 1936. *J. Am. Chem. Soc.* 58:1863. [27] Fischer, R. H., and C. S. Vestling. 1955. *Ibid.* 77:5703. [28] Fredericq, E. 1953. *Nature* 171:570. [29] Frieden, E. H. 1951. *Arch. Biochem.* 30:138. [30] Gordon, W. G., and W. F. Semmett. 1953. *J. Am. Chem. Soc.* 75:328. [31] Gralén, N. 1939. *Biochem. J.* 33:1907. [32] Gralén, N., and T. Svedberg. 1938. *Ibid.* 32:1375. [33] Gutfreund, H. 1948. *Ibid.* 42:544. [34] Harfenist, E. J. 1953. *J. Am. Chem. Soc.* 75:5528. [35] Harfenist, E. J., and L. C. Craig. 1952. *Ibid.* 74:3087. [36] Heidelberger, M., and K. O. Pedersen. 1935. *J. Gen. Physiol.* 19:95. [37] Herbert, D., and J. Pinsent. 1948. *Biochem. J.* 43:203. [38] Hill, R. J., and W. Konigsberg. 1961. *J. Biol. Chem.* 236:PC7. [39] Howitt, F. O., and E. B. R. Prideaux. 1932. *Proc. Roy. Soc. (London), B*, 112:13. [40] Jansen, E. F., and A. K. Balls. 1952. *J. Biol. Chem.* 194:721. [41] Joubert, F. J., and W. H. Cook. 1958. *Can. J. Biochem. Physiol.* 36:399. [42] Kabat, E. A. 1939. *J. Exptl. Med.* 69:103. [43] Kabat, E. A., M. Heidelberger, and A. E. Bezer. 1947. *J. Biol. Chem.* 168:629. [44] Kekwick, R. A., and K. O. Pedersen. 1936. *Biochem. J.* 30:2201. [45] Klostergaard, H., and R. A. Pasternak. 1957. *J. Am. Chem. Soc.* 79:5671. [46] Krejci, L., and T. Svedberg. 1935. *Ibid.* 57:946. [47] Kunitz, M., and M. R. McDonald. 1946. *J. Gen. Physiol.* 29:393. [48] Kunitz, M., and J. H. Northrop. 1936. *Ibid.* 19:991. [49] Lamm, O., and A. Polson. 1936. *Biochem. J.* 30:528. [50] Lauffer, M. A. 1944. *J. Am. Chem. Soc.* 66:1188. [51] Lauffer, M. A., and W. M. Stanley. 1940. *J. Biol. Chem.* 135:463. [52] Li, C. H. 1947. *Ann. Rev. Biochem.* 16:291. [53] Li, C. H. 1952. *Arch. Biochem. Biophys.* 36:462. [54] Li, C. H., H. M. Evans, and M. E. Simpson. 1943. *J. Biol. Chem.* 149:413. [55] Li, C. H., H. M. Evans, and M. E. Simpson. 1945. *Ibid.* 159:353. [56] Li, C. H., and H. Fraenkel-Conrat. 1942. *J. Am. Chem. Soc.* 64:1586. [57] Li, C. H., W. R. Lyons, and H. M. Evans. 1940. *Ibid.* 62:2925. [58] Li, C. H., W. R. Lyons, and H. M. Evans. 1941. *J. Biol. Chem.* 140:43. [59] Li, C. H., and M. Moskowitz. 1949. *Ibid.* 178:203. [60] Li, C. H., M. E. Simpson, and H. M. Evans. 1942. *J. Am. Chem. Soc.* 64:367. [61] Longworth, L. G. 1941. *Ann. N. Y. Acad. Sci.* 41:267. [62] Low, B. W. 1952. *J. Am. Chem. Soc.* 74:4830. [63] Lundgren, H. P., et al. 1942. *J. Biol. Chem.* 142:367. [64] McCalla, A. G., and N. Gralén. 1945. *Can. J. Res., C*, 20:130. [65] McFarlane, A. S., and R. A. Kekwick. 1938. *Biochem. J.* 32:1607. [66] McMeekin, T. L., and K. Marshall. 1952. *Science* 116:142. [67] Mallette, M. F., and C. R. Dawson. 1949. *Arch. Biochem.* 23:29. [68] Martin, W. G., and W. H. Cook. 1958.

*continued*

105. PROTEINS: PHYSICAL AND CHEMICAL CHARACTERISTICS

- Can. J. Biochem. Physiol. 36:153. [69] Martin, W. G., J. E. Vandegaer, and W. H. Cook. 1957. *Ibid.* 35:241. [70] Michaelis, L., and H. Pechstein. 1912. *Biochem. Z.* 47:260. [71] Miyake, S. 1927. *Z. Physiol. Chem.* 172:225. [72] Munro, M. P., and F. L. Munro. 1943. *J. Biol. Chem.* 150:427. [73] Neurath, H., and G. R. Cooper. 1940. *Ibid.* 135:455. [74] Northrop, J. H. 1930. *J. Gen. Physiol.* 13:739. [75] Northrop, J. H. 1930. *Ibid.* 13:767. [76] Oncley, J. L., G. Scatchard, and A. Brown. 1947. *J. Phys. Colloid Chem.* 51:184. [77] Oncley, J. L., et al. 1952. *J. Phys. Chem.* 56:85. [78] Palmer, K. J., M. Ballantyne, and J. A. Galvin. 1948. *J. Am. Chem. Soc.* 70:906. [79] Pappenheimer, A. M., Jr. 1942. *J. Bacteriol.* 43:273. [80] Pappenheimer, A. M., Jr., H. P. Lundgren, and J. W. Williams. 1940. *J. Exptl. Med.* 71:247. [81] Pauling, L., et al. 1949. *Science* 110:543. [82] Pedersen, K. O. 1936. *Biochem. J.* 30:948. [83] Pedersen, K. O. 1936. *Ibid.* 30:961. [84] Pedersen, K. O. 1947. *J. Phys. Colloid Chem.* 51:164. [85] Petermann, M. L., and N. V. Hakala. 1942. *J. Biol. Chem.* 145:701. [86] Petermann, M. L., and A. M. Pappenheimer, Jr. 1941. *J. Phys. Chem.* 45:1. [87] Philpot, J. St. L. 1935. *Biochem. J.* 29:2458. [88] Portzehl, H. 1950. *Z. Naturforsch.* 5b:75. [89] Portzehl, H., G. Schramm, and H. H. Weber. 1950. *Ibid.* 5b:61. [90] Putnam, F. W., C. Lammana, and D. G. Sharp. 1948. *J. Biol. Chem.* 176:401. [91] Putnam, F. W., et al. 1946. *Ibid.* 166:603. [92] Quensel, O. 1942. *Untersuchungen über die Gerstenglobuline.* Almqvist and Wiksell, Stockholm. [93] Rothen, A. 1940. *J. Gen. Physiol.* 24:203. [94] Sarker, N. K. 1947. *J. Indian Chem. Soc.* 24:61. [95] Schroeder, A., et al. 1961. *Proc. Natl. Acad. Sci. U.S.* 47:811. [96] Schwert, G. W. 1949. *J. Biol. Chem.* 179:655. [97] Schwert, G. W. 1951. *Ibid.* 190:799. [98] Schwert, G. W., and S. Kauffman. 1951. *Ibid.* 190:807. [99] Schwimmer, S., and A. K. Balls. 1949. *Ibid.* 179:1063. [100] Seibert, F. B., K. O. Pedersen, and A. Tiselius. 1938. *J. Exptl. Med.* 68:413. [101] Senti, F. R., and R. C. Warner. 1948. *J. Am. Chem. Soc.* 70:3318. [102] Sjögren, B., and T. Svedberg. 1931. *Ibid.* 53:2657. [103] Smith, E. L. 1946. *J. Biol. Chem.* 164:345. [104] Smith, E. L. 1946. *Ibid.* 165:665. [105] Smith, E. L., and D. M. Brown. 1950. *Ibid.* 183:241. [106] Smith, E. L., D. M. Brown, and H. T. Hanson. 1949. *Ibid.* 180:33. [107] Smith, E. L., et al. 1949. *Ibid.* 177:305. [108] Steiner, R. F., K. Laki, and S. Spicer. 1952. *J. Polymer Sci.* 8:23. [109] Stenhagen, E. 1938. *Biochem. J.* 32:714. [110] Sumner, J. B., N. Gralén, and I.-B. Eriksson-Quensel. 1938. *J. Biol. Chem.* 125:37. [111] Sumner, J. B., N. Gralén, and I.-B. Eriksson-Quensel. 1938. *Ibid.* 125:45. [112] Sumner, J. B., and D. B. Hand. 1929. *J. Am. Chem. Soc.* 51:1255. [113] Svedberg, T. 1937. *Nature* 139:1051. [114] Svedberg, T., and K. O. Pedersen. 1940. *The ultracentrifuge.* Clarendon Press, Oxford. [115] Svedberg, T., and B. Sjögren. 1929. *J. Am. Chem. Soc.* 51:3594. [116] Svedberg, T., and B. Sjögren. 1930. *Ibid.* 52:279. [117] Svedberg, T., and A. J. Stamm. 1929. *Ibid.* 51:2170. [118] Tague, E. L. 1925. *Ibid.* 47:418. [119] Taylor, J. F. 1950. *Federation Proc.* 9:237. [120] Taylor, J. F., A. A. Green, and G. T. Cori. 1948. *J. Biol. Chem.* 173:591. [121] Theorell, H. 1935. *Biochem. Z.* 279:463. [122] Theorell, H. 1936. *Ibid.* 285:207. [123] Theorell, H., and Å. Åkeson. 1941. *J. Am. Chem. Soc.* 63:1804. [124] Theorell, H., and Å. Åkeson. 1956. *Arch. Biochem. Biophys.* 65:439. [125] Tiselius, A. 1930. *Nova Acta Regiæ Soc. Sci. Upsaliensis*, IV, 7:1. [126] Townend, R., C. A. Kiddy, and S. N. Timasheff. 1961. *J. Am. Chem. Soc.* 83:1419. [127] Townend, R., and S. N. Timasheff. 1957. *Ibid.* 79:3613. [128] Velick, S. F. 1949. *J. Phys. Colloid Chem.* 53:135. [129] Velick, S. F., and S. Udenfriend. 1951. *J. Biol. Chem.* 191:233. [130] Wagman, J. 1954. *Arch. Biochem. Biophys.* 50:104. [131] Wagman, J., and J. B. Bateman. 1951. *Ibid.* 31:424. [132] Wagman, J., and J. B. Bateman. 1953. *Ibid.* 35:375. [133] Warner, R. C., and I. Weber. 1951. *J. Biol. Chem.* 191:173. [134] Wetter, L. R., and H. F. Deutsch. 1951. *Ibid.* 192:237. [135] Williams, J. W., and C. C. Watson. 1938. *Cold Spring Harbor Symp. Quant. Biol.* 6:208. [136] Wintersteiner, O., and H. A. Abramson. 1933. *J. Biol. Chem.* 99:741.

## 106. AMINO ACIDS: PHYSICAL AND CHEMICAL CHARACTERISTICS

**Solubility** (columns E and F): a. = acid; ac. = acetic; acet. = acetone; al. = alcohol; alk. = alkali; aq. = aqueous; deliq. = deliquescent; dil. = dilute; e. = ethyl; eth. = ether; i. = insoluble; me. = methyl; s. = soluble; sl. = slightly; v. = very.

Amino Acid	Chemical Formula	Molecular Weight	Melting Point <sup>1</sup> °C	Solubility <sup>2</sup>		Specific Rotation				Isoelectric Point pH
				Water 25°C	Other Solvents	Solvent	g/100 ml	Temp. °C	[α] <sub>D</sub>	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
1 L-Alanine	C <sub>3</sub> H <sub>7</sub> O <sub>2</sub> N	89.09	297	16.51	sl. s. al.; i. acet., eth.	1.0 N HCl	5.79	15	+14.7	6.11 <sup>3</sup>
2 β-Alanine	C <sub>3</sub> H <sub>7</sub> O <sub>2</sub> N	89.09	196	v. s.	v. sl. s. al.; i. eth.	.....	.....	...	0	6.90
3 L-2-Aminobutyric acid	C <sub>4</sub> H <sub>9</sub> O <sub>2</sub> N	103.12	285	28	0.18, al.; i. eth.	20% HCl	.....	20	+14.1	5.98
4 L-Anserine	C <sub>10</sub> H <sub>16</sub> O <sub>3</sub> N <sub>4</sub>	240.26	238-239	s.	s. me. al.; sl. s. e. al.	H <sub>2</sub> O	5.0	20	+12.2	8.27
5 L-Arginine	C <sub>6</sub> H <sub>14</sub> O <sub>2</sub> N <sub>4</sub>	174.20	238	v. s.	i. al., eth.	6.0 N HCl	1.65	23	+26.9	10.76
6 L-Asparagine	C <sub>4</sub> H <sub>8</sub> O <sub>3</sub> N <sub>2</sub>	132.12	236	2.46	s. dil. NH <sub>4</sub> OH; v. sl. s. al.; i. eth.	3.4 N HCl	2.24	20	+34.3	5.41
7 L-Aspartic acid	C <sub>4</sub> H <sub>7</sub> O <sub>4</sub> N	133.10	269-271	0.50	s. dil. HCl; v. sl. s. al.; i. eth.	6.0 N HCl	2.0	24	+24.6	2.98
8 L-Canaline	C <sub>4</sub> H <sub>10</sub> O <sub>3</sub> N <sub>2</sub>	134.14	214	s.	.....	H <sub>2</sub> O	1.6	21	-8.1	....
9 L-Canavanine	C <sub>5</sub> H <sub>12</sub> O <sub>3</sub> N <sub>4</sub>	176.18	184	s.	.....	H <sub>2</sub> O	3.2	20	+8.1	8.2
10 L-Carnosine	C <sub>6</sub> H <sub>14</sub> O <sub>3</sub> N <sub>4</sub>	226.23	246-250	s.	.....	H <sub>2</sub> O	2.0	20	+20.5	8.17
11 L-Citrulline	C <sub>6</sub> H <sub>13</sub> O <sub>3</sub> N <sub>3</sub>	175.19	222	v. sl. s.	i. al.	1.0 N HCl	2.0	27	+24.3	5.92
12 L-Cystathionine	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub> N <sub>2</sub> S	222.26	270-312	.....	s. HCl	1.0 N HCl	1.0	22	+23.7	....
13 L-Cysteic acid	C <sub>3</sub> H <sub>7</sub> O <sub>5</sub> NS	169.17	260	s.	s. a., alk.; i. al.	H <sub>2</sub> O	.....	...	+8.7	1.6
14 L-Cysteine	C <sub>3</sub> H <sub>7</sub> O <sub>2</sub> NS	121.16	175-178	v. s.	s. a., alk.	H <sub>2</sub> O	2.0	21	-10.1	5.07
15 L-Cystine	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub> N <sub>2</sub> S <sub>2</sub>	240.30	258-261	0.011	s. a. <sup>4</sup> , NH <sub>4</sub> OH; i. al., eth.	1.0 N HCl	1.0	24	-214.4	5.02
16 L-3,5-Dibromotyrosine	C <sub>9</sub> H <sub>9</sub> O <sub>3</sub> NBr <sub>2</sub>	338.99	245 <sup>5</sup>	.....	.....	0.3 N HCl	.....	20	-2.4	4.30
17 L-3,4-Dihydroxyphenylalanine	C <sub>9</sub> H <sub>11</sub> O <sub>4</sub> N	197.19	280	0.50	s. a., alk.; i. al., eth.	4% HCl	1.0	25	-12.0	....
18 L-3,5-Diiodotyrosine	C <sub>9</sub> H <sub>9</sub> O <sub>3</sub> NI <sub>2</sub>	432.99	194	0.62	.....	1.1 N HCl	5.1	20	+2.9	4.29 <sup>3</sup>
19 L-Djenkolic acid	C <sub>7</sub> H <sub>14</sub> O <sub>4</sub> N <sub>2</sub> S <sub>2</sub>	254.33	300-350	0.10	.....	1% HCl	2.0	26	-44.5	....
20 L-Ergothioneine	C <sub>9</sub> H <sub>15</sub> O <sub>2</sub> N <sub>3</sub> S	229.30	290	.....	.....	H <sub>2</sub> O	5.0	21	+116.0	....
21 L-Ethionine	C <sub>6</sub> H <sub>13</sub> O <sub>2</sub> NS	163.24	272-284	s.	.....	0.2 N HCl	0.8	25	+23.5	....
22 L-Glutamic acid	C <sub>5</sub> H <sub>9</sub> O <sub>4</sub> N	147.13	247	0.86	.....	6.0 N HCl	1.0	22	+31.2	3.22 <sup>3</sup>
23 L-Glutamine	C <sub>5</sub> H <sub>10</sub> O <sub>3</sub> N <sub>2</sub>	146.15	185-186	4.25	v. sl. s. al.; i. eth.	H <sub>2</sub> O	.....	19	+8.0	5.65
24 Glycine	C <sub>2</sub> H <sub>5</sub> O <sub>2</sub> N	75.07	233	24.99	0.043, 90% al.	.....	.....	...	0	6.20
25 L-Histidine	C <sub>6</sub> H <sub>9</sub> O <sub>2</sub> N <sub>3</sub>	155.16	277	4.19	v. sl. s. al.; i. eth.	H <sub>2</sub> O	1.1	25	-39.0	7.64
26 L-Homocysteine	C <sub>4</sub> H <sub>9</sub> O <sub>2</sub> NS	135.19	232-233 <sup>3</sup>	s.	.....	.....	.....	...	.....	.....
27 L-Homocystine	C <sub>8</sub> H <sub>16</sub> O <sub>4</sub> N <sub>2</sub> S <sub>2</sub>	268.36	260-265 <sup>3</sup>	v. sl. s.	.....	1.0 N HCl	1.0	26	+77	5.53
28 L-δ-Hydroxylysine	C <sub>6</sub> H <sub>14</sub> O <sub>3</sub> N <sub>2</sub>	162.20	220	s.	s. a.; i. al.	6.0 N HCl	.....	25	+17.8	9.15
29 L-γ-Hydroxyproline	C <sub>5</sub> H <sub>9</sub> O <sub>3</sub> N	131.13	238-241	36.11	v. sl. s. al.; i. eth.	H <sub>2</sub> O	1.0	22	-75.2	5.82
30 L-Isoleucine	C <sub>6</sub> H <sub>13</sub> O <sub>2</sub> N	131.18	283-284	4.12	0.09, al.; s. hot ac. a.; i. eth.	6.1 N HCl	5.1	20	+40.6 <sup>5</sup>	6.04 <sup>3</sup>

<sup>1/1</sup> Most amino acids decompose when melting. <sup>2/2</sup> Grams of amino acid soluble in 100 ml of solvent. <sup>3/3</sup> Racemic mixture (DL). <sup>4/4</sup> Mixture of acetonitrile and perchloric acid. <sup>5/5</sup> Dihydrate.

*continued*

106. AMINO ACIDS: PHYSICAL AND CHEMICAL CHARACTERISTICS

Amino Acid	Chemical Formula	Molec- ular Weight	Melting Point <sup>1</sup> °C	Solubility <sup>2</sup>		Specific Rotation				Iso- elec- tric Point pH
				Water 25°C	Other Solvents	Solvent	g/100 ml	Temp. °C	[α] <sub>D</sub>	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
31 L-Lanthionine	C <sub>6</sub> H <sub>12</sub> O <sub>4</sub> N <sub>2</sub> S	208.24	245-295	i.	s. aq. NH <sub>3</sub> , aq. HCl	2.4 N NaOH	5.0	22	+8.6	....
32 L-Leucine	C <sub>6</sub> H <sub>13</sub> O <sub>2</sub> N	131.18	295	2.19	0.022, al.; s. ac. a.; i. eth.	6.0 N HCl	2.0	26	+15.1	6.04 <sup>3</sup>
33 L-Lysine	C <sub>6</sub> H <sub>14</sub> O <sub>2</sub> N <sub>2</sub>	146.19	224	v. s.	v. sl. s. al.; i. eth.	6.0 N HCl	2.0	23	+25.9	9.47
34 L-Methionine	C <sub>5</sub> H <sub>11</sub> O <sub>2</sub> NS	149.21	283	5.75	i. eth.	0.2 N HCl	0.8	25	+21.2	5.74 <sup>3</sup>
35 L-Norleucine	C <sub>6</sub> H <sub>13</sub> O <sub>2</sub> N	131.18	301	1.149 <sup>3</sup>	0.017, al. <sup>3</sup>	6.0 N HCl	4.3	20	+21.3	6.08 <sup>3</sup>
36 L-Norvaline	C <sub>5</sub> H <sub>11</sub> O <sub>2</sub> N	117.15	291-292	10.7 <sup>6</sup>	sl. s. al.; i. eth.	20% HCl	5	20	+22.8	6.04
37 D-Octapine	C <sub>9</sub> H <sub>18</sub> O <sub>4</sub> N <sub>4</sub>	246.27	229-230	s.	.....	H <sub>2</sub> O	.....	17	+20.9	5.51
38 L-Ornithine	C <sub>5</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub>	132.16	225	v. de- liq.	v. s. al.; sl. s. eth.	H <sub>2</sub> O	4.0	27	+16.5 <sup>6</sup>	9.70
39 L-Phenylal- anine	C <sub>9</sub> H <sub>11</sub> O <sub>2</sub> N	165.19	283	2.96	sl. s. al.; i. eth.	H <sub>2</sub> O	1.9	20	-35.1	5.91 <sup>3</sup>
40 L-Proline	C <sub>5</sub> H <sub>9</sub> O <sub>2</sub> N	115.13	220-222	162.3	1.55, al.; i. eth.	0.5 N HCl	0.6	20	-52.6	6.3
41 Sarcosine	C <sub>3</sub> H <sub>7</sub> O <sub>2</sub> N	89.1	210	v. s.	sl. s. al.; i. eth.	.....	.....	...	0	6.12
42 L-Serine	C <sub>3</sub> H <sub>7</sub> O <sub>3</sub> N	105.09	228	5.023 <sup>5</sup>	i. al., eth.	1.0 N HCl	9.3	25	+14.5	5.68 <sup>3</sup>
43 L-Thiolhisti- dine	C <sub>6</sub> H <sub>9</sub> O <sub>2</sub> N <sub>3</sub> S	187.2	310 <sup>7</sup>	s.	s. a.; i. al., or- ganic solvents	1.0 N HCl	1.0	25	-9.5	5.16
44 L-Threonine	C <sub>4</sub> H <sub>9</sub> O <sub>3</sub> N	119.12	229-230	20.1 <sup>3</sup>	i. al., eth.	H <sub>2</sub> O	1.0	26	-28.4	5.59
45 L-Thyroxine	C <sub>15</sub> H <sub>11</sub> O <sub>4</sub> N <sub>4</sub>	776.88	235-236	0.001	i. al., eth.	0.13 N NaOH in 70% al- cohol	3	...	-4.4	....
46 L-Tryptophan	C <sub>11</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub>	204.23	289	1.14	sl. s. al.; i. eth.	H <sub>2</sub> O	1.0	20	-31.5	5.88
47 L-Tyrosine	C <sub>9</sub> H <sub>11</sub> O <sub>3</sub> N	181.19	295	0.045	0.01, al.; s. alk.; i. eth., acet.	6.3 N HCl	4.4	20	-8.6	5.63
48 L-Valine	C <sub>5</sub> H <sub>11</sub> O <sub>2</sub> N	117.15	293	8.85	0.019, al. <sup>3</sup>	6.0 N HCl	3.4	20	+28.8	6.00 <sup>3</sup>

/1/ Most amino acids decompose when melting. /2/ Grams of amino acid soluble in 100 ml of solvent. /3/ Racemic mixture (DL). /5/ Dihydrochloride. /6/ At 50°C. /7/ Decomposes without melting.

Contributors: (a) Sauberlich, H. E., (b) Ward, Wilfred H., (c) Evans, Robert John

References: [1] Andrews, S., and C. L. A. Schmidt. 1927. J. Biol. Chem. 73:651. [2] Ashley, J. N., and C. B. Harington. 1930. J. Chem. Soc., p. 2586. [3] Bergel, F. 1948. Biochem. Soc. Symp. (Cambridge, Engl.) 1:78. [4] Block, R. J., R. LeStrance, and G. Zweig. 1952. Paper chromatography: a laboratory manual. Academic Press, New York. [5] California Foundation for Biochemical Research. 1955. Properties of the L- (natural) amino acids. Rev. ed. Los Angeles. [6] Cohn, E. J., and J. T. Edsall. 1943. Proteins, amino acids and peptides. Reinhold, New York. [7] Dunn, M. S. 1960-61. In C. D. Hodgman, ed. Handbook of chemistry and physics. Ed. 42. Chemical Rubber, Cleveland, p.1760. [8] Du Vigneaud, V., et al. 1942. J. Biol. Chem. 143:59. [9] Dyer, H. M. 1938. Ibid. 124:519. [10] Greenstein, J. P., and M. Winitz. 1961. Chemistry of the amino acids. J. Wiley, New York. v. 1-3. [11] Heilbron, I., and H. M. Bunbury. 1953. Dictionary of organic compounds. Eyre and Spottiswoode, London. [12] Howe, E. E. 1951. Amino acids and proteins. C. C. Thomas, Springfield, Ill. p. 3. [13] Riegel, B., and V. du Vigneaud. 1935. J. Biol. Chem. 112:149. [14] Schmidt, C. L. A. 1945. The chemistry of the amino acids and proteins. C. C. Thomas, Springfield, Ill. [15] West, E. S., and W. R. Todd. 1951. Textbook of biochemistry. Macmillan, New York. [16] Wichers, E. 1952. J. Am. Chem. Soc. 74:2447.



## 107. VITAMINS AND PROVITAMINS:

*Abbreviations* (columns H, I): a. = acid; abs. = absolute; ac. = acetic; acet. = acetone; al. = alcohol; alk. = alkali; uble; me. = methyl; pet. = petroleum; s. = soluble; sl. = slightly; v. = very; w. = water.

Vitamin or Provitamin	Synonyms	Systematic Name	Chemical Formula	Physical State	Melting or Boiling Point °C	
(A)	(B)	(C)	(D)	(E)	(F)	
Vitamins						
1	Vitamin A <sub>1</sub>	Retinol; antixerophthalmia factor	3,7-Dimethyl-9-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6,8-nonatetraen-1-ol	C <sub>20</sub> H <sub>30</sub> O	Pale yellow prisms	62-64; distills at 120-125 at 5 x 10 <sup>-3</sup> mm
2	Vitamin A <sub>2</sub>	Dehydroretinol		C <sub>20</sub> H <sub>28</sub> O	Yellow oil	
3	Vitamin A, neo-		2- <i>cis</i> -Vitamin A	C <sub>20</sub> H <sub>30</sub> O	Yellow needles	59-60
4	Vitamin A aldehyde	Retinene; retinal; axerophthal	All- <i>trans</i> form	C <sub>20</sub> H <sub>28</sub> O	Orange crystals	61-64
5	<i>p</i> -Aminobenzoic acid	PABA		C <sub>7</sub> H <sub>7</sub> NO <sub>2</sub>	Monoclinic prisms from dilute alcohol	187.0-187.5
6	Ascorbic acid	Vitamin C; anti-scorbutic factor	<i>L</i> - <i>threo</i> -2,3,4,5-6-Pentahydroxy-2-hexeno-γ-lactone	C <sub>6</sub> H <sub>8</sub> O <sub>6</sub>	Crystals, plates or needles; monoclinic, colorless	190-192 (some decomposition)
7	Biotin	Vitamin H; coenzyme R; factor S, W, X; bios II G; antiegg-white injury factor	<i>cis</i> -Hexahydro-2-oxo-1H-thieno [3,4]imidazole-4-valeric acid	C <sub>10</sub> H <sub>16</sub> N <sub>2</sub> O <sub>3</sub> S	Fine long needles or white crystalline powder	232-233 (some decomposition)
8	Choline		(β-Hydroxyethyl)-trimethylammonium hydroxide	C <sub>5</sub> H <sub>15</sub> NO <sub>2</sub>	Colorless, viscous, hygroscopic alkaline liquid	
9	Cobalamin	Cyanocobalamin (vitamin B <sub>12</sub> ); hydroxycobalamin (vitamin B <sub>12a'</sub> , B <sub>12b</sub> )	5,6-Dimethylbenzimidazolyl cyanocobamide	C <sub>63</sub> H <sub>90</sub> CoN <sub>14</sub> O <sub>14</sub> P	Hygroscopic, dark red needles; birefringent	Darkens at 210-220; not melted at 300
10	Vitamin D <sub>2</sub>	Calciferol; ergocalciferol; activated ergosterol; anti-rachitic factor; oleovitamin D <sub>2</sub> ; viosterol	9,10-Secoergosta-5,7,10(19),22-tetraen-3-ol	C <sub>28</sub> H <sub>44</sub> O	Prisms from acetone	115-118; sublimes in very high vacuum without decomposition
11	Vitamin D <sub>3</sub>	Activated 7-dehydrocholesterol; cholecalciferol; oleovitamin D <sub>3</sub>	22,23-Dihydro-24-demethylcalciferol	C <sub>27</sub> H <sub>44</sub> O	Fine needles from dilute acetone	84-85
12	Vitamin E	α-Tocopherol, anti-sterility factor	5,7,8-Trimethyltolcol	C <sub>29</sub> H <sub>50</sub> O <sub>2</sub>	Slightly viscous, pale yellow oil	Boils at 200-220 at 0.1 mm
13		β-Tocopherol <sup>b</sup>	5,8-Dimethyltolcol	C <sub>28</sub> H <sub>48</sub> O <sub>2</sub>	Yellow oil	
14		γ-Tocopherol <sup>b</sup>	7,8-Dimethyltolcol	C <sub>28</sub> H <sub>48</sub> O <sub>2</sub>	Crystals	
15		σ-Tocopherol <sup>b</sup>	8-Methyltolcol	C <sub>27</sub> H <sub>46</sub> O <sub>2</sub>	Yellow oil	
16	Inositol	<i>meso</i> -Inositol; <i>i</i> -inositol; bios I	Hexahydroxycyclohexane	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	Efflorescent crystals (dihydrate)	218, dihydrate; 225-227, anhydrous
17	Vitamin K <sub>1</sub>	Antihemorrhagic vitamin	2-Methyl-3-phytyl-1,4-naphthoquinone	C <sub>31</sub> H <sub>46</sub> O <sub>2</sub>	Yellow, viscous oil	-20; decomposes above 100-120
18	Vitamin K <sub>2</sub>		2-Methyl-3-difarneysyl-1,4-naphthoquinone	C <sub>41</sub> H <sub>56</sub> O <sub>2</sub>	Yellow crystals	53.5-54.5
19	Nicotinic acid	Niacin; P.P. factor	Pyridine-3-carboxylic acid	C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub>	Colorless needles	236.5

<sup>1</sup>/<sub>1</sub> Measurement based on sodium light of wavelength 589 mμ, unless otherwise indicated. <sup>1</sup>/<sub>1</sub> E<sub>1 cm</sub><sup>1%</sup> = extinc-α-tocopherol.

## PHYSICAL AND CHEMICAL CHARACTERISTICS

bz. = benzene; chl. = chloroform; dil. = dilute; eth. = ether; glac. = glacial; gly. = glycerol; hex. = hexane; i. = insol-

Stability (G)	Solubility g/100 ml (H)	Specific Rotation <sup>1</sup>			Absorption Maxima		
		Solvent (I)	Temp. °C (J)	[α] <sub>D</sub> (K)	Wavelength, mμ, (E <sub>1</sub> <sup>1%</sup> 1 cm) <sup>2</sup> (L)	Solvent (M)	
<b>Vitamins</b>							
Inactivated by ultraviolet; sensitive to air oxidation.	s. most organic solvents, fats, oils; i. w.	Inactive			325 (1,835)	Isopropanol	1
					288 (773), 352 (1,450)	Ethanol	2
					328 (1,686)	Ethanol	3
	s. most organic solvents, fats, oils; i. w.				373	Cyclohexane	4
Incompatible with ferric salts and oxidizing agents.	v.s. al., eth., glac. ac. a.; 0.5, w. (25°C); sl.s. bz.	Inactive			265 (1,090)	0.1 N NaOH	5
					284 (1,020)	At pH 3.75	5
Stable to air when dry; impure preparations and natural products oxidized by air and light.	3,5, al.; 2, abs. al.; 33, w.; 1, gly.	me. al. w.	23 25	+48 +20.5-21.5	243 (560) 266 (855)	0.1 N H <sub>2</sub> PO <sub>3</sub> At pH 7	6
Stable to air, temperature; moderately acid and neutral solution stable several months; alkaline solution less stable.	0.080, al. (25°C); 0.022, w. (25°C); more s. hot w. or dil. alk.	0.1 N NaOH	21	+91			7
Dilute aqueous solution stable to boiling; decomposes in hot alkali.	v.s. al., w.; i. eth.						8
Heat-stable in aqueous solution; inactivated slowly by weak acid, alkali.	s. al.; 1.25, w.; i. acet., chl., eth.	dil. aqueous solution	23	-59±9 <sup>3</sup>	278 (115), 361 (204), 550 (64)	Water	9
Crystals stable 9 months in amber-evacuated ampules at 4°C; propylene glycol solution stable to air for long periods.	6,95, acet. (7°C); s. most organic solvents; sl.s. vegetable oils; i. w.	acet. al. chl. eth.	20 20 20 20	+82.6 +103 +52 +91.2	264 (459)	Hexane	10
At least as stable as vitamin D <sub>2</sub> .	s. most fat solvents; i. w.	acet.	20	+84.8	264 (450-490)	Hexane	11
Very stable to heat, acid; slowly oxidized by atmospheric O <sub>2</sub> , rapidly by ferric and silver salts.	v.s. al., acet., chl., eth., fats, oils; i. w.	al. bz.	25 25	+0.32 <sup>4</sup> -3.0 <sup>4</sup>	294 (71-76)	Ethanol	12
		al.	25	+2.9 <sup>4</sup>	295		13
		al.	25	+2.2 <sup>4</sup>	295		14
		al.	25	+3.4 <sup>4</sup>	298		15
Becomes anhydrous at 100°C; decomposes at 250°C.	14, w.; i. abs. al., eth.	Inactive					16
Stable to air, moisture; decomposes in sunlight; stable to dilute acid; labile to alkaline hydroxides.	s. acet., al., bz., chl., eth.; sl.s. me. al.; i. w.		20	-0.4	243 (410), 249 (425), 260 (395), 269 (395), 325 (75)	Hexane	17
					243 (320), 249 (329), 260 (305), 269 (305), 325 (58)	Hexane	18
Stable to air, light, pH; non-hygroscopic.	0.73, al. (25°C); 1.67, w. (25°C); i. eth.	Inactive			260.5 (432) 263 (260)	0.1 N HCl At pH 11	19

tion coefficients of 1% solutions of 1-cm thickness. /<sub>3</sub>/ D = 656 mμ. /<sub>4</sub>/ D = 546.1 mμ. /<sub>5</sub>/ Much less active than

*continued*

## 107. VITAMINS AND PROVITAMINS:

Vitamin or Provitamin	Synonyms	Systematic Name	Chemical Formula	Physical State	Melting or Boiling Point °C	
(A)	(B)	(C)	(D)	(E)	(F)	
Vitamins						
20	Nicotinamide	Niacinamide	Pyridine-3-carboxamide	$C_6H_6N_2O$	Colorless needles	129-131
21	Pantothenic acid	Chick antidermatitis factor; factor II	$\alpha$ -( <i>d</i> )- <i>N</i> -( $\alpha$ , $\gamma$ -Dihydroxy- $\beta$ , $\beta$ -dimethylbutyryl)- $\beta$ '-alanine	$C_9H_{17}NO_5$	Colorless, viscous oil	Unstable; calcium salt decomposes at 195-196
22	Pteroylglutamic acid	PGA; folic acid; folacin, vitamin M; <i>Lactobacillus casei</i> factor; vitamin B <sub>9</sub>	<i>N</i> -[4-[(2-Amino-4-hydroxy-6-pteridyl)methyl]-amino]-benzoyl]-glutamic acid	$C_{19}H_{19}N_7O_6$	Yellowish-orange platelets	Darkens and chars from approximately 250
23	Pyridoxine <sup>6</sup>	Vitamin B <sub>6</sub> -HCl; anti-acrodynia factor; adermine	5-Hydroxy-6-methyl-3,4-pyridine dimethanol hydrochloride	$C_8H_{11}NO_3 \cdot HCl$	Colorless platelets	Decomposes: 205-212; sublimes: free base 160
24	Riboflavin	Vitamin B <sub>2</sub> ; vitamin G; lactoflavin; ovoflavin; hepatoflavin	6,7-Dimethyl-9-( $\alpha$ -1'-ribityl)-isoalloxazine	$C_{17}H_{20}N_4O_6$	Yellow to orange-yellow polymorphic crystals	277-291 (some decomposition)
25	Thiamine	Vitamin B <sub>1</sub> ; aneurine; antineuritic factor	3-(4-Amino-2-methylpyrimidyl-5-methyl)-4-methyl-5- $\beta$ -hydroxyethylthiazolium chloride hydrochloride <sup>7</sup>	$C_{12}H_{17}N_4OSCl \cdot HCl$	Monoclinic plates in rosette clusters, or white powder	246-250 decomposes
Provitamins						
26	$\beta$ -Carotene <sup>9</sup>	Provitamin A		$C_{40}H_{56}$	Red crystals	180 corrected
27	Ergosterol	Provitamin D <sub>2</sub>	$\Delta^5, 7, 22$ -Ergostatrien-3 $\beta$ -ol	$C_{28}H_{44}O$	Small white plates from alcohol	168
28	7-Dehydrocholesterol	Provitamin D <sub>3</sub>	$\Delta^5, 7$ -Cholestadien-3 $\beta$ -ol	$C_{27}H_{44}O$	Crystals	150
29	22,23-Dihydroergosterol	Provitamin D <sub>4</sub>		$C_{28}H_{46}O$	Solvated needles from ethyl acetate and methyl alcohol	152-153
30	7-Dehydrositosterol	Provitamin D <sub>5</sub>		$C_{29}H_{48}O$	Platelets from alcohol	144-145
31	7-Dehydrostigmasterol	Provitamin D		$C_{29}H_{46}O$	Crystals	154
32	<i>epi</i> -7-Dehydrocholesterol	Provitamin D <sub>1</sub>		$C_{27}H_{44}O$	Crystals	124-126
33	Kitol	Dimer of vitamin A		$C_{40}H_{60}O_2$	Prisms from alcohol	88-90
34	Lumisterol	Provitamin D <sup>10</sup>		$C_{28}H_{44}O$	Needles from acetone methanol	118
35	Pantotheryl alcohol	Provitamin of pantothenic acid; pantotherylol; <i>N</i> -pantoyl-3-propanolamine	2,4-Dihydroxy- <i>N</i> -(3-hydroxypropyl)-3,3-dimethylbutyramide	$C_9H_{19}NO_4$	Colorless, viscous oil	Racemizes at boiling point
36	Tachysterol	Provitamin <sup>10</sup>		$C_{28}H_{44}O$		

<sup>1/1</sup> Measurement based on sodium light of wavelength 589 m $\mu$ , unless otherwise indicated. <sup>2/2</sup>  $E_{1\%}^{1\text{cm}}$  = extinc-tion, with different properties and biological activity. <sup>3/3</sup> The mononitrate is a less hygroscopic form. <sup>4/4</sup> Soret ing vitamin A activity:  $\alpha$ -carotene,  $\gamma$ -carotene, neo- $\beta$ -carotene, and cryptoxanthin. <sup>5/5</sup> Intermediates between

## PHYSICAL AND CHEMICAL CHARACTERISTICS

Stability	Solubility g/100 ml	Specific Rotation <sup>1</sup>			Absorption Maxima		
		Solvent	Temp. °C	[α] <sub>D</sub>	Wavelength, mμ, (E <sub>1</sub> <sup>1%</sup> <sub>1 cm</sub> ) <sup>2</sup>	Solvent	
(G)	(H)	(I)	(J)	(K)	(L)	(M)	
<b>Vitamins</b>							
Stable to air, light, pH; non-hygroscopic.	66.6, al.; 100, w.; sl.s. eth.	Inactive			260.5 (435)	0.1 N HCl	20
					262 (250)	At pH 11	
Very hygroscopic; labile to acid, alkali, heat; calcium salt stable to air, light.	v.s. glac. ac. a., w.; sl.s. eth.; i. bz., chl.	Ca salt	25	+37.5			21
			25	+28.2			
Very labile to heat in acid media; sunlight causes deterioration.	s. ac. a.; sl.s. me. al., w.; i. acet., bz., chl., eth.				256 (603), 282 (600), 365 (215)	At pH 11	22
Fairly stable to light, air. Acid solution stable; may be heated 120° for 30 minutes.	22, w.; l.l. al., eth.; sl.s. acet.; i. eth.	Inactive			245 (306)	0.1 N NaOH	23
					254 (179)	Phosphate buffer, pH 7	
					291 (420)	0.1 N HCl	
					310 (333)	0.1 N NaOH	
					325 (337)	Phosphate buffer, pH 7	
When dry, stable to diffused light; very labile to alkali solution, especially in light; stable to mineral acids in dark.	0.045, al. (27.5°C); 0.019, w. (40°C); i. acet., bz., chl., eth.	dil. alcoholic NaOH (50 mg in 2 ml)		-112 to -122	223 (800), 266 (870), 375 (288), 444 (310)	0.1 N HCl	24
Stable in acid solution and increasingly unstable as pH increases.	1, al.; 0.3, abs. al.; 5, gly.; 100, w.; i. bz., chl., eth., hex.				246 (410), 263S <sup>3</sup> (350)	0.1 N HCl	25
<b>Provitamins</b>							
Sensitive to O <sub>2</sub> , autooxidation in light; stable to heat.	s. bz., CS <sub>2</sub> , pet. eth.; sl.s. al., eth.; i. w.				450, 485, 520	CS <sub>2</sub>	26
Destroyed by ultraviolet; decomposes oxidizing agents.	v.s. most fat solvents; i. w.	chl.		-130	280	Ether	27
	v.s. most fat solvents; i. w.	chl.		-113	280	Ether	28
	v.s. most fat solvents; i. w.	chl.		-109			29
Browns on contact with air.	v.s. most fat solvents; i. w.	chl.		-116	262, 271, 282, 293	Alcohol	30
	v.s. most fat solvents; i. w.	bz.		-113.1			31
	v.s. most fat solvents; i. w.	chl.		-70.5			32
	v.s. most fat solvents; i. w.	chl.		-1.35	286		33
	v.s. most fat solvents; i. w.	acet.		+192	265, 280		34
Labile to acid, alkali.	v.s. al., w.; sl.s. eth.	3% aqueous	20	+29.7			35
	v.s. most fat solvents; i. w.	bz.		-70	280		36

tion coefficients of 1% solutions of 1-cm thickness. /<sup>a</sup>/ The vitamin B<sub>6</sub> group also includes pyridoxal and pyridox-effect, i.e., the concentration varies with the temperature gradient within the solution. /<sup>s</sup>/ Other carotenoids have provitamin and vitamin.

*continued*



**107. VITAMINS AND PROVITAMINS: PHYSICAL AND CHEMICAL CHARACTERISTICS**

*Contributors:* (a) Bird, Orson D., and Vandenberg, J. M., (b) Oser, Bernard L., (c) DeRitter, E. and Rubin, Saul H.

*References:* [1] Deuel, H. J., Jr. 1951-57. The lipids; their chemistry and biochemistry. Interscience, New York. [2] Harris, R. S., and D. J. Ingle, ed. 1960. Vitamins Hormones, v. 18. [3] Sebrell, W. H., Jr., and R. S. Harris. 1954. The vitamins. Academic Press, New York. v. 1-3. [4] Stecher, P. G., et al., ed. 1960. The Merck index. Ed. 7. Merck, Rahway, N. J.

**108. VARIOUS CELLS AND CELL PARTS: CHEMICAL COMPOSITION**

For information on additional species and tissues, consult reference 7. **Chemical Constituent** (column B): DNA = deoxyribonucleic acid; RNA = ribonucleic acid; N = nitrogen; P = phosphorus.

Cell or Cell Part	Chemical Constituent	Value	Reference	Cell or Cell Part	Chemical Constituent	Value	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
<b>Man</b>				<b>Pancreas</b>					
1	Bone marrow, whole cell	DNA-P, $\mu\text{g}/\text{cell}$	0.87	12	30	Whole cell	DNA-P, mg/g fresh tissue	0.22	11
2		RNA-P, $\mu\text{g}/\text{cell}$	0.69		31	cell	RNA-P, mg/g fresh tissue	1.77	
3	Brain, whole cell	DNA-P, $\mu\text{g}/\text{cell}$	0.68	5	32	Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	6.6	6,40
4		RNA-P, $\mu\text{g}/\text{cell}$	2.63		<b>Sperm</b>				
5	Leukocyte, whole cell	DNA-P, $\mu\text{g}/\text{cell}$	0.73	12	33	Whole cell	DNA, $\mu\text{g}/\text{cell}$	2.82-3.40	6,31
6		RNA-P, $\mu\text{g}/\text{cell}$	0.25		34	Head	Total nucleic acid, % dry wt	48.0	10
7	Kidney, whole cell	DNA-P, $\mu\text{g}/\text{cell}$	0.83	13	35		DNA, $\mu\text{g}/\text{head}$	3.3	6,40
8		RNA-P, $\mu\text{g}/\text{cell}$	1.10		36		Basic protein, % dry wt	28.7	10
9	Liver Whole cell	DNA-P, $\mu\text{g}/\text{cell}$	1.0	13	37		Acidic protein, "lipo-," % dry wt	19.6	10
10		RNA-P, $\mu\text{g}/\text{cell}$	2.48		38	Spleen, nucleus	Total nucleic acid, % dry wt	32.6-33.6	29
11		Total N, $\mu\text{g}/\text{cell}$	75.3		39		DNA, $\mu\text{g}/\text{nucleus}$	6.8	6,40
12	Nucleus	Nucleoprotein, %	42-59	15	40		RNA, % dry wt	0.7-1.1	29
13		Acidic protein, %	35-51		<b>Thymus</b>				
14		Other protein, "residual," %	4.7-7.5		41	Whole cell (calf)	DNA-P, mg/g fresh tissue	2.24-2.50	11
15	Sperm	DNA-P, $\mu\text{g}/\text{sperm}$	0.31	13	42	cell	RNA-P, mg/g fresh tissue	0.80-1.00	
16		RNA-P, $\mu\text{g}/\text{sperm}$	0.24		43	Nucleus	Total nucleic acid, % total N	31.0	30
17	Spleen, whole cell	DNA-P, mg/g fresh tissue	0.77	11	44		Basic protein, % total N	35.0	30
18		RNA-P, mg/g fresh tissue	0.36		45		Acidic protein, % total N	14.0	30
<b>Cattle</b>				46			DNA, $\mu\text{g}/\text{nucleus}$	6.4	6,40
19	Liver Whole cell	DNA-P, mg/g fresh tissue	0.34	11	<b>Dog</b>				
20		RNA-P, mg/g fresh tissue	0.70		47	Liver Whole cell	Total lipid, % dry wt	17.2	44
21	Nucleus	Total nucleic acid, % dry wt	27.5-30.7	29	48		Phospholipid, % dry wt	9.2	
22		DNA, $\mu\text{g}/\text{nucleus}$	6.4	6,40	49		Cholesterol, % dry wt	1.07	
23		RNA, % dry wt	0.9-1.9	29	50		Neutral fat, % dry wt	6.9	
24	Ribosome (calf)	RNA, %	40	32	51	Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	5.3	42
25	Heart, nucleus	DNA, % dry wt	30.0	4	52		Total lipid, % dry wt	16.5	44
26		Total lipid, % dry wt	26.0	38	53		Phospholipid, % dry wt	10.7	44
27		Phospholipid, % dry wt	15.7	38	54		Cholesterol, % dry wt	1.2	44
28		Cholesterol, % dry wt	3.6	38	55		Fatty acid, % dry wt	4.6	44
29		Fatty acid, % dry wt	6.5	38	56	Sperm, head	Total nucleic acid, % dry wt	55.3	10
<b>Guinea Pig</b>				57			Basic protein, % dry wt	25.0	
59	Liver Whole cell	DNA-P, mg/g fresh tissue	0.42	34	58		Acidic protein, "lipo-," % dry wt	17.0	
60		RNA-P, mg/g fresh tissue	0.97	34	<b>Guinea Pig</b>				
61		Total protein, % dry wt	15.0	26	59	Liver Whole cell	DNA-P, mg/g fresh tissue	0.42	34
					60	cell	RNA-P, mg/g fresh tissue	0.97	34
					61		Total protein, % dry wt	15.0	26

*continued*

**108. VARIOUS CELLS AND CELL PARTS: CHEMICAL COMPOSITION**

Cell or Cell Part	Chemical Constituent	Value	Reference	Cell or Cell Part	Chemical Constituent	Value	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
Guinea Pig				Heart					
62	Liver Mitochondria	Total N, % dry wt of fraction <sup>1</sup>	10.0-12.0	8	Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	6.46	39	
		Total lipid, % dry wt of fraction <sup>1</sup>	25.0		Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.267	39
		Phospholipid, % dry wt of fraction <sup>1</sup>	16.0			cell	RNA-P, mg/g fresh tissue	0.657	
					Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	6.72	39	
63	Microsome	Total N, % dry wt of fraction <sup>2</sup>	9.15	8	Liver				
		Total lipid, % dry wt of fraction <sup>2</sup>	40.0-51.0	8,27	Whole cell	DNA, mg/g fresh tissue	1.92	33	
		Phospholipid, % dry wt of fraction <sup>2</sup>	28.0-29.0	8,27		RNA, mg/g fresh tissue	5.88	33	
						DNA-P, mg/g fresh tissue	0.21-0.25	11	
			RNA-P, mg/g fresh tissue	0.77-1.10		11			
64	Microsome	Phospholipid, % total lipid	58	27	Total protein, mg/g fresh tissue	129.0	33		
					Total lipid, % dry wt	15.2	44		
					Phospholipid, % dry wt	8.3	44		
					Cholesterol, % dry wt	2.4	44		
Mouse				Nucleus					
69	Liver Whole cell	DNA, mg/g fresh tissue	2.85	3	Total nucleic acid, % dry wt	11.4-27.5	20,29		
		RNA, mg/g fresh tissue	9.0	3	DNA, mg/g fresh tissue	1.84	33		
		DNA-P, mg/g fresh tissue	0.232	25	DNA, % dry wt	4.4-30.0	20,43		
		RNA-P, mg/g fresh tissue	0.927	25	RNA, mg/g fresh tissue	0.64	33		
70	Microsome	Total protein, mg/g fresh tissue	126.3	3	RNA, % dry wt	2.9-7.6	20,29		
		Phospholipid, mg/g fresh tissue	30.1	3	Nucleoprotein, mg/g fresh tissue	20.0	33		
					Total lipid, % dry wt	10.5-18.1	17,43, 44		
					Total lipid, %	3.2-10.0	18		
71	Nucleus	DNA, %	27.0	3	Mitochondria	DNA, % total nucleic acid	11.7 <sup>5</sup>	23	
		RNA, %	3.4		RNA, % total nucleic acid	19.0-46.0	22,23		
		Nucleoprotein, %	66.0		RNA-P, $\mu\text{g}/\text{mg N}$	11.0	37		
		Phospholipid, %	3.4		Total N, %	23.0-38.6	22,23, 37		
72	Mitochondria	DNA, % total nucleic acid	5.6 <sup>3</sup>	36	Total protein, %	30.0-33.0	37		
		RNA, % total nucleic acid	16.8	36	Total protein, mg/g fresh tissue	35.0-40.0	33		
		RNA, % dry wt of fraction <sup>1</sup>	3.7	2	Total lipid, % dry wt of fraction <sup>1</sup>	25.0-30.0	37		
		Total N, %	23.5	35	Phospholipid, % total lipid	66.0	37		
73	Mitochondria	Total N, % dry wt of fraction <sup>1</sup>	12.1	2	Microsome	RNA, % total nucleic acid	50.0	37	
		Total lipid, % dry wt of fraction <sup>1</sup>	27.4	2	Total N, %	18.0-20.0	37		
		Phospholipid, % total lipid	56.6	2	Total protein, mg/g fresh tissue	19.0-21.0	33		
		Cholesterol <sup>4</sup> , % total lipid	12.6	2	Total lipid, % dry wt of fraction <sup>2</sup>	40.0	37		
74	Microsome	Neutral fat, % total lipid	30.8	2	Lung				
		DNA, % total nucleic acid	14.2 <sup>5</sup>	36	Whole cell	DNA-P, mg/g fresh tissue	0.921	39	
		RNA, % total nucleic acid	52.4	36		cell	RNA-P, mg/g fresh tissue	0.520	
		RNA, % dry wt of fraction <sup>2</sup>	9.1	2	Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	6.71	39	
75	Microsome	Total N, %	23.1	35	Spleen				
		Total N, % dry wt of fraction <sup>2</sup>	10.3	2	Whole cell	DNA-P, mg/g fresh tissue	1.40	39	
		Total lipid, % dry wt of fraction <sup>2</sup>	35.1	2,26		cell	RNA-P, mg/g fresh tissue	0.499	
		Phospholipid, % total lipid	62.7	2,26	Nucleus	DNA, $\mu\text{g}/\text{nucleus}$	6.52	39	
76	Microsome	Cholesterol <sup>4</sup> , % total lipid	14.5	26	Rabbit				
		Neutral fat, % total lipid	22.8	26	Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
77	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
78	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
79	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
80	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
81	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
82	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
83	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
84	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
85	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
86	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
87	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
88	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
89	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
90	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
91	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
92	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
93	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
94	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
95	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
96	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
97	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
98	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
99	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	
100	Microsome				Rabbit				
					Kidney	Whole cell	DNA-P, mg/g fresh tissue	0.125	28
							cell	RNA-P, mg/g fresh tissue	0.167
					Nucleus	Total nucleic acid, % dry wt	26.0	29	
101	Microsome				RNA, % dry wt	1.2			
					Liver				
					Whole cell	DNA-P, mg/g fresh tissue	0.16-0.29	11	
						cell	RNA-P, mg/g fresh tissue	0.44-0.76	

/1/ Large granule fraction obtained by differential centrifugation of liver cytoplasm extract. /2/ Small granule fraction obtained by differential centrifugation of liver cytoplasm extract. /3/ Contamination with nuclear material cannot be excluded. /4/ Unsaponifiable.

*continued*

## 108. VARIOUS CELLS AND CELL PARTS: CHEMICAL COMPOSITION

Cell or Cell Part	Chemical Constituent	Value	Reference	Cell or Cell Part	Chemical Constituent	Value	Reference		
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
Rabbit				Fish					
147	Liver Nucleus	Total nucleic acid, % dry wt	26.2	29	172	Cod sperm, head	Total nucleic acid, % dry wt	30.3	29
148		RNA, % dry wt	2.0		173		RNA, % dry wt	0.3	
149	Mitochondria	Nucleic acid P, $\mu\text{g}/\text{mg}$ total N	70.0	1	174	Herring sperm, head	Total nucleic acid, % dry wt	38.8-59.0	29
150		Total N, % dry wt of fraction <sup>1</sup>	10.5		175		RNA, % dry wt	0-0.2	
151		Total lipid, % dry wt of fraction <sup>1</sup>	29.6		176	Salmon sperm, head	Total nucleic acid, % dry wt	60.8	29
152		Phospholipid, % dry wt of fraction <sup>1</sup>	17.5		177		RNA, % dry wt	0.1	
153	Microsome	Nucleic acid P, $\mu\text{g}/\text{mg}$ total N	80.0	1	Sea Urchin <sup>b</sup>				
154		Total N, % dry wt of fraction <sup>2</sup>	9.0		178	Ovum	DNA, $\mu\text{g}/\text{cell}$	28.0	21
155		Total lipid, % dry wt of fraction <sup>2</sup>	43.4		179		DNA, % dry wt	0.01	
156		Phospholipid, % dry wt of fraction <sup>2</sup>	31.2		180	Sperm	DNA, $\mu\text{g}/\text{cell}$	1.0	21
157	Reticulocyte, ribosome	RNA, %	50	16	181		DNA, % dry wt	15.0	
Fowl				Bacteria					
158	Erythrocytes, nucleus	Total nucleic acid, % dry wt	33.9-38.1	29	182	<i>Bacillus anthracis</i>	Total nucleic acid, % dry wt	4.35	41
159		DNA, $\mu\text{g}/\text{nucleus}$	2.34-2.49	14,31	183		DNA, % dry wt	1.15	
160		DNA, %	45.0	19	184		RNA, % dry wt	3.20	
161		RNA, % dry wt	0.7-2.5	29	185		Total N, % dry wt	10.0	
162		Nucleoprotein, %	50.0-60.0	24	186		Total protein, % dry wt	58.1	
163		Acidic protein, %	33-40	24	187	<i>Escherichia coli</i>	Total nucleic acid, % dry wt	12.84	41
164	Liver Whole cell	DNA-P, mg/g fresh tissue	0.31-0.41	9	188		DNA, % dry wt	3.72	
165		RNA-P, mg/g fresh tissue	0.76-0.84		189		RNA, % dry wt	9.12	
166	Nucleus	Total nucleic acid, % dry wt	29.4-31.2	29	190		Total N, % dry wt	14.61	
167		DNA, $\mu\text{g}/\text{nucleus}$	2.39-2.54	14,31	191		Total protein, % dry wt	78.5	
168		RNA, % dry wt	2.0-2.2	29	192	<i>Salmonella typhosa</i>	Total nucleic acid, % dry wt	13.12	41
169	Sperm, nucleus	DNA, $\mu\text{g}/\text{nucleus}$	1.26	31	193		DNA, % dry wt	4.40	
170	Thymus, nucleus	Total nucleic acid, % dry wt	32.0-36.3	29	194		RNA, % dry wt	8.72	
171		RNA, % dry wt	1.3-1.4		195		Total N, % dry wt	14.40	
					196		Total protein, % dry wt	76.8	
					197	<i>Staphylococcus</i> (strain 72)	Total nucleic acid, % dry wt	11.57	41
					198		DNA, % dry wt	2.82	
					199		RNA, % dry wt	8.75	
					200		Total N, % dry wt	13.95	
					201		Total protein, % dry wt	75.5	

<sup>1</sup>/ Large granule fraction obtained by differential centrifugation of liver cytoplasm extract. <sup>2</sup>/ Small granule fraction obtained by differential centrifugation of liver cytoplasm extract. <sup>b</sup>/ *Paracentrotus lividus*.

Contributors: (a) Kirkham, William R., (b) Allfrey, Vincent G.

References: [1] Ada, G. L. 1949. *Biochem. J.* 45:422. [2] Barnum, C. P., and R. A. Huseby. 1948. *Arch. Biochem.* 19:17. [3] Barnum, C. P., et al. 1950. *Ibid.* 25:376. [4] Behrens, M. 1932. *Z. Physiol. Chem.* 209:59. [5] Bieth, R., and P. Mandel. 1953. *Experientia* 9:185. [6] Boivin, A., R. Vendrely, and C. Vendrely. 1948. *Compt. Rend.* 226:1061. [7] Chargaff, E., and J. N. Davidson, ed. 1955. *The nucleic acids*, Academic Press, New York. v. 2. [8] Claude, A. 1946. *J. Exptl. Med.* 84:51. [9] Common, R. H., D. G. Chapman, and W. A. Maw. 1951. *Can. J. Zool.* 29:265. [10] Dallam, R. D., and L. E. Thomas. 1953. *Biochim. Biophys. Acta* 11:79. [11] Davidson, J. N. 1947. *Cold Spring Harbor Symp. Quant. Biol.* 12:50. [12] Davidson, J. N., I. Leslie, and J. C. White. 1951. *J. Pathol. Bacteriol.* 63:471. [13] Davidson, J. N., I. Leslie, and J. C. White. 1951.

continued



## 108. VARIOUS CELLS AND CELL PARTS: CHEMICAL COMPOSITION

Lancet 260:1287. [14] Davidson, J. N., et al. 1950. Biochem. J. 46:xl. [15] Debov, S. S., 1951. Chem. Abstr. 45:10374. [16] Dintzis, H. M., H. Borsook, and J. Vinograd. 1958. In R. B. Roberts, ed. Microsomal particles and protein synthesis. Pergamon Press, New York. p. 95. [17] Dounce, A. L. 1943. J. Biol. Chem. 147:685. [18] Dounce, A. L. 1943. Ibid. 151:221. [19] Dounce, A. L., and T. H. Lan. 1943. Science 97:584. [20] Dounce, A. L., et al. 1950. J. Gen. Physiol. 33:629. [21] Elson, D., and E. Chargaff. 1952. Experientia 8:143. [22] Hogeboom, G. H., W. C. Schneider, and G. E. Pallade. 1948. J. Biol. Chem. 172:619. [23] Hogeboom, G. H., W. C. Schneider, and M. J. Striebich. 1952. Ibid. 196:111. [24] Jeener, R. 1946. Compt. Rend. Soc. Belge Biol. 140:1103. [25] Johnson, R. M., and S. Albert. 1953. J. Biol. Chem. 200:335. [26] Kretchmer, N., and C. P. Barnum. 1951. Arch. Biochem. Biophys. 31:141. [27] Lazarow, A. 1946. Biol. Symp. 10:17. [28] Lowe, C. U., W. L. Williams, and L. Thomas. 1951. Proc. Soc. Exptl. Biol. Med. 78:818. [29] Mauritzen, C. M., A. B. Roy, and E. Stedman. 1952. Proc. Roy. Soc. (London), B, 140:18. [30] Mayer, D. T., and A. Gulick. 1943. J. Biol. Chem. 146:433. [31] Mirsky, A. E., and H. Ris. 1949. Nature 163:666. [32] Peterman, M. L., and M. G. Hamilton. 1957. J. Biol. Chem. 224:725. [33] Price, J. M., et al. 1950. Cancer Res. 10:18. [34] Rerabek, J. 1947. Arkiv Kemi Mineral. Geol. 24:1. [35] Schneider, W. C., and G. H. Hogeboom. 1950. J. Natl. Cancer Inst. 10:969. [36] Schneider, W. C., and G. H. Hogeboom. 1950. Ibid. 10:977. [37] Schneider, W. C., and G. H. Hogeboom. 1951. Cancer Res. 11:1. [38] Stoneburg, C. A. 1939. J. Biol. Chem. 129:189. [39] Thomson, R. Y., et al. 1953. Biochem. J. 53:460. [40] Vendrely, C. 1952. Bull. Biol. France Belg. 86:1. [41] Vendrely, R., and Y. Lehault. 1946. Compt. Rend. 222:1357. [42] Vendrely, R., and C. Vendrely. 1949. Experientia 5:327. [43] Villela, G. G., and F. Ubatuba. 1948. Rev. Brasil. Biol. 8:35. [44] Williams, H. H., et al. 1945. J. Biol. Chem. 160:227.

## 109. ANIMAL TISSUES AND ORGANS: WATER CONTENT

Values are for adult animals, unless otherwise indicated. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Tissue or Organ	% water	Refer- ence	Tissue or Organ	% Water	Refer- ence		
(A)	(B)	(C)	(A)	(B)	(C)		
Man							
Nervous tissue			Cardiac tissue				
1	Whole brain	77(76-78)	39	21	Right atrium	81.2	11
2	White matter	70(68-73)	39	22	Septum	79.2	11
3	Gray matter	84(82-85)	39	23	Kidney	78.4(77.7-79.0)	50
4	Spinal cord	71(63-75)	39	Reproductive tissue			
5	Peripheral nerve	66(62-68)	39	24	Testis	84.0	50
6	Eye lens	67.6	57	25	Prostate gland	82.5	50
Dental tissue			26	Ovary <sup>2</sup>	80.5	44	
7	Whole teeth	9.2(4.0-14.3)	13	27	Uterus	79.9	50
8	Enamel <sup>1</sup>	2.8	6	28	Muscle	76.0	10
9	Dentin <sup>1</sup>	11.1	6	29	Bone	43.9	60
Alimentary tract			30	Skin <sup>2</sup>	71.8(67.8-75.8) <sup>b</sup>	23	
10	Cardiac stomach	73.4	50	31	Hair	4.1(4.0-4.2)	4
11	Pyloric stomach	68.6	50	Cat			
12	Small intestine	71.0(60.2-81.8)	50	Nervous tissue			
13	Large intestine	72.7	50	32	Whole brain	72.2	41
14	Liver	75.0(72.9-77.3)	50	33	White matter	69	30
15	Pancreas	74.8	46	34	Gray matter	82	30
16	Spleen	78.7(76.5-81.1)	50	35	Spinal cord	67.7	42
17	Lung	81.3(79.5-82.7)	50	36	Sciatic nerve	(66.2-68.9)	35
Cardiac tissue			37	Eye lens <sup>3</sup>	74.5	7	
18	Whole heart	77.6(71.2-80.3)	50	38	Liver	70.7(68.7-72.7) <sup>b</sup>	65
19	Left ventricle	79.2	11	Cardiac tissue			
20	Right ventricle	80.7	11	39	Whole heart	78.7	56
				40	Left ventricle	77.7	61

<sup>1</sup>/ Deciduous teeth. <sup>2</sup>/ Fat-free basis. <sup>3</sup>/ Young animal.

continued



**109. ANIMAL TISSUES AND ORGANS: WATER CONTENT**

Tissue or Organ	% Water	Reference	Tissue or Organ	% Water	Reference
(A)	(B)	(C)	(A)	(B)	(C)
<b>Cat</b>			<b>Horse</b>		
41	Cardiac tissue		90	Nervous tissue	
	Right ventricle	61		Whole brain	25
42	Muscle	65	91	Peripheral nerve	2
<b>Cattle</b>			92	Eye lens	7
43	Nervous tissue		93	Heart	5
	Whole brain	51	<b>Rabbit</b>		
44	White matter	16	94	Nervous tissue	
45	Gray matter	16		Whole brain	27,41
46	Spinal cord	39	95	White matter	39
47	Obturator nerve	16	96	Gray matter	3,39
	Eye		97	Spinal cord	42,44
48	Cornea	17	98	Peripheral nerve	39
49	Lens	7	<b>Alimentary tract</b>		
50	Retina	12	99	Stomach	24,44
51	Liver	20	100	Small intestine	44
52	Lung	51	101	Duodenum	24
	Cardiac tissue		102	Cecum	24
53	Whole heart	51	103	Ileum	24
54	Left ventricle	16	104	Colon	44
55	Right ventricle	16	105	Liver	24
56	Left atrium	16	106	Spleen	24
57	Right atrium	16	107	Lung	24,44
58	Tricuspid valve	16	<b>Cardiac tissue</b>		
59	Kidney	51	108	Whole heart	44
60	Testis	55	109	Left ventricle	19
61	Muscle	51	110	Right ventricle	19
<b>Dog</b>			111	Kidney	24,44
62	Nervous tissue		<b>Reproductive tissue</b>		
	Whole brain	22	112	Testis	24,44
63	White matter	28,39	113	Ovary	24,44
64	Gray matter	28,39	114	Muscle	24
65	Spinal cord	39	115	Bone	8
66	Peripheral nerve	39	116	Skin	24,44
67	Intestine	38	117	Hair	24
68	Liver	38	<b>Rat</b>		
69	Pancreas	38	118	Nervous tissue	
70	Spleen	38		Whole brain	44
71	Lung	62	119	White matter	14
	Cardiac tissue		120	Gray matter	53
72	Whole heart	63	<b>Alimentary tract</b>		
73	Left ventricle	59	121	Stomach	44
74	Right ventricle	59	122	Small intestine	44
75	Left atrium	59	123	Large intestine	44
76	Right atrium	59	124	Liver	44
77	Kidney <sup>2</sup>	21	125	Spleen	44
78	Testis	33	126	Lung	44
79	Muscle	32	127	Heart	37
	Bone		128	Kidney	44
80	Humerus	9	<b>Reproductive tissue</b>		
81	Femur	9	129	Testis <sup>2</sup>	44
82	Tibia	9	130	Ovary	29
83	Skin	18	131	Muscle	44
<b>Guinea Pig</b>			132	Bone	8
84	Nervous tissue		133	Skin	64
	Whole brain	41	<b>Sheep</b>		
85	Spinal cord	42	134	Nervous tissue	
86	Liver	45		White matter	15
87	Lung	52	135	Gray matter	15
88	Heart	51	136	Liver <sup>3</sup>	51
89	Muscle	51	137	Muscle	54

<sup>2</sup>/ Fat-free basis. <sup>3</sup>/ Young animal.

*continued*

**109. ANIMAL TISSUES AND ORGANS: WATER CONTENT**

Tissue or Organ			Tissue or Organ				
(A)	% Water (B)	Refer-ence (C)	(A)	% Water (B)	Refer-ence (C)		
Sheep			Pigeon				
138	Bone	17.0(12.3-22.0)	66	Brain	(78.4-80.0)	34	
139	Wool	17.0(9.0-28.0)	47	149	Muscle	74.0	51
Swine			Frog				
140	Brain	77.0	25	Nervous tissue			
141	Tooth pulp	89.8	49	150	Whole brain	84	1
142	Liver	78.0	58	151	Peripheral nerve	85	26
143	Spleen	81.5	58	152	Muscle	(78.9-81.6)	48
144	Muscle	65.3	31	Carp			
145	Hair	11.4	43	153	Brain	74	1
Chicken			154	Eye lens	52	7	
146	Brain	78	36	Eel			
147	Feathers	9.1	43	155	Liver	77.9	40
				156	Muscle	57.1(53.8-59.1)	40
				157	Skin	66.5	40

*Contributors:* (a) Love, R. M., (b) McKibbin, John M., (c) Clarke, Norman E., (d) Logan, J. E., (e) Himwich, Williamina A.

*References:* [1] Abderhalden, E., and A. Weil. 1913. Z. Physiol. Chem. 83:425. [2] Alcock, N. H., and A. R. Lynch. 1907. J. Physiol. (London) 36:93. [3] Apreson, M. H., A. Lukenhill, and W. E. Segar. 1960. J. Neurochem. 5:150. [4] Bagchi, K. N., and H. D. Ganguly. 1941. Ann. Biochem. Exptl. Med. (Calcutta) 1:83. [5] Bertrand, G., and R. Vladesco. 1920. Compt. Rend. 171:744. [6] Bird, M. J., et al. 1940. J. Dental Res. 19:413. [7] Brückner, R. 1941. Ophthalmologica 100:203. [8] Burns, C. M. 1929. Biochem. J. 23:860. [9] Burns, C. M., and N. Henderson. 1946. Ibid. 40:501. [10] Chou, T. P., and W. H. Adolph. 1935. Ibid. 29:476. [11] Clarke, N. E., and R. E. Mosher. 1952. Circulation 5:907. [12] Collins, F. D., R. M. Love, and R. A. Morton. 1952. Biochem. J. 51:670. [13] Crowell, C. D., H. C. Hodge, and W. R. Line. 1934. J. Dental Res. 14:251. [14] Davenport, V. D. 1949. Am. J. Physiol. 156:322. [15] Davidson, J. N., and C. Waymouth. 1944. Biochem. J. 38:39. [16] Davies, F., et al. 1952. J. Physiol. (London) 118:276. [17] Davson, H. 1949. Brit. J. Ophthalmol. 33:175. [18] DeBoer, B. 1946. Am. J. Physiol. 147:49. [19] Decherd, G. M., Jr., G. Herrmann, and E. H. Schwab. 1936. Proc. Soc. Exptl. Biol. Med. 34:864. [20] Eggleton, W. G. E. 1939. Biochem. J. 33:403. [21] Eichelberger, L., and W. G. Bibler. 1940. J. Biol. Chem. 132:645. [22] Eichelberger, L., and R. B. Richter. 1944. Ibid. 154:21. [23] Eisele, C. W., and L. Eichelberger. 1945. Proc. Soc. Exptl. Biol. Med. 58:97. [24] Fore, H., and R. A. Morton. 1952. Biochem. J. 51:600. [25] Frankel, S., and K. Linnert. 1910. Biochem. Z. 26:44. [26] Gerard, R. W. 1932. Physiol. Rev. 12:469. [27] Graves, J., and H. E. Himwich. 1955. Am. J. Physiol. 180:205. [28] Gregersen, M. I., C. Pallavicini, and S. Chien. 1962. Radiation Res. 17:209. [29] Haldi, J., and G. Giddings. 1939. Am. J. Physiol. 128:537. [30] Halliburton, W. D. 1894. J. Physiol. (London) 15:90. [31] Hankins, O. G., A. J. Ernst, and W. R. Kauffman. 1946. Food Res. 11:501. [32] Hastings, A. B., and L. Eichelberger. 1937. J. Biol. Chem. 117:73. [33] Huggins, C., and L. Eichelberger. 1944. Cancer Res. 4:447. [34] Koch, M. L., and O. Riddle. 1919. J. Comp. Neurol. 31:83. [35] Krnjevic, K. 1955. J. Physiol. (London) 128:473. [36] Lajtha, A. 1956. J. Neurochem. 1:216. [37] Lemley, J. M., and G. R. Meneely. 1952. Am. J. Physiol. 169:61. [38] Lepore, M. J. 1932. Arch. Internal Med. 50:488. [39] Logan, J. E. 1961. In Blood and other body fluids. Federation of American Societies for Experimental Biology, Washington, D. C. p. 326. [40] McCance, R. A. 1944. Biochem. J. 38:474. [41] McColl, J. D., and R. J. Rossiter. 1952. J. Exptl. Biol. 29:196. [42] McColl, J. D., and R. J. Rossiter. 1952. Ibid. 29:203. [43] McComb, E. A. 1948. Anal. Chem. 20:1219. [44] Manery, J. F., and A. B. Hastings. 1939. J. Biol. Chem. 127:657. [45] Marble, A., A. L. Graflin, and R. M. Smith. 1940. Ibid. 134:253. [46] Marx, H. 1926. Biochem. Z. 179:414. [47] Miller, L. 1937.

*continued*

**109. ANIMAL TISSUES AND ORGANS: WATER CONTENT**

Przemysl Chem. 21:15. [48] Moran, T. 1929. Proc. Roy. Soc. (London), B, 105:177. [49] Motomura, S. 1941. Z. Physiol. Chem. 270:33. [50] Neufeld, A. H. 1937. Can. J. Res., B, 15:132. [51] Okey, R. 1945. J. Am. Dietet. Assoc. 21:341. [52] Okey, R. Unpublished. Univ. California, Berkeley, 1954. [53] Pappius, H. M., and K. A. C. Elliott. 1956. Can. J. Biochem. Physiol. 34:1007. [54] Peterson, W. H., J. T. Skinner, and F. M. Strong. 1943. Elements of food biochemistry. Prentice-Hall, New York. [55] Rewald, B. 1928. Biochem. Z. 202:99. [56] Robertson, W. V. B., and P. Peyser. 1951. Am. J. Physiol. 166:277. [57] Salit, P. W. 1943. Arch. Ophthalmol. (Chicago) 30:255. [58] Schultze, M. O., C. A. Elvehjem, and E. B. Hart. 1936. J. Biol. Chem. 116:93. [59] Sherrod, T. R. 1947. Proc. Soc. Exptl. Biol. Med. 65:89. [60] Shohl, A. T. 1939. Mineral metabolism. Reinhold, New York. [61] Wedd, A. M. 1939. J. Pharmacol. Exptl. Therap. 65:268. [62] Wood, E. H. 1942. J. Biol. Chem. 143:165. [63] Wood, E. H., and G. K. Moe. 1936. Am. J. Physiol. 136:515. [64] Wynn, W., and J. Haldi. 1944. Ibid. 142:508. [65] Yannet, H., and D. C. Darrow. 1938. J. Biol. Chem. 123:295. [66] Young, M. W. 1936. New Zealand J. Sci. Technol. 18:391.

**110. CELL SAP: CHEMICAL COMPOSITION**

Species (Common Name)	Plant Part	Growth Stage	Constituent	Value mg/100 g	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 <i>Avena sativa</i> (common oat)	Stem	Mature	NO <sub>3</sub> -N	41	3
2 <i>Beta saccharifera</i> (sugar beet)	Leaf	Mature	Mg	80-200	7
3		Mature	P	10.5-251.0	7
4		Mature	K	97-516	7
5		44-79 da	NO <sub>3</sub> -N	3-52	5
6	Stem	Mature	Mg	70-130	7
7			P	4.5-24.2	
8			K	34.0-41.4	
9 <i>Daucus carota</i> (carrot)	Root	2-3 mo	NO <sub>3</sub> -N	3.1-31.8	5
10			P	0.04-1.00	
11 <i>Fagopyrum esculentum</i> (buckwheat)	Leaf	.....	P	22-105	8
12	Stem	.....	P	24-145	8
13	Shoot	.....	NO <sub>3</sub> -N	12.6-17.2	5
14 <i>Glycine soja</i> (soybean)	Entire plant	Fruiting	NO <sub>3</sub> -N	2.4-3.6	11
15			P	2.6-9.0	
16			K	96-885	
17 <i>Hordeum vulgare</i> (barley)	Leaf	Mature	NO <sub>3</sub> -N	20	3
18			P	6.4-55.6	7
19			K	310-860	7
20	Stem	Mature	NO <sub>3</sub> -N	23	3
21			P	8.9-77.7	7
22			K	260-800	7
23	Entire plant	36 da	P	10	7
24			K	250	
25 <i>Lactuca sativa</i> (lettuce)	Leaf	Mature	NO <sub>3</sub> -N	6-45	5
26 <i>Lycopersicon esculentum</i> (tomato)	Petiole	Fruiting	NO <sub>3</sub> -N	6.9	6
27	.....	.....	P	30	
28	Stem	Immature	Ca	18	6
29 <i>Lycopersicon esculentum</i> (tomato)	Stem	Immature	NO <sub>3</sub> -N	5.8-11.5	6
30		Immature	P	12.5-20.0	
31		Immature	K	200-300	
32		.....	Mg	6-41	
33	Shoot	Immature	P	9.7-19.2	1
34 <i>Phaseolus vulgaris</i> (kidney bean)	Leaf	Mature	Ca	515-690	7
35			Mg	42-109	
36			P	3.5-17.7	
37			K	28-200	
38	Stem	Mature	Ca	137-283	7
39			Mg	76-105	
40 <i>Solanum tuberosum</i> (potato)	Lower stem	Mature	Ca	100	2
41		Mature	P	6	2
42		Mature	K	254-622	5
43		.....	NO <sub>3</sub> -N	16.1-18.4	1
44 <i>Triticum aestivum</i> (wheat)	Shoot	Immature	P	35-81	7
45		Mature	Ca	68-106	
46		Mature	NO <sub>3</sub> -N	1.4-5.8	
47		.....	Mg	27-107	
48	Entire plant	Immature	Ca	43	4
49			Mg	39	
50			NO <sub>3</sub> -N	2.8	
51			P	80	
52			K	411	
53 <i>Vigna sinensis</i> (cowpea)	Petiole	.....	Mg	19-42	1
54			NO <sub>3</sub> -N	2.2-22.0	
55			P	1.2-4.4	
56			K	250-415	
57 <i>Zea mays</i> (corn)	Stem	.....	Ca	4.7-11.6	10
58	Lower stem	Mature	NO <sub>3</sub> -N	8-41	4
59			P	2.7-3.8	5
60			K	166	9
61	Entire plant	Mature	NO <sub>3</sub> -N	0.5-25.0	4
62		Mature	P	1.6-12.2	9
63		.....	K	390-650	9

Contributors: (a) Giddens, Joel, (b) Samuels, George

References: [1] Carolus, R. L. 1936. Proc. Am. Soc. Hort. Sci. 33:579. [2] Carolus, R. L. 1937. Am. Potato J.

continued

### 110. CELL SAP: CHEMICAL COMPOSITION

14:141. [3] Cook, R. L. 1930. J. Am. Soc. Agron. 22:393. [4] Giddens, J. Unpublished. Univ. Georgia, Athens, 1955. [5] Gilbert, B. E., and L. J. Hardin. 1927. J. Agr. Res. 35:185. [6] Hester, J. B. 1941. Com. Fertilizer 63:10. [7] McCool, M. M., and M. D. Weldon. 1928. J. Am. Soc. Agron. 20:778. [8] Neller, J. R. 1935. J. Agr. Res. 51:287. [9] Pettinger, N. A. 1931. Ibid. 43:95. [10] Pierre, W. H., and G. G. Pohlman. 1933. J. Am. Soc. Agron. 25:144. [11] Poehlman, J. M. 1935. Ibid. 27:195.

### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

**Plant Part** (column B): veg. = plants in vegetative condition; fl. = plants flowering; fr. = plants fruiting; yg. = young; e. = early; im. = immature; un. = unripe.

#### Part I. MAJOR ELEMENTS

Values are g/100 g of dry weight.

Species (Common Name)	Plant Part	Potassium	Phosphorus	Calcium	Magnesium	Sulfur	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Monocotyledoneae								
1	<i>Allium cepa</i> (gar- den onion)	Shoots 1.03-1.92	0.187 0.17-0.74	2.59 0.260-0.863	0.262 0.11-0.16	0.177 0.120-0.600	11 7,12	
3	<i>Asparagus officinalis</i> (garden as- paragus)	Shoots	1.45-3.45	0.140-0.783	0.249-1.358	0.097-0.190	0.132-0.260	11,12
4		Shoots, yg.	2.665	0.529	0.329	0.048	0.176	12
5		Roots	2.914	0.105	1.043	0.078	0.172	12
6		Fruits	0.155-1.669	0.293-0.382	0.088-0.114	0.036-0.129	0.101-0.172	12
7	<i>Avena sativa</i> (common oat)	Shoots, veg.	2.780-2.880	0.244-0.365	0.380-0.534	0.150-0.218	0.136-0.170	10,11
8		Shoots, fl.	2.01-2.17	0.212-0.480	0.314-0.730	0.133-0.330	0.067-0.570	9-11
9		Shoots, fr.	0.78-2.20	0.16-0.40	0.21-0.51	0.13-0.41	0.07-0.28	7,72
10		Straw	0.59-3.52	0.02-0.36	0.15-0.67	0.06-0.54	0.08-0.51	7,12
11		Grain	0.280-1.086	0.150-0.955	0.019-0.190	0.060-0.356	0.020-0.294	7,12,35,62
12	<i>Elodea canadensis</i> (Canada water- weed)	Shoots	1.48-3.01	0.275-0.744	2.91-8.23	0.468-1.168	0.147-0.968	11
13	<i>Hordeum vulgare</i> (barley)	Shoots	3.88	0.347	0.677	0.250	0.192	11
14		Shoots, veg.	2.44	0.327	0.357	0.119	0.097	11
15		Shoots, fl.	1.37	0.291	0.267	0.118	0.076	11
16		Straw	1.08-1.96	0.04-0.56	0.15-0.60	0.040-0.287	0.018-0.230	7,12
17		Grain	0.270-0.923	0.150-0.620	0.011-0.150	0.018-0.273	0.019-0.366	7,12,35,62
18	<i>Oryza sativa</i> (rice)	Straw	0.930-1.640	0.066-0.130	0.193-0.300	0.060-0.110	0.100-0.138	1,12,49,75,82
19		Roots	0.755	0.079	0.307	0.121	.....	12
20		Grain	0.243-0.480	0.190-0.430	0.050-0.138	0.092-0.170	0.001-0.138	1,7,12,19,75
21	<i>Phleum pratense</i> (timothy)	Shoots	0.79-3.84	0.080-0.600	0.040-1.200	0.030-0.380	0.072-0.320	3,7,11
22		Shoots, fl.	0.92-2.32	0.17-0.41	0.15-0.44	0.09-0.17	0.02-0.27	7,8
23		Shoots, fr.	1.58	0.17	0.16	0.07	0.16	7
24		Grain	0.458	0.347	0.162	0.122	0.026	12
25	<i>Phoenix dactylife- ra</i> (date palm)	Pinnae	0.232-1.640	0.060-0.186	0.295-1.118	0.051-0.244	.....	37,70
26	<i>Poa pratensis</i> (Kentucky blue- grass)	Shoots	1.35-4.33	0.189-0.952	0.130-1.200	0.089-0.230	0.055-0.656	3,11,17,27,59, 62,78,79
27		Shoots, veg.	.....	0.193-0.611	0.302-0.871	.....	.....	8,16,69
28		Shoots, fl.	1.41-2.85	0.164-0.403	0.130-0.424	0.11	.....	7,27
29		Shoots, fr.	1.52-2.07	0.200-0.299	0.189-0.300	0.115	0.18	33,72
30		Grain	0.670	0.850	0.315	0.197	0.202	59
31	<i>Triticum aestivum</i> (wheat)	Shoots	2.88	0.249	0.368	0.174	0.120	11
32		Shoots, veg.	2.81	0.315	0.845	0.085	0.111	11
33		Shoots, fl.	1.61	0.233	0.194	0.112	0.052	11
34		Straw	0.26-1.54	0.03-0.17	0.08-0.43	0.03-0.22	0.07-0.30	7,28
35		Grain	0.237-0.971	0.150-0.540	0.005-0.296	0.090-0.290	0.003-0.290	7,12,28,35, 62, 80
36	<i>Zea mays</i> (corn)	Leaves	0.24-1.57	0.052-0.256	0.11-0.91	0.17-0.29	0.23-0.25	12,32,53,94
37		Stems	0.260-2.433	0.026-0.202	0.140-0.629	0.140-0.290	0.05-0.17	12,32,53,94
38		Shoots, fl.	1.669-1.790	0.136-0.550	0.592-0.681	0.230-0.392	0.075-0.370	11,12

continued



### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

#### Part I. MAJOR ELEMENTS

Species (Common Name)	Plant Part	Potassium	Phosphorus	Calcium	Magnesium	Sulfur	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Monocotyledoneae								
39	<i>Zea mays</i> (corn)	Shoots, fr.	0.26-1.89	0.04-0.42	0.10-0.84	0.08-0.51	0.08-0.31	7,11,28,95
40		Roots	0.270-1.277	0.030-0.141	0.129-0.720	0.094-0.180	0.033-0.280	12,53
41		Flowers, ♂	1.264	0.146	0.569	0.268	.....	12
42		Flowers, ♀	1.482	0.146	0.630	0.343	.....	12
43		Kernels	0.22-0.92	0.23-0.80	0.006-0.060	0.09-0.27	0.004-0.300	7,12,28,53,62
44		Cob	0.46	0.094	0.022	0.11	0.021	53
Dicotyledoneae								
45	<i>Acer saccharum</i> (sugar maple)	Leaves	0.95-1.58	0.24-0.46	0.57-2.42	0.24-0.35	0.01-0.24	61
46	<i>Alnus glutinosa</i>	Pollen	1.708	0.532	0.264	.....	.....	30
47	(European alder)	Fruits	0.412	0.105	0.358	0.121	0.027	12
48	<i>Beta vulgaris</i>	Leaves	1.68-6.45	0.089-0.436	0.78-3.12	0.17-1.74	0.345-0.845	7,11,12,46,62
49	(common beet)	Shoots	1.01-7.64	0.08-0.38	0.39-2.83	0.26-1.07	0.31-0.61	7
50		Crowns	.....	0.091-0.169	0.81-1.50	0.24-0.65	.....	7,46
51		Roots	0.370-4.539	0.035-0.620	0.09-2.83	0.013-0.498	0.03-0.23	7,12,46,62,72
52		Pollen	1.141	0.346	.....	.....	0.119	12
53		Fruits	0.878-1.530	0.284-0.402	0.631-1.062	0.516-0.645	0.095-0.168	12
54		Fruit coats	2.60	0.173	1.74	1.644	0.278	12
55		Seeds	0.952	0.441	0.948	0.477	0.101	12
56	<i>Betula populifolia</i>	Bark	0.159	0.052	0.485	0.072	0.003	12
57	(gray birch)	Wood	0.066	0.021	0.069	0.033	0.002	12
58		Fruits	0.944	0.200	0.715	0.233	0.081	12
59	<i>Carya illinoensis</i> (pecan)	Leaves	0.337-0.924	0.097-0.148	1.129-1.583	0.376-0.403	.....	40
60	<i>Catalpa speciosa</i> (northern catalpa)	Leaves	1.31-2.47	0.30-0.58	1.17-2.26	0.34-0.51	0.22-0.48	61
61	<i>Chrysanthemum</i>	Shoots	1.596	0.388	0.588	0.390	0.212	11
62	<i>segetum</i> (corn chrysanthemum)	Shoots, fl.	2.00-5.37	0.279-0.682	1.110-1.133	0.226-0.465	0.055-0.266	11
63	<i>Cinchona ledgeri- ana</i> (ledger-bark cinchona)	Leaves	0.82-1.21	0.20-0.49	0.43-0.70	0.21-0.29	.....	58
64	<i>Citrus limon</i>	Leaves	1.55-7.43	.....	3.621-4.365	0.397-0.619	.....	24,38
65	(lemon)	Bark	0.142-0.750	.....	2.088-2.864	0.151-0.460	.....	38
66		Root bark	0.159-0.578	.....	1.878-2.671	0.155-0.303	.....	38
67		Small roots	0.142-0.806	.....	0.613-0.883	0.325-0.580	.....	38
68		Fruits	1.08-1.46	0.16-0.22	0.39-0.92	0.09-0.14	0.04-0.05	7
69		Fruit rinds	0.388-0.830	.....	0.826-1.047	0.132-0.164	.....	38
70		Fruit pulp	0.560-2.041	.....	0.260-0.348	0.116-0.133	.....	38
71	<i>C. sinensis</i> (sweet orange)	Leaves	0.217-2.860	0.112-0.178	2.41-6.02	0.194-0.576	.....	25,38,76,77
72		Leaves, yg.	1.55	0.18	4.34	0.12	0.230	22,23
73		Leaves, old	0.08	0.11	8.17	0.09	0.26	22,23
74		Twig bark	0.62	0.28	5.22	0.12	0.27	22,23
75		Twig wood	0.24	0.22	1.26	0.08	0.12	22,23
76		Trunk bark	0.66	0.24	4.40	0.35	0.18	22,23
77		Trunk wood	0.21	0.16	0.69	0.08	0.11	22,23
78		Root bark	0.75	0.24	3.26	0.18	0.20	22,23
79		Root wood	0.18	0.16	0.73	0.09	0.08	22,23
80		Small roots	0.158-1.248	0.25	0.457-0.507	0.220-0.779	0.14	22,23,38
81		Fruit rinds	0.225-1.310	.....	0.544-1.010	0.08-0.16	.....	38,39
82		Fruit pulp	0.695-2.950	.....	0.060-0.394	0.088-0.180	.....	38,39
83	<i>Cornus florida</i> (flowering dog- wood)	Leaves	0.37-1.13	0.18-0.32	2.71-4.21	0.27-0.51	0.38-0.70	61
84	<i>C. mas</i> (cornelian cherry dogwood)	Fruits	1.685	0.118	0.224	0.075	0.154	12
85	<i>Cucumis sativus</i> (cucumber)	Fruits	4.46-4.51	0.450-1.153	0.518-0.689	0.307-0.331	0.366	12,21,94
86	<i>Cucurbita pepo</i>	Fruits	0.711-2.86	0.240-0.633	0.244-0.500	0.09-0.26	0.042	7,12,94
87	(pumpkin)	Seed em- bryos	0.573	0.917	0.029	0.422	.....	73

continued

111. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

Part I. MAJOR ELEMENTS

	Species (Common Name)	Plant Part	Potassium	Phosphorus	Calcium	Magnesium	Sulfur	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Dicotyledoneae								
88	<i>Daucus carota</i>	Leaves	1.326	0.234	3.181	0.280	0.395	12
89	(carrot)	Shoots	2.43-3.37	0.197-0.289	1.29-3.29	0.219-0.386	0.073-0.477	62,78
90		Roots	1.677-5.920	0.306-0.468	0.376-0.502	0.121-0.235	0.141-0.156	12,62,72
91		Fruits	1.349	0.586	2.362	0.344	0.193	12
92	<i>Fagopyrum escul-</i>	Shoots	1.66-2.55	0.127-0.540	1.68-2.65	0.442-0.607	0.103-0.114	11,50
93	<i>lentum</i> (buck-	Shoots, fl.	2.11-3.17	0.219-1.02	1.97-3.15	0.29-1.22	0.121-0.298	7,9,11
94	wheat)	Shoots, fr.	2.92	0.39	2.59	0.27	.....	33
95		Straw	0.750-3.728	0.030-0.509	0.841-1.214	0.262-0.349	0.078-0.108	2,12
96		Fruits	0.262-0.539	0.240-0.447	0.044-0.171	0.103-0.229	0.012-0.037	2,12
97	<i>Fagus sylvatica</i>	Leaves	0.450-0.822	0.176-0.178	1.022-2.048	0.173-0.221	0.037-0.103	12
98	(European	Stems	0.128	0.029	0.262	0.034	0.007	12
99	beech)	Bark	0.144-0.429	0.016-0.044	1.394-2.310	0.087-0.189	0.001-0.0048	12
100		Heartwood	0.082-0.128	0.003-0.007	0.095-0.113	0.031-0.044	0.0012-0.0062	12
101		Sapwood	0.085-0.160	0.009-0.024	0.090-0.104	0.041-0.051	0.002-0.0077	12
102		Wood	0.014-0.141	0.011-0.030	0.071-0.096	0.029-0.039	0.0046-0.0054	12
103		Shoots	0.419	0.194	0.786	0.135	0.042	11
104		Involucres	0.016-0.886	0.013-0.066	0.152-0.503	0.030-0.062	0.010-0.037	12
105		Fruits	0.520-0.879	0.315-0.487	0.371-0.650	0.175-0.311	0.036-0.076	12
106		Fruit coats	0.228-0.276	0.019-0.036	0.497-0.965	0.048-0.068	0.018-0.040	12
107		Seeds	0.980-1.212	0.441-0.493	0.307-0.515	0.194-0.252	0.076-0.095	12
108	<i>Fraxinus excel-</i>	Leaves	1.087	0.692	1.973	0.343	0.197	12
109	<i>sior</i> (European	Bark	0.285	0.070	2.355	0.058	0.025	12
110	ash)	Wood	0.040	0.0107	0.160	0.0128	0.0033	12
111		Seeds	1.56	0.282	0.661	0.168	0.154	12
112	<i>Glycine soja</i>	Leaves	0.80	0.16	3.18	0.79	0.25	54
113	(soybean)	Stems	0.67	0.20	0.89	0.42	0.27	54
114		Shoots	0.54-2.31	0.09-0.74	0.52-2.18	0.16-0.86	0.125-0.520	7,32,57,62
115		Shoots, fr.	0.93	0.31	2.10	0.76	.....	33
116		Shoots & roots	2.13-4.34	0.323-0.454	0.38-0.70	0.356-0.668	.....	41
117		Roots	1.44-1.56	0.85-1.00	2.60-5.84	1.07-3.18	.....	57
118		Fruits	1.95	0.51	0.56	0.41	0.22	54
119		Seeds	0.81-2.39	0.50-1.80	0.119-0.339	0.169-0.340	0.002-0.450	7,12
120	<i>Gossypium hir-</i>	Shoots	0.875-2.110	0.230-0.429	0.97-2.17	0.108-0.690	0.260-0.332	26,28
121	<i>sutum</i> (upland	Shoots, veg.	2.583-3.297	0.297-0.332	2.223-2.837	0.537-1.303	.....	67
122	cotton)	Shoots, fr.	1.013-1.503	0.166-0.205	1.444-1.515	0.362-0.585	.....	67
123		Burs	1.42-5.74	0.07-0.21	0.44-1.02	0.19-0.34	.....	42
124		Lint	0.46-0.75	0.025-0.124	0.013-0.27	0.07-0.11	0.04-0.06	7,26
125		Seeds	0.94-1.86	0.48-1.79	0.063-0.310	0.18-0.44	0.050-0.764	7,26,28
126	<i>Helianthus annuus</i>	Leaves	1.620-1.899	0.237-0.350	6.324-7.640	1.097-3.150	0.430-0.660	12,93
127	(common sun-	Petioles	0.84	0.118	1.49	0.591	0.092	62
128	flower)	Stems	3.23	0.07	1.72	1.20	0.14	93
129		Upper	2.523	0.133	0.650	0.241	0.088	12
130		Lower	1.386	0.085	0.507	0.157	0.168	12
131		Shoots	0.46-2.76	0.14-0.25	0.485-3.160	0.219-0.501	0.037-0.238	10-12,57,62
132		Roots	1.361-3.800	0.10-0.34	0.371-2.160	0.132-1.270	0.34	12,57
133		Flowers	1.55	0.406	0.82	0.344	0.084	62
134		Fruits	0.98-3.24	0.326-0.497	0.300-0.979	0.162-0.300	0.188	12
135		Fruit coats	0.92	0.07	0.35	0.23	0.05	93
136		Head minus fruits	9.43	0.41	2.49	1.26	0.46	93
137		Seeds	0.96	1.01	0.129-0.21	0.398-0.40	0.02	73,93
138	<i>Ilex aquifolium</i>	Leaves	0.51	0.068	0.783	0.381	0.013	12
139	(English holly)							
139	<i>Ipomoea batatas</i>	Leaves	1.61-2.37	0.19-0.28	0.71-1.18	0.45-0.54	.....	32,55
140	(sweet potato)	Shoots	3.15-4.62	.....	0.71	0.20	.....	32,74
141		Roots	0.68-1.74	0.06-0.22	0.040-0.218	0.063-0.210	0.059-0.120	7,12,26,32
142	<i>Juglans nigra</i>	Leaves	1.98-2.45	0.32-0.54	1.06-3.23	0.35-0.50	0.01-0.23	61
143	(black walnut)							
143	<i>J. regia</i> (Persian	Seeds	0.188-0.550	0.407-0.450	0.071-0.131	0.168-0.178	0.009	12
144	walnut)							
144	<i>Lactuca sativa</i>	Leaves	2.69-7.91	0.19-1.05	0.330-1.892	0.040-0.709	0.25-0.31	7,12
	(lettuce)							

continued

## III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

### Part I. MAJOR ELEMENTS

Species (Common Name)	Plant Part	Potassium	Phosphorus	Calcium	Magnesium	Sulfur	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Dicotyledoneae								
145	<i>Lycopersicon esculentum</i> (tomato)	Leaves	0.518-3.760	0.160-0.724	2.280-8.702	0.620-1.547	0.164	6,11,48,56,86,87
		Leaf blades						
146		Upper	2.670-3.492	0.800-0.808	1.610-1.747	0.670-0.681	0.98	65,66,68
147		Middle	3.763	0.822	3.633	1.210	.....	68
148		Lower	2.67-3.20	0.500-0.780	3.840-6.033	0.990-1.954	1.45	65,66,68
		Petioles						
149		Upper	6.27-7.22	0.730-0.752	1.080-1.557	0.920-1.078	0.38	65,66,68
150		Middle	6.734	0.891	2.421	1.669	.....	68
151		Lower	3.200-6.471	0.690-1.099	2.230-3.252	1.640-2.646	0.49	65,66,68
152		Stems	1.35-5.30	0.099-0.540	1.360-2.321	0.470-0.652	0.18	11,48,56,86,87
153		Upper	5.250-6.006	0.620-0.700	0.670-0.910	0.552-0.690	0.33	65,66,68
154		Middle	4.753	0.689	1.366	0.750	.....	68
155		Lower	2.270-3.450	0.410-0.703	0.990-2.084	0.500-0.995	0.28	65,66,68
156		Shoots, veg.	4.466	0.786	2.738	1.205	.....	68
157		Roots	0.796-3.41	2.34-2.45	1.26	0.46	0.75	56,65,66
158		Fruits	1.88-5.90	0.29-0.84	0.08-0.48	0.13-0.59	0.14-0.45	6,7,12,45,56,87
159		Fruits, un.	1.895	0.354	0.129	0.152	.....	6,7,12,45,56
160		Seeds	0.238-0.465	0.686	0.20	0.410	.....	45,65
161	<i>Magnolia macrophylla</i> (big-leaf magnolia)	Leaves	1.25-3.30	0.18-0.48	0.09-2.38	0.30-0.41	0.02-0.29	61
162	<i>Malus sylvestris</i> (apple)	Leaves	0.49-3.92	0.090-0.749	0.61-2.67	0.11-0.78	.....	4,13-15,18,50,51,64,71,81,88-90
163		Stems	0.71-1.39	.....	0.60-1.14	0.09-0.33	.....	18,88
164		Fruits	0.427-1.410	0.019-0.680	0.023-0.129	0.018-0.098	0.022-0.090	7,12,72
165		Fruit flesh	0.62-0.90	0.055-0.113	0.021-0.177	0.047	0.021	7,12
166		Seeds	0.828	0.705	0.250	0.355	0.047	12
167	<i>Medicago sativa</i> (alfalfa)	Shoots	0.702-4.030	0.153-0.713	0.55-3.49	0.060-1.020	0.188-0.400	5,6,11,29,34,62,78,83
168		Shoots, veg.	.....	0.315	2.05	.....	.....	10
169		Shoots, e.fl.	1.443	0.248-0.410	1.63-3.12	0.219	0.170	10,11,36
170		Shoots, full fl.	0.55-4.29	0.14-0.51	0.59-4.15	0.17-0.37	0.20-0.32	7,10,36,84
171	<i>Nicotiana tabacum</i> (common tobacco)	Leaves	0.51-7.81	0.12-0.55	1.20-6.07	0.03-2.74	0.18-1.19	7,12,26,28,32
172		Upper	3.40	0.291	4.18	0.93	0.185	12
173		Lower	4.55	0.260	3.87	0.91	0.185	12
174		Stems	1.900-4.026	0.140-0.491	0.54-2.97	0.039-0.920	0.109-0.310	12,26,28,32
175		Seeds	1.285	0.622	0.243	0.334	0.018	12
176	<i>Pastinaca sativa</i> (garden parsnip)	Roots	0.86-2.62	0.20-0.84	0.250-0.395	0.120-0.166	0.101	7,12
177	<i>Phaseolus vulgaris</i> (kidney bean)	Fruits	1.49-3.23	0.20-0.84	0.41-1.21	0.05-0.52	.....	7
178		Fruit coats	2.58	0.553	0.671	0.341	0.191	60
179		Seeds, im.	2.00	0.744	0.274	0.324	0.868	60
180		Seeds	1.176-1.890	0.412-0.500	0.104-0.179	0.148-0.200	0.052-0.227	12,60
181	<i>Pisum sativum</i> (garden pea)	Shoots	1.69	0.336	1.54	0.489	0.328	62
182		Shoots, fl.	2.31	0.358	1.34	0.459	0.246	11
183		Fruit coats	1.959	0.214	2.51	0.307	0.268	62
184		Seeds	0.814-0.977	0.380-0.607	0.057-0.114	0.127-0.163	0.037-0.156	12,62
185	<i>Populus tremula</i> (European aspen)	Leaves	1.353	0.341	3.147	0.212	0.101	12
186		Bark	0.214-0.351	0.021-0.045	0.641-1.732	0.143-0.156	0.0098-0.031	12
187		Wood	0.039-0.167	0.007-0.0076	0.069-0.202	0.009-0.034	0.0016-0.0051	12
188	<i>Prunus amygdalus</i> (almond)	Seeds	0.223-1.137	0.442-0.934	0.206-0.309	0.225-0.522	0.007-0.046	12
189	<i>P. domestica</i> (garden plum)	Fruits	0.895	0.120	0.131	0.060	0.028	12
190		Fruit without seed	1.126-1.426	0.107-0.118	0.061-0.108	0.050-0.054	0.021-0.057	12
191		Fruit skin	1.158	0.138	0.140	0.133	0.018	12
192		Fruit flesh	0.765-1.109	0.087-0.181	0.081-0.144	0.059-0.077	0.030	12
193		Seed	0.902	0.624	0.249	0.400	0.117	12
194		Endocarp	0.047	0.003	0.052	0.006	0.0007	12

continued

111. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

Part I. MAJOR ELEMENTS

Species (Common Name)	Plant Part	Potassium	Phosphorus	Calcium	Magnesium	Sulfur	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Dicotyledoneae							
195	<i>Prunus persica</i>	Leaves	0.76-2.35	0.092-0.720	1.06-2.71	0.410-1.450	50
196	(peach)	Fruit with- out seed	1.269-1.644	0.161-0.227	0.041-0.091	0.057-0.069	12
197		Fruit flesh	1.53-2.44	0.184-0.205	0.079-0.229	0.094-0.109	12
198	<i>Pyrus communis</i>	Fruits	0.894-0.938	0.105-0.131	0.066-0.112	0.052-0.062	12
199	(pear)	Fruit with- out seed	0.790	0.094	0.061	0.047	12
200		Fruit flesh	0.512	.....	0.074	0.055	12
201		Seeds	0.688	0.628	0.215	0.335	12
202	<i>Quercus robur</i>	Leaves	0.963	0.187	0.652	0.286	12
203	(English oak)	Bark	1.191	0.060	1.722	0.095	12
204		Bast and cambium	0.298-0.395	0.012-0.016	2.79-3.93	0.021-0.092	12
205		Outer bark	0.096-0.190	0.0064-0.012	1.865-5.50	0.082-0.435	12
206		Heartwood	0.048-0.088	0.0015-0.0056	0.029-0.058	0.003-0.0074	12
207		Sapwood	0.075-0.162	0.0114-0.031	0.050-0.094	0.0095-0.023	12
208		Wood	0.140	0.036	0.078	0.048	12
209		Fruits	1.161	0.142	0.108	0.069	12
210	<i>Raphanus sativus</i>	Leaves	2.270	0.272	3.095	0.452	12
211	(garden radish)	Roots	1.924-2.859	0.343	0.773-0.983	0.135-0.334	12
212	<i>Rheum palmatum</i>	Leaf blades	0.829	0.106	0.196	0.105	12
213	(sorrel rhubarb)	Petioles	0.953-7.146	0.929-1.079	0.224-1.036	0.267	12
214		Leaves	7.144	0.891	1.036	0.109	12
215		Leaves and stems	3.39	0.099	2.75	0.893	12
216		Roots	1.72	0.102	1.39	0.376	12
217	<i>Ribes nigrum</i>	Leaves	0.91-1.06	.....	.....	0.26-0.45	90,92
218	(European black currant)	Fruits	0.96-1.67	0.275-0.374	0.190-0.531	0.080-0.146	12
219	<i>Rosa centifolia</i>	Petals	1.277-1.369	0.194-0.253	0.151-0.226	0.133-0.151	12
220	(cabbage rose)						
220	<i>Salix viminalis</i>	Leaves	1.013	0.052	1.722	0.585	12
221	(basket willow)	Stems	0.272-0.332	0.050-0.073	0.489-0.538	0.040-0.082	12
222		Bark	0.732-1.024	0.017-0.170	0.886-1.247	0.095-0.205	12
223		Wood	0.307	0.026	0.207	0.084	12
224	<i>Solanum tuberosum</i>	Leaves	2.10-6.79	.....	1.08	0.231-0.860	20,32,47,52, 64,91
225		Shoots	0.03-4.19	0.083-0.260	0.58-4.12	0.220-1.15	26,32,62,85
226		Tubers	1.05-3.96	0.11-0.49	0.017-0.290	0.05-0.23	7,12,26,32,44, 62,63,72
227	<i>Trifolium pra- tense</i>	Leaves	1.846-1.969	0.231-0.337	2.030-2.759	0.468-0.710	12,31
228	(red clo- ver)	Petioles	2.630	0.481	2.157	0.612	12
229		Stems	1.678-1.958	0.122-0.320	1.086-1.284	0.389-0.454	12,31
230		Shoots	0.66-2.82	0.11-0.52	0.61-3.07	0.13-0.75	7,62,78
231		Shoots, veg.	2.16-2.99	0.320-0.529	1.92-2.02	0.510-0.559	11,43
232		Shoots, fl.	1.11-3.41	0.210-0.290	1.07-2.12	0.35-0.68	7,11,72
233		Shoots, fr.	0.974-1.840	0.226-0.392	1.33-2.03	0.494-0.602	7,72
234		Flowers	1.497-2.111	0.376-0.495	1.163-1.193	0.404-0.455	12,31
235		Seeds	1.321	0.746	0.206	0.350	12
236	<i>Ulmus americana</i>	Leaves	0.59-2.03	0.13-0.59	1.40-2.45	0.41-0.57	61
237	(American elm)						
237	<i>Vicia faba</i>	Shoots	2.09	0.267	0.917	0.211	62
238	(broad bean)	Shoots, veg.	.....	0.277-0.357	1.38-1.70	.....	10
239		Shoots, fl.	.....	0.226-0.246	1.24-1.35	.....	10
240		Fruit coats	2.291-3.450	0.109-0.138	0.567-0.829	0.187-0.422	12,62
241		Seeds	1.250-1.312	0.585-0.616	0.129-0.143	0.145-0.157	12,62
242	<i>Vitis vinifera</i>	Leaves	0.585-0.984	0.077-0.198	1.822-2.846	0.238-0.426	12
243	(European grape)	Stems	0.759	0.135	0.642	0.099	12
244		Fruits	0.838-2.422	0.191-0.353	0.078-0.399	0.040-0.132	12
245		Seeds	0.594-0.648	0.333-0.343	0.451-0.680	0.097-0.122	12

Contributor: McIlrath, Wayne J.

continued



III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

Part I. MAJOR ELEMENTS

- References:* [1] Aiyar, S. P. 1945. *Current Sci. (India)* 14:10. [2] Alway, F. J., W. N. Shaw, and W. J. Methley. 1926. *J. Agr. Res.* 33:701. [3] Archibald, J. G., E. Bennett, and W. S. Ritchie. 1943. *Ibid.* 66:341. [4] Batjer, L. P., and E. S. Degman. 1940. *Ibid.* 60:101. [5] Bear, F. E., and A. Wallace. 1950. *New Jersey Agr. Expt. Sta. Bull.* 748. [6] Bear, F. E., et al. 1951. *Ibid.* 760. [7] Beeson, K. C. 1941. *U.S. Dept. Agr. Misc. Publ.* 369. [8] Beeson, K. C., L. Gray, and M. B. Adams. 1947. *J. Am. Soc. Agron.* 39:356. [9] Bertrand, G., and V. Ghitescu. 1934. *Compt. Rend.* 199:1269. [10] Bondi, A., and H. Meyer. 1951. *Bull. Res. Council Israel* 1:126. [11] Boresch, K. 1935. *Tabulae Biologicae* 10:315. [12] Boresch, K. 1936. *Ibid.* 11:136. [13] Boynton, D. 1945. *Proc. Am. Soc. Hort. Sci.* 46:1. [14] Boynton, D., and A. B. Burrell. 1944. *Soil Sci.* 58:441. [15] Boynton, D., J. C. Cain, and O. C. Compton. 1944. *Proc. Am. Soc. Hort. Sci.* 44:15. [16] Brown, M. E. 1943. *Missouri Univ. Agr. Expt. Sta. Res. Bull.* 360. [17] Buckner, G. D., and A. H. Henry. 1945. *Kentucky Agr. Expt. Sta. Bull.* 473. [18] Cain, J. C. 1948. *Proc. Am. Soc. Hort. Sci.* 51:1. [19] Capen, R. G., and J. A. LeClerc. 1948. *J. Agr. Res.* 77:65. [20] Carolus, R. L. 1933. *Am. Potato J.* 10:147. [21] Carolus, R. L. 1935. *Proc. Am. Soc. Hort. Sci.* 31:610. [22] Chapman, H. D., and S. M. Brown. 1941. *Hilgardia* 14:161. [23] Chapman, H. D., and S. M. Brown. 1941. *Ibid.* 14:183. [24] Chapman, H. D., and S. M. Brown. 1943. *Soil Sci.* 55:87. [25] Chapman, H. D., S. M. Brown, and D. S. Rayner. 1947. *Hilgardia* 17:619. [26] Cooper, H. P., W. R. Paden, and W. H. Garman. 1947. *Soil Sci.* 63:27. [27] Cooper, H. P., and J. H. Wilson. 1930. *Ibid.* 30:421. [28] Cooper, H. P., et al. 1948. *Soil Sci. Soc. Am. Proc.* 13:323. [29] Drake, M., and E. H. Stewart. 1950. *Soil Sci.* 69:459. [30] Elser, E., and J. Ganzmüller. 1931. *Z. Physiol. Chem.* 194:21. [31] Fagan, T. W. 1928. *Welsh J. Agr.* 4:92. [32] Garner, W. W., et al. 1930. *J. Agr. Res.* 40:145. [33] Goessmann, C. A. 1891. *Mass. Agr. Expt. Sta. Ann. Rept.* 8:159. [34] Goss, A. 1903. *New Mexico Agr. Expt. Sta. Bull.* 44. [35] Greaves, J. E., and C. T. Hirst. 1929. *Utah Agr. Expt. Sta. Bull.* 210. [36] Grizzard, A. L. 1935. *J. Am. Soc. Agron.* 27:81. [37] Haas, A. R. C. 1947. *Proc. Am. Soc. Hort. Sci.* 50:200. [38] Haas, A. R. C. 1949. *Plant Physiol.* 24:395. [39] Haas, A. R. C., and L. J. Klotz. 1935. *Hilgardia* 9:179. [40] Hammar, H. E., and J. H. Hunter. 1949. *Plant Physiol.* 24:16. [41] Hampton, H. E., and W. A. Albrecht. 1944. *Missouri Univ. Agr. Expt. Sta. Res. Bull.* 381. [42] Harper, H. J., H. A. Daniels, and G. W. Volk. 1938. *J. Am. Soc. Agron.* 30:827. [43] Haselhoff, E., and S. Werner. 1913. *Landwirtsch. Jahrb. Schweiz* 44:651. [44] Headden, W. P. 1924. *Colo. Agr. Expt. Sta. Bull.* 291. [45] Hester, J. B. 1938. *Am. Fertilizer* 89:5. [46] Hirst, C. T., and J. E. Greaves. 1944. *Soil Sci.* 58:25. [47] Jones, J. O., and W. Plant. 1943. *Bristol Agr. Hort. Res. Sta. Ann. Rept.* 1942:44. [48] Kalin, E. W. 1943. *Proc. Am. Soc. Hort. Sci.* 43:235. [49] Kelley, W. P., and A. R. Thompson. 1910. *Hawaii Agr. Expt. Sta. Bull.* 21. [50] Kenworthy, A. L. 1950. *Proc. Am. Soc. Hort. Sci.* 55:41. [51] Kidson, E. B., H. O. Askew, and E. Chittenden. 1940. *J. Pomol. Hort. Sci.* 18:119. [52] Large, E. C. 1945. *Gt. Brit. Agr. Res. Council Rept.* 7849:37. [53] Latshaw, W. L., and E. C. Miller. 1924. *J. Agr. Res.* 27:845. [54] Lechartier, G. 1902-03. *Ann. Sci. Agron.*, II, 1:380. [55] Leonard, O. A., W. S. Anderson, and M. Gieger. 1949. *Proc. Am. Soc. Hort. Sci.* 53:387. [56] Lewis, A. H., and F. B. Marmoy. 1939. *J. Pomol. Hort. Sci.* 17:275. [57] Loehwing, W. F. 1934. *Plant Physiol.* 9:567. [58] Loustalot, A. J., and H. F. Winters. 1948. *Ibid.* 23:343. [59] McHargue, J. S. 1927. *Ind. Eng. Chem.* 19:274. [60] McHargue, J. S., and W. R. Roy. 1931. *J. Nutr.* 3:479. [61] McHargue, J. S., and W. R. Roy. 1932. *Botan. Gaz.* 94:381. [62] Mach, F., and R. Herrmann. 1934. *Landwirtsch. Vers. Sta.* 119:1. [63] Mangels, C. E. 1921. *J. Ind. Eng. Chem.* 13:418. [64] Nicholas, D. J. D., and J. D. Jones. 1945. *Bristol Agr. Hort. Res. Sta. Ann. Rept.* 1944:84. [65] Nightingale, G. T., L. G. Schermerhorn, and W. R. Robbins. 1930. *New Jersey Agr. Expt. Sta. Bull.* 499. [66] Nightingale, G. T., et al. 1931. *Plant Physiol.* 6:605. [67] Olson, L. C., and R. P. Bledsoe. 1942. *Georgia Agr. Expt. Sta. Bull.* 222. [68] Phillips, T. G., T. O. Smith, and R. B. Dearborn. 1934. *New Hampshire Agr. Expt. Sta. Tech. Bull.* 59. [69] Pierre, W. H., and R. R. Robinson. 1937. *J. Am. Soc. Agron.* 29:477. [70] Reuther, W. 1948. *Proc. Am. Soc. Hort. Sci.* 51:137. [71] Reuther, W., and D. Boynton. 1940. *Ibid.* 37:32. [72] Robinson, W. O., L. A. Steinkoenig, and C. F. Miller. 1917. *U.S. Dept. Agr. Bull.* 600. [73] Schulze, E., and C. Godet. 1908. *Z. Physiol. Chem.* 58:156. [74] Scott, L. E. 1950. *Proc. Am.*

*continued*

### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

#### Part I. MAJOR ELEMENTS

Soc. Hort. Sci. 56:248. [75] Sen, A. T. 1938. Burma Dept. Agr. Rept. 1937-38:35. [76] Smith, P. F., and W. Reuther. 1951. Plant Physiol. 26:110. [77] Smith, P. F., W. Reuther, and A. W. Specht. 1950. Ibid, 25:496. [78] Strigel, A. 1912. Landwirtsch. Jahrb. Schweiz 43:349. [79] Strutzer, A. 1907. Landwirtsch. Vers. Sta. 65:264. [80] Sullivan, B., and C. Near. 1927. Ind. Eng. Chem. 19:498. [81] Thompson, A. H., R. S. Marsh, and O. E. Schubert. 1952. West Va. Univ. Agr. Expt. Sta. Bull. 356. [82] Thompson, A. R. 1908. Hawaii Agr. Expt. Sta. Ann. Rept., p. 51. [83] Vandecaveye, S. C., and G. O. Baker. 1944. J. Agr. Res. 68:191. [84] Vandecaveye, S. C., and L. V. Bond. 1936. J. Am. Soc. Agron. 28:491. [85] Von Daszewski, A., and B. Tollens. 1900. J. Landwirtsch. 48:223. [86] Wall, M. E. 1939. Soil Sci. 47:143. [87] Wall, M. E. 1940. Ibid. 49:315 [88] Wallace, T. 1929. J. Pomol. Hort. Sci. 8:23. [89] Wallace, T. 1940. Ibid. 18:145. [90] Wallace, T. 1940. Ibid. 18:261. [91] Wallace, T., J. O. Jones, and W. Plant. 1942. Bristol Agr. Hort. Res. Sta. Ann. Rept. 1941:39. [92] Wallace, T., and D. A. Osmond. 1941. Ibid. 1940:13. [93] Wiley, H. W. 1901. U.S. Dept. Agr. Div. Chem. Bull. 60. [94] Wilkins, L. K. 1917. New Jersey Agr. Expt. Sta. Bull. 310. [95] Wimer, D. C. 1937. Illinois Univ. Agr. Expt. Sta. Bull. 437.

#### Part II. MINOR ELEMENTS

Many of the ranges cover a vast literature, but because of space limitations only the more recent references (which frequently cite the earlier literature) have been included. Values are mg/kg of dry weight.

	Species (Common Name)	Plant Part	Boron	Cop- per	Iron	Manga- nese	Zinc	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Monocotyledoneae								
1	<i>Allium cepa</i> (garden onion)	Bulbs	.....	2-24	24-265	2-96	100	21,93,121,143,144
2	<i>Asparagus officinalis</i> (garden as- paragus)	Shoots, yg.	.....	7-17	60-979	12-29	52	21,40,88,121
3	<i>Avena sativa</i> (common oat)	Leaves	.....	.....	.....	12-240	19-276	2,66,94,95,169
4		Shoots	.....	0.7-17	.....	5-93	4-29	93,139,140
5		Shoots, veg.	.....	4-12	.....	79-90	18-40	138,139
6		Shoots, fl.	15-50	3-4	50-270	5-82	12-25	27,28,44,82,138,139
7		Shoots, fr.	2-17	1-9	154	5-116	12-13	41,127,138,139,148
8		Straw	8	2-54	61-860	4-1,656	4-193	45,53,84,109,139
9	<i>Hordeum vulgare</i> (barley)	Grain	1-19	0.7-51	7-350	14-76	22-40	45,53,84,139,169
10		Shoots	4-53	14	180-450	.....	.....	44,93,127,130
11	<i>Lilium</i> spp. (lily)	Grain	0.6-13	1-70	14-350	7-38	21-132	8,15,60,114,151
12		Stigmas	13-14	.....	.....	2	.....	31-35
13	<i>Oryza sativa</i> (rice)	Anthers	8-13	.....	.....	2	.....	31-35
14		Roots	.....	8-16	.....	430	.....	45,149
15	<i>Phleum pratense</i> (timothy)	Grain	9.4	3-4	76-350	18-70	30	18,45,105,149,151
16		Shoots	10-16	2-7	30-287	11-165	30-60	13,43,127,155
17	<i>Phoenix dactylifera</i> (date palm)	Grain	.....	.....	61-410	72	.....	45,98
18		Pinnae	73-172	.....	.....	.....	.....	70
19	<i>Poa pratensis</i> (Kentucky blue- grass)	Fruit flesh	8-18	2-5	35-70	2-54	4	17,18,21,52,70,91,92, 135
20		Leaves, veg.	3	14	.....	80	170	60,102
21		Shoots	6-12	.....	60-425	29-216	17-28	100,104,108,154,155, 158
22	<i>Triticum aestivum</i> (wheat)	Grain	.....	60	350-460	85-110	360	45,98,102
23		Shoots	3-10	3-12	290-580	.....	13-25	38,39,41,44,93,139
24		Straw	9-10	1-18	60-630	22-150	7-25	13,21,61,84,108,114
25	<i>Zea mays</i> (corn)	Grain	1-11	3-24	3-420	5-260	19-105	8,13,15,79,84,108, 123,160
26		Leaves	27-72	.....	41-810	230-440	.....	14,60,80,87
27		Stems	.....	.....	400-740	100-230	.....	87
28		Roots	.....	.....	500-760	450-880	.....	87
29		Shoots	15-18	.....	312-321	52-200	.....	126,158
30		Stover	.....	2-9	94-345	54-270	5-80	13,61,151,155
31	Kernels	.....	1-9	4-30	13-550	5-500	20	8,17,18,21,121,151
32		Cobs	.....	.....	250	310	.....	87

continued

### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

#### Part II. MINOR ELEMENTS

	Species (Common Name)	Plant Part	Boron	Cop- per	Iron	Manga- nese	Zinc	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Dicotyledoneae								
33	<i>Acer saccharum</i> (sugar maple)	Leaves	.....	11-12	150-440	40-220	24-54	107,145
34		Twigs	.....	.....	55	170	.....	145
35		Bark	.....	.....	86	261	.....	145
36		Sapwood	.....	.....	27	59	.....	145
37		Heartwood	.....	.....	18	55	.....	145
38	<i>Beta vulgaris</i> (beet)	Leaves	7-29	9-18	142-1,932	16-205	.....	91,121,126,143
39		Roots	2-46	6-27	69-290	19-104	25-69	13,21,54,84,143
40	<i>Capsicum frutescens</i> (bush red pepper)	Leaves	34-118	.....	.....	.....	.....	60
41		Fruits	21	8-18	.....	13-70	.....	54,60
42		Fruit flesh	.....	16	57-630	18-19	.....	7,92,117,121,135,137
43	<i>Carya illinoensis</i> (pecan)	Leaves	.....	21-28	144-185	.....	4-202	1,64,65
44		Seeds	3	14	26	36	.....	13,104,135,137
45	<i>Catalpa speciosa</i> (northern catalpa)	Leaves	.....	18-21	330-680	80-130	28-50	107
46	<i>Citrus limon</i> (lemon)	Leaves	19-200	2-13	288	14-75	.....	51,73,74
47		Fruit rinds	23-35	3-4	60	.....	.....	71,74,135
48		Fruit pulp	12-26	5	420	3	.....	29,30,71,74,117
49	<i>C. sinensis</i> (sweet orange)	Leaves	17-386	7-18	38-345	24-46	24-47	50,71,73,89,156,157
50		Roots	95	.....	.....	.....	240	50
51		Fruits	22-27	3-4	.....	.....	.....	71,74
52		Fruit rinds	10-27	6	.....	.....	.....	71,73,74,104,152
53		Fruit pulp	10-38	1-31	19-70	3	12	21,30,71,73,74,120
54	<i>Cornus florida</i> (flowering dogwood)	Leaves	23	7-9	240-380	30-50	3-28	104,107
55	<i>Cucumis sativus</i> (cucumber)	Fruits	.....	5-30	33-420	24-48	44	21,91,121,137
56	<i>Cucurbita pepo</i> (pumpkin)	Fruits	.....	11	.....	13	.....	67,81
57		Fruit flesh	.....	.....	.....	4	44	20,21,137
58	<i>Daucus carota</i> (carrot)	Shoots	20-45	5-10	355-765	23-199	26	21,57,127,143
59		Roots	21-57	5-20	39-490	6-91	10	13,21,54,60,93,121,143
60	<i>Fagopyrum esculentum</i> (buck- wheat)	Shoots	.....	.....	100	14	15	27,28
61		Fruits	.....	.....	34-170	.....	12	20,21,45,135
62	<i>Fragaria</i> spp. (strawberry)	Fruits & floral receptacles	.....	2-8	68-267	6-40	4	21,91,92,120
63	<i>Fraxinus americana</i> (white ash)	Leaves	.....	.....	195	94	.....	146
64		Twigs	.....	.....	55	27	.....	146
65		Bark	.....	.....	78	96	.....	146
66		Sapwood	.....	.....	15	Trace	.....	146
67		Heartwood	.....	.....	16	Trace	.....	146
68	<i>Glycine soja</i> (soybean)	Leaves	30-148	8	336	29-192	110	78,99,127,158,159
69		Stems	8-38	.....	.....	7-20	.....	78,127,158
70		Shoots	1-13	4-12	100-570	45-280	28-80	13,55,61,125,155
71		Roots	14-16	.....	.....	17-147	.....	127,158,159
72		Seeds	6-41	12-23	57-161	14-41	18	61,99,121,127,158
73	<i>Gossypium</i> spp. (cotton)	Leaves	60-795	.....	1,754	80-100	.....	45,60,86,113
74		Stems	6-186	.....	610	40-50	.....	45,86,113
75		Lint	.....	.....	190	11-190	.....	86,101
76		Seeds	27-130	54	150-590	13-31	320	45,86,101,113
77	<i>Helianthus annuus</i> (common sun- flower)	Shoots	12-150	.....	.....	72-1,268	.....	114,128,161
78		Fruits	.....	.....	34	23	19	20,21,98
79	<i>Ilex opaca</i> (American holly)	Leaves	.....	6-14	200-270	260-540	130-240	107
80	<i>Ipomoea batatas</i> (sweet potato)	Tuberous roots	4-44	3-9	11-140	3-28	5	21,54,60,121,122,143
81	<i>Juglans</i> spp. (walnut)	Seeds	15	8	200	19-33	.....	8,45,81,92,104
82	<i>J. nigra</i> (black walnut)	Leaves	40-67	11	280-780	60-190	26-49	107,147
83	<i>Lactuca sativa</i> (garden lettuce)	Leaves	13-75	3-33	65-4,830	15-523	105-119	13,19,54,93,118,121, 143
84	<i>Lycopersicon esculentum</i> (common tomato)	Leaflets	92-147	12	277-542	70-398	17-30	96,97,134
85		Leaves	21-150	12-21	106-840	53-4,930	.....	14,47,60,90,127,147
86		Stems	21-26	6-13	110-230	14-45	.....	90,147
87		Shoots	24	11-12	2,000	32-100	.....	9,68,77
88		Fruits	13-36	4-34	32-800	2-410	2-67	13,54,60,90,97,103, 143
89	<i>Magnolia macrophylla</i> (big-leaf magnolia)	Leaves	.....	6-8	150-230	29-30	19-62	107
90	<i>Malus sylvestris</i> (apple)	Leaves	11-43	3-12	65-507	20-156	4-345	12,46,48,164
91		Stems	9-21	.....	.....	.....	16-80	49,110,170
92		Fruits	3-76	5-7	1-69	1-22	3-9	10,21,76,91,92,117

continued



### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

#### Part II. MINOR ELEMENTS

	Species (Common Name)	Plant Part	Boron	Cop- per	Iron	Manga- nese	Zinc	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
	Dicotyledoneae							
93	<i>Medicago sativa</i> (alfalfa)	Leaves	4-654	.....	.....	45-76	.....	59,60,83,111
94		Shoots	12-128	4-61	110-675	10-124	13-112	11,13,63,100
95		Shoots, veg.	29	80	140-410	29-101	37	41,139,148,167
96		Shoots, fl.	25-50	.....	161-1,000	14-936	14	16,27,28,167
97		Seeds	10	.....	.....	6-15	.....	17,18,36,80,98,142
98	<i>Nicotiana</i> spp. (tobacco)	Leaves	6-93	17	70-2,100	48-2,262	.....	6,14,61,80,124,129
99		Seeds	6	.....	240	70	.....	98,104
100	<i>Pastinaca sativa</i> (garden parsnip)	Roots	.....	7	46-210	.....	.....	91,117,135,136
101	<i>Persea</i> spp. (persea)	Fruits	36-132	.....	12-575	.....	.....	69,117,120,165
102	<i>Phaseolus vulgaris</i> (kidney bean)	Shoots, fr.	.....	.....	1,350	124	.....	148
103		Plant	6-12	.....	.....	32-68	.....	60,133
104		Fruits	2-37	6-20	52-769	11-57	8	13,21,54,143,163
105		Fruit coats	22-40	12	270	39	53	106,127
106		Seeds, im.	.....	10	80-270	16-19	46	13,98,106
107		Seeds	17	.....	120	20	.....	17,18,36,148
108	<i>Pisum sativum</i> (garden pea)	Leaves	16	.....	.....	38	.....	60,83
109		Shoots	17-22	.....	.....	15	.....	22-26,81
110		Fruits	.....	7-12	.....	21-27	.....	81,83,116,144
111		Seeds	2-8	6-15	70-282	4-25	40-48	57,85,104,142,166
112	<i>Prunus amygdalus</i> (almond)	Seeds	15-57	11-13	42-190	13	21	21,72,85,91,137
113	<i>P. domestica</i> (garden plum)	Leaves	.....	7-10	.....	55-93	.....	3,62
114	<i>P. persica</i> (peach)	Leaves	17-81	.....	.....	17-325	6-345	4,49,60,62,112
115		Stems	7-44	.....	.....	.....	11-50	49,60,112
116		Fruits	38-52	.....	140	.....	2	20,21,58,104,117
117	<i>Pyrus communis</i> (pear)	Leaves	.....	5-41	28-94	.....	.....	14,80,119,131,132
118		Bark	.....	4-17	53-120	.....	.....	119,132
119		Wood	.....	2-12	5-20	.....	.....	119,132
120		Fruit	19	6-13	29-140	2-4	9	21,58,91,92,117
121	<i>Quercus velutina</i> (black oak)	Leaves	.....	7-9	250-280	490-1,870	36-68	107
122	<i>Raphanus sativus</i> (garden radish)	Leaves	20-196	.....	141-224	.....	.....	60,126
123		Roots	17-152	29	97-825	14-45	23	21,60,89,117,126,137
124	<i>Rheum</i> spp. (rhubarb)	Petioles	.....	9	154-680	31	22	20,21,91,117,135,137
125	<i>Ribes</i> spp. (currant)	Fruits	19-58	17	80-420	4	.....	45,58,91,92
126	<i>Solanum tuberosum</i> (potato)	Leaves	20-98	30-89	.....	.....	.....	56,60,84
127		Shoots	14-39	11	.....	86-108	.....	38,39,41,43,57,114
128		Tubers	2-22	0-28	28-363	3-94	11-14	13,21,54,84,114,141
129	<i>Trifolium pratense</i> (red clover)	Leaves	57	.....	.....	40-84	.....	42,83
130		Stems	28	.....	.....	15-20	.....	42,83
131		Shoots	31-36	6-20	100-1,300	25-542	24-80	13,63,93,104
132		Shoots, veg.	23-58	.....	.....	.....	.....	41
133		Shoots, fl.	19-109	.....	.....	287	.....	41,148
134		Flowers	40	.....	.....	30-66	.....	42,83
135		Seeds	.....	17	21-336	6-38	76	13,83,98,100
136	<i>Ulmus americana</i> (American elm)	Leaves	277	7-16	245-810	39-130	10-22	60,107,146
137		Twigs	14	.....	68	20	.....	60,146
138		Bark	.....	.....	145	30	.....	146
139		Sapwood	.....	.....	48	9	.....	146
140		Heartwood	.....	.....	22	2	.....	146
141	<i>Vicia faba</i> (broad bean)	Shoots	.....	.....	280	36	.....	114,168
142		Straw	.....	.....	63	23	.....	114,168
143		Seeds	11-223	10-11	21	14-15	.....	37,67,114
144	<i>Vitis</i> spp. (grape)	Leaves	16-2,084	3-47	190-220	180-220	.....	75,115,162
145		Stems	5-50	15-16	28-33	37-46	.....	60,115,153
146		Fruits	15-34	5-10	13-530	Trace	2	5,13,21,92,150

Contributor: McIlrath, Wayne J.

References: [1] Alben, A. O., and H. E. Hammar. 1939. Proc. Texas Pecan Growers' Assoc. 19:48. [2] Albert, W. B. 1932. S. Carolina Agr. Expt. Sta. Ann. Rept. 1931-32:46. [3] Anderssen, F. G. 1932. J. Pomol. Hort. Sci. 10:130. [4] Archibald, E., and F. B. Wann. 1942. Am. J. Botany 29:694. [5] Askew, H. O. 1944. Better Crops Plant Food 29(5):21. [6] Askew, H. O. 1949. New Zealand J. Sci. Technol., A, 28:161. [7] Askew, H. O.,

continued



## III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

### Part II. MINOR ELEMENTS

- R. H. K. Thomson, and E. Chittenden. 1938. *Ibid.*, A, 20:74. [8] Bagchi, K., and S. Chowdhury. 1949. *Ann. Biochem. Exptl. Med. (Calcutta)* 9:107. [9] Bailey, L. F., and J. S. McHargue. 1943. *Am. J. Botany* 30:558. [10] Batjer, L. P., and M. H. Haller. 1942. *Proc. Am. Soc. Hort. Sci.* 40:29. [11] Bear, F. E., and A. Wallace. 1950. *New Jersey Agr. Expt. Sta. Bull.* 748. [12] Beattie, J. M., and C. W. Ellenwood. 1950. *Proc. Am. Soc. Hort. Sci.* 55:47. [13] Beeson, K. C. 1941. *U.S. Dept. Agr. Misc. Publ.* 369. [14] Bennett, J. P. 1945. *Soil Sci.* 60:91. [15] Bergh, H. 1948. *Kgl. Norske Videnskab. Selskabs Skrifter* 1942-45:1. [16] Bertrand, G. 1941. *Ann. Agron.* 11:1. [17] Bertrand, G. 1942. *Ibid.* 12:189. [18] Bertrand, G. 1942. *Ann. Inst. Pasteur* 68:457. [19] Bertrand, G., and M. Andreitcheva. 1934. *Ibid.* 52:249. [20] Bertrand, G., and B. Benzon. 1928. *Bull. Soc. Sci. Hyg. Aliment.* 16:457. [21] Bertrand, G., and B. Benzon. 1929. *Ann. Inst. Pasteur* 43:386. [22] Bertrand, G., and H. L. deWaal. 1936. *Ann. Agron.* 6:536. [23] Bertrand, G., and H. L. deWaal. 1936. *Ann. Inst. Pasteur* 57:121. [24] Bertrand, G., and H. L. deWaal. 1936. *Bull. Soc. Chim. France, Ser. 5,* 3:875. [25] Bertrand, G., and H. L. deWaal. 1936. *Compt. Rend. Acad. Agr. France* 22:321. [26] Bertrand, G., and H. L. deWaal. 1936. *Compt. Rend.* 202:605. [27] Bertrand, G., and V. Ghitescu. 1934. *Compt. Rend. Acad. Agr. France* 20:1052. [28] Bertrand, G., and V. Ghitescu. 1934. *Compt. Rend.* 199:1269. [29] Bertrand, G., and M. Rosenblatt. 1921. *Ann. Inst. Pasteur* 35:815. [30] Bertrand, G., and M. Rosenblatt. 1921. *Compt. Rend.* 173:333. [31] Bertrand, G., and M. Rosenblatt. 1921. *Ibid.* 173:1118. [32] Bertrand, G., and L. Silberstein. 1938. *Ann. Inst. Pasteur* 61:102. [33] Bertrand, G., and L. Silberstein. 1938. *Bull. Soc. Chim. France, Ser. 5,* 5:1069. [34] Bertrand, G., and L. Silberstein. 1938. *Compt. Rend.* 206:796. [35] Bertrand, G., and L. Silberstein. 1938. *Compt. Rend. Acad. Agr. France* 24:597. [36] Bertrand, G., and L. Silberstein. 1941. *Compt. Rend.* 213:221. [37] Bertrand, G., and L. Silberstein. 1944. *Ann. Agron.* 14:257. [38] Bertrand, G., L. Silberstein, and H. L. deWaal. 1936. *Ibid.* 6:183, 537. [39] Bertrand, G., L. Silberstein, and H. L. deWaal. 1937. *Ibid.* 7:333, 505. [40] Bishop, W. B. S. 1928. *Australian J. Exptl. Biol. Med. Sci.* 5:125. [41] Bobko, E. V. 1940. *Compt. Rend. Acad. Sci. URSS* 29:510. [42] Bobko, E. V., and V. V. Zerling. 1938. *Ann. Agron.* 8:174. [43] Bolin, D. W. 1934. *J. Agr. Res.* 48:657. [44] Boresch, K. 1935. *Tabulae Biologicae* 10:315. [45] Boresch, K. 1936. *Ibid.* 11:136. [46] Bould, C. D., et al. 1950. *Nature* 165:920. [47] Brennan, E. G., and J. W. Shive. 1948. *Soil Sci.* 66:65. [48] Chabannes, J., S. Trocme, and G. Barbier. 1950. *Ann. Agron.* 20:362. [49] Chandler, W. H., D. R. Hoagland, and P. L. Hibbard. 1933. *Proc. Am. Soc. Hort. Sci.* 30:70. [50] Chapman, H. D., S. M. Brown, and D. S. Rayner. 1947. *Hilgardia* 17:619. [51] Chapman, H. D., G. F. Liebig, and E. R. Parker. 1939. *Calif. Citrograph* 25:11. [52] Cleveland, M. M., and C. R. Fellers. 1932. *Ind. Eng. Chem., Anal. Ed.* 4:267. [53] Coic, Y., M. Coppenet, and S. Voix. 1950. *Compt. Rend.* 230:1610. [54] Coleman, J. M., and R. W. Ruprecht. 1935. *J. Nutr.* 9:51. [55] Cook, F. C. 1916. *J. Agr. Res.* 5:877. [56] Cook, F. C. 1921. *Ibid.* 22:281. [57] Cunningham, I. J. 1931. *Biochem. J.* 25:1267. [58] Dodd, A. S. 1929. *Analyst* 54:15. [59] Dregne, H. E., and W. L. Powers. 1942. *J. Am. Soc. Agron.* 34:902. [60] Eaton, F. M. 1944. *J. Agr. Res.* 69:237. [61] Elvehjem, C. A., and E. B. Hart. 1929. *J. Biol. Chem.* 82:473. [62] Epstein, E., and O. Lilleland. 1942. *Proc. Am. Soc. Hort. Sci.* 41:11. [63] Evans, H. J., and E. R. Purvis. 1948. *J. Am. Soc. Agron.* 40:1046. [64] Finch, A. H. 1936. *J. Agr. Res.* 52:363. [65] Finch, A. H., and A. F. Kinnison. 1933. *Ariz. Univ. Agr. Expt. Sta. Tech. Bull.* 47. [66] Gerretsen, F. C. 1949. *Plant Soil* 1:347. [67] Guerithault, B. 1920. *Compt. Rend.* 171:196. [68] Gum, O. B., H. D. Brown, and R. C. Burrell. 1945. *Plant Physiol.* 20:267. [69] Haas, A. R. C. 1943. *Calif. Avocado Soc. Yearbook,* p. 41. [70] Haas, A. R. C. 1944. *Proc. Am. Soc. Hort. Sci.* 44:34. [71] Haas, A. R. C. 1945. *Plant Physiol.* 20:323. [72] Haas, A. R. C. 1945. *Proc. Am. Soc. Hort. Sci.* 46:69. [73] Haas, A. R. C. 1945. *Soil Sci.* 59:465. [74] Haas, A. R. C., and H. J. Quayle. 1935. *Hilgardia* 9:143. [75] Hansen, C. J. 1945. *Proc. Am. Soc. Hort. Sci.* 46:781. [76] Heinicke, A. J., W. Reuther, and J. C. Cain. 1942. *Ibid.* 40:31. [77] Hester, J. B. 1938. *Am. Fertilizer* 89:5. [78] Hodgkiss, S. W., R. H. Hageman, and J. S. McHargue. 1942. *Plant Physiol.* 17:652. [79] Hoffman, C., T. R. Schweitzer, and G. Dalby. 1943. *Cereal Chem.* 20:328. [80] Jacobson, L. 1945. *Plant Physiol.* 20:233. [81] Jadin, F., and A. Astruc. 1913. *J. Pharm. Chim.* 7:155. [82] Jones, H. E., and G. D. Scarseth. 1944. *Soil*

*continued*

### III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION

#### Part II. MINOR ELEMENTS

- Sci. 57:15. [83] Jones, J. S., and D. E. Bullis. 1921. J. Ind. Eng. Chem. 13:524. [84] Katalymov, M. V. 1946. Compt. Rend. Acad. Sci. URSS 53:821. [85] Kohler, G. O., C. A. Elvehjem, and E. B. Hart. 1936. J. Biol. Chem. 113:49. [86] Kruglova, E. K. 1943. Chem. Abstr. 37:4191. [87] Latshaw, W. L., and E. C. Miller. 1924. J. Agr. Res. 27:845. [88] Levine, H., F. B. Culp, and C. B. Anderson. 1932. J. Nutr. 5:295. [89] Levitt, E. C., and R. I. Nicholson. 1941. Agr. Gaz. N.S. Wales 52:283. [90] Lewis, A. H., and F. B. Marmoy. 1939. J. Pomol. Hort. Sci. 17:275. [91] Lindow, C. W., C. A. Elvehjem, and W. H. Peterson. 1929. J. Biol. Chem. 82:465. [92] Lindow, C. W., and W. H. Peterson. 1927. Ibid. 75:169. [93] Lucas, R. E. 1948. Soil Sci. 65:461. [94] Lundegårdh, H. 1931. Kgl. Landbruks-Akad. Handl. Tidskr. 70:1021. [95] Lundegårdh, H. 1931. Medd. Centralanstalt. Foersoksvaesendet Jordbruks. 403:1. [96] Lyon, C. B., and K. C. Beeson. 1943. J. Am. Soc. Agron. 35:166. [97] Lyon, C. B., K. C. Beeson, and G. H. Ellis. 1943. Botan. Gaz. 104:495. [98] McHargue, J. S. 1923. J. Agr. Res. 23:395. [99] McHargue, J. S. 1925. Ibid. 30:193. [100] McHargue, J. S. 1925. J. Am. Soc. Agron. 17:368. [101] McHargue, J. S. 1926. Ibid. 18:1076. [102] McHargue, J. S. 1927. Ind. Eng. Chem. 19:274. [103] McHargue, J. S., and R. K. Calfee. 1937. J. Am. Soc. Agron. 29:385. [104] McHargue, J. S., W. S. Hodgkiss, and E. B. Offutt. 1940. Ibid. 32:622. [105] McHargue, J. S., and W. R. Roy. 1931. Am. J. Physiol. 99:221. [106] McHargue, J. S., and W. R. Roy. 1931. J. Nutr. 3:479. [107] McHargue, J. S., and W. R. Roy. 1932. Botan. Gaz. 94:381. [108] McHargue, J. S., W. R. Roy, and J. G. Pelphrey. 1932. J. Am. Soc. Agron. 24:562. [109] McHargue, J. S., and O. M. Shedd. 1930. Ibid. 22:739. [110] McLarty, H. R., J. C. Wilcox, and C. G. Woodbridge. 1936. Proc. Wash. State Hort. Assoc. 32:142. [111] McLarty, H. R., J. C. Wilcox, and C. G. Woodbridge. 1937. Sci. Agr. 17:515. [112] McLarty, H. R., and C. G. Woodbridge. 1950. Ibid. 30:392. [113] McLean, R. C., and W. L. Hughes. 1936. Ann. Appl. Biol. 23:231. [114] Mach, F., and R. Herrmann. 1934. Landwirtsch. Vers. Sta. 119:1. [115] Magoon, C. A., et al. 1938. Proc. Am. Soc. Hort. Sci. 36:485. [116] Maquenne, L., and E. Demoussy. 1920. Bull. Soc. Chim. France, Ser. 4, 27:266. [117] Marañon, J. 1935. Philippine J. Sci. 58:317. [118] Midgley, A. R., and D. E. Dunklee. 1946. Better Crops Plant Food 30:17. [119] Milad, Y. 1924. Proc. Am. Soc. Hort. Sci. 21:93. [120] Miller, C. D., K. Bazore, and R. C. Robbins. 1937. Hawaii Agr. Expt. Sta. Bull. 77. [121] Miller, C. D., W. Ross, and L. Louis. 1947. Hawaii Agr. Expt. Sta. Tech. Bull. 5. [122] Mitchell, J. H., and W. T. Mattison. 1933. S. Carolina Agr. Expt. Sta. Ann. Rept. 46:51. [123] Moran, T. 1945. Nature 155:205. [124] Morgan, M. F., and O. E. Street. 1938. Conn. Agr. Expt. Sta. New Haven Bull. 410. [125] Morris, H. D., and W. H. Pierre. 1949. J. Am. Soc. Agron. 41:107. [126] Muhr, G. R. 1942. Soil Sci. 54:55. [127] Munsell, R. I., and B. A. Brown. 1943. J. Am. Soc. Agron. 35:401. [128] Neidig, R. E., and R. S. Snyder. 1925. J. Agr. Res. 31:1165. [129] New Zealand Department of Scientific and Industrial Research. 1939. Annual report. Wellington. v. 13, p. 75. [130] Odellien, M. 1932. Mededeel. Inst. Suikerbieteneteelt 2:43. [131] Oserkowsky, J., and H. E. Thomas. 1933. Science 78:315. [132] Oserkowsky, J., and H. E. Thomas. 1938. Plant Physiol. 13:451. [133] Parbery, N. H. 1943. Agr. Gaz. N.S. Wales 54:14. [134] Parks, R. Q., C. B. Lyon, and S. L. Hood. 1944. Plant Physiol. 19:404. [135] Peterson, W. H., and C. A. Elvehjem. 1928. J. Biol. Chem. 78:215. [136] Peterson, W. H., and C. A. Hoppert. 1925. J. Home Econ. 17:265. [137] Peterson, W. H., and J. T. Skinner. 1931. J. Nutr. 4:419. [138] Piper, C. S. 1942. J. Agr. Sci. 32:143. [139] Piper, C. S., and A. Walkley. 1943. Australia J. Council Sci. Ind. Res. 16:217. [140] Rademacher, B. 1940. Bodenkunde Pflanzenernaehr. 19:80. [141] Ranninger, R. 1944. Biol. Generalis 18:126. [142] Rao, A. L. S. 1940. J. Indian Chem. Soc. 17:351. [143] Remington, R. E., and H. E. Shiver. 1930. J. Assoc. Offic. Agr. Chemists 13:129. [144] Richards, M. B. 1930. Biochem. J. 24:1572. [145] Riou, P., and G. Delorme. 1938. Compt. Rend. 207:300. [146] Riou, P., and G. Delorme. 1938. Ibid. 207:1244. [147] Robinson, W. O., and G. Edgington. 1942. Soil Sci. 53:309. [148] Robinson, W. O., L. A. Steinkoenig, and C. F. Miller. 1917. U.S. Dept. Agr. Bull. 600. [149] Sarata, U. 1938. Japan. J. Med. Sci., II, 4:193. [150] Satterfield, G. H., and S. O. Jones. 1932. J. Elisha Mitchell Sci. Soc. 48:16. [151] Schaible, P. J., S. L. Bandemer, and J. A. Davidson. 1938. Mich. State Univ. Agr. Expt. Sta. Tech. Bull. 159. [152] Scofield, C. S., and L. V. Wilcox. 1930. Science 71:542.

*continued*

**III. PLANT TISSUES AND ORGANS: MINERAL COMPOSITION****Part II. MINOR ELEMENTS**

- [153] Scott, L. E. 1944. Soil Sci. 57:55. [154] Shorland, F. B. 1934. Trans. Proc. Roy. Soc. New Zealand 64:35. [155] Skinner, J. T., and W. H. Peterson. 1928. J. Biol. Chem. 79:679. [156] Smith, P. F., and W. Reuther. 1951. Plant Physiol. 26:110. [157] Smith, P. F., W. Reuther, and A. W. Specht. 1950. Ibid. 25:496. [158] Snider, H. J. 1943. Soil. Sci. 56:187. [159] Somers, I. I., and J. W. Shive. 1942. Plant Physiol. 17:582. [160] Sullivan, B. 1933. Cereal Chem. 10:503. [161] Tanada, T., and L. A. Dean. 1942. Hawaiian Planters' Record 46:65. [162] Teakle, L. J. H., H. K. Johns, and A. G. Turton. 1943. J. Dept. Agr. W. Australia 20:171. [163] Terlikowski, F., and B. Nowicki. 1932. Roczniki Nauk Rolniczych Lesnych 28:135. [164] Thomas, W., W. B. Mack, and F. N. Fagan. 1947. Proc. Am. Soc. Hort. Sci. 50:1. [165] Tilt, J., and M. Winfield. 1928. J. Home Econ. 20:43. [166] Walsh, T., and S. J. Cullinan. 1945. Proc. Roy. Irish Acad., B, 50:279. [167] Weathers, E. K. 1938. Tenn. Univ. Agr. Expt. Sta. Bull. 166. [168] Wester, D. H. 1921. Biochem. Z. 118:158. [169] Wood, J. G., and P. M. Sibly. 1950. Australian J. Sci. Res., B, 3:14. [170] Woodbridge, C. G. 1937. Sci. Agr. 18:41.

## XI. ENVIRONMENT AND SURVIVAL

### 112. HIBERNATION: MAMMALS AND BIRDS

Hibernation in homiotherms is a lethargic condition characterized by a lowering of the temperature of the body to approximately the temperature of the environment--with a concurrent reduction in metabolism--and the resumption of the elevated temperature at some future time without the aid of heat from external sources.

	Species (Common Name)	Distribution	Temperature °C		Heart Rate beats/ min	Respi- ration Rate breaths/ min	O <sub>2</sub> Con- sumption ml/g body wt/hr	CO <sub>2</sub> Pro- duction ml/g body wt/hr	RQ <sup>1</sup>	Ref- erence
			Air	Rectal						
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Mammalia										
1	<i>Citellus citellus</i> (sus- lik)	Central Asia, south- ern Russia to Aus- tria	.....	6	.....	.....	0.320	0.230 <sup>2</sup>	0.72	15
2			7	7.2	.....	.....	0.015	.....	...	14
3			11	11.7	.....	5	.....	.....	...	9
4			13	15.5	.....	.....	0.034	.....	...	20
5	<i>C. tridecemlineatus</i>	Central United States, Canada	.....	3-10 <sup>3</sup>	5-20	14-15	0.081-0.191	.....	...	1.10
6	(thirteen-lined ground squirrel)		4.0	5.7	.....	1	0.081	.....	...	11
7			8.6	10.2	.....	1.6	0.125	.....	...	11
8			12.5	13.6	.....	1.8	0.197	.....	...	11
9	<i>C. undulatus</i> (Arctic ground squirrel)	Alaska, northern Can- ada, northeastern Siberia	.....	5.2	68 <sup>4</sup>	10	.....	.....	...	8
10			5.9	5.9	.....	6	.....	.....	...	8
11	<i>Eptesicus fuscus</i> (big brown bat)	United States, south- ern Canada	8	9	.....	3-10 <sup>5</sup>	.....	.....	...	21
12			22-26	.....	.....	.....	0.8	.....	...	8
13	<i>Erinaceus europaeus</i>	Great Britain to Spain, Italy, Greece	2-3	6.2-7.7	18-24	.....	0.014-0.033	.....	...	26,27
14	(European hedgehog)		3.5	5	.....	.....	0.88	0.83	0.68	5
15			6	.....	.....	.....	0.40	0.29 <sup>2</sup>	0.73	15
16			9.7	12.0	.....	.....	0.126	0.056	...	5
17	<i>Glis glis</i> (fat dormouse)	Central and southern Europe	6	.....	.....	.....	0.029	0.021 <sup>2</sup>	0.72	15
18			11.8	.....	.....	.....	0.024	.....	...	14
19	<i>Marmota marmota</i> (Eur- asian marmot)	Alps	10	10.5	.....	0.35 <sup>5</sup>	0.018	0.012	0.68	14
20	<i>M. monax</i> (woodchuck)	Eastern United States, Canada, Alaska	.....	4-7	4-5	.....	0.008-0.034	.....	...	2
21			1.2	5.8	.....	16	.....	.....	...	8
22			8	8	.....	6	.....	.....	...	8
23	<i>Mesocricetus auratus</i>	Rumania, eastern Asia Minor, Syria, Pales- tine, northwestern Iran	5	5-6	.....	.....	0.183	0.132 <sup>2</sup>	0.72	16
24	(golden hamster)		5.5	5.5 <sup>3</sup>	.....	.....	0.032	.....	...	18
25			5.8	6.4 <sup>3</sup>	.....	.....	0.06	.....	...	18
26			5	5	4-15	.....	0.060-0.080	.....	...	18,19
27	<i>Muscardinus avellanarius</i>	Southern Italy to En- gland and Sweden	6	.....	.....	9-10	.....	.....	...	25
28	(common dormouse)		10.1	.....	.....	.....	0.80	0.57 <sup>2</sup>	0.71	14
29			11.6	.....	10-12	.....	.....	.....	...	28
30	<i>Myotis keenii</i> (long-eared little brown bat)	Eastern United States, British Columbia	21.5	22.7 <sup>e</sup>	.....	140-168	0.85	.....	...	8
31	<i>M. lucifugus</i> (little brown bat)	Northern United States, southern Canada, southern Alaska	0.5	.....	.....	.....	0.113	.....	...	7
32			.....	2	7-10	.....	0.022-0.039	.....	...	7,24
33			10	.....	.....	.....	0.071	.....	...	7
34		23	23.2 <sup>e</sup>	.....	72-80	0.45	.....	...	7	
35	<i>M. myotis</i> (common brown bat)	Europe to China to Af- ghanistan	1.7	.....	.....	.....	0.020	0.009	...	6
36			2.5	.....	.....	.....	0.051	0.033	0.65	6
37	<i>Nyctalus noctula</i> (noctule bat)	Europe to Siberia, Japan to Palestine	4.3	.....	.....	.....	0.51	0.38 <sup>2</sup>	0.75	15
38			12.5	.....	.....	.....	3.49 <sup>7</sup>	2.58 <sup>2</sup>	0.74	15
39			20	.....	.....	.....	0.403 <sup>2</sup>	0.314	0.78	13
40			30	.....	.....	.....	0.682 <sup>2</sup>	0.484	0.71	13
41	<i>Pipistrellus pipistrellus</i>	Europe to northern Asia, Japan	.....	5	.....	.....	0.247	0.175 <sup>2</sup>	0.71	15
42	(European brown bat)		5	.....	.....	.....	0.053	0.038 <sup>2</sup>	0.72	15

/1/ Respiratory quotient. Probably does not reflect actual exchange of gases or the true nature of combustion of foods during hibernation. /2/ Calculated. /3/ Oral temperature. /4/ Feeble heart beat in deep hibernation, becoming more evident as awakening progresses. /5/ Respiration rates are very irregular in deep hibernation, and there may be several minutes with no respiration followed by several respirations. Cheyne-Stokes respiration is not uncommon; range is average of several minutes. /6/ Subcutaneous temperature. /7/ During awakening from hibernation.

*continued*



## 112. HIBERNATION: MAMMALS AND BIRDS

	Species (Common Name)	Distribution	Temperature °C		Heart Rate beats/ min	Respi- ration Rate breaths/ min	O <sub>2</sub> Con- sumption ml/g body wt/hr	CO <sub>2</sub> Pro- duction ml/g body wt/hr	RQ <sup>1</sup>	Ref- er- ence
			Air	Rectal						
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Mammalia										
43	<i>Plecotus auritus</i> (long-eared bat)	Europe to Japan, eastern Siberia to Sudan	0	.....	.....	.....	0.037	.....	...	4
44			5	6.5	.....	.....	0.069	0.049 <sup>2</sup>	0.71	13
45			10	10.7	.....	.....	0.094	0.079 <sup>2</sup>	0.84	13
46			19.7	.....	.....	.....	0.255	.....	...	12
47	<i>Rhinolophus ferrum-equinum</i> (greater horseshoe bat)	England to Korea, Japan to Morocco	13	13	.....	.....	0.150	0.089	...	3
48			19	19	.....	.....	0.426	0.366	0.77	3
49	<i>R. hipposideros</i> (lesser horseshoe bat)	Europe to Asia Minor, northwestern India to Sudan	15	15	.....	.....	2.23	1.80	0.80	3
50	<i>Ursus americanus</i> (black bear) <sup>3</sup>	N. America (north of Mexico)	4.4	35.5	.....	2-3	.....	.....	...	29
51	<i>Vespertilio murinus</i> (parti-colored bat)	Europe to Japan, northwestern India	0	.....	.....	.....	.....	0.037	...	4
52			7.05	7.05	50-55	.....	.....	.....	...	23
53			8	.....	.....	.....	.....	0.020	...	4
Aves										
54	<i>Apus apus</i> (swift)	N. America, Europe, Africa, Asia	19	23 <sup>2</sup>	.....	8-10 <sup>2</sup>	0.7	0.31	...	17
55	<i>Calypte anna</i> (Anna's hummingbird)	United States (California)	24	.....	.....	.....	0.84	.....	...	22
56	<i>Selasphorus sasin</i> (Allen's hummingbird)	United States (California)	22	.....	.....	.....	1.24	.....	...	22

<sup>1/1</sup> Respiratory quotient. Probably does not reflect actual exchange of gases or the true nature of combustion of foods during hibernation. <sup>2/2</sup> Calculated. <sup>3/3</sup> Respiration rates are very irregular in deep hibernation, and there may be several minutes with no respiration followed by several respirations. Cheyne-Stokes respiration is not uncommon; range is average of several minutes. <sup>4/4</sup> Not a true hibernator, as indicated by the discrepancy between rectal and ambient temperature. <sup>5/5</sup> Proventricular temperature, taken orally.

Contributors: (a) Hock, Raymond J., (b) Lyman, Charles P.

References: [1] Baldwin, F. M., and K. L. Johnson. 1941. J. Mammal. 22:130. [2] Benedict, F. G., and R. C. Lee. 1938. Carnegie Inst. Wash. Publ. 497. [3] Burbank, R. C., and J. Z. Young. 1934. J. Physiol. (London) 82:459. [4] Delsaux, E. 1887. Arch. Biol. (Liege) 7:207. [5] Gorer, P. A., and M. S. Pembrey. 1929. J. Physiol. (London) 67:xxi. [6] Hari, P. 1909. Arch. Ges. Physiol. 130:112. [7] Hock, R. J. 1951. Biol. Bull. 101:289. [8] Hock, R. J. Unpublished. U.S. Public Health Service, Arctic Health Research Center, Anchorage, Alaska, 1955. [9] Horvath, A. 1872. Zentr. Med. Wiss. 10:865. [10] Johnson, G. E. 1928. J. Exptl. Zool. 50:15. [11] Johnson, K. L. 1940. Master's Thesis. Univ. Southern California, Los Angeles. [12] Kayser, C. 1938. Compt. Rend. Soc. Biol. 128:1204. [13] Kayser, C. 1939. Ann. Physiol. Physicochim. Biol. 15:1087. [14] Kayser, C. 1942. Biol. Abstr. 16:19728. [15] Kayser, C. 1950. Arch. Sci. Physiol. 4:361. [16] Kayser, C. 1952. Compt. Rend. Soc. Biol. 146:929. [17] Koskimies, J. 1950. Ann. Acad. Sci. Fennicae, A IV, 15:1. [18] Lyman, C. P. 1948. J. Exptl. Zool. 109:55. [19] Lyman, C. P. 1951. Am. J. Physiol. 167:638. [20] Mares, F. 1892. Compt. Rend. Soc. Biol. 44:313. [21] Pearson, O. P. 1947. Ecology 28:127. [22] Pearson, O. P. 1950. Condor 52:145. [23] Prunelle, M. 1811. Ann. Museum Hist. Nat. 18:20. [24] Rawson, K. S. Unpublished. Dept. of Biology, Swarthmore College, Penna., 1955. [25] Saissy, J. A. 1811. Mem. Acad. Sci. Turin (2):1. [26] Sarajas, H. S. S. 1954. Acta Physiol. Scand. 32(1):28. [27] Schenk, P. 1922. Arch. Ges. Physiol. 197:66. [28] Spallanzani, L. 1803. Opere. U. Hoepli, Milan. [29] Svihla, A., and H. S. Bowman. 1954. Am. Midland Naturalist 52:248.

## 113. DIAPAUSE: INSECTS AND MITES

Diapause in insects and mites may be facultative (influenced by the environment and not present in each generation) or obligate (occurring in virtually every individual and in each generation without respect to environment). Species experiencing facultative diapause generally complete two or more generations annually (bivoltine or multivoltine cycles), whereas those with obligate diapause produce one generation (univoltine cycle). **Dormant Stage** (column B): I = instar; FI = final instar; FD = fully developed; S = small; HG = half-grown.

	Species (Common Name)	Dormant Stage	Diapause		Refer- ence
			Type	Duration	
	(A)	(B)	(C)	(D)	(E)
Insecta					
1	<i>Aedes hexodontus</i> (mosquito)	Embryo (FD)	Obligate	Throughout winter	5
2	<i>A. triseriatus</i> (mosquito)	Embryo (FD)	Facultative	Up to 6 months	3
3	<i>Anax imperator</i> (darnier)	Larva (FI)	Facultative	2-3 months	13
4	<i>Anthrenus verbasci</i> (varied carpet beetle)	Larva (S-FI)	Obligate	Many months	7
5	<i>Apanteles glomeratus</i> (little braconid)	Prepupa	Facultative	.....	19
6	<i>Bombyx mori</i> (silkworm)	Embryo (S or HG)	Facultative <sup>1</sup> or obligate <sup>2</sup>	5-9 months	22
7	<i>Cephus cinctus</i> (wheat stem sawfly)	Prepupa	Obligate	6-8 months	31
8	<i>Ceratophyllus fasciatus</i> (European rat flea)	Larva (FI)	Obligate	2-12 months	2
9	<i>Drosophila deflexa</i> (fruit fly)	Larva (FI)	Facultative	Several months	4
10	<i>Dytiscus marginalis</i> (diving beetle)	Adult	Obligate <sup>3</sup>	.....	21
11	<i>Ephestia elutella</i> (tobacco moth)	Larva	Facultative	8-9 months	34
12	<i>Epilachna corrupta</i> (ladybird beetle)	Adult	Facultative	Throughout winter	17
13	<i>Eurydema ornatum</i> (ornate vegetable bug)	Adult	Facultative	.....	10
14	<i>Eurygaster integriceps</i> (soun pest)	Adult	Obligate	.....	18
15	<i>Leptinotarsa decemlineata</i> (Colorado potato beetle)	Adult	Facultative <sup>5</sup>	Many months	11,12
16	<i>Lestes sponsa</i> (damselfly)	Egg	Obligate	15 weeks	14
17	<i>Locusta migratoria gallica</i> (migratory locust)	Egg	Obligate	14-156 days	23
18	<i>Malacosoma disstria</i> (forest tent caterpillar)	Embryo (FD)	Obligate	6-9 months	20
19	<i>Melanoplus bivittatus</i> (two-striped grasshopper)	Embryo (FD)	Obligate	Few weeks to many months	32
20	<i>M. differentialis</i> (differential grasshopper)	Egg	Obligate	Few weeks to many months	8,9
21	<i>Melittobia chalybii</i> (chalcid)	Larva (FI)	Facultative	Several months	33
22	<i>Phaenicia sericata</i> (greenbottle fly)	Larva (FI)	Facultative	Several weeks to many months	15,29
23	<i>Pieris rapae</i> (imported cabbageworm)	Pupa	Facultative	6-52 weeks	28
24	<i>Popillia japonica</i> (Japanese beetle)	Larva (3rd I)	Facultative	50 days	25,26
25	<i>Pyrausta nubilalis</i> (European corn borer)	Larva (FI)	Facultative <sup>1</sup> or obligate <sup>2</sup>	6-9 months	1
26	<i>Reduvius personatus</i> (masked hunter)	Larva	Obligate	Few weeks to several months	30
27	<i>Samia cynthia</i> (cynthia moth)	Pupa	Facultative	.....	16
28	<i>Trichogramma cacaeciae</i> (fairfly)	Larva (FI)	Facultative	6-8 months	27
Acari					
29	<i>Metatetranychus ulmi</i> (European red mite)	Egg	Facultative	.....	6,24
30	<i>Tetranychus telarius</i> (two-spotted spider mite)	Adult	Facultative	.....	24

<sup>1/2</sup> Bivoltine strains. <sup>2/2</sup> Univoltine strains. <sup>3/2</sup> Reproductive diapause involving corpus allatum.

Contributors: (a) Andrewartha, H. G., (b) Lees, A. D., (c) Wilkes, A.

References: [1] Arbuthnott, K. D. 1949 U.S. Dept. Agr. Tech. Bull. 869. [2] Bacot, A. 1914. J. Hyg., Plague Suppl. 3:447. [3] Baker, F. C. 1935. Can. Entomologist 67:149. [4] Basden, E. B. 1954. Proc. Roy. Entomol. Soc. London, A, 29:114. [5] Beckel, W. E. 1958. Can. J. Zool. 36:541. [6] Blair, C. A., and J. R. Groves. 1952. J. Hort. Sci. 27:14. [7] Blake, G. M. 1958. Bull. Entomol. Res. 49:751. [8] Bodine, J. H. 1929. Physiol. Zool. 2:459. [9] Bodine, J. H., et al. 1939. J. Cellular Comp. Physiol. 14:173. [10] Bonnemaison, L. 1948. Compt. Rend. 227:985. [11] Breitenbrecher, J. K. 1918. Carnegie Inst. Wash. Publ. 263:341. [12] Busnel, R. G., and A. Drilhon. 1937. Compt. Rend. Soc. Biol. 124:916. [13] Corbet, P. S. 1954. Ph. D. Thesis. Cambridge Univ., England. [14] Corbet, P. S. 1956. Proc. Roy. Entomol. Soc. London, A, 31:45. [15] Cragg, J. B., and P. Cole. 1952. J. Exptl. Biol. 29:600. [16] Danilyevsky, A. S. 1939. Zool. Zh. 18:1926. [17] Douglass, J. R. 1928. J.

continued

## 113. DIAPAUSE: INSECTS AND MITES

Econ. Entomol. 21:203. [18] Fedotov, D. M. 1944. Compt. Rend. Acad. Sci. URSS 42(9):408. [19] Gayspitz, K. F., and I. I. Kyao. 1953. Entomol. Obozrenie 33:32. [20] Hodson, A. C., and C. J. Weinman. 1945. Univ. Minn. Agr. Expt. Sta. Tech. Bull. 170. [21] Joly, P. 1945. Arch. Zool. Exptl. Gen. 84:49. [22] Kogure, M. 1933. J. Dept. Agr. Kyushu Imp. Univ. 4:1. [23] LeBerre, J. R. 1951. Rev. Zool. Agr. Appl. 10-12. [24] Lees, A. D. 1953. Ann. Appl. Biol. 40:449. [25] Ludwig, D. 1932. Physiol. Zool. 5:431. [26] Ludwig, D. 1953. J. Gen. Physiol. 36:751. [27] Marchal, P. 1936. Ann. Epiphyt. Phytogenet. 2:447. [28] Mazaki, S. 1955. Oyo Dobutsugaku Zasshi 20:98. [29] Melanby, K. 1938. Parasitology 30:392. [30] Readio, P. A. 1931. Ann. Entomol. Soc. Am. 24:19. [31] Salt, R. W. 1947. Can. J. Res., D, 25:66. [32] Salt, R. W. 1949. Ibid., D, 27:236. [33] Schmieder, R. G. 1933. Biol. Bull. 65:338. [34] Waloff, N. 1949. Trans. Roy. Entomol. Soc. London 100:147.

## 114. DISPERSION OF SMALL ORGANISMS

Units Dispersed (columns C-G): Values are means.

### Part I. INVERTEBRATES

Dispersion by flight may possibly be aided by air movements.

Invertebrate (Common Name) [ Means of Dispersion ]		Distances and Units Dispersed					Refer- ence	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
1	<i>Aedes</i> spp. (mosquito) [flight]	Miles from original case Yellow fever cases	17.5 14.6	32.5 4.6	47.5 1.7	62.5 0.8	92.5 0.6	91
2	<i>A. aegypti</i> (yellow-fever mosquito) [flight]	Meters from release point Eggs in traps	250 1,100	750 46	1,250 40	1,750 5	2,250 2	98
3	<i>A. albopictus</i> (mosquito) [flight]	Yards from release point Mosquitoes recovered	25 92	125 43	225 2	325 1	475 2	5
4	<i>A. communis</i> (mosquito) [flight]	Feet from release point (subarctic) Mosquitoes recovered	75 80	300 24	1,000 6	2,000 1	5,000 1	43
5	<i>A. leucocelaenus</i> & <i>Haemagogus spe-gazzinii</i> (forest mosquitoes) [flight]	Kilometers from release point Mosquitoes recovered/100 man-hr	1.0 7	2.0 1	2.4 0.5	2.7 0.2	4.0 0.1	12
6	<i>A. polynesiensis</i> (mosquito) [flight]	Yards from release point Mosquitoes recovered	0 37	50 8	100 3	150 0		41
7	<i>Agriotes obscurus</i> (click beetle) [walking, running]	Feet from release point Beetles/trap	3 12	9 3	18 0.6	36 0.1		79
8	<i>Anastatus bifasciatus</i> (gypsy moth egg parasite) [flight]	Yards from release point Eggs parasitized, % (north) Eggs parasitized, % (south)	100 23 32	900 12 20	1,500 9 18	3,900 4 13	5,100 3 12	16
9	<i>Anopheles funestus</i> (mosquito) [flight]	Yards from riverbank Mosquitoes caught	100 179	600 14	1,200 5	2,400 4		20
10	<i>A. gambiae</i> (African malaria mosquito) [flight]	Miles from release point Mosquitoes recovered	0.75 3.0	1.25 2.2	1.75 1.6	2.25 1.2		20
11	<i>A. pseudopunctipennis</i> (mosquito) [flight]	Miles from release point Mosquitoes recovered	0-1 16	1-2 10	2-3 8	3-4 6		77
12	<i>A. quadrimaculatus</i> (malaria mosquito) [flight]	Feet from source Mosquitoes caught	500 71	1,500 43	2,500 30	3,500 22	6,500 6	81
13		Feet from mosquito source Malarial infections, %	500 36	1,500 22	3,500 11	6,000 4		81
14		Miles from breeding-site reservoir Females caught/wk	0.5 542	1.5 79	2.5 5	3.5 3	4.5 1	27
15	<i>Anthonomus grandis</i> (boll weevil) [flight]	Yards from overwinter area Weevils trapped	220 59	440 43	880 26	1,320 17	1,760 9	32
16	<i>A. grandis</i> [crawling]	Feet from release point Females recovered Males recovered	25 17 28	75 8 6	125 3 2	175 0 1	225 0 0	75
17	<i>Apis mellifera</i> (honeybee) [flight]	Yards from apiary Honeybees found	5.5 6.1	11 3.9	16.5 2.6	22 1.6	33 0.3	26

continued



## 114. DISPERSION OF SMALL ORGANISMS

### Part I. INVERTEBRATES

	Invertebrate (Common Name) [ Means of Dispersion ]	Distances and Units Dispersed					Refer- ence	
		(A)	(B)	(C)	(D)	(E)		(F)
Horizontal Dispersion								
18	<i>Apis mellifera</i> (honeybee) [flight]	Yards from apiary	200	500	833	1,126	1,426	57
		Pollination/2 sq yd	8	7	7	6	6	
19		Feet from apiary	130	630	1,130	1,730	2,330	6
		Honeybees/2 sq yd	21	13	9	7	5	
		Yield of red clover seed, lb/2 sq yd	65	34	22	14	8	
20	<i>Bruchus pisorum</i> (pea weevil) [flight]	Feet from field margin	300	600	1,400			51
		Weevils found	51	50	48			
21		Miles from overwinter area	1	2	3	4	5	90
		Weevils found	13	8	5	3	2	
22	<i>Calendra maidis</i> (maize billbug) [crawling]	Rows from field margin	3	9	33	50		11
		Beetles found	19	11	3	1		
23	<i>Camnula pellucida</i> (clear-winged grasshopper) & <i>Melanoplus mexicanus</i> (migratory grasshopper) [crawling <sup>1</sup> ]	Yards from release point	10	50	90	160		78
		Grasshoppers recovered	179	85	51	18		
24	<i>Carpocapsa pomonella</i> (codling moth) [flight]	Feet from release point	75	189	264	332		87
		Moths recovered, %	57	25	13	5		
25	<i>Catocala</i> spp. (moth) [flight]	Feet from release point	13	88	163	238	363	7
		Moths recovered	8.1	3.7	2.4	1.5	0.6	
26	<i>Chalcodermus aeneus</i> (cowpea curculio) [unknown]	Feet from field margin	1.0	11.5	37.5	105.0	138.5	4
		Curculios found	1.49	0.71	0.40	0.14	0.07	
27	<i>Chionaspis furfura</i> (scurfy scale) [crawling]	Inches from dispersal point	1.5	7.5	13.5	19.5	22.5	39
		Mature scales recovered	44	7	3	0.7	0.1	
28	<i>Circulifer tenellus</i> (beet leafhopper) [flight]	Miles from breeding area	15	35	52	105	215	31
		Leafhoppers caught	499	151	76	15	4	
29	<i>Cochliomyia hominivorax</i> (screw-worm) [crawling]	Inches from carcass	3	9	15	21		86
		Larvae/sq ft	320	47	4	2		
30	<i>C. macellaria</i> (secondary screwworm) [flight]	Miles from release point (rural)	0.5	2.5	4.5	6.5	9.0	70
		Flies recovered	2.2	0.2	0.1	0.1	0.1	
31	<i>C. macellaria</i> <sup>2</sup> [flight]	Miles from release point	0.25	1	2	3		69
		Traps containing flies, %	70	56	42	34		
32	<i>Conotrachelus nemuphar</i> (plum curculio) [flight]	Feet from release point	50	136	235	335	478	83
		Curculios recovered	48	13	5	2	1	
33	<i>Culex quinquefasciatus</i> (southern house mosquito) [flight]	Miles from release point	0.2	0.5	0.75	1.0	2.5	74
		Mosquitoes recovered	47	3	2	5	1	
34	<i>C. tarsalis</i> (mosquito) [flight]	Miles from release point	0.25	0.75	1.25	1.75	2.5	74
		Mosquitoes recovered	13	12	2	0	1	
35	<i>Culicoides grahamii</i> (punkie) [flight]	Yards from edge of breeding site	5	25	50	100	150	62
		Punkies caught biting	20	12	9	6	4	
36	<i>C. impunctatus</i> (punkie) [flight]	Yards from breeding center	0	28	57	80	89	47,48
		Males caught	1,163	311	130	43	16	
		Females caught	832	258	135	77	58	
37	<i>Cylas formicarius elegantulus</i> (sweet potato weevil) [flight]	Yards from source	440	880	940	1,320	1,760	14
		Sweet potato plants infested, %	42	32	31	26	21	
38	<i>Daphnia pulex</i> (water flea) [crawling]	Inches from source	0.5	2.5	4.5	6.5	8.5	8
		Fleas found	21	9	5	2	0	
39	<i>Dendroctonus monticolae</i> (mountain pine beetle) [flight]	Yards from road	2	5	10	15		3
		Trees killed, %	63	26	10	1		
40	<i>D. valens</i> (red turpentine beetle) [flight]	Feet from source logs	50	200	400			36
		Infestation	63	27	9			
41	<i>Diabrotica duodecimpunctata</i> (spotted cucumber beetle) [flight]	Feet from field margin	11.5	37.5	105.0	138.0	273.0	4
		Beetles found	0.32	0.27	0.22	0.20	0.17	
42	<i>D. vittata</i> (striped cucumber beetle) [flight]	Miles from release point	0.25	0.75	1.25	1.75		25
		Beetles recovered	31	11	4	1		
43	<i>Dissosteira longipennis</i> (high plains grasshopper) & <i>Melanoplus mexicanus</i> (migratory grasshopper) [flight]	Miles from release point	25	75	125	175	225	93
		Grasshoppers recovered	8	6	6	5	5	
44	<i>Drosophila funebris</i> (fruit fly) [flight]	Meters from release point	5	25	45	65	75	85
		Flies recovered	17.1	1.7	0.5	0	0.1	

1/-/ Some flight by adults. 2/ Yellow-eyed mutant strain.

continued



## 114. DISPERSION OF SMALL ORGANISMS

### Part I. INVERTEBRATES

Invertebrate (Common Name) [ Means of Dispersion ]		Distances and Units Dispersed						Refer- ence
		(A)	(B)	(C)	(D)	(E)	(F)	
Horizontal Dispersion								
45	<i>Drosophila melanogaster</i> (fruit fly) [ flight ]	Meters from release point	5	15	25	35	45	85
		Flies recovered	12.7	5.7	2.1	0.7	0.3	
46	<i>D. repleta</i> (fruit fly) [ flight ]	Feet from release point	25	150	250	350	450	67
		Flies recovered	30	6	2	1	0	
47	<i>Empoasca fabae</i> (potato leafhopper) [ flight ]	Miles from nearest land	3	6	9	10		82
		Leafhoppers caught	38	27	21	20		
48	<i>Epitrix cucumeris</i> (potato flea beetle) [ flight ]	Feet from field margin	100	200	300	400	500	94
		Injuries/tuber	9	8	7	7	6	
49	<i>Glossina</i> sp. (tsetse fly) [ flight ]	Miles from release point	0	0.67	1.33	2.67	3.33	42
		Males recovered	593	208	136	51	3	
50		Yards from thicket to host animals	25	50	100	125		58
		Flies caught during dry season	52	13	7	2		
51		Yards from thicket to host animals	80	180	230	280		58
		Flies caught during wet season	35	14	8	3		
52		Miles from fly belt	0.13	0.63	1.3	3.0	4.0	59
		Compounds ruined, %	83	40	29	17	13	
53	<i>C. morsitans</i> (tsetse fly) [ flight ]	Yards (following man)	50	1,000	2,000	4,000	6,000	84
		Flies found	30	13	9	5	3	
54	<i>Grapholitha molesta</i> (oriental fruit moth) [ flight ]	Feet from orchard	100	1,320	2,640			30
		Moths caught	194	14	8			
55	<i>Harmolita grandis</i> (wheat strawworm), spring form [ crawling ]	Feet from wheat stubble	10	20	30	40	60	50
		Infestation, %	4.2	3.0	2.1	1.4	0.5	
56	<i>H. grandis</i> , summer form [ flight ]	Feet from wheat stubble	1	50	100	150		50
		Infestation, %	52	19	13	10		
57	<i>H. tritici</i> (wheat jointworm) [ flight ]	Yards from wheat stubble	58	174	290	450		13
		Adults caught	18	9	4	0		
58	<i>Heliothis armigera</i> (corn earworm) [ flight ]	Rows from light traps (convergence)	1	3	5	10		10
		Plants infested, %	32.4	31.9	31.6	31.3		
59	<i>Hippodamia convergens</i> (convergent lady beetle) [ flight ]	Miles from release point	0.5	1.5	2.5	3.5	5.5	18
		Beetles recovered	1.1	0.6	0.4	0.3	0.1	
60	<i>Hylurgopinus rufipes</i> (elm bark beetle) [ flight ]	Feet from source	60	120	320	579	816	96
		Beetles/sq ft	14	9	5	2	1	
61	<i>Laemophloeus minutum</i> (flat grain beetle) [ flight ]	Feet from probable source	50	100	200	400		24
		Beetles caught	11	8	6	3		
62	<i>Liriomyza pusilla</i> (serpentine leaf miner) [ flight ]	Feet from field margins	20	100	140	180	260	95
		Mines/leaf	76	64	50	31	17	
63		Feet from field margins	9	100	300	400	600	95
		Potato yield, bushels/acre	171	188	202	206	211	
64	<i>Littorina</i> sp. (periwinkle) [ crawling ]	Inches from release point	0.5	1.5	3.5	4.5	5.5	8
		Periwinkles found	20	9.5	3.0	2.0	1.0	
65	<i>Lydella stabulans griseescens</i> (tachinid fly) [ flight ]	Miles from release point	0.5	1.5	2.5	4.0	6.0	2
		European corn borers parasitized, %	27	18	13	9	5	
66	<i>L. stabulans griseescens</i> [ flight <sup>3</sup> ]	Miles from colonization point	0.5	2	4	6		55
		European corn borer larvae parasitized, %	25	16	12	9		
67	<i>Macrosteles divinus</i> (six-spotted leaf- hopper) [ flight, crawling ]	Feet from release point	50	100	150	200		54
		Leafhoppers recovered	9	3	1	1		
68		Feet from release point	30	225	450			54
		Days to first leafhopper recovery	4	13	16			
69	<i>M. divinus</i> [ flight ]	Miles from nearest land	3	6	9	10		82
		Leafhoppers caught	516	145	37	18		
70	<i>Meianoplus</i> spp. <sup>4</sup> (grasshopper) [ crawling ]	Feet from release point	100	200	300	400		60
		Grasshoppers on bare ground	17	10	6	3		
71	<i>Merodon equestris</i> (narcissus bulb fly) [ flight ]	Feet from old planting	7	85	150	200	300	23
		Plants infested, %	42	21	16	13	10	
72	<i>Musca domestica</i> (housefly) [ flight ]	Yards from release point	35	110	220	330		97
		Flies recovered	572	189	49	28		
73		Miles from release point	0.43	1.5	2.5	3.5	4.5	71
		Tagged flies/trap	106	12	3	2	0.3	
74	<i>Myzus persicae</i> (green peach aphid) [ flight ]	Rows from field margin	10	20	40	80	160	49
		Aphids/100 late potato leaves	3,076	1,700	1,009	661	484	

<sup>3</sup>/ Except for passive transportation by other insects. <sup>4</sup>/ See also lines 23 and 43.

continued

114. DISPERSION OF SMALL ORGANISMS

Part I. INVERTEBRATES

Invertebrate (Common Name) [ Means of Dispersion]		Distances and Units Dispersed					Refer- ence	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
75	<i>Pectinophora gossypiella</i> (pink boll-worm) [flight]	Feet from moth source Worms/boll	1,000 1.33	2,750 1.22	3,500 1.20	6,250 1.16	63	
76	<i>Phaenicia</i> sp. (blowfly) [flight]	Miles from release point Flies recovered	0.25 31	0.5 11	0.75 5	1 2	1.25 0.5	80
77	<i>P. sericata</i> (greenbottle fly) [flight]	Miles from release point Flies recovered, %	0.5 1.8	1 1.3	2 0.3	4 0.1	6 0	53
78	<i>Phormia regina</i> (black blowfly) [flight]	Miles from release point Flies recovered	0.8 1.8	1.4 1.3	2.4 0.8	3.4 0.5	4.4 0.3	80
79		Miles from release point Flies recovered	0.25 0.12	0.5 0.02	0.75 0.03	1.13 0		80
80		Miles from release point Flies recovered, %	0.5 9.1	2 0.8	4 0.1	7 0	12 0	53
81	<i>Phytophaga destructor</i> (Hessian fly) [flight]	Feet from wheat field Flies caught	100 95	400 16	600 12			56
82		Feet from hibernation Plants infested, %	7.5 44	27.5 16	47.5 9	97.5 3		46
83	<i>Popillia japonica</i> (Japanese beetle) [crawling]	Inches from release point Larvae recovered	22 79	42 44	62 23	84 7		37
84	<i>P. japonica</i> [flight]	Feet from field margin Damage to corn, %	0 3.4	38.5 0.8	73.5 0.4	118.5 0		99
85	<i>Porthetria dispar</i> (gypsy moth) [flight]	Feet from source Males caught	330 67	660 22	1,320 5	2,640 3		15
86	<i>P. dispar</i> [carried by wind]	Feet from source Larvae found	50 16	150 4.4	250 1.7	350 1.3	600 0.5	9
87	<i>Psila rosae</i> (carrot rust fly) [flight]	Yards inside field Larval mines/100 carrots	1 78	5 48	15 28	30 15	50 6	100
88	<i>P. rosae</i> [burrowing in soil]	Yards from plants Pupae found	15 94	25 42	35 15	45 0		65
89	<i>Psorophora</i> sp. (rice-field mosquito) [flight]	Miles from release point Mosquitoes recovered	0.5 8	2.5 0.6	4.5 0.2	6.5 0.1		72
90	<i>P. confinis</i> (rice-field mosquito) [flight]	Miles from release point Mosquitoes/trap	0.5 3.7	2.5 1.8	4.5 1.1	8.5 0.4	11.5 0	40
91	<i>Pyrausta nubilalis</i> (European corn borer) [flight]	Rows from light trap convergence Plants infested, %	1 85	2 62	3 51	4 41	5 36	38
92	<i>P. nubilalis</i> [crawling]	Inches from source Larvae found	40 56	56 34	80 13	89 7	103 0	61
93	<i>Rhabdocnemis obscura</i> (New Guinea sugarcane weevil) [flight, crawling]	Feet from release point Weevils recovered Days to recovery	265 26 52	405 16 60	600 8 70	740 4 76	825 1 78	88
94	<i>Rhagoletis pomonella</i> (apple maggot) [flight]	Feet from release point Maggots recovered	37.5 47	87.5 33	137.5 26	187.5 21	237.5 17	66
95	<i>Saissetia oleae</i> (black scale) [air currents]	Feet from source Scales caught	13 564	35 433	100 293	250 172	450 92	73
96	<i>Scolytus multistriatus</i> (smaller European elm bark beetle) [flight]	Feet from source Twig crotches wounded, %	50 39	100 23	200 14	400 8	1,000 3	89
97	<i>Simulium</i> sp. (blackfly) [flight]	Miles from release point Flies recovered/visit	0.5 2.3	2.5 0.4	4.5 0.1	9.5 0.1	10.5 0	17
98		Days between release and recovery Flies caught	5 30	25 0	45 1	65 2		17
99	<i>Sitophilus oryza</i> (rice weevil) [probably crawling]	Feet from source Weevils found	50 3	100 2	200 1			24
100	<i>Tiphia vernalis</i> (Japanese beetle parasite) [flight]	Feet from feeding area Eggs/larva	7 3.7	21 3.2	49 2.7	77 2.5	90 2.4	34
101	<i>Trioza tribunctata</i> (blackberry psyllid) [crawling]	Rows from field margin Psyllids/10 bushes	1 10	3 5	5 4	10 3	20 1	64
102	<i>Tyloclerma fragariae</i> (strawberry crown borer) [crawling]	Yards from release point Borers recovered	25 6	50 6	68 5	125 4	300 1	76
Vertical Dispersion								
103	<i>Anopheles</i> spp. (mosquito) [flight]	Altitude, meters (in forest) Distribution, % Distribution, % Distribution, %	0 51 3 4	5 18 3 7.3	10 19 39 24.6	15 12 55 64.1		19

continued

## 114. DISPERSION OF SMALL ORGANISMS

### Part I. INVERTEBRATES

	Invertebrate (Common Name) [ Means of Dispersion ]	Distances and Units Dispersed						Refer- ence
		(A)	(B)	(C)	(D)	(E)	(F)	
Vertical Dispersion								
104	<i>Anopheles quadrimaculatus</i> (malaria mosquito) [flight]	Altitude, feet	1.5	3.0	6.0	7.5		28
		Mosquitoes caught	14	11	5	3		
105	Aphididae (aphid) [flight]	Altitude, feet	50	250	500	1,000	1,500	44
		Aphids caught	270	228	75	21	6	
106	Aphididae [air currents, flight]	Altitude, miles	0.17	1.15	2.17	2.68		68
		Aphids caught	2	9	16	19		
107	Aphididae, 7-hr collection [flight]	Altitude, feet	10	150	500	1,000		45
		Aphids caught/hr	18	3	2	1		
108	<i>Circulifer tenellus</i> (beet leafhopper) [flight]	Altitude, feet	8	16	25			21
		Leafhoppers caught	276	211	169			
109		Altitude, feet	2.5	15	32			52
		Parasitism, %	3.5	1.7	0.9			
110	Coccinellidae (ladybird beetle) [flight]	Altitude, miles	0.17	1.15	2.17	2.68		68
		Beetles caught	40	13	4	1		
111	<i>Drosophila</i> sp. (fruit fly) [flight]	Altitude, miles	0.54	2.04	3.54	5.04		68
		Flies caught	413	237	164	117		
112	<i>Epitrix hirtipennis</i> (tobacco flea beetle) [flight]	Altitude, feet	12	19	23			1
		Beetles caught	96	41	18			
113		Altitude, feet	1.5	5	14	19	23	22
		Beetles caught/day	56	32	12	6	2	
114	<i>Heterodera rostochiensis</i> (golden nematode of potato) [wind currents]	Height of trap, inches	22	38	55			92
		Viable cysts	2,077	435	103			
115	Insects, miscellaneous [flight, air currents]	Altitude, feet	10	177	277			29
		Insects caught/cu ft air	239	51	21			
116		Altitude, feet	20	1,000	4,000	7,000		35
		Insects caught/10 min flight	26	6	3	1		
117	<i>Psallus seriatus</i> (cotton fleahopper) [flight, air currents]	Altitude, feet	5.5	11.5	17.5	23.5		22
		Fleahoppers caught	61	97	156	250		
118	<i>Pyrausta nubilalis</i> (European corn borer) [flight]	Altitude, feet	5	10	15			33
		Moths caught	914	547	332			

Contributor: Wolfenbarger, D. O.

References: [1] Annand, P. N. 1932. U.S. Dept. Agr. Circ. 244. [2] Baker, W. A., et al. 1949. U.S. Dept. Agr. Tech. Bull. 983. [3] Bedard, W. D. Unpublished. U.S. Dept. of Agriculture, Bureau of Entomology and Plant Quarantine, Washington, D. C., 1939. [4] Bissell, T. L. 1939. J. Econ. Entomol. 32:546. [5] Bonnet, D. D., and D. J. Worcester. 1946. Am. J. Trop. Med. 26:465. [6] Braun, E., et al. 1953. Sci. Agr. 33:437. [7] Brower, A. E. 1930. Entomol. News 41:10,44. [8] Brownlee, J. 1911. Proc. Roy. Soc. Edinburgh, B, 31:262. [9] Burgess, A. F. 1913. U.S. Dept. Agr. Bur. Entomol. Bull. 119. [10] Carruth, L. A., and T. W. Kerr. 1937. J. Econ. Entomol. 30:297. [11] Cartwright, O. L. 1929. S. Carolina Agr. Expt. Sta. Bull. 257. [12] Causey, O. R., and H. W. Kumm. 1948. Am. J. Trop. Med. 28:469. [13] Chamberlain, T. R. 1941. U.S. Dept. Agr. Tech. Bull. 784. [14] Cockerham, K. L., et al. 1954. Louisiana Agr. Expt. Sta. Tech. Bull. 483. [15] Collins, C. W., and S. F. Potts. 1932. U.S. Dept. Agr. Tech. Bull. 336. [16] Crossman, S. S. 1917. J. Econ. Entomol. 10:177. [17] Dalmat, H. T., and C. L. Gibson. 1952. Ann. Entomol. Soc. Am. 45:605. [18] Davidson, W. M. 1925. Trans. Am. Entomol. Soc. 50:163. [19] Deane, L. M., et al. 1953. Folia Clin. Biol. (Sao Paulo) 20:101. [20] De Meillon, B. 1934. Publ. S. African Inst. Med. Res. 6:195. [21] Dobzhansky, T., and S. Wright. 1943. Genetics 28:304. [22] Dominick, C. B. 1943. Virginia Agr. Expt. Sta. Bull. 355. [23] Doucette, C. F. 1942. U.S. Dept. Agr. Tech. Bull. 809. [24] Douglas, W. A. 1941. U.S. Dept. Agr. Circ. 602. [25] Dudley, J. E., and E. M. Searls. 1923. J. Econ. Entomol. 16:363. [26] Echert, J. E. 1933. J. Agr. Res. 47:257. [27] Eyles, D. E., et al. 1945. U.S. Public Health Rept. 60:1265. [28] Ficht, G. A., and T. E. Heinton. 1941. J. Econ. Entomol. 34:599. [29] Freeman, J. A. 1938. Nature 142:153. [30] Frost, S. W. 1933. Penna. State Univ. Agr. Expt. Sta. Bull. 301. [31] Fulton, R. A., and V. E. Romney. 1940. J. Agr. Res. 61:737. [32] Gaines, J. C. 1932. J. Econ. Entomol. 25:1181.

continued



114. DISPERSION OF SMALL ORGANISMS

Part I. INVERTEBRATES

- [33] Gaines, J. C., and K. P. Ewing. 1938. *Ibid.* 31:674. [34] Gardner, T. R. 1938. *Ibid.* 31:204. [35] Glück, P. A. 1939. U.S. Dept. Agr. Tech. Bull. 673. [36] Graham, S. A. 1922. Rept. Minn. State Entomologist 19:15. [37] Hawley, I. M. 1934. *J. Econ. Entomol.* 27:503. [38] Hervey, G. E. R., and C. E. Palm. 1935. *Ibid.* 28:670. [39] Hill, C. B. 1952. Virginia Agr. Expt. Sta. Tech. Bull. 119. [40] Horsfall, W. R. 1942. Arkansas Univ. (Fayetteville) Agr. Expt. Sta. Bull. 427. [41] Jachowski, L. A., Jr. 1954. *Am. J. Hyg.* 60:186. [42] Jackson, C. H. N. 1940. *Ann. Eugenics* 10:332. [43] Jenkins, D. W., and C. C. Hassett. 1951. *Can. J. Zool.* 29:178. [44] Johnson, C. G. 1951. *Sci. Progr. (London)* 39(153):41. [45] Johnson, C. G., and H. L. Penman. 1951. *Nature* 168:337. [46] Jones, E. T. Unpublished. U.S. Dept. of Agriculture, Bureau of Entomology and Plant Quarantine, 1937. [47] Kettle, D. S. 1951. *Bull. Entomol. Res.* 42:239. [48] Kettle, D. S. 1951. *Proc. Roy. Entomol. Soc. London, A*, 26:59. [49] Klostermeyer, C. E. 1953. Wash. State Univ. Agr. Expt. Sta. Tech. Bull. 9. [50] Larrimer, W. H., and A. L. Ford. 1919. *J. Econ. Entomol.* 12:417. [51] Larson, A. O., et al. 1933. *Ibid.* 26:1063. [52] Lawson, F. R., et al. 1951. U.S. Dept. Agr. Tech. Bull. 1030. [53] Lindquist, A. W., et al. 1951. *J. Econ. Entomol.* 44:397. [54] Linn, M. B. 1940. Cornell Univ. Agr. Expt. Sta. Bull. 742. [55] MacCreary, D., and P. L. Rice. 1949. *Ann. Entomol. Soc. Am.* 42:141. [56] McCullough, J. W. 1917. *J. Econ. Entomol.* 10:162. [57] MacVicar, R. M., et al. 1952. *Sci. Agr.* 32:67. [58] Moggridge, J. Y. 1949. *Bull. Entomol. Res.* 40:43. [59] Morris, K. R. S. 1952. *Ibid.* 43:375. [60] Munro, J. A., and H. S. Telford. 1942. N. Dakota Agr. Expt. Sta. Bull. 309. [61] Neiswander, C. R., and J. R. Savage. 1931. *J. Econ. Entomol.* 23:389. [62] Nicholas, W. L. 1953. *Am. Trop. Med. Parasitol.* 47:309. [63] Ohlendorf, W. 1926. U.S. Dept. Agr. Bull. 1374. [64] Peterson, A. 1923. New Jersey Agr. Expt. Sta. Bull. 378. [65] Petherbridge, F. R., and D. W. Wright. 1943. *Ann. Appl. Biol.* 30:348. [66] Phipps, C. R., and C. O. Dirks. 1932. *J. Econ. Entomol.* 25:576. [67] Pimental, D., and R. W. Fay. 1955. *Ibid.* 48:19. [68] Profft, J. 1939. *Arb. Physiol. Angew. Entomol. Berlin-Dahlem* 6:119. [69] Quarterman, K. D., et al. 1949. *Am. J. Med.* 29:973. [70] Quarterman, K. D., et al. 1954. *J. Econ. Entomol.* 47:405. [71] Quarterman, K. D., et al. 1954. *Ibid.* 47:413. [72] Quarterman, K. D., et al. 1955. *Ibid.* 48:30. [73] Quayle, H. J. 1911. *Ibid.* 4:301. [74] Reeves, W. C., et al. 1948. *Mosquito News* 8:61. [75] Reinhard, H. J., and F. L. Thomas. 1933. Texas Agr. Expt. Sta. Bull. 475. [76] Richter, P. O. 1939. Kentucky Agr. Expt. Sta. Bull. 389. [77] Rickard, E. R. 1928. *Inst. Clin. Quirugica (Buenos Aires) Bol.* 4:133. [78] Riegert, P. W., et al. 1954. *Can. Entomologist* 86:223. [79] Roebuck, A., et al. 1947. *Ann. Appl. Biol.* 34:186. [80] Schoof, H. F., and R. R. Siverly. 1954. *J. Econ. Entomol.* 47:830. [81] Smith, G. E., R. B. Watson, and R. L. Crowell. 1941. *Am. J. Hyg.* 34(C):102. [82] Stearns, L. A., and D. MacCreary. 1938. *J. Econ. Entomol.* 31:226. [83] Steiner, H. M., and H. N. Worthley. 1941. *Ibid.* 34:249. [84] Swynnerton, C. F. M. 1936. *Trans. Roy. Entomol. Soc. London* 84:1. [85] Timofeeff-Ressovsky, N. V., and E. A. Timofeeff-Ressovsky. 1940. *Z. Induktive Abstammungs- Vererbungslehre* 79:44. [86] Travis, B. V., et al. 1940. *J. Econ. Entomol.* 33:847. [87] Van Leeuwen, E. R. 1940. *Ibid.* 33:162. [88] Van Zwaluwenburg, R. H., and J. S. Rosa. 1940. Hawaiian Planters' Record 44:3. [89] Wadley, F. M., and D. O. Wolfenbarger. 1944. *J. Agr. Res.* 69:299. [90] Wakeland, C. R. 1934. *J. Econ. Entomol.* 28:981. [91] Walcott, A. M., et al. 1937. *Am. J. Trop. Med.* 17:677. [92] White, J. H. 1953. *Nature* 172:686. [93] Willis, H. R. 1939. *J. Econ. Entomol.* 32:401. [94] Wolfenbarger, D. O. 1940. *Ann. Entomol. Soc. Am.* 33:391. [95] Wolfenbarger, D. O. 1948. *Florida Entomologist* 31:15. [96] Wolfenbarger, D. O. Unpublished. U.S. Dept. of Agriculture, Bureau of Entomology and Plant Quarantine, Washington, D. C., 1941. [97] Wolfensohn, M. 1953. *Bull. Res. Council Israel, B*, 3:263. [98] Wolfensohn, M., and R. Galun. 1953. *Ibid.*, B, 2:433. [99] Woodside, A. M. 1954. *J. Econ. Entomol.* 47:349. [100] Wright, D. W., and D. G. Ashby. 1946. *Ann. Appl. Biol.* 33:69.

*continued*



## 114. DISPERSION OF SMALL ORGANISMS

### Part II. VIRUSES, BACTERIA, AND FUNGI

Disease (Organism) [ Means of Dispersion ]		Distances and Units Dispersed						Ref- erence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
<b>Viruses</b>								
1	Beet mosaic [black bean aphid]	Yards from steckling bed	220	1,320	1,560			31
		Plants infected, %	95	57	12			
2	Cabbage mosaic [cabbage aphid]	Miles from fields	0.06	0.6	7	22		30
		Plants infected, %	72.6	46.9	1.0	0.3		
3	Celery mosaic [insects]	Feet from harbored plants	3	15	28	75	120	43
		Diseased plants, %	100	89	52	28	16	
4	Cucumber cucurbit mosaic [insects]	Yards from harbored plant	1	140	225	350	500	7
		Days to first symptoms	17	42	45	47	49	
5	Eastern X-disease of peach [certain leafhoppers]	Feet from chokecherry plants	50	125	187.5			40
		Plants infected, %	89.3	8.1	0.6			
6	Mild streak of black raspberry [presumably insects]	Feet from wild brambles	25	75	125			16
		Infections, %	66	29	12			
7	Potato calico [insects]	Rows from source	1	2	3	4	5	29
		Diseased plants	26	27	22	17	13	
8	Potato leaf roll [insects]	Inches from inoculum source	18	36	54	72	90	11
		Plants infected	41	25	17	12	8	
9	Potato leaf roll [aphids]	Rows from infected plants	1	2	3	4	6	22
		Diseased plants, %	21	12	7	5	1	
10	Potato mosaic [insects]	Rows from diseased plants	1	2	3	4	6	21
		Diseased plants, %	36	24	18	13	6	
11	Potato yellow dwarf [six-spotted leafhopper]	Feet from old meadow	1	30	60	90	135	9
		Diseased plants, %	80	23	14	9	4	
12	Rugose mosaic of potato [insects]	Inches from inoculum source	18	36	54	72	90	11
		Infected plants	37.7	11.7	5.3	3.4	2.8	
13	Severe streak of raspberry [insects]	Rows from wild brambles	3	8	13	18	23	5
		Diseased plants	165	86	47	21	1	
14	Sudden death of clove [probably insects]	Tree intervals	1	2	3	4	5	24
		Life expectancy, months	30.2	38.1	42.7	46.0	48.5	
15	Sugar beet curly top [beet leafhopper]	Miles from breeding ground	57	265	315	385	430	33
		Diseased plants, %	100	15	10	4	1	
16	Tristeza disease of citrus [aphids]	Feet from inoculum source	5	25	45	65	73	1
		Diseased trees, %	51	29	20	15	13	
17	Wheat streak mosaic [eriophyid mite]	Yards from source	0	8	18	50	90	34
		Plants infected, %	64	26	19	10	5	
<b>Bacteria</b>								
18	(Colonies on sea-water medium) [air currents]	Miles from land (over water)	5	80	275			46
		Bacterial colonies	41	58	65			
19		Miles from sea (over land)	0	0.06	0.25	0.50	1.0	46
		Bacterial colonies	548	292	215	177	138	
<b>Fungi</b>								
20	(Air-borne spores) [wind]	Degrees north of equator	57°30'	64°20'	68°55'	71°05'		27
		Fungus colonies on plate	3.61	0.49	0.48	0.72		
21	Beet downy mildew ( <i>Peronospora</i> sp.) [wind]	Meters from seed plants	10	150	1,000			13
		Plants injured, %	28	8	1			
22	Blossom infection ( <i>Sclerotinia taxa</i> ) [air currents]	Feet from center of nearest source row	22	44	66	88		44
		Blossom infection, %	55.7	39.1	29.3	22.4		
23	( <i>Bovista plumbea</i> ) [air currents]	Meters from release point	5	10	15	20		39
		Spores caught	912	323	165	102		
24	Cedar and apple rust ( <i>Gymnosporangium</i> sp.) [air currents]	Yards from infected trees	0	55	110	220	440	18
		Leaf infections	64	40	33	26	19	
25	Chestnut blight ( <i>Endothia parasitica</i> ) [air currents]	Feet from spore source	27	85	180	266		12
		Ascospores found	23	11	8	8		
26	Crown rust of oats ( <i>Puccinia coronata</i> ) [wind]	Feet from inoculum source	3	5	7.7	10.3	13	45
		Infections, %	92.9	53.4	35	19.5	0.7	
27	Damping-off disease [mycelial growth]	Centimeters from inoculum source	1	3	5	7	9	2
		Plants damped-off, %	100	8	7	16	0	
28	Downy mildew ( <i>Pseudoperonospora humilis</i> ) [air currents]	Feet from spore source	10	50	100	200	400	20
		Leaves infected, %	26	16	12	7	3	

continued

## 114. DISPERSION OF SMALL ORGANISMS

### Part II. VIRUSES, BACTERIA, AND FUNGI

Disease (Organism) [Means of Dispersion]		Distances and Units Dispersed					Ref- er- ence	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
Fungi								
29	Dutch elm disease ( <i>Ceratostomella ulmi</i> ) [elm bark beetles]	Feet from inoculum source	200	600	1,000	1,800	2,300	19
		Diseased trees/acre	2.52	0.38	0.47	0	0.09	
30		Feet from inoculum source	12.5	62.5	300	575		4
		Diseased trees	27	19	11	8		
31	Eyespot disease of wheat ( <i>Helminthosporium sacchari</i> ) [water]	Meters from spore source	5	20	50	70		25
		Culms with eye spots, %	26.7	12.3	7.7	6.4		
32	Hollyhock rust [gravity and ejection]	Spore dispersal, millimeters	0.15	0.35	0.55	0.75		3
		Spores/0.1 sq mm	14.6	7.6	2.2	0.1		
33	Leaf spots on tulips [raindrop splash and wind]	Centimeters from conidia source	15.2	34.6	58.0	79.8	102.0	42
		Lesions/plant	31.6	20.1	12.9	8.5	5.1	
34	Loose smut of wheat ( <i>Ustilago tritici</i> ) [air currents]	Meters from spore source	2	4	24	80		26
		Smutted heads	241	234	114	0		
35	Maize rust ( <i>Puccinia sorghi</i> ) [wind]	Kilometers from spore source	0.5	2.5	4.5	6.5		47
		Plants attacked, %	100	3	0.3	0		
36	Onion mildew ( <i>Peronospora destructor</i> ) [air currents]	Feet from onion sets	120	780	1,750	2,000		23
		Lesions/100 ft row	1,138	98	1	0		
37	Potato late blight ( <i>Phytophthora infestans</i> ) [wind]	Centimeters from edge of infective group	30	90	150	210	270	10
		Plants infected, %	89	63	43	22	5	
38	Powdery mildew on barley ( <i>Erysiphe graminis</i> ) [wind]	Meters from source	1.5	3.5	5.5	7.5	8.5	28
		Plants affected, %	99	84	76	70	68	
39	Stem rust ( <i>Puccinia graminis</i> ) [wind]	Feet from barberry hedge	15	125	225	325	425	17
		Grass infected, %	100	41	5	1	0.5	
40	Stem rust on rye ( <i>P. graminis secalis</i> ) [wind]	Meters from source plant	50	300	1,000	3,000		8
		Yield/100 ears, grams	47.6	92.3	122.3	149.7		
41	( <i>Tilletia tritici</i> ) [air currents]	Meters from release point	5	10	15	20		39
		Spores caught	800	168	49	30		
42	Tobacco blue mold ( <i>Peronospora tabacina</i> ) [wind]	Yards from source	0	4	8	12		41
		Plant lesions/1,000 sq in. of field	140	8	1	0.5		
43	Wheat stem rust ( <i>Puccinia graminis</i> ) [air currents]	Miles from known source	200	360	580	740	940	37
		Spores collected	13,092	10,768	8,883	7,920	6,975	
44	White pine blister rust ( <i>Cronartium ribicola</i> ) [air currents]	Feet from gooseberry bush	50	150	350	450	650	36
		Diseased trees, %	75	55	40	36	29	
Vertical Dispersion								
45	(Bacteria, miscellaneous) [air currents]	Altitude, feet	1,500	6,000	12,000	15,000		32
		Bacteria	113	48	15	5		
Fungi								
46	Azalea flower spot ( <i>Oculinia azalaeae</i> ) [air currents, water drip]	Inches above ground	4	10	18	48		35
		Infections	42	28	17	0		
47	Onion mildew ( <i>Peronospora destructor</i> ) [air currents]	Altitude, feet	100	200	700	1,200		23
		Spores/cu ft air	32	102	451	801		
48	Wheat stem rust ( <i>Puccinia graminis</i> ) [air currents]	Feet above barberry bushes	1,000	2,000	7,000	12,000		38
		Aeciospores caught	19	14	5	1		
49		Altitude, feet	1,000	5,000	10,000	14,000		6
		Uredospores	48,200	7,730	144	40		
50		Elevation, meters	30	400	600	800		15
		Spores/sq cm/min	1,458	490	339	231		

Contributor: Wolfenbarger, D. O.

References: [1] Bitancourt, A. A., and A. J. Rodriguez. 1948. Arquiv. Inst. Biol. (Sao Paulo) 18:313. [2] Blair, I. D. 1943. Ann. Appl. Biol. 30:118. [3] Buller, A. H. 1924. Researches on fungi. Longmans, Green; London. v. 3, p. 533. [4] Collins, D. L., et al. 1940. Cornell Univ. Agr. Expt. Sta. Bull. 740. [5] Cooley, L. M. 1936. N. Y. State Agr. Expt. Sta. (Geneva) Bull. 665. [6] Craigie, J. H. 1945. Sci. Agr. 25:285. [7] Doolittle, S. P.

continued

## 114. DISPERSION OF SMALL ORGANISMS

### Part II. VIRUSES, BACTERIA, AND FUNGI

1925. J. Agr. Res. 31:1. [8] Fisher, H. 1950. Z. Pflanzenkrankh. Pflanzenschutz 57:1. [9] Frampton, V. L. 1942. Phytopathology 32:799. [10] Gregory, P. H. 1945. Brit. Mycol. Soc. Trans. 28:26. [11] Gregory, P. H., and D. R. Read. 1949. Ann. Appl. Biol. 36:475. [12] Heald, F. D., et al. 1915. J. Agr. Res. 3:493. [13] Höchapel, H. 1950. Nachrbl. Deut. Pflanzenschutzdienst (Braunschweig) 2:124. [14] Hodgson, H. J. 1949. Agron. J. 41:337. [15] Hubert, K. 1932. Fortschr. Landwirtsch. 7:195. [16] Jeffers, W. F., and W. W. Woods. 1948. Phytopathology 38:222. [17] Johnson, A. G., and J. G. Dickson. 1919. Wisconsin Univ. Agr. Expt. Sta. Bull. 304. [18] Jones, L. R., and E. T. Bartholomew. 1915. Ibid. 257. [19] Liming, O. N., et al. 1951. Phytopathology 41:146. [20] Magie, R. O. 1942. N. Y. State Agr. Expt. Sta. (Geneva) Tech. Bull. 267. [21] Murphy, P. A. 1921. Can. Exptl. Farms Bull. 44. [22] Murphy, P. A., and E. J. Worthley. 1920. Phytopathology 10:407. [23] Newhall, A. G. 1938. Ibid. 28:257. [24] Nutman, F. J., and F. M. L. Sheffield. 1949. Ann. Appl. Biol. 36:419. [25] Oort, A. J. P. 1936. Tijdschr. Plantenziekten 42:179. [26] Oort, A. J. P. 1940. Ibid. 46:1. [27] Pady, S. M., et al. 1950. Phytopathology 40:632. [28] Pape, H., and B. Rademacher. 1934. Angew. Botan. 16:115. [29] Porter, D. R. 1935. Hilgardia 9:383. [30] Pound, G. S. 1946. Phytopathology 36:1035. [31] Pound, G. S. 1947. J. Agr. Res. 75:31. [32] Proctor, B. E. 1934. Proc. Am. Acad. Arts Sci. 69:315. [33] Romney, V. E. 1939. U.S. Dept. Agr. Circ. 518. [34] Slykhuis, J. T. 1955. Phytopathology 45:116. [35] Smith, F. F., and F. Weiss. 1942. U.S. Dept. Agr. Tech. Bull. 798. [36] Snell, W. H. 1941. J. Forestry 39:537. [37] Stakman, E. C., and L. M. Hamilton. 1939. Plant Disease Repr., Suppl., 117:69. [38] Stakman, E. C., et al. 1923. J. Agr. Res. 24:599. [39] Stepanov, K. M. 1935. Tr. Zashchite Rastenii, II, 8:1. [40] Stoddard, E. M. 1947. Conn. Agr. Expt. Sta. New Haven Bull. 506. [41] Waggoner, P. E., and G. S. Taylor. 1955. Plant Disease Repr. 39:79. [42] Wallace, E. R. 1934. Holland County (Engl.) Council Bulb Res. Subcommittee Rept., 1933, p. 37. [43] Wellman, F. L. 1935. Phytopathology 25:289. [44] Wilson, E. E., and G. A. Baker. 1946. J. Agr. Res. 72:301. [45] Wilson, E. E., and G. A. Baker. 1946. Phytopathology 36:418. [46] Zobel, C. E. 1942. Publ. Am. Assoc. Advan. Sci. 17:55. [47] Zogg, H. 1949. Phytopathol. Z. 15:143.

### Part III. POLLEN AND SEEDS

	Spermatophyte (Common Name) [Means of Dispersion]	Distances and Units Dispersed						Reference	
		(A)	(B)	(C)	(D)	(E)	(F)		(G)
Horizontal Dispersion									
1	<i>Abies alba</i> (silver fir) [air currents]	Yards from seed trees	55	165	275				19
		Seedlings/acre	22	9	3				
2	<i>Agropyron cristatum</i> (crested wheat-grass) [wind]	Rods from field	5	15	25				23
		Pollen grains	72	29	10				
3	<i>A. intermedium</i> (intermediate wheat-grass) [wind]	Rods from field	5	12	25				23
		Pollen grains	44	17	4				
4	<i>Beta</i> sp. (beet) [wind]	Meters from seed fields	0	300	500	800			21
		Pollen grains/sq cm	11,613	1,941	1,075	278			
5		Feet from contaminant	2.3	20.7	43.2	73.2			5
		Hybrids, %	5.6	0.3	0.2	0			
6	<i>Brassica rapa</i> (turnip) [insects]	Feet from contaminating plants	1	4.5	15	30	42.5		6
		Proportion of hybrid seed, %	42	14	2.5	0.2	0.5		
7	<i>Bromus</i> sp. (brome-grass) [wind]	Rods from field	5	15	25	40	60		23
		Pollen grains	146	41	21	10	4		
8	<i>Cedrus atlantica</i> (atlas cedar) [wind]	Feet from source tree	40	120	240	325	700		37
		Pollen grains	189	116	71	51	0.1		
9	<i>C. libani</i> (cedar of Lebanon) [wind]	Feet from source tree	15	75	135	195			37
		Pollen grains	127	62	37	22			
10	<i>Citrullus vulgaris</i> (watermelon) [honey-bees]	Feet from field margin	50	250	450	650			28
		Melons/acre	734	653	623	605			

continued



## 114. DISPERSION OF SMALL ORGANISMS

### Part III. POLLEN AND SEEDS

Spermatophyte (Common Name) [Means of Dispersion]		Distances and Units Dispersed						Refer- ence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
11	<i>Dactylis</i> sp. (orchard grass) [wind]	Meters from field	0	200	400	600	800	21
		Pollen grains/sq cm	3,096	447	172	120	86	
12	<i>Fraxinus</i> sp. (ash) [wind]	Feet from source tree	25	50	150	400		37
		Pollen grains	2,545	1,008	141	29		
13	<i>Gilia</i> sp. (gilia) [insects]	Feet from white flowers	25	250	2,640	10,560		12
		Hybridization of blue-flowered plants, frequency	100	74	44	26		
14	<i>Gossypium</i> sp. (cotton) [wind]	Feet from marker plants	7.5	20.0	32.5	45.0	70.0	1
		Natural crossing, %	29.7	8.5	9.1	5.1	0.8	
15		Feet from red cotton	16	35	51	99	189	15
		Hybrids, %	6.9	3.0	2.0	0.9	0.3	
16	<i>G. arboreum</i> (Asiatic tree cotton) [bumblebees]	Feet from flowers dusted with methylene blue	1.5	4.5	7.5	10.5	13.5	33
		Flowers with dye particles, %	60	49	44	40	38	
17	<i>G. hirsutum</i> (upland cotton) [bumblebees]	Feet from flowers dusted with methylene blue	1.5	4.5	7.5	10.5	13.5	33
		Flowers with dye particles, %	94	85	81	78	76	
18	<i>G. hirsutum</i> [wind or insects]	Feet from contaminant	1	10	100	700	1,800	29
		Cross-pollination, %	18	12	7	3	1	
19	<i>Helianthus</i> sp. (sunflower) [honeybees]	Feet from apiary	0	200	600	1,000		13
		Seed yield, lbs/acre	1,285	981	918	889		
20	<i>Juglans regia</i> (Persian walnut) [air currents]	Feet from pollen source	60	150	500	1,000	1,600	9
		Pollen grains/sq mm/24 hr	4	2.8	1.4	0.6	0	
21	<i>Juniperus scopularum</i> (western red cedar) [air currents]	Yards from seed source	22	44	66	88		19
		Seedlings/acre	5,588	259	192	0		
22	<i>Leontodon</i> sp. (hawkbit) [wind]	Feet from source	2	12	20	28	32	7
		Seeds	184	21	9	3.5	1	
23	<i>Lolium</i> sp. (ryegrass) [wind]	Meters from ryegrass field	0	200	500	700	900	21
		Pollen grains/sq cm	4,045	1,053	535	345	204	
24	<i>L. perenne</i> (perennial ryegrass) [wind]	Centimeters from rough clone contaminant	40	120	200	280		34
		Rough plants, %	40.1	13.8	7.2	3.8		
25	<i>Lycopersicon esculentum</i> (tomato) [air currents; insects?]	Feet from contaminant	6	18	30	42	54	11
		Cross-pollination, %	1.1	0.6	0.4	0.2	0.1	
26	<i>Malus pumila</i> (common apple) [wind]	Feet from source tree	0	165	330			37
		Pollen grains	13	2	0.9			
27		Feet from pollen source	8	19	42			30
		Fruit set/100 blossom spurs	52	34	18			
28	<i>M. pumila</i> [honeybees]	Yards from bee colonies	25	50	75	100	150	20
		Fruit set, %	7	7	6	6	4	
29	<i>Oryza sativa</i> (rice) [dehiscence and wind]	Centimeters from pollen source	25	50	100	150	200	31
		Pollen grains	22	9	3	1	0.4	
30	<i>Panicum virgatum</i> (switch grass) [wind]	Rods from field	5	15	25	40	60	23
		Switchgrass pollen	27	7	4	2	0.5	
31	<i>Parthenium argentatum</i> (guayule parthenium) [wind]	Yards from guayule plants	100	400	850	1,200		14
		Pollen grains/sq in.	89	49	27	17		
32	<i>Paspalum notatum</i> (Pensacola Bahia grass) [wind]	Rods from albino seedling isolation blocks	0	5	10	15		18
		Albinos, %	14.0	19.3	21.7	23.0		
33	<i>Pennisetum glaucum</i> (pearl millet) [wind]	Yards from release point	4	50	200	400		18
		Pollen, %	100.0	8.9	0.8	0.4		
34	<i>Persea</i> sp. (persea) [honeybees]	Feet from apiary	125	562.5	1,062.5			36
		Fruit yield/tree, bushels	2.38	1.26	0.94			
35	<i>P. americana</i> , Taylor variety (American avocado) [insects]	Rows from nearest reciprocating variety	1	2	3	4	5	35
		Fruit set/tree	61	54	50	47	44	
36	<i>Phaseolus limensis</i> (lima bean) [insects]	Feet from pollen parent	2.5	5.0	7.5	12.5	32.5	2
		Natural hybrids, %	7.3	2.7	1.4	0.6	0.5	
37	<i>P. lunatus</i> (sieva bean) [air currents; insects?]	Yards from kidney beans	1	2	3	5	9	4
		Cross-pollination, %	5	4	3	2	1	
38	<i>P. vulgaris</i> (kidney bean) [air currents; insects?]	Yards from sieva beans	1	2	3	5	9	4
		Cross-pollination, %	9	7	6	4	3	

continued



## 114. DISPERSION OF SMALL ORGANISMS

### Part III. POLLEN AND SEEDS

Spermatophyte (Common Name) [Means of Dispersion]		Distances and Units Dispersed						Ref- er- ence
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
Horizontal Dispersion								
39	<i>Phleum pratense</i> (timothy) [wind]	Meters from timothy field	0	100	200	300	500	21
		Pollen grains/sq cm	2,613	781	505	343	140	
40	<i>Picea</i> sp. (spruce) [wind]	Feet from source tree	0	165	330			37
		Pollen grains	9.7	0.1	0.7			
41	<i>P. mariana</i> (black spruce) [air currents]	Feet from seed trees	10	80	160	240		3
		Seedlings/acre	71,260	47,180	18,540	0		
42	<i>Pinus</i> spp. (pine) [air currents]	Yards from seed tree stand	3,344	6,248	8,426			24
		Seedlings/acre	2,002	991	507			
43	<i>P. cembroides</i> (Mexican piñon pine) [wind]	Feet from source tree	10	75	150	225	300	37
		Pollen grains	8,479	462	86	38	52	
44	<i>P. echinata</i> (shortleaf pine)	Miles from forest margin	0	0.1	0.2	0.25		8
		Pollen, %	100	17	15	10		
45	<i>P. monticola</i> (western white pine) [air currents]	Yards from seed source	22	44	66	88		19
		Seedlings/acre	616	177	57	9		
46	<i>Populus</i> sp. (poplar) [wind]	Feet from source tree	50	500	1,400	3,200	4,200	37
		Pollen grains	107	86	76	69	66	
47	<i>P. deltoides</i> (eastern poplar) [wind]	Feet from source tree	25	250	500	1,550	3,550	37
		Pollen grains	115	62	46	20	0.3	
48	<i>Pseudotsuga taxifolia</i> (Douglas fir) [air currents]	Feet from seed trees	2	4	6	8		19
		Seedlings/acre	304	170	91	35		
49	<i>Raphanus</i> sp. (radish) [insects]	Feet from contaminant	1	3	4.2	5.0	6.4	6
		Contamination, %	11.9	5.3	1.7	1.6	0.7	
50	<i>R. sativus</i> (garden radish) [wind or insects]	Feet from contaminant	1	95	191	335	420	10
		Cross-pollination, %	75	18	10	3	0	
51	<i>Secale cereale</i> (rye) [wind]	Rods from rye field	5	15	25	40	60	23
		Pollen grains	453	232	124	52	11	
52		Meters from rye field	100	300	500	700		21
		Pollen grains/sq cm	4,181	2,579	1,834	1,343		
53	<i>S. cereale</i> [air currents]	Feet from pollen source	0.3	2.6	5.2	7.9	10.5	32
		Cross-pollination, %	24	18	13	9	7	
54	<i>Trifolium hybridum</i> (alsike clover) [honeybee]	Yards from bee colonies	5.5	440	880			26
		Seeds/head	38	33	32			
55		Miles from bee yards	0.125	0.625	1.5	2.5		16
		Seed yield, lbs/acre	128	95	77	67		
56	<i>T. pratense</i> (red clover) [honeybees]	Yards from apiary	200	500	833	1,126	1,426	25
		Seed yield, lbs/acre	166	149	139	133	128	
57	<i>T. repens</i> (white clover) [honeybees]	Miles from bee yards	0.125	0.625	1.5	2.5		16
		Seed yield, lbs/acre	206	102	46	13		
58	<i>Tsuga heterophylla</i> (western hemlock) [air currents]	Yards from seed trees	22	44	66	88		19
		Seedlings/acre	1,434	169	101	0		
59	<i>Ulmus</i> sp. (elm) [wind]	Feet from source tree	500	1,100	2,700	5,500		37
		Pollen grains	115	152	12	8		
60	<i>Zea mays</i> (corn) [wind]	Rods from field	5	15	25	40	60	23
		Pollen grains	18	6	3	2	0.8	
61		Feet from pollen source	10	30	50	70		5
		Pollen grains	7,330	341	121	30		
62		Feet from pollen source	4	16	28	40	44	17
		Seed set	256	197	122	75	31	
63		Feet from contaminating plants	13	29	45	61	77	5
		Hybridization, %	7	6	3	1.3	0.3	
64		Rods north of contaminating field	5	25	60	100		22
		Outcrossed seeds, %	16.5	0.8	0.2	0.2		
Vertical Dispersion								
65	<i>Beta vulgaris</i> (beet) [air currents]	Altitude, feet	1,000	2,000	3,000	4,000		27
		Pollen grains	56	26	14	9		

Contributor: Wolfenbarger, D. O.

References: [1] Afzal, M., and A. H. Khan. 1950. Agron. J. 42:89. [2] Allard, R. W. 1954. Proc. Am. Soc.

continued

**114. DISPERSION OF SMALL ORGANISMS**

**Part III. POLLEN AND SEEDS**

Hort. Sci. 64:410. [3] Anonymous. 1939. U.S. Dept. Agr. Forest Serv. Lake States Forest Expt. Sta. Tech. Note 147. [4] Barrons, K. 1938. Proc. Am. Soc. Hort. Sci. 36:637. [5] Bateman, A. J. 1947. Heredity 1:235. [6] Bateman, A. J. 1947. J. Genet. 48:257. [7] Brownlee, J. 1911. Proc. Roy. Soc. Edinburgh, B, 31:262. [8] Buell, M. F. 1947. J. Elisha Mitchell Sci. Soc. 63:163. [9] Crane, H. L., et al. 1938. Better plants and animals. Yearbook of agriculture. U.S. Govt. Printing Office, Washington, D. C. [10] Crane, M. B., and K. Mather. 1943. Ann. Appl. Biol. 30:301. [11] Currence, T. M., and J. M. Jenkins, Jr. 1942. Proc. Am. Soc. Hort. Sci. 41:273. [12] Epling, C., and T. Dobzhansky. 1942. Genetics 27:317. [13] Furgala, B. 1954. Gleanings Bee Cult. 82:532. [14] Gardner, E. J. 1946. J. Am. Soc. Agron. 38:264. [15] Green, J. M., and M. D. Jones. 1953. Agron. J. 45:366. [16] Harrison, C. M., et al. 1945. Mich. State Univ. Agr. Expt. Sta. Quart. Bull. 28:85. [17] Haskell, G., and P. Dow. 1951. Empire J. Exptl. Agr. 19:45. [18] Hodgson, H. J. 1949. Agron. J. 41:337. [19] Hoffmann, J. V. 1911. J. Agr. Res. 11:1. [20] Hutson, R. 1926. New Jersey Agr. Expt. Sta. Bull. 434. [21] Jensen, L., and H. Bøgh. 1941. Tidsskr. Planteavl 46:238. [22] Jones, M. D., and J. S. Brooks. 1950. Oklahoma Agr. Expt. Sta. Tech. Bull. 38. [23] Jones, M. D., and L. C. Newell. 1946. Nebraska Univ. Agr. Expt. Sta. Res. Bull. 148. [24] McQuilken, W. E. 1940. Ecology 21:135. [25] MacVicar, R. M., et al. 1952. Sci. Agr. 32:67. [26] Megee, C. R., and R. H. Kilty. 1932. Mich. State Univ. Agr. Expt. Sta. Quart. Bull. 14:271. [27] Meier, F. C., and F. Artschwager. 1938. Science 88:507. [28] Parris, G. K., and J. D. Haynie. 1950. Florida Dept. Agr. (Tallahassee) Bull. 135:45. [29] Pope, O. A., et al. 1944. J. Agr. Res. 68:347. [30] Roberts, R. H. 1945. Proc. Am. Soc. Hort. Sci. 46:87. [31] Rodrigo, P. A. 1925. Philippine Agriculturist 14:155. [32] Roemer, T. 1931. Z. Zuecht., A, 17:14. [33] Stephens, S. G., and M. D. Finkner. 1953. Econ. Botany 7:257. [34] Wit, F. 1952. Euphytica 1:95. [35] Wolfe, H. S., et al. 1934. Florida Univ. Agr. Expt. Sta. (Gainesville) Bull. 272. [36] Wolfenbarger, D. O. 1954. Florida Univ. Agr. Expt. Sta. (Gainesville) Ann. Rept., 1953, p. 290. [37] Wright, J. W. 1952. U.S. Dept. Agr. Forest Serv. Northeastern Forest Expt. Sta. Paper 46.

**115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES**

**Part I. ANIMAL VIRUSES**

	Virus	Substrate	Inactivation		Reference		Virus	Substrate	Inactivation		Reference
			Temp. °C	Time min					Temp. °C	Time min	
	(A)	(B)	(C)	(D)	(E)		(A)	(B)	(C)	(D)	(E)
1	Adenovirus		56	2.5-5.0	17	18	Herpes simplex	Aqueous suspension	52	30	12
2			56	30	10	19		Moist	52	30	3
3	Common cold		56	30	2,3	20		Dry	90	30	3
4	Coxsackie A		53-55	30	4	21	Hog cholera		60-70	60	19
5	Coxsackie C	Aqueous suspension	60	30	17	22	Infectious hepatitis		56	30	17
6		Milk, cream	70-80	30	17	23			60	60	
7		Tissue suspension <sup>1</sup>	55	30	5	24	Influenza A	Allantoic preparation	55	5-15	9
8		Water	60	30	13	25			56	30 <sup>3</sup>	3
9	Dengue		50	30	21	26			67	30	1
10	Distemper, canine		58	20	20	27	Influenza B	Allantoic preparation	56	15-30	9
11			60	30	15	28	Louping ill		56	30	7
12	Encephalitis, western equine	Filtrate	56	30	8	29		Tissue suspension <sup>1</sup>	58	10	17
13			60	10	17	30			60	2	17
14	Foot-and-mouth disease	Defibrinated blood	55	20	16	31			80	0.5	17
15		Epithelium suspension <sup>2</sup>	85	360	6	32	Lymphogranuloma venereum		56	10	17
16		Vesicular fluid	60	5	16	33	Measles		55	15	19
17	Fowl plague		55	30	15	34			56	60	12
						35	Mumps		55	20	17
						36			56	20	3

<sup>1</sup>/ Mouse brain. <sup>2</sup>/ Cattle tongue. <sup>3</sup>/ Some strains 90 minutes.

*continued*

## 115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

### Part I. ANIMAL VIRUSES

	Virus	Substrate	Inactivation		Ref-er-ence	Virus	Substrate	Inactivation		Ref-er-ence	
			Temp. °C	Time min				Temp. °C	Time min		
	(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	
37	Newcastle disease		56	5 <sup>4</sup>	1	50	Rabies	60	5	3,12	
38			56	120	14	51		100	2-3	3	
39	Poliomyelitis	Aqueous suspension	45-55	30	18	52		Aqueous suspension	54-60	60	17
40			50-55	30	12	53		Dry	54-56	1,440	17
41		Milk, cream, ice cream	62	30	17	54	Rift Valley fever	Blood	56	40	20
42		Milk	61.7	30	11	55	Small pox		55	30	17
43			71.1	0.25		56		Moist heat	60	10	3
44		Tissue suspension <sup>1</sup>	55	30	5	57		Dry heat	100	10	3
45		Water	50-55	30	11	58	Vaccinia	Dry	100	10	17
46			75	30	3	59		Fluid suspension	60	10	
47	Psittacosis		60	10	17	60	Trachoma		45	15	17
48	Rabies		50	60	3,12	61	Yellow fever		55	5	19
49			56	30	19	62			65	10	12

<sup>1</sup>/ Mouse brain. <sup>4</sup>/ Some strains 360 minutes.

Contributors: (a) Dupre, Margaret V., (b) Frobisher, Martin

References: [1] Anderson, S. G. 1959. The viruses. Academic Press, New York. v. 3. [2] Andrews, C. H. 1960. Sci. Am. 203(6):88. [3] Cruickshank, R., ed. 1960. Mackie and McCartney's Handbook of bacteriology. Ed. 10. E. and S. Livingstone, London. [4] Dalldorf, G. 1950. Bull. N. Y. Acad. Med. 26:329. [5] Dalldorf, G. 1955. Ann. Rev. Microbiol. 9:277. [6] Dimopoulos, G. T. 1960. Ann. N. Y. Acad. Sci. 83:706. [7] Edward, D. G. 1949. Brit. J. Exptl. Pathol. 30:582. [8] Fastier, L. B. 1952. J. Immunol. 68(5):531. [9] Francis, T., Jr. 1947. Ann. Rev. Microbiol. 1:351. [10] Huebner, R. J., W. P. Rowe, and R. M. Chanock. 1958. Ibid. 12:49. [11] Kaplan, H. S., and J. L. Melnick. 1952. Am. J. Public Health 42:525. [12] Jawetz, E., J. L. Melnick, and E. A. Adelberg. 1960. Review of medical microbiology. Ed. 4. Lange Medical Publications, Los Altos, Calif. [13] Melnick, J. L. 1950. Bull. N. Y. Acad. Med. 26:342. [14] Nelson, C. B., et al. 1952. Am. J. Public Health 42:672. [15] Porter, J. R. 1946. Bacterial chemistry and physiology. J. Wiley, New York. [16] Reddish, G. F., ed. 1954. Antiseptics, disinfectants, fungicides, and chemical and physical sterilization. Lea and Febiger, Philadelphia. [17] Rivers, T. M., and F. L. Horsfall, Jr., ed. 1959. Viral and rickettsial infections of man. Ed. 3. J. B. Lippincott, Philadelphia. [18] Schultz, E. W. 1948. Ann. Rev. Microbiol. 2:335. [19] Smith, D. T., and N. F. Conant, ed. 1960. Zinsser's Microbiology. Ed. 12. Appleton-Century-Crofts, New York. [20] Smith, D. T., and D. S. Martin, ed. 1948. Zinsser's Textbook of bacteriology. Ed. 9. Appleton-Century-Crofts, New York. [21] Stitt, E. R., P. W. Clough, and S. E. Branham. 1948. Practical bacteriology, hematology, and animal parasitology. Ed. 10. Blakiston, New York.

### Part II. PLANT VIRUSES

Medium (column D): dil. = diluted. Inactivation (column G): Temperature at which infectivity is lost in 10 minutes.

	Virus	Plant Source of Virus	Species	Test Conditions		Survival Time	Inactivation, °C	Ref-er-ence
				Medium	Temp., °C			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Alfalfa mosaic	<i>Cucumis sativus</i>	<i>Nicotiana tabacum</i>	Dry tissue <sup>1</sup>	1-2	>303 da		44
2		<i>Nicotiana tabacum</i> <sup>2</sup>	<i>Phaseolus vulgaris</i>	Plant juice	4	>7 da		56
3			(Early Golden Cluster)	Phosphate buffer <sup>3</sup>	24	>4 da		

<sup>1</sup>/ Desiccated above freezing temperature. <sup>2</sup>/ Necrotic type, young. <sup>3</sup>/ Purified virus in 0.1 M phosphate buffer.

continued

115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

Part II. PLANT VIRUSES

	Virus	Plant Source of Virus	Species	Test Conditions		Survival Time	Inactivation, °C	Reference
				Medium	Temp., °C			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
4	Alfalfa mosaic	<i>Pisum sativum</i> & <i>Vicia faba</i>	<i>Phaseolus vulgaris</i> (Stringless Green Refugee)	Plant juice	20 (dark)	<5 da	70	76
5	Dolichos lablab strain	<i>Vicia faba minor</i>	<i>V. faba minor</i>	Plant juice	25	>24-<48 hr	65-71	51
6				Crushed dried leaves	Room	>3 mo		
7	Israel strain	<i>Nicotiana glutinosa</i>	<i>Phaseolus vulgaris bulgarit</i>	Plant juice dil. 1:1 <sup>4</sup>	10-15	4 hr		47
8	Pierce strain	<i>Medicago sativa</i>	<i>Phaseolus vulgaris</i> (Stringless Green Refugee)	Plant juice	Room ?	9 da	64	53
9	Vein necrosis strain	<i>Glycine soja</i>	<i>Phaseolus vulgaris</i> (Pinto U.I. 111)	Plant juice	18	>30-<32 hr	62-64	78
10		<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Dry tissue	Room	>50-<95 da		
11	Common bean mosaic	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Stringless Green Refugee)	Plant juice	18	>28-<32 hr	56-58	53
12	Southern bean mosaic	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Plant juice	18	32 wk	90-95	77
13	Bean-pod mottle	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Plant juice	18	>62-<93 da	70-75	79
14	Bean yellow mosaic	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Stringless Green Refugee)	Plant juice	18	>24-<32	58-60	81
15	Bean yellow stipple	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Plant juice	18	5 da	72-75	80
16	Cucumber mosaic	<i>Cucumis sativus</i>		Plant juice	Room	<22 da	60-70	17
17				Dry tissue	Room	>40 da		
18		<i>Nicotiana glutinosa</i>	<i>Chenopodium amaranticolor</i>	Plant juice	10-15	6 da		47
19			<i>N. tabacum</i>	Plant juice <sup>5</sup>	26-32	>3 da	>60-<65	35
20		<i>Nicotiana tabacum</i>	<i>N. tabacum</i>	Plant juice <sup>5</sup>	26-32	>1-<2 da	>65-<70	35
21		<i>Zea mays</i>	<i>Nicotiana tabacum</i>	Leaf powder <sup>1</sup>	1-2	>669 da		44
22		<i>Z. mays</i> , <i>Nicotiana tabacum</i>	Leaf powder <sup>6</sup>	23 (over CaCl <sub>2</sub> )	>58 da		44	
23	Pea mosaic	<i>Pisum sativum</i>	<i>P. sativum</i> (Prince of Wales)	Plant juice	18	>48-<72 hr	60-64	22
24	Pea enation mosaic	<i>Pisum sativum</i>	<i>P. sativum</i> (Prince of Wales)	Plant juice	18	>72-<96 hr	56-58	72
25	Pea mottle	<i>Pisum sativum</i>		Plant juice	18	>24-<32 hr	56-58	81
26	Pea streak	<i>Pisum sativum</i>	<i>P. sativum</i> (Perfected Prince of Wales)	Plant juice	18	>16-<32 da	58-60	27
27	Pea stunt	<i>Pisum sativum</i>	<i>P. sativum</i> (Perfected Prince of Wales)	Plant juice	18	>48-<72 hr	58-60	26
28	Potato A	<i>Nicandra physalodes</i> <sup>7</sup>	<i>Solanum demissum</i>	Plant juice, undiluted	18	<18 hr		45
29		<i>Nicotiana tabacum</i> ?	<i>N. tabacum</i> ?	Plant juice	Room	Few hours	ca. 50	28
30	Veinbanding strain	<i>Nicotiana tabacum</i> ?	<i>N. tabacum</i>	Plant juice	Room	4 da	>58-<60	39
31	Potato M	<i>Solanum tuberosum</i> (seedling EK)	<i>Datura metel</i>	Plant juice	20	>2-<4 da	65-70	2
32	Potato S	<i>Solanum tuberosum</i> (seedling 41956)	<i>Nicotiana debneyi</i>	Plant juice	20	>4-<8 da	>55-<60	2
33	Potato X	<i>Lycopersicon esculentum</i>	<i>Nicotiana tabacum</i> (Connecticut Havana) & <i>L. esculentum</i> (John Baer)	Leaves, air-dried	Room	50 da		13

/1/ Desiccated above freezing temperature. /4/ With water. /5/ Kept in a darkened drawer. /6/ Dried at 35°C. /7/ Inoculated 14 days before test.

continued



115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

Part II. PLANT VIRUSES

Virus	Plant Source of Virus	Species	Test Conditions		Survival Time	Inactivation, °C	Reference	
			Medium	Temp., °C				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
34	Potato X	<i>Lycopersicon esculentum</i> <sup>8</sup>	<i>Nicotiana tabacum</i> (Connecticut Havana) & <i>L. esculentum</i> (John Baer)	Leaves, air-dried	Room	1,251 da	13	
35		<i>Nicotiana tabacum</i>	<i>N. tabacum</i> (Connecticut Havana) & <i>Lycopersicon esculentum</i> (John Baer)	Leaves, air-dried	Room	>286 da	13	
36		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	14-17	>234 da	>70-<75	74
37		<i>Nicotiana rustica</i>	<i>N. rustica</i>	Plant juice	16-20	>360 da	72	42
38	Heat-resistant strains	<i>Nicotiana tabacum</i> (Samsun)	<i>N. tabacum</i> (Samsun)	Plant juice	Room		>72-<76	40
39		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	Room	4 mo	70	57
40	Heat-susceptible strains	<i>Nicotiana tabacum</i> (Samsun)	<i>N. tabacum</i> (Samsun)	Plant juice	Room		68	40
41		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	Room	4-5 mo	68	57
42	Mottle strain	<i>Nicotiana tabacum</i> (Connecticut Havana No. 38)	<i>N. tabacum</i> (Connecticut Havana No. 38)	Plant juice	Room	>28 da	>68-<70	38
43	Ring spot strain	<i>Nicotiana tabacum</i> (Connecticut Havana No. 38)	<i>N. tabacum</i> (Connecticut Havana No. 38)	Plant juice	Room	>28 da	>65-<68	38
44	Potato Y	<i>Datura metel</i>	<i>D. metel</i>	Plant juice	Room ?	2-3 da	55-60	12
45		<i>Nicotiana tabacum</i>	<i>N. tabacum</i>	Leaf tissue <sup>9</sup>	1-2	>78 da		44
46		<i>Nicotiana tabacum</i> ?	<i>N. tabacum</i> ?	Plant juice	15	>3-<4 da	57	23
47		<i>Nicotiana tabacum</i> & <i>Solanum tuberosum</i>	<i>N. tabacum</i> (Connecticut Havana No. 38)	Plant juice	Room ?	>6 da	60	39
48		<i>Nicotiana tabacum</i> <sup>10</sup>	<i>N. tabacum</i> <sup>11</sup>	Dried leaves	4	>16 mo		20
49				Plant juice	20-22	>6-<18 da	56-62	
50	Necrotic strain	<i>Nicotiana tabacum</i> (Samsun)	<i>N. tabacum</i> (Samsun)	Plant juice	21-23	>50 da	62	37
51		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	20	70 da	61	1
52	Pepper veinbanding mosaic strain	<i>Nicotiana tabacum</i> (Turkish)	<i>N. tabacum</i> (Turkish)	Plant juice, undiluted	23	>10-<15 da	61-65	66
53	Standard strain	<i>Nicotiana repanda</i>	<i>N. repanda</i>	Plant juice	10-15	3 da		47
54		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	20	12 da	60	1
55	Vein-necrosis strain	<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (White Burley)	Plant juice	Room	<8 da	58	48
56	Potato aucuba mosaic	<i>Solanum tuberosum</i> (President)	<i>S. nodiflorum</i>	Plant juice	15 (dark)	>3-<4 da	>63-<65	16
57		<i>Nicotiana tabacum</i> ?	<i>N. sylvestris</i>	Plant juice	15	>4 da	68	24
58		<i>Nicotiana tabacum</i> (Samsun)	<i>Capsicum frutescens</i> (Early Calwonder)	Plant juice, dil. 1:9 <sup>4</sup>	18-20 (dark)	30-90 da		41
59				Plant juice, dil. 1:4 <sup>4</sup>			>65-<70	
60	Potato stem mottle, tobacco rattle strain	<i>Nicotiana tabacum</i> (Samsun)	<i>N. tabacum</i> (Samsun)	Plant juice, unpurified	20-22	>260 da		60
61				Dried leaves	20-22	>120 da		
62	Type M	<i>Nicotiana tabacum</i>	<i>Phaseolus vulgaris</i>	Plant sap, dil. 1:99 <sup>4</sup>	20	>6 wk		14
63				Plant sap, dil. 1:4 <sup>4</sup>			>80-<85	
64				Frozen plant juice	-10	>15 mo		
65	Potato yellow dwarf	<i>Nicotiana rustica</i>	<i>N. rustica</i>	Plant juice	23-27	>12 hr	52	10

<sup>4</sup>/ With water. <sup>8</sup>/ Also infected with tobacco mosaic virus. <sup>9</sup>/ Finely cut and dried over CaCl<sub>2</sub>. <sup>10</sup>/ Recently infected. <sup>11</sup>/ Infected with potato X virus.

continued

115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

Part II. PLANT VIRUSES

	Virus	Plant Source of Virus	Species	Test Conditions		Survival Time	Inactivation, °C	Reference
				Medium	Temp., °C			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
66	Potato yellow dwarf	<i>Nicotiana rustica</i>	<i>N. rustica</i>	Juice from frozen leaves	13-15	>1-<7 mo		11
67				Juice in KH <sub>2</sub> PO <sub>4</sub> buffer	0	>7 da		11
68				Buffer <sup>12</sup>	0	>4 wk		11
69	Sugar beet curly top	<i>Beta vulgaris</i>	<i>B. vulgaris</i>	Phloem exudate, dil. <sup>13</sup>			80	7
70				Plant leaf juice	Room	>7-<14 da		
71		<i>Beta vulgaris</i> <sup>14</sup>	<i>B. vulgaris</i>	Dried natural exudate	Room	>10 mo		7
72		<i>Beta vulgaris</i>	<i>B. vulgaris</i>	Exudate <sup>15</sup>	Room	>5 mo		7
73				Dried leaves <sup>16</sup>	Room <sup>17</sup>	8 yr		8
74	Tobacco etch virus	<i>Nicotiana glutinosa</i>	<i>Physalis peruviana</i>	Dried plant juice <sup>18</sup>	Room	>10 da		31
75		<i>Nicotiana tabacum</i>	<i>Physalis peruviana</i>	Plant juice, dil. 1:10 <sup>19</sup>	Above freezing point	>10 da		31
76			<i>N. tabacum</i>	Plant leaf tissue <sup>1</sup>	1-2	>301 da		44
77			<i>N. tabacum</i> (starch-iodine method)	Plant juice	Room	13 da	58	5
78				Dried leaves	Room	24 da		
79	Tobacco mosaic	<i>Nicotiana tabacum</i> (Turkish)	<i>N. glutinosa</i> & <i>Phaseolus vulgaris</i> (Early Golden Cluster)	Plant juice, dil. 1:20 <sup>4</sup>	68	20 da		54
80				Plant juice, dil. 1:20 <sup>4</sup>			93	
81			<i>N. glutinosa</i>	Plant juice	10-15	49 da		47
82		<i>Nicotiana tabacum</i> (White Burley)	<i>N. tabacum</i> (Turkish)	Whole leaves, dried	Room	52 yr		33
83	Tobacco necrosis	<i>Nicotiana tabacum</i>	<i>Vigna sinensis</i> ?	Plant juice, dil. 1:30	29	>9-<40 da		71
84				Plant juice, dil. 1:5	2	>40 da		
85		<i>Nicotiana tabacum</i> ?	<i>Vigna sinensis</i> ?	Plant juice <sup>20</sup>	Room	>6 mo		69
86	Bean strain	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Plant juice	18	22 da	85-90	6
87	Tobacco ring spot	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i>	Plant juice	Room	7-9 da	66	53
88		<i>Nicotiana rustica</i>	<i>N. tabacum</i>	Plant juice, kept dark	26-32	>6 da	>65-<70	35
89		<i>Nicotiana sylvestris</i>	<i>N. tabacum</i>	Plant juice, kept dark	26-32	>1-<2 da	>60-<65	35
90		<i>Nicotiana tabacum</i>	<i>N. tabacum</i>	Plant juice, kept dark	26-32	6 da	>65-<70	35
91			<i>N. tabacum</i> <sup>16</sup>	Plant juice	Room	>12-<24 hr	>60	30
92			<i>N. tabacum</i>	Plant juice	15	>1-<3 da		55
93				Plant juice	5	>17-<19 da		
94		<i>Nicotiana tabacum</i> & <i>Petunia hybrida</i>	<i>N. tabacum</i> <sup>10</sup>	Plant juice	-18	>22 mo		30
95		<i>Nicotiana tabacum</i>	<i>N. tabacum</i>	Leaf tissue <sup>9</sup>	1-2	>393 da		44
96	Common strain	<i>Lactuca sativa</i>	<i>Nicotiana tabacum</i> ?	Plant juice	23	36-48 hr		25
97	Eucharis strain	?	?	Plant juice	Room	6-8 da	65	36
98	Lettuce calico strain	<i>Nicotiana tabacum</i>	<i>N. tabacum</i> ?	Plant juice	23	72-96 hr		25
99	Tobacco streak Standard strain	<i>Phaseolus vulgaris</i> (Pinto U.I. No. 78)	<i>P. vulgaris</i> (Pinto U.I. No. 78)	Plant juice	18	<24 hr	54-56	73
100				Plant leaves <sup>21</sup>	18	>90 da		
101		<i>Nicotiana tabacum</i> (Havana)	<i>N. tabacum</i> (Havana)	Plant juice	22	>24-<36 hr	53	34

/1/ Desiccated above freezing temperature. /4/ With water. /5/ Finely cut and dried over CaCl<sub>2</sub>. /12/ Partially purified virus in buffer. /13/ With sugar solution. /14/ Petioles. /15/ Dried alcoholic precipitate. /16/ Young plant. /17/ Kept over CaCl<sub>2</sub> in airtight container. /18/ Resuspended residue. /19/ In presence of acid buffer. /20/ Dried precipitate or in absolute alcohol. /21/ Air-dried at 20-25°C.

*continued*

## 115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

### Part II. PLANT VIRUSES

	Virus	Plant Source of Virus	Species	Test Conditions		Survival Time	Inactivation, °C	Reference
				Medium	Temp., °C			
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
102	Tobacco streak							
	Bean red node strain	<i>Phaseolus vulgaris</i> (Pinto U.I. No. 78)	<i>P. vulgaris</i> (Pinto U.I. No. 78)	Plant juice	18	>24-<48 hr		73
103				Plant leaves <sup>21</sup>	18	>30-<90 da		
104	Brazilian strain	<i>Nicotiana tabacum</i>	<i>N. tabacum</i>	Plant juice	Room	>12-<24 hr	50-55	19
105		<i>Nicotiana tabacum</i> (recovered tissue)	<i>N. tabacum</i> (Turkish)	Plant juice <sup>22</sup>	Room ?	>9-<27 hr	>60-<65	18
106	Canadian strain	<i>Nicotiana tabacum</i> ?	<i>N. tabacum</i> ?	Plant juice	Room	<24 hr	54	9
107	Pea strain	<i>Phaseolus vulgaris</i>	<i>P. vulgaris</i> (Pinto U.I. 111)	Plant juice	22	>26-<27 hr	64	52
108				Plant leaves <sup>23</sup>	22	>98 da		
109	Tomato aspermy	<i>Nicotiana tabacum</i>		Plant juice	23-25	>12 hr	70	49
110	Tomato bunchy top	<i>Lycopersicon esculentum</i>	<i>Nicotiana glutinosa</i>	Plant juice	Room	>24 hr	70	43
111	Tomato bushy stunt	<i>Lycopersicon esculentum</i>	<i>Vigna sinensis</i>	Plant juice	Room	33 da	80	68
112	Tomato mosaic	<i>Lycopersicon esculentum</i>	<i>L. esculentum</i>	Plant juice	Room	>138 da	85-90	75
113				Dry tissue	Room	Indefinite	85-90	
114	Tomato ring spot	<i>Lycopersicon esculentum</i>	<i>Datura stramonium</i>	Plant juice	Room	>21-<27 hr		58
115			<i>L. esculentum</i>	Plant juice	Room		>56-<58	58
116				Dry tissue	Room	<300 hr	58	
117	Beet ring spot strain	<i>Nicotiana tabacum</i> (White Burley)	<i>Phaseolus vulgaris</i> ?	Plant juice	20	>2-<3 wk	>63-<66	29
118	Brazilian strain	<i>Lycopersicon esculentum</i>	<i>Nicandra physalodes</i>	Plant juice	25	>13-<20 da		65
119	Tomato black ring strain	<i>Nicotiana tabacum</i> ?	<i>N. tabacum</i> ?	Plant juice	Room	>7 da	>58-<62	70
120	Tomato ring spot strain & peach yellow bud strain	<i>Petunia hybrida</i>	<i>P. hybrida</i>	Plant juice, dil. 1:4 <sup>4</sup>	-10	>5 da	>60-<65	15
121	Tomato spotted wilt	<i>Lycopersicon esculentum</i> <sup>10</sup>	<i>L. esculentum</i>	Plant juice, dil. <sup>4</sup>	16	>4.5-<6 hr	>39-<42	3
122		<i>Lycopersicon esculentum</i> <sup>10,18</sup>	<i>L. esculentum</i> <sup>16</sup>	Plant juice, untreated	21	>2 hr	>44-<46	50
123		<i>Lycopersicon esculentum</i>	<i>Nicotiana glutinosa</i>	Plant juice	Room	5 hr	42	59
124		<i>Lycopersicon esculentum</i> (top leaves)	<i>Nicotiana tabacum</i> (Blue Pryor)	Plant juice, dil. <sup>4</sup>	20-22	<7 hr		4
125				Plant juice <sup>24</sup>	20-22	>36 hr		
126		<i>Nicotiana tabacum</i> (White Burley)	<i>N. glutinosa</i> or <i>Petunia hybrida</i>	Plant juice, dil. <sup>4</sup>	Room	3.5 hr		67
127			<i>N. tabacum</i> (White Burley)	Plant juice, dil. <sup>4</sup>	Room	2 hr		
128	Corcova strain	<i>Lycopersicon esculentum</i>	<i>L. esculentum</i> & <i>Nicotiana glutinosa</i>	Plant juice	21	<2.5 hr	>46-<48	21
129	Tomato tip blight strain	<i>Lycopersicon esculentum</i>	<i>L. esculentum</i>	Plant juice, undiluted	18	<1 hr	41.5	46
130	Tropaeolum mosaic	<i>Tropaeolum majus</i>	<i>T. majus</i>	Plant juice	25	<24 hr	58	64
131	Brazilian strain	<i>Tropaeolum majus</i>	<i>T. majus</i>	Plant juice	3	>3 da		64
132	California strain	<i>Tropaeolum majus</i>	<i>T. majus</i>	Plant juice	Room	4 da	55	32
133	Ring mosaic strain	<i>Chenopodium quinoa</i>	<i>C. quinoa</i>	Plant juice	Room	>11-<18 da	>66-<68	61
134		<i>Nicotiana glutinosa</i>	<i>N. glutinosa</i>	Plant juice	Room	<3 da		61
135				Dried leaves	Room	>8-<9 da		
136		<i>Nicotiana alata</i>	<i>N. alata</i>	Plant juice	Room	5 da	>65-<68	62
137				Dried leaves	Room	>2-<3 wk		
138	Ring spot mosaic strain	<i>Nicotiana tabacum</i> (Samsun)	<i>Chenopodium quinoa</i>	Plant juice, purified	Room (dark)	>24-<30 da	62-65	63

<sup>4</sup>/ With water. <sup>10</sup>/ Recently infected. <sup>16</sup>/ Young plant. <sup>21</sup>/ Air-dried at 20-25°C. <sup>22</sup>/ In 0.02 M phosphate buffer plus sodium sulfite. <sup>23</sup>/ Air-dried and powdered. <sup>24</sup>/ In 0.2% solution of sodium sulfite.

Contributors: (a) Silberschmidt, Karl M., (b) Zaumeyer, William J., (c) Webb, Raymon E.

continued



# 115. EFFECT OF TEMPERATURE ON INACTIVATION AND SURVIVAL: VIRUSES

## Part II. PLANT VIRUSES

- References:* [1] Aubert, O. 1960. Mem. Soc. Vaudoise Sci. Nat. 12(77):153. [2] Bagnall, R. H., R. H. Larson, and J. C. Walker. 1956. Wisconsin Univ. Agr. Expt. Sta. Res. Bull. 198. [3] Bald, J. G., and G. Samuel. 1931. Australia Council Sci. Ind. Res. Bull. 54. [4] Bald, J. G., and G. Samuel. 1934. Ann. Appl. Biol. 21:179. [5] Bawden, F. C., and B. Kassanis. 1941. Ibid. 28:107. [6] Bawden, F. C., and J. P. H. van der Want. 1945. Tijdschr. Plantenziekten 55:142. [7] Bennett, C. W. 1935. J. Agr. Res. 50:211. [8] Bennett, C. W. 1942. Phytopathology 32:826. [9] Berkeley, G. H., and J. H. H. Phillips. 1943. Can. J. Res., C, 21:181. [10] Black, L. M. 1938. Phytopathology 28:863. [11] Black, L. M. 1951. Ibid. 41:213. [12] Borges, M. V. 1958. Agron. Lusitana 20(4):287. [13] Burnett, G. 1934. Phytopathology 24:215. [14] Cadman, C. H., and B. D. Harrison. 1959. Ann. Appl. Biol. 47:542. [15] Cadman, C. H., and R. M. Lister. 1961. Phytopathology 51:29. [16] Clinch, P. E. M., J. B. Loughnane, and P. A. Murphy. 1936. Sci. Proc. Roy. Dublin Soc., N.S. 21:431. [17] Cohen, S., and F. E. Nitzany. 1963. Phytopathology 53:193. [18] Costa, A. S., and A. M. B. Carvalho. 1961. Phytopathol. Z. 42:113. [19] Costa, A. S., A. R. Lima, and R. Forster. 1940. J. Agron. São Paulo 3:1. [20] Darby, J. F., R. H. Larson, and J. C. Walker. 1951. Wisconsin Univ. Agr. Expt. Sta. Res. Bull. 177. [21] Delle Coste, A. C., and S. Zabala. 1946. Publ. Inst. Sanidad Vegetal (Buenos Aires), A, 17. [22] Doolittle, S. P., and F. R. Jones. 1925. Phytopathology 15:763. [23] Dykstra, T. P. 1939. Ibid. 29:40. [24] Dykstra, T. P. 1939. Ibid. 29:917. [25] Grogan, R. G., and W. C. Schnathorst. 1955. Plant Disease Repr. 39:803. [26] Hagedorn, D. J., and J. C. Walker. 1949. J. Agr. Res. 78:617. [27] Hagedorn, D. J., and J. C. Walker. 1949. Phytopathology 39:837. [28] Hansen, H. P. 1937. Tidsskr. Planteavl 42:631. [29] Harrison, B. D. 1957. Ann. Appl. Biol. 45:462. [30] Henderson, R. G., and S. A. Wingard. 1931. J. Agr. Res. 43:191. [31] Holmes, F. O. 1942. Phytopathology 32:1058. [32] Jensen, D. D. 1950. Ibid. 40:967. [33] Johnson, E. M., and W. D. Valleau. 1935. Kentucky Agr. Expt. Sta. Res. Bull. 361:264. [34] Johnson, J. 1936. Phytopathology 26:285. [35] Johnson, J., and T. J. Grant. 1932. Ibid. 22:741. [36] Kahn, R. P., and H. A. Scott. 1962. Ibid. 52:16. [37] Klinkowski, M., and K. Schmelzer. 1957. Phytopathol. Z. 28:285. [38] Koch, K. L. 1933. Phytopathology 23:319. [39] Koch, K. L., and J. Johnson. 1935. Ann. Appl. Biol. 22:37. [40] Koehler, E. 1937. Phytopathol. Z. 10:31. [41] Kollmer, G. F., and P. H. Larson. 1960. Wisconsin Univ. Agr. Expt. Sta. Res. Bull. 223. [42] Ladenburg, R. C., R. H. Larson, and J. C. Walker. 1950. Ibid. 165. [43] McClean, A. P. D. 1931. Union S. Africa Dept. Agr. Sci. Bull. 100. [44] McKinney, H. H. 1947. Phytopathology 37:139. [45] MacLachlan, D. S., R. H. Larson, and J. C. Walker. 1953. Wisconsin Univ. Agr. Expt. Sta. Res. Bull. 180. [46] Milbrath, J. A. 1939. Phytopathology 29:156. [47] Nitzany, F. E., and S. Friedman. 1963. Ibid. 53:548. [48] Nobrega, N. R., and K. Silberschmidt. 1944. Arquiv. Inst. Biol. (São Paulo) 15:307. [49] Nooroom, D. 1952. Tijdschr. Plantenziekten 58:121. [50] Norris, D. O. 1946. Australia Council Sci. Ind. Res. Bull. 202. [51] Nour, M. A., and J. J. Nour. 1962. Phytopathology 52:427. [52] Patino, G., and W. J. Zaumeyer. 1959. Ibid. 49:43. [53] Pierce, W. H. 1934. Ibid. 24:87. [54] Price, W. C. 1933. Ibid. 23:749. [55] Priode, C. N. 1928. Am. J. Botany 15:88. [56] Ross, A. F. 1941. Phytopathology 31:394. [57] Salaman, R. N. 1938. Phil. Trans. Roy. Soc. London, B, 229:137. [58] Samson, R. W., and E. P. Imle. 1942. Phytopathology 32:1037. [59] Samuel, G., J. G. Bard, and H. A. Pittman. 1930. Australia Council Sci. Ind. Res. Bull. 44. [60] Schmelzer, K. 1957. Phytopathol. Z. 30:281. [61] Schmelzer, K. 1960. Z. Pflanzenkrankh. Pflanzenschutz 67:193. [62] Schumann, K. 1963. Phytopathol. Z. 48:135. [63] Schwarz, R. 1958. Ibid. 33:375. [64] Silberschmidt, K. 1953. Phytopathology 43:304. [65] Silberschmidt, K. 1963. Phytopathol. Z. 46:209. [66] Simon, J. N. 1956. Phytopathology 46:53. [67] Smith, K. M. 1932. Ann. Appl. Biol. 19:305. [68] Smith, K. M. 1935. Ibid. 22:731. [69] Smith, K. M. 1937. Parasitology 29:86. [70] Smith, K. M. 1946. Ibid. 37:126. [71] Smith, K. M., and J. G. Bald. 1935. Ibid. 27:231. [72] Stubbs, M. W. 1937. Phytopathology 27:242. [73] Thomas, R. R., and W. J. Zaumeyer. 1950. Ibid. 40:832. [74] Vasudeva, R. S., and T. B. Lal. 1945. Indian J. Agr. Sci. 14:288. [75] Walker, M. N. 1926. Phytopathology 16:431. [76] Zaumeyer, W. J. 1938. J. Agr. Res. 56:747. [77] Zaumeyer, W. J., and L. L. Harter. 1943. Ibid. 67:297. [78] Zaumeyer, W. J., and G. Patino. 1960. Phytopathology 50:226. [79] Zaumeyer, W. J., and H. R. Thomas. 1948. J. Agr. Res. 77:81. [80] Zaumeyer, W. J., and H. R. Thomas. 1950. Phytopathology 40:847. [81] Zaumeyer, W. J., and B. L. Wade. 1935. J. Agr. Res. 51:715.



## 116. EFFECT OF TEMPERATURE ON GROWTH AND SURVIVAL: RICKETTSIA AND BACTERIA

Values are for data obtained under diverse conditions by many investigators. Data may differ for various species within the same genus, and even for various cultures of the same species.

### Part I. OPTIMUM TEMPERATURE FOR GROWTH

Bacteria may be grouped on the basis of their optimum growth temperatures; psychrophiles (optima below 20°C), mesophiles (optima from 21°-50°C), and thermophiles (optima above 50°C). These groupings are arbitrary and not mutually exclusive. Some bacteria, with optima below 50°C but which grow well at temperatures above 50°C, are not true thermophiles, as defined above, and are called thermoduric. Values in parentheses are for minimum and maximum temperatures at which growth can occur.

Species	Temp. °C	Refer- ence	Species	Temp. °C	Refer- ence
(A)	(B)	(C)	(A)	(B)	(C)
Rickettsia			13 <i>Diplococcus pneumoniae</i>	37 (25-42)	3
1 <i>Bartonella bacilliformis</i>	28 (max. 37)	2,5,9, 13	14 <i>Erwinia carotovora</i>	25-30 (4-39)	6
2 <i>Coxiella burneti</i>	35	11	15 <i>Escherichia coli</i>	30-37 (10-45)	2
3 <i>Miyagawanella lymphogranu- lomatis</i>	35-37	2	16 <i>Lactobacillus acidophilus</i>	37 (22-48)	2
4 <i>Rickettsia prowazekii</i>	34-36 (30-37)	11	17 <i>Mycobacterium tuberculosis</i>	37 (30-42)	2,3
Bacteria			18 <i>Neisseria gonorrhoeae</i>	37 (30-39)	3,4
5 <i>Actinomyces bovis</i>	37 (20-40)	4	19 <i>Photobacterium fischeri</i>	25-28 (min. 5-10)	2
6 <i>Aerobacter aerogenes</i>	37 (2.5-45)	7	20 <i>Proteus vulgaris</i>	30-37 (good at 20)	3
7 <i>Agrobacterium tumefaciens</i>	25-30 (0-37)	6,14	21 <i>Pseudomonas aeruginosa</i>	37 (5-42)	13
8 <i>Azotobacter chroococcum</i>	25-28	2	22 <i>Rhizobium leguminosarum</i>	25	2,8,10
9 <i>Bacillus subtilis</i>	28-40 (max. 50-55)	2	23 <i>Salmonella typhosa</i>	37 (4-40)	12
10 <i>Brucella melitensis</i>	37 (6-45)	3	24 <i>Shigella dysenteriae</i>	37 (10-40)	5,13
11 <i>Clostridium botulinum</i>	25 (20-35)	10	25 <i>Staphylococcus aureus</i>	35-37 (10-42)	4
12 <i>Corynebacterium diphtheriae</i>	34-36 (15-40)	3	26 <i>Streptococcus pyogenes</i>	37 (10-42)	3
			27 <i>Streptomyces griseus</i>	30-37	1,2
			28 <i>Vibrio comma</i>	37 (16-42)	3
			29 <i>Xanthomonas campestris</i>	30-32 (5-39)	6,14

Contributor: Dupré, Margaret V.

- References: [1] Bradley, S. G. 1959. Appl. Microbiol. 7:89. [2] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. Bergey's Manual of determinative bacteriology. Ed. 7. Williams and Wilkins, Baltimore. [3] Burrows, W. 1963. Textbook of microbiology. Ed. 18. W. B. Saunders, Philadelphia. [4] Cruickshank, R., ed. 1960. Mackie and McCartney's Handbook of bacteriology. Ed. 10. E. and S. Livingstone, Edinburgh. [5] Dubos, R. J., ed. 1958. Bacterial and mycotic infections of man. Ed. 3. J. B. Lippincott, Philadelphia. [6] Elliott, C. 1951. Manual of bacterial plant pathogens. Ed. 2. Chronica Botanica, Waltham, Mass. [7] Foster, E. M., et al. 1957. Dairy microbiology. Prentice-Hall, Englewood Cliffs, N. J. [8] Hawker, L. E., et al. 1960. An introduction to the biology of microorganisms. W. Clowes, London. [9] Jawetz, E., J. L. Melnick, and E. A. Adelberg. 1960. Review of medical microbiology. Ed. 4. Lange Medical Publications, Los Altos, Calif. [10] Pelczar, M. J., Jr., and R. D. Reid. 1958. Microbiology. McGraw-Hill, New York. [11] Rivers, T. M., and F. L. Horsfall, Jr., ed. 1959. Viral and rickettsial infections of man. Ed. 3. J. B. Lippincott, Philadelphia. [12] Smith, A. L., ed. 1960. Carter's Microbiology and pathology. Ed. 7. C. V. Mosby, St. Louis. [13] Smith, D. T., and N. F. Conant, ed. 1960. Zinsser's Microbiology. Ed. 12. Appleton-Century-Crofts, New York. [14] Stapp, C. 1961. Bacterial plant pathogens. Oxford Univ. Press, London.

continued

## 116. EFFECT OF TEMPERATURE ON GROWTH AND SURVIVAL: RICKETTSIA AND BACTERIA

### Part II. THERMAL DEATH TIME

Bacteria which do not form endospores are generally killed within 20 minutes when directly exposed in fluid to temperatures of 70°C or over (moist heat), but thermophiles are somewhat more resistant. Bacterial endospores may resist moist heat at 100°C for two minutes to many hours. However, no known living organism can survive compressed steam at 121°C (routine autoclaving) for 20 minutes. The thermal death times listed below are generally based on exposure to moist heat, unless otherwise specified. Inconsistencies in the values may be attributed to different experimental methods and to the fact that all of the variables, especially pH (a crucial factor), were not always reported. **Substrate** (column B): PS = phosphate solution; PW = peptone water.

	Species	Substrate	Temp. °C	Time min	Reference		Species	Substrate	Temp. °C	Time min	Reference	
	(A)	(B)	(C)	(D)	(E)		(A)	(B)	(C)	(D)	(E)	
	Rickettsia						47	<i>Lactobacillus thermophilus</i>	82	2.5	2	
1	<i>Coxiella burnetii</i>		50	30	23	48	<i>Mycobacterium tuberculosis</i>		58	30	5	
2		Egg yolk sac	63-65	30	22	49			59	20	5	
3		Milk	62.2	30	7	50			65	2	5	
4		Milk <sup>1</sup>	62.7	30-40	14	51		Ice cream	62.6	6	20	
5		Milk <sup>2</sup>	67.7-68.7	0.25	17	52		Milk	55	60	10	
6		Milk <sup>3</sup>	71.6	0.25	18	53			60	10	10	
7		Milk	72.2	0.25	7	54			65	2	10	
8	<i>Miyagawanella lymphogranulomatis</i>		60	10	15	55			71	3	10	
9	<i>Rickettsia prowazekii</i>		56	30	23	56			82	<3	10	
						57			100	<1	10	
	Bacteria					58		Sputum	100	15	29	
10	<i>Actinomyces bovis</i>		62-64	3-10	28	59		Sputum (dried)	100	60	5	
11	<i>Aerobacter aerogenes</i>	Milk	57.2	<60	10	60	<i>Neisseria gonorrhoeae</i>	Infected pus <sup>7</sup>	42	300-900	6	
12			60	30	2	61			50	5	13	
13	<i>Bacillus subtilis</i>	PS, pH 4.4	100	2	11	62		Moist	55	5	25	
14		PS, pH 5.6	100	7	11	63			55	5	7,8	
15		PS, pH 6.8-7.6	100	11	11	64	<i>Proteus vulgaris</i>		55	60	3	
16		PS, pH 8.4	100	9	11	65	<i>Pseudomonas aeruginosa</i>		55	60	8,25	
17		1% PW <sup>4</sup>	100	11	11	66	<i>Salmonella typhosa</i>	Broth	55	23.8	27	
18	1% PW <sup>5</sup>	100	16	11	67			56	4-9	19		
19	<i>Brucella melitensis</i>	Moist	60	10	3	68			60	4.3	11	
20			65	5	3	69		Ice cream	57.2	10	20	
21		Aqueous emulsion	57.5	10	13	70			62.8	3	20	
22		Milk	61.1-62.7	30	5	71	Milk	55	30	24		
23		0.85% NaCl	62.5	10	1	72		60	20	13		
24	<i>Clostridium botulinum</i>	2% aminoids	100	65	9	73		61.1-62.7	4-8	27		
25		Corn (canned)	100	105	12	74						
26			115	15	12	75						
27		Pears (canned)	100	30	12	76	<i>Shigella dysenteriae</i>	Milk	60	6	27	
28			115	5	12	77			Milk <sup>6</sup>	60	10	21
29		PS, pH 7	100	330	9	78			Water	60	1	27
30		110	32	9	79	<i>Staphylococcus aureus</i>		60	30-60	8		
31		115	10	9	80			62	30	7		
32		120	4	9	81		Broth	65.6	2	26		
33	<i>Corynebacterium diphtheriae</i>	Broth	58	10	5		82	Broth <sup>6</sup>	58	10	13	
34		Ice cream	65.6	0.5	20		83	Broth	65	18.8	27	
35		Milk	55-60	2	24		84	Custard filling	88	10	11	
36		Water	58	10	25		85	Skimmed milk & 14% sugar	60	30	16	
37		100	1	25	86		Dry heat	190.6-218.3	30	11		
38	<i>Diplococcus pneumoniae</i>		52	10	25	87						
39			55	20	3	88	<i>Streptococcus pyogenes</i>		54	30	7	
40		Blood broth	56	5-7	19	89		Milk	60	15	4	
41		Broth <sup>6</sup>	60	30	13	90		Cream	57.2	5	20	
42		Melted dextrose agar	60	15	13	91			61.1	1	20	
43	<i>Escherichia coli</i>	Milk	57.2	>60	10	92		Diluted salt-gelatin	60	10	13	
44			60	30	5,8	93	Melted dextrose agar	60	15	13		
45			62.5	30	27	94	Milk	62	30	25		
46	<i>Lactobacillus thermophilus</i>		71	30	2	95	<i>Vibrio comma</i>		55	10	5,25	

<sup>1/1</sup>/ Sealed tubes in water. <sup>2/2</sup>/ Naturally infected. <sup>3/3</sup>/ Artificially infected. <sup>4/4</sup>/ Incubated at 21°-23°C. <sup>5/5</sup>/ Incubated at 37°C. <sup>6/6</sup>/ In sealed tubes. <sup>7/7</sup>/ In sealed pipette.

continued

**116. EFFECT OF TEMPERATURE ON GROWTH AND SURVIVAL:  
RICKETTSIA AND BACTERIA**

**Part II. THERMAL DEATH TIME**

*Contributors:* (a) Frobisher, Martin, (b) Dupré, Margaret V.

*References:* [1] Boak, R., and C. M. Carpenter. 1928. *J. Infect. Diseases* 43:327. [2] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. *Bergey's Manual of determinative bacteriology*. Ed. 7. Williams and Wilkins, Baltimore. [3] Bryan, A. H., C. A. Bryan, and C. G. Bryan. 1962. *Bacteriology*. Ed. 6. Barnes and Noble, New York. [4] Buchanan, R. E., and E. D. Buchanan. 1951. *Bacteriology*. Ed. 5. Macmillan, New York. [5] Burrows, W. 1963. *Textbook of microbiology*. Ed. 18. W. B. Saunders, Philadelphia. [6] Carpenter, C. M., et al. 1933. *J. Lab. Clin. Med.* 18:981. [7] Cruickshank, R., ed. 1960. *Mackie and McCartney's Handbook of bacteriology*. Ed. 10. E. and S. Livingstone, Edinburgh. [8] Dubos, R. J., ed. 1958. *Bacterial and mycotic infections of man*. Ed. 3. J. B. Lippincott, Philadelphia. [9] Esty, J. R., and K. F. Meyer. 1922. *J. Infect. Diseases* 31:650. [10] Foster, E. M., et al. 1957. *Dairy microbiology*. Prentice-Hall, Englewood Cliffs, N. J. [11] Frazier, W. C. 1958. *Food microbiology*. McGraw-Hill, New York. [12] Halversen, W. V., and G. L. Hays. 1936. *J. Bacteriol.* 32:466. [13] Hampil, B. 1932. *Quart. Rev. Biol.* 7:171. [14] Huebner, R. J., and J. A. Bell. 1951. *J. Am. Med. Assoc.* 145:301. [15] Jawetz, E., J. L. Melnick, and E. A. Adelberg. 1960. *Review of medical microbiology*. Ed. 4. Lange Medical Publications, Los Altos, Calif. [16] Kalan, R. S., W. H. Martin, and R. Mickelsen. 1963. *Appl. Microbiol.* 11:45. [17] Lennette, E. H., et al. 1952. *Am. J. Hyg.* 55:246. [18] Marmion, B. P., et al. 1951. *Min. Health (Gr. Brit.) Monthly Bull.* 10:119. [19] Morton, H. E. Unpublished. School of Medicine, Univ. of Pennsylvania, Philadelphia, 1953. [20] Oldenbusch, C., M. Frobisher, and J. H. Shrader. 1930. *Am. J. Public Health* 20:615. [21] Park, W. H. 1927. *Ibid.* 17:36. [22] Ransom, S. E., and R. J. Huebner. 1951. *Am. J. Hyg.* 53(1):110. [23] Rivers, T. M., and F. L. Horsfall, Jr., ed. 1959. *Viral and rickettsial infections of man*. Ed. 3. J. B. Lippincott, Philadelphia. [24] Rosenau, E. C. 1908. *U.S. Public Health Serv. Hyg. Lab. Bull.* 42. [25] Smith, D. T., and N. F. Conant, ed. 1960. *Zinsser's Microbiology*. Ed. 12. Appleton-Century-Crofts, New York. [26] Stritar, J. E. 1941. *Am. Meat Inst. 36th Ann. Conv.*, p. 15. [27] Tanner, F. W. 1944. *The microbiology of foods*. Ed. 2. Garrard Press, Champaign, Ill. [28] Waksman, S. A., and H. A. Lechevalier. 1959-62. *The actinomycetes*. Williams and Wilkins, Baltimore. v. 1-3. [29] Willis, H. S., and M. M. Cummings. 1952. *Diagnostic and experimental methods in tuberculosis*. C. C. Thomas, Springfield, Ill.

**117. EFFECT OF TEMPERATURE ON GROWTH AND SURVIVAL: FUNGI**

Values in parentheses are for minimum (min.) and maximum (max.) temperatures at which growth can occur.

	Species	Temperature for Growth <sup>1</sup> °C	Thermal Death		Refer- ence
			Temp. °C	Time min	
	(A)	(B)	(C)	(D)	(E)
Animal Pathogens					
1	<i>Aspergillus niger</i>	37 (max. <60)	.....	...	8
2	<i>Candida albicans</i>	30-37 (<20->40)	60	10	2-4,7
3	<i>Cladosporium mansonii</i>	30-32	.....	...	1,5
4	<i>Rhizopus equinus</i>	37-39 (min. >5)	100	20	1
5	<i>Torulopsis famata</i>	25-37	.....	...	1
Plant Pathogens					
6	<i>Alternaria brassicae</i>	33-35 (1-46) <sup>2</sup> ; 25-27 (2-36)	55	10	6
7	<i>Aspergillus niger</i> <sup>3</sup>	30-39 (7-46)	99 <sup>4</sup> ; 62 <sup>5</sup>	10	6
8	<i>Cladosporium fulvum</i> <sup>3</sup>	18-26 (0-33) <sup>2</sup> ; 16-26 (0-34)	70 <sup>4</sup>	10	6
9	<i>Fusarium oxysporum</i> <sup>6</sup>	15-32 (4-40)	57	10	6
10	<i>Helminthosporium turcicum</i>	28-30 (7-35)	.....	...	6

<sup>1</sup>/ In culture, unless otherwise specified. <sup>2</sup>/ Spore germination. <sup>3</sup>/ Fungus exhibits variability among different strains or in different hosts. <sup>4</sup>/ Dry. <sup>5</sup>/ Wet. <sup>6</sup>/ Fungus exhibits extreme variability among different strains or in different hosts.

*continued*

### 117. EFFECT OF TEMPERATURE ON GROWTH AND SURVIVAL: FUNGI

Species	Temperature for Growth <sup>1</sup> °C	Thermal Death		Reference
		Temp. °C	Time min	
(A)	(B)	(C)	(D)	(E)
Plant Pathogens				
11 <i>Penicillium expansum</i>	25-27 (0-30)	.....	...	6
12 <i>Puccinia graminis</i>	5-25 (2-35)	.....	...	6
13 <i>Rhizopus nigricans</i> <sup>2</sup>	19-41 (2-41) <sup>2</sup> ; 20-36 (2-40)	55 <sup>7</sup>	10	6
14 <i>Ustilago hordei</i> <sup>3</sup>	10-30 (0-35) <sup>2</sup> ; 16-26 (<1-<35)	43-48	10	6
15 <i>Venturia inaequalis</i>	13-25 (0-35) <sup>2</sup> ; 20 (<4-<32)	.....	...	6

<sup>1</sup>/ In culture, unless otherwise specified. <sup>2</sup>/ Spore germination. <sup>3</sup>/ Fungus exhibits variability among different strains or in different hosts. <sup>7</sup>/ Spores.

**Contributors:** (a) Gordon, Morris A., (b) Rossetti, Victoria

**References:** [1] Dodge, C. W. 1935. Medical mycology. C. V. Mosby, St. Louis. [2] Kadisch, E. 1930. Dermatol. Z. 60:48. [3] MacKinnon, J. E. 1946. El siglo ilustrado. Zimologia medica, Montevideo. [4] McClary, D. O. 1952. Ann. Missouri Botan. Garden 39:137. [5] Simons, R. D. G., ed. 1954. Medical mycology. Elsevier, Amsterdam. [6] Togashi, K. 1949. Biological characters of plant pathogens. Temperature relations. Meibundo, Tokyo. [7] Wickerham, L. J., and L. F. Rettger. 1939. J. Trop. Med. Hyg. 42:174, 187, 204. [8] Wolf, F. A., and F. T. Wolf. 1947. The fungi. J. Wiley, New York. v. 2.

### 118. TEMPERATURE TOLERANCES: ALGAE

Most values were based on observations of algae growing in their natural habitat. Since it is difficult to determine true temperature under such conditions, and since light absorption may raise the temperature of an algal mass above that of its surroundings, the data should be interpreted with caution. Values in parentheses are temperatures for maximum growth rate.

Species	Temperature, °C		Reference	Species	Temperature, °C		Reference
	Minimum	Maximum			Minimum	Maximum	
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
Cyanophyta							
1 <i>Anabaena variabilis</i>	.....	(35)	4	9 <i>Chlorella pyrenoidosa</i> (7-11-05)	.....	42 (38-39)	8
2 <i>Chroococcus yellowstonensis</i>	.....	41	2	10 <i>Cladophora hamosa</i>	<-7	>35	1
3 <i>Nostoc muscorum</i>	.....	(33)	4	11 <i>Protococcus botryoides</i>	.....	80	2
4 <i>Oscillatoria filiformis</i>	59	85 (73)	2	12 <i>Ulothrix</i> sp.	.....	17	7
5 <i>Phormidium bijahensis</i>	38	85 (60-62)	2	Phaeophyta			
6 <i>Synechococcus eximius</i>	70	84 (79)	2	13 <i>Fucus vesiculosus</i>	-18 to -20 <sup>1</sup>	30	5,6
Chlorophyta				14 <i>Laminaria saccharina</i>	-4 to -6 <sup>2</sup> ;	.....	6
7 <i>Chlamydomonas nivalis</i>	-36	4 (0)	3		-11 to -17 <sup>3</sup>		
8 <i>Chlorella pyrenoidosa</i> (Emerson)	.....	29 (25-26)	8	Rhodophyta			
				15 <i>Polysiphonia pulvinata</i>	<-7	>30	1
				16 <i>Porphyra leucosticta</i>	<-7	>30	1

<sup>1</sup>/ Based on observations in polar seas; uncertain that algae were actually growing. <sup>2</sup>/ One-year-old algae. <sup>3</sup>/ More than one year old.

**Contributors:** (a) Allen, Mary Belle, (b) Pisek, A., (c) Sorokin, Constantine

**References:** [1] Biebl, R. 1939. Jahrb. Wiss. Botan. 88:389. [2] Copeland, J. J. 1936. Ann. N. Y. Acad. Sci. 36:1. [3] Huber-Pestalozzi, G. 1926. In C. Schröter, ed. Das Pflanzenleben der Alpen. Raustein, Zurich. p. 942. [4] Kratz, W. A., and J. Myers. 1955. Am. J. Botany 42:282. [5] Kylin, H. 1910. Arkiv Botan. 10(1):1. [6] Kylin, H. 1917. Ber. Deut. Botan. Ges. 35:370. [7] Oltmans, F. 1923. Morphologie und Biologie der Algen. G. Fischer, Jena. v. 3. [8] Sorokin, C. 1959. Nature 184:613.



## 119. SOIL pH: SPERMATOPHYTES

With good management, and if other factors are favorable, many of the plants listed will grow and develop satisfactorily outside the pH range indicated. Field crops and vegetables generally are not as sensitive to soil pH as are flowers and shrubs.

Species (Common Name)	pH	Ref- er- ence	Species (Common Name)	pH	Ref- er- ence
(A)	(B)	(C)	(A)	(B)	(C)
Gymnospermae					
1 <i>Abies</i> spp. (fir)	4.5-6.5	4	32 <i>Citrus limon</i> (lemon)	6.0-7.5	3
2 <i>Ginkgo biloba</i> (ginkgo)	5.5-7.0	4	33 <i>C. sinensis</i> (sweet orange)	6.0-7.5	3
3 <i>Juniperus virginiana</i> (eastern red cedar)	5.0-8.0	4	34 <i>Cornus florida</i> (flowering dogwood)	5.0-6.5	4
4 <i>Larix</i> spp. (larch)	4.5-7.5	4	35 <i>Cucumis sativus</i> (cucumber)	5.5-7.0	3
5 <i>Picea</i> spp. (spruce)	4.5-6.5	4	36 <i>Cucurbita pepo</i> (pumpkin)	5.5-7.0	1
6 <i>Pinus</i> spp. (pine)	4.5-6.5	4	37 <i>Daucus carota</i> (carrot)	5.5-7.0	3
7 <i>Taxus</i> spp. (yew)	5.0-7.5	4	38 <i>Fagopyrum esculentum</i> (buckwheat)	5.5-7.0	2,3
8 <i>Thuja occidentalis</i> (northern white cedar)	6.0-7.5	3	39 <i>Fagus sylvatica</i> (European beech)	6.0-7.5	4
9 <i>Tsuga canadensis</i> (eastern hemlock)	4.5-6.0	4	40 <i>Fragaria</i> spp. (strawberry)	5.0-6.5	1,3
Angiospermae (Monocotyledoneae)			41 <i>Glycine soja</i> (soybean)	6.0-7.5	4
10 <i>Allium cepa</i> (garden onion)	6.0-7.5	4	42 <i>Gossypium hirsutum</i> (upland cotton)	5.0-6.5	4
11 <i>Asparagus officinalis</i> (garden asparagus)	6.0-8.0	3	43 <i>Helianthus annuus</i> (common sunflower)	6.0-7.5	2,3
12 <i>Avena sativa</i> (common oat)	5.0-7.5	2,3	44 <i>Ilex aquifolium</i> (English holly)	5.0-6.5	4
13 <i>Gladiolus</i> spp. (gladiolus)	6.0-8.0	1	45 <i>Ipomoea batatas</i> (sweet potato)	5.0-6.5	4
14 <i>Hordeum vulgare</i> (barley)	6.0-7.5	4	46 <i>Juglans</i> spp. (walnut)	6.0-7.5	4
15 <i>Iris</i> spp. (iris)	6.0-8.0	1	47 <i>Lactuca sativa</i> (lettuce)	6.0-7.5	4
16 <i>Lilium longiflorum</i> (Easter lily)	6.0-7.0	3	48 <i>Lycopersicon esculentum</i> (tomato)	5.5-7.5	3
17 <i>Oryza sativa</i> (rice)	5.0-6.5	2,3	49 <i>Magnolia grandiflora</i> (southern magnolia)	5.0-7.0	1
18 <i>Phleum pratense</i> (timothy)	6.0-8.0	3	50 <i>Malus pumila</i> (common apple)	5.0-6.5	3
19 <i>Poa pratensis</i> (Kentucky bluegrass)	5.5-7.5	2,3	51 <i>Medicago sativa</i> (alfalfa)	6.2-7.8	2,3
20 <i>Tradescantia virginiana</i> (Virginia spiderwort)	5.0-7.5	3	52 <i>Nicotiana tabacum</i> (common tobacco)	5.5-7.5	2,3
21 <i>Triticum aestivum</i> (wheat)	5.5-7.5	2	53 <i>Oenothera biennis</i> (common evening primrose)	6.0-8.0	3
22 <i>Zea mays</i> (corn)	5.5-7.5	2,3	54 <i>Phaseolus vulgaris</i> (kidney bean)	6.0-7.5	2,3
Angiospermae (Dicotyledoneae)			55 <i>Pisum sativum</i> (garden pea)	6.0-8.0	1
23 <i>Acer</i> spp. (maple)	5.5-7.5	4	56 <i>Populus tremuloides</i> (quaking aspen)	4.5-5.5	4
24 <i>Alnus</i> spp. (alder)	6.0-7.5	4	57 <i>Prunus persica</i> (peach)	6.0-7.5	3
25 <i>Antirrhinum majus</i> (snapdragon)	6.0-7.5	3	58 <i>Pyrus communis</i> (pear)	6.0-7.5	3
26 <i>Beta vulgaris</i> (common beet)	6.0-7.5	3	59 <i>Quercus alba</i> (white oak)	6.0-8.0	1
27 <i>Betula lenta</i> (sweet birch)	4.5-6.0	4	60 <i>Raphanus sativus</i> (garden radish)	5.5-7.0	4
28 <i>Capsicum frutescens</i> (bush red pepper)	5.5-7.0	1	61 <i>Rhododendron obtusum amoenum</i> (amoena azalea)	4.5-6.0	3
29 <i>Carya ovata</i> (shagbark hickory)	6.0-6.5	4	62 <i>Rosa</i> sp. (hybrid tea rose)	5.5-7.0	3
30 <i>Catalpa</i> spp. (cataiba)	6.0-7.5	4	63 <i>Salix</i> spp. (willow)	5.5-7.5	4
31 <i>Chrysanthemum morifolium</i> (florist's chrysanthemum)	6.0-7.5	3	64 <i>Solanum tuberosum</i> (potato)	5.0-6.5	2,3
			65 <i>Trifolium pratense</i> (red clover)	6.0-7.5	2,3
			66 <i>Ulmus americana</i> (American elm)	6.0-8.0	1
			67 <i>Vicia faba equina</i> (horsebean)	6.0-7.0	3
			68 <i>Vitis</i> spp. (grape)	6.0-8.0	1

Contributors: (a) Walker, Richard B., (b) Wherry, Edgar T., (c) Bennett, W. F.

References: [1] Bennett, W. F. 1953. Texas Agr. Expt. Sta. Ext. Serv. Leaflet 164. [2] Ignatieff, V. 1949. Food Agr. Organ. U. N. Agr. Studies 9:108. [3] Spurway, C. H. 1941. Mich. State Univ. Agr. Expt. Sta. Spec. Bull. 306. [4] Wherry, E. T. Unpublished. Univ. Pennsylvania, Philadelphia, 1954.

## 120. SHADE TOLERANCE: VASCULAR PLANTS

**Tolerance** (column B): T = highly tolerant; t = moderately tolerant; I = intermediate; f = moderately intolerant; T = highly intolerant.

Species (Common Name)	Toler- ance	Ref- erence	Species (Common Name)	Toler- ance	Ref- erence
(A)	(B)	(C)	(A)	(B)	(C)
Pteridophyta			19 <i>Tradescantia virginiana</i> (Virginia spiderwort)	T	7
Gymnospermae			Dicotyledoneae		
1 <i>Adiantum pedatum</i> (American maidenhair)	T	4	20 <i>Acer saccharinum</i> (silver maple)	t	1
2 <i>Equisetum hyemale</i> (scouring rush)	T	2	21 <i>Beta vulgaris</i> (common beet)	I	6
3 <i>Lycopodium lucidulum</i> (shining club moss)	T	2	22 <i>Betula lenta</i> (sweet birch)	I	1
4 <i>Polypodium virginianum</i> (rock polypody)	t	4	23 <i>Carya illinoensis</i> (pecan)	f	1,8
5 <i>Abies concolor</i> (white fir)	t	1,4	24 <i>Catalpa bignonioides</i> (southern catalpa)	f	2
6 <i>Ginkgo biloba</i> (ginkgo)	f	2	25 <i>Chrysanthemum</i> spp. (chrysanthemum)	I	7
7 <i>Juniperus virginiana</i> (eastern red cedar)	f	1,8	26 <i>Cornus florida</i> (flowering dogwood)	T	1
8 <i>Larix</i> spp. (larch)	T	1,8	27 <i>Digitalis purpurea</i> (common foxglove)	t	2
9 <i>Picea glauca</i> (white spruce)	t	1,8	28 <i>Fagus grandifolia</i> (American beech)	T	1,8
10 <i>Pinus strobus</i> (eastern white pine)	I	1,8	29 <i>Fraxinus americana</i> (white ash)	I	1,2
11 <i>Sequoia gigantea</i> (giant sequoia)	I	1	30 <i>Ilex opaca</i> (American holly)	T	1
12 <i>Taxodium distichum</i> (bald cypress)	I	1,8	31 <i>Juglans nigra</i> (black walnut)	f	1,2
13 <i>Taxus canadensis</i> (Canada yew)	T	4	32 <i>Lactuca sativa</i> (lettuce)	f	6
14 <i>Thuja occidentalis</i> (northern white cedar)	t	1,8	33 <i>Lycopersicon esculentum</i> (tomato)	I	6
15 <i>Tsuga canadensis</i> (eastern hemlock)	T	1,8	34 <i>Malus coronaria</i> (sweet crab apple)	I	2
Monocotyledoneae			35 <i>Phlox divaricata</i> (Sweet William phlox)	T	7
16 <i>Iris cristata</i> (crested iris)	I	4	36 <i>Populus deltoides</i> (eastern poplar)	T	1,8
17 <i>Lilium</i> spp. (lily)	f	5	37 <i>Prunus serotina</i> (black cherry)	f	1
18 <i>Poa trivialis</i> (roughstalk bluegrass)	t	3	38 <i>Quercus alba</i> (white oak)	I	1,8
			39 <i>Rhododendron</i> spp. (rhododendron)	T	4
			40 <i>Ribes americanum</i> (American black currant)	T	2
			41 <i>Salix</i> spp. (willow)	T	1,2
			42 <i>Solanum melongena</i> (eggplant)	I	6
			43 <i>Ulmus americana</i> (American elm)	I	1,2

**Contributors:** (a) Clapp, Grace L., (b) Kramer, Paul J., (c) Roe, Eugene I.

**References:** [1] Baker, F. S. 1949. *J. Forestry* 47:179. [2] Clapp, G. L. Unpublished. Windsor, Connecticut, 1952. [3] Curtis, R. W., and J. F. Cornman. 1941. *N. Y. State Agr. Expt. Sta. (Geneva) Bull.* 465. [4] Morse, H. K. 1962. *Gardening in the shade*. Rev. ed. Scribner's Sons, New York. [5] Taylor, N. 1961. *Encyclopedia of gardening*. Houghton Mifflin, Boston. [6] Went, F. W. 1946. *Proc. Am. Soc. Hort. Sci.* 48:374. [7] Williams, F. R. 1949. *J. N. Y. Botan. Garden* 50:201. [8] Zon, R., and H. S. Graves. 1911. *U.S. Dept. Agr. Forest Serv. Bull.* 92.

## 121. EFFECT OF LIGHT ON DEVELOPMENT: ANGIOSPERMS

### Part I. VARIOUS WAVELENGTHS

Data present the effectiveness of brief interruption of the dark period to control flowering and certain vegetative expressions. Plants were grown under radiation from carbon arc and incandescent filament lamps for a daily period of approximately 12 hours. Midpoint in the dark period they were exposed to radiation of known energy and wavelength.

Effect and Species (Common Name)	Relative Energy Normalized to Maximum Response at Wavelength (Angstrom Units) of											Con- version Factor <sup>1</sup>	Ref- erence
	4400	4800	5000	5200	5400	5600	5800	6200	6600	6800	7000		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)
Inhibition of flowering													
1 <i>Glycine soja</i> (soybean)	18	27	17	6	3.7	2	1.3	I	1.3	1.6	3.5	30	8,9
2 <i>Xanthium pensylvanicum</i> (cocklebur)	125	173	92	40	8	5.4	2.6	1 <sup>2</sup>	1.5	3.1	7	40	3,8

<sup>1</sup>/ Relative energy (columns B-L) may be converted to kiloergs per sq cm by multiplying by the appropriate factor (column M). <sup>2</sup>/ 7200-7600 Å reverses the response caused by red (6200-6600 Å).

*continued*

121. EFFECT OF LIGHT ON DEVELOPMENT: ANGIOSPERMS

Part I. VARIOUS WAVELENGTHS

Effect and Species (Common Name)	Relative Energy Normalized to Maximum Response at Wavelength (Ångstrom Units) of											Con- version Factor <sup>1</sup>	Ref- er- ence
	4400	4800	5000	5200	5400	5600	5800	6200	6600	6800	7000		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)
3 Promotion of flowering <i>Hordeum vulgare</i> (barley)	218	185	85	35	4	1.8	1.3	1 <sup>2</sup>	1.5	4	7	35	1,5,6
4 <i>Hyoscyamus niger</i> (black henbane)	...	...	...	...	4	1.8	1.3	1 <sup>2</sup>	1.5	4	7	300	5-7
5 Promotion of germination <i>Lactuca sativa</i> (lettuce)	...	...	...	...	...	18	10	3 <sup>2</sup>	1	1.2	50	2	4
6 Promotion of leaf elongation <i>Pisum sativum</i> (garden pea)	100	190	200	95	24	10	6.5	1	1	1	1.3	0.16	9
7 Inhibition of stem elongation <i>Hordeum vulgare</i> (barley)	250	...	200	40	20	5	2	1.3 <sup>2</sup>	1	2	6	100	2,5,6
8 Production of pigmentation <i>Lycopersicon esculentum</i> (tomato)	30	30	30	30	20	10	3	1 <sup>2</sup>	1	1.2	7	200	10

<sup>1/</sup> Relative energy (columns B-L) may be converted to kiloergs per sq cm by multiplying by the appropriate factor (column M). <sup>2/</sup> 7200-7600 Å reverses the response caused by red (6200-6600 Å).

Contributor: Downs, R. J.

References: [1] Borthwick, H. A., S. B. Hendricks, and M. W. Parker. 1948. *Botan. Gaz.* 110:103. [2] Borthwick, H. A., S. B. Hendricks, and M. W. Parker. 1951. *Ibid.* 113:95. [3] Borthwick, H. A., S. B. Hendricks, and M. W. Parker. 1952. *Proc. Natl. Acad. Sci. U.S.* 38(1):929. [4] Borthwick, H. A., et al. 1952. *Ibid.* 38(8):662. [5] Downs, R. J. 1955. *Plant Physiol.* 30:468. [6] Downs, R. J. 1956. *Ibid.* 31:279. [7] Parker, M. W., S. B. Hendricks, and H. A. Borthwick. 1950. *Botan. Gaz.* 111:242. [8] Parker, M. W., et al. 1945. *Science* 102:152. [9] Parker, M. W., et al. 1946. *Botan. Gaz.* 108:1. [10] Piringer, A. A., and P. H. Heinze. 1954. *Plant Physiol.* 29:467.

Part II. VARIOUS EXPOSURES

Species (Common Name)	Beginning of Test	Photoperiodic Classification	Light Exposure hr	Development		
				Budding da	Flowering da	Height cm
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 <i>Agrostis nebulosa</i> (cloud bent grass)	May 14	Long day	13.0	49	58	25.4
2			13.5	48	58	33.0
3			14.0	37	45	27.9
4			14.5	35	41	40.6
5			24.0	34	48	27.9
6 <i>Antirrhinum majus</i> (snapdragon)	May 31	Indeterminate	10.0	28	45	40.6
7			12.0	24	35	30.5
8			13.0	20	33	35.6
9			14.0	20	35	40.6
10			24.0	20	33	33.0
11 <i>Chrysanthemum</i> sp. (chrysanthemum) <sup>1</sup>	.....	Indeterminate	10.0	21	54	40.6
12			12.0	24	54	33.0
13			12.5	24	54	40.6
14			13.0	25	59	35.6
15			13.5	24	56	50.8
16			14.0	38	73	48.3
17			14.5	38	78	50.8
18			24.0	46	71	45.7
19 <i>Convolvulus sepium</i> (hedge glorybind)	May 19	Long day	13.5	44	59	101.6
20			24.0	35	49	114.3
21 <i>Glycine soja</i> (soybean)	May 25	Short day	10.0	20	23	22.9
22			12.0	21	27	35.6

<sup>1/</sup> Yellow Normandie variety.

*continued*

121. EFFECT OF LIGHT ON DEVELOPMENT: ANGIOSPERMS

Part II. VARIOUS EXPOSURES

	Species (Common Name)	Beginning of Test	Photoperiodic Classification	Light Exposure hr	Development		
					Budding da	Flowering da	Height cm
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
23	<i>Glycine soja</i> (soybean)	May 25	Short day	12.5	24	27	45.7
24				13.0	25	31	40.6
25				13.5	34	37	48.3
26				14.0	42	48	68.6
27				14.5	50	60	76.2
28				24.0	81	90	96.5
29	<i>Hibiscus syriacus</i> (shrub althea)	March 27	Long day	12.5	95	117	77.5
30				13.0	96	123	120.7
31				13.5	95	126	115.6
32				14.0	103	136	111.8
33				14.5	106	144	132.1
34				24.0	103	130	111.8
35	<i>Nicotiana tabacum</i> (common tobacco) <sup>2</sup>	May 14	Indeterminate	10.0	18	30	101.6
36				12.0	19	30	114.3
37				13.0	18	30	114.3
38				14.0	18	30	111.8
39				24.0	18	32	106.7
40				<i>N. tabacum</i> <sup>3</sup>	June 7	Short day	10.0
41	12.0	...	73				162.6
42	13.0	...	73				154.9
43	14.0	...	88				167.6
44	24.0	...	90				149.9
45	<i>Oenothera speciosa</i> (white evening prim-rose)	April 16	Indeterminate				10.0
46				12.0	13	35	.....
47				12.5	13	36	14.0
48				13.0	15	50	25.4
49				13.5	15	10	22.9
50				14.0	15	37	22.9
51				14.5	15	40	22.9
52				24.0	15	41	22.9
53	<i>Phaseolus coccineus</i> (scarlet runner bean)	May 21	Indeterminate	10.0	13	55	66.0
54				12.0	14	22	53.3
55				12.5	14	21	78.7
56				13.0	14	22	78.7
57				13.5	14	25	116.8
58				14.0	14	28	167.6
59				14.5	20	28	116.8
60				24.0	14	32	142.2
61	<i>Salix humilis</i> (prairie willow)	April 10	Short day	10.0	68	...	.....
62				12.0	55	...	.....
63				12.5	60	88	55.9
64				13.0	61	...	.....
65				13.5	61	...	.....
66				14.0	61	76	.....

<sup>1</sup>/a/ Extra Early variety. <sup>1</sup>/3/ Maryland Mammoth variety.

Contributor: Williams, Bert C.

Reference: Garner, W. W., and H. A. Allard. 1940. U.S. Dept. Agr. Tech. Bull. 727.



## 122. PHOTOPERIOD, WITH TEMPERATURE INTERACTIONS, FOR FLOWERING: ANGIOSPERMS

Varietal differences account for multiple photoperiodic classifications (column B); the common classification has been given priority in listing. Classification is followed in parentheses by the light period for flowering (>12 hr should be interpreted as 12 hours or more, <12 hr as up to 12 hours). **Temperature Interactions** (column C): Th = photoperiodic response occurring at relatively high temperatures (plant may also flower at other day lengths at lower temperatures), or reproductive development promoted by high temperatures during photoperiodic induction; Tl = photoperiodic response occurring at relatively low temperatures (plant may also flower at other day lengths at higher temperatures), or reproductive development promoted by low temperatures during photoperiodic induction; Tp = thermoperiodic (i.e., development affected by alternation of temperature between day and night periods); Tq = quantitative effect of temperature on critical day length (i.e., an increase in temperature lowers the minimum limits for long-day plants and raises the maximum limits for short-day plants), or on the degree of photoperiodic response; Ve = vernalization essential for complete reproductive development (or other low-temperature preconditioning of embryo plants, seedlings, buds, or plants), prior to photoperiodic induction; Va = vernalization not essential but promotes reproductive development; Vo = vernalization not effective. For additional information, consult references 5, 15, 30, 35, 45, and 54.

	Species (Common Name)	Photoperiodic Class and Light Period	Temperature Interactions	Reference
	(A)	(B)	(C)	(D)
Monocotyledoneae				
1	<i>Allium cepa</i> (garden onion)	Long day favorable; short day favorable; day neutral	Tl	24,31,46
2	<i>Avena sativa</i> (common oat)	Long day required (>9 hr)	Tl; Va (for winter varieties); Vo (for spring varieties)	24
3	<i>Hordeum vulgare</i> (barley)			
	Spring	Long day favorable	Vo	40,41
4	Winter	Long day required (>12 hr)	Va (7°-9°C)	12,24
5	<i>Oryza sativa</i> (rice)			
	Summer	Day neutral	Th	47,48
6	Winter	Short day required (<12 hr)	Th	47,48
7	<i>Phleum pratense</i> (timothy)	Long day required (>12 hr)	.....	3,52
8	<i>Poa pratensis</i> (Kentucky bluegrass)	Long day favorable	Th; Tl (day neutral or short day favorable); Ve	3,37,42,49
	<i>Triticum aestivum</i> (wheat)			
9	Spring	Long day favorable	Vo	24,33,34,55
10	Winter	Long day required (>12 hr)	Va	24,33,34,55
11	<i>Zea mays</i> (corn)	Day neutral; short day required	.....	25,32
Dicotyledoneae				
12	<i>Antirrhinum majus</i> (snapdragon)	Long day favorable	Th; Tl (day neutral)	29,39
13	<i>Beta saccharifera</i> (sugar beet)	Long day required	Tl (7°-9°C); Ve	22
14	<i>B. vulgaris</i> (common beet)	Long day favorable	Th; Tl (long day required)	41
15	<i>Capsicum frutescens</i> (bush red pepper)	Day neutral; short day favorable	Tp	14,18,20
16	<i>Chrysanthemum maximum</i> (Pyrenees chrysanthemum)	Long day required	.....	41
17	<i>Cucumis sativus</i> (cucumber)	Day neutral	.....	16,51
18	<i>Daucus carota</i> (carrot)	Day neutral	Ve (4°-10°C)	44
19	<i>Digitalis purpurea</i> (common foxglove)	Long day favorable	Ve	8
20	<i>Fagopyrum esculentum</i> (buckwheat)	Day neutral	.....	6,24
21	<i>Fragaria chiloensis</i> (chiloe strawberry)	Short day required (<10 hr)	Tq	17,26
22	Everbearing	Long day favorable; day neutral	.....	17
23	<i>Glycine soja</i> (soybean)	Short day required to short day favorable	Th; Tq	11,23,36
24	<i>Gossypium hirsutum</i> (upland cotton)	Day neutral; short day favorable	Tq	10,28
25	<i>Helianthus annuus</i> (common sunflower)	Short day favorable; day neutral	.....	19
26	<i>Ilex aquifolium</i> (English holly)	Day neutral	.....	43
27	<i>Ipomoea batatas</i> (sweet potato)	Short day required; short day favorable	.....	32
28	<i>Lactuca sativa</i> (lettuce)	Long day favorable	Th; Tl (day neutral)	7,13,50
29	<i>Lycopersicon esculentum</i> (tomato)	Day neutral; long day favorable; short day favorable	Tp	1,53

/-/ = applicable to most varieties.

continued

## 122. PHOTOPERIOD, WITH TEMPERATURE INTERACTIONS, FOR FLOWERING: ANGIOSPERMS

Species (Common Name)	Photoperiodic Class and Light Period	Temperature Interactions	Reference
(A)	(B)	(C)	(D)
Dicotyledoneae			
30 <i>Medicago sativa</i> (alfalfa)	Long day favorable	Th; Tl (day neutral) <sup>2</sup>	42
31 <i>Nicotiana tabacum</i> (common tobacco)	Day neutral <sup>1</sup>	.....	23
32 Common tobacco (Havana)	Long day favorable	.....	41
33 Common tobacco (Maryland Mammoth)	Short day required (<14 hr)	Th; <13°C (day neutral)	23,42
34 <i>Oenothera biennis</i> (common evening primrose)	Long day favorable	Tl	42
35 <i>Phaseolus vulgaris</i> (kidney bean)	Day neutral; short day re- quired <sup>9</sup>	.....	4,23,32
36 <i>Phlox paniculata</i> (summer phlox)	Long day required	Th	41
37 <i>Pisum sativum</i> (garden pea)	Day neutral; long day fa- vorable	.....	9,42
38 <i>Raphanus sativus</i> (garden radish)	Long day required	.....	23,38
39 <i>Rhododendron</i> sp. (rhododendron)	Day neutral	.....	43
40 <i>Solanum tuberosum</i> (potato)	Long day favorable; short day favorable; day neutral	.....	2,24,27
41 <i>Trifolium pratense</i> <sup>4</sup> (red clover)	Long day required (>12 hr)	.....	52
42 <i>Vicia faba</i> (broad bean)	Day neutral	Va	21

/1/ Data applicable to most varieties. /2/ Vegetative in warm nights. /3/ Photoperiod influences fruit development, but floral initiation is not affected. /4/ English Montgomery variety; for American Medium, long day favorable (>9 hr).

**Contributors:** (a) Greulach, Victor A., (b) Cooper, J. P., and Calder, D. M., (c) Roberts, R. H., and Struckmeyer, Burdean E., (d) Hagen, Charles W., Jr.

**References:** [1] Adams, J. 1924. *Am. J. Botany* 11:229. [2] Allard, H. A. 1938. *J. Agr. Res.* 57:775. [3] Allard, H. A., and M. W. Evans. 1941. *Ibid.* 62:193. [4] Allard, H. A., and W. J. Zaumeyer. 1944. U.S. Dept. Agr. Tech. Bull. 867. [5] Altman, P. L., and D. S. Dittmer, ed. 1962. *Growth, including reproduction and morphological development.* Federation of American Societies for Experimental Biology. Washington, D. C. [6] Arthur, J. M., and J. D. Guthrie. 1927. *Mem. Hort. Soc. N. Y.* 3:73. [7] Arthur, J. M., J. D. Guthrie, and J. M. Newell. 1930. *Am. J. Botany* 16:338. [8] Arthur, J. M., and E. K. Harvill. 1941. *Contrib. Boyce Thompson Inst.* 12:111. [9] Aso, K., and U. Muari. 1924. *J. Sci. Agr. Soc. (Japan)* 254:31. [10] Berkeley, E. E. 1931. *Ann. Missouri Bot. Garden* 18:573. [11] Borthwick, H. A., and M. W. Parker. 1939. *Botan. Gaz.* 101:341. [12] Borthwick, H. A., M. W. Parker, and P. H. Heinze. 1941. *Ibid.* 103:326. [13] Bremer, A. H. 1931. *Gartenbauwissenschaft* 4:469. [14] Cochran, H. L. 1936. *Cornell Univ. Agr. Expt. Sta. Mem.* 190. [15] Cooper, J. P. 1960. *Herbage Abstr.* 30:71. [16] Danielson, L. L. 1944. *Plant Physiol.* 19:638. [17] Darrow, G. M., and G. F. Waldo. 1934. U.S. Dept. Agr. Tech. Bull. 453. [18] Dorland, R. E., and F. W. Went. 1947. *Am. J. Botany* 34:393. [19] Dyer, H. J., J. Skok, and N. J. Scully. 1959. *Botan. Gaz.* 121:50. [20] Eguchi, T. 1937. *Proc. Imp. Acad. (Tokyo)* 13:332. [21] Evans, L. T. 1959. *Ann. Botany (London), N.S.* 23:521. [22] Fife, J. M., and C. Price. 1953. *Plant Physiol.* 28:475. [23] Garner, W. W., and H. A. Allard. 1920. *J. Agr. Res.* 18:553. [24] Garner, W. W., and H. A. Allard. 1923. *Ibid.* 23:871. [25] Gerhard, E. 1940. *J. Landwirtsch.* 87:161. [26] Hartman, H. T. 1947. *Plant Physiol.* 22:407. [27] Jones, H. A., and H. A. Borthwick. 1938. *Am. Potato J.* 15:331. [28] Konstantinov, P. N. 1938. U.S. Office Expt. Sta. Record 78:170. [29] Laurie, A. 1930. *Proc. Am. Soc. Hort. Sci.* 27:319. [30] Leopold, A. C. 1951. *Quart. Rev. Biol.* 26:247. [31] Magruder, R., and H. A. Allard. 1937. *J. Agr. Res.* 54:719. [32] McClelland, T. B. 1928. *Ibid.* 37:603. [33] McKinney, H. H., and W. J. Sandow. 1933. *J. Heredity* 24:169. [34] McKinney, H. H., and W. J. Sandow. 1935. *J. Agr. Res.* 51:621. [35] Murneek, A. E., and R. O. Whyte, ed. 1948. *Vernalization and photoperiodism.* Chronica Botanica, Waltham, Mass. [36] Parker, M. W., and H. A. Borthwick. 1943. *Botan. Gaz.* 104:612. [37] Peterson, M. L., and W. E. Loomis. 1949. *Plant Physiol.* 24:31. [38] Plitt, T. M. 1932. *Ibid.* 7:337. [39] Post, K., and C. L. Weddle. 1940. *Proc. Am. Soc. Hort. Sci.* 37:1037. [40] Purvis, O. N. 1934. *Ann. Botany (London)* 48:919. [41] Roberts, R. H., and B. E. Struckmeyer. 1938. *J. Agr. Res.* 56:633. [42] Roberts, R. H., and B. E. Struckmeyer. 1939. *Ibid.* 59:699. [43] Roberts, R. H. Unpublished. Univ. Wisconsin, Madison, 1953. [44] Sakr, E. S., and H. C. Thompson. 1942. *Proc. Am. Soc.*

*continued*

**122. PHOTOPERIOD, WITH TEMPERATURE INTERACTIONS,  
FOR FLOWERING: ANGIOSPERMS**

Hort. Sci. 41:343. [45] Samygin, G. A. 1946. Tr. Inst. Fiziol. Rast. Akad. Nauk SSSR 3:129. [46] Scully, N. J., H. A. Borthwick, and M. W. Parker. 1945. Botan. Gaz. 107:52. [47] Sircar, S. M. 1946. Proc. Natl. Inst. Sci. India 12:191. [48] Sircar, S. M., and B. Pariji. 1945. Nature 155:395. [49] Sprague, V. G. 1948. J. Am. Soc. Agron. 40:144. [50] Thompson, H. C., and J. E. Knott. 1933. Proc. Am. Soc. Hort. Sci. 30:507. [51] Tiedjens, V. A. 1928. J. Agr. Res. 36:721. [52] Tincker, M. A. H. 1925. Ann. Botany (London) 39:721. [53] Went, F. W. 1945. Am. J. Botany 32:469. [54] Withrow, R. B., ed. 1959. Publ. Am. Assoc. Advan. Sci. 55. [55] Wort, D. J. 1941. Botan. Gaz. 102:725.

**123. FACTORS AFFECTING PROTOPLASMIC STREAMING: PLANTS**

Protoplasmic streaming as considered in this table includes shuttle-type flow of protoplasm in slime molds, protoplasmic streaming within rigid cell walls of Algae, and cyclosis in Monocotyledoneae. Many techniques were used to study streaming rates, and the results vary according to the methods and conditions of measurement. Values were interpolated where necessary from graphic and tabular data in the literature.

**PART I. TEMPERATURE**

	Species (Common Name)	Rate of Streaming, $\mu$ /sec								Refer- ence
		5°C	10°C	15°C	20°C	25°C	30°C	35°C	40°C	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Algae										
1	<i>Chara foetida</i> (stonewort)	20.4	25.6	37.6	41.8	51.5	76.3	92.7	71.5	4
2		11.2	24.5	40.9	56.2	74.4	90.0			3
3	<i>Nitella</i> sp. (nitella)	10.5	15.3	20.0	26.7	34.3	43.3	47.3	37.0	2
4	<i>N. mucronata</i> (nitella)	12.7	28.0	41.2	59.6	75.3	96.2	109.1		3
Monocotyledoneae										
5	<i>Elodea canadensis</i> (Canada waterweed)	4.8	7.7	8.8	10.5	11.5	13.3	16.7	0	4
6	<i>Vallisneria spiralis</i> (spiral wild celery)	4.7	8.7	15.1	20.0	26.3	31.2	38.5		4
	<i>Avena</i> sp. (oat), coleoptile									
7	90 hr old	3.4	4.8	7.0	8.2	8.3	8.5			1
8	200 hr old	3.6	5.7	7.9	10.4	12.8	15.8			

Contributor: Olson, Rodney A.

References: [1] Bottelier, H. P. 1934. Rec. Trav. Botan. Neerl. 31:474. [2] Ganong, W. F. 1908. A laboratory course in plant physiology. Ed. 2. H. Holt, New York. [3] Lambers, M. H. R. 1926. Dissertation. Utrecht Univ., Netherlands. [4] Velten, W. 1876. Flora (Jena) 59:209.

**Part II. SUDDEN CHANGES OF TEMPERATURE**

$R_0$  (column B) = average streaming rate ( $\mu$ /sec) at initial temperature.

	Initial Temp., °C	$R_0$	Final <sup>1</sup> Temp., °C	Rate of Streaming, $\mu$ /sec, at Specified Interval in minutes											Temperature Sensitivity <sup>2</sup>
				20	30	40	50	60	70	80	90	100	110	120	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)
<i>Nitella flexilis</i> (Pliant Nitella)															
1	13.8	28.5	24.5	55	55	55	55	55	55	55	55	55	55	55	
2	13.8	28.5	31.5	80	80	80	80	80	80	80	80	80	80	80	
3	13.8	28.5	37.5	99	99	99	99	99	99	99	99	99	99	99	
4	19	55	12.8	36	36	36	36	36	36	36	36	36	36	36	
5	22	62.5	12.8	36	36	36	36	36	36	36	36	36	36	36	
6	24.3	70	12.8	36	36	36	36	36	36	36	36	36	36	36	$\frac{100}{12} = 8.3$
7	26	74	12.8	24	28	32	34	36	36	36	36	36	36	36	

<sup>1</sup>/ Temperature prevailing during test. <sup>2</sup>/ Equals 100 divided by the maximum temperature difference (at which no rate lag occurs).

*continued*



**123. FACTORS AFFECTING PROTOPLASMIC STREAMING: PLANTS**

**Part II. SUDDEN CHANGES OF TEMPERATURE**

	Initial Temp., °C	R <sub>0</sub>	Final <sup>1</sup> Temp., °C	Rate of Streaming, μ/sec, at Specified Interval in minutes												Temperature Sensitivity <sup>2</sup>
				20	30	40	50	60	70	80	90	100	110	120		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	
<i>Nitella flexilis</i> (Pliant Nitella)																
8	25	86	20	71	71	71	71	71	71	71	71	71	71	71	$\frac{100}{6} = 16.6$	
9	27.5	93	20	64	68	69	70	71	71	71	71	71	71	71		
10	30	100	20	62	65	66	68	69	70	71	71	71	71	71		
11	35	114	20	45	50	52	54	56	58	60	62	63	64	65	$\frac{100}{9} = 11.1$	
12	30	100	25	83	83	83	83	83	83	83	83	83	83	83		
13	32.5	104	25	83	83	83	83	83	83	83	83	83	83	83		
14	33	106	25	83	83	83	83	83	83	83	83	83	83	83		
15	35	113	25	77	80	82	82	83	83	83	83	83	83	83		
16	38.5	125	25	71	74	76	77	78	80	81	81	82	83	83		
17	40	131	25	25	40	48	54	58	62	65	68	70	71	72		
<i>Avena</i> (Oat) Coleoptile																
18	21	12	13	6.2	6.4	6.6	6.6	6.6	6.6	6.6	6.6	6.6	6.6	6.6	$\frac{100}{7.5} = 13$	
19	24	12.8	13	0.1	2.1	3.2	4.4	5.1	5.8	6.3	6.9	7.5	7.6	7.6		
20	28	12.6	13	0	0.4	1.2	2.6	4.5	5.8	6.6	7.3	7.7	7.7	7.7	$\frac{100}{3.5} = 30$	
21	25	11	21	9.1	9.9	10.2	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.3		
22	27	10	21	4.5	9.9	11.9	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1		
23	36	3.4	21	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.2	0.4	1.4	3.6		

<sup>1</sup>/ Temperature prevailing during test. <sup>2</sup>/ Equals 100 divided by the maximum temperature difference (at which no rate lag occurs).

*Contributor:* Olson, Rodney A.

*Reference:* Romijn, G. 1931. Koninkl. Ned. Akad. Wetenschap. Proc., C, 34:1.

**Part III. LIGHT INTENSITY: AVENA COLEOPTILE**

Nonchlorophyll-containing cells of *Avena* coleoptile were observed in the orange-red, phototropically inactive spectral region after exposure to various doses of blue light (4,360 Å), at 23°C. Data show the fitness of the "product rule" (intensity x time), as well as the effect of total energy (dosage). **Reaction** (column D) = the algebraic sum of percent departures from normal rate; negative values indicate a decrease in streaming, and positive values, an increase.

	Illumination			Reac-tion <sup>2</sup>	Dosage <sup>1</sup> ergs per sq cm	Illumination			Reac-tion <sup>2</sup>	Dosage <sup>1</sup> ergs per sq cm	Illumination			Reac-tion <sup>2</sup>
	Intensity ergs per sq cm per sec	Time sec	(D)			Intensity ergs per sq cm per sec	Time sec	(D)			Intensity ergs per sq cm per sec	Time sec	(D)	
	(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)		
1	10	2	5	-29	14	110	11	10	-118	26	440	11	40	-84
2	12	23.6	0.5	-54	15	118	23.6	5	-92	27	472	23.6	20	-54
3	20	2	10	-49	16	120	2	60	-112	28	480	2	240	-107
4	20	5	4	-67	17	142	23.6	6	-97	29	600	5	120	-69
5	22	11	2	-74	18	160	5	32	-103	30	660	11	60	-22
6	24	23.6	1	-75	19	180	2	90	-160	31	746	23.6	32	-15
7	40	2	20	-72	20	189	23.6	8	-136	32	944	23.6	40	+32
8	47	23.6	2	-87	21	220	11	20	-169	33	1,320	11	120	+132
9	55	11	5	-70	22	236	23.6	10	-105	34	1,420	23.6	60	+42
10	66	11	6	-74	23	240	2	120	-118	35	2,100	23.6	90	+28
11	80	2	40	-91	24	360	2	180	-104	36	2,800	23.6	120	+26
12	88	11	8	-78	25	378	23.6	16	-81	37	4,200	23.6	180	+4
13	94	23.6	4	-77										

<sup>1</sup>/ Intensity (column B) x time (column C). <sup>2</sup>/ Average of 1-20 separate experiments for each energy value.

*Contributor:* Olson, Rodney A.

*Reference:* Bottelier, H. P. 1934. Rec. Trav. Botan. Neerl. 31:474.

*continued*



## 123. FACTORS AFFECTING PROTOPLASMIC STREAMING: PLANTS

### Part IV. VARIOUS WAVELENGTHS: AVENA COLEOPTILE

**Reaction** (column C) = the algebraic average of percent departures from normal rate; negative values indicate a decrease in streaming, and positive values, an increase.

Wavelength Å	Dosage <sup>1</sup>	Reaction	Equivalent Energy <sup>2</sup>	Spectral Sensitivity	Wavelength Å	Dosage <sup>1</sup>	Reaction	Equivalent Energy <sup>2</sup>	Spectral Sensitivity
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)
1 3,660	270	-46	16	6	7 5,460	230	-16	7	3
2	760	-85	90	12	8	240	-15	7	3
3 4,360	See Part III				9	1,790	-60	30	1.7
4 4,550	21	-15	7	33	10 5,780	9,000	+1		<1
5	67	-33	13	20	11 6,200	See Fn. 3	+1		<1
6	190	-90	95	50					

<sup>1</sup>/ Intensity x time. <sup>2</sup>/ Equivalent energy required at 4,360 Å for comparable effect. <sup>3</sup>/ Approximately 70 ergs per sq cm per sec.

*Contributor:* Olson, Rodney A.

*Reference:* Bottelier, H. P. 1934. Rec. Trav. Botan. Neerl. 31:474.

### Part V. OXYGEN

	Material	Temp. °C	O <sub>2</sub> Concentration		Streaming Rate μ/sec	% of Normal Rate		Refer- ence	
			% Saturation	ml/L		Streaming	Respiration		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
1	<i>Physarum polycephalum</i>	22	21	6.27	Normal	100	100	1	
2			2.4	0.72	Normal	....	60	1	
3			1.0	0.30	Reduced	....	30	1	
4			0.3	0.08	Reduced	....	9	1	
5			0.1	0.03	Reduced	....	3	1	
6			0	0	Reduced	0	0	1	
7	<i>Avena</i> sp., coleoptile	21	100	30.4	9.9	99	...	2	
8			100	27.8	10.1	132	...	2	
9			21	21	6.32	10.0	100	...	2
10			26	21	5.85	10.1	100	...	2
11		260 hr old	21	21	6.32	10.2	100	...	2
12				21	5.85	13.4	100	...	2
13				4	1.23	9.0	88	...	2
14				4	1.02	9.1	68	...	2
15		80 hr old	25	10.8-14.6 <sup>1</sup>	3.1-4.1	.....	100	100 <sup>2</sup>	3
16				14.6 <sup>1</sup> -21.0	4.1-5.7	.....	....	50 <sup>2</sup>	3
17		95 hr old	25	10.8-18.2 <sup>1</sup>	3.1-5.3	.....	100	100 <sup>2</sup>	3
18				18.8 <sup>1</sup> -21.0	5.3-5.7	.....	....	50 <sup>2</sup>	3
19		130 hr old	25	11.6-12.4 <sup>1</sup>	3.2-3.5	.....	100	100 <sup>2</sup>	3
20				12 <sup>1</sup> -21	3.5-5.7	.....	....	50 <sup>2</sup>	3

<sup>1</sup>/ Critical O<sub>2</sub> tension. <sup>2</sup>/ Rates are uniform within indicated O<sub>2</sub> ranges; a sharp change in rate occurs at the critical O<sub>2</sub> tension.

*Contributor:* Olson, Rodney A.

*References:* [1] Allen, P. J., and W. Price. 1950. Am. J. Botany 37:393. [2] Bottelier, H. P. 1935. Rec. Trav. Botan. Neerl. 32:287. [3] DuBuy, H. G., and R. A. Olson. 1940. Am. J. Botany 27:392.

## 124. FACTORS AFFECTING TRANSPIRATION RATES: ANGIOSPERMS

### Part I. VARIOUS CONDITIONS

Water loss was calculated from data obtained in experiments using intact plants growing in soil, unless otherwise stated. **Specifications** (columns B and C): SM = soil moisture; SWC = wilting coefficient of soil; EA = evaporation from atmometer; AT = air temperature; RH = relative humidity; WV = wind velocity; SS = possible sunshine; ES = evaporation from free-water surface; ST = soil temperature; WHC = soil moisture in percent of water-holding capacity on dry-weight basis; LIE = light intensity energy; LI = light intensity; SD = saturation deficit of air; RI = total radiation intensity.

	Species (Common Name)	Specifications		Water Loss mg/sq cm leaf surface/hr	Ref- er- ence
		Constants	Variables		
	(A)	(B)	(C)	(D)	(E)
1	<i>Aeonium haworthii</i>	1 plant	WV = 1,100 cm/sec	0.58	12
2	(Haworth aeonium)		WV = 570 cm/sec	0.21	
3	<i>Ananas comosus</i>	19 plants; 1.5 mo old	Hr ending 3 p.m.; AT = 31°C; RH = 59%; WV = 128 cm/sec; SS = 100%; EA = 3 cc/hr	1.96	5
4	(pineapple) <sup>1</sup>		Hr ending 4 a.m.; AT = 24°C; RH = 85%; WV = 6 cm/sec; SS = 0; EA = 0.5 cc/hr	0.03	
5	<i>Bellis perennis</i> (En- glish daisy)	3 plants; several yr old; AT = 20°-22°C; RH = 40- 65%; WV = 90 cm/sec; ES = 0.0209 g/sq cm	Continental type plants	15.3	13
6			Maritime type plants	12.2	
7	<i>Capsicum frutescens</i>	6 plants; 8 inches high	WHC = 50	3.5	3
8	(bush red pepper)		WHC = 25	1.1	
9	<i>Citrus limon</i> (lemon)	3 plants; 6± mo old	ST = 31°C	4.9	4
10			ST = 19°C	3.5	
11	<i>C. paradisi</i> (grape- fruit)	3 plants; 6± mo old	ST = 27°C	4.6	4
12			ST = 35°C	3.1	
13	<i>Euphorbia capitellata</i>	1 plant	10 a.m.-2 p.m.; ES = 0.050 g/sq cm	4.5	8
14	(head euphorbia)		9:30 p.m.-7 a.m.; ES = 0.010 g/sq cm	0.1	
15	<i>Gossypium herbaceum</i>	6± wk old	Untreated soil of low salt concentration	15.0	10
16	(Levant cotton)		0.8% calcium nitrate added to soil	1.6	
17	<i>Helianthus annuus</i>	5 plants; few wk old	Hr ending 1 p.m.; RI = 0.960 cal/sq cm/ min; AT = 23.9°C; RH = 42%; WV = 215 cm/sec	23.8	9
18	(common sunflower)		Hr ending 8 a.m.; RI = 0.223 cal/sq cm/ min; AT = 13.3°C; RH = 69%; WV = 4.5 cm/sec	2.7	
19		4 plants; 6 wk old; AT = 26.7°C; RH = 27%; WV = 19.0 cm/sec	ST = 37.8°C	27.2	2
20			ST = 2.2°C	4.7	
21	<i>Hieracium pilosella</i>	1 plant; LI = 30,000 lux	Hr ending 7 a.m.; AT = 19.2°C; SD = 7.1%	5.5	6
22	(mouse-ear hawk- weed)		Hr ending 4 p.m.; AT = 21.1°C; SD = 8.1%	3.5	
23	<i>Lychmis dioica</i>	3 plants; several yr old; AT = 19.5°-20.5°C; RH = 40-65%; WV = 90 cm/sec; ES = 0.0285 g/sq cm	Beech-forest type plants	10.5	13
24	(red campion)		Maritime type plants	6.3	
25	<i>Malus pumila</i> (com- mon apple)	2 plants; 11 yr old	Grown with straw mulch	17.2	3
26			Grown in sod without mulch	14.3	
27	<i>Nicotiana tabacum</i>	9 plants; 3-4 leaves; RH = 68%	Exposed to visible light: LIE = 0.72 cal/ sq cm/min; AT = 36.7°-37.8°C	36.4	1
28	(common tobacco)		Exposed to infrared light: LIE = 0.65 cal/ sq cm/min; AT = 22.8°-25.6°C	5.6	
29	<i>Rumex acetosa</i> (gar- den sorrel)	3 plants; several yr old; AT = 17°-24°C; RH = 40- 65%; WV = 90 cm/sec; ES = 0.0334 g/sq cm	Alpine type plants	28.0	13
30			Lowland type plants	11.6	
31	<i>Triticum aestivum</i>	6 plants; 1+ mo old	Grown in good loam soil	2.1	7
32	(wheat)		Grown in poor soil	1.1	
33	<i>Veronica beccabunga</i>	2 plants; LI = 30,000 lux	Hr ending 2 p.m.; AT = 21.0°C; SD = 10.0%	6.75	6
34	(beccabunga speed- well)		Hr ending 6 p.m.; AT = 21.5°C; SD = 10.6%	3.17	
35	<i>Zea mays</i> (corn)	1 plant; 2 ft high; 9 fully, 7 partly unfolded leaves; SM = 22% of dry wt; SWC = 15.1	2-hr periods ending 3 p.m.; EA = 7.7 cc/hr	28.29	11
36			2-hr periods ending 7 a.m.; EA = 0.8 cc/hr	0.86	

<sup>1/1</sup> Growing in nutrient solution.

continued

## 124. FACTORS AFFECTING TRANSPIRATION RATES: ANGIOSPERMS

### Part I. VARIOUS CONDITIONS

*Contributor:* Krauss, Beatrice

*References:* [1] Arthur, J. M., and W. D. Stewart. 1933. *Contrib. Boyce Thompson Inst.* 5:491. [2] Clements, F. E., and E. V. Martin. 1934. *Plant Physiol.* 9:621. [3] Cullinen, F. P. 1920. *Proc. Am. Soc. Hort. Sci.* 17:237. [4] Haas, A. R. C. 1936. *Calif. Citrograph* 21:479. [5] Krauss, B. H. 1930. M.S. Thesis. Univ. Hawaii, Honolulu. p. 82. [6] Lachenmeier, J. 1932. *Jahrb. Wiss. Botan.* 76:825. [7] Livingston, B. E. 1905. *Botan. Gaz.* 40:189. [8] Livingston, B. E. 1906. *Carnegie Inst. Wash. Publ.* 50:45. [9] Martin, E. V. 1935. *Plant Physiol.* 10:344. [10] Meyer, B. S. 1931. *Am. J. Botany* 18:85. [11] Miller, E. C. 1918. *J. Agr. Res.* 13:585. [12] Seybold, A. 1929. *Die physikalische Komponente der pflanzlichen Transpiration.* J. Springer, Berlin. p. 91. [13] Turesson, G. 1928. *Hereditas* 11:199.

### Part II. VARIATION IN SOIL CONDITIONS: CORN

	Soil Condition	Leaf Area per Plant sq cm	Dry Matter per Plant, g		per Plant kg	Water Loss	
			Grain	Total		per Gram Grain, g <sup>1</sup>	per Gram Dry Matter, g <sup>1</sup>
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
<b>8-Year Average</b>							
No manure added							
1	Infertile	2,948	17	72	38.3	8,263	531
2	Intermediate	3,573	26	108	51.3	2,485	489
3	Fertile	4,612	74	213	77.4	1,100	368
2.4 lb manure added/plant							
4	Infertile	4,244	52	180	68.9	1,518	396
5	Intermediate	4,638	67	207	75.6	1,173	369
6	Fertile	5,121	108	289	94.2	879	327
<b>3-Year Average</b>							
7	Too dry (50% saturation)	6,818	165	379	105.5	695	289
8	Favorable (70% saturation)	8,056	230	522	162.2	716	316
9	Too wet (95% saturation)	7,031	168	422	142.1	854	341

<sup>1/1</sup> Average of ratios.

*Contributor:* Kiesselbach, T. A.

*Reference:* Kiesselbach, T. A. 1929. *Proc. Intern. Congr. Plant Sci.*, 1st, Ithaca, 1926, 1:87.

### Part III. DIURNAL VARIATION: CORN

Left Column						Right Column							
Period	Temp. °C	RH <sup>1</sup> %	Wind mi/hr	Water Loss per Plant <sup>2</sup> g	Water Evapo- ration <sup>3</sup> g	Period	Temp. °C	RH <sup>1</sup> %	Wind mi/hr	Water Loss per Plant <sup>2</sup> g	Water Evapo- ration <sup>3</sup> g		
(A)	(B)	(C)	(D)	(E)	(F)	(A)	(B)	(C)	(D)	(E)	(F)		
1	8 a.m.	23.1	80	6	84	4.8	9	4 p.m.	32.8	50	9	343	22.2
2	9	25.5	73	7	111	7.7	10	5	32.2	51	8	294	18.5
3	10	27.6	67	8	167	11.6	11	6	31.2	53	8	217	14.1
4	11	29.3	63	8	215	15.0	12	7	28.4	58	7	132	10.2
5	12	30.8	58	9	279	19.2	13	8	27.6	64	7	64	6.4
6	1 p.m.	31.9	55	9	329	23.7	14	Day average	29.8	59.8	8	226	15.5
7	2	32.7	53	9	356	24.5	15	Night average	22.7	81.5	6	16	2.3
8	3	32.9	52	9	354	23.8							

<sup>1/1</sup> Mean relative humidity. <sup>2/2</sup> Water transpired from one plant. <sup>3/3</sup> From 36 square inches free-water surface, under identical conditions.

*continued*

## 124. FACTORS AFFECTING TRANSPIRATION RATES: ANGIOSPERMS

### Part III. DIURNAL VARIATION: CORN

Contributor: Kiesselbach, T. A.

Reference: Kiesselbach, T. A. 1916. Nebraska Univ. Agr. Expt. Sta. Res. Bull. 6.

### Part IV. ANNUAL VARIATION

Averages (column I) are comparable only when derived from data for identical years.

	Species	Common Name	Seasonal Water Requirement, g <sup>2</sup>						Average
			1912	1913	1914	1915	1916	1917	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
South Dakota [1]									
1	<i>Medicago sativa</i>	Alfalfa	735	735	1,038	696	673	866	790
2	<i>Setaria italica</i>	Foxtail millet	239	293	311	171	233	278	254
3	<i>Sorghum vulgare sudanense</i>	Sudan grass	....	....	....	272	314	344	310
4	<i>Triticum aestivum</i>	Wheat	463	436	528	333	352	487	433
Colorado [2]									
5	<i>Amaranthus retroflexus</i>	Redroot amaranth	....	320	306	229	340	307	300
6	<i>Avena sativa</i>	Common oat	449	617	615	445	809	636	595
7	<i>Bouteloua gracilis</i>	Blue grama	....	....	389	312	336	290	332
8	<i>Gossypium hirsutum</i>	Upland cotton	488	657	574	443	612	522	549
9	<i>Hordeum vulgare</i>	Barley	443	....	501	404	664	522	507
10	<i>Medicago sativa</i>	Alfalfa	657	834	890	695	1,047	822	824
11	<i>Secale cereale</i>	Rye	....	....	622	469	800	625	629
12	<i>Setaria italica</i>	Foxtail millet	187	286	295	202	367	284	273
13	<i>Sorghum vulgare sudanense</i>	Sudan grass	....	....	394	260	426	378	365
14	<i>Triticum aestivum</i>	Wheat	394	496	518	405	636	471	487
15	<i>Vigna sinensis</i>	Cowpea	....	571	659	413	767	481	578
16	<i>Zea mays</i>	Corn	280	399	368	253	495	346	357

/1/ Ratio of weight of water transpired to weight of dry matter produced.

Contributor: Bailey, Lowell F.

References: [1] Dillman, A. C. 1931. J. Agr. Res. 42:187. [2] Shantz, H. L., and I. N. Piemeisel. 1927. Ibid. 34:1093.

## 125. FACTORS AFFECTING OSMOTIC POTENTIAL: VASCULAR PLANTS

Osmotic pressure, a direct function of cell sap concentration, causes movement of water into roots and throughout plant. Osmotic potential is the maximum pressure that could be developed under ideal conditions. Values are given in atmospheres.

### Part I. SPECIES VARIATION: LEAVES

Species (Common Name)			Osmotic Potential	Ref-er-ence	Species (Common Name)			Osmotic Potential	Ref-er-ence
(A)	(B)	(C)	(B)	(C)	(A)	(B)	(C)	(B)	(C)
1	<i>Acer rubrum</i> (red maple)		11.2-16.7	3	10	<i>Gossypium</i> sp. (cotton)		22.0	1
2	<i>Arctium minus</i> (smaller burdock)		9.8-13.7	3	11	<i>Helianthus annuus</i> (common sunflower)		13.8-18.0	3
3	<i>Beta vulgaris</i> (common beet)		14.0	1	12	<i>Impatiens biflora</i> (spotted snapweed)		4.6-8.4	3
4	<i>Betula lutea</i> (yellow birch)		12.6-16.0	3	13	<i>Iris germanica</i> (German iris)		13.1	2
5	<i>Chenopodium album</i> (lamb's-quarters)		13.2	3	14	<i>Juglans nigra</i> (black walnut)		12.6-18.3	3
6	<i>Citrus limon</i> (lemon)		15.1-21.4	3	15	<i>Liquidambar styraciflua</i> (American sweet gum)		13.3-15.5	3
7	<i>Cornus florida</i> (flowering dogwood)		11.1-16.7	3	16	<i>Liriodendron tulipifera</i> (yellow poplar)		11.3-16.4	3
8	<i>Fraxinus americana</i> (white ash)		16.4	3	17	<i>Osmunda cinnamomea</i> (cinnamon fern)		9.8	3
9	<i>Galium aparine</i> (catchweed bedstraw)		9.6	3					

continued



## 125. FACTORS AFFECTING OSMOTIC POTENTIAL: VASCULAR PLANTS

### Part I. SPECIES VARIATION: LEAVES

Species (Common Name)	Osmotic Potential	Ref- er- ence	Species (Common Name)	Osmotic Potential	Ref- er- ence
(A)	(B)	(C)	(A)	(B)	(C)
18 <i>Phytolacca americana</i> (pokeberry)	8.5-9.5	3	26 <i>Quercus alba</i> (white oak)	15.8-18.4	3
19 <i>Picea engelmanni</i> (Engelmann spruce)	11.5-23.5	3	27 <i>Q. coccinea</i> (scarlet oak)	19.1	3
20 <i>Pinus</i> spp. (pine)	16.0-18.4	3	28 <i>Robinia pseudoacacia</i> (black locust)	9.8-14.3	3
21 <i>Pisum sativum</i> (garden pea)	9.2	5	29 <i>Salix alba</i> (white willow)	12.3-14.2	3
22 <i>Platanus occidentalis</i> (American sycamore)	13.5	3	30 <i>Solidago</i> sp. (goldenrod)	10.3	3
23 <i>Poa pratensis</i> (Kentucky bluegrass)	12.6-18.6	4	31 <i>Taraxacum officinale</i> (dandelion)	8.5-10.8	3
24 <i>Populus alba</i> (white poplar)	19.7-20.1	3	32 <i>Typha latifolia</i> (cattail)	9.7-11.8	3
25 <i>P. deltoides</i> (eastern poplar)	21.3	4	33 <i>Verbascum thapsus</i> (flannel mullein)	8.0-10.0	3
			34 <i>Xanthium</i> sp. (cocklebur)	8.4-10.7	3

Contributors: (a) Anderson, Donald B., (b) Howell, Robert W.

References: [1] Bennet-Clark, T. A., and D. Bexon. 1940. *New Phytologist* 39:337. [2] Dixon, H. H. 1914. *Transpiration and the ascent of sap in plants*. Macmillan, New York. [3] Harris, J. A. 1934. *The physico-chemical properties of plant saps in relation to phytogeography*. Univ. Minnesota Press, Minneapolis. [4] Meyer, B. S., and D. B. Anderson. 1939. *Plant physiology*. Van Nostrand, New York. [5] Thatcher, F. S. 1939. *Am. J. Botany* 26:449.

### Part II. PHYSICAL AND ENVIRONMENTAL VARIATION

Species (Common Name)	Plant Part	Specification	Osmotic Potential	Ref- er- ence	Species (Common Name)	Plant Part	Specification	Osmotic Potential	Ref- er- ence	
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	
Temperature										
1	<i>Potamogeton crispus</i> (curly pondweed)	Leaves 0°C	14.3	1	25	<i>Ambrosia trifida</i> (giant ragweed)	2nd leaf from top	6 a.m.	12.9	
2		13°C	13.5		26		10 a.m.	15.0		
3		23°C	12.7		27		2 p.m.	17.1		
4		30°C	11.1		28		5 p.m.	15.7		
				29	8 p.m.		14.9			
Height on Tree										
5	<i>Juglans nigra</i> (black walnut)	Leaves 2.4 cm	16.8	2	30		Lowest leaf	6 a.m.	10.1	
6		6.3 cm	17.8		31			10 a.m.	13.0	
7		9.7 cm	18.2		32			2 p.m.	15.9	
8		11.6 cm	17.2		33			5 p.m.	14.3	
9		13.4 cm	18.3		34	8 p.m.		14.0		
10	15.9 cm	18.3								
Distance from Growing Point										
11	<i>Vicia faba</i> (broad bean)	Root 1.5 mm	11.3	7	35	<i>Andropogon scoparius</i> (little blue-stem)		Shoots	12 noon	23.0
12		3.0 mm	11.7		36			2 p.m.	25.0	
13		5.0 mm	12.7		37			4 p.m.	27.0	
14		8.0 mm	9.8		38		6 p.m.	26.0		
				39	8 p.m.		21.5			
				40	10 p.m.		20.5			
				41	4 a.m.		21.0			
				42	8 a.m.		25.0			
Diurnal										
15	<i>Ambrosia trifida</i> (giant ragweed)	Top leaf 6 a.m.	12.5	3	43		<i>Rumex patientia</i> (patience dock)	Leaf guard cells	8:30 a.m.	16.6
16		10 a.m.	15.3		44	1:30 p.m.		21.0		
17		2 p.m.	17.4		45	3:30 p.m.		20.2		
18		5 p.m.	16.5		46	7:30 p.m.		13.2		
19		8 p.m.	16.3		47	Leaf subsidiary cells		8:30 a.m.	16.6	
20		1st leaf from top	6 a.m.		12.5	48		1:30 p.m.	16.6	
21			10 a.m.		15.6	49		3:30 p.m.	17.8	
22			2 p.m.		17.2	50		7:30 p.m.	15.5	
23			5 p.m.		15.7	51		Leaf epidermal cells	8:30 a.m.	14.4
24			8 p.m.		14.9	52		1:30 p.m.	14.4	
					53	3:30 p.m.	14.4			
				54	7:30 p.m.	13.2				

1/1 During extreme drought.

continued

## 125. FACTORS AFFECTING OSMOTIC POTENTIAL: VASCULAR PLANTS

### Part II. PHYSICAL AND ENVIRONMENTAL VARIATION

Species (Common Name)	Plant Part	Specification	Osmotic Potential	Reference	Species (Common Name)	Plant Part	Specification	Osmotic Potential	Reference		
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)		
Seasonal					100	<i>Triticum aestivum</i> (wheat)	Entire plant	8 da	6.0	12	
55	<i>Linnaea borealis</i> (twin-flower)	Oct	19.6	5	101			9 da	5.4		
56		Dec	25.0	102	10 da			5.1			
57		Mar	25.6	103	13 da			4.9			
58		May	14.3	104	15 da			5.4			
59		<i>Picea glauca</i> (white spruce)	Oct	17.1	5	105	25 da	6.0			
60	Dec		20.3	106	Shoots	10 da	6.56				
61	Jan		20.0	107	15 da	7.04					
62	Mar		24.9	108	Roots	10 da	3.06				
63	Apr		20.1	109	15 da	3.03					
64	<i>Populus tremuloides</i> (quaking aspen)	May	21.0	5	Tissue						
65		June	19.7		110	<i>Castanea sativa</i> (European chestnut)	Sieve-tubes	15.6-17.1	8		
66		Oct	15.0		111		Roots	Cambium		9.7-11.0	
67		Dec	16.8		112	Wood	6.3				
68		Jan	16.2		113	Stems	Cambium	11.1-12.9			
69	Feb	13.7	114	Wood	7.5-11.8						
70	Mar	17.0	115	<i>Fagus sp.</i> (beech)	Leaves	Epidermis	15.0	11			
71	May	10.6	116			Spongy parenchyma	22.4				
72	<i>Pyrola rotundifolia</i> (European pyrola)	Dec	24.6		5	Palisade parenchyma	37.7				
73		Feb	23.9		117	Stems	Cambium		24.6		
74		Apr	17.2		118		Xylem parenchyma		36.6		
75	June	12.6	119	Wood lays	35.2						
Soil Water Content					120	<i>Helleborus sp.</i> (hellebore)	Leaves	Cortex	26.1	11	
76	<i>Zea mays</i> (corn)	Shoots	31%	22.1	4			Phloem parenchyma	22.5		
77			16%	24.4	121			Stems	Cambium		21.9
78			14%	25.0	122				Xylem parenchyma		22.2
79	11%	26.5	123	Pith	20.6						
80	Roots	31%	5.9	124	Cortex	20.8					
81			16%	7.8	125	Pulvini	Pith	6.5			
82			14%	9.2	126	<i>Tradescantia sp.</i> (spiderwort)	Starch sheath	7.7			
83	11%	12.0	127	<i>Urtica sp.</i> (nettle)	Cortex		6.3				
Osmotic Concentration of Soil Solution					130		Leaves	Epidermis	18.8	11	
84	<i>Zea mays</i> (corn)	Shoots	1.2 atm		6.2	6		Spongy parenchyma	24.7		
85			2.0 atm	7.1	131	Palisade parenchyma		37.7			
86			3.4 atm	7.0	132	Stems	Cambium	21.5			
87			5.0 atm	7.2	133		Xylem parenchyma	23.0			
88			7.2 atm	7.3	134		Pith	18.7			
89	Roots	1.2 atm	4.6	135	Cortex	19.2					
90			2.0 atm	5.5	136	Phloem parenchyma	20.4				
91			3.4 atm	6.6	137	Age					
92	5.0 atm	7.5	138	<i>Triticum aestivum</i> (wheat)	Entire plant	2 da	10.7	12			
93	7.2 atm	8.2	139			3 da	9.8				
94	2 da	10.7	12			4 da	8.0				
95	3 da	9.8	12			5 da	7.8				
96	4 da	8.0	12			6 da	7.1				
97	5 da	7.8	12	7 da	6.6						
98	6 da	7.1	12								
99	7 da	6.6	12								

Contributors: (a) Levitt, J., (b) Howell, Robert W., (c) Yocum, L. Edwin

References: [1] Gamuna, H. 1932. Protoplasma 16:489. [2] Harris, J. A., R. A. Gortner, and J. V. Lawrence. 1917. Bull. Torrey Botan. Club 44:267. [3] Herrick, E. M. 1933. Am. J. Botany 20:18. [4] Hibbard, R. P., and O. E. Harrington. 1916. Physiol. Res. 1:441. [5] Lewis, F. J., and G. M. Tuttle. 1920. Ann. Botany (London) 34:405. [6] McCool, M. M., and C. E. Miller. 1917. Soil Sci. 3:113. [7] Molz, F. J. 1926. Am. J. Botany 13:413.

continued

## 125. FACTORS AFFECTING OSMOTIC POTENTIAL: VASCULAR PLANTS

### Part II. PHYSICAL AND ENVIRONMENTAL VARIATION

[8] Pfeiffer, M. 1933. *Planta* 19:272. [9] Sayre, J. D. 1926. *Ohio J. Sci.* 26:233. [10] Stoddart, L. A. 1935. *Plant Physiol.* 10:661. [11] Ursprung, A., and G. Blum. 1916. *Ber. Deut. Botan. Ges.* 34:88. [12] Yocum, L. E. 1925. *J. Agr. Res.* 31:727.

### Part III. VARIATION IN DEPTH OF ROOTING

	Species	Common Name	Osmotic Potential								
			Apr 19	Apr 24	May 8	May 28	June 8	June 20	July 2	July 9	July 31
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
Shallow-rooted Plants											
1	<i>Astragalus crassicaarpus</i>	Ground plum milk vetch	10.9	13.2	15.4	18.7	....	....	....	....	....
2	<i>Koeleria cristata</i>	Prairie June grass	16.4	17.4	24.1	35.0	....	....	....	....	....
3	<i>Lomatium foeniculaceum</i>	Lomatium	11.5	14.6	16.1	20.0	....	....	....	....	....
Moderately Deep-rooted Plants											
4	<i>Andropogon scoparius</i>	Little bluestem	8.1	9.6	9.9	25.3	18.1	17.4	31.3	....	....
5	<i>Helianthus rigidus</i>	Stiff sunflower	....	....	12.6	19.9	15.2	22.9	36.9	....	....
6	<i>Solidago glaberrima</i>	Goldenrod	....	12.0	....	12.8	14.3	16.8	....	....	....
Deep-rooted Plants											
7	<i>Amorpha canescens</i>	Leadplant	....	11.6	13.1	20.8	16.2	15.1	17.2	20.0	22.1
8	<i>Baptisia leucophaea</i>	Plains wild indigo	8.6	10.8	12.0	14.8	16.0	16.7	17.3	....	....
9	<i>Psoralea tenuiflora</i>	Slim flower scurf pea	....	12.2	14.2	17.8	16.0	16.8	15.2	17.2	15.8

Contributor: Yocum, L. Edwin

Reference: Stoddart, L. A. 1935. *Plant Physiol.* 10:661.

### Part IV. VARIATION IN HABITAT

	Habitat	Osmotic Potential			Habitat	Osmotic Potential	
		Woody Plants	Herbaceous Plants			Woody Plants	Herbaceous Plants
	(A)	(B)	(C)		(A)	(B)	(C)
Jamaica				Arizona			
1	Ruinat	13	10	5	Rocky slopes	22	16
2	Ridge forest	12	9	6	Canyons	21	13
3	Leeward ravines	11	8	7	Arroyos	17	13
4	Windward habitat	10	8	8	Bajada slopes	30	20
				9	Salt spots	45	24

Contributor: Yocum, L. Edwin

Reference: Harris, J. A., and J. V. Lawrence. 1917. *Am. J. Botany* 4:268.

### Part V. VARIATION IN ECOLOGIC GROUPS

	Plant Group	Osmotic Potential	Refer-ence		Plant Group	Osmotic Potential	Refer-ence
1	Summer ephemerals	8-42	4	6	Hydrophytes		
2	Succulents and winter ephemerals	4-24	4	7	Water leaves	8-9	2
3	Xerophytes	14-57	4	8	Epiphytes	3-6	1
4	Hydrophytes	8-13	1	9	Halophytes	30-115	5
5	Air leaves	18-21	2	10	Parasites	14-17	3
					Hosts	11-14	3

[1] *Phoradendron flavescens* (American mistletoe), 15.8; host, 11.6.

continued

## 125. FACTORS AFFECTING OSMOTIC POTENTIAL: VASCULAR PLANTS

### Part V. VARIATION IN ECOLOGIC GROUPS

*Contributor:* Levitt, J.

*References:* [1] Gamma, H. 1932. Protoplasma 16:489. [2] Gessner, F. 1940. Ber. Deut. Botan. Ges. 58:2. [3] Harris, J. A. 1934. The physico-chemical properties of plant saps in relation to phytogeography. Univ. Minnesota Press, Minneapolis. [4] Huber, B. 1951. Fortschr. Botan. 13:227. [5] Zohary, M., and G. Orshansky. 1949. Palestine J. Botany, Jerusalem Ser., 4:177.

## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part I. DOSE EQUIVALENT TO BODY ORGANS

Values are the recommended permissible doses of ionizing radiation to the various organs of the body of the occupational worker, and are in addition to doses from medical and background exposure. The values apply to both external and internal exposure. The unit of dose equivalent used in this table is the rem. No. of rem = no. of rad x RBE x *n*. (Rad = unit of absorbed dose; 1 rad corresponds to 100 ergs/g of medium. RBE = relative biological effectiveness, i.e., ratio of absorbed dose, in rads, from reference X rays to the absorbed dose, in rads, from the given radiation field required to produce the same effect as the reference X rays. Reference X rays in most cases have been those from 200-250 kilovolts X-radiation or  $\gamma$ -radiation from Co<sup>60</sup>. *n* = relative damage factor.) DE = dose equivalent.

Body Organ	Maximum DE in Any 13 Wk <sup>1</sup> rem/13 wk	Average DE in 1 Yr <sup>2</sup> rem/yr	Accumulated DE for Ages >18 Yr <sup>3</sup> rem	Body Organ	Maximum DE in Any 13 Wk <sup>1</sup> rem/13 wk	Average DE in 1 Yr <sup>2</sup> rem/yr	Accumulated DE for Ages >18 Yr <sup>3</sup> rem
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
1 Total body	3	5	5(age-18)	6 Bone	10	30	30(age-18)
2 Head and trunk	3	5	5(age-18)	7 Skin	8-10	30	30(age-18)
3 Lenses of eyes	3	5	5(age-18)	8 Thyroid	8-10	30	30(age-18)
4 Blood-forming organs	3	5	5(age-18)	9 Feet, ankles, hands, & forearms	20-25	75	75(age-18)
5 Gonads	3	5	5(age-18)	10 Other single organs	4-5	15	15(age-18)

<sup>1/</sup> These values may be used for the accumulated short-term exposures in any 13-week interval. <sup>2/</sup> These values may be used for a planned emergency exposure. <sup>3/</sup> To determine accumulated dose equivalent, multiply value times age minus 18 years.

*Contributor:* Morgan, Karl Z.

*Reference:* Morgan, K. Z. 1963. Science 139:565.

### Part II. TYPE OF RADIATION

All values in columns C and D may be increased by a factor of 6 if the exposure is primarily to the bone, skin, or thyroid. They may be increased by a factor of 3 if the exposure is limited to organs other than the eyes, gonads, or blood-forming organs.

Type of Radiation	QF <sup>1</sup>	Average Exposure Rate <sup>2</sup> mrad/wk	Approximate Flux to Give a Maximum Permissible Exposure in an 8-Hour Day <sup>3</sup>
(A)	(B)	(C)	(D)
1 X and $\gamma$ rays	1	100	$\frac{1400}{E}$ photons per sq cm per sec in free air at 0°C (error <13% for E = 0.07-2 Mev)

<sup>1/</sup> Quality factor, a term used to express the modification of RBE due to LET (linear energy transfer of the radiation), *n* (relative damage factor), and other conditions. <sup>2/</sup> Permissible to eyes, gonads, and blood-forming organs (essentially total body exposure) of individuals 18 years or older. These values may be averaged over a year, provided the dose equivalent in any 13 weeks does not exceed 3 rem (rem = rad x QF). <sup>3/</sup> Rate based on a 20-mrem dose equivalent delivered to tissue in an 8-hour day (= 2.5 per QF mrad per hr). The rad in soft tissue is considered to correspond to an energy absorption of 100 ergs/g. Mev = one million electron volts.

*continued*



## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part II. TYPE OF RADIATION

Type of Radiation	QF <sup>1</sup>	Average Exposure Rate <sup>2</sup> mrad/wk	Approximate Flux to Give a Maximum Permissible Exposure in an 8-Hour Day <sup>3</sup>
(A)	(B)	(C)	(D)
2   β rays and electrons	1	100	$\frac{4.3 \times 10^7}{(QF)P}$ electrons or β rays per sq cm per sec incident on tissue ( $\approx 23$ electrons or 15 β per sq cm per sec of 1 Mev energy)
3   Thermal neutrons	2.5	40	700 thermal neutrons per sq cm per sec incident on tissue
4   Fast neutrons	10	10	19 neutrons of 2 Mev energy per sq cm per sec incident on tissue
5   α particles	10	10	$\frac{4.3 \times 10^7}{(QF)P}$ α particles per sq cm per sec incident on tissue ( $\approx 0.005$ α particles of 5 Mev per sq cm per sec)
6   Protons	10	10	$\frac{4.3 \times 10^7}{(QF)P}$ protons per sq cm per sec incident on tissue ( $\approx 0.06$ protons of 5 Mev per sq cm per sec)
7   Heavy ions	20	5	$\frac{4.3 \times 10^7}{(QF)P}$ heavy ions per sq cm per sec ( $\approx 0.0002$ oxygen ions of 5 Mev per sq cm per sec)

<sup>1/1</sup> Quality factor, a term used to express the modification of RBE due to LET (linear energy transfer of the radiation), *n* (relative damage factor), and other conditions. <sup>2/2</sup> Permissible to eyes, gonads, and blood-forming organs (essentially total body exposure) of individuals 18 years or older. These values may be averaged over a year, provided the dose equivalent in any 13 weeks does not exceed 3 rem (rem = rad × QF). <sup>3/3</sup> Rate based on a 20-mrem dose equivalent delivered to tissue in an 8-hour day (= 2.5 per QF mrad per hr). The rad in soft tissue is considered to correspond to an energy absorption of 100 ergs/g. The *P* is the stopping power in units of electron volts per g per sq cm of soft tissue. Mev = one million electron volts.

*Contributor:* Morgan, Karl Z.

*Reference:* National Research Council, Division of Physical Sciences. 1962. Nuclear instruments and their uses. J. Wiley, New York.

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Values are for radionuclides ingested (in water) or inhaled (in air). Any mixture of the radionuclides listed is considered permissible if the accumulated body burden in any organ, or the concentration in the contents of the gastrointestinal tract, does not reach a value that delivers a dose exceeding the maximum permissible dose-rate of 0.3 rem per week. **Type of Decay** (column C): α = alpha particle; β<sup>-</sup> = negatron; β<sup>+</sup> = positron; γ = gamma ray; e<sup>-</sup> = internal conversion electron; ε = orbital electron capture; SF = spontaneous fission. **Radionuclide** (column D): s = soluble compounds of the radionuclide; i = insoluble compounds of the radionuclide. **Critical Organ** (columns F and I): GI = gastrointestinal tract; (S) = stomach; (SI) = small intestine; (ULI) = upper large intestine; (LLI) = lower large intestine. μc = microcurie, one millionth of a curie or 3.7 × 10<sup>4</sup> disintegrations per second.

Radionuclide				q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i		In Water			In Air			
					Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
1	1	H <sup>3</sup> (HTO or H <sub>2</sub> <sup>3</sup> O)	β <sup>-</sup>	s	1,000	Body tissue	0.1	0.03	Body tissue	5 × 10 <sup>-6</sup>	2 × 10 <sup>-6</sup>
2	4	Be <sup>7</sup>	ε, γ	s	600 <sup>4</sup>	GI (LLI)	0.05	0.02	Total body	6 × 10 <sup>-6</sup>	2 × 10 <sup>-6</sup>
3				i	.....	GI (LLI)	0.05	0.02	Lung	10 <sup>-6</sup>	4 × 10 <sup>-7</sup>
4	6	C <sup>14</sup> (CO <sub>2</sub> )	β <sup>-</sup>	s	300	Fat	0.02	8 × 10 <sup>-3</sup>	Fat	4 × 10 <sup>-6</sup>	10 <sup>-6</sup>
5	9	F <sup>18</sup>	β <sup>+</sup>	s	.....	GI (SI)	0.02	8 × 10 <sup>-3</sup>	GI (SI)	5 × 10 <sup>-6</sup>	2 × 10 <sup>-6</sup>
6				i	.....	GI (ULI)	0.01	5 × 10 <sup>-3</sup>	GI (ULI)	3 × 10 <sup>-6</sup>	9 × 10 <sup>-7</sup>
7	11	Na <sup>22</sup>	β <sup>+</sup> , γ	s	10	Total body	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Total body	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>
8				i	.....	GI (LLI)	9 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	9 × 10 <sup>-9</sup>	3 × 10 <sup>-9</sup>
9	11	Na <sup>24</sup>	β <sup>-</sup> , γ	s	.....	GI (SI)	6 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	GI (SI)	10 <sup>-6</sup>	4 × 10 <sup>-7</sup>
10				i	.....	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	5 × 10 <sup>-8</sup>

<sup>1/1</sup> Z = atomic number. <sup>2/2</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3/3</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4/4</sup> For total body.

*continued*

## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide		Type of Decay	s o r i	q <sup>a</sup>	Maximum Permissible Concentrations of Radionuclides						
					In Water			In Air			
Z <sup>b</sup>	Symbol and Mass No.				Critical Organ <sup>c</sup>	40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>c</sup>	40-hr wk μc/ml	168-hr wk μc/ml	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
11	14	Si <sup>31</sup>	β <sup>-</sup> , γ	s	.....	GI (S)	0.03	9 × 10 <sup>-3</sup>	GI (S)	6 × 10 <sup>-6</sup>	2 × 10 <sup>-6</sup>
12				i	.....	GI (ULI)	6 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	GI (ULI)	10 <sup>-6</sup>	3 × 10 <sup>-7</sup>
13	15	P <sup>32</sup>	β <sup>-</sup>	s	6	Bone	5 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	Bone	7 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
14				i	.....	GI (LLI)	7 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	Lung	8 × 10 <sup>-8</sup>	3 × 10 <sup>-8</sup>
15	16	S <sup>35</sup>	β <sup>-</sup>	s	90	Testis	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	Testis	3 × 10 <sup>-7</sup>	9 × 10 <sup>-8</sup>
16				i	.....	GI (LLI)	8 × 10 <sup>-3</sup>	3 × 10 <sup>-3</sup>	Lung	3 × 10 <sup>-7</sup>	9 × 10 <sup>-8</sup>
17	17	Cl <sup>36</sup>	β <sup>-</sup>	s	80	Total body	2 × 10 <sup>-3</sup>	8 × 10 <sup>-4</sup>	Total body	4 × 10 <sup>-7</sup>	10 <sup>-7</sup>
18				i	.....	GI (LLI)	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-8</sup>	8 × 10 <sup>-9</sup>
19	17	Cl <sup>38</sup>	β <sup>-</sup> , γ	s	.....	GI (S)	0.01	4 × 10 <sup>-3</sup>	GI (S)	3 × 10 <sup>-6</sup>	9 × 10 <sup>-7</sup>
20				i	.....	GI (S)	0.01	4 × 10 <sup>-3</sup>	GI (S)	2 × 10 <sup>-6</sup>	7 × 10 <sup>-7</sup>
21	19	K <sup>42</sup>	β <sup>-</sup> , γ	s	.....	GI (S)	9 × 10 <sup>-3</sup>	3 × 10 <sup>-3</sup>	GI (S)	2 × 10 <sup>-6</sup>	7 × 10 <sup>-7</sup>
22				i	.....	GI (LLI)	6 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	4 × 10 <sup>-8</sup>
23	20	Ca <sup>45</sup>	β <sup>-</sup>	s	30	Bone	3 × 10 <sup>-4</sup>	9 × 10 <sup>-5</sup>	Bone	3 × 10 <sup>-8</sup>	10 <sup>-8</sup>
24				i	.....	GI (LLI)	5 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 × 10 <sup>-8</sup>
25	20	Ca <sup>47</sup>	β <sup>-</sup> , γ	s	5	Bone	10 <sup>-3</sup>	5 × 10 <sup>-4</sup>	Bone	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>
26				i	.....	GI (LLI)	10 <sup>-3</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	2 × 10 <sup>-7<sup>5</sup></sup>	6 × 10 <sup>-8<sup>5</sup></sup>
27	21	Sc <sup>46</sup>	β <sup>-</sup> , γ	s	10 <sup>6</sup>	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Liver	2 × 10 <sup>-7<sup>7</sup></sup>	8 × 10 <sup>-8<sup>7</sup></sup>
28				i	.....	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-8</sup>	8 × 10 <sup>-9</sup>
29	21	Sc <sup>47</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	3 × 10 <sup>-3</sup>	9 × 10 <sup>-4</sup>	GI (LLI)	6 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>
30				i	.....	GI (LLI)	3 × 10 <sup>-3</sup>	9 × 10 <sup>-4</sup>	GI (LLI)	5 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>
31	21	Sc <sup>48</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>
32				i	.....	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	5 × 10 <sup>-8</sup>
33	23	V <sup>48</sup>	β <sup>+</sup> , ε, γ	s	.....	GI (LLI)	9 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>
34				i	.....	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	6 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
35	24	Cr <sup>51</sup>	ε, γ	s	800 <sup>4</sup>	GI (LLI)	0.05	0.02	Total body	10 <sup>-5<sup>7</sup></sup>	4 × 10 <sup>-6<sup>7</sup></sup>
36				i	.....	GI (LLI)	0.05	0.02	Lung	2 × 10 <sup>-6</sup>	8 × 10 <sup>-7</sup>
37	25	Mn <sup>52</sup>	β <sup>+</sup> , ε, γ	s	.....	GI (LLI)	10 <sup>-3</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	2 × 10 <sup>-7</sup>	7 × 10 <sup>-8</sup>
38				i	.....	GI (LLI)	9 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	10 <sup>-7</sup>	5 × 10 <sup>-8<sup>7</sup></sup>
39	25	Mn <sup>54</sup>	ε, γ	s	20 <sup>6</sup>	GI (LLI)	4 × 10 <sup>-3</sup>	10 <sup>-3</sup>	Liver	4 × 10 <sup>-7</sup>	10 <sup>-7</sup>
40				i	.....	GI (LLI)	3 × 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	4 × 10 <sup>-8</sup>	10 <sup>-8</sup>
41	25	Mn <sup>56</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	4 × 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 × 10 <sup>-7</sup>	3 × 10 <sup>-7</sup>
42				i	.....	GI (LLI)	3 × 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	5 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>
43	26	Fe <sup>55</sup>	ε	s	1,000	Spleen	0.02	8 × 10 <sup>-3</sup>	Spleen	9 × 10 <sup>-7</sup>	3 × 10 <sup>-7</sup>
44				i	.....	GI (LLI)	0.07	0.02	Lung	10 <sup>-6</sup>	3 × 10 <sup>-7</sup>
45	26	Fe <sup>59</sup>	β <sup>-</sup> , γ	s	20 <sup>5</sup>	GI (LLI)	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	Spleen	10 <sup>-7</sup>	5 × 10 <sup>-8</sup>
46				i	.....	GI (LLI)	2 × 10 <sup>-3</sup>	5 × 10 <sup>-4</sup>	Lung	5 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
47	27	Co <sup>57</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	0.02	5 × 10 <sup>-3</sup>	GI (LLI)	3 × 10 <sup>-6</sup>	10 <sup>-6</sup>
48				i	.....	GI (LLI)	0.01	4 × 10 <sup>-3</sup>	Lung	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>
49	27	Co <sup>58m</sup>	β <sup>+</sup> , ε, γ	s	.....	GI (LLI)	0.08	0.03	GI (LLI)	2 × 10 <sup>-5</sup>	6 × 10 <sup>-6</sup>
50				i	.....	GI (LLI)	0.06	0.02	Lung	9 × 10 <sup>-6</sup>	3 × 10 <sup>-6</sup>
51	27	Co <sup>58</sup>	β <sup>+</sup> , ε	s	.....	GI (LLI)	4 × 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 × 10 <sup>-7</sup>	3 × 10 <sup>-7<sup>6</sup></sup>
52				i	.....	GI (LLI)	3 × 10 <sup>-3</sup>	9 × 10 <sup>-4</sup>	Lung	5 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
53	27	Co <sup>60</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	10 <sup>-3</sup>	5 × 10 <sup>-4</sup>	GI (LLI)	3 × 10 <sup>-7</sup>	10 <sup>-7<sup>9</sup></sup>
54				i	.....	GI (LLI)	10 <sup>-3</sup>	3 × 10 <sup>-4</sup>	Lung	9 × 10 <sup>-9</sup>	3 × 10 <sup>-9</sup>
55	28	Ni <sup>59</sup>	ε	s	1,000	Bone	6 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	Bone	5 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>
56				i	.....	GI (LLI)	0.06	0.02	Lung	8 × 10 <sup>-7</sup>	3 × 10 <sup>-7</sup>
57	28	Ni <sup>63</sup>	β <sup>-</sup>	s	200	Bone	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Bone	6 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
58				i	.....	GI (LLI)	0.02	7 × 10 <sup>-3</sup>	Lung	3 × 10 <sup>-7</sup>	10 <sup>-7</sup>
59	28	Ni <sup>65</sup>	β <sup>-</sup> , γ	s	.....	GI (ULI)	4 × 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (ULI)	9 × 10 <sup>-7</sup>	3 × 10 <sup>-7</sup>
60				i	.....	GI (ULI)	3 × 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (ULI)	5 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>
61	29	Cu <sup>64</sup>	β <sup>-</sup> , β <sup>+</sup> , ε	s	.....	GI (LLI)	0.01	3 × 10 <sup>-3</sup>	GI (LLI)	2 × 10 <sup>-6</sup>	7 × 10 <sup>-7</sup>
62				i	.....	GI (LLI)	6 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 × 10 <sup>-7</sup>
63	30	Zn <sup>65</sup>	β <sup>+</sup> , ε, γ	s	60	Total body	3 × 10 <sup>-3</sup>	10 <sup>-3<sup>10,11</sup></sup>	Total body	10 <sup>-7<sup>10,11</sup></sup>	4 × 10 <sup>-8<sup>10</sup></sup>
64				i	.....	GI (LLI)	5 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	Lung	6 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>
65	30	Zn <sup>69m</sup>	γ, e <sup>-</sup> , β <sup>-</sup>	s	0.7 <sup>12</sup>	GI (LLI)	2 × 10 <sup>-3</sup>	7 × 10 <sup>-4</sup>	Prostate	4 × 10 <sup>-7</sup>	10 <sup>-7</sup>
66				i	.....	GI (LLI)	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	GI (LLI)	3 × 10 <sup>-7</sup>	10 <sup>-7</sup>

<sup>1/1</sup> Z = atomic number. <sup>1/2</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote.  
<sup>1/3</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>1/4</sup> For total body.  
<sup>1/5</sup> Also lung. <sup>1/6</sup> For liver. <sup>1/7</sup> Also lower large intestine. <sup>1/8</sup> For spleen. <sup>1/9</sup> Also total body. <sup>1/10</sup> Also prostate. <sup>1/11</sup> Also liver. <sup>1/12</sup> For prostate.

continued

## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide				q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i		(E)	In Water			In Air		
						Critical Organ <sup>3</sup>	40-hr wk µc/ml	168-hr wk µc/ml	Critical Organ <sup>3</sup>	40-hr wk µc/ml	168-hr wk µc/ml
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
67	30	Zn <sup>69</sup>	β <sup>-</sup>	s	0.8 <sup>12</sup>	GI (S)	0.05	0.02	Prostate	7 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
68				i	.....	GI (S)	0.05	0.02	GI (S)	9 x 10 <sup>-6</sup>	3 x 10 <sup>-6</sup>
69	31	Ga <sup>72</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
70				i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
71	32	Ge <sup>71</sup>	ε	s	.....	GI (LLI)	0.05	0.02	GI (LLI)	10 <sup>-5</sup>	4 x 10 <sup>-6</sup>
72				i	.....	GI (LLI)	0.05	0.02	Lung	6 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
73	33	As <sup>73</sup>	ε, γ	s	300 <sup>4</sup>	GI (LLI)	0.01	5 x 10 <sup>-3</sup>	Total body	2 x 10 <sup>-6</sup>	7 x 10 <sup>-7</sup>
74				i	.....	GI (LLI)	0.01	5 x 10 <sup>-3</sup>	Lung	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
75	33	As <sup>74</sup>	β <sup>-</sup> , β <sup>+</sup> , ε, γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
76				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
77	33	As <sup>76</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	6 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
78				i	.....	GI (LLI)	6 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	3 x 10 <sup>-8</sup>
79	33	As <sup>77</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	8 x 10 <sup>-4</sup>	GI (LLI)	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
80				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	8 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
81	34	Se <sup>75</sup>	ε, γ	s	90	Kidney	9 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup> <sup>7</sup>	Kidney	10 <sup>-6</sup> <sup>9</sup>	4 x 10 <sup>-7</sup>
82				i	.....	GI (LLI)	8 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
83	35	Br <sup>82</sup>	β <sup>-</sup> , γ	s	10	Total body	8 x 10 <sup>-3</sup> <sup>13</sup>	3 x 10 <sup>-3</sup> <sup>12</sup>	Total body	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
84				i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
85	37	Rb <sup>86</sup>	β <sup>-</sup> , γ	s	30 <sup>5</sup>	Pancreas	2 x 10 <sup>-3</sup> <sup>5</sup>	7 x 10 <sup>-4</sup> <sup>5</sup>	Pancreas	3 x 10 <sup>-7</sup> <sup>7</sup>	10 <sup>-7</sup> <sup>7,11</sup>
86				i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	7 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
87	37	Rb <sup>87</sup>	β <sup>-</sup>	s	200 <sup>5,11</sup>	Pancreas	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Pancreas	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup> <sup>7,11</sup>
88				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	7 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
89	38	Sr <sup>85m</sup>	ε, γ	s	.....	GI (SI)	0.2	0.07	GI (SI)	4 x 10 <sup>-5</sup>	10 <sup>-5</sup>
90				i	.....	GI (SI)	0.2	0.07	GI (SI)	3 x 10 <sup>-5</sup>	10 <sup>-5</sup>
91	38	Sr <sup>85</sup>	ε, γ	s	60	Total body	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Total body	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
92				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
93	38	Sr <sup>89</sup>	β <sup>-</sup>	s	4	Bone	3 x 10 <sup>-4</sup>	10 <sup>-4</sup>	Bone	3 x 10 <sup>-8</sup>	10 <sup>-8</sup>
94				i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	4 x 10 <sup>-8</sup>	10 <sup>-8</sup>
95	38	Sr <sup>90</sup>	β <sup>-</sup>	s	2	Bone	10 <sup>-5</sup>	4 x 10 <sup>-6</sup>	Bone	10 <sup>-9</sup>	4 x 10 <sup>-10</sup>
96				i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	Lung	5 x 10 <sup>-9</sup>	2 x 10 <sup>-9</sup>
97	38	Sr <sup>91</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
98				i	.....	GI (LLI)	10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	9 x 10 <sup>-8</sup>
99	38	Sr <sup>92</sup>	β <sup>-</sup> , γ	s	.....	GI (ULI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	GI (ULI)	4 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
100				i	.....	GI (ULI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (ULI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
101	39	Y <sup>90</sup>	β <sup>-</sup>	s	.....	GI (LLI)	6 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
102				i	.....	GI (LLI)	6 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	3 x 10 <sup>-8</sup>
103	39	Y <sup>91m</sup>	β <sup>-</sup> , γ	s	.....	GI (SI)	0.01	0.03	GI (SI)	2 x 10 <sup>-5</sup>	8 x 10 <sup>-6</sup>
104				i	.....	GI (SI)	0.01	0.03	GI (SI)	2 x 10 <sup>-5</sup>	6 x 10 <sup>-6</sup>
105	39	Y <sup>91</sup>	β <sup>-</sup> , γ	s	5 <sup>14</sup>	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Bone	4 x 10 <sup>-8</sup>	10 <sup>-8</sup>
106				i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	3 x 10 <sup>-8</sup>	10 <sup>-8</sup>
107	39	Y <sup>92</sup>	β <sup>-</sup> , γ	s	.....	GI (ULI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (ULI)	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
108				i	.....	GI (ULI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (ULI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
109	39	Y <sup>93</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
110				i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
111	40	Zr <sup>93</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	100 <sup>14</sup>	GI (LLI)	0.02	8 x 10 <sup>-3</sup>	Bone	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
112				i	.....	GI (LLI)	0.02	8 x 10 <sup>-3</sup>	Lung	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
113	40	Zr <sup>95</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	20 <sup>4</sup>	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	Total body	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
114				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	Lung	3 x 10 <sup>-8</sup>	10 <sup>-8</sup>
115	40	Zr <sup>97</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
116				i	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	9 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
117	41	Nb <sup>93m</sup>	γ, e <sup>-</sup>	s	200 <sup>14</sup>	GI (LLI)	0.01	4 x 10 <sup>-3</sup>	Bone	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
118				i	.....	GI (LLI)	0.01	4 x 10 <sup>-3</sup>	Lung	2 x 10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
119	41	Nb <sup>95</sup>	β <sup>-</sup> , γ	s	40 <sup>6</sup>	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Total body	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
120				i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	3 x 10 <sup>-8</sup>
121	41	Nb <sup>97</sup>	β <sup>-</sup> , γ	s	.....	GI (ULI)	0.03	9 x 10 <sup>-3</sup>	GI (ULI)	6 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
122				i	.....	GI (ULI)	0.03	9 x 10 <sup>-3</sup>	GI (ULI)	5 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>

<sup>1/1</sup> Z = atomic number. <sup>1/2</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote.

<sup>1/3</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>1/4</sup> For total body.

<sup>1/5</sup> Also lower large intestine. <sup>1/6</sup> Also total body. <sup>1/11</sup> Also liver. <sup>1/12</sup> For prostate. <sup>1/13</sup> Also small intestine.

<sup>1/14</sup> For bone. <sup>1/15</sup> For kidney.

*continued*



## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide		Maximum Permissible Concentrations of Radionuclides								
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i	q <sup>2</sup>	Critical Organ <sup>3</sup>	In Water 40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>2</sup>	In Air 40-hr wk μc/ml	168-hr wk μc/ml
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
123	42 Mo <sup>99</sup>	β <sup>-</sup> , γ	s	8	Kidney	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup> <sup>7</sup>	Kidney	7 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
124			i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	7 x 10 <sup>-8</sup>
125	43 Tc <sup>96m</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	0.4	0.1	GI (LLI)	8 x 10 <sup>-5</sup>	3 x 10 <sup>-5</sup>
126			i	.....	GI (LLI)	0.3	0.1	Lung	3 x 10 <sup>-5</sup>	10 <sup>-5</sup>
127	43 Tc <sup>96</sup>	ε, γ	s	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
128			i	.....	GI (LLI)	10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
129	43 Tc <sup>97m</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	0.01	4 x 10 <sup>-3</sup>	GI (LLI)	2 x 10 <sup>-6</sup>	8 x 10 <sup>-7</sup>
130			i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	2 x 10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
131	43 Tc <sup>97</sup>	ε	s	60 <sup>15</sup>	GI (LLI)	0.05	0.02	Kidney	10 <sup>-5</sup> <sup>7</sup>	4 x 10 <sup>-6</sup> <sup>7</sup>
132			i	.....	GI (LLI)	0.02	8 x 10 <sup>-3</sup>	Lung	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
133	43 Tc <sup>99m</sup>	β <sup>-</sup> , γ	s	.....	GI (ULI)	0.2	0.06	GI (ULI)	4 x 10 <sup>-5</sup>	10 <sup>-5</sup>
134			i	.....	GI (ULI)	0.08	0.03	GI (ULI)	10 <sup>-5</sup>	5 x 10 <sup>-6</sup>
135	43 Tc <sup>99</sup>	β <sup>-</sup>	s	.....	GI (LLI)	0.01	3 x 10 <sup>-3</sup>	GI (LLI)	2 x 10 <sup>-6</sup>	7 x 10 <sup>-7</sup>
136			i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	6 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
137	44 Ru <sup>97</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	0.01	4 x 10 <sup>-3</sup>	GI (LLI)	2 x 10 <sup>-6</sup>	8 x 10 <sup>-7</sup>
138			i	.....	GI (LLI)	0.01	3 x 10 <sup>-3</sup>	GI (LLI)	2 x 10 <sup>-6</sup> <sup>5</sup>	6 x 10 <sup>-7</sup>
139	44 Ru <sup>103</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	8 x 10 <sup>-4</sup>	GI (LLI)	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
140			i	.....	.....	.....	.....	Lung	8 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
141	44 Ru <sup>105</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (ULI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (ULI)	7 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
142			i	.....	GI (ULI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (ULI)	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
143	44 Ru <sup>106</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	4 x 10 <sup>-4</sup>	10 <sup>-4</sup>	GI (LLI)	8 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
144			i	.....	GI (LLI)	3 x 10 <sup>-4</sup>	10 <sup>-4</sup>	Lung	6 x 10 <sup>-9</sup>	2 x 10 <sup>-9</sup>
145	45 Rh <sup>103m</sup>	γ, e <sup>-</sup>	s	.....	GI (S)	0.4	0.1	GI (S)	8 x 10 <sup>-5</sup>	3 x 10 <sup>-5</sup>
146			i	.....	GI (S)	0.3	0.1	GI (S)	6 x 10 <sup>-5</sup>	2 x 10 <sup>-5</sup>
147	45 Rh <sup>105</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
148			i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
149	46 Pd <sup>103</sup>	ε, γ, e <sup>-</sup>	s	20 <sup>15</sup>	GI (LLI)	0.01	3 x 10 <sup>-3</sup>	Kidney	10 <sup>-6</sup>	5 x 10 <sup>-7</sup>
150			i	.....	GI (LLI)	8 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup>	Lung	7 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
151	46 Pd <sup>109</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	3 x 10 <sup>-3</sup>	9 x 10 <sup>-4</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
152			i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
153	47 Ag <sup>105</sup>	ε, γ	s	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
154			i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	8 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
155	47 Ag <sup>110m</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	9 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	7 x 10 <sup>-8</sup>
156			i	.....	GI (LLI)	9 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	10 <sup>-8</sup>	3 x 10 <sup>-9</sup>
157	47 Ag <sup>111</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
158			i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
159	48 Cd <sup>109</sup>	ε, γ, e <sup>-</sup>	s	20 <sup>6,15</sup>	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Liver	5 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup> <sup>8</sup>
160			i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	7 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
161	48 Cd <sup>115m</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	3 <sup>3</sup>	GI (LLI)	7 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Liver	4 x 10 <sup>-8</sup> <sup>15</sup>	10 <sup>-8</sup>
162			i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	4 x 10 <sup>-8</sup>	10 <sup>-8</sup>
163	48 Cd <sup>115</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	10 <sup>-3</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
164			i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
165	49 In <sup>113m</sup>	γ, e <sup>-</sup>	s	.....	GI (ULI)	0.04	0.01	GI (ULI)	8 x 10 <sup>-6</sup>	3 x 10 <sup>-6</sup>
166			i	.....	GI (ULI)	0.04	0.01	GI (ULI)	7 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
167	49 In <sup>114m</sup>	β <sup>-</sup> , ε, γ, e <sup>-</sup>	s	6 <sup>15</sup>	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Kidney	10 <sup>-7</sup> <sup>7,17</sup>	4 x 10 <sup>-8</sup> <sup>7,17</sup>
168			i	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	2 x 10 <sup>-8</sup>	7 x 10 <sup>-9</sup>
169	49 In <sup>115m</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (ULI)	0.01	4 x 10 <sup>-3</sup>	GI (ULI)	2 x 10 <sup>-6</sup>	8 x 10 <sup>-7</sup>
170			i	.....	GI (ULI)	0.01	4 x 10 <sup>-3</sup>	GI (ULI)	2 x 10 <sup>-6</sup>	6 x 10 <sup>-7</sup>
171	49 In <sup>115</sup>	β <sup>-</sup>	s	30 <sup>15</sup>	GI (LLI)	3 x 10 <sup>-3</sup>	9 x 10 <sup>-4</sup>	Kidney	2 x 10 <sup>-7</sup>	9 x 10 <sup>-8</sup>
172			i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	9 x 10 <sup>-4</sup>	Lung	3 x 10 <sup>-8</sup>	10 <sup>-8</sup>
173	50 Sn <sup>113</sup>	ε, γ, e <sup>-</sup>	s	30 <sup>14</sup>	GI (LLI)	2 x 10 <sup>-3</sup>	9 x 10 <sup>-4</sup>	Bone	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
174			i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	8 x 10 <sup>-4</sup>	Lung	5 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
175	50 Sn <sup>125</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
176			i	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	8 x 10 <sup>-8</sup>	3 x 10 <sup>-8</sup>
177	51 Sb <sup>122</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
178			i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
179	51 Sb <sup>124</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup> <sup>8</sup>	5 x 10 <sup>-8</sup>
180			i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	2 x 10 <sup>-8</sup>	7 x 10 <sup>-9</sup>

<sup>1</sup>/ Z = atomic number. <sup>2</sup>/ Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3</sup>/ That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4</sup>/ Also lung. <sup>5</sup>/ For liver. <sup>6</sup>/ Also lower large intestine. <sup>7</sup>/ Also total body. <sup>8</sup>/ For bone. <sup>9</sup>/ For kidney. <sup>10</sup>/ Also kidney. <sup>11</sup>/ Also spleen.

continued



## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide				q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i		In Water			In Air			
					Critical Organ <sup>3</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$	Critical Organ <sup>3</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
181	51	Sb <sup>125</sup>	$\beta^-$ , $\gamma$ , e	s	40 <sup>1a</sup>	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	Lung	$5 \times 10^{-7}$	$2 \times 10^{-7}$
182				i	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	Lung	$3 \times 10^{-8}$	$9 \times 10^{-9}$
183	52	Te <sup>125m</sup>	$\gamma$ , e <sup>-</sup>	s	20 <sup>2c</sup>	Kidney	$5 \times 10^{-37}$	$2 \times 10^{-37,20}$	Kidney	$4 \times 10^{-7}$	$10^{-7}$
184				i	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	Lung	$10^{-7}$	$4 \times 10^{-8}$
185	52	Te <sup>127m</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	7 <sup>20</sup>	Kidney	$2 \times 10^{-37,20}$	$6 \times 10^{-4}$	Kidney	$10^{-7,20}$	$5 \times 10^{-8,20}$
186				i	.....	GI (LLI)	$2 \times 10^{-3}$	$5 \times 10^{-4}$	Lung	$4 \times 10^{-8}$	$10^{-8}$
187	52	Te <sup>127</sup>	$\beta^-$	s	.....	GI (LLI)	$8 \times 10^{-3}$	$3 \times 10^{-3}$	GI (LLI)	$2 \times 10^{-6}$	$6 \times 10^{-7}$
188				i	.....	GI (LLI)	$5 \times 10^{-3}$	$2 \times 10^{-3}$	GI (LLI)	$9 \times 10^{-7}$	$3 \times 10^{-7}$
189	52	Te <sup>129m</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	3 <sup>15,20</sup>	GI (LLI)	$10^{-316,20}$	$3 \times 10^{-4}$	Kidney	$8 \times 10^{-8}$	$3 \times 10^{-8,20}$
190				i	.....	GI (LLI)	$6 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$3 \times 10^{-8}$	$10^{-8}$
191	52	Te <sup>129</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	.....	GI (S)	0.02	$8 \times 10^{-3}$	GI (S)	$5 \times 10^{-6}$	$2 \times 10^{-6}$
192				i	.....	GI (ULI)	0.02	$8 \times 10^{-3}$	GI (ULI)	$4 \times 10^{-6}$	$10^{-6}$
193	52	Te <sup>131m</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	GI (LLI)	$4 \times 10^{-7}$	$10^{-7}$
194				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$6 \times 10^{-8}$
195	52	Te <sup>132</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
196				i	.....	GI (LLI)	$6 \times 10^{-4}$	$2 \times 10^{-4}$	GI (LLI)	$10^{-7}$	$4 \times 10^{-8}$
197	53	I <sup>126</sup>	$\beta^-$ , $\epsilon$ , $\gamma$	s	1	Thyroid	$5 \times 10^{-5}$	$2 \times 10^{-5}$	Thyroid	$8 \times 10^{-9}$	$3 \times 10^{-9}$
198				i	.....	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	Lung	$3 \times 10^{-7}$	$10^{-7}$
199	53	I <sup>129</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	3	Thyroid	$10^{-5}$	$4 \times 10^{-6}$	Thyroid	$2 \times 10^{-9}$	$6 \times 10^{-10}$
200				i	.....	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$7 \times 10^{-8}$	$2 \times 10^{-8}$
201	53	I <sup>131</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	0.7	Thyroid	$6 \times 10^{-5}$	$2 \times 10^{-5}$	Thyroid	$9 \times 10^{-9}$	$3 \times 10^{-9}$
202				i	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	GI (LLI)	$3 \times 10^{-7}$	$10^{-7}$
203	53	I <sup>132</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	0.3	Thyroid	$2 \times 10^{-3}$	$6 \times 10^{-4}$	Thyroid	$2 \times 10^{-7}$	$8 \times 10^{-8}$
204				i	.....	GI (ULI)	$5 \times 10^{-3}$	$2 \times 10^{-3}$	GI (ULI)	$9 \times 10^{-7}$	$3 \times 10^{-7}$
205	53	I <sup>133</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	0.3	Thyroid	$2 \times 10^{-4}$	$7 \times 10^{-5}$	Thyroid	$3 \times 10^{-8}$	$10^{-8}$
206				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
207	53	I <sup>134</sup>	$\beta^-$ , $\gamma$	s	0.2	Thyroid	$4 \times 10^{-3}$	$10^{-3}$	Thyroid	$5 \times 10^{-7}$	$2 \times 10^{-7}$
208				i	.....	GI (S)	0.02	$6 \times 10^{-3}$	GI (S)	$3 \times 10^{-6}$	$10^{-6}$
209	53	I <sup>135</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	0.3	Thyroid	$7 \times 10^{-4}$	$2 \times 10^{-4}$	Thyroid	$10^{-7}$	$4 \times 10^{-8}$
210				i	.....	GI (LLI)	$2 \times 10^{-3}$	$7 \times 10^{-4}$	GI (LLI)	$4 \times 10^{-7}$	$10^{-7}$
211	55	Cs <sup>131</sup>	$\epsilon$	s	700	Total body	0.07	0.02	Total body	$10^{-5,11}$	$4 \times 10^{-6,11}$
212				i	.....	GI (LLI)	0.03	$9 \times 10^{-3}$	Lung	$3 \times 10^{-6}$	$10^{-6}$
213	55	Cs <sup>134m</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	.....	GI (S)	0.02	0.06	GI (S)	$4 \times 10^{-5}$	$10^{-5}$
214				i	.....	GI (ULI)	0.03	0.01	GI (ULI)	$6 \times 10^{-6}$	$2 \times 10^{-6}$
215	55	Cs <sup>134</sup>	$\beta^-$ , $\gamma$	s	20	Total body	$3 \times 10^{-4}$	$9 \times 10^{-5}$	Total body	$4 \times 10^{-8}$	$10^{-8}$
216				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Lung	$10^{-8}$	$4 \times 10^{-9}$
217	55	Cs <sup>135</sup>	$\beta^-$	s	200	Liver	$3 \times 10^{-3}$	$10^{-3,17}$	Liver	$5 \times 10^{-7,17}$	$2 \times 10^{-7,17}$
218				i	.....	GI (LLI)	$7 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$9 \times 10^{-8}$	$3 \times 10^{-8}$
219	55	Cs <sup>136</sup>	$\beta^-$ , $\gamma$	s	30	Total body	$2 \times 10^{-3}$	$9 \times 10^{-4}$	Total body	$4 \times 10^{-7}$	$10^{-7}$
220				i	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	Lung	$2 \times 10^{-7}$	$6 \times 10^{-8}$
221	55	Cs <sup>137</sup>	$\beta^-$ , $\gamma$ , e <sup>-</sup>	s	30	Total body	$4 \times 10^{-4}$	$2 \times 10^{-4,21}$	Total body	$6 \times 10^{-8}$	$2 \times 10^{-8}$
222				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Lung	$10^{-8}$	$5 \times 10^{-9}$
223	56	Ba <sup>131</sup>	$\epsilon$ , $\gamma$	s	.....	GI (LLI)	$5 \times 10^{-3}$	$2 \times 10^{-3}$	GI (LLI)	$10^{-6}$	$4 \times 10^{-7}$
224				i	.....	GI (LLI)	$5 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$4 \times 10^{-7}$	$10^{-7}$
225	56	Ba <sup>140</sup>	$\beta^-$ , $\gamma$	s	4 <sup>14</sup>	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Bone	$10^{-7}$	$4 \times 10^{-8}$
226				i	.....	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$4 \times 10^{-8}$	$10^{-8}$
227	57	La <sup>140</sup>	$\beta^-$ , $\gamma$	s	.....	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$5 \times 10^{-8}$
228				i	.....	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	GI (LLI)	$10^{-7}$	$4 \times 10^{-8}$
229	58	Ce <sup>141</sup>	$\beta^-$ , $\gamma$	s	3 <sup>19</sup>	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	Liver	$4 \times 10^{-7}$	$2 \times 10^{-7,12}$
230				i	.....	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	Lung	$2 \times 10^{-7}$	$5 \times 10^{-8}$
231	58	Ce <sup>143</sup>	$\beta^-$ , $\gamma$	s	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$3 \times 10^{-7}$	$9 \times 10^{-8}$
232				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
233	58	Ce <sup>144</sup>	$\alpha$ , $\beta^-$ , $\gamma$	s	5 <sup>14</sup>	GI (LLI)	$3 \times 10^{-4}$	$10^{-4}$	Bone	$10^{-8,11}$	$3 \times 10^{-9}$
234				i	.....	GI (LLI)	$3 \times 10^{-4}$	$10^{-4}$	Lung	$6 \times 10^{-9}$	$2 \times 10^{-9}$
235	59	Pr <sup>142</sup>	$\beta^-$ , $\gamma$	s	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
236				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$5 \times 10^{-8}$

<sup>1/</sup> Z = atomic number. <sup>2/</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3/</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4/</sup> Also lung. <sup>5/</sup> For liver. <sup>7/</sup> Also lower large intestine. <sup>9/</sup> Also total body. <sup>11/</sup> Also liver. <sup>14/</sup> For bone. <sup>15/</sup> For kidney. <sup>16/</sup> Also kidney. <sup>17/</sup> Also spleen. <sup>18/</sup> For lung. <sup>19/</sup> Also bone. <sup>20/</sup> Also testis. <sup>21/</sup> Also liver, spleen, and muscle.

continued

## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Z <sup>1</sup>	Radionuclide			q <sup>3</sup>	Maximum Permissible Concentrations of Radionuclides						
	Symbol and Mass No.	Type of Decay	s or i		In Water			In Air			
					Critical Organ <sup>2</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$	Critical Organ <sup>2</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
237	59	Pr <sup>143</sup>	$\beta^-$	s	.....	GI (LLI)	$10^{-3}$	$5 \times 10^{-4}$	GI (LLI)	$3 \times 10^{-7}$	$10^{-7}$
238				i	.....	GI (LLI)	$10^{-3}$	$5 \times 10^{-4}$	Lung	$2 \times 10^{-7}$	$6 \times 10^{-8}$
239	60	Nd <sup>144</sup>	$\alpha$	s	0.1	Bone	$2 \times 10^{-37}$	$7 \times 10^{-4}$	Bone	$8 \times 10^{-11}$	$3 \times 10^{-11}$
240				i	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	Lung	$3 \times 10^{-10}$	$10^{-10}$
241	60	Nd <sup>147</sup>	$\alpha, \beta^-, \gamma$	s	$10^0$	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	Liver	$4 \times 10^{-77}$	$10^{-77}$
242				i	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	Lung	$2 \times 10^{-7}$	$8 \times 10^{-8}$
243	60	Nd <sup>149</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$8 \times 10^{-3}$	$3 \times 10^{-3}$	GI (LLI)	$2 \times 10^{-6}$	$6 \times 10^{-7}$
244				i	.....	GI (ULI)	$8 \times 10^{-3}$	$3 \times 10^{-3}$	GI (ULI)	$10^{-6}$	$5 \times 10^{-7}$
245	61	Pm <sup>147</sup>	$\alpha, \beta^-$	s	$60^{14}$	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Bone	$6 \times 10^{-8}$	$2 \times 10^{-8}$
246				i	.....	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$10^{-7}$	$3 \times 10^{-8}$
247	61	Pm <sup>149</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$3 \times 10^{-7}$	$10^{-7}$
248				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$8 \times 10^{-8}$
249	62	Sm <sup>147</sup>	$\alpha$	s	0.1	Bone	$2 \times 10^{-37}$	$6 \times 10^{-4}$	Bone	$7 \times 10^{-11}$	$2 \times 10^{-11}$
250				i	.....	GI (LLI)	$2 \times 10^{-3}$	$7 \times 10^{-4}$	Lung	$3 \times 10^{-10}$	$9 \times 10^{-11}$
251	62	Sm <sup>151</sup>	$\beta^-, \gamma$	s	$100^{14}$	GI (LLI)	0.01	$4 \times 10^{-3}$	Bone	$6 \times 10^{-8}$	$2 \times 10^{-8}$
252				i	.....	GI (LLI)	0.01	$4 \times 10^{-3}$	Lung	$10^{-7}$	$5 \times 10^{-8}$
253	62	Sm <sup>153</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	GI (LLI)	$5 \times 10^{-7}$	$2 \times 10^{-7}$
254				i	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	GI (LLI)	$4 \times 10^{-7}$	$10^{-7}$
255	63	Eu <sup>152</sup> (9.2 hr)	$\beta^-, \epsilon, \gamma$	s	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	GI (LLI)	$4 \times 10^{-7}$	$10^{-7}$
256				i	.....	GI (LLI)	$2 \times 10^{-3}$	$6 \times 10^{-4}$	GI (LLI)	$3 \times 10^{-7}$	$10^{-7}$
257	63	Eu <sup>152</sup> (13 yr)	$\beta^-, \epsilon, \gamma$	s	$20^{15}$	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	Kidney	$10^{-8}$	$4 \times 10^{-9}$
258				i	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	Lung	$2 \times 10^{-8}$	$6 \times 10^{-9}$
259	63	Eu <sup>154</sup>	$\beta^-, \epsilon, \gamma$	s	$5^{15,10}$	GI (LLI)	$6 \times 10^{-4}$	$2 \times 10^{-4}$	Kidney	$4 \times 10^{-9,9}$	$10^{-9,9}$
260				i	.....	GI (LLI)	$6 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$7 \times 10^{-9}$	$2 \times 10^{-9}$
261	63	Eu <sup>155</sup>	$\beta^-, \gamma$	s	$70^{15}$	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Kidney	$9 \times 10^{-8}$	$3 \times 10^{-8,14}$
262				i	.....	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$7 \times 10^{-8}$	$3 \times 10^{-8}$
263	64	Gd <sup>153</sup>	$\epsilon, \gamma, e^-$	s	$90^{14}$	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Bone	$2 \times 10^{-7}$	$8 \times 10^{-8}$
264				i	.....	GI (LLI)	$6 \times 10^{-3}$	$2 \times 10^{-3}$	Lung	$9 \times 10^{-8}$	$3 \times 10^{-8}$
265	64	Gd <sup>159</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	GI (LLI)	$5 \times 10^{-7}$	$2 \times 10^{-7}$
266				i	.....	GI (LLI)	$2 \times 10^{-3}$	$8 \times 10^{-4}$	GI (LLI)	$4 \times 10^{-7}$	$10^{-7}$
267	65	Tb <sup>160</sup>	$\beta^-, \gamma$	s	$20^{14}$	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Bone	$10^{-7,12}$	$3 \times 10^{-8}$
268				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Lung	$3 \times 10^{-8}$	$10^{-8}$
269	66	Dy <sup>165</sup>	$\beta^-, \gamma$	s	.....	GI (ULI)	0.01	$4 \times 10^{-3}$	GI (ULI)	$3 \times 10^{-6}$	$9 \times 10^{-7}$
270				i	.....	GI (ULI)	0.01	$4 \times 10^{-3}$	GI (ULI)	$2 \times 10^{-6}$	$7 \times 10^{-7}$
271	66	Dy <sup>166</sup>	$\beta^-, \gamma, e^-$	s	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$8 \times 10^{-8}$
272				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
273	67	Ho <sup>166</sup>	$\beta^-, \gamma, e^-$	s	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$7 \times 10^{-8}$
274				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$6 \times 10^{-8}$
275	68	Er <sup>169</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	GI (LLI)	$6 \times 10^{-7}$	$2 \times 10^{-7}$
276				i	.....	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	Lung	$4 \times 10^{-7}$	$10^{-7}$
277	68	Er <sup>171</sup>	$\beta^-, \gamma, e^-$	s	.....	GI (ULI)	$3 \times 10^{-3}$	$10^{-3}$	GI (ULI)	$7 \times 10^{-7}$	$2 \times 10^{-7}$
278				i	.....	GI (ULI)	$3 \times 10^{-3}$	$10^{-3}$	GI (ULI)	$6 \times 10^{-7}$	$2 \times 10^{-7}$
279	69	Tm <sup>170</sup>	$\beta^-, \epsilon, \gamma, e^-$	s	$9^{14}$	GI (LLI)	$10^{-3}$	$5 \times 10^{-4}$	Bone	$4 \times 10^{-8}$	$10^{-8}$
280				i	.....	GI (LLI)	$10^{-3}$	$5 \times 10^{-4}$	Lung	$3 \times 10^{-8}$	$10^{-8}$
281	69	Tm <sup>171</sup>	$\beta^-$	s	$90^{14}$	GI (LLI)	0.01	$5 \times 10^{-3}$	Bone	$10^{-7}$	$4 \times 10^{-8}$
282				i	.....	GI (LLI)	0.01	$5 \times 10^{-3}$	Lung	$2 \times 10^{-7}$	$8 \times 10^{-8}$
283	70	Yb <sup>175</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$7 \times 10^{-7}$	$2 \times 10^{-7}$
284				i	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$6 \times 10^{-7}$	$2 \times 10^{-7}$
285	71	Lu <sup>177</sup>	$\beta^-, \gamma$	s	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$6 \times 10^{-7}$	$2 \times 10^{-7}$
286				i	.....	GI (LLI)	$3 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$5 \times 10^{-7}$	$2 \times 10^{-7}$
287	72	Hf <sup>181</sup>	$\beta^-, \gamma$	s	$4^8$	GI (LLI)	$2 \times 10^{-3}$	$7 \times 10^{-4}$	Spleen	$4 \times 10^{-8}$	$10^{-8}$
288				i	.....	GI (LLI)	$2 \times 10^{-3}$	$7 \times 10^{-4}$	Lung	$7 \times 10^{-8}$	$3 \times 10^{-8}$
289	73	Ta <sup>182</sup>	$\beta^-, \gamma$	s	$7^6$	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Liver	$4 \times 10^{-8}$	$10^{-8}$
290				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Lung	$2 \times 10^{-8}$	$7 \times 10^{-9}$
291	74	W <sup>181</sup>	$\epsilon, \gamma$	s	.....	GI (LLI)	0.01	$4 \times 10^{-3}$	GI (LLI)	$2 \times 10^{-6}$	$8 \times 10^{-7}$
292				i	.....	GI (LLI)	0.01	$3 \times 10^{-3}$	Lung	$10^{-7}$	$4 \times 10^{-8}$

<sup>1/</sup> Z = atomic number. <sup>2/</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3/</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4/</sup> For liver. <sup>7/</sup> Also lower large intestine. <sup>8/</sup> For spleen. <sup>10/</sup> Also total body. <sup>14/</sup> For bone. <sup>15/</sup> For kidney. <sup>16/</sup> Also kidney. <sup>17/</sup> Also bone.

*continued*

## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide				q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i		(E)	In Water			In Air		
						Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
293	74	W <sup>185</sup>	β <sup>-</sup>	s	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
294				i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
295	74	W <sup>187</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
296				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
297	75	Re <sup>183</sup>	ε, γ	s	80 <sup>4</sup>	GI (LLI)	0.02 <sup>2</sup>	6 x 10 <sup>-3</sup>	Total body	3 x 10 <sup>-6</sup>	9 x 10 <sup>-7</sup>
298				i	.....	GI (LLI)	8 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup>	Lung	2 x 10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
299	75	Re <sup>186</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	3 x 10 <sup>-3</sup>	9 x 10 <sup>-4</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
300				i	.....	GI (LLI)	10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
301	75	Re <sup>187</sup>	β <sup>-</sup>	s	300 <sup>22</sup>	GI (LLI)	0.07	0.03 <sup>23</sup>	Skin	9 x 10 <sup>-6</sup>	3 x 10 <sup>-6</sup>
302				i	.....	GI (LLI)	0.04	0.02	Lung	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
303	75	Re <sup>188</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
304				i	.....	GI (LLI)	9 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	6 x 10 <sup>-8</sup>
305	76	Os <sup>185</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	GI (LLI)	5 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
306				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	Lung	5 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
307	76	Os <sup>191m</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	0.07	0.03	GI (LLI)	2 x 10 <sup>-5</sup>	6 x 10 <sup>-6</sup>
308				i	.....	GI (LLI)	0.07	0.02	Lung	9 x 10 <sup>-6</sup>	3 x 10 <sup>-6</sup>
309	76	Os <sup>191</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
310				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
311	76	Os <sup>193</sup>	β <sup>-</sup>	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	GI (LLI)	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
312				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	9 x 10 <sup>-8</sup>
313	77	Ir <sup>190</sup>	ε, γ	s	.....	GI (LLI)	6 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
314				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Lung	4 x 10 <sup>-7</sup>	10 <sup>-7</sup>
315	77	Ir <sup>192</sup>	β <sup>-</sup> , γ	s	6 <sup>18</sup>	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	Kidney	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
316				i	.....	GI (LLI)	10 <sup>-3</sup>	4 x 10 <sup>-4</sup>	Lung	3 x 10 <sup>-8</sup>	9 x 10 <sup>-9</sup>
317	77	Ir <sup>194</sup>	β <sup>-</sup>	s	.....	GI (LLI)	10 <sup>-3</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
318				i	.....	GI (LLI)	9 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
319	78	Pt <sup>191</sup>	ε, γ	s	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
320				i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
321	78	Pt <sup>193m</sup>	ε, γ	s	.....	GI (LLI)	0.03	0.01	GI (LLI)	7 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
322				i	.....	GI (LLI)	0.03	0.01	GI (LLI)	5 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
323	78	Pt <sup>193</sup>	ε	s	70	Kidney	0.03	9 x 10 <sup>-3</sup>	Kidney	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
324				i	.....	GI (LLI)	0.05	0.02	Lung	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
325	78	Pt <sup>197m</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (ULI)	0.03	0.01	GI (ULI)	6 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
326				i	.....	GI (ULI)	0.03	9 x 10 <sup>-3</sup>	GI (ULI)	5 x 10 <sup>-6</sup>	2 x 10 <sup>-6</sup>
327	78	Pt <sup>197</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
328				i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
329	79	Au <sup>196</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
330				i	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
331	79	Au <sup>198</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	2 x 10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	3 x 10 <sup>-7</sup>	10 <sup>-7</sup>
332				i	.....	GI (LLI)	10 <sup>-3</sup>	5 x 10 <sup>-4</sup>	GI (LLI)	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
333	79	Au <sup>199</sup>	β <sup>-</sup> , γ	s	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
334				i	.....	GI (LLI)	4 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
335	80	Hg <sup>197m</sup>	ε, γ, e <sup>-</sup>	s	4	Kidney	6 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Kidney	7 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
336				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
337	80	Hg <sup>197</sup>	ε, γ, e <sup>-</sup>	s	20	Kidney	9 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup>	Kidney	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
338				i	.....	GI (LLI)	0.01	5 x 10 <sup>-3</sup>	GI (LLI)	3 x 10 <sup>-6</sup>	9 x 10 <sup>-7</sup>
339	80	Hg <sup>203</sup>	β <sup>-</sup> , γ, e <sup>-</sup>	s	4	Kidney	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Kidney	7 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
340				i	.....	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
341	81	Tl <sup>200</sup>	ε, γ	s	.....	GI (LLI)	0.01	4 x 10 <sup>-3</sup>	GI (LLI)	3 x 10 <sup>-6</sup>	9 x 10 <sup>-7</sup>
342				i	.....	GI (LLI)	7 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
343	81	Tl <sup>201</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	9 x 10 <sup>-3</sup>	3 x 10 <sup>-3</sup>	GI (LLI)	2 x 10 <sup>-6</sup>	7 x 10 <sup>-7</sup>
344				i	.....	GI (LLI)	5 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (LLI)	9 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
345	81	Tl <sup>202</sup>	ε, γ, e <sup>-</sup>	s	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	GI (LLI)	8 x 10 <sup>-7</sup>	3 x 10 <sup>-7</sup>
346				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	7 x 10 <sup>-4</sup>	Lung	2 x 10 <sup>-7</sup>	8 x 10 <sup>-8</sup>
347	81	Tl <sup>204</sup>	β <sup>-</sup>	s	10 <sup>18</sup>	GI (LLI)	3 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Kidney	6 x 10 <sup>-7</sup>	2 x 10 <sup>-7</sup>
348				i	.....	GI (LLI)	2 x 10 <sup>-3</sup>	6 x 10 <sup>-4</sup>	Lung	3 x 10 <sup>-8</sup>	9 x 10 <sup>-9</sup>

<sup>1/1</sup> Z = atomic number. <sup>1/2</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote.  
<sup>1/3</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>1/4</sup> For total body.  
<sup>1/5</sup> Also lower large intestine. <sup>1/6</sup> Also total body. <sup>1/5</sup> For kidney. <sup>1/7</sup> Also spleen. <sup>1/8</sup> For skin. <sup>1/9</sup> Also skin.

continued



## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Radionuclide				q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
Z <sup>1</sup>	Symbol and Mass No.	Type of Decay	s or i		Critical Organ <sup>2</sup>	In Water			In Air		
						40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>2</sup>	40-hr wk μc/ml	168-hr wk μc/ml	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
349	82	Pb <sup>203</sup>	ε, γ	s	GI (LLI)	0.01	4 × 10 <sup>-3</sup>	GI (LLI)	3 × 10 <sup>-6</sup>	9 × 10 <sup>-7</sup>	
350				i	GI (LLI)	0.01	4 × 10 <sup>-3</sup>	GI (LLI)	2 × 10 <sup>-6</sup>	6 × 10 <sup>-7</sup>	
351	82	Pb <sup>210</sup>	α, β <sup>-</sup> , γ	s	Kidney	4 × 10 <sup>-6<sup>2</sup></sup>	10 <sup>-6<sup>3</sup></sup>	Kidney	10 <sup>-10</sup>	4 × 10 <sup>-11</sup>	
352				i	GI (LLI)	5 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	Lung	2 × 10 <sup>-10</sup>	8 × 10 <sup>-11</sup>	
353	82	Pb <sup>212</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Kidney	6 × 10 <sup>-4<sup>7</sup></sup>	2 × 10 <sup>-4<sup>7</sup></sup>	Kidney	2 × 10 <sup>-8</sup>	6 × 10 <sup>-9</sup>	
354				i	GI (LLI)	5 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-8</sup>	7 × 10 <sup>-9</sup>	
355	83	Bi <sup>206</sup>	ε, γ	s	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Kidney	2 × 10 <sup>-7<sup>7</sup></sup>	6 × 10 <sup>-8</sup>	
356				i	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Lung	10 <sup>-7</sup>	5 × 10 <sup>-8</sup>	
357	83	Bi <sup>207</sup>	ε, γ	s	GI (LLI)	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	Kidney	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>	
358				i	GI (LLI)	2 × 10 <sup>-3</sup>	6 × 10 <sup>-4</sup>	Lung	10 <sup>-8</sup>	5 × 10 <sup>-9</sup>	
359	83	Bi <sup>210</sup>	α, β <sup>-</sup>	s	GI (LLI)	0.04 <sup>15</sup>	4 × 10 <sup>-4</sup>	Kidney	6 × 10 <sup>-9</sup>	2 × 10 <sup>-9</sup>	
360				i	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Lung	6 × 10 <sup>-9</sup>	2 × 10 <sup>-9</sup>	
361	83	Bi <sup>212</sup>	α, β <sup>-</sup> , γ	s	GI (S)	0.01 <sup>16</sup>	4 × 10 <sup>-3</sup>	Kidney	10 <sup>-7</sup>	3 × 10 <sup>-8</sup>	
362				i	GI (S)	0.01	4 × 10 <sup>-3</sup>	Lung	2 × 10 <sup>-7</sup>	7 × 10 <sup>-8</sup>	
363	84	Po <sup>210</sup>	α	s	Spleen	2 × 10 <sup>-5<sup>16</sup></sup>	7 × 10 <sup>-6</sup>	Spleen	5 × 10 <sup>-10<sup>12</sup></sup>	2 × 10 <sup>-10<sup>15</sup></sup>	
364				i	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-10</sup>	7 × 10 <sup>-11</sup>	
365	85	At <sup>211</sup>	α, ε, γ	s	Thyroid	5 × 10 <sup>-5<sup>24</sup></sup>	2 × 10 <sup>-5<sup>24</sup></sup>	Thyroid	7 × 10 <sup>-9<sup>24</sup></sup>	2 × 10 <sup>-9</sup>	
366				i	GI (ULI)	2 × 10 <sup>-3</sup>	7 × 10 <sup>-4</sup>	Lung	3 × 10 <sup>-8</sup>	10 <sup>-8</sup>	
367	86	Rn <sup>220<sup>25</sup></sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	...	...	...	...	Lung	3 × 10 <sup>-7</sup>	10 <sup>-7</sup>	
368	86	Rn <sup>222<sup>25</sup></sup>	α, β <sup>-</sup> , γ	...	...	...	...	Lung	3 × 10 <sup>-8</sup>	10 <sup>-8</sup>	
369	88	Ra <sup>223</sup>	α, β <sup>-</sup> , γ	s	Bone	2 × 10 <sup>-5</sup>	7 × 10 <sup>-6</sup>	Bone	2 × 10 <sup>-9</sup>	6 × 10 <sup>-10</sup>	
370				i	GI (LLI)	10 <sup>-4</sup>	4 × 10 <sup>-5</sup>	Lung	2 × 10 <sup>-10</sup>	8 × 10 <sup>-11</sup>	
371	88	Ra <sup>224</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Bone	7 × 10 <sup>-5</sup>	2 × 10 <sup>-5</sup>	Bone	5 × 10 <sup>-9</sup>	2 × 10 <sup>-9</sup>	
372				i	GI (LLI)	2 × 10 <sup>-4</sup>	5 × 10 <sup>-5</sup>	Lung	7 × 10 <sup>-10</sup>	2 × 10 <sup>-10</sup>	
373	88	Ra <sup>226</sup>	α, β <sup>-</sup> , γ	s	Bone	4 × 10 <sup>-7</sup>	10 <sup>-7</sup>	Bone	3 × 10 <sup>-11</sup>	10 <sup>-11</sup>	
374				i	GI (LLI)	9 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>	
375	88	Ra <sup>228</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Bone	8 × 10 <sup>-7</sup>	3 × 10 <sup>-7</sup>	Bone	7 × 10 <sup>-11</sup>	2 × 10 <sup>-11</sup>	
376				i	GI (LLI)	7 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	4 × 10 <sup>-11</sup>	10 <sup>-11</sup>	
377	89	Ac <sup>227</sup>	α, β <sup>-</sup> , γ	s	Bone	6 × 10 <sup>-5</sup>	2 × 10 <sup>-5</sup>	Bone	2 × 10 <sup>-12</sup>	8 × 10 <sup>-13</sup>	
378				i	GI (LLI)	9 × 10 <sup>-3</sup>	3 × 10 <sup>-3</sup>	Lung	3 × 10 <sup>-11</sup>	9 × 10 <sup>-12</sup>	
379	89	Ac <sup>228</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	GI (ULI)	0.05 <sup>8</sup>	9 × 10 <sup>-4</sup>	Liver	8 × 10 <sup>-8</sup>	3 × 10 <sup>-8<sup>13</sup></sup>	
380				i	GI (ULI)	3 × 10 <sup>-3</sup>	9 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-8</sup>	6 × 10 <sup>-9</sup>	
381	90	Th <sup>227</sup>	α, β <sup>-</sup> , γ	s	GI (LLI)	0.02 <sup>14</sup>	5 × 10 <sup>-4</sup>	Bone	3 × 10 <sup>-10</sup>	10 <sup>-10</sup>	
382				i	GI (LLI)	5 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	Lung	2 × 10 <sup>-10</sup>	6 × 10 <sup>-11</sup>	
383	90	Th <sup>228</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Bone	2 × 10 <sup>-4</sup>	7 × 10 <sup>-5</sup>	Bone	9 × 10 <sup>-12</sup>	3 × 10 <sup>-12</sup>	
384				i	GI (LLI)	4 × 10 <sup>-4</sup>	10 <sup>-4</sup>	Lung	6 × 10 <sup>-12</sup>	2 × 10 <sup>-12</sup>	
385	90	Th <sup>230</sup>	α, γ	s	Bone	5 × 10 <sup>-5</sup>	2 × 10 <sup>-5</sup>	Bone	2 × 10 <sup>-12</sup>	8 × 10 <sup>-13</sup>	
386				i	GI (LLI)	9 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	GI (LLI)	10 <sup>-11</sup>	3 × 10 <sup>-12</sup>	
387	90	Th <sup>231</sup>	α, β <sup>-</sup> , γ	s	GI (LLI)	7 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	5 × 10 <sup>-7</sup>	
388				i	GI (LLI)	7 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	GI (LLI)	10 <sup>-6</sup>	4 × 10 <sup>-7</sup>	
389	90	Th <sup>232</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Bone	5 × 10 <sup>-5</sup>	2 × 10 <sup>-5</sup>	Bone	2 × 10 <sup>-12<sup>28</sup></sup>	7 × 10 <sup>-13<sup>26</sup></sup>	
390				i	GI (LLI)	10 <sup>-3</sup>	4 × 10 <sup>-4</sup>	Lung	10 <sup>-11</sup>	4 × 10 <sup>-12</sup>	
391	90	Th <sup>234</sup>	β <sup>-</sup> , γ	s	GI (LLI)	4 <sup>14</sup>	5 × 10 <sup>-4</sup>	Bone	6 × 10 <sup>-8</sup>	2 × 10 <sup>-8</sup>	
392				i	GI (LLI)	5 × 10 <sup>-4</sup>	2 × 10 <sup>-4</sup>	Lung	3 × 10 <sup>-8</sup>	10 <sup>-8</sup>	
393	90	Th-nat	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	Bone	3 × 10 <sup>-5</sup>	10 <sup>-5</sup>	Bone	2 × 10 <sup>-12<sup>28</sup></sup>	6 × 10 <sup>-13<sup>26</sup></sup>	
394				i	GI (LLI)	3 × 10 <sup>-4</sup>	10 <sup>-4</sup>	Lung	4 × 10 <sup>-12</sup>	10 <sup>-12</sup>	
395	91	Pa <sup>230</sup>	α, β <sup>-</sup> , ε, γ	s	GI (LLI)	0.07 <sup>14</sup>	7 × 10 <sup>-3</sup>	Bone	2 × 10 <sup>-9</sup>	6 × 10 <sup>-10</sup>	
396				i	GI (LLI)	7 × 10 <sup>-3</sup>	2 × 10 <sup>-3</sup>	Lung	8 × 10 <sup>-10</sup>	3 × 10 <sup>-10</sup>	
397	91	Pa <sup>231</sup>	α, β <sup>-</sup> , γ	s	Bone	3 × 10 <sup>-5</sup>	9 × 10 <sup>-6</sup>	Bone	10 <sup>-12</sup>	4 × 10 <sup>-13</sup>	
398				i	GI (LLI)	8 × 10 <sup>-4</sup>	3 × 10 <sup>-4</sup>	Lung	10 <sup>-10</sup>	4 × 10 <sup>-11</sup>	
399	91	Pa <sup>233</sup>	β <sup>-</sup> , γ	s	GI (LLI)	40 <sup>15</sup>	4 × 10 <sup>-3</sup>	Kidney	6 × 10 <sup>-7</sup>	2 × 10 <sup>-7</sup>	
400				i	GI (LLI)	3 × 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	2 × 10 <sup>-7</sup>	6 × 10 <sup>-8</sup>	
401	92	U <sup>230</sup>	α, β <sup>-</sup> , γ	s	Kidney	0.01	7 × 10 <sup>-5</sup>	Kidney	3 × 10 <sup>-10</sup>	10 <sup>-10</sup>	
402				i	GI (LLI)	10 <sup>-4</sup>	5 × 10 <sup>-5</sup>	Lung	10 <sup>-10</sup>	4 × 10 <sup>-11</sup>	

<sup>1/</sup> Z = atomic number. <sup>2/</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3/</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4/</sup> For liver. <sup>5/</sup> Also lower large intestine. <sup>6/</sup> Also total body. <sup>7/</sup> For bone. <sup>8/</sup> For kidney. <sup>9/</sup> Also kidney. <sup>10/</sup> Also bone. <sup>11/</sup> Also ovary. <sup>12/</sup> The daughter elements of Rn<sup>220</sup> and Rn<sup>222</sup> are assumed present to the extent they occur in unfiltered air; for all other isotopes the daughter elements are not considered as part of the intake, and if present they must be considered on the basis of rules for mixtures. <sup>13/</sup> Provisional values.

continued



## 126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN

### Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES

Z <sup>1</sup>		Radionuclide			q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides					
		Symbol and Mass No.	Type of Decay	s or i		In Water			In Air		
						Critical Organ <sup>3</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$	Critical Organ <sup>3</sup>	40-hr wk $\mu\text{c/ml}$	168-hr wk $\mu\text{c/ml}$
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
403	92	U <sup>232</sup>	$\alpha, \beta^-, \gamma, e^-$	s	0.01	Bone	$2 \times 10^{-5}$	$8 \times 10^{-6}$	Bone	$10^{-10}$	$3 \times 10^{-11}$
404				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$3 \times 10^{-11}$	$9 \times 10^{-12}$
405	92	U <sup>233</sup>	$\alpha, \gamma$	s	0.05	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$5 \times 10^{-10}$	$2 \times 10^{-10}$
406				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
407	92	U <sup>234</sup>	$\alpha, \gamma$	s	0.05	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$6 \times 10^{-10}$	$2 \times 10^{-10}$
408				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
409	92	U <sup>235</sup>	$\alpha, \beta^-, \gamma$	s	0.03	Kidney	$10^{-4}$ <sup>15</sup>	$4 \times 10^{-5}$	Kidney	$5 \times 10^{-10}$	$2 \times 10^{-10}$ <sup>15</sup>
410				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
411	92	U <sup>236</sup>	$\alpha, \gamma$	s	0.06	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$6 \times 10^{-10}$	$2 \times 10^{-10}$
412				i	.....	GI (LLI)	$10^{-3}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
413	92	U <sup>238</sup>	$\alpha, \gamma, e^-$	s	0.005	Kidney	$2 \times 10^{-5}$	$6 \times 10^{-6}$	Kidney	$7 \times 10^{-11}$	$3 \times 10^{-11}$
414				i	.....	GI (LLI)	$10^{-3}$	$4 \times 10^{-4}$	Lung	$10^{-10}$	$5 \times 10^{-11}$
415	92	U-nat	$\alpha, \beta^-, \gamma, e^-$	s	0.005	Kidney	$2 \times 10^{-5}$	$6 \times 10^{-6}$	Kidney	$7 \times 10^{-11}$	$3 \times 10^{-11}$
416				i	.....	GI (LLI)	$5 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$6 \times 10^{-11}$	$2 \times 10^{-11}$
417	92	U <sup>240</sup> +	$\alpha, \beta^-, \gamma, e^-$	s	.....	GI (LLI)	$10^{-3}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$8 \times 10^{-8}$
418		Np <sup>240</sup>		i	.....	GI (LLI)	$10^{-3}$	$3 \times 10^{-4}$	GI (LLI)	$2 \times 10^{-7}$	$6 \times 10^{-8}$
419	93	Np <sup>237</sup>	$\alpha, \beta^-, \gamma$	s	0.06	Bone	$9 \times 10^{-5}$	$3 \times 10^{-5}$	Bone	$4 \times 10^{-12}$	$10^{-12}$
420				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
421	93	Np <sup>239</sup>	$\alpha, \beta^-, \gamma$	s	.....	GI (LLI)	$4 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$8 \times 10^{-7}$	$3 \times 10^{-7}$
422				i	.....	GI (LLI)	$4 \times 10^{-3}$	$10^{-3}$	GI (LLI)	$7 \times 10^{-7}$	$2 \times 10^{-7}$
423	94	Pu <sup>238</sup>	$\alpha, \gamma$	s	0.04	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$2 \times 10^{-12}$	$7 \times 10^{-13}$
424				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$3 \times 10^{-11}$	$10^{-11}$
425	94	Pu <sup>239</sup>	$\alpha, \gamma$	s	0.04	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$2 \times 10^{-12}$	$6 \times 10^{-13}$
426				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$4 \times 10^{-11}$	$10^{-11}$
427	94	Pu <sup>240</sup>	$\alpha, \gamma$	s	0.04	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$2 \times 10^{-12}$	$6 \times 10^{-13}$
428				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$4 \times 10^{-11}$	$10^{-11}$
429	94	Pu <sup>241</sup>	$\alpha, \beta^-, \gamma$	s	0.9	Bone	$7 \times 10^{-3}$	$2 \times 10^{-3}$	Bone	$9 \times 10^{-11}$	$3 \times 10^{-11}$
430				i	.....	GI (LLI)	0.04	0.01	Lung	$4 \times 10^{-8}$	$10^{-8}$
431	94	Pu <sup>242</sup>	$\alpha$	s	0.05	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$2 \times 10^{-12}$	$6 \times 10^{-13}$
432				i	.....	GI (LLI)	$9 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$4 \times 10^{-11}$	$10^{-11}$
433	94	Pu <sup>243</sup>	$\alpha, \beta^-, \gamma, e^-$	s	.....	GI (ULI)	0.01	$3 \times 10^{-3}$	GI (ULI)	$2 \times 10^{-6}$	$6 \times 10^{-7}$
434				i	.....	GI (ULI)	0.01	$3 \times 10^{-3}$	GI (ULI)	$2 \times 10^{-6}$	$8 \times 10^{-7}$
435	94	Pu <sup>244</sup>	$\alpha, \beta^-, \gamma, e^-$ (99.7%)	s	0.04	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$2 \times 10^{-12}$	$6 \times 10^{-13}$
436			SF (0.3%)	i	.....	GI (LLI)	$3 \times 10^{-4}$	$10^{-4}$	Lung	$3 \times 10^{-11}$	$10^{-11}$
437	95	Am <sup>241</sup>	$\alpha, \gamma$	s	0.1	Kidney	$10^{-4}$ <sup>15</sup>	$4 \times 10^{-5}$	Kidney	$6 \times 10^{-12}$ <sup>15</sup>	$2 \times 10^{-12}$ <sup>15</sup>
438				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
439	95	Am <sup>242m</sup>	$\alpha, \beta^-, \gamma, \epsilon, e^-$	s	0.07	Bone	$10^{-4}$ <sup>16</sup>	$4 \times 10^{-5}$	Bone	$6 \times 10^{-12}$ <sup>16</sup>	$2 \times 10^{-12}$ <sup>16</sup>
440				i	.....	GI (LLI)	$3 \times 10^{-3}$	$9 \times 10^{-4}$	Lung	$3 \times 10^{-10}$	$9 \times 10^{-11}$
441	95	Am <sup>242</sup>	$\alpha, \beta^-, \gamma, \epsilon, e^-$	s	0.06 <sup>3</sup>	GI (LLI)	$4 \times 10^{-3}$	$10^{-3}$	Liver	$4 \times 10^{-8}$	$10^{-8}$
442				i	.....	GI (LLI)	$4 \times 10^{-3}$	$10^{-3}$	Lung	$5 \times 10^{-8}$	$2 \times 10^{-8}$
443	95	Am <sup>243</sup>	$\alpha, \beta^-, \gamma$	s	0.05	Bone	$10^{-4}$ <sup>15</sup>	$4 \times 10^{-5}$	Bone	$6 \times 10^{-12}$ <sup>15</sup>	$2 \times 10^{-12}$ <sup>15</sup>
444				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
445	95	Am <sup>244</sup>	$\alpha, \beta^-, \gamma, e^-$	s	0.2 <sup>14, 16</sup>	GI (SI)	0.1	0.05	Bone	$4 \times 10^{-6}$ <sup>16</sup>	$10^{-6}$ <sup>16</sup>
446				i	.....	GI (SI)	0.1	0.05	GI (SI)	$2 \times 10^{-5}$ <sup>6</sup>	$8 \times 10^{-6}$ <sup>6</sup>
447	96	Cm <sup>242</sup>	$\alpha, \gamma$	s	0.05 <sup>3</sup>	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	Liver	$10^{-10}$	$4 \times 10^{-11}$
448				i	.....	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$2 \times 10^{-10}$	$6 \times 10^{-11}$
449	96	Cm <sup>243</sup>	$\alpha, \gamma$	s	0.09	Bone	$10^{-4}$	$5 \times 10^{-5}$	Bone	$6 \times 10^{-12}$	$2 \times 10^{-12}$
450				i	.....	GI (LLI)	$7 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$10^{-10}$	$3 \times 10^{-11}$
451	96	Cm <sup>244</sup>	$\alpha, \gamma$	s	0.1	Bone	$2 \times 10^{-4}$	$7 \times 10^{-5}$	Bone	$9 \times 10^{-12}$	$3 \times 10^{-12}$
452				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$3 \times 10^{-11}$
453	96	Cm <sup>245</sup>	$\alpha, \beta^-, \gamma$	s	0.04	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$5 \times 10^{-12}$	$2 \times 10^{-12}$
454				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
455	96	Cm <sup>246</sup>	$\alpha$	s	0.05	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$5 \times 10^{-12}$	$2 \times 10^{-12}$
456				i	.....	GI (LLI)	$8 \times 10^{-4}$	$3 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$
457	96	Cm <sup>247</sup>	$\alpha, \beta^-, \gamma, e^-$	s	0.04	Bone	$10^{-4}$	$4 \times 10^{-5}$	Bone	$5 \times 10^{-12}$	$2 \times 10^{-12}$
458				i	.....	GI (LLI)	$6 \times 10^{-4}$	$2 \times 10^{-4}$	Lung	$10^{-10}$	$4 \times 10^{-11}$

<sup>1/1</sup> Z = atomic number. <sup>2/2</sup> Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3/3</sup> That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4/4</sup> Also lung. <sup>5/5</sup> For liver. <sup>6/6</sup> For bone. <sup>7/7</sup> Also kidney. <sup>8/8</sup> Also bone.

*continued*

**126. MAXIMUM PERMISSIBLE OCCUPATIONAL EXPOSURE TO RADIATION: MAN**

**Part III. INTERNAL CONCENTRATION OF RADIONUCLIDES**

Z <sup>1</sup>	Radionuclide			q <sup>2</sup>	Maximum Permissible Concentrations of Radionuclides						
	Symbol and Mass No.	Type of Decay	s or i		In Water			In Air			
					Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml	Critical Organ <sup>3</sup>	40-hr wk μc/ml	168-hr wk μc/ml	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	
459	96	Cm <sup>248</sup>	α (89%)	s	0.005	Bone	10 <sup>-5</sup>	4 x 10 <sup>-6</sup>	Bone	6 x 10 <sup>-13</sup>	2 x 10 <sup>-13</sup>
460			SF (11%)	i	.....	GI (LLI)	4 x 10 <sup>-5</sup>	10 <sup>-5</sup>	Lung	10 <sup>-11</sup>	4 x 10 <sup>-12</sup>
461	96	Cm <sup>249</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	1 <sup>14</sup>	GI (S)	0.06	0.02	Bone	10 <sup>-5</sup> <sup>27</sup>	4 x 10 <sup>-6</sup>
462				i	.....	GI (S)	0.06	0.02	GI (S)	10 <sup>-5</sup>	4 x 10 <sup>-6</sup>
463	97	Bk <sup>249</sup>	α, β <sup>-</sup> , γ	s	0.7 <sup>14</sup>	GI (LLI)	0.02	6 x 10 <sup>-3</sup>	Bone	9 x 10 <sup>-10</sup>	3 x 10 <sup>-10</sup>
464				i	.....	GI (LLI)	0.02	6 x 10 <sup>-3</sup>	Lung	10 <sup>-7</sup>	4 x 10 <sup>-8</sup>
465	97	Bk <sup>250</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	0.05 <sup>14</sup>	GI (ULI)	6 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	Bone	10 <sup>-7</sup>	5 x 10 <sup>-8</sup>
466				i	.....	GI (ULI)	6 x 10 <sup>-3</sup>	2 x 10 <sup>-3</sup>	GI (ULI)	10 <sup>-6</sup>	4 x 10 <sup>-7</sup>
467	98	Cf <sup>249</sup>	α, γ	s	0.04	Bone	10 <sup>-4</sup>	4 x 10 <sup>-5</sup>	Bone	2 x 10 <sup>-12</sup>	5 x 10 <sup>-13</sup>
468				i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	10 <sup>-10</sup>	3 x 10 <sup>-11</sup>
469	98	Cf <sup>250</sup>	α	s	0.04	Bone	4 x 10 <sup>-4</sup>	10 <sup>-4</sup>	Bone	5 x 10 <sup>-12</sup>	2 x 10 <sup>-12</sup>
470				i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	10 <sup>-10</sup>	3 x 10 <sup>-11</sup>
471	98	Cf <sup>251</sup>	α, γ	s	0.04	Bone	10 <sup>-4</sup>	4 x 10 <sup>-5</sup>	Bone	2 x 10 <sup>-12</sup>	6 x 10 <sup>-13</sup>
472				i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	10 <sup>-10</sup>	3 x 10 <sup>-11</sup>
473	98	Cf <sup>252</sup>	α, γ, SF	s	0.01 <sup>14</sup>	GI (LLI)	2 x 10 <sup>-4</sup>	7 x 10 <sup>-5</sup>	Bone	6 x 10 <sup>-12</sup>	2 x 10 <sup>-12</sup>
474				i	.....	GI (LLI)	2 x 10 <sup>-4</sup>	7 x 10 <sup>-5</sup>	Lung	3 x 10 <sup>-11</sup>	10 <sup>-11</sup>
475	98	Cf <sup>253</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	0.04 <sup>14</sup>	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Bone	8 x 10 <sup>-10</sup>	3 x 10 <sup>-10</sup>
476				i	.....	GI (LLI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	8 x 10 <sup>-10</sup>	3 x 10 <sup>-10</sup>
477	98	Cf <sup>254</sup>	SF	s	0.0007 <sup>14</sup>	GI (LLI)	4 x 10 <sup>-6</sup>	10 <sup>-6</sup>	Bone	5 x 10 <sup>-12</sup>	2 x 10 <sup>-12</sup>
478				i	.....	GI (LLI)	4 x 10 <sup>-6</sup>	10 <sup>-6</sup>	Lung	5 x 10 <sup>-12</sup>	2 x 10 <sup>-12</sup>
479	99	Es <sup>253</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	0.04 <sup>14</sup>	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Bone	8 x 10 <sup>-10</sup>	3 x 10 <sup>-10</sup>
480				i	.....	GI (LLI)	7 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	6 x 10 <sup>-10</sup>	2 x 10 <sup>-10</sup>
481	99	Es <sup>254m</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	0.02 <sup>14</sup>	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Bone	5 x 10 <sup>-9</sup>	2 x 10 <sup>-9</sup>
482				i	.....	GI (LLI)	5 x 10 <sup>-4</sup>	2 x 10 <sup>-4</sup>	Lung	6 x 10 <sup>-9</sup>	2 x 10 <sup>-9</sup>
483	99	Es <sup>254</sup>	α, β <sup>-</sup> , γ, e <sup>-</sup>	s	0.02 <sup>14</sup>	GI (LLI)	4 x 10 <sup>-4</sup>	10 <sup>-4</sup>	Bone	2 x 10 <sup>-11</sup>	6 x 10 <sup>-12</sup>
484				i	.....	GI (LLI)	4 x 10 <sup>-4</sup>	10 <sup>-4</sup>	Lung	10 <sup>-10</sup>	4 x 10 <sup>-11</sup>
485	99	Es <sup>255</sup>	α, β <sup>-</sup> , γ	s	0.04 <sup>14</sup>	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Bone	5 x 10 <sup>-10</sup>	2 x 10 <sup>-10</sup>
486				i	.....	GI (LLI)	8 x 10 <sup>-4</sup>	3 x 10 <sup>-4</sup>	Lung	4 x 10 <sup>-10</sup>	10 <sup>-10</sup>
487	100	Fm <sup>254</sup>	α, γ, e <sup>-</sup>	s	0.02 <sup>14</sup>	GI (ULI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Bone	6 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
488			SF (99.9448%)	i	.....	GI (ULI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	7 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
			SF (5.52 x 10 <sup>-2</sup> %)	i	.....	GI (ULI)	4 x 10 <sup>-3</sup>	10 <sup>-3</sup>	Lung	7 x 10 <sup>-8</sup>	2 x 10 <sup>-8</sup>
489	100	Fm <sup>255</sup>	α, γ	s	0.04 <sup>14</sup>	GI (LLI)	10 <sup>-3</sup>	3 x 10 <sup>-4</sup>	Bone	2 x 10 <sup>-8</sup>	6 x 10 <sup>-9</sup>
490				i	.....	GI (LLI)	10 <sup>-3</sup>	3 x 10 <sup>-4</sup>	Lung	10 <sup>-8</sup>	4 x 10 <sup>-9</sup>
491	100	Fm <sup>256</sup>	SF	s	0.0008 <sup>14</sup>	GI (ULI)	3 x 10 <sup>-5</sup>	9 x 10 <sup>-6</sup>	Bone	3 x 10 <sup>-9</sup>	10 <sup>-9</sup>
492				i	.....	GI (ULI)	3 x 10 <sup>-5</sup>	9 x 10 <sup>-6</sup>	Lung	2 x 10 <sup>-9</sup>	6 x 10 <sup>-10</sup>

<sup>1</sup>/ Z = atomic number. <sup>2</sup>/ Maximum permissible burden in the total body resulting from maximum permissible concentration of the radionuclide in water or food when deposited in the critical organ (columns F and I). When other footnote numbers appear in column E, "q" pertains only to the critical organ specified in the footnote. <sup>3</sup>/ That organ receiving the radiation dose that results in the greatest damage to the body. <sup>4</sup>/ For bone. <sup>5</sup>/ Also stomach.

*Contributor:* Morgan, Karl Z.

*References:* [1] International Commission on Radiological Protection. 1959. Radiation protection, II. Pergamon Press, New York. [2] International Commission on Radiological Protection. 1964. Radiation protection, IV. Pergamon Press, New York.

## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

**Radiation** (columns A and B): c = curie, the quantity of radioactive nuclide in which the number of disintegrations is  $3.7 \times 10^{10}$ /sec; mc = millicurie, 1/1,000 of a curie or  $3.7 \times 10^7$  disintegrations/sec;  $\mu\text{c}$  = microcurie, one millionth of a curie or  $3.7 \times 10^4$  disintegrations/sec; ev = electron volt, the energy imparted to an electron when it is accelerated through a potential difference of one volt ( $1.602 \times 10^{-12}$  ergs); Mev = one million electron volts, a unit of energy equal to  $1.6 \times 10^{-6}$  ergs; kv = kilovolt, a unit of electrical potential equal to 1,000 volts; n = neutron, a nuclear particle of zero charge and mass number 1; r = roentgen, the quantity of X- or  $\alpha$ -radiation such that the associated corpuscular emission per 0.001293 g of air produces, in air, ions carrying one electrostatic unit of electrical charge of either sign; rep = roentgen equivalent physical, the equivalent of 93 ergs/g energy absorption; rem = roentgen equivalent man (or mammal), the quantity of radiation absorbed in tissue, which gives the same observable effect as one rep of X or  $\gamma$  rays; rad = unit of absorbed dose (1 rad corresponds to 100 ergs/g of medium); RBE = relative biological effectiveness, i.e., ratio of absorbed dose, in rads, from reference X rays to the absorbed dose, in rads, from the given radiation field required to produce the same effect as the reference X rays (reference X rays in most cases have been those from 200-250 kv X-radiation or  $\gamma$ -radiation from  $\text{Co}^{60}$ ). **Late Effects** (column D): figures separated by a slash (e.g., 13/16) give the number of individuals affected and number exposed. **Symbols**: > = greater than;  $\geq$  = greater than or equal to;  $\leq$  = less than or equal to.

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Man					
1	A-bomb	Epilation dose	2 yr	A-bomb cataract: Japan, 10 cases; USA, 1 case.	13,15
2	Hiroshima	Irradiated in utero, 1st half of pregnancy	5 yr	Microcephaly and mental retardation, 7/11 children within 1,200 m of hypocenter.	58
3	Japan	Area within 2,000 m of hypocenter	3-5 yr	Leukemia incidence 9.3 times that of nonexposed population of Hiroshima and Nagasaki (1948-1950, inclusive).	23
4		Area within 1,000 m of hypocenter	3-5 yr	Leukemia incidence 32 times that of nonexposed population (1948-1950, inclusive).	
5	Cyclotron 16 Mev neutrons, fast	Epilation dose	2 yr	Cyclotron-induced cataract, 2 cases.	13,15
6		400-500 n	2 mo-5 yr	Severe epidermolytic reaction, 13/16: skin atrophy and fibrosis, persistent ulcerations, diminished repair by normal tissues. Radiation osteitis. Severe bowel reactions.	69
7	0-20 Mev neutrons, fast (small $\gamma$ components)	10-135 n	2-10 yr	Cataracts: severe, 3/10; slight to moderate, 4/10; minimal, 3/10. Chronic irradiation: no blood changes; mild epilation, 2 cases.	1
8	$\gamma$ rays and fission neutrons	22.8-365 rads to total body	29 mo	Gross chromosome aberrations of leukocyte cultures, % abnormal cells: 22.8 rads, 4%; 339 rads, 28.2%; 365 rads, 18%. Minor aberrations of grossly normal cells: 22.8 rads, 83%; 339 rads, 53%; 365 rads, 67%.	4,5
9			42 mo	Gross chromosome aberrations of leukocyte cultures, % abnormal cells: 22.8 rads, 20%; 339 rads, 19%; 365 rads, 23%. Minor aberrations of grossly normal cells: 22.8 rads, 0%; 339 rads, 50%; 365 rads, 30%.	
10	Nuclear and allied radiations		6-8 yr	Peak incidence of leukemia.	49
11			After 10th yr	Diminishing incidence of leukemia; increasing incidence of other cancers.	
12	Radium (external)	1,000-1,500 mg-hr		Cessation of ovarian function, 77%/63.	57
13		1,500-2,000 mg-hr		Cessation of ovarian function, 6/7.	
	Radium <sup>1</sup>	Residual (ingested)			2,48, 75, 76
14		0.02-0.5 $\mu\text{g}$	14-48 yr	Radiation osteitis, 25%.	
15		0.5-2.0 $\mu\text{g}$	1-32 yr	Osteomyelitis of jaw and loss of teeth, 5/9; radiation osteitis, 8/9; pathological fractures, 3/9; giant cell tumor, 1/9; osteogenic sarcoma, 1/9; epidermoid carcinoma of nasopharynx, 1/9. High incidence of deafness and arthritis.	
16		2.7 $\mu\text{g}$	8-32 yr	Osteomyelitis of jaw and loss of teeth, 5/9; radiation osteitis, 8/9; pathological fracture, 1/9; fibrosarcoma, 1/9; epidermoid carcinoma of nasopharynx, 1/9.	
17		2-20 $\mu\text{g}$	6-8 yr	Radiation osteitis (osteosclerosis); osteogenic sarcoma deaths, 5/18; pathological fractures.	

<sup>1</sup>/ Variable amounts of mesothorium also present in some instances.

*continued*



## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Man					
18	Radium <sup>1</sup>	Residual (ingested) 8-23 µg	7-21 yr	Osteomyelitis of jaw and loss of teeth, 6/8; radiation osteitis, 7/8; epidermoid carcinoma of nasopharynx, 1/8; osteogenic carcinoma, 1/8; pathological fractures, 3/8; leukemia, 1/8.	2,48, 75, 76
19		10-180 µg	1-8 yr	Anemia with hyperplastic marrow, jaw necrosis (13 cases).	
20		Residual, 1.0-10.0 µg (equivalent to 100-800 µg original dose)	20-30 yr posttreatment	24 cases: changes in bone density (similar to those in dead or dying bone) in patients having at least 1 µg; minimal changes with 0.5 µg; dental changes in all with at least 4 µg. Greatly enlarged Haversian canals. Distortion of normal bone configuration, 7/24; edentia, mandibular lesions, 3/19; aseptic necrosis of bone, 7/24; fibrosarcoma, honeycombed teeth (pink tooth).	43
21	Uranium, radium ores	Variable doses	13-23 yr	Lung cancer in uranium miners of Joachimsthal and Schneeberg.	28,40
22	X ray	Variable doses	Months to years	Skin atrophy, telangiectasis, sclerosis, pigmentation, alopecia, altered vasomotion, diminished sweat and sebaceous function, loss of cutaneous ridges and fingerprints, ulcers and keratoses, malignancies, hyperkeratotic warty growths, deformed and brittle dry nails, loss of nails, fissures, subungual hyperkeratoses.	3,28, 51, 75, 76
23		Varied exposures (diagnostic radiology)		One leukemic death for every 46,000 X-ray examinations in Great Britain.	72
24		Irradiation in utero		Incidence of leukemia and cancer per 10,000 children	45
25		No exposure		Leukemia--firstborn, 4.7; others, 3.6; all, 4.0. Cancer of CNS, 1.6; other cancers, 1.8.	
26		Exposed		Leukemia--firstborn, 6.9; others, 4.6; all, 6.1. Cancer of CNS, 2.5; other cancers, 2.5.	
27		1 or 2 films		Leukemia, 3.8; cancer of CNS, 2.7; other cancers, 2.7.	
28		Pelvimetry		Leukemia, 6.9; cancer of CNS, 2.3; other cancers, 2.1.	
29		100-300 r (200 & 1,000 kv)	60-680 da	Temporary drop in WBC count.	70
30		500-624 r	100-600 da	Temporary macrocytic anemia.	
31		625 r to ovaries		94% castrated.	57
32		≤1,000 r to center of vertebrae	Followed 13 yr post-treatment	Permanent cessation of menstruation, 72 patients. Vertebrae normal (irrespective of age), 45 children.	50,75, 76
33		1,000-2,000 r to spine	Followed 3-7 yr	"Transverse-line" growth disturbance of vertebrae (irrespective of age up to 6 yr), 45 children.	
34		Approximately 2,000 r to spine	Followed 2-13 yr	Contour irregularity of vertebrae, 44/45 children.	
35		1,500-8,500 r (therapeutic dose, eye tumor)	2 yr	Cataract, 7 cases.	13,15
36		3,500-5,000 r	8-28 mo	Lens opacities, 4 cases.	41
37		1,500-25,000 r (130-200 kv)	6-22 yr	Osteogenic sarcoma, 11 cases.	12
38		1,700-3,000 r (half-value layer 1.5 cu mm) to lower abdomen	6-18 mo	Nephrosclerosis, hypertension, elevated albuminuria, 22/55. Edema, anuria. Death from congestive heart failure and/or uremia, 7/55. Over 2,300 r, high risk of renal failure.	33
39	X ray, radium	4,000-6,700 r	5-20 yr	Sarcoma; osteogenic sarcoma. Therapy for lupus vulgaris, papillomata of bladder, actinomycosis, tubercular psoriasis.	30

/1/ Variable amounts of mesothorium also present in some instances.

continued



## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Man					
40	X-ray and radium equipment	Unknown	Leukemia in radiologists (4.68%), 9 times the incidence in nonradiological physicians (0.51%).	20,47, 74	
Dog					
41	Neutrons	0.012, 0.06, 0.11 n/da	1 yr	Reduction of lymphocytes only observed effect (dose given 6 da/wk).	7
42		1.7 n/da	1 yr	Mucoid conjunctivitis, keratoconjunctivitis and corneal opacities; reduced size of spleen and testes; increased incidence of infection; hypoplasia of bone marrow and regional lymph nodes; hemorrhage of lymph nodes, heart, stomach, small bowel and kidney; reduction of lymphocytes, neutrophils, and erythrocytes.	
43	Neutrons, fast	150 n	2 yr	Destruction and chronic inflammation of cornea, and changes in lens capsules and fibers, but no cataracts.	53
44		800-900 n	2 yr	Cataracts 60-75% of subjects.	
45	X ray	0.1 r/da (6 times/wk)	2 yr	Lowered sperm count, increase in abnormal sperm.	7
46		0.5 r/da	11 mo	Lowered sperm count, increase in abnormal sperm.	
47			2 yr	Partial testicular atrophy; slight reduction of lymphocytes.	
48		1.0 r/da	6 mo	Lymphopenia.	
49			9 mo	50% aspermic <sup>a</sup> .	
50			1 yr	100% aspermic <sup>a</sup> ; neutrophil reduction.	
51			2 yr	Severe injury of testes <sup>a</sup> .	
52		3.0 r/da	2 mo	Lymphocyte and platelet reduction.	
53		6.0 r/da	9 mo	50% aspermic.	
54			1.5 yr	Severe injury of testes.	
55		10.0 r/da	2 mo	Lymphocyte, platelet, & erythrocyte depression.	
56			3.5 mo	50% aspermic.	
57			1.5 yr	Bone marrow hypoplasia; focal bowel and lymph node hemorrhages.	
58			6 mo	50% survival.	
Guinea Pig					
59	Cobalt-60	15 r/da	106 da	50% survival.	73
60	(γ rays), external	30 r/da	63 da		
61		60 r/da	41 da		
62		90 r/da	20 da		
63		120 r/da	18 da		
64	Phosphorus-32	7,750 rep (external)	2 mo	Alopecia.	78
65	Radium, filtered	Nonradiated (controls)	38 mo	75% survival.	29,44,
66	(γ rays), repeated low dose	0.11 r/da	38 mo	75% survival.	70
67		1.1 r/da (1,050 r)	32 mo	50% survival.	
68		2.2 r/da (2,100 r)	32 mo	50% survival.	
69		4.4 r/da (2,400 r)	18 mo	50% survival; reduction of WBC.	
70		8.8 r/da (2,300 r)	5 mo	50% survival; recurrent anemia.	
Mouse					
71	Carbon-14	100 μc sodium formate of C-14, intraperitoneally during gestation	Until parturition	Normal litters; offspring show no differences from controls in weight, longevity, or tumor incidence. No lowered fertility.	68
72	Cesium-137 (γ rays)	258 r to total body before mating	Until parturition	Reduced size of litters: age at mating, 60-79 da--controls, 14.08; irradiated, 9.16; age at mating, 80-99 da--controls, 30.8; irradiated, 8.63; age at mating, 110-119 da--controls, 12.8; irradiated, 7.68; age at mating, 120-139 da--controls, 12.05; irradiated, 6.59.	26
73	Cobalt-60	90 r/da	52 da	50% survival.	73
74	(γ rays), external	115 r/da	37 da		
75		140 r/da	22 da		

<sup>a</sup>/ If irradiation is stopped after one year, there is partial recovery of sperm four months later.

continued

127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference
Type	Exposure			
(A)	(B)	(C)	(D)	(E)
Mouse				
76	Deuteron beams Brain region 1-mm beam at 8,000 rads to cerebellum	2-7 mo	Significant variation in locomotor activity wheel tests & escape-avoidance conditioning.	56
77	1-mm beam at 8,000 rads to both parietal regions	7 mo	Large lesions at end of Bragg zone owing to disappearance of nerve cell bodies, leaving fluid-filled cavities without structural detail.	
78	Neutrons, fast 2 rads/da, 2.24 neutron high, 0.287 neutron low; $\gamma$ -irradiation--16 rads/da, 15.8 high, 2.26 low		Life shortening; RBE 10.	55
79	Neutrons, fast, or Cobalt-60 ( $\gamma$ rays) 3-50 rems daily		Life shortening; RBE independent of exposure time and may vary with dose rate.	52
80	Neutrons, fast (4 Mev); X ray (200 kv) 1 exposure		Cataractogenesis; RBE 4.	18,62,
81	3 exposures		Cataractogenesis; RBE 6.	63
82	Neutrons, fast, $10^3$ ev to $\geq 4$ Mev, and $\gamma$ rays (divided doses) Nonradiated (controls)	51 wk	50% survival for controls and each dosage increment. For shortening life span, 1 n (divided small doses) is equivalent to 35 r. (For acute killing, 1 n is equivalent to 9 r.) Threshold for shortening life span is approximately 1 r/da and less than 0.1 n.	27
83	1 r/da	61 wk		
84	8.6 r/da	48 wk		
85	Nonradiated (controls)	70 wk		
86	0.115 n/da (total 32.2 n)	40 wk		
87	1.15 n/da (total 241.5 n)	30 wk		
88	4.3 n/da (total 301 n)	10 wk		
89	13 n/da (total 273 n)	3 wk		
90	Neutrons, fast, and X rays (single exposure) Nonradiated (controls)	64 wk	50% survival for controls and each dosage increment. Terminal changes generalized. Atrophy. Increased incidence of mediastinal lymphomatosis.	27
91	500 r	58 wk		
92	700 r	39 wk		
93	26 n	52 wk		
94	50 n	48 wk		
95	78 n	42 wk		
96	90 n	6 wk		
97	Phosphorus-32 3,000-4,000 rep (external)	30 da	General epilation, skin atrophy, loss of ear tips, ulcerations, keratosis.	61,78
98		3-4 mo		
99		8 mo		
100		6-8 mo		
101	Plutonium-239 3.1-15.6 $\mu\text{c/kg}$ (iv)	190-250 da	Osteogenic sarcoma.	11,75,
102	>6.1 $\mu\text{c/kg}$ (iv)		Marked shortening of life span.	76
103	Radium-226 >12 $\mu\text{c/kg}$ (iv)	250-300 da	Increased osteogenic sarcoma.	11,75,
104	>50 $\mu\text{c/kg}$ (0.6-4,170 $\mu\text{c/kg}$ , iv)		Marked shortening of life span; debilitation and increased incidence of infection.	76
105	Radium, filtered ( $\gamma$ rays), repeated low dose Nonradiated (controls)	24 mo	50% survival for controls and each dosage increment; increase in lymphatic leukemia, mammary and ovarian carcinoma.	44
106	1.1 r/da (760 r)	24 mo		
107	2.2 r/da (1,390 r)	22 mo		
108	4.4 r/da (2,640 r)	21 mo		
109	8.8 r/da (4,400 r)	17 mo		
110	Strontium-89 ( $\beta$ rays) Monthly injections 0.05 $\mu\text{c/g}$	500 da	Bone tumors begin to develop.	10
111	0.1 $\mu\text{c/g}$	425 da		
112	0.2 $\mu\text{c/g}$	350 da		
113	0.5 $\mu\text{c/g}$	220 da		
114	1.0 $\mu\text{c/g}$	160 da		
115	Single injection 2.5 $\mu\text{c/g}$	200 da		
116	5.0 $\mu\text{c/g}$	150 da		
117	Uranium-232 0.1-1.0 $\mu\text{c/kg}$ (iv)	575 da	Osteogenic sarcoma (probably significant); used 0.1-10.0 $\mu\text{c/kg}$ .	11,75,
118	>1.0 $\mu\text{c/kg}$ (iv)	575 da		
119	Uranium-233 2-5 $\mu\text{c/kg}$ (iv)	575 da	Osteogenic sarcoma (probably significant); used 0.1-100 $\mu\text{c/kg}$ .	11,75,
120	>5 $\mu\text{c/kg}$ (iv)	575 da		
121	>53 $\mu\text{c/kg}$ (iv)	575 da	Shortening of life span.	
122	Uranium-238 >3.6 $\times 10^{-7}$ $\mu\text{c/kg}$ (iv)	575 da	Possibly significant increase in osteogenic sarcoma.	11,75,
				76

*continued*

## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Mouse					
123	X ray	25-200 r to fetus early in pregnancy (equivalent to 2-6 wk, human)	Until birth	At birth, severe skeletal and other abnormalities, with clearly defined critical periods.	65
124		Irradiation of fetus late in pregnancy (equivalent to after 6 wk, human)	Until birth	Several weeks after birth: cataracts, hydrocephalus, behavior disturbances, skin lesions.	
125		400 r (single dose)	15-23 mo	Ovarian tumors 15 times that of controls.	25
126		600-1,000 r (divided doses)	9-14 mo	Myelosis 8 times that of controls.	
127			6-12 mo	Mediastinal lymphosarcoma 7 times that of controls.	
128	X ray (120 kv)	Nonradiated (controls)	526-570 da	Lymphomas, 3/36	31
129		600 r (150 r alternate weeks) to whole body	420-580 da	Lymphomas, 33/63.	
130		600 r (150 r alternate weeks) to whole body, mediastinum shielded	118-525 da	No lymphomas.	
131		1,200 r (300 r alternate weeks) to upper half of body	324-365 da	Lymphomas, 3/40.	
132		1,200 r (300 r alternate weeks) to alternate halves of body	420-508 da	Lymphomas, 3/37.	
133	X ray (250 kv)	50 r	17 wk	Lens opacities, 59% at 30 wk <sup>a</sup> .	71
134		200 r	14 wk	Lens opacities, 100% at 22 wk <sup>a</sup> .	
135		500 r	6 wk	Lens opacities, 100% at 17 wk <sup>a</sup> .	
136		100 r to whole body	3 mo	Shortening of life span, 7%.	17
137		100-300 r to whole body	3 mo	Increase in thymic lymphoma.	
138		300 r to whole body	3 mo	Increase in myeloid leukemia; shortening of life span, 21%.	
139		300 r to head	3 mo	Ocular lens cataracts, 40%; shortening of life span, 2%.	
140		300 r to middle third of body	3 mo	Shortening of life span, 3%.	
141		300 r to lower third of body	3 mo	Increase of ovarian tumors; shortening of life span, 4%.	
142		260-520 r (1-4 exposures, males & females 164-730 da old)	Death	Life span shortened 4.6-26.2% for males, 8.8-36.5% for females.	32
143		550 r to whole body	5-6 mo	Decrease in fat of skin of female; permanent depression of DNA synthesis relative to RNA in kidneys of both sexes.	24
144		500 r, 550 r, 600 r (females 10 wk old)		Increased incidence of thymic lymphomas and ovarian tumors; greater severity of lens opacities than in controls.	16
145		750 r to whole body	6-10 mo	Weight reduction in body, heart, kidneys, and testes; increase in fat content of carcass.	22
Rat					
146	Cerium-144	1-3 mc/kg	200 da	Osteogenic sarcoma, liver atrophy with ascites and jaundice.	42
147	Cobalt-60	20 r/da	332 da	50% survival; increased tumor frequency.	73
148	(γ rays), external	60 r/da	236 da		
149		70 r/da	72 da		
150		80 r/da	53 da		
151		90 r/da	48 da		
152		120 r/da	38 da		
153	Cobalt-60 (γ rays)	110 r (13-15th & 16-18th da of gestation)	Until parturition	Neonatal death rate higher.	54
154		150 r (14th & 20th da of gestation)			

<sup>a</sup>/s/ The RBE of thermal neutrons calculated on the basis of the production of lens opacities was approximately nine times the RBE calculated on the basis of the production of lethality at 30 days.

*continued*

## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Rat					
155	Cobalt-60 (γ rays)	220 r (12-16th & 17-20th da of gestation)	Until parturition	Neonatal death rate higher; average weaning weight lower in all irradiated groups; ovulation 20% in irradiated females; mature body and testes weight lower in irradiated males.	54
156	Neutrons	0.012-0.060 n/da	1 yr	No effects observed.	7
157		0.11 n/da	1 yr	Number of neoplasms 3 times that of controls (including leukemias).	
158		1.7 n/da	1 yr	Increased infection and incidence of neoplasms (high leukemia incidence); bilateral cataracts; hypoplasia of spleen; atrophy of testes and ovarian follicles; early reduction of lymphocytes; reduction of erythrocytes.	
159	Phosphorus-32	4,000-5,000 rep (external)	4-5 mo	Tumors, keratoses, lens opacities (low incidence), ulceration of scrotum and base of tail, alopecia.	78
160	Praseodymium-144	1-3 mc/kg	200 da	Osteogenic sarcoma, liver atrophy with ascites and jaundice.	42
161	Plutonium-239	0.02 μg/g (0.0031-0.0062 c/g)	300-400 da	Osteogenic sarcomas.	60
162		0.03 μc/g	3-7 mo	Areas of dead bone and calcified cartilage; resumption of normal bone growth at epiphysis; destruction of spermatogenic cells, atrophy of ovary.	
163	Radium	0.125 μc/g	5 mo	Damage to epiphyseal cartilage, overgrowth with atypical bone, loss of normal bone cells; atrophy of ovary.	8,75, 76
164		0.5 μc/g	3-6 mo	Degenerative changes in ovary; damage to blood vessels.	
165		0.6 μc/g	5 mo	Damage to epiphyseal cartilage and marrow; production of abnormal bone.	
166	Strontium-89	0.05 μc/g	500 da	Osteogenic sarcomas.	60
167		0.1 μc/g	400 da		
168		0.5 μc/g (single injection)	400 da		
169		1.0 μc/g level (monthly injections)	250 da		
170		5.0 μc/g (single injection)	200 da		
171	Strontium-90	10-30 da in drinking water Total dose, 33 μc; skeletal burden, 1 μc (425 da old)	380 da	No leukemia; no osteogenic sarcoma.	26
172		Total dose, 650 μc; skeletal burden, 2 μc (346 da old)	372 da	0.0007 leukemia/rat/wk; no osteogenic sarcoma.	
173		Total dose, 790 μc; skeletal burden, 11 μc (117 da old)	254 da	Moderate bone marrow damage; 0.0017 leukemia/rat/wk; 0.0076 osteogenic sarcoma/rat/wk.	
174		Total dose, 464 μc; skeletal burden, 33 μc (40 da old)	106 da	Atrophy of bone marrow cells; no leukemia; 0.023 osteogenic sarcoma/rat/wk.	
175	Thorium dioxide (α and γ rays)	0.3 ml	14 mo	Produced fibroblastic tumors, 14/60.	28
176		2.5 ml	10-17 mo	Produced sarcoma, 33/50.	
177		5.0 ml	10-17 mo	Produced sarcoma, 50/50.	
178	X ray	0.1 r/da (total 49 r)	81.7 wk	50% survival.	7
179		0.5 r/da (total 230 r)	76.7 wk	50% survival; increase in fibroadenoma of mammary gland.	
180		1.0 r/da (total 460 r)	76.9 wk	50% survival.	
181		10.0 r/da (total 3,500 r)	58.3 wk	50% survival; increase in leukemia.	
182		12.5-100.0 r (8th da of gestation)		Retardation of growth.	
183		12.5-100.0 r (9th & 10th da of gestation)		Malformations, increased mortality, growth retardation.	
184		≥375 r to whole body		Enamel hypoplasia, retarded enamel dentin formation.	3,75, 76

continued



## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference
Type	Exposure			
(A)	(B)	(C)	(D)	(E)
Rat				
185	X ray	≥2,000 r (locally to teeth)	Retardation of eruption of incisors.	3,75.
186		4,000 r (locally to teeth)	Stoppage of growth of dentin (lengthwise) in incisors.	76
187		500 r to whole body	5 months required for complete restoration of epithelium in testes (Vanderbilt's strain, adult males).	66
188	X ray (250 kv)	25-400 r to whole body (40 da old)	10.5-11 mo Incidence of mammary gland neoplasia linear with doses between 25 and 400 r; above 400 r, incidence is either decreased or remains constant.	9
189		120-480 r to whole body (1-6 exposures, 3 mo old)	Death Mean longevity: controls, 28.6 mo; 120 r, 24.9 mo; 240 r, 23.1 mo; 480 r, 20.2 mo.	39
190		400 r to whole body (either castrated or intact male, 40 da old)	16 mo <sup>4</sup> 50% incidence of breast neoplasms (more fibrosarcomas); total incidence lower than in irradiated females.	67
191		400 r (female 40 da old)	At end of 10 mo 79% tumor incidence, with one or more breast neoplasms.	19
192		773±88 r to abdomen only	420 da 50% mortality.	21
193		943±60 r to head only	168 da	
194		1,062±12 r to thorax only	90 da	
195		1,000 r (under 5% oxygen pressure) to whole body	Up to 500 da Shortening of life span, nephrosclerosis, generalized arteriosclerosis, hypertension, thrombocytopenia, anemia, increased susceptibility to oral hypertonic sodium chloride, earlier appearance of neoplasms.	6,34-38
196	Yttrium-91	2.0 µc/g	1-3 mo Damage to epiphyseal cartilage, production of atypical bone; increase in spleen hematopoiesis.	8
197		20-30 mc/kg (1 dose orally), or 1-2 mc/kg/da, 6 da/wk for 3 mo	200-400 da A variety of intestinal lesions with obstruction. Yttrium essentially not absorbed from intestinal tract.	42
Rabbit				
198	Neutrons	0.012-0.11 n/da, 6 doses/wk	1 yr No observed effects.	7
199		1.7 n/da	1 yr Increased incidence of infection; hypoplasia of bone marrow; atrophy of testes; reduction of lymphocytes; neutrophils depressed after 8 wk, erythrocytes after 32-35 wk.	
200		3.7 n/wk (52.7-83.7 n, total)	4-12 mo No lens changes.	53
201		33-100 n (single doses)	2-5 mo Cataracts of posterior lens.	
202	Fission	3 x 10 <sup>10</sup> particles/ml	125 da Cataracts (2 x 10 <sup>9</sup> particles/ml, or less, is threshold for lens opacities).	14
203	14 Mev	8 x 10 <sup>10</sup> particles/ml	125 da	
204	Phosphorus-32 (external)	2,500-3,000 rep	2 mo Graying of fur.	78
205		5,000 rep	Epilation; recovery by 10 wk.	
206		7,500 rep	Permanent epilation, ulcerations.	
207		15,000 rep	Emaciation.	
208	Radium	100 µg in 90 da (RaSO <sub>4</sub> in glycerine, orally)	2 mo Mottling and shortening of incisors.	28,64
209			5-18 mo Pathological fractures, alopecia.	
210			9 mo Jaw abscesses; rarefaction of mandible and other bones (generalized osteoporosis and necrosis); weight loss; mild, progressive, regenerative anemia, with hyperplasia and fibrosis of marrow, lymph nodes, spleen.	
211	Strontium-90	100 µc/kg (iv)	90 da % of controls: total body wt, 92%; wt of femur, 81%; wt of tibia, 91%; length of tibia, 98%.	46
212		600 µc/kg (iv)	90 da % of controls: total body wt, 34%; wt of femur, 71%; wt of tibia, 70%; length of tibia, 69%.	
213	Thorium dioxide (α and γ rays)	Intravenous injection	Osteosarcoma; hematopoietic depression and damage to liver and spleen. Deposited in reticuloendothelial system.	59

<sup>4</sup>/ Longer interval than for irradiated females.

continued

## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiation		Latent Period	Late Effects	Reference	
Type	Exposure				
(A)	(B)	(C)	(D)	(E)	
Rabbit					
214	X ray	0.1-0.5 r/da	1 yr	No changes detected.	7
215		1.0 r/da	1 yr	Possible testicular injury.	
216		10 r/da	8 wk	Lymphocytes significantly decreased.	
217			3 mo	Platelets decreased.	
218			1 yr	Erythrocytes decreased; testicular injury; ovarian follicles disappear.	
219		≤250 r (1.2 Mev)	150 da	Threshold for production of lens opacities.	14
220		500 r	125 da	Cataracts.	

*Contributors:* (a) Bennett, L. R., and Chastain, Sarah, (b) Billings, Marta S., (c) Thomson, John F.

*References:* [1] Abelson, P. H., and P. G. Kruger. 1949. *Science* 110:655. [2] Aub, J. C., et al. 1952. *Medicine* 31:221. [3] Behrens, C. F., ed. 1959. *Atomic medicine*. Ed. 3. Williams and Wilkins, Baltimore. [4] Bender, M. A., et al. 1962. *Radiation Res.* 16:44. [5] Bender, M. A., et al. 1963. *Ibid.* 18:389. [6] Bennett, L. R., et al. 1953. *Radiology* 61:411. [7] Blair, H. A., ed. 1954. *Natl. Nucl. Energy Ser. VI-2*. [8] Bloom, W., ed. 1948. *Ibid.* IV-22 I. [9] Bond, U. P., et al. 1960. *Radiation Res.* 12:276. [10] Brues, A. M. 1949. *J. Clin. Invest.* 28:1286. [11] Brues, A. M. 1953. *Nucl. Sci. Abstr.* 7:10. [12] Cahan, W. G., et al. 1948. *Cancer* 1:3. [13] Cogan, D. G., D. D. Donaldson, and A. B. Reese. 1952. *Arch. Ophthalmol. (Chicago)* 47:55. [14] Cogan, D. G., J. L. Goff, and E. Graves. 1952. *Ibid.* 47:584. [15] Cogan, D. G., S. F. Martin, and S. J. Kimura. 1949. *Science* 110:654. [16] Conklin, J. W., et al. 1963. *Radiation Res.* 19:156. [17] Cosgrove, G. E., et al. 1962. *Intern. J. Radiation Biol.* 4:97. [18] Cronkite, E. P., et al. 1956. *Ann. Rev. Physiol.* 18:483. [19] Cronkite, E. P., et al. 1960. *Radiation Res.* 12:811. [20] Dublin, L. I., and M. Spiegelman. 1948. *J. Am. Med. Assoc.* 137:1519. [21] Dunjic, A., et al. 1960. *Radiation Res.* 12:155. [22] Esnouf, M. P., et al. 1961. *Intern. J. Radiation Biol.* 3:459. [23] Folley, J. H., W. Borges, and T. Yamawaki. 1952. *Am. J. Med.* 13:311. [24] Franklin, T. J., et al. 1961. *Intern. J. Radiation Biol.* 3:467. [25] Furth, J., and O. B. Furth. 1936. *Am. J. Cancer* 28:54. [26] Harris, R. J. C., ed. 1963. *Cellular basis and aetiology of late somatic effects of ionizing radiations*. Academic Press, New York. [27] Henshaw, P. S., E. F. Riley, and G. E. Stapleton. 1947. *Radiology* 49:348. [28] Hueper, W. C. 1942. *Occupational tumors and allied diseases*. C. C. Thomas, Springfield, Ill. [29] Jacobson, L. O., and E. K. Marks. 1947. *Radiology* 49:286. [30] Jones, A. 1953. *Brit. J. Radiol.* 26:273. [31] Kaplan, H. S., and M. B. Brown. 1951. *J. Natl. Cancer Inst.* 12:427. [32] Kohn, H., et al. 1963. *Radiation Res.* 18:348. [33] Kunkler, P. B., R. F. Farr, and R. W. Luxton. 1952. *Brit. J. Radiol.* 25:190. [34] Lamson, B. G., et al. 1957. *Arch. Pathol.* 64:505. [35] Lamson, B. G., et al. 1958. *Ibid.* 66:311. [36] Lamson, B. G., et al. 1958. *Ibid.* 66:322. [37] Lamson, B. G., et al. 1959. *J. Natl. Cancer Inst.* 22:1059. [38] Lamson, B. G., et al. 1962. *Radiation Res.* 16:54. [39] Lamson, B. G., et al. 1963. *Ibid.* 18:255. [40] Lawrence, J. H., and J. G. Hamilton, ed. 1951. *Advan. Biol. Med. Phys.* 2. [41] Leinfelder, P. J., and H. D. Kerr. 1936. *Am. J. Ophthalmol.* 19:739. [42] Lisco, H., M. P. Finkel, and A. M. Brues. 1947. *Radiology* 49:361. [43] Looney, W. B. 1951. *Nucl. Sci. Abstr.* 5:6046. [44] Lorenz, E., et al. 1947. *Radiology* 49:274. [45] McMahon, B. 1962. *J. Natl. Cancer Inst.* 28:1173. [46] McPherson, S. 1961. *Intern. J. Radiation Biol.* 3:515. [47] March, H. C. 1950. *Am. J. Med. Sci.* 220:282. [48] Martland, H. S. 1931. *Am. J. Cancer* 15:2435. [49] Medical Research Council. 1960. *The hazards to man of nuclear and allied radiations*. H. M. Stationery Office, London. [50] Meuhauser, E. B. D., et al. 1952. *Radiology* 59:637. [51] Mohs, F. E. 1952. *J. Am. Dental Assoc.* 45:160. [52] Mole, T. 1961. *Intern. J. Radiation Biol.* 3:493. [53] Moses, C., J. G. Linn, Jr., and A. J. Allen. 1953. *Arch. Ophthalmol. (Chicago)* 50:609. [54] Murphree, R. L., et al. 1960. *Radiation Res.* 12:495. [55] Neary, G. J., et al. 1962. *Intern. J. Radiation Biol.* 4:239. [56] Ordy, J. M., et al. 1963. *Radiation Res.* 20:30. [57] Peck, W. S., et al. 1940. *Radiology* 34:176. [58] Plummer, G. 1952. *Pediatrics* 10:687. [59] Prezyna, A. P., W. W. Ayres, and W. C. Mulry. 1953.

*continued*

## 127. LATE EFFECTS OF IRRADIATION: MAMMALS

Radiology 60:573. [60] Prosser, C. L., et al. 1947. Ibid. 49:299. [61] Raper, J. R. 1947. Ibid. 49:314. [62] Riley, E. F., et al. 1954. Radiation Res. 1:556. [63] Riley, E. F., et al. 1955. Ibid. 3:342. [64] Rosenthal, M., and E. Grace. 1936. Am. J. Med. Sci. 191:607. [65] Russell, L. B., and W. L. Russell. 1952. Radiology 58:369. [66] Shaver, S. L. 1953. Am. J. Anat. 92:391, 433. [67] Shellabarger, C. J., et al. 1960. Radiation Res. 12:94. [68] Simpson, L., et al. 1962. Ibid. 17:145. [69] Stone, R. S. 1948. Am. J. Roentgenol. Radium Therapy 59:771. [70] Stone, R. S., ed. 1951. Natl. Nucl. Energy Ser. IV-20. [71] Storer, J., and P. Harris. 1953. Nucl. Sci. Abstr. 7:15. [72] Stover, M. 1958. Proc. Soc. Exptl. Biol. Med. 99:201. [73] Thomson, J. F., et al. 1953. Am. J. Roentgenol. Radium Therapy Nucl. Med. 69:830. [74] Ulrich, H. 1946. New Engl. J. Med. 234:45. [75] Warren, S. 1942. Arch. Pathol. 34:443, 562, 749, 917, 1070. [76] Warren, S. 1943. Ibid. 35:121, 304. [77] Wilson, J. G., R. L. Brent, and H. C. Jordan. 1953. Proc. Soc. Exptl. Biol. Med. 82:67. [78] Zirkle, R. E., ed. 1951. Natl. Nucl. Energy Ser. IV-22 E.

## XII. PARASITISM

### 128. ARTHROPOD PARASITES: MAMMALS AND BIRDS

Many of the arthropods listed are known to be parasites of man. Some of these are identified by an asterisk (\*).

Species (Common Name); Distribution	Free Stage Location	Host [Location and Stage in Host]	Effect on Host	Ref- er- ence
(A)	(B)	(C)	(D)	(E)
Pentastomida				
1 <i>Linguatula serrata</i> (tongue worm); worldwide	Eggs swallowed, hatched in alimentary tract of herbivores; larvae, nymphs develop in mesenteries	Mammals, birds, [Eggs expelled in respiratory tract; adults in nasal passages]	Severe irritation and blockage of nasal passages.	11
Arachnida				
2 <i>Amblyomma americanum</i> * (lone-star tick); N., Cen., & S. America	Eggs in soil; unfed larvae, nymphs, adults on grass	Cattle, dog, horse, goat, sheep, occasionally birds. [External--on host only while feeding]	Damage to hide, milk reduction. Vector of organisms causing Rocky Mt. spotted fever, Q fever. Larvae, nymphs, adults are bloodsuckers.	1,13
3 <i>Argas persicus</i> (fowl tick); warm & temperate semi-arid regions of world	All stages in crevices, cracks of housing, under bark of trees	Domestic fowl, occasionally wild birds. [External--on host only while feeding]	Anemia, leg weakness, egg reduction, occasionally death. Vector of organisms causing fowl spirochetosis, spiroplasmosis. Nymphs, adults are bloodsuckers.	1,3,13
4 <i>Bdellonyssus sylviarum</i> (northern fowl mite); United States, Canada, Mexico, Europe, S. Africa	Eggs on feathers, in nests; other stages on surroundings	Poultry, wild birds. [External; on body and feathers]	Skin lesions, egg reduction, retarded growth, anemia. Harbinger of neurotropic viruses. Larvae, nymphs, adults are bloodsuckers.	1,3
5 <i>Boophilus annulatus</i> * (cattle tick); N. America	Eggs on soil; unfed larvae on grass	Principally ungulates. [External]	Damage to hide, milk reduction. Vector of organisms causing Texas cattle fever. Larvae, nymphs, adults are bloodsuckers.	1,2, 11, 13
6 <i>Demodex canis</i> (dog follicle mite); worldwide	None	Dog. [Eggs, nymphs, adults in hair follicles and sebaceous glands]	Follicle inflammation, mange, thickened skin, alopecia, emaciation; sometimes death.	1,11
7 <i>Dermacentor andersoni</i> * (Rocky Mountain wood tick); Western N. America	Eggs on soil	Most mammals. [External; larvae, nymphs on most small animals; adults usually on larger animals during feeding period]	Paralysis, particularly in sheep. Vector of organisms causing Rocky Mt. spotted fever, tularemia, equine encephalomyelitis, anaplasmosis. Larvae, nymphs, adults are bloodsuckers.	11,13
8 <i>D. variabilis</i> (American dog tick); N. America	Eggs on soil; unfed larvae, nymphs, adults on vegetation until host is found	Principally dog; other domestic and wild animals. [External--on host only while feeding; larvae, nymphs mainly attack rodents and other small animals]	Skin damage. Vector of organisms causing bovine anaplasmosis, tularemia, Rocky Mt. spotted fever. Larvae, nymphs, adults are bloodsuckers.	1,13
9 <i>Dermanyssus gallinae</i> (chicken mite); worldwide	Eggs, non-feeding larvae, nymphs, adults in crevices of coops, roosts	Poultry, other birds. [External]	Decreased egg production, retarded growth, anemia; sometimes death. Vector of organisms causing spirochetosis, fowl cholera. Larvae, nymphs, adults are bloodsuckers; nocturnal feeders only.	1,3
10 <i>Eutrombicula alfreddugesi</i> (chigger); N. & S. America, W. Indies	Active forms in grasses, shrubs, brambles	Domestic and wild vertebrates. [External]	Irritation due to toxins; sometimes death (small poultry). Larvae are bloodsuckers.	1,11, 13

continued



## 128. ARTHROPOD PARASITES: MAMMALS AND BIRDS

Species (Common Name); Distribution	Free Stage Location	Host [Location and Stage in Host]	Effect on Host	Ref- er- ence
(A)	(B)	(C)	(D)	(E)
Arachnida				
11 <i>Ixodes ricinus scapularis</i> (black-legged tick); principally Europe	Eggs on soil; larvae, nymphs, adults on grass and shrubbery until host is found	Principally dog; other domestic and wild animals. [External--on host only while feeding; adults on head, neck of dog; on flank, leg, under tail of other animals]	Anemia. Vector of cattle red-water fever, louping ill virus, tick-borne fever virus of sheep. Larvae, nymphs are bloodsuckers in ear, eyelid, head, rarely body.	1,11
12 <i>Knemidokoptes mutans</i> (scaly-leg mite); worldwide	None	Chicken, turkey, other domestic birds. [External; all active stages in tunnels between scales, feet, legs, neck, comb]	Inflammation, keratinization between scales of feet, legs; lameness.	1,3,11
13 <i>Otobius megnini</i> * (ear tick); United States, Mexico, S. America, S. Africa	Eggs on ground, in cracks; adults, unattached larvae in out-buildings	Domestic animals. [Inside ears]	Ear inflammation, anorexia, dullness; sometimes death. Larvae, nymphs are bloodsuckers.	1,11, 13
14 <i>Otodectes cyanotis</i> (ear mite); worldwide	None	Dog, cat, ferret. [All stages in ears; sometimes external]	Inflammation, ear scabs, head-shaking, scratching, droopy ears with discharge; epileptiform fits (severe cases).	1,11
15 <i>Psoroptes equi ovis</i> (sheep-scab mite); worldwide	None	Sheep. [External; all active stages on skin around edge of lesions]	Scabbing; wool loss from biting, scratching; emaciation; sometimes death.	1,11, 13
16 <i>Rhipicephalus sanguineus</i> * (brown dog tick); worldwide	Active forms near habitat of dog	Principally dog. [External]	Vector of organisms causing canine piroplasmosis, cattle gall sickness. Larvae, nymphs, adults are bloodsuckers.	1,11
17 <i>Sarcoptes scabiei</i> * (itch mite); worldwide	None	Most mammals, sheep (on head). [External; all active stages in skin tunnels]	Scratching, papules, vesicles, keratinization, alopecia, mange, emaciation; sometimes death.	1,11, 13
Insecta				
18 <i>Aedes dorsalis</i> * (mosquito); N. America, Europe, northern Africa	Immature forms in moist soil; eggs survive long periods of drying in soil	Warm-blooded animals. [External]	Adult females are bloodsuckers. Vectors of organisms causing equine encephalomyelitis.	9,11
19 <i>Anopheles punctipennis</i> * (mosquito); N. America	Immature forms in streams, ponds of hilly country	Warm-blooded animals. [External; where hair or feathers are thinnest]	Adult females are bloodsuckers. Vectors of organisms causing dog heartworm.	9
20 <i>Bovicola bovis</i> (cattle-biting louse); N. & S. America, Europe, Africa, Australia	None	Cattle. [External; eggs on hair; nymphs, adults feed on skin]	Reduced vigor, irritation, scaly skin.	11,13
21 <i>Cimex lectularius</i> * (bedbug); worldwide	All stages in cracks, crevices, similar hiding places	Domestic animals, poultry. [External]	Skin irritation, welts. Nymphs, adults are bloodsuckers.	11,13
22 <i>Chrysops discalis</i> (deerfly); Western N. America	Eggs near water, larvae in water, pupae in mud	Principally horse, cattle. [External; mainly on underside of abdomen, neck, withers]	Vector of tularemia, surra. Adults are bloodsuckers.	5,11
23 <i>Cochliomyia hominivorax</i> * (screw-worm); tropical & subtropical areas of western hemisphere	Pupae in soil, adults in pastures	Obligatory parasite of warm-blooded animals, including livestock, wild mammals, dog, cat. [Eggs deposited on edges of wounds]	Infection and extension of wounds; untreated host invariably dies.	4,8,10

1/1 Other varieties infest various domestic animals. 1/2 Adult stage of *C. macellaria* (secondary screwworm) resembles *C. hominivorax* in appearance, but differs by being a secondary invader (facultative parasite), and by breeding in carcasses. Larvae occasionally infest wool or necrotic wounds.

continued

## 128. ARTHROPOD PARASITES: MAMMALS AND BIRDS

Species (Common Name); Distribution	Free Stage Location	Host [Location and Stage in Host]	Effect on Host	Ref- er- ence
(A)	(B)	(C)	(D)	(E)
Insecta				
24	<i>Ctenocephalides canis</i> (dog flea); worldwide	Cat, dog, swine, other animals. [External]	Coat damaged from biting, scratching. <i>C. canis</i> and <i>C. felis</i> are vectors of dog and dwarf tapeworms, heartworm, plague, epidemic typhus. Adults are bloodsuckers.	6,11, 17
25				
26	<i>Cuclotogaster heterographus</i> (chicken head louse); worldwide	Chicken, partridge. [External; nymphs, adults on skin and feathers of head, neck; eggs on feathers]	Irritation.	3,11
27	<i>Culex quinquefasciatus</i> * (southern house mosquito); worldwide from 60°N to 50°S latitude	Warm-blooded animals, especially birds. [External; where hair or feathers are thinnest]	Adult females are bloodsuckers. Vectors of organisms causing avian malaria, fowl pox.	3,9,11
28	<i>Dermatobia hominis</i> * (human botfly); S. America, W. Indies, tropical N. America	Dog, swine, mule, cattle, wild animals. [Larvae leave transport arthropod on contact with host; penetrate skin]	Boil-like skin lesions, reduced milk production, damage to hide, decreased growth rate.	8,10, 14
29	<i>Echidnophaga gallinacea</i> * (sticktight flea); worldwide, especially warm climates	Poultry, domestic animals, rodents. [External; skin, comb, wattles; around eyes and ears]	Anemia; sometimes death. Adults are bloodsuckers.	3,13, 16
30	<i>Gasterophilus intestinalis</i> <sup>3</sup> (horse botfly); worldwide	Ass, horse, mule; rarely other animals. [Eggs on foreleg fetlock hairs; larvae (maggots) in mouth, pharynx, stomach]	Extension and infection of wounds.	8,10, 14
31	<i>Glossina morsitans</i> * (tsetse fly); central Africa	Cattle, other animals. [External]	Vector of organisms causing cattle and horse nagana, sleeping sickness to man. Adults are bloodsuckers.	11,17
32	<i>Haematopinus eurysternus</i> <sup>4</sup> (short-nosed cattle louse); worldwide	Cattle. [External; eggs on shaft or at base of hairs]	Hair damage from rubbing; stunting; reduced milk production. Nymphs, adults are bloodsuckers.	11,13
33	<i>H. suis</i> (hog louse); worldwide	Swine. [External; eggs on hair]	Dermatitis, skin sores, retarded growth. Vector of organism causing swine pox. Nymphs, adults are bloodsuckers.	13
34	<i>Hypoderma lineatum</i> <sup>5</sup> (common cattle grub); prevalent in America, Europe, India, northern Asia	Cattle, rarely horse. [Eggs on hair of legs, body; larger larvae form tumor under skin of back]	Skin perforation, hide and flesh damage, milk reduction.	4,11, 13
35	<i>Melophagus ovinus</i> (sheep ked); most parts of world	Sheep, occasionally goat. [External; pupae attached to wool]	Anemia; wool stained, damaged from rubbing. Adults are bloodsuckers.	11,13, 17
36	<i>Menacanthus stramineus</i> (chicken body louse); worldwide	All domestic fowl. [External; nymphs, adults on skin around vent; eggs attached to feathers]	Scabbing of skin; wasting; reduced egg production.	3,11
37	<i>Menopon gallinae</i> (shaft louse); worldwide	Chicken, guinea fowl. [External; eggs, nymphs, adults feed on scales, scabs, feathers]	Scaling, scabbing.	3,11

[2] *G. haemorrhoidalis* (nose botfly) and *G. nasalis* (throat botfly) are similar in many respects to *G. intestinalis*.

[4] Information also applicable to *Linognathus vituli* (long-nosed cattle louse). [5] Information also applicable to *H. bovis* (northern cattle grub).

continued

## 128. ARTHROPOD PARASITES: MAMMALS AND BIRDS

Species (Common Name); Distribution	Free Stage Location	Host [Location and Stage in Host]	Effect on Host	Refer- ence
(A)	(B)	(C)	(D)	(E)
Insecta				
38 <i>Musca domestica</i> (housefly); world- wide	Immature forms in manure and decayed matter, adults in buildings	Any larger animal with lesions or secretions. [External; adults acci- dentally ingested by host]	Adults cause decreased produc- tivity of livestock. Vector of several tapeworm species; me- chanical vector of many bacterial and protozoan pathogens and hel- minth eggs.	3,11, 13
39 <i>Oestrus ovis</i> (sheep botfly); worldwide	Pupae on ground, adults in warm corners, crevices	Sheep, rarely goat. [Lar- vae in nasal cavities, sinuses]	Mucosal irritation, nasal dis- charge, emaciation; sometimes death.	8,11, 13
40 <i>Phaenicia sericata</i> (greenbottle fly); worldwide, except S. America and Pacific Islands	Adults free-living, de- posit eggs on flesh, soiled wool; pupae in soil	Sheep, goat, other ani- mals. [Larvae on skin, in wounds]	Invasion of wounds, suppuration.	7,11
41 <i>Phlebotomus papa- tasii</i> * (sand fly); Mediterranean region, southern Europe, Asia	Immature forms in dark moist places, manure	Warm-blooded animals. [External]	Swelling at site of bite. Vector of organisms causing pappataci fever. Adults are nocturnal bloodsuckers.	11,13, 16
42 <i>Phormia regina</i> <sup>e</sup> (black blowfly); worldwide	Pupae in soil, adults in pastures	Sheep, other mammals. [Eggs in hair or wool, larvae in wounds; eggs and larvae also in carcasses]	Extension and infection of wounds.	7,13
43 <i>Pulex irritans</i> * (hu- man flea); world- wide	Eggs, larvae, and pu- pae in soil, adults on ground part of time	Man, dog, swine, other animals. [External]	Irritation, poor condition, coat damaged from biting and scratch- ing. Adults are bloodsuckers.	11,17
44 <i>Simulium</i> spp. <sup>?</sup> (blackfly); world- wide in temperate to subarctic cli- mates	Immature forms on under sides of stones in moderate-running streams	All warm-blooded ani- mals. [External; on bare parts of head, body, legs; under wings]	Red swelling, vesicles, anemia, toxemia; death. <i>S. occidentale</i> and <i>S. slossonae</i> are vectors of turkey leucocytozoan disease; some species are vectors of on- chocerciasis to man and cattle. Adults are bloodsuckers.	3,11, 13, 16
45 <i>Siphona irritans</i> (horn fly); Amer- ica, Europe	Eggs, maggots in fresh dung, pupae in dung or soil	Cattle, other animals. [External]	Weight loss; milk reduction. Adults are bloodsuckers.	11,13, 17
46 <i>Stomoxys calci- trans</i> * (stable fly); worldwide	Immature forms in manure and other moist organic waste	Most mammals and birds. [External]	Weight loss; milk reduction. Vec- tor of poultry tapeworms, fila- riae, spiruroids; mechanical vector of surra. Adults are bloodsuckers.	3,7,11
47 <i>Tabanus atratus</i> (black horsefly); N. America	Immature forms in leaves and mud in and near streams, ponds	Most warm-blooded ani- mals. [External]	Vector of organism causing ana- plasmiasis. Adults are blood- suckers during day.	13,15
48 <i>Triatoma sangui- suga</i> <sup>3</sup> (cone-nose bug); worldwide	All stages commonly found in or close to rodent nests or habi- tats	All domestic animals, poultry; wood rat usual host. [External]	Swelling, anemia. Nymphs, adults are bloodsuckers.	12,13, 17
49 <i>Trichodectes canis</i> (dog-biting louse); worldwide	None	Dog. [External; eggs on hair; nymphs and adults feed on skin]	Scaly skin from rubbing, scratch- ing.	11
50 <i>Wohlfahrtia vigil</i> * (flesh fly); N. America	Pupae on ground, lar- vae in woods	Rabbit, mink, guinea pig, young of domestic and wild animals. [Larvae in wounds]	Mild to extensive subcutaneous pustular lesions; occasionally death.	7,11

[<sup>3</sup>] *Chrysomya chloropyga* is similar to *P. regina* in its parasitism of sheep. [<sup>7</sup>] *S. articum*, *S. occidentale*, *S. ornatum*, *S. vittatum* are the important blackfly pests of livestock. [<sup>1</sup>] Sixteen species of *Triatoma* found in the western hemisphere are as important and as widely distributed as *T. sanguisuga*.

Contributors: (a) Edgar, S. A., (b) Furman, Deane P.

continued



## 128. ARTHROPOD PARASITES: MAMMALS AND BIRDS

*References:* [1] Baker, E. W., and G. W. Wharton. 1952. An introduction to acarology. Macmillan, New York. [2] Belding, D. L. 1952. Textbook of clinical parasitology. Ed. 2. Appleton-Century-Crofts, New York. [3] Biester, H. E., and L. H. Schwarte. 1959. Diseases of poultry. Ed. 4. Iowa State College Press, Ames. [4] Bishopp, F. C., et al. 1926. U.S. Dept. Agr. Farmers' Bull. 857. [5] Dickmans, G. 1945. Am. J. Vet. Res. 6:211. [6] Ewing, H. E. 1929. A manual of external parasites. C. C. Thomas, Springfield, Ill. [7] Hall, D. G. 1948. The blowflies of North America. Thomas Say Foundation, Baltimore. [8] Herms, W. B. 1961. Medical entomology. Ed. 5. Macmillan, New York. [9] Horsfall, W. R. 1955. Mosquitoes, their bionomics and relation to disease. Ronald Press, New York. [10] James, M. T. 1947. U.S. Dept. Agr. Misc. Publ. 631. [11] Lapage, G., ed. 1962. Mönnig's Veterinary helminthology and entomology. Ed. 5. Williams and Wilkins, Baltimore. [12] Matheson, R. 1950. Medical entomology. Ed. 2. Comstock, Ithaca. [13] Metcalf, C. L., and W. P. Flint. 1962. Destructive and useful insects: their habits and control. Ed. 4. McGraw-Hill, New York. [14] Neel, W. W. 1954. J. Econ. Entomol. 47(3):540. [15] Oldroyd, H. 1954. The horse-flies of the Ethiopian region. British Museum, London. v. 2. [16] Patton, W. S., and K. M. Evans. 1929. Insects, ticks, mites and venomous animals. H. R. Grubb, Croydon, England. pt. 1. [17] Smart, J. 1956. A handbook for the identification of insects of medical importance. Ed. 3. British Museum, London.

## 129. ARTHROPOD PESTS: PLANTS AND PLANT PRODUCTS

Species	Common Name	Stage <sup>1</sup>	General Distribution	Host	Destructive Activity
(A)	(B)	(C)	(D)	(E)	(F)
Arachnida					
1 <i>Eriophyes pyri</i>	Pear leaf blister mite	Adult; immature	All pear-growing regions	Pear, apple	Sucking causes blisters on underside of leaves
2 <i>Rhizoglyphus echinopus</i>	Bulb mite	Adult; immature	N. America, Europe, Asia	Ornamental bulbs, onion	Bores into bulbs
3 <i>Steneotarsonemus pallidus</i>	Cyclamen mite	Adult; immature	N. America, Europe	Greenhouse ornamentals, strawberry	Sucks plant juices, distorts buds and leaves
4 <i>Tetranychus telarius</i>	Two-spotted spider mite	Adult; immature	United States, Europe, Africa, Asia, Australia	Cultivated plants	Sucks plant juices, causing loss of vigor, dropping of leaves
Crustacea					
5 <i>Porcellio laevis</i>	Sowbug	Adult; immature	Worldwide	Vegetables, ornamentals	Chews roots, growths near ground
Insecta					
6 <i>Acanthoscelides obtectus</i>	Bean weevil	Larva	Worldwide	Bean, pea, cowpea	Devours inside of bean in field and in storage
7 <i>Agriotes, Horistonotus, Limonius, Melanotus</i> spp.	Wireworms	Larva	Worldwide	Truck, cereal, and forage crops	Devours or bores into roots, seeds
8 <i>Alabama argillacea</i>	Cotton leafworm	Larva	N. & S. America	Cotton only	Devours leaves
9 <i>Altica, Phyllotreta</i> spp.	Flea beetles	Adult; larva <sup>2</sup>	Worldwide	Vegetable crops	Adult makes holes in leaves, larva often feeds on roots
10 <i>Amphibolips confluenta</i>	Oak gall wasp	Larva	Worldwide	Oak	Causes galls on oak leaves
11 <i>Anabrus simplex</i>	Mormon cricket	Adult; nymph	Western United States	Hay, grain, many plants	Devours hay, grain, leaves of plants
12 <i>Anasa tristis</i>	Squash bug	Adult; nymph	N. & Cen. America	Squash, other cucurbits	Sap sucking causes plants to wilt and die

1/ Destructive stage of arthropod. 2/ Overwinters as adult.

*continued*



## 129. ARTHROPOD PESTS: PLANTS AND PLANT PRODUCTS

Species	Common Name	Stage <sup>1</sup>	General Distribution	Host	Destructive Activity	
(A)	(B)	(C)	(D)	(E)	(F)	
Insecta						
13	<i>Anthonomus grandis</i>	Boll weevil	Adult; larva	Southern United States, Mexico	Cotton	Destroys buds, devours squares and bolls
14	<i>Aphis pomi</i>	Apple aphid	Adult; nymph	N. America	Apple	Causes wilting
15	<i>Aspidiotus perniciosus</i>	San Jose scale	Adult; nymph	Worldwide	Deciduous fruit trees, ornamentals	Secreted toxins cause wilting, kill infested trees
16	<i>Blissus leucopterus</i>	Chinch bug	Adult; nymph	N. America	Corn, grains, grasses	Sap sucking causes wilting, death
17	<i>Carpocapsa pomonella</i>	Codling moth	Larva	Apple-growing regions of N. & S. America, Europe, Asia, S. Africa, southern Australia	Apple, pear, quince, walnut, apricot, similar fruits	Bores into and destroys fruit, or reduces its value
18	<i>Cephus pygmaeus</i>	European wheat stem sawfly	Larva	Northeastern United States, Europe, Near East	Wheat, rye, barley, timothy, other grasses	Mines stems, causing breakage
19	<i>Ceratitis capitata</i>	Mediterranean fruit fly	Adult; larva	Mediterranean region, Hawaii, S. Africa, S. America, western Australia	Fruits, vegetables	Adult makes egg punctures, larva burrows through fruit
20	<i>Choristoneura fumiferana</i>	Spruce budworm	Larva	Northern United States, Canada	Fir, spruce, hemlock, larch, white pine	Causes partial-to-complete defoliation
21	<i>Chrysobothris femorata</i>	Flatheaded apple tree borer	Larva	United States, Canadian fruit-growing areas	Fruit trees, many shade trees	Bores into trunk of weakened trees, branches, twigs; kills tree
22	<i>Cladius isomerus</i>	Rose slug	Larva	Worldwide	Rosebush	Skeletonizes and causes browning of leaves
23	<i>Coccus hesperidum</i>	Brown scale	Adult; nymph	Worldwide in greenhouses (subtropical spp.)	Citrus, ornamentals	Sap sucking causes plants to die back <sup>3</sup>
24	<i>Conotrachelus nemphar</i>	Plum curculio	Adult; larva	Eastern United States, Canada	Plum, apple, peach, cherry, deciduous stone fruits	Adult punctures fruit, larva feeds within and destroys fruit
25	<i>Corythuca arcuata</i>	Oak lace bug		Worldwide	Various trees, shrubs	Speckling of leaves, stunting
26	<i>Dendroctonus monticolae</i>	Mountain pine beetle	Adult; larva	Western United States	Western, lodgepole, sugar, ponderosa, white bark, and limber pines	Bores into bark and cambial region; may girdle and kill tree
27	<i>Diabrotica undecimpunctata</i>	Spotted cucumber beetle	Adult; larva	N. America	Corn, cucurbits, weeds, grasses, other plants	Larva feeds on roots, adult devours foliage <sup>1</sup>
28	<i>Diprion hercyniae</i>	European spruce sawfly	Larva	Europe, northeastern United States, Canada	Spruce	Devours leaves
29	<i>Drosophila melanogaster</i>	Fruit fly	Larva	Worldwide	Ripe or decaying fruit	Breeds in ripe fruit
30	<i>Empoasca fabae</i>	Potato leafhopper	Adult; nymph	N. & S. America	Potato, alfalfa, bean, celery, other plants	Leaf sucking <sup>2</sup> causes wilting, drying of leaves, stunting
31	<i>Ephestia kuehniella</i>	Mediterranean flour moth	Larva	Worldwide	Mill products	Destroys grain products
32	<i>Epicauta vittata</i>	Blister beetle	Adult	Worldwide	Potato, legumes	Devours plants

<sup>1</sup>/ Destructive stage of arthropod. <sup>2</sup>/ Honeydew is formed or excreted. <sup>3</sup>/ Also vector of bacterial wilt of cucurbits, and of viral yellow disease of asters. <sup>4</sup>/ Also transmits hopperburn disease.

continued

## 129. ARTHROPOD PESTS: PLANTS AND PLANT PRODUCTS

Species	Common Name	Stage <sup>1</sup>	General Distribution	Host	Destructive Activity	
(A)	(B)	(C)	(D)	(E)	(F)	
Insecta						
33	<i>Epilachna varivestis</i>	Mexican bean beetle	Adult; larva	United States, Mexico	Bean, soybean, cowpea, other legumes	Devours leaves, pods, stems
34	<i>Epitrix hirtipennis</i>	Tobacco flea beetle	Adult	Worldwide	Tobacco	Devours leaves, especially those of young plants
35	<i>Eriosoma lanigerum</i>	Woolly apple aphid	Adult; nymph	N. & S. America, Europe, S. Africa, Asia, Australia	Apple, pear, elm	Branch and root sucking causes deformed twigs, knotty roots, stunting
36	<i>Forficula auricularia</i>	European earwig		Worldwide	Growing plants, stored grain, decayed vegetation	Chewing
37	<i>Gryllus</i> spp.	Field cricket	Adult; nymph	N., Cen., & S. America	Hay crops, cotton, linen	Devours hay, plants, cotton, linen
38	<i>Harmolita tritici</i>	Wheat joint-worm	Larva	Eastern & central United States	Wheat, some grasses	Causes gall in wheat, breaking of stems
39	<i>Heliothis zea</i>	Corn earworm	Larva	Worldwide	Cotton, corn, tomato, alfalfa, other plants	Bores into and feeds on bolls, ears, buds; stunts plants, reduces yield
40	<i>Hylemya antiqua</i>	Onion maggot	Larva	Europe, N. America	Onion, garlic	Mines out bulbs <sup>3</sup>
41	<i>Hyphantria cunea</i>	Fall webworm	Larva	United States, southern Canada	Broadleaf fruit, shade and nut trees	Webs branches, devours foliage
42	<i>Lampetia equestris</i>	Narcissus bulb fly	Larva	Europe, N. America	Narcissus, other bulbs	Bores into bulbs
43	<i>Lasius alienus americanus</i>	Cornfield ant	Adult	United States	Corn	Symbiotic, with aphids attacking corn roots
44	<i>Lepisma saccharina</i>	Silverfish	Adult; nymph	Worldwide	Starchy substances	Devours bookbindings, fabrics, wallpaper
45	<i>Leptinotarsa decemlineata</i>	Colorado potato beetle	Adult; larva	N. America, Europe	Potato, tomato, tobacco, eggplant, other solanaceous plants	Devours leaves, terminates growth
46	<i>Liposcelis divinatorius</i>	Book louse	Adult; immature	Worldwide	Cereals, vegetables	Contaminates food, destroys bookbindings
47	<i>Lygus lineolaris</i>	Tarnished plant bug	Adult; nymph	N. America	Many plants, trees	Leaf sucking and toxins cause bud drop, distortion, stunting
48	<i>Magicicada septendecim</i>	Periodical cicada	Adult	Eastern & southern United States	Many deciduous trees, shrubs	Oviposition punctures injure or kill twigs
49	<i>Malacosoma disstria</i>	Forest tent caterpillar	Larva	N. America	Aspen, sugar maple, oak, birch, basswood, ash, gum, other trees	Defoliates trees in summer
50	<i>Megachile latimanus</i>	Leaf-cutting bee	Adult	Worldwide	Various trees	Cuts off leaf fragments for nests
51	<i>Melanoplus femurrubrum</i>	Red-legged grasshopper	Adult; nymph	Worldwide	Hay crops (range and pasture)	Devours hay, grasses, vegetation
52	<i>Microcentrum rhombifolium</i>	Broad-winged katydid	Adult; nymph	N. America	Many broad-leaved trees, shrubs	Chews leaves
53	<i>Murgantia histrionica</i>	Harlequin cabbage bug	Adult; nymph	Southern United States, Mexico, Cen. America	Cabbage, related crops, other plants	Sap sucking causes plants to wilt, brown, and die
54	<i>Myzus persicae</i>	Green peach aphid	Adult; nymph	Warm regions of world	Many trees, shrubs	Leaf sucking causes curling, distortion of leaves <sup>3</sup>

<sup>1</sup>/ Destructive stage of arthropod. <sup>3</sup>/ Honeydew is formed or excreted.

continued

## 129. ARTHROPOD PESTS: PLANTS AND PLANT PRODUCTS

Species	Common Name	Stage <sup>1</sup>	General Distribution	Host	Destructive Activity	
(A)	(B)	(C)	(D)	(E)	(F)	
Insecta						
55	<i>Oryzaephilus surinamensis</i>	Saw-toothed grain beetle	Adult; larva	Worldwide	Grain, grain products, dried fruit	Infests and devours grain, grain products, dried fruit
56	<i>Oscinella frit</i>	Frit fly	Larva	N. America, Europe, Asia	Cereals, grasses	Bores into stems, eats central shoots
57	<i>Paleacrita vernata</i>	Spring cankerworm	Larva	Eastern United States, Canada	Fruit and shade trees	Defoliates trees in spring
58	<i>Pectinophora gossypiella</i>	Pink bollworm	Larva	Southern United States, S. America, Africa, Europe, Asia, Australia	Cotton, okra, other malvaceous plants	Bores into and feeds on squares and bolls, cutting fiber, reducing yield
59	<i>Peridroma saucia</i>	Variegated cutworm	Larva	Worldwide	Many plants	Cuts down seedlings, transplants
60	<i>Philaenus leucophthalmus</i>	Meadow spittlebug	Nymph	Eastern United States	Legumes, hay crops	Sap sucking causes wilting, stunting, reduced forage yield
61	<i>Phyllophaga</i> spp. <sup>6</sup>	June beetle	Adult; larva	N. America	Most plants	Adult defoliates trees; larva devours roots, underground parts
62	<i>Phylloxera vitifoliae</i>	Grape phylloxera	Adult; nymph	N. America, Europe	Grape vines	Root and leaf sucking causes galls, eventual death of vines
63	<i>Phytophaga destructor</i>	Hessian fly	Larva	Europe, Asia, N. America, New Zealand	Wheat, barley, rye	Feeds on stems, causing breaking and stunting
64	<i>Pieris rapae</i>	Imported cabbageworm	Larva	N. America, Asia, Australia, Europe	Cabbage, other crucifers	Devours foliage
65	<i>Plodia interpunctella</i>	Indian meal moth	Larva	Worldwide	Grain, grain products, dried fruit, nuts	Destroys and webs grain, grain products; infests fruit, nuts
66	<i>Plutella maculipennis</i>	Diamondback moth	Larva	Worldwide	All cruciferous plants	Eats small holes in outer leaves
67	<i>Popillia japonica</i>	Japanese beetle	Adult; larva	Eastern United States, Japan, China	Fruit trees, ornamentals, vegetables, grasses	Destroys turf, foliage, blossoms, fruit
68	<i>Porthetria dispar</i>	Gypsy moth	Larva	Northeastern United States, Europe, Asia	Most deciduous and evergreen trees, shrubs	Devours leaves
69	<i>Protoparce quinque-maculata</i>	Tomato hornworm	Larva	N. & S. America, Europe, Hawaii	Tomato, tobacco, other solanaceous plants	Devours foliage
70	<i>Pseudaletia unipuncta</i>	Armyworm	Larva	Worldwide	Grains, grasses, some legumes	Devours foliage
71	<i>Pseudococcus citri</i>	Mealybug	Adult; nymph	Tropical & subtropical areas	Citrus, ornamental plants	Sap sucking causes plants to die back <sup>3</sup>
72	<i>Psila rosae</i>	Carrot rust fly	Larva	Europe, northern N. America	Carrot, celery, umbelliferous plants	Bores into and eats fibrous roots
73	<i>Psyllia pyricola</i>	Pear psylla	Adult; nymph	Europe, United States	Pear	Leaf sucking causes leaf drop <sup>3</sup>
74	<i>Pyrausta nubilalis</i>	European corn borer	Larva	Eastern United States, Europe, Asia	Corn is main host; also many vegetables, weeds, ornamentals	Bores into stalks and ears, causing breakage, reduced yield and quality
75	<i>Ramosia tipuliformis</i>	Currant borer	Larva	N. America, Asia, Europe, Australia	Currant, gooseberry, black elder, sumac	Burrows through canes
76	<i>Reticulitermes flavipes</i>	Eastern subterranean termite	Adult; nymph	Eastern United States	Wood, dead wood, cellulose products	Riddles, weakens, destroys wood and cellulose materials

<sup>1/1</sup> Destructive stage of arthropod. <sup>3/3</sup> Honeydew is formed or excreted. <sup>6/6</sup> Other important June beetles belong to *Melolontha* and *Polyphylla* spp.

continued



**129. ARTHROPOD PESTS: PLANTS AND PLANT PRODUCTS**

Species	Common Name	Stage <sup>1</sup>	General Distribution	Host	Destructive Activity	
(A)	(B)	(C)	(D)	(E)	(F)	
Insecta						
77	<i>Rhagoletis pomonella</i>	Apple maggot	Larva	Northeastern & north central United States	Apple, blueberry	Bores into and destroys fruit
78	<i>Saminoidea exitiosa</i>	Peach tree borer	Larva	All peach-growing areas	Peach, other stone-fruit trees	Bores into trunk at ground-level roots, girdles tree trunk, kills tree
79	<i>Saperda candida</i>	Round-headed apple tree borer	Larva	Eastern United States, Canada	Apple, pear, quince trees	Bores into trunk
80	<i>Schistocerca gregaria</i> <sup>2</sup>	Desert locust	Adult; nymph	India, Iran, Arabia, N. Africa	Many plants	Chews leaves
81	<i>Sitophilus oryza</i>	Rice weevil	Adult; larva	Worldwide	Stored grains, cereal products	Larva grows in kernels, destroys stored grain
82	<i>Sminthurus viridis</i>	Lucerne flea	Adult; immature	Europe, Australia	Legumes	Surface feeding causes scorching of leaves
83	<i>Spissistilus festinus</i>	Three-cornered alfalfa bug	Adult	Worldwide	Alfalfa	Stunting
84	<i>Tenebrio molitor</i>	Yellow mealworm	Larva	Worldwide	Grain products, refuse	Destroys grain, grain products
85	<i>Tenebroides mauritanicus</i>	Cadelle	Adult; larva	Worldwide	Stored grain, grain products	Infests and destroys grain, grain products
86	<i>Thermobia domestica</i>	Firebrat	Adult; nymph	Worldwide	Starchy substances	Devours bookbindings, fabrics, wall-paper
87	<i>Thrips tabaci</i>	Onion thrips	Adult; larva; nymph	N. & S. America, Europe, Asia, S. Africa, Australia	Onion, bean, cabbage, tomato, cotton	Sap sucking causes leaves and buds to pucker and silver
88	<i>Thyridopteryx ephemeraeformis</i>	Bagworm	Larva	Eastern United States	Cedar, other trees	Devours foliage
89	<i>Trialeurodes vaporariorum</i>	Greenhouse whitefly	Nymph	Worldwide	Most plants	Leaf sucking causes wilting <sup>3</sup>
90	<i>Tribolium confusum</i>	Confused flour beetle	Adult; larva	Worldwide	Flour, grain products	Infests and contaminates flour, mixes, bread
Symphyla						
91	<i>Scutigrella immaculata</i>	Garden symphylid	Adult; immature	N. & S. America, Europe, Africa	Garden plants, flowers	Chews tender plants, rootlets
Diplopoda						
92	<i>Julus heseris</i>	Millipede	Adult; immature	Worldwide	Vegetables, ornamentals	Chews young roots, stems

<sup>1</sup>/ Destructive stage of arthropod. <sup>2</sup>/ Honeydew is formed or excreted. <sup>3</sup>/ Has a migratory phase.

Contributors: (a) Allen, William W., (b) Stanley, W. W., and Dozier, Byrd K.

References: [1] Craighead, F. C. 1949. U.S. Dept. Agr. Misc. Publ. 657. [2] Essig, E. O. 1958. Insects and mites of western North America. Macmillan, New York. [3] Imms, A. D., O. W. Richards, and R. G. Davies. 1960. A general textbook of entomology. Ed. 9. E. P. Dutton, New York. [4] Little, W. A. 1963. General and applied entomology. Ed. 2. Harper and Row, New York. [5] Metcalf, C. L., and W. P. Flint. 1962. Destructive and useful insects, their habits and control. Ed. 4. McGraw-Hill, New York. [6] Peairs, L. M., and R. H. Davidson. 1956. Insect pests of farm, garden, and orchard. Ed. 5. Chapman and Hall, New York. [7] Pfadt, R. E. 1962. Fundamentals of applied entomology. Macmillan, New York. [8] U.S. Department of Agriculture. 1952. Insects. Yearbook of agriculture. U.S. Govt. Printing Office, Washington, D. C.



Species (Common Name)	Geographic Distribution	Reservoir Host of Definitive Stage	Vector, or Obligate Host Other than Man	Infective Stage
(A)	(B)	(C)	(D)	(E)
<b>Nematoda</b>				
1 <i>Ancylostoma braziliense</i> (hookworm)	Limited distribution in warm climates	Cat, dog	None	Filariform larva
2 <i>A. duodenale</i> (hookworm)	Tropical & subtropical Africa, Asia, Europe, United States; western S. America	None	None	Filariform larva
3 <i>Ascaris lumbricoides</i> (large roundworm)	Worldwide; more common in warm climates	Swine ?	None	Fully embryonated egg
4 <i>Brugia malayi</i> (Malayan filarial worm)	Warm climates in Asia	Cat, monkey	Mosquito ( <i>Armi-geres, Mansonia</i> )	Filariform larva
5 <i>Dracunculus medinensis</i> (guinea worm)	Warm climates of eastern hemisphere	Fur-bearing mammals	<i>Cyclops</i>	3rd stage larva in <i>Cyclops</i>
6 <i>Enterobius vermicularis</i> (pinworm)	Worldwide; common in children	None	None	Fully embryonated egg
7 <i>Loa loa</i> (African filarial worm)	Tropical Africa	None	Mango fly ( <i>Chrys-ops</i> )	Filariform larva
8 <i>Necator americanus</i> (hookworm)	Warm climates	None	None	Filariform larva
9 <i>Onchocerca volvulus</i> (convoluted filarial worm)	Tropical Africa, Mexico, Guatemala, eastern Venezuela, Dutch Guiana ?	None	Blackfly ( <i>Simulium</i> )	Filariform larva
10 <i>Strongyloides stercoralis</i> (intestinal threadworm)	Warm, moist climates	Chimpanzee, dog	None	Filariform larva
11 <i>Trichinella spiralis</i> (trichina worm)	Worldwide; common in United States	Bear, swine, walrus	None	Larva encysted in pork muscle
12 <i>Trichuris trichiura</i> (human whipworm)	Warm, moist climates	Ape, monkey; swine ?	None	Fully embryonated egg
13 <i>Wuchereria bancrofti</i> (Bancroft's filarial worm)	Warm climates	None	Mosquito ( <i>Aedes, Culex, etc.</i> )	Filariform larva
<b>Cestoda</b>				
14 <i>Diphyllobothrium latum</i> (fish tapeworm)	North temperate & sub-arctic zones; lakes in Argentina, Chile	Bear, cat, dog	<i>Cyclops, Diaplo-mus</i> ; freshwater fish	Sparganum larva in fish flesh
15 <i>Dipylidium caninum</i> (double-pored dog tapeworm)	Warm climates	Cat, dog	Cat flea, dog flea, human flea	Larva in hemocoel of dog flea
16 <i>Echinococcus granulosus</i> (hydatid tapeworm)	Worldwide; common in southern S. America	Dog, wild relatives	Cattle, sheep, swine (alternating with dog)	Eggs in dog's excreta
17 <i>E. multiocularis</i> (multi-ocular hydatid tapeworm)	North temperate zones	Wild canids	Small rodents	Eggs in excreta of wild canids
18 <i>Hymenolepis diminuta</i> (rat tapeworm)	Warm & temperate climates	Mouse, rat	Grain beetle, meal moth, rodent flea, etc.	Larva in hemocoel of insect
19 <i>H. nana</i> (dwarf tapeworm)	Warm & temperate climates	Mouse, rat	None; may develop in grain beetle	Egg
20 <i>Taenia saginata</i> (beef tapeworm)	Worldwide	None	Cattle	Cysticercus larva in beef
21 <i>T. solium</i> (pork tapeworm)	Worldwide	None	Swine	Cysticercus larva in pork; egg

/1/ By direct or indirect contact with body excreta containing parasite. /2/ From proboscis of insect vector at time mouth.

## PARASITES: MAMMALS AND BIRDS

### MAN

Portal of Infection	In Man			Identification of Parasite	
	Immature Stage	Definitive Stages			
		Primary Site	Secondary Site		
(F)	(G)	(H)	(I)	(J)	
<b>Nematoda</b>					
Skin <sup>-</sup>	Larva migrates from skin via blood and lungs to epiglottis and GI tract	Attached to small intestine	None	Egg in feces; larva in cutaneous tunnels	1
Skin <sup>1</sup>	Larva migrates from skin via blood and lungs to epiglottis and GI tract	Attached to small intestine	None	Egg in feces	2
Mouth <sup>1</sup>	Larva migrates	Lumen of small intestine	Various viscera	Egg in feces	3
Skin <sup>2</sup>	Larva migrates in lymphatics	Lymphatics of lower trunk	Lymphatics of upper trunk	Microfilaria (sheathed) in peripheral blood (nocturnal and sub-periodic)	4
Mouth <sup>2</sup>	In viscera	Gravid female migrates to skin	None known	Gravid female in ruptured skin blister	5
Mouth <sup>1</sup>	In transit down small intestine	Attached to cecum, appendix	Female genital tract, perianal folds	Egg or adult in anal swab, anus	6
Skin <sup>2</sup>	Migrates in subcutaneous tissues	Migrates in subcutaneous tissues	Orbit, conjunctiva of eye	Microfilaria (sheathed) in diurnal blood	7
Skin <sup>1</sup>	Larva migrates	Attached to small intestine	None	Egg in feces	8
Skin <sup>2</sup>	Larva in skin, may invade eye tissues	Adult in subcutaneous nodules; larva in skin, may invade eye tissues	None known	Microfilaria (unsheathed) in skin biopsy	9
Skin <sup>4</sup>	Larva migrates	Within intestinal mucosa	Lungs	Larva in feces or duodenal aspirate	10
Mouth <sup>3</sup>	Enters duodenal mucosa	In duodenal mucosa	Larva migrates; encysts in striped muscle	Larva in compressed or digested muscle	11
Mouth <sup>1</sup>	In transit down small intestine	Attached to cecum, appendix	Colon, rectum	Egg in feces	12
Skin <sup>2</sup>	Larva migrates in lymphatics	Lymphatics of lower trunk, legs	Lymphatics of upper trunk	Microfilaria (sheathed) in blood (usually nocturnal)	13
<b>Cestoda</b>					
Mouth	Develops in small intestine	Attached to small intestine	None known	Egg in feces	14
Mouth	Develops in small intestine	Attached to small intestine	None	Proglottid in feces	15
Mouth	Develops in liver, lungs	Attached to small intestine	Hydatid cysts in viscera	Hydatid cysts with scolices during aspiration or exploratory operation	16
Mouth	Develops unconfined in liver	Attached to small intestine	Hydatid cysts in liver	Hydatid cysts with scolices during postmortem examination	17
Mouth	Develops in duodenum, small intestine	Attached to small intestine	None	Proglottid or egg in feces	18
Mouth	Develops in duodenal villi	Attached to small intestine	None known	Egg in feces	19
Mouth	Develops in small intestine	Attached to small intestine	None known	Proglottid or egg in feces	20
Mouth	Develops in small intestine	Attached to small intestine	Cysticercus larva in various stages	Proglottid or egg in feces	21

of skin puncture to obtain blood or tissue juice from host. /a/ From infected food or contaminated water taken into

*continued*

## 130. HELMINTH AND PROTOZOAN

Part I.

Species (Common Name)	Geographic Distribution	Reservoir Host of Definitive Stage	Vector, or Obligate Host Other than Man	Infective Stage
(A)	(B)	(C)	(D)	(E)
Trematoda				
22 <i>Clonorchis sinensis</i> (Chinese liver fluke)	Sino-Japanese & Indo-Chinese areas	Many fish-eating mammals	Freshwater fishes, snail	Larva encysted in flesh of freshwater fish
23 <i>Fasciola hepatica</i> (liver fluke)	Sheep-raising countries	Herbivores	Snail, moist vegetation	Larva encysted on water plants
24 <i>Fasciolopsis buski</i> (intestinal fluke)	Oriental countries	Swine	Snail, water plants	Larva encysted on water plants
25 <i>Paragonimus westermani</i> (lung fluke)	Sino-Japanese areas, Southwest Pacific islands, northern S. America	Cat, dog, swine, other animals	Crab, crayfish, and snail; sputum of man	Larva encysted in soft tissues of crabs, crayfish
26 <i>Schistosoma haematobium</i> (human blood fluke)	Africa, Near East, Middle East, southern Portugal	Gerbil	Bulinid snail	Cercaria free in freshwater
27 <i>S. japonicum</i> (blood fluke)	China, Japan, Philippines, Formosa, Celebes	Many mammals	Oncomelaniid snail	Cercaria free in freshwater
28 <i>S. mansoni</i> (blood fluke)	Africa, Arabia, Brazil, Guianas, Venezuela, West Indies	Monkey (rarely)	Planorbid snail	Cercaria free in freshwater
Protozoa				
29 <i>Balantidium coli</i>	Worldwide; most common in warm climates	Monkey (?), swine	None	Mature cyst
30 <i>Entamoeba coli</i>	Worldwide; most common in warm climates	Monkey ?	None	Four-nucleate cyst
31 <i>E. histolytica</i>	Worldwide; most common in warm climates	Dog, monkey; rat ?	None	Four-nucleate cyst
32 <i>Giardia lamblia</i>	Worldwide; most common in warm climates	None	None	Four-nucleate cyst
33 <i>Leishmania braziliensis</i>	Western hemisphere from southern Mexico to northern Argentina	Dog, possibly other mammals	Sand fly ( <i>Phlebotomus</i> )	Leptomonad
34 <i>L. donovani</i>	China, India, Africa, Mediterranean area, S. America	Dog, rodents	Sand fly ( <i>Phlebotomus</i> )	Leptomonad
35 <i>L. tropica</i>	Western India, Middle East, Near East, N. Africa	Dog, rodents	Sand fly ( <i>Phlebotomus</i> )	Leptomonad
36 <i>Plasmodium falciparum</i> , <i>P. malariae</i> , <i>P. vivax</i>	Temperate or warm climates	None	Mosquito ( <i>Anopheles</i> )	Sporozoite
37 <i>Toxoplasma gondii</i>	Worldwide	Many mammals and birds	None known	Trophozoite
38 <i>Trichomonas vaginalis</i>	Worldwide (relatively common in both sexes)	None	None	Trophozoite (only stage known)
39 <i>Trypanosoma cruzi</i>	Western hemisphere from United States to northern Argentina	Many mammals	Triatomid bug	Metacyclic trypanosome
40 <i>T. gambiense</i>	Western & central Africa	Cattle ?	Tsetse fly ( <i>Glossina</i> )	Metacyclic trypanosome
41 <i>T. rhodesiense</i>	Central & eastern Africa	Mammals, wild game	Tsetse fly ( <i>Glossina</i> )	Metacyclic trypanosome

<sup>1</sup>/ By direct or indirect contact with body excreta containing parasite. <sup>2</sup>/ From proboscis of insect vector at time mouth. <sup>4</sup>/ In contact with infested water. <sup>5</sup>/ From feces of insect vector while feeding on blood or tissue juice of

Contributor: Faust, Ernest Carroll

Reference: Faust, E. C., Beaver, P. C., and Jung, R. C. 1962. Animal agents and vectors of human disease. Ed. 2.

## PARASITES: MAMMALS AND BIRDS

### MAN

In Man					Identification of Parasite
Portal of Infection	Immature Stage	Definitive Stages		(J)	
		Primary Site	Secondary Site		
(F)	(G)	(H)	(I)		
<b>Trematoda</b>					
Mouth <sup>3</sup>	In transit from duodenum to bile ducts	Distal bile ducts	Pancreatic ducts (rare)	Egg in feces	22
Mouth <sup>3</sup>	In transit from duodenum to bile ducts	Proximal bile ducts	Abdominal wall (?), lungs, brain	Egg in feces	23
Mouth <sup>3</sup>	Develops in duodenum, jejunum	Attached to duodenum, jejunum	None	Egg in feces	24
Mouth <sup>3</sup>	In transit from duodenum to lungs	Lungs, near bronchioles	Abdominal viscera, brain	Egg in sputum or feces	25
Skin <sup>4</sup>	Migrates in blood vessels	Vesical venous plexus	Pelvic organs, rectum, lungs, central nervous system	Egg in urine or feces	26
Skin <sup>4</sup>	Migrates in blood vessels	Mesenteric venules	Liver, lungs, brain	Egg in feces	27
Skin <sup>4</sup>	Migrates in blood vessels	Mesenteric venules	Liver, lungs, brain	Egg in feces	28
<b>Protozoa</b>					
Mouth <sup>1</sup>	None described	Wall of large intestine	None	Trophozoite or cyst in feces	29
Mouth <sup>1</sup>	None described	Lumen of large intestine	None	Trophozoite or cyst in feces	30
Mouth <sup>1</sup>	None described	Wall of large intestine	Other viscera, skin	Trophozoite or cyst in feces, visceral abscesses, or skin abscesses	31
Mouth <sup>1</sup>	None described	Duodenal crypts	Gallbladder ?	Trophozoite or cyst in feces	32
Skin <sup>2</sup>	None described	Skin	Mucous membranes	Leishmanial stage in reticuloendothelial cells, skin, or viscera Leptomonad stage in culture	33
Skin <sup>2</sup>	None described	Skin	Reticuloendothelium (fundamental)	Leishmanial stage in reticuloendothelial cells, skin, or viscera Leptomonad stage in culture	34
Skin <sup>2</sup>	None described	Skin	Mucous membranes (rare)	Leishmanial stage in reticuloendothelial cells, skin, or viscera Leptomonad stage in culture	35
Skin <sup>2</sup>	Schizonts in hepatic parenchyma	Exoerythrocytic foci	Erythrocytes	Trophozoite, schizont, or gametocyte in blood	36
Unknown	None known	Reticuloendothelium, many parenchymal cells	Brain, retina	Trophozoite, pseudocyst, or cyst in focal areas of necrosis	37
Vulva <sup>1</sup> or urethra	None described	Vaginal fold	Bladder	Trophozoite in urine or vaginal smear	38
Skin <sup>5</sup> , conjunctiva	None described	Skin	In tissues, blood	Trypanosomal stage in blood or tissues Leishmanial stage in macrophage Crithidial stage in culture	39
Skin <sup>2</sup>	None described	Skin	Blood, lymph nodes, central nervous system	Trypanosomal stage in blood, gland juice, or spinal fluid	40
Skin <sup>2</sup>	None described	Skin	Blood, lymph nodes, central nervous system	Trypanosomal stage in blood, gland juice, or spinal fluid	41

of skin puncture to obtain blood or tissue juice from host. /<sup>3</sup>/ From infected food or contaminated water taken into host.

Lea and Febiger, Philadelphia.

*continued*



## 130. HELMINTH AND PROTOZOAN

### Part II. VERTEBRATES

	Species (Common Name)	Geographic Distribution	Intermediate Host
	(A)	(B)	(C)
Acanthocephala			
1	<i>Macracanthorhynchus hirudinaceus</i> (thorny-headed worm)	Worldwide	Beetle ( <i>Cotinis</i> , <i>Phyllophaga</i> )
Nematoda			
2	<i>Ancylostoma caninum</i> (dog hookworm)	Worldwide	None
3	<i>Ascaridia galli</i> (large roundworm of chicken)	Worldwide	None
4	<i>Ascaris lumbricoides suum</i> (large roundworm)	Worldwide	None
5	<i>Dictyocaulus filaria</i> (thread lungworm of sheep)	Worldwide	None
6	<i>D. viviparus</i> (lungworm of cattle)	Worldwide	None
7	<i>Dirofilaria immitis</i> (dog heartworm)	Worldwide	Mosquito ( <i>Aedes</i> , <i>Anopheles</i> , <i>Culex</i> )
8	<i>Haemonchus contortus</i> (twisted stomach worm)	Worldwide	None
9	<i>Heterakis gallinae</i> (cecal worm)	Worldwide	None
10	<i>Metastrongylus elongatus</i> (swine lungworm)	Worldwide	Earthworm (several genera including <i>Eisenia</i> , <i>Lumbricus</i> )
11	<i>Oesophagostomum columbianum</i> (sheep nodular worm)	Worldwide	None
12	<i>Ostertagia circumcincta</i> (brown stomach worm of sheep)	Worldwide	None
13	<i>O. ostertagi</i> (brown stomach worm of cattle)	Worldwide	None
14	<i>Parascaris equorum</i> (large roundworm)	Worldwide	None
15	<i>Strongyloides stercoralis</i> (intestinal threadworm)	Cosmopolitan in warm climates	None
16	<i>Strongylus vulgaris</i> (single-toothed strongyle)	Worldwide	None
17	<i>Toxascaris leonina</i> (dog roundworm)	Worldwide	None
18	<i>Toxocara canis</i> (dog roundworm)	Worldwide	None
19	<i>Trichinella spiralis</i> (trichina worm)	Worldwide	Same individual both definitive and intermediate host <sup>2</sup>
20	<i>Trichostrongylus axei</i> (minute stomach worm)	Worldwide	None
21	<i>T. colubriformis</i> (hairworm)	Worldwide	None
22	<i>Trichuris suis</i> (swine whipworm)	Worldwide	None
23	<i>T. vulpis</i> (dog whipworm)	Worldwide	None
Cestoda			
24	<i>Dibothriocephalus latus</i> (fish tapeworm)	N. America, Argentina, Chile, Europe, Australia, Manchuria, Siberia, Japan	First: copepod ( <i>Cyclops</i> , <i>Diaplo-mus</i> ); second: fish
25	<i>Dipylidium caninum</i> (double-pored dog tapeworm)	Worldwide	Flea ( <i>Ctenocephalides</i> , <i>Pulex</i> ); louse ( <i>Trichodectes</i> )
26	<i>Echinococcus granulosus</i> (hydatid tapeworm)	N. & S. America, Europe, Iceland, Australia, northern Asia, Africa	Camel, cow, dog, goat, horse, monkey, moose, rabbit, sheep, rodents, etc. <sup>1</sup>
27	<i>Hymenolepis carioca</i> (thread tapeworm)	Worldwide	Many beetles ( <i>Anisotarsus</i> , <i>Aphodius</i> , and others); stable fly ( <i>Stomoxys</i> )
28	<i>Moniezia expansa</i> (sheep tapeworm)	Worldwide	Grass mite ( <i>Galumna</i> , <i>Oribatula</i> , and others)

[1] Also man. [2] Reservoir host: swine for man.

PARASITES: MAMMALS AND BIRDS

OTHER THAN MAN

Definitive Host	Primary Location in Definitive Host	Disease or Disorder	
(D)	(E)	(F)	
Acanthocephala			
Swine	Small intestine	Nodule formation	1
Nematoda			
Cat, coyote, dog, fox <sup>1</sup>	Small intestine	Anemia, emaciation, skin reactions	2
Chicken, goose, guinea fowl, turkey, wild birds	Small intestine	Emaciation	3
Swine <sup>2</sup>	Small intestine	Pneumonia, abdominal discomfort and obstruction, emaciation	4
Goat, sheep, some wild ruminants	Bronchi, bronchioles	Catarrhal inflammation, coughing, emaciation	5
Cattle, deer	Bronchi, bronchioles	Catarrhal inflammation, coughing, emaciation	6
Cat, coyote, dog, fox, wolf	Heart, pulmonary artery; microfilariae in blood	Emaciation, cough, edema, dyspnea	7
Cattle, goat, sheep, other ruminants	Abomasum	Anemia, emaciation	8
Chicken, guinea fowl, pheasant, quail, turkey, other birds	Cecum	None; egg carries <i>Histomonas</i>	9
Swine	Bronchi, bronchioles	Bronchitis, pneumonia; transmits swine influenza virus	10
Antelope, goat, sheep	Large intestine; larvae in nodules throughout intestine	Diarrhea, emaciation, nodules in intestine	11
Goat, sheep	Abomasum	Anemia, emaciation	12
Cattle, sheep, rarely horse	Abomasum	Anemia, edema, emaciation	13
Horse, other equids	Small intestine	Pneumonia, digestive disturbances, emaciation	14
Cat, dog, fox <sup>3</sup>	Small intestine mucosa	Diarrhea	15
Horse, other equids	Large intestine	Anemia, edema, digestive disturbances, emaciation; larvae form aneurysms in anterior mesenteric arteries	16
Cat, dog, fox, wild canids and felids	Small intestine	Emaciation, poor growth	17
Coyote, dog, fox	Small intestine	Emaciation, poor growth	18
Badger, rat, swine, many other mammals <sup>4</sup>	Small intestine; larvae in muscles	Trichinosis, toxemia, muscle pains	19
Cattle, deer, goat, horse, sheep, swine <sup>5</sup>	Abomasum	Emaciation	20
Antelope, camel, cattle, goat, sheep	Small intestine	Emaciation	21
Ape, monkey, swine <sup>1</sup>	Cecum	Toxemia	22
Dog, fox	Cecum	Emaciation, low-grade inflammation	23
Cestoda			
Cat, dog, fox, polar bear, other fish-eating mammals <sup>1</sup>	Small intestine	Toxemia, anemia	24
Cat, dog, fox, wolf, other carnivores <sup>1</sup>	Small intestine	Enteritis, anal pruritus	25
Dog, fox, wolf, other canids	Small intestine	Slight, if any, enteritis; hydatid cysts in liver, lungs, etc., of intermediate hosts cause serious damage	26
Chicken, quail, turkey	Small intestine	Slight damage	27
Cattle, goat, sheep, other ruminants	Small intestine	Emaciation	28

*continued*

## 130. HELMINTH AND PROTOZOAN

### Part II. VERTEBRATES

	Species (Common Name)	Geographic Distribution	Intermediate Host
(A)	(B)	(C)	
Cestoda			
29	<i>Railletina cesticillus</i> (broad-headed tapeworm)	Worldwide	Ground and dung beetles (several genera)
30	<i>Taenia pisiformis</i> (dog tapeworm)	Worldwide	Hare, rabbit, rat, squirrel
Trematoda			
31	<i>Fasciola hepatica</i> (liver fluke)	Worldwide	Freshwater snail ( <i>Fossaria</i> , <i>Galba</i> , <i>Lymnaea</i> , <i>Pseudosuccinea</i> , and others) <sup>3</sup>
32	<i>Nanophyetus salmincola</i> ("salmon-poisoning" fluke)	Pacific Northwest	First: snail ( <i>Goniobasis</i> ); second: fish (usually <i>Salmo</i> , also <i>Onco-rhynchus</i> , <i>Salvelinus</i> )
Protozoa			
33	<i>Babesia bigemina</i>	N., Cen., & S. America, Europe, Australia, Asia, Africa, Pacific islands	Tick ( <i>Boophilus</i> , <i>Rhipicephalus</i> )
34	<i>Balantidium coli</i>	Worldwide	None
35	<i>Eimeria ahsala</i>	Worldwide	None
36	<i>E. necatrix</i>	Worldwide	None
37	<i>E. tenella</i>	Worldwide	None
38	<i>E. zurnii</i>	Worldwide	None
39	<i>Histomonas meleagridis</i>	Worldwide	None; transmitted in <i>Heterakis gallinae</i> eggs <sup>4</sup>
40	<i>Iodamoeba buetschlii</i>	Worldwide	None
41	<i>Isospora bigemina</i>	Worldwide	None
42	<i>Leishmania donovani</i>	Cen. & S. America, Mediterranean basin, Balkan states, Near East, India, China, Russia, Africa	Sand fly ( <i>Phlebotomus</i> ) <sup>5</sup>
43	<i>L. tropica</i>	Mediterranean basin, Near East, India, Russia, Africa	Sand fly ( <i>Phlebotomus</i> ) <sup>5</sup>
44	<i>Theileria annulata</i>	Southern Europe, Asia, Africa	Tick ( <i>Hyalomma</i> )
45	<i>Toxoplasma gondii</i>	Worldwide	None? <sup>7</sup>
46	<i>Trichomonas gallinae</i>	Worldwide	None
47	<i>Tritrichomonas foetus</i>	Worldwide	None
48	<i>Trypanosoma brucei</i>	Africa	Tsetse fly ( <i>Glossina</i> ) <sup>6</sup>
49	<i>T. cruzi</i>	Southwestern United States, Cen. & S. America	Kissing bug ( <i>Panstrongylus</i> , <i>Triatomu</i> ); assassin bug ( <i>Rhodinus</i> ) <sup>8</sup>
50	<i>T. evansi</i>	Cen. & S. America, southeastern Europe, Asia, Africa	Stable fly ( <i>Stomoxys</i> ), horsefly ( <i>Tabanus</i> ) <sup>9</sup>

<sup>1</sup>/ Also man. <sup>2</sup>/ Reservoir host: wild rabbit for ruminants. <sup>3</sup>/ Reservoir hosts: chicken and wild gallinaceous <sup>4</sup>/ Reservoir hosts: wild rodents. <sup>5</sup>/ Reservoir hosts: wild ruminants and equids. <sup>6</sup>/ Reservoir hosts: wild ani-

Contributor: Levine, Norman D.

References: [1] Chandler, A. C., and C. P. Read. 1961. Introduction to parasitology. Ed. 10. J. Wiley, New York, Baltimore. [2] Levine, N. D. 1961. Protozoan parasites of domestic animals and of man. Burgess, Minneapolis. and J. A. McLeod. 1952. The zoology of tapeworms. Univ. Minnesota Press, Minneapolis.

## PARASITES: MAMMALS AND BIRDS

### OTHER THAN MAN

Definitive Host	Primary Location in Definitive Host	Disease or Disorder	
(D)	(E)	(F)	
Cestoda			
Chicken, pheasant, quail, turkey, wild galliforms	Small intestine	Slight damage	29
Cat, coyote, dog, fox, wolf	Small intestine	Slight, if any, enteritis; anal pruritus	30
Trematoda			
Cat, dog, elephant, hare, horse, kangaroo, rabbit, swine, rodents; cattle, goat, sheep, other ruminants <sup>1</sup>	Proximal bile duct	Liver necrosis, cirrhosis, calcification of bile ducts	31
Coyote, dog, fox, lynx, mink, raccoon	Small intestine	Enteritis; parasite carries <i>Neorickettsia helminthoeca</i> , the cause of "salmon poisoning"	32
Protozoa			
Cattle	Erythrocytes	Fever, anemia, hemoglobinuria; causes Texas fever	33
Monkey, swine <sup>1</sup>	Large intestine	Secondary invader of mucosa	34
Goat, sheep	Small intestine cells	Diarrhea, emaciation	35
Chicken	Small intestine cells	Hemorrhagic enteritis	36
Chicken	Cecal cells	Hemorrhagic enteritis	37
Cattle	Intestinal cells	Enteritis, hemorrhagic dysentery	38
Chicken, partridge, peafowl, pheasant, quail, ruffed grouse	Cecum, liver	Enterohepatitis, necrosis, ulceration	39
Monkey, swine <sup>1</sup>	Large intestine	None	40
Cat, dog, fox, mink	Small intestine cells	Diarrhea	41
Dog <sup>1</sup>	Reticuloendothelial system	Kala azar; reticuloendotheliosis, splenomegaly	42
Dog, gerbil <sup>1</sup>	Cutaneous tissues	Oriental sore, skin ulcer	43
Cattle, zebu	Lymphocytes, erythrocytes	Fever, anemia, emaciation	44
Birds; cat, dog, rodents, other mammals <sup>1</sup>	Endothelial cells, leucocytes	Chorioretinitis, cerebral calcification, pneumonia	45
Chicken, dove, hawk, pigeon, turkey	Crop, esophagus	Caseous nodules, necrosis	46
Cattle	Uterus, genital system, prepuce and penile membranes	Abortion, estrus irregularity, macerated fetus, contaminated semen	47
Cat, dog, swine, equids, ruminants	Blood	Anemia, emaciation, edema; causes nagana	48
Armadillo, bat, cat, dog, monkey, opossum, wood rat <sup>1</sup>	Blood, myocardium, and other tissues	Chagas' disease; tissue destruction	49
Cat, dog, elephant, swine, equids, ruminants	Blood	Urticaria, edema, emaciation; surra	50

birds for turkey. /s/ Reservoir host: dog for man. /s/ Reservoir hosts: dog, gerbil and *Rhombomys* for man.

[2] Lapage, G., ed. 1962. Mönning's Veterinary helminthology and entomology. Ed. 5. Williams and Wilkins.

[4] Morgan, B. B., and P. A. Hawkins. 1949. Veterinary helminthology. Burgess, Minneapolis. [5] Wardle, R. A.,



## 131. NEMATODE

Most of the nematode parasites of plants are found in close association with the roots, or in the upper 16 inches of ceous forms, also found in the soil, by the presence of a protrusile spear or stylet used to puncture and feed on plant others live only in the ocean, and many are parasites of animals and man.

Species (Common Name)	Geographic Distribution <sup>1</sup>	Host <sup>2</sup>
(A)	(B)	(C)
1 <i>Anguina</i> spp. (wheat gall eel-worm)	N. America, Europe	Several <i>Agrostis</i> species; other grasses and cereals
2 <i>A. tritici</i> (wheat nematode)	Southern Atlantic states, Europe, southern & eastern Asia, Egypt, Australia	Emmer, rye, spelt, wheat
3 <i>Aphelenchoides besseyi</i> (summer crimp nematode of strawberry)	Southeastern United States (Maryland to Texas)	Rice, strawberry
4 <i>A. cocophilus</i> (coconut palm nematode)	West Indies, Honduras, Panama, British Guiana, Venezuela	Coconut, date, and oil palms
5 <i>A. fragariae</i> (spring crimp nematode of strawberry)	Massachusetts, Connecticut, Delaware, Maryland, Europe	Strawberry
6 <i>A. ritzei-bosi</i> (chrysanthemum nematode)	N. America, Europe	About 50 different plants, including chrysanthemum, larkspur, phlox, strawberry, verbena, zinnia
7 <i>Belonolaimus gracilis</i> (sting nematode)	Southern Atlantic states	Bean, beet, cabbage, celery, citrus, corn, cotton, cowpea, grass, millet, okra, onion, peanut, pine seedling, soybean, strawberry
8 <i>Criconeoides</i> spp. (ring nematode)	Widespread	Many plants; reported as injuring peach trees and peanut vines
9 <i>Ditylenchus destructor</i> (potato rot nematode)	Idaho, Wisconsin, Prince Edward Island, Europe	Carrot, iris, potato, sweet potato, tulip
10 <i>D. dipsaci</i> (bulb and stem nematode)	Widespread in temperate zones	Over 300 different plants, including alfalfa, clover, hyacinth, iris, narcissus, oats, onion, phlox, potato
11 <i>Dolichodorus heterocephalus</i> (awl nematode)	Florida, Georgia, North Carolina, Michigan	Bean, celery, Chinese water chestnut, corn, tomato; many other plants growing in wet locations
12 <i>Helicotylenchus</i> spp., <i>Rotylenchus</i> spp. (spiral nematode)	Widespread in sub-tropical and tropical regions	Many plants, including bean, cotton, cowpea, grass, pineapple, soybean, and ornamentals
13 <i>Heterodera glycine</i> (soybean cyst nematode)	Midwestern & southern United States, China, Japan	Adzuki bean, annual lespedeza, kidney bean, snap bean, soybean, vetch
14 <i>H. rostochiensis</i> (golden nematode of potato)	Long Island, N. Y.; Bolivia, Peru, Europe	Potato, tomato, several other solanaceous plants
15 <i>H. schachtii</i> (sugar beet nematode)	United States, Canada, Europe, Australia	Over 100 plants, including broccoli, cabbage, cauliflower, kale, mangel-wurzel, mustard, rutabaga, sugar beet, table beet, turnip
16 <i>Hoplotylenchus tylenchiformis</i> (lance nematode)	N. America, Philippines, Europe	Many plants, including corn, cotton, pine tree, sugarcane, St. Augustine and other lawn grasses

<sup>1/</sup> Information on geographic distribution of plant parasitic nematodes is fragmentary and incomplete, even for the genus vary in ability to attack plants; some have a rather wide host range, others are highly host-specific, attacking stage of development of the host and parasite. <sup>2/</sup> Symptoms of nematode damage are often difficult to distinguish find the nematode in the diseased tissue or soil adjacent to the roots of affected plants.

## PARASITES: PLANTS

soil formerly occupied by the roots. In general, plant nematodes can be distinguished from saprophagous or predaceous. The soil is not the only habitat, however, for nematodes: some live in freshwater rivers, lakes, and ponds,

Feeding Habits <sup>3</sup>	Symptoms <sup>4</sup>	Control	Reference	
(D)	(E)	(F)	(G)	
Larvae, ectoparasites around growing point; adults, endoparasites of flower primordia	Abnormal flowers developing into galls	Crop rotation; planting of gall-free seed	21,24	1
Larvae, ectoparasites around growing point; adults, endoparasites of flower primordia	Stunted plants, distorted foliage, galls instead of seed	Planting of gall-free seed (galls may be removed by salt brine flotation or mechanical separators)	18, 34	2
Vagrant ectoparasites of buds and growing point between young developing leaves	Small, crinkled, distorted foliage	Setting of new beds with uninfested plants; hot-water treatment or methyl bromide fumigation for rice seed	13,22	3
Vagrant endoparasites of roots, trunk (near periphery), leaf petioles	Disintegration of trunk tissues (causing "red ring") and of root cortex	No established control methods	40	4
Vagrant ectoparasites of buds between young developing leaves	Small, crinkled, distorted foliage	Setting of new beds with uninfested plants	5,13	5
Vagrant endoparasites of buds and foliage	Crinkled, distorted leaves and leaf spots	Hot-water treatment of dormant plants; parathion sprays	23,25,26, 46	6
Vagrant ectoparasites of root tips, sides of succulent roots, other underground parts	Devitalized root tips, root lesions, causing many short stubby branched roots, severely stunted plants	Soil fumigation	15,16,31, 39	7
Semi-sedentary ectoparasites of roots, other underground parts	Small lesions, stunting of plant	Soil fumigation	11,37	8
Vagrant endoparasites of tubers and, to some extent, of roots	Destruction of tuber tissues causing sunken areas, followed by rot	Crop rotation; planting of clean seed; soil fumigation	4,52	9
Vagrant endoparasites of bulbs, stems, leaves, occasionally roots	Twisting, wrinkling, distortion of stems and flowers; necrosis and destruction of bulb tissues	Hot-water treatment of bulbs, corms; crop rotation; field sanitation; methyl bromide fumigation of infected onion and clover seeds; planting of resistant varieties	1,20,48	10
Vagrant ectoparasites of root tips, sides of succulent roots	Devitalized root tips, small lesions on sides of roots; sometimes extensive root destruction	Soil fumigation	41,50	11
Vagrant ectoparasites, occasionally endoparasites of roots and other underground parts	Stunting of plant from retarded root growth; lesions may occur	Soil fumigation	29,47	12
Sedentary parasites of roots, internal in early stages, external as adults	General stunting of plants, reduction in size of root system; causes disease known as "yellow dwarf" in Japan and China	Crop rotation; soil fumigation; planting of resistant varieties	32,57	13
Sedentary parasites, internal in early stages, becoming largely external as adults; attack roots, other underground parts	Stunting of plant, decrease in size of root system; often increase in number of small branch rootlets	Crop rotation; soil fumigation; planting of resistant varieties	12,38	14
Sedentary parasites of roots, other underground parts; internal in early stages, external as adults	Stunting of plant, overall decrease in size of root system; often increase in number of small branch rootlets	Crop rotation; soil fumigation with dichloropropene-dichloropropane mixture before planting (sugar beet)	19,27,42, 53	15
Vagrant internal or partly external parasites of roots	Lesions leading to complete destruction and sloughing off of cortex	Soil fumigation	33	16

best known species. Undoubtedly distribution is far wider than indicated. /a/ Species of nematodes within a given only one or two crop plants. /a/ Feeding habits and particular tissues attacked vary with the species, host plant, and from those caused by other organisms or by poor growing conditions; hence it is important in making a diagnosis to

*continued*

## 131. NEMATODE

Species (Common Name)	Geographic Distribution <sup>1</sup>	Host <sup>2</sup>
(A)	(B)	(C)
17 <i>Meloidogyne</i> spp. (root-knot nematode)	Worldwide; most common in warm climates	Over 2,000 plants; hosts of individual species more restricted
18 <i>Paratylenchus</i> spp. (pin nematode)	N. America, Hawaii, British Isles, Netherlands, western Africa	Many plants, including alfalfa, cabbage, celery, cowpea, cucumber, fig tree, oats, okra, pineapple, radish, wheat
19 <i>Pratylenchus</i> spp. (lesion nematode)	Worldwide	Many plants, including alfalfa, corn, cotton, small grains, strawberry, tobacco, trees and shrubs
20 <i>Radopholus oryzae</i> (rice-root nematode)	Louisiana, Texas, Indonesia, Japan, rice-growing areas of southeastern Asia	Rice, various grasses, and related monocotyledonous plants
21 <i>R. similis</i> (burrowing nematode)	Florida, Louisiana, Jamaica, West Indies, Cen. America, Peru, Brazil, Hawaii, Philippines, Formosa, Indonesia, India	About 50 different plants, including avocado, banana, black pepper, canna, citrus, coffee, rice, sugarcane, tea
22 <i>Trichodorus</i> spp. (stubby root nematode)	Widespread; important in southeastern United States, southern California, Nicaragua, Tunisia	Many plants, including beet, cabbage, cauliflower, celery, chayote, corn, cotton, fig
23 <i>Tylenchorhynchus</i> spp. (stunt nematode)	Apparently widespread	Many plants, including azalea, cotton, oats, sugarcane, tobacco, wheat
24 <i>Tylenchulus semipenetrans</i> (citrus nematode)	Florida, Texas, California, most citrus fruit-growing regions; southern Europe	Most <i>Citrus</i> and closely related genera; olive
25 <i>Xiphinema</i> spp. (dagger nematode)	Worldwide	Many plants, shrubs, trees, including clove, corn, laurel oak, oats, pecan, rose, strawberry, some grasses

<sup>1</sup>/ Information on geographic distribution of plant parasitic nematodes is fragmentary and incomplete, even for the genus vary in ability to attack plants; some have a rather wide host range, others are highly host-specific, attacking stage of development of the host and parasite. <sup>2</sup>/ Symptoms of nematode damage are often difficult to distinguish find the nematode in the diseased tissue or soil adjacent to the roots of affected plants. <sup>3</sup>/ All species in the genus wood and potato tubers, respectively.

**Contributors:** (a) Christie, Jesse R., (b) Sasser, J. N.

**References:** [1] Anonymous. 1951. Natl. Inst. Agr. Botany (Gr. Brit.), Seed Notes 38. [2] Atkins, J. G., M. J. 1948. Phytopathology 38(11):912. [4] Baker, A. D. 1946. Sci. Agr. 26(3):138. [5] Ballard, E., and G. S. Peren. K. E., and R. W. Hanks. 1954. Ibid. 67:83. [8] Brooks, T. L. 1954. Ibid. 67:81. [9] Buhner, E. M. 1938. Plant B. G. 1949. Proc. Helminthol. Soc. Wash. D. C. 16(1):6. [12] Chitwood, B. G., and E. M. Buhner. 1946. Phyto-Sci. Soc. Florida 12:30. [15] Christie, J. R. 1953. Down Earth 9(1):8. [16] Christie, J. R., A. N. Brooks, and [18] Chu, V. M. 1945. Phytopathology 35(5):288. [19] Corder, M. N., E. M. Buhner, and G. Steiner. 1936. Plant W. D., and H. B. Howell. 1952. Plant Disease Repr. 36(3):75. [22] Cralley, E. M. 1952. Arkansas Farm Res. J. R. Christie. 1937. Ibid. 21(9):144. [25] Dimock, A. W., and C. H. Ford. 1950. Phytopathology 40(1):7. Heterodera. Commonwealth Bureau of Agricultural Parasitology, Farnham Royal, England. [28] Garriss, H. R. [30] Graham, T. W. 1954. Ibid. 44(6):332. [31] Holderman, Q. L. 1955. Plant Disease Repr. 39(1):5. [32] Ichinohe, Assoc. Southern Agr. Workers, 52nd, p. 143. [34] Leukel, R. W. 1929. U.S. Dept. Agr. Farmers' Bull. 1607. Stoddard, and J. W. Lownsbery. 1952. Phytopathology 42(12):651. [37] Machmer, J. H. 1953. Plant Disease 1952. Phytopathology 42(9):470. [40] Nowell, W. 1919. West Indian Bull. 17(4):189. [41] Perry, V. G. 1953. H. W., and M. M. Evans. 1953. Plant Disease Repr. 37(11):540. [44] Schindler, A. F. 1954. Phytopathology L. N. 1947. Agriculture (Engl.) 54(6):278. [47] Steiner, G. 1938. J. Agr. Res. 56(1):1. [48] Steiner, G., and [50] Tarjan, A. C. 1952. Phytopathology 42(2):114. [51] Thomas, E. E. 1923. Calif. Univ. Agr. Expt. Sta. Tech. Agr. Farmers' Bull. 2054. [54] Thorne, G., and M. W. Allen. 1950. Proc. Helminthol. Soc. Wash. D. C. 17(1):27. Bergman. 1952. Ibid. 131. [57] Winstead, N. N., C. B. Skotland, and J. N. Sasser. 1955. Plant Disease Repr.



## PARASITES: PLANTS

Feeding Habits <sup>3</sup>	Symptoms <sup>4</sup>	Control	Reference
(D)	(E)	(F)	(G)
Sedentary endoparasites of roots, other underground parts	Swellings, galls, often local necrosis of tissues; increase or reduction of branch rootlets	Annual crops: rotation and fumigation; planting of resistant varieties; hot-water treatment of bulbs, corms, tubers	9,10,28 17
Vagrant ectoparasites of roots and other underground structures	Stunting of plants from root injury and retarded root growth	Fumigation somewhat effective	35,36,54 18
Vagrant endoparasites of roots and tubers <sup>5</sup>	Small brown root lesions; causes "brown root rot" of tobacco	Crop rotation, tobacco; row fumigation with dichloropropene-dichloropropane mixture	45 19
Vagrant endoparasites of roots	Root lesions, destruction of cortex, root hairs; in Indonesia associated with "mentek," a rice root rot	No established control measures	2,55,56 20
Vagrant endoparasites of roots	Root lesions and disintegration	Hot-water treatment of infected citrus nursery stock; pulling of affected trees, then soil fumigation	6-8,49 21
Vagrant ectoparasites of root tips	Devitalized root tips, causing numerous short, stubby branch rootlets	No satisfactory control known	17 22
Mostly external, occasionally internal vagrant parasites of roots	Stunting of plant	Soil fumigation	30,43 23
Females are sedentary, partly external parasites of roots	Extensive necrosis, discoloration of cortex of small roots	Planting of uninfected stock on clean land	3,51 24
Vagrant ectoparasites of root tips, sides of succulent roots	Devitalized root tips, necrosis of small roots, gall-like swellings, clusters of small stubby branches	Soil fumigation	14,44,54 25

best known species. Undoubtedly distribution is far wider than indicated. /2/ Species of nematodes within a given only one or two crop plants. /3/ Feeding habits and particular tissues attacked vary with the species, host plant, and from those caused by other organisms or by poor growing conditions; hence it is important in making a diagnosis to *Pratylenchus* are root parasites, with the exception of *P. mahogani* and *P. scribneri* observed in diseased mahogany

Fielding, and J. P. Hollis. 1955. Plant Disease Repr. 39(3):221. [3] Baines, R. C., O. F. Clark, and W. P. Bitters, 1923. J. Pomol. Hort. Sci. 3:142. [6] Birchfield, W. 1954. Proc. Florida State Hort. Soc. 67:94. [7] Bragdon, Disease Repr. 22(12):216. [10] Buhner, E. M., C. Cooper, and G. Steiner. 1933. Ibid. 17(7):64. [11] Chitwood, pathology 36(3):180. [13] Christie, J. R. 1943. U.S. Dept. Agr. Circ. 681. [14] Christie, J. R. 1952. Proc. Soil V. G. Perry. 1952. Phytopathology 42(4):173. [17] Christie, J. R., and V. G. Perry. 1951. Science 113:491. Disease Repr. 20(3):38. [20] Courtney, W. D. 1948. Proc. Bulb Growers' Short Course, p. 7. [21] Courtney, 1(1):5. [23] Crossman, L., and J. R. Christie. 1936. Plant Disease Repr. 20(10):155. [24] Crossman, L., and [26] Franklin, M. T. 1950. Ann. Appl. Biol. 37(1):1. [27] Franklin, M. T. 1951. The cyst-forming species of 1953. N. Carolina State Coll. Agr. Ext. Serv. Circ. 374. [29] Golden, A. M. 1954. Phytopathology 44(7):389. M. 1955. Hokkaido Noji Shikensho Hokoku 48:1. [33] Krusberg, L. R., and J. N. Sasser. 1955. Proc. Ann. Conv. [35] Linford, M. B., J. M. Oliveira, and M. Ishii. 1949. Pacific Sci. 3(2):111. [36] Lownsbery, B. F., E. M. Repr. 37(3):156. [38] Mai, W. F., and B. Lear. 1953. Cornell Univ. Agr. Ext. Bull. 870. [39] Miller, L. I. Proc. Helminthol. Soc. Wash. D.C. 20(1):21. [42] Raski, D. J. 1950. Phytopathology 40(2):135. [43] Reynolds, 44(7):389. [45] Sher, S. A., and M. W. Allen. 1953. Univ. Calif. (Berkeley) Publ. Zool. 57(6):441. [46] Staniland, E. M. Buhner. 1932. Plant Disease Repr. 16(8):76. [49] Suit, R. F. 1954. Proc. Florida State Hort. Soc. 67:85. Paper 2. [52] Thorne, G. 1945. Proc. Helminthol. Soc. Wash. D.C. 12(2):27. [53] Thorne, G. 1952. U.S. Dept. [55] Van der Vecht, J. 1953. Contrib. Gen. Agr. Res. Sta. (Bogor) 137. [56] Van der Vecht, J., and B. H. II. 39(1):9.



# Contrails

## 132. VIRAL DISEASES: ANIMALS

Abbreviations: NP = nasopharyngeal; CNS = central nervous system; CSF = cerebrospinal fluid; RBC = red blood cells.

	Virus	Natural Host	Location in Natural Infection	Natural Transmission	Experimental Host	Tissue Culture (Growth in Egg <sup>1</sup> )	Estimated Size <sup>2</sup> μm	Remarks
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Adenovirus	Man, cattle, chicken, dog, monkey, mouse	Respiratory and intestinal tracts	NP secretions	Man (?), dog (?), suckling hamster	Human tumor, monkey kidney, dog and chick embryo	85 (em)	Variable pathogenicity; may agglutinate RBC
2	Bluetongue	Cattle, sheep	Blood, all organs	<i>Culicoides</i> midge	Goat, hamster, suckling mouse	Sheep kidney (+)	100-150 (mf)	Several antigenic types
3	Chicken pox (varicella)	Man	Fluid and crusts of cutaneous lesions	Air-borne contact		Human foreskin and embryo	210-243 (em)	Antigenically identical to herpes zoster
4	Cowpox	Man, cattle	Cutaneous lesions on teat and udder	Contact with discharge from lesion, hands of milker	Rabbit, guinea pig	Cattle fetus, skin (+)		Antigenically related to vaccinia
5	Coxsackie	Man	Feces, blood, pharynx, saliva, CNS	Ingestion	Suckling mouse; hamster, monkey, chimpanzee	Newborn mouse; human, chick, monkey, and mouse embryo (+)	37 (em)	Thirty or more antigenic types
6	Dengue	Man, mosquito	Blood	<i>Aedes</i> mosquito	Mouse	Monkey kidney (+)	17-25 (mf)	Four antigenic types
7	Distemper, canine	Dog, ferret, other carnivores	Blood, secretions, excretions	Contact with secretions, excretions	Ferret, hamster, suckling mouse	Dog kidney (+)	20-22 (em)	Classical, neurotrophic, hard-pad types
8	Encephalitis, western equine	Man, horse, domestic and wild birds	CNS, blood, spleen	<i>Culex</i> mosquito	Mouse, many domestic and wild animals	Human and chick embryo, hamster and monkey kidney (+)	25 (mf); 40 (c, em)	Antigenically distinct; Arbor group A
9	Foot-and-mouth disease	Man, cattle, goat, sheep, swine	Blood, saliva, milk, urine	Contact with secretions, excretions; raw garbage	Rabbit, guinea pig, young dog, mouse, rat, chick	Guinea pig and cattle epithelium (+)	22 (em)	Three or more antigenic types; agglutinates RBC
10	Fowl plague	Domestic fowl, some wild birds	Blood, secretions, excretions	Ingestion	Starling, canary, mouse, rat, rabbit, ferret	Chick embryo (+)	60-140 (mf, c, em)	Antigenic variants; agglutinates RBC
11	Herpes simplex	Man	Skin, cornea, blood, mucosa, CNS	Contact ?	Rabbit, mouse	Rabbit and chick embryo (+)	100-150 (mf)	Antigenically related to pseudorabies, B-viruses
12	Hog cholera	Swine	Blood, all secretions, excretions	Contact with secretions, excretions, raw garbage	Rabbit, guinea pig	Swine marrow, testis, spleen, kidney, white blood cells	27 (em)	Antigenic, neurotrophic variants
13	Infectious hepatitis	Man	Blood, feces	Ingestion	Man ?	HeLa and other human cells ?	Passes Seitz	Two antigenic types
14	Influenza	Man	Respiratory tract	Contact with NP secretions	Ferret, mouse	Chick embryo (+)	80-120 (mf, c, em)	Antigenic types A-C; agglutinates RBC
15	Louping ill	Man, cattle, sheep	CNS, blood; CSF (man)	Tick; contact with infected animals (man)	Mouse, vole, swine, cattle, monkey	Chick embryo (+)	15-20 (mf); 22-27 (c)	Antigenically related to Russian Far East virus
16	Lymphogranuloma venereum	Man	Genital lesion, CSF, inguinal lymph node	Direct contact with lesion or exudate	Mouse, guinea pig, monkey	Chick and mouse embryo; guinea pig and mouse testis (+)	300 (dm); 438 (em)	Antigenically related to psittacosis; produces toxin

<sup>1</sup>/ Growth in embryonated chicken egg indicated by plus sign in parentheses. <sup>2</sup>/ Method of determination given in parentheses: (c) = centrifugation; (dm) = direct microscopy; (em) = electron microscopy; (mf) = membrane filtration.

continued

## 132. VIRAL DISEASES: ANIMALS

Virus	Natural Host	Location in Natural Infection	Natural Transmission	Experimental Host	Tissue Culture (Growth in Egg <sup>1</sup> )	Estimated Size <sup>2</sup> m $\mu$	Remarks
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
17 Measles (rubella)	Man	Blood, skin, respiratory tract, urine	Contact with NP secretions	Monkey	Chick embryo, human and monkey kidney (+)	120-150 (em)	
18 Measles, German (rubella)	Man	Respiratory tract, blood, urine	Contact with NP secretions	Man ?	Human amnion, monkey kidney	<800 (mf)	May produce fetal malformation
19 Molluscum contagiosum	Man	Skin	Direct contact		HeLa cells	190-250 (em)	
20 Mumps	Man	Salivary glands, blood, gonads, CSF	Droplets of saliva	Monkey, hamster	Chick and mouse embryo (+)	90-135 (mf); 140-268 (em)	Agglutinates RBC; produces hemolysin
21 Newcastle disease	Man, cattle, domestic fowl, some wild birds	Conjunctival and nasal secretions (man); lungs (calf); blood, secretions, excretions (ovipara)	Contact with secretions, excretions, sick animals; raw garbage (poultry scraps)	Common laboratory animals, cattle, quail, sparrow	Chick embryo, chicken trachea cells, swine embryo lymph node (+)	70-180 (em)	Neural pneumonic visceral strains; agglutinates RBC
22 Parainfluenza	Man, cattle	Respiratory tract	Contact with NP secretion		Monkey and cattle kidney	80-150 (em)	Four or more antigenic types
23 Poliomyelitis	Man	CNS, intestinal tract, blood	Ingestion	Chimpanzee, monkey, hamster, mouse	Human, monkey	28 (em)	Three antigenic types
24 Polyoma	Mouse	Blood, tissues, excretions	Contact	Suckling mouse, hamster, and rat	Mouse embryo	45 (em)	Tumorigenic for newborn mice; agglutinates RBC
25 Psittacosis	Man, psittacine birds, duck, chicken	Lung, spleen, liver, sputum (man); respiratory secretions, cloacal contents	Inhalation of dried secretions, droppings, contact with infected tissues	Mouse, guinea pig, monkey	Chick embryo, mouse (+)	455 (em)	Produces toxin
26 Rabies	All mammals	CNS, salivary and lacrimal glands, kidney, pancreas, saliva	Through broken epithelium contaminated with saliva	Mammals, domestic fowl	Rabbit, rat, mouse, chick embryo (+)	100-150 (mf)	
27 Reovirus	Man, cattle, monkey, mouse	Intestinal tract	Ingestion ?	Man ?	Human, monkey	60-90 (em)	Three antigenic types; agglutinates RBC
28 Rift Valley fever	Man, sheep	Blood, liver	Mosquito ?	Cattle, goat, monkey, mouse	Chick embryo (+)	23-35 (mf)	
29 Rinderpest	Domestic animals, deer	Blood, secretions, excretions	Ingestion	Rabbit, guinea pig, Chinese pig	Chick embryo (+)	Passes V Berkefeld	
30 Rous sarcoma	Chicken	Tumor, blood	Contact, egg	Chick, turkey	Chick embryo (+)	65-90 (em)	One of several related tumorigenic chicken viruses
31 Shope papilloma	Rabbit	Skin		Domestic rabbit, hamster		22-35 (mf); 32-50 (c)	Malignancy becomes carcinomatous
32 Silkworm jaundice	<i>Bombyx mori</i>	Fat, hypodermis, blood cells; other tissues ?	Oral; egg ?	Gypsy moth (?), nun moth ?		40-288 (em)	Inclusion bodies: polyhedral

<sup>1</sup>/ Growth in embryonated chicken egg indicated by plus sign in parentheses. <sup>2</sup>/ Method of determination given in parentheses: (c) = centrifugation; (dm) = direct microscopy; (em) = electron microscopy; (mf) = membrane filtration.

continued

**132. VIRAL DISEASES: ANIMALS**

Virus	Natural Host	Location in Natural Infection	Natural Transmission	Experimental Host	Tissue Culture (Growth in Egg <sup>1</sup> )	Estimated Size <sup>2</sup> m $\mu$	Remarks
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
33 Smallpox (variola)	Man; monkey ?	Cutaneous lesions, NP secretions, mucosa, blood	Inhalation of secretions, scabs	Monkey, rabbit	Human embryo, skin, kidney, tumor, etc. (+)	244-302 (em)	Antigenically related to vaccinia; agglutinates RBC
34 Trachoma	Man	Conjunctiva, conjunctival exudate	Contact with conjunctival exudate	Ape; man ?	(+)	250 (dm)	
35 Vaccinia	Man (?), rabbit (?), cattle ?	Cutaneous lesions ?		Man, cattle, monkey, rabbit, mouse	Rabbit, chicken; guinea pig adult and embryo (+)	236-252 (mf)	Antigenically related to smallpox; agglutinates RBC
36 Wart (verruca)	Man	Cutaneous lesions	Contact ?	Man ?		Passes N Berkefeld	
37 Yellow fever	Man, monkey	Blood, liver, CNS, spleen, lymph nodes	<i>Aedes aegypti</i> and other culicines	Mouse, monkey, hedgehog	Chick and mouse embryo (+)	12-19 (c); 22 (mf)	Arbor group B

<sup>1/1</sup> Growth in embryonated chicken egg indicated by plus sign in parentheses. <sup>2/2</sup> Method of determination given in parentheses: (c)= centrifugation; (dm) = direct microscopy; (em) = electron microscopy; (mf) = membrane filtration.

Contributors: (a) Thompson, Randall L., (b) Moses, Harold E., (c) Minton, Sherman A., Jr.

References: [1] Andrewes, C. H. 1962. *Advan. Virus Res.* 9:271. [2] Bass, E. P., and J. D. Ray. 1963. *J. Am. Vet. Med. Assoc.* 142:1112. [3] Burnet, F. M. 1960. *Principles of animal virology*. Ed. 2. Academic Press, New York. [4] Burnet, F. M., and W. M. Stanley. 1959. *The viruses*. Academic Press, New York. v. 1. [5] Hagan, W. A., and D. W. Bruner. 1961. *The infectious diseases of domestic animals*. Ed. 4. Comstock, Ithaca. [6] Johnstone, M. C., and H. Koprowski, ed. 1962. *Ann. N. Y. Acad. Sci.* 101:327. [7] Loan, R. L., and D. P. Gustafson. 1961. *Am. J. Vet. Res.* 22:741. [8] Merchant, I. A., and R. A. Packer. 1961. *Veterinary bacteriology and virology*. Ed. 6. Iowa State Univ. Press, Ames. [9] Miner, R. W. 1953. *Ann. N. Y. Acad. Sci.* 56:381. [10] Morehouse, L. G., D. P. Gustafson, and H. E. Moses. 1963. *Am. J. Vet. Res.* 24:588. [11] Morehouse, L. G., H. E. Moses, and D. P. Gustafson. 1963. *Ibid.* 24:580. [12] Rhodes, A. J., and C. E. van Rooyen. 1962. *Textbook of virology*. Ed. 4. Williams and Wilkins, Baltimore. [13] Rivers, T. M., and F. L. Horsfall, Jr. 1959. *Viral and rickettsial infections of man*. Ed. 3. J. B. Lippincott, Philadelphia. [14] Sanders, M., I. Kiem, and D. Lagunoff. 1953. *Arch. Pathol.* 56:143.

**133. VIRAL DISEASES: PLANTS**

Host Plant and Disease	Distribution	Principal Insect Vector	Other Means of Transmission	Symptoms
(A)	(B)	(C)	(D)	(E)
1 <i>Allium cepa</i> (garden onion) Onion yellow dwarf	United States, Germany, New Zealand, USSR	Aphids (including <i>Aphis maidis</i> , <i>A. rumicis</i> )	Leaf rubbing	Leaves yellowed, crinkled; plants dwarfed; bulbs small; few seeds
2 <i>Beta vulgaris</i> (beet) Beet yellows	Belgium, Netherlands, Denmark, England	Aphids (including <i>Aphis fabae</i> , <i>Myzus persicae</i> )	.....	Leaves yellow, thick, brittle; necrosis in secondary phloem
3 Sugar beet curly top	Western North America	Leafhopper ( <i>Circulifer tenellus</i> )	Grafting; dodder	Leaves curled, enations on veins; plant stunted; many rootlets

*continued*

## 133. VIRAL DISEASES: PLANTS

	Host Plant and Disease	Distribution	Principal Insect Vector	Other Means of Transmission	Symptoms
	(A)	(B)	(C)	(D)	(E)
4	<i>Cucumis sativus</i> (cucumber) Cucumber mosaic	Almost world-wide	Aphids (including <i>Aphis gossypii</i> , <i>Myzus persicae</i> )	Leaf rubbing; dodder	Leaves mottled, distorted, small; plant stunted; fruits mottled
5	Cucurbit mosaic	England	.....	Leaf rubbing	Chlorotic mottling and distortion of foliage; plant stunted
6	<i>Fragaria hybrida</i> (strawberry) Strawberry stunt	United States	Aphid ( <i>Capitophorus fragaeifolii</i> )	Grafting	Leaves green, luster dull; leaflets cupped; plant stunted; fruits small
7	<i>Gossypium hirsutum</i> (upland cotton) Cotton leaf curl	Sudan, Nigeria	Whitefly ( <i>Bemisia gossypiperda</i> )	Grafting	Leaves pale-spotted, puckered, unsymmetrical; internodes shortened
8	<i>Lactuca sativa</i> (lettuce) Lettuce mosaic	Worldwide	Aphids ( <i>Macrosiphum gei</i> , <i>Myzus persicae</i> )	Leaf abrasion; seeds	Clearing of veins, followed by systemic chlorotic mottling
9	<i>Lycopersicon esculentum</i> (tomato) Tomato bushy stunt	British Isles	.....	Leaf rubbing; dodder	Foliage yellowed; plant stunted; axillary buds stimulated
10	Tomato spotted wilt	Almost world-wide	<i>Frankliniella moultoni</i> , <i>F. schultzei</i> , <i>Thrips tabaci</i>	Leaf abrasion	Bronze-ring lesions, necrosis or mottling; fruit discolored
11	<i>Medicago sativa</i> (alfalfa) Alfalfa mosaic	United States	Aphids ( <i>Macrosiphum pisi</i> , <i>M. solanifolii</i> )	Leaf rubbing	Systemic chlorotic mottling, often masked
12	<i>Nicotiana tabacum</i> (common tobacco) Tobacco etch	United States	Aphids (especially <i>Myzus persicae</i> )	Leaf rubbing	Systemic chlorotic mottling with traces of whitish etching
13	Tobacco leaf curl	Africa, India, Sumatra, Formosa	Whitefly ( <i>Bemisia gossypiperda</i> )	Grafting	Leaves curled, wrinkled; veins thick, enations; plant stunted
14	Tobacco mosaic	Worldwide	.....	Leaf rubbing; dodder; soil	Systemic chlorotic mottling; some distortion of leaves
15	Tobacco necrosis	United States, British Isles, Australia	.....	Leaf rubbing; soil; fungus vector ( <i>Olpidium</i> spp.)	Necrosis in midrib and veins of lower leaves in winter
16	Tobacco ring spot	United States	.....	Leaf rubbing; seeds; nematode vector ( <i>Xiphinema americanum</i> )	Necrotic, ringlike primary and secondary lesions; later recovery
17	<i>Oryza sativa</i> (rice) Rice dwarf	Japan, Philippine Islands	Leafhoppers (including <i>Nephotettix apicalis</i> )	.....	Leaves chlorotic, spotted, streaked; internodes and roots short
18	<i>Phaseolus vulgaris</i> (kidney bean) Bean mosaic	Almost world-wide	Aphids (including <i>Aphis rumicis</i> )	Leaf abrasion; seeds	Systemic chlorotic mottling
19	Southern bean mosaic	Southern United States	.....	Leaf rubbing; seeds	Mottling in some varieties, localized or systemic necrosis in others

continued



## 133. VIRAL DISEASES: PLANTS

	Host Plant and Disease	Distribution	Principal Insect Vector	Other Means of Transmission	Symptoms
	(A)	(B)	(C)	(D)	(E)
20	<i>Prunus persica</i> (peach) Peach phony disease	Southeastern United States	Leafhoppers (including <i>Homalodisca triquetra</i> )	Root grafting	Foliage abnormally green; tree dwarfed, fruit small
21	Peach rosette	United States	.....	Budding; dodder	Stems short with dwarfed leaves; veins thickened; tree dies soon
22	Peach X-disease	United States	Leafhoppers (including <i>Colladonus clitellarius</i> )	Budding	Leaves light green, tattered; old leaves drop; fruit bitter
23	Peach yellows	Eastern United States, eastern Canada	Leafhopper ( <i>Macropsis trimaculata</i> )	Budding	Leaves chlorotic; shoots erect, thin, numerous; tree dies soon
24	<i>Solanum tuberosum</i> (potato) Potato aucuba mosaic	United States, Europe	Probably aphid ( <i>Myzus persicae</i> )	Leaf rubbing	Yellow mottling of foliage; some necrosis in tubers
25	Potato leaf roll	Wherever potatoes are grown	Aphids (especially <i>Myzus persicae</i> )	Grafting	Leaves thick, leathery, rolled, starchy; plant small; few tubers
26	Potato mild mosaic	United States, England, Holland	Aphids ( <i>Aphis abbreviata</i> , <i>Myzus persicae</i> )	Leaf rubbing	Mild chlorotic mottling or masked symptoms in most varieties
27	Potato mottle	Worldwide	.....	Leaf rubbing; root and leaf contacts	No obvious disease, or very mild chlorotic mottling
28	Potato spindle tuber	United States, Canada	Aphids ( <i>Macrosiphum solanifolii</i> , <i>Myzus persicae</i> )	Leaf rubbing; seed-piece cutting	Leaves small, erect, dark green; plant brittle; tubers tapered
29	Potato veinbanding	United States, Brazil, England, France	Aphids (especially <i>Myzus persicae</i> )	Leaf rubbing	Chlorotic mottling, necrotic stemstreak, or no obvious disease
30	Potato witches'-broom	United States, USSR	.....	Tuber core graft, stem graft	Leaves small, pale; branches numerous, spindly; tubers small
31	Potato yellow dwarf	United States, Canada	Leafhopper ( <i>Acerata-gallia sanguinolenta</i> )	Leaf abrasion in some hosts; grafting	Leaves yellowed, plant dwarfed; tubers few, small, often cracked
32	<i>Trifolium incarnatum</i> (crimson clover) Clover club leaf	United States (New Jersey)	Leafhopper ( <i>Agalliopsis novella</i> )	.....	Plant dwarfed; leaves small, yellowed or reddened at margins
33	Wound tumor	United States	Leafhoppers ( <i>Agallia constricta</i> , <i>Agalliopsis novella</i> )	.....	Experimentally: veins thickened, enations; plant dwarfed
34	<i>Triticum aestivum</i> (wheat) Wheat mosaic	United States, Japan	.....	Leaf abrasion; soil	Systemic chlorotic mottling; dwarfing; vacuolate inclusions
35	Wheat streak mosaic	United States, Canada	Mite ( <i>Aceria tulipae</i> )	Leaf abrasion	Systemic chlorotic mottling, streaking of leaves
36	Winter-wheat mosaic	USSR	Leafhopper ( <i>Deltoccephalus striatus</i> )	.....	Chlorotic mottling; phloem necrosis, vacuolate inclusions
37	<i>Zea mays</i> (corn) Maize streak	Africa	Leafhoppers ( <i>Cicadulina mbila</i> , <i>C. storeyi</i> , <i>C. zeae</i> )	.....	Chlorotic spotting, streaking of leaves

Contributor: Holmes, Francis O.

References: [1] Bawden, F. C. 1950. Plant viruses and virus diseases. Ed. 3. Chronica Botanica, Waltham.

continued

## 133. VIRAL DISEASES: PLANTS

Mass. [2] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. Bergey's Manual of determinative bacteriology. Ed. 7. Williams and Wilkins, Baltimore. p. 985. [3] Cook, M. T. 1947. Viruses and virus diseases of plants. Burgess, Minneapolis. p. 985. [4] Smith, K. M. 1957. A textbook of plant virus diseases. Little and Brown, Boston. [5] U.S. Agricultural Research Service, Crops Research Division. 1960. U.S. Dept. Agr. Handbook 165.

## 134. RICKETTSIAL PARASITES: MAMMALS AND BIRDS

Host (column B): Animals are listed in order of decreasing susceptibility.

	Species (A)	Host (B)	Disease or Disorder (C)	Method of Transmission (D)
1	<i>Anaplasma centrale</i>	Cattle	Benign anaplasmosis	Tick to host
2	<i>A. marginale</i>	Cattle	Malignant anaplasmosis	Tick to host
3	<i>A. ovis</i>	Sheep, goat	Ovine anaplasmosis	Tick to host
4	<i>Bartonella bacilliformis</i>	Man	Oroya fever	Sand fly to man
5	<i>Chlamydia oculogenitalis</i>	Man	Inclusion blennorrhoea, inclusion conjunctivitis, neonatal conjunctivitis	Contact; contaminated swimming pools
6	<i>C. trachomatis</i>	Man, ape, monkey	Trachoma	Contact
7	<i>Colesiotea conjunctivae</i>	Sheep, cattle, goat	Ophthalmia	Uncertain
8	<i>Colletsia pecoris</i>	Cattle, sheep, goat	Conjunctivitis ?	Uncertain
9	<i>Coudria ruminantium</i>	Goat, sheep, cattle	Heartwater	Tick feces; tick to host
10	<i>Coxiella burneti</i>	Man, cattle, sheep, goat	Q fever	Tick feces; host to tick to host; contact with domestic animals; inhalation of infected dust; contaminated milk
11	<i>Ehrlichia bovis</i>	Cattle	Bovine rickettsiosis	Tick to host
12	<i>E. canis</i>	Dog	Canine rickettsiosis	Tick to host
13	<i>E. ovina</i>	Sheep	Ovine rickettsiosis	Presumed tick to host
14	<i>E. phagocytophila</i>	Sheep	Ovine rickettsiosis	Tick to host
15	<i>Eperythrozoon</i> spp.	Cattle, mouse, sheep, swine	Eperythrozoonosis, anemia	Mouse louse to mouse (other arthropods suspected)
16	<i>Grahamella talpae</i>	Mole	Grahamellosis, anemia	Uncertain
17	<i>Haemobartonella</i> spp.	Animals	Haemobartonellosis, anemia	Flea to host
18	<i>Miyagawanella bronchopneumoniae</i>	Mouse	Mouse pneumonitis	Contact
19	<i>M. felis</i>	Cat	Feline pneumonitis	Contact
20	<i>M. lymphogranulomatosis</i>	Man	Lymphogranuloma venereum (lymphogranuloma inguinale)	Venereal contact
21	<i>M. ornithosis</i>	Man, nonpsittacine birds	Ornithosis	Inhalation of infected dust; bird to man, man to man
22	<i>M. psittaci</i>	Man, psittacine birds	Psittacosis	Inhalation of infected dust; bird to man, man to man
23	<i>Neorickettsia helminthoeca</i>	Dog, fox, coyote	Salmon poisoning	Intestinal parasitic fluke to dog
24	<i>Rickettsia akari</i>	Man	Rickettsial pox	Mite to man
25	<i>R. conorii</i>	Man; dog as reservoir	Boutonneuse (Marseilles, Mediterranean) fever	Tick to man
26	<i>R. prowazekii</i>	Man	Epidemic (classic Old World) typhus	Louse feces; man to louse to man
27	<i>R. rickettsii</i>	Man, rabbit, squirrel	Rocky Mountain spotted fever	Tick to man
28	<i>R. siberica</i>	Man, rodents	Siberian tick typhus	Tick to host
29	<i>R. tsutsugamushi</i>	Man, monkey, rodents	Tsutsugamushi fever (scrub typhus)	Mite to man
30	<i>R. typhi</i>	Man, rodents	Murine or endemic typhus	Flea feces; rat to flea to man
31	<i>Ricolesia bovis</i>	Cattle	Infectious conjunctivitis	Contact
32	<i>R. caprae</i>	Goat	Infectious conjunctivitis	Possibly contact
33	<i>R. conjunctivae</i>	Fowl	One form of ocular roup	Possibly contact
34	<i>R. lestoquardii</i>	Swine	Infectious conjunctivitis	Possibly contact
35	<i>Rochalimaea quintana</i>	Man	Trench fever	Body louse to man

Contributor: Philip, Cornelius B.

continued

**134. RICKETTSIAL PARASITES: MAMMALS AND BIRDS**

*References:* [1] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. *Bergey's Manual of determinative bacteriology*. Ed. 7. Williams and Wilkins, Baltimore. [2] Burrows, W. 1963. *Textbook of microbiology*. Ed. 18. W. B. Saunders, Philadelphia. [3] Hagan, W. A., and D. W. Bruner. 1961. *The infectious diseases of domestic animals*. Ed. 4. Comstock, Ithaca. [4] Hull, T. G. 1955. *Diseases transmitted from animals to man*. Ed. 4. C. C. Thomas, Springfield, Ill. [5] Merchant, I. A., and R. A. Packer. 1961. *Veterinary bacteriology and virology*. Ed. 6. Iowa State Univ. Press, Ames. [6] Moulton, F. R., ed. 1948. *Rickettsial diseases of man*. Symposium. American Association for the Advancement of Science, Washington, D. C. [7] Philip, C. B. 1950. In R. L. Pullen, ed. *Communicable diseases*. Lea and Febiger, Philadelphia. p. 781. [8] Rivers, T. M., and F. L. Horsfall, Jr., ed. 1959. *Viral and rickettsial infections of man*. Ed. 3. J. B. Lippincott, Philadelphia.

**135. BACTERIAL PARASITES: MAMMALS AND BIRDS**

Host (column B): Animals are listed in order of decreasing susceptibility.

Species (A)	Host (B)	Disease or Disorder (C)	Method of Transmission (D)
1 <i>Actinobacillus lignieresii</i>	Cattle, swine	Actinobacillosis (wooden tongue)	Not definitely known
2 <i>A. mallei</i>	Horse, man	Glanders	Contact; contaminated feed, water
3 <i>Actinomyces bovis</i>	Cattle, swine, horse, man	Actinomycosis (lumpy jaw)	Buccal cavity; transmission not definitely known
4 <i>Bacillus anthracis</i>	Man	Anthrax	Soil-borne; contact with infected animal by-products, carcasses
5	Cattle, sheep, horse, mule, swine	Anthrax (splenic fever)	Soil-borne; infected feed, water, carcasses
6 <i>Bordetella pertussis</i>	Man	Whooping cough	Flora of respiratory tract; droplet infection
7 <i>Borrelia anserina</i>	Fowl	Spirochetosis	Arthropod vector; feces-borne
8 <i>B. recurrentis</i>	Man	European relapsing fever	Arthropod vector
9 <i>B. theileri</i>	Cattle	Spirochetosis	Arthropod vector
10 <i>B. vincentii</i>	Man	Associated with <i>Fusobacterium fusiforme</i> in Vincent's angina	Buccal cavity; transmission not definitely known
11 <i>Brucella abortus</i>	Cattle, man	Brucellosis (undulant fever)	Ingestion of infected milk; contact
12 <i>B. melitensis</i>	Goat, sheep, man	Brucellosis (undulant fever)	Ingestion of infected milk; contact
13 <i>B. suis</i>	Swine, man	Brucellosis (undulant fever)	Contact
14 <i>Clostridium botulinum</i>	Man, chicken, duck, horse, mule, cattle	Botulism, food intoxication, limberneck of fowl	Ingestion of toxin in food
15 <i>C. chauvoei</i>	Cattle, man	Blackleg, symptomatic anthrax	Soil-borne; wound infection
16 <i>C. haemolyticum</i>	Cattle, sheep	Icterohemoglobinuria	Soil-borne; contaminated feed, water
17 <i>C. novyi</i>	Sheep, man	Infectious necrotic hepatitis	Soil-borne; associated with liver fluke infection
18 <i>C. perfringens</i>	Man	Gas gangrene	Soil-borne; wound infection
19	Sheep	Dysentery in lamb, infectious enterotoxemia	Soil-borne; wound infection
20 <i>C. septicum</i>	Horse, sheep, cattle, man	Malignant edema	Soil-borne; wound infection
21 <i>C. tetani</i>	Man, sheep, cattle, goat, swine, horse	Tetanus (lockjaw)	Soil-borne; wound infection
22 <i>Corynebacterium diphtheriae</i>	Man	Diphtheria	Carrier contact
23 <i>C. equi</i>	Horse	Pneumonia of foal	Possible in utero
24	Swine	Submaxillary gland infection	Contaminated soil, feed
25 <i>C. pseudotuberculosis</i>	Sheep, goat	Caseous lymphadenitis	Contaminated feed, water
26	Horse, deer, elk, moose, mountain sheep	Ulcerative lymphangitis (pseudoglanders)	Contact; wound infection
27 <i>C. pyogenes</i>	Cattle, swine, sheep, goat, deer	Mastitis, purulent infections, arthritis	Inhabits mucous membranes; contact; wound infection

*continued*

## 135. BACTERIAL PARASITES: MAMMALS AND BIRDS

Species (A)	Host (B)	Disease or Disorder (C)	Method of Transmission (D)
28 <i>Corynebacterium renale</i>	Cattle, swine	Pyelonephritis	Inhabits mucous membranes; transmission not definitely known
29 <i>Diplococcus pneumoniae</i>	Man	Lobar pneumonia, meningitis, endocarditis	Carrier contact
30 <i>Erysipelothrix insidiosa</i>	Man	Erysipeloid	Wound infection; contact with infected carcasses
31	Swine, sheep, fowl	Erysipelas	Feces-borne; contaminated soil, feed, water
32 <i>Escherichia coli</i>	Man, domestic animals	Genitourinary and intestinal infections	Feces-borne; normal flora of intestinal tract.
33 <i>Fusobacterium fusiforme</i>	Man	Associated with ulcerative stomatitis (Vincent's angina)	Buccal cavity; transmission not definitely known
34 <i>Haemophilus ducreyi</i>	Man	Soft chancre, chancroid	Direct genital contact
35 <i>H. gallinarum</i>	Chicken	Infectious coryza	Contact
36 <i>H. haemoglobinophilus</i>	Dog	Prepuccial infection	Direct sexual contact
37 <i>H. influenzae</i>	Man	Meningitis, obstructive respiratory infections	Flora of respiratory tract; droplet infection
38 <i>H. suis</i>	Swine	Associated with viral influenza	Contact; flora of respiratory tract
39 <i>Klebsiella pneumoniae</i>	Man, horse, cattle	Respiratory infection; mastitis (cattle)	Contact; flora of respiratory tract
40 <i>Leptospira canicola</i>	Dog, man, swine, cattle	Leptospirosis (Stuttgart's disease of dog)	Direct contact; contamination of water by infected urine
41 <i>L. icterohaemorrhagiae</i>	Man, dog, swine, cattle, rodents	Leptospirosis (Weil's disease)	Direct contact; contamination of water by infected urine
42 <i>L. pomona</i>	Cattle, swine, man	Leptospirosis (swineherd's disease)	Direct contact; contamination of water by infected urine
43 <i>Listeria monocytogenes</i>	Man, domestic animals, fowl	Listeriosis, meningoenzephalitis, abortion	Not definitely known
44 <i>Moraxella bovis</i>	Cattle	Infectious keratitis	Contact
45 <i>M. lacunata</i>	Man	Conjunctivitis	Not definitely known
46 <i>Mycobacterium avium</i>	Fowl, swine	Tuberculosis	Contact; feces-borne; contaminated feed, water
47 <i>M. bovis</i>	Cattle, man, swine	Tuberculosis	Contact; feces-borne; contaminated feed, water, milk
48 <i>M. leprae</i>	Man	Hansen's disease (leprosy)	Not definitely known
49 <i>M. paratuberculosis</i>	Cattle, sheep	Johne's disease	Contact; feces-borne; contaminated feed, water
50 <i>M. tuberculosis</i>	Man	Tuberculosis	Contact; via respiratory or alimentary tract
51 <i>Neisseria catarrhalis</i>	Man	Catarrhal inflammations	Flora of respiratory tract
52 <i>N. gonorrhoeae</i>	Man	Gonorrhea	Direct sexual contact
53 <i>N. meningitidis</i>	Man	Epidemic cerebrospinal meningitis	Infection from respiratory tract of carrier
54 <i>Pasteurella multocida</i>	Domestic animals, fowl, man	Hemorrhagic septicemia, bronchiectasis, conjunctivitis	Contact; contaminated feed, water; bites
55 <i>P. pestis</i>	Man, rodents	Bubonic and pneumonic plague	Flea bite from infected rat; droplet infection
56 <i>P. pseudotuberculosis</i>	Guinea pig, rabbit, horse, cattle, goat, rodents, dog, cat, monkey, fowl	Pseudotuberculosis	Soil-borne; contaminated water, fodder, milk
57 <i>P. tularensis</i>	Man, rodents, lagomorphs	Tularemia	Contact with contaminated carcasses; insect vector
58 <i>Proteus vulgaris</i>	Man	Genitourinary and intestinal infections	Feces-borne
59 <i>Pseudomonas aeruginosa</i>	Man, animals	Suppurative processes, septicemia, meningitis, genitourinary infections	Contaminated water, soil, feces; wound infection
60 <i>Salmonella enteritidis</i>	Man, rodents	Food poisoning, gastroenteritis	Feces-borne
61 <i>S. gallinarum</i>	Fowl	Fowl typhoid, bacillary white diarrhea	Feces-borne; ovarian transmission
62 <i>S. hirschfeldii</i>	Man	Paratyphoid fever	Feces-borne
63 <i>S. paratyphi A</i>	Man	Paratyphoid fever	Feces-borne
64 <i>S. paratyphi B</i>	Man	Paratyphoid fever	Feces-borne
65 <i>S. typhimurium</i> <sup>1</sup>	Man, rodents	Food poisoning, gastroenteritis	Feces-borne

/1/ A natural pathogen for all warm-blooded animals

continued



## 135. BACTERIAL PARASITES: MAMMALS AND BIRDS

	Species (A)	Host (B)	Disease or Disorder (C)	Method of Transmission (D)
66	<i>Salmonella typhosa</i>	Man	Typhoid fever	Feces-borne
67	<i>Shigella dysenteriae</i>	Man	Dysentery	Feces-borne; contaminated food, water
68	<i>S. equirulis</i>	Horse	Joint infection, nephritis	Possible in utero
69	<i>Sphaerophorus necrophorus</i>	Man ?	Associated with ulcerative colitis ?	Feces-borne; associated with unsanitary conditions
70		Horse	Gangrenous dermatitis (scratches)	Feces-borne; associated with unsanitary conditions
71		Cattle	Calf diphtheria	Feces-borne; associated with unsanitary conditions
72		Sheep, goat	Lip-and-leg ulceration, ulcerative stomatitis, foot rot	Feces-borne; associated with unsanitary conditions
73		Swine	Ulcerative stomatitis, enteritis	Feces-borne; associated with unsanitary conditions
74	<i>Spirillum minus</i>	Man, rodents	Rat-bite fever	Rat bite
75	<i>Staphylococcus aureus</i>	Man, animals	Suppurative processes, food poisoning, septicemia	Wound infection; contaminated food; flora of skin and mucous membranes
76	<i>Streptococcus agalactiae</i>	Cattle	Mastitis	Contaminated milking equipment
77	<i>S. dysgalactiae</i>	Cattle	Mastitis	Contaminated milking equipment
78	<i>S. equi</i>	Horse	Strangles	Contact; contaminated water, feed
79	<i>S. pyogenes</i>	Man, vole	Scarlet fever, septic sore throat	Direct personal contact; contaminated milk
80	<i>S. uberis</i>	Cattle	Mastitis	Contaminated milking equipment
81	<i>Treponema pallidum</i>	Man	Syphilis	Direct sexual contact
82	<i>Vibrio comma</i>	Man	Cholera	Feces-borne from carrier; contaminated food, water
83	<i>V. fetus</i>	Cattle, sheep	Vibrionic abortion, sterility	Direct sexual contact
84	<i>V. jejuni</i>	Cattle	Dysentery	Feces-borne; contaminated feed and water

Contributor: Cunningham, Charles H.

References: [1] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. Bergey's Manual of determinative bacteriology. Ed. 7. Williams and Wilkins, Baltimore. [2] Burrows, W. 1963. Textbook of microbiology. Ed. 18. W. B. Saunders, Philadelphia. [3] Dubos, R. J. 1958. Bacterial and mycotic infections of man. Ed. 3. J. B. Lippincott, Philadelphia. [4] Hagan, W. A., and D. W. Bruner. 1961. The infectious diseases of domestic animals. Ed. 4. Comstock, Ithaca. [5] Hull, T. G. 1955. Diseases transmitted from animals to man. Ed. 4. C. C. Thomas, Springfield, Ill. [6] Merchant, I. A., and R. A. Packer. 1961. Veterinary bacteriology and virology. Ed. 6. Iowa State Univ. Press, Ames.

## 136. BACTERIAL PARASITES: PLANTS

	Host Plant and Pathogen (A)	Disease (B)		Host Plant and Pathogen (A)	Disease (B)
	<i>Acer</i> spp. (maple)		13	<i>Beta vulgaris</i> (common beet)	
1	<i>Pseudomonas aceris</i>	Leaf spot	14	<i>Erwinia scabiegena</i>	Blister scab
2	<i>Xanthomonas acerrea</i>	Leaf blight	15	<i>Pseudomonas aptata</i>	Blight
	<i>Allium cepa</i> (garden onion)		16	<i>P. wieringae</i>	Ring rot
3	<i>Pseudomonas alliicola</i>	Scale rot		<i>Xanthomonas beticola</i>	Bacterial pocket
4	<i>P. cepacia</i>	Sour skin		<i>Capsicum frutescens</i> (bush red pepper)	
5	<i>P. cichorii</i> , <i>P. marginalis</i>	Soft rot	17	<i>Erwinia carotovora</i>	Soft rot
6	<i>Xanthomonas striiformans</i>	Stripe	18	<i>Pseudomonas solanacearum</i>	Wilt
	<i>Antirrhinum majus</i> (snapdragon)		19	<i>Xanthomonas vesicatoria</i>	Bacterial spot
7	<i>Xanthomonas antirrhini</i>	Leaf spot		<i>Chrysanthemum</i> spp. (chrysanthemum)	
	<i>Avena sativa</i> (common oat)		20	<i>Erwinia chrysanthemi</i>	Bacterial blight
8	<i>Pseudomonas coronafaciens</i>	Halo blight	21	<i>Pseudomonas cichorii</i>	Leaf spot
9	<i>P. coronafaciens atropurpurea</i>	Purple spot		<i>Citrus</i> spp. (citrus)	
10	<i>P. striafaciens</i>	Stripe blight	22	<i>Erwinia citrimaculans</i>	Fruit spot
11	<i>Xanthomonas translucens</i>	Blight	23	<i>Pseudomonas syringae</i>	Blast and black pit
	<i>Beta vulgaris</i> (common beet)		24	<i>Xanthomonas citri</i>	Canker
12	<i>Corynebacterium betae</i>	Silvering disease			

continued

## 136. BACTERIAL PARASITES: PLANTS

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)	(B)		(A)	(B)	
25	<i>Cucumis sativus</i> (cucumber)		70	<i>Medicago sativa</i> (alfalfa)	
	<i>Erwinia tracheiphila</i>	Wilt		<i>Pseudomonas medicaginis</i>	Stem blight
26	<i>Pseudomonas lachrymans</i>	Angular leaf spot	71	<i>Xanthomonas alfalfae</i>	Leaf spot
27	<i>Xanthomonas cucurbitae</i>	Leaf spot		<i>Nicotiana</i> spp. (tobacco)	
	<i>Cucurbita pepo</i> (pumpkin)		72	<i>Erwinia aroideae</i>	Hollow stalk and barn rot
28	<i>Erwinia tracheiphila</i>	Wilt			
	<i>Daucus carota</i> (carrot)		73	<i>Pseudomonas angulata</i>	Angular leaf spot
29	<i>Erwinia carotovora</i>	Soft rot	74	<i>P. mellea</i>	Rust
30	<i>Xanthomonas carotae</i>	Leaf blight	75	<i>P. polycolor</i>	Leaf spot and wet rot
	<i>Fragaria virginiana</i> (Virginia strawberry)		76	<i>P. pseudozoogloeae</i>	Black rust
31	<i>Corynebacterium fuscians</i>	Cauliflower disease	77	<i>P. solanacearum</i>	Granville disease
32	<i>Xanthomonas fragariae</i>	Angular leaf spot	78	<i>P. tabaci</i>	Wildfire
	<i>Fraxinus</i> spp. (ash)		79	<i>Xanthomonas heterocea</i>	Rust
33	<i>Pseudomonas savastanoi</i>	Canker		<i>Oryza sativa</i> (rice)	
	<i>fraxini</i>		80	<i>Pseudomonas oryzaicola</i>	Leaf spot
	<i>Gladiolus</i> spp. (gladiolus)		81	<i>P. setariae</i>	Stripe
34	<i>Pseudomonas gladioli</i>	Rot	82	<i>Xanthomonas oryzae</i>	Blight
35	<i>P. marginata</i>	Corm scab and leaf blight	83	<i>X. oryzaicola</i>	Leaf streak
	<i>Xanthomonas gummisudans</i>	Blight		<i>Pastinaca sativa</i> (parsnip)	
36			84	<i>Pseudomonas pastinacae</i>	Brown rot
	<i>Glycine soja</i> (soybean)			<i>Phaseolus vulgaris</i> (kidney bean)	
37	<i>Pseudomonas glycinea</i>	Blight	85	<i>Corynebacterium flaccum-faciens</i>	Wilt
38	<i>P. tabaci</i>	Wildfire	86	<i>Pseudomonas flectens</i>	Pod twist
39	<i>Xanthomonas phaseoli so-jensis</i>	Pustule	87	<i>P. phaseolicola</i>	Halo blight
	<i>Gossypium</i> spp. (cotton)		88	<i>P. stizolobii</i>	Leaf spot
40	<i>Xanthomonas malvacearum</i>	Angular leaf spot	89	<i>P. syringae</i>	Lilac blight
	<i>Helianthus annuus</i> (sunflower)		90	<i>P. viridiflava</i>	Leaf spot and blight
41	<i>Pseudomonas helianthi</i>	Leaf spot	91	<i>Xanthomonas phaseoli</i>	Common blight
	<i>Hordeum vulgare</i> (barley)		92	<i>X. phaseoli fuscans</i>	Fuscous blight
42	<i>Pseudomonas coronafaciens</i>	Halo blight		<i>Phleum pratense</i> (timothy)	
43	<i>P. striafaciens</i>	Stripe blight	93	<i>Xanthomonas translucens phleipratensis</i>	Streak
44	<i>Xanthomonas translucens</i>	Blight		<i>Pisum sativum</i> (garden pea)	
	<i>Ipomea batatas</i> (sweet potato)		94	<i>Pseudomonas pisi</i>	Blight
45	<i>Pseudomonas viciae</i>	Leaf and stem spot	95	<i>Xanthomonas pisi</i>	Stem rot
46	<i>Streptomyces ipomoea</i>	Soil rot		<i>Populus</i> spp. (poplar)	
	<i>Iris</i> spp. (iris)		96	<i>Corynebacterium humiferum</i>	Wetwood
47	<i>Pseudomonas cichorii</i>	Leaf blight	97	<i>Pseudomonas rimaefaciens</i>	Canker
48	<i>P. iridicola</i>	Leaf blight		<i>Prunus domestica</i> (garden plum)	
49	<i>P. marginata</i>	Corm scab and leaf blight	98	<i>Pseudomonas morsprunorum</i>	Canker and leaf spot
50	<i>Xanthomonas tardicrescens</i>	Leaf blight	99	<i>P. syringae</i>	Canker and gummosis
	<i>Juglans</i> spp. (walnut)		100	<i>Xanthomonas pruni</i>	Fruit and leaf spot
51	<i>Erwinia nigrifluens</i>	Bark canker		<i>Prunus persica</i> (peach)	
52	<i>Xanthomonas juglandis</i>	Blight	101	<i>Pseudomonas syringae</i>	Canker and gummosis
	<i>Lactuca sativa</i> (lettuce)		102	<i>Xanthomonas pruni</i>	Leaf and fruit spot
53	<i>Pseudomonas cichorii</i>	Head rot		<i>Pyrus communis</i> (pear)	
54	<i>P. marginalis</i>	Marginal blight & rot	103	<i>Erwinia amylovora</i>	Fire blight
55	<i>P. viridilivida</i>	Leaf spot and wilt	104	<i>Pseudomonas barkeri</i>	Blossom blight
56	<i>Xanthomonas vitians</i>	Wilt and rot	105	<i>P. nectarophila</i>	Blossom blight
	<i>Lycopersicon esculentum</i> (tomato)		106	<i>P. syringae</i>	Twig and blossom blight
57	<i>Corynebacterium michiganense</i>	Canker		<i>Raphanus sativus</i> (garden radish)	
	<i>Pseudomonas gardeneri</i>	Fruit spot and scab			
58	<i>P. solanacearum</i>	Wilt	107	<i>Pseudomonas maculicola</i>	Black spot
59	<i>P. tomato</i>	Bacterial speck	108	<i>Xanthomonas campestris</i>	Black rot
60	<i>Xanthomonas vesicatoria</i>	Bacterial spot	109	<i>X. vesicatoria raphani</i>	Leaf spot
	<i>Malus pumila</i> (common apple)			<i>Rheum rhaponticum</i> (garden rhubarb)	
62	<i>Agrobacterium rhizogenes</i>	Hairy root	110	<i>Erwinia rhapontici</i>	Crown rot
63	<i>A. tumefaciens</i>	Crown gall	111	<i>Pseudomonas marginalis</i>	Soft rot
64	<i>Erwinia amylovora</i>	Fire blight		<i>Ribes aureum</i> (golden currant)	
65	<i>Pseudomonas melophthora</i>	Brown rot of fruit	112	<i>Pseudomonas ribicola</i>	Leaf spot
66	<i>P. papulans</i>	Blisters spot		<i>Rosa multiflora</i> (Japanese rose)	
67	<i>P. pomi</i>	Fruit rot	113	<i>Agrobacterium rhizogenes</i>	Hairy root
68	<i>P. syringae</i>	Twig blight			
	<i>Medicago sativa</i> (alfalfa)				
69	<i>Corynebacterium insidiosum</i>	Wilt			

continued

## 136. BACTERIAL PARASITES: PLANTS

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
	<i>Rosa multiflora</i> (Japanese rose)		136	<i>Vicia faba</i> (broad bean) <i>Pseudomonas viciae</i>	Leaf and stem spot
114	<i>Agrobacterium tumefaciens</i>	Crown gall		<i>Vitis</i> spp. (grape)	
115	<i>Pseudomonas syringae</i>	Leaf spot and blast	137	<i>Agrobacterium tumefaciens</i>	Crown gall
	<i>Salix</i> spp. (willow)		138	<i>Erwinia vitivora</i>	Blight
116	<i>Erwinia salicis</i>	Watermark	139	<i>Xanthomonas vitis-carnosae</i>	Leaf spot
117	<i>Pseudomonas saliciperda</i>	Blight		<i>Zea mays</i> (corn)	
	<i>Solanum tuberosum</i> (potato)		140	<i>Erwinia carotovora zeae</i>	Stalk rot and leaf blight
118	<i>Bacillus</i> spp.	Storage rots	141	<i>E. dissolvens</i>	Stalk rot
119	<i>Corynebacterium sepedonicum</i>	Ring rot	142	<i>Pseudomonas alboprecipitans</i>	Leaf blight and stalk rot
120	<i>Erwinia</i> spp.	Soft rot	143	<i>P. andropogonis</i>	Stripe
121	<i>E. atroseptica</i>	Blackleg	144	<i>P. desaiiana</i>	Stinking rot
122	<i>Pseudomonas solanacearum</i>	Brown rot	145	<i>P. lapsa</i>	Leaf and stalk rot
123	<i>Streptomyces scabies</i>	Scab	146	<i>Xanthomonas stewartii</i>	Wilt
	<i>Trifolium</i> spp. (clover)			Fleshy vegetables <sup>1</sup>	
124	<i>Pseudomonas cichorii</i>	Leaf blight	147	<i>Erwinia aroideae</i>	Soft rots
125	<i>P. radiciperda</i>	Root rot	148	<i>E. atroseptica</i>	Soft rots
126	<i>P. stizolobii</i>	Leaf spot	149	<i>E. carotovora</i>	Soft rots
127	<i>P. syringae</i>	Leaf spot and blight	150	<i>E. chrysanthemi</i>	Soft rots
	<i>Triticum aestivum</i> (wheat)			Numerous plants	
128	<i>Corynebacterium iranicum</i>	Spike blight	151	<i>Agrobacterium tumefaciens</i>	Crown gall
129	<i>C. tritici</i>	Spike blight	152	<i>Corynebacterium fascians</i>	Fasciation
130	<i>Pseudomonas atrofaciens</i>	Basal glume rot	153	<i>Erwinia amylovora</i>	Fire blight of Rosaceae
131	<i>Xanthomonas translucens undulosa</i>	Black chaff	154	<i>E. aroideae</i>	Soft rot
	<i>Ulmus</i> spp. (elm)		155	<i>E. atroseptica</i>	Soft rot
132	<i>Erwinia nimipressuralis</i>	Wetwood	156	<i>E. carotovora</i>	Soft rot
133	<i>Pseudomonas lignicola</i>	Black streak of wood	157	<i>E. chrysanthemi</i>	Soft rot
134	<i>P. ulmi</i>	Leaf spot	158	<i>Pseudomonas solanacearum</i>	Brown rot
	<i>Vicia faba</i> (broad bean)		159	<i>P. syringae</i>	Blight
135	<i>Pseudomonas fabae</i>	Blight			

<sup>1</sup>/ Plants of tuberous and fleshy roots.

Contributors: (a) Dickey, Robert S., (b) Burkholder, W. H.

References: [1] Anderson, H. W. 1956. Diseases of fruit crops. McGraw-Hill, New York. [2] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. Bergey's Manual of determinative bacteriology. Ed. 7. Williams and Wilkins, Baltimore. [3] Chupp, C., and A. F. Sherf. 1960. Vegetable diseases and their control. Ronald Press, New York. [4] Dickson, J. G. 1956. Diseases of field crops. Ed. 2. McGraw-Hill, New York. [5] Dowson, W. J. 1957. Plant diseases due to bacteria. Ed. 2. Cambridge Univ. Press, London. [6] Elliott, C. 1951. Manual of bacterial plant pathogens. Ed. 2. Chronica Botanica, Waltham, Mass. [7] U.S. Department of Agriculture. 1953. Plant diseases. Yearbook of Agriculture. U.S. Govt. Printing Office, Washington, D. C.

## 137. FUNGAL PARASITES: PLANTS

### Part I. FIELD, FRUIT, AND VEGETABLE CROPS

Most vegetable crops have damping-off and root rot caused by *Phytophthora* spp., *Pythium* spp., or *Rhizoctonia solani*.

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
	<i>Allium</i> spp. (onion)			<i>Asparagus officinalis</i> (garden asparagus)	
1	<i>Botrytis</i> spp.	Neck rot	5	<i>Fusarium oxysporum</i>	Wilt, root rot
2	<i>Peronospora destructor</i>	Downy mildew	6	<i>Puccinia asparagi</i>	Rust
3	<i>Pyrenochaeta terrestris</i>	Pink rot			
4	<i>Urocystis cepulae</i>	Smut			

continued



## 137. FUNGAL PARASITES: PLANTS

### Part I. FIELD, FRUIT, AND VEGETABLE CROPS

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
7	<i>Avena sativa</i> (common oat)		54	<i>Glycine soja</i> (soybean)	
8	<i>Leptosphaeria avenaria</i>	Black stem	55	<i>Cephalosporium gregatum</i>	Brown stem rot
9	<i>Puccinia coronata</i>	Crown rust	56	<i>Cercospora sojina</i>	Frogeye leaf spot
10	<i>P. graminis</i>	Stem rust	57	<i>Colletotrichum truncatum</i>	Anthracnose
11	<i>Ustilago avenae</i>	Loose smut	58	<i>Corynespora cassicola</i>	Target spot
	<i>U. kolleri</i>	Covered smut		<i>Diaporthe phaseolorum batatis</i>	Stem canker
12	<i>Beta vulgaris</i> (common beet)		59	<i>D. phaseolorum sojiae</i>	Pod and stem blight
13	<i>Aphanomyces cochlioides</i>	Black root, tip rot	60	<i>Fusarium oxysporum tracheiphilum</i>	Fusarium wilt
14	<i>Cercospora beticola</i>	Cercospora leaf spot	61	<i>Glomerella glycines</i>	Anthracnose
	<i>Fusarium</i> spp.	Root rot, storage rot, wilt	62	<i>Macrophomina phaseoli</i>	Ashy stem blight
15	<i>Peronospora schachtii</i>	Downy mildew	63	<i>Pellicularia rolfsii</i>	Southern wilt
16	<i>Pleospora betae</i>	Leaf spot, root rot	64	<i>Peronospora manshurica</i>	Downy mildew
17	<i>Uromyces betae</i>	Rust	65	<i>Phyllosticta sojicola</i>	Leaf spot
	<i>Capsicum frutescens</i> (bush red pepper)		66	<i>Phymatotrichum omnivorum</i>	Texas root rot
18	<i>Cercospora capsici</i>	Frogeye leaf spot	67	<i>Sclerotinia sclerotiorum</i>	Stem rot
19	<i>Gloeosporium piperatum</i>	Anthracnose	68	<i>Septoria glycines</i>	Brown spot
20	<i>Phytophthora capsici</i>	Phytophthora blight		<i>Gossypium</i> spp. (cotton)	
	<i>Carya illinoensis</i> (pecan)		69	<i>Fusarium oxysporum</i>	Fusarium wilt
21	<i>Cercospora fusca</i>	Brown leaf spot	70	<i>Glomerella gossypii</i>	Anthracnose
22	<i>Cladosporium effusum</i>	Scab	71	<i>Pellicularia filamentosa</i>	"Sore shin" seedling stem canker
23	<i>Gnomonia caryae</i>	Liver spot	72	<i>Phymatotrichum omnivorum</i>	Root rot
24	<i>G. nerviseda</i>	Vein spot	73	<i>Puccinia stakmanii</i>	Rust
25	<i>Mycosphaerella caryigena</i>	Downy spot	74	<i>Verticillium albo-atrum</i>	Wilt
26	<i>M. dendroides</i>	Leaf blotch		<i>Hordeum vulgare</i> (barley)	
27	<i>Phymatotrichum cactorum</i>	Cotton root rot	75	<i>Erysiphe graminis</i>	Powdery mildew
	<i>Citrus</i> spp. (citrus)		76	<i>Gibberella zeae</i>	Fusarium blight (scab)
28	<i>Clitocybe tabescens</i>	Root rot	77	<i>Helminthosporium gramineum</i>	Stripe disease
29	<i>Diaporthe citri</i>	Melanose, stem-end rot	78	<i>H. sativum</i>	Spot blotch, root rot, foot rot, kernel blight
30	<i>Diplodia natalensis</i>	Twig and branch die-back, stem-end rot	79	<i>Puccinia graminis</i>	Stem rust
31	<i>Elsinoe fawcettii</i>	Scab	80	<i>P. hordei</i>	Leaf rust
32	<i>Glomerella cingulata</i>	Withertip, anthracnose	81	<i>Pyrenophora teres</i>	Net blotch
33	<i>Phytophthora citrophthora</i>	Foot rot, brown rot	82	<i>Rhynchosporium secalis</i>	Scald
	Cucurbitaceae (cucumber, gourd, muskmelon, pumpkin, squash, watermelon)		83	<i>Typhula itoana</i>	Snow mold
34	<i>Alternaria cucumerina</i>	Leaf blight	84	<i>Ustilago hordei</i>	Covered smut
35	<i>Cladosporium cucumerinum</i>	Scab	85	<i>U. nigra</i>	Black or semiloose smut
36	<i>Colletotrichum lagenarium</i>	Anthracnose <sup>1</sup>	86	<i>U. nuda</i>	Loose smut
37	<i>Erysiphe cichoracearum</i>	Powdery mildew		<i>Ipomoea batatas</i> (sweet potato)	
38	<i>Fusarium oxysporum</i>	Fusarium wilt <sup>2</sup>	87	<i>Endoconidiophora fibriata</i>	Black rot
39	<i>F. oxysporum niveum</i>	Fusarium wilt <sup>3</sup>	88	<i>Fusarium oxysporum</i>	Wilt, stem rot
40	<i>F. solani</i>	Fusarium root rot	89	<i>Monilochaetes infuscans</i>	Scurf
41	<i>Mycosphaerella melonis</i>	Black rot	90	<i>Plenodomus destruens</i>	Foot rot
42	<i>Pseudoperonospora cubensis</i>	Downy mildew	91	<i>Streptomyces ipomoea</i>	Pox or soil rot
43	<i>Verticillium albo-atrum</i>	Verticillium wilt		<i>Juglans regia</i> (Persian walnut)	
	<i>Daucus carota</i> (carrot)		92	<i>Armillaria mellea</i>	Root rot
44	<i>Alternaria dauci</i>	Leaf blight	93	<i>Ascochyta juglandis</i>	Ring spot
45	<i>A. radicina</i>	Black rot	94	<i>Dothiorella gregaria</i>	Dieback, black sap
46	<i>Cercospora carotae</i>	Blight, leaf spot	95	<i>Exosporina fawcetti</i>	Branch wilt, canker
	<i>Fragaria chiloensis</i> (chiloe strawberry)		96	<i>Gnomonia leptostyla</i>	Leaf blotch
47	<i>Botrytis cinerea</i>	Gray mold rot	97	<i>Phytophthora cactorum</i>	Crown rot
48	<i>Dendrophoma obscurans</i>	Leaf blight, stem-end rot		<i>Lactuca sativa</i> (lettuce)	
49	<i>Diplocarpon earliana</i>	Leaf scorch	98	<i>Botrytis cinerea</i>	Gray mold
50	<i>Gnomonia fragariae</i>	Leaf and stem blight	99	<i>Bremia lactucae</i>	Downy mildew
51	<i>Mycosphaerella fragariae</i>	Leaf spot		<i>Lycopersicon esculentum</i> (tomato)	
52	<i>Phytophthora fragariae</i>	Red stele	100	<i>Alternaria solani</i>	Early blight
53	<i>Verticillium albo-atrum</i>	Verticillium wilt			

/1/ Disease of cucurbitaceous plants other than squash. /2/ Disease of muskmelon. /3/ Disease of watermelon.

continued



## 137. FUNGAL PARASITES: PLANTS

### Part I. FIELD, FRUIT, AND VEGETABLE CROPS

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)	(B)	(B)	(A)	(B)	(B)
101	<i>Lycopersicon esculentum</i> (tomato) <i>Colletotrichum phomoides</i>	Anthracnose	144	<i>Phaseolus vulgaris</i> (kidney bean) <i>Fusarium phaseoli</i> , <i>F. solani</i>	Root rot
102	<i>Fusarium oxysporum</i>	Fusarium wilt	145	<i>Macrophomina phaseoli</i>	Ashy stem blight, charcoal rot, leaf spot, root rot
103	<i>Phoma destructiva</i>	Phoma rot	146	<i>Sclerotinia sclerotiorum</i>	White mold (sclerotinia wilt)
104	<i>Phytophthora infestans</i>	Late blight	147	<i>Sclerotium rolfsii</i>	Southern blight
105	<i>Septoria lycopersici</i>	Leaf spot	148	<i>Uromyces phaseoli</i>	Rust
106	<i>Stemphylium solani</i>	Gray leaf spot		<i>Pisum sativum</i> (garden pea)	
107	<i>Malus pumila</i> (common apple) <i>Botryosphaeria ribis</i>	Botryosphaeria canker and fruit rot	149	<i>Aphanomyces euteiches</i>	Root rot
108	<i>Clitocybe tabescens</i>	Root rot	150	<i>Ascochyta pinodella</i>	Ascochyta foot rot
109	<i>Corticium galactinum</i>	White root rot	151	<i>A. pisi</i>	Ascochyta leaf and pod spot
110	<i>Gloeodes pomigena</i>	Sooty blotch	152	<i>Colletotrichum pisi</i>	Anthracnose
111	<i>Glomerella cingulata</i>	Bitter rot of fruit, stem canker	153	<i>Erysiphe polygoni</i>	Powdery mildew
112	<i>Gymnosporangium juniperi</i>	Rust	154	<i>Fusarium oxysporum pisi</i> , strain 1	Fusarium wilt
113	<i>Phyllosticta solitaria</i>	Blotch, leaf spot, canker	155	<i>F. oxysporum pisi</i> , strain 2	Near wilt
114	<i>Podosphaera leucotricha</i>	Powdery mildew	156	<i>F. solani</i>	Root rot
115	<i>Venturia inaequalis</i>	Scab	157	<i>Mycosphaerella pinodes</i>	Mycosphaerella blight
116	<i>Xylaria mali</i>	Black root rot	158	<i>Peronospora viciae</i>	Downy mildew
117	<i>Medicago sativa</i> (alfalfa) <i>Leptosphaeria pratensis</i>	Leaf spot, root rot, crown rot		<i>Prunus amygdalus</i> (almond)	
118	<i>Peronospora trifoliorum</i>	Downy mildew	159	<i>Armillaria mellea</i>	Root rot
119	<i>Phoma herbarum medicaginis</i>	Spring black stem	160	<i>Coryneum carpophilum</i>	Blight, shot hole
120	<i>Pseudopeziza jonesii</i>	Yellow leaf blotch	161	<i>Monilinia fructicola</i>	Peach brown rot
121	<i>P. medicaginis</i>	Common leaf spot	162	<i>M. laxa</i>	Brown rot, blossom blight
122	<i>Sclerotinia trifoliorum</i>	Crown rot, root rot		<i>Prunus domestica</i> (garden plum)	
123	<i>Uromyces striatus</i>	Rust	163	<i>Armillaria mellea</i>	Root rot
124	<i>Urophlyctis alfalfae</i>	Crown wart	164	<i>Dibotryon morbosum</i>	Black knot
	<i>Nicotiana tabacum</i> (common tobacco)		165	<i>Monilinia fructicola</i>	Brown rot, blossom blight
125	<i>Cercospora nicotianae</i>	Frogeye leaf spot	166	<i>M. laxa</i>	European brown rot, blossom and twig blight
126	<i>Fusarium oxysporum</i>	Fusarium wilt		<i>Prunus persica</i> (peach)	
127	<i>Macrophomina phaseoli</i>	Charcoal rot	167	<i>Armillaria mellea</i>	Root rot
128	<i>Pellicularia filamentosa</i>	Stem canker	168	<i>Clitocybe tabescens</i>	Root rot
129	<i>Peronospora tabacina</i>	Blue mold	169	<i>Coryneum carpophilum</i>	Blight, shot hole
130	<i>Phytophthora parasitica</i>	Black shank	170	<i>Glomerella cingulata</i>	Ripe rot, twig blight
131	<i>Thielaviopsis basicola</i>	Black rot	171	<i>Monilinia fructicola</i>	Brown rot, twig canker
	<i>Oryza sativa</i> (rice)		172	<i>Taphrina deformans</i>	Leaf curl
132	<i>Cercospora oryzae</i>	Cercospora spot		<i>Pyrus communis</i> (pear)	
133	<i>Cochliobolus miyabeanus</i>	Helminthosporium blight	173	<i>Botrytis cinerea</i>	Gray mold
	<i>Persea americana</i> (American avocado)		174	<i>Clitocybe tabescens</i>	Root rot
134	<i>Botryosphaeria ribis</i>	Branch canker, fruit rot	175	<i>Neofabraea malicorticis</i>	Black spot canker
135	<i>Cercospora purpurea</i>	Cercospora spot or blotch	176	<i>N. perennans</i>	Perennial canker
136	<i>Colletotrichum gloeosporioides</i>	Anthracnose or black spot	177	<i>Podosphaera leucotricha</i>	Powdery mildew
137	<i>Diplodia theobromae</i>	Stem-end rot	178	<i>Venturia pyrina</i>	Scab
138	<i>Phomopsis</i> spp.	Stem-end rot		<i>Solanum tuberosum</i> (potato)	
139	<i>Phytophthora cinnamomi</i>	Phytophthora root rot	179	<i>Alternaria solani</i>	Early blight
140	<i>Sphaceloma perseae</i>	Scab <sup>4</sup>	180	<i>Fusarium</i> spp.	Wilt and tuber rot
141	<i>Verticillium albo-atrum</i>	Verticillium wilt	181	<i>Phytophthora infestans</i>	Late blight
	<i>Phaseolus vulgaris</i> (kidney bean)		182	<i>Streptomyces scabies</i>	Common scab
142	<i>Colletotrichum lindemuthianum</i>	Anthracnose	183	<i>Verticillium albo-atrum</i>	Verticillium wilt
143	<i>Erysiphe polygoni</i>	Powdery mildew		<i>Trifolium</i> spp. (clover)	
			184	<i>Colletotrichum trifolii</i>	Southern anthracnose

<sup>4</sup>/ Disease of fruit and foliage.

continued

## 137. FUNGAL PARASITES: PLANTS

### Part I. FIELD, FRUIT, AND VEGETABLE CROPS

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
185	<i>Trifolium</i> spp. (clover)		203	<i>Vitis</i> spp. (grape)	
186	<i>Cymadothea trifolii</i>	Sooty blotch <sup>5</sup>	204	<i>Cryptosporella viticola</i>	Dead-arm
187	<i>Erysiphe polygoni</i>	Powdery mildew <sup>6</sup>	205	<i>Elsinoe ampelina</i>	Anthracnose
188	<i>Kabatiella caulivora</i>	Northern anthracnose <sup>7</sup>	206	<i>Guignardia bidwellii</i>	Black rot
189	<i>Phoma herbarum medicaginis</i>	Spring black stem <sup>6</sup>	207	<i>Plasmopara viticola</i>	Downy mildew
	<i>Pseudoplea trifolii</i>	Leaf spot		<i>Uncinula necator</i>	Powdery mildew
190	<i>Triticum aestivum</i> (wheat)		208	<i>Zea mays</i> (corn)	
191	<i>Erysiphe graminis</i>	Powdery mildew	209	<i>Cochliobolus heterostrophus</i>	Southern leaf blight, seedling blight
192	<i>Gibberella zeae</i>	Fusarium blight (scab)	210	<i>Diplodia macrospora, D. zeae</i>	Stalk rot, dry ear rot
193	<i>Helminthosporium sativum</i>	Crown rot, root rot	211	<i>Gibberella fujikuroi</i>	Pink ear rot, seedling blight
194	<i>Leptosphaeria nodorum</i>	Glume blotch, node canker	212	<i>G. zeae</i>	Stalk rot, red ear rot, seedling blight, root rot
195	<i>Ophiobolus graminis</i>	Take-all	213	<i>Helminthosporium carbonum</i>	Northern leaf spot, charred ear, seedling blight
196	<i>Puccinia glumarum</i>	Stripe rust	214	<i>H. turcicum</i>	Northern leaf blight, seedling blight
197	<i>P. graminis</i>	Stem rust	215	<i>Physalospora zeae</i>	Gray ear rot
198	<i>P. rubigo-vera</i>	Leaf rust	216	<i>Puccinia sorghi</i>	Rust
199	<i>Septoria tritici</i>	Leaf blotch	217	<i>Sclerospora graminicola</i>	Downy mildew
200	<i>Tilletia brevipaciens</i>	Dwarf bunt		<i>Ustilago maydis</i>	Smut
201	<i>T. caries, T. foetida</i>	Bunt (stinking smut)			
202	<i>Urocystis tritici</i>	Flag smut			
	<i>Ustilago tritici</i>	Loose smut			

/s/ Disease of red, white, and alsike clover. /e/ Disease of red clover. /7/ Disease of crimson and red clover.

Contributor: Andersen, Axel L.

References: [1] Beattie, J. H. 1951. U.S. Dept. Agr. Farmers' Bull. 1468. [2] Boswell, V. R., S. P. Doolittle, and L. M. Pultz. 1952. Ibid. 2051. [3] Demaree, J. B. 1948. Ibid. 1891. [4] Demaree, J. B., and G. W. Still. 1951. Ibid. 1893. [5] Dickson, J. G. 1947. Diseases of field crops. McGraw-Hill, New York. [6] Doolittle, S. P. 1948. U.S. Dept. Agr. Farmers' Bull. 1934. [7] Dykstra, T. P. 1948. Ibid. 1881. [8] Harter, L. L., W. J. Zaumeyer, and B. L. Wade. 1945. Ibid. 1735. [9] Rhoads, A. S., and E. F. DeBusk. 1931. Florida Univ. Agr. Expt. Sta. (Gainesville) Bull. 229. [10] Stevens, H. E., and R. B. Piper. 1941. U.S. Dept. Agr. Circ. 582. [11] Thompson, R. C. 1945. U.S. Dept. Agr. Farmers' Bull. 1646. [12] U.S. Department of Agriculture. 1953. Plant diseases. Yearbook of agriculture. U.S. Govt. Printing Office, Washington, D.C. [13] Walker, J. C. 1947. U.S. Dept. Agr. Farmers' Bull. 1060. [14] Walker, J. C. 1952. Diseases of vegetable crops. McGraw-Hill, New York. [15] Weiss, F. 1950. U.S. Dept. Agr. Plant Disease Surv. Spec. Publ. 1(1-3). [16] Weiss, F., and M. J. O'Brien. 1952-53. Ibid. 1(4). [17] Zaumeyer, W. J., and H. R. Thomas. 1949. U.S. Dept. Agr. Farmers' Bull. 1692.

### Part II. FOREST TREES

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
	<i>Abies</i> spp. (fir)			<i>Abies</i> spp. (fir)	
1	<i>Adelopus gammamii</i>	Swiss needle cast	9	<i>Cytispora abietis</i>	Cytispora canker
2	<i>Aleurodiscus amorphus</i>	Aleurodiscus canker	10	<i>Dasyscypha resinaria</i>	Dasyscypha canker
3	<i>Armillaria mellea</i>	Shoestring root rot	11	<i>Fomes amosus</i>	Root and butt rot
4	<i>Bifusella abietis, B. faullii</i>	Needle cast	12	<i>F. pini</i>	Red ring rot
5	<i>Caecoma faulliana</i>	Needle rust	13	<i>F. pinicola</i>	Brown cubical rot
6	<i>Calyptospora goeppertiana</i>	Needle rust	14	<i>Hyalospora aspidiotus</i>	Needle cast
7	<i>Cephalosporium</i> sp.	Cephalosporium canker	15	<i>Hypoderma robustum</i>	Needle cast
8	<i>Corticium galactinum</i>	Corticium rot			

continued

## 137. FUNGAL PARASITES: PLANTS

### Part II. FOREST TREES

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
16	<i>Abies</i> spp. (fir)		64	<i>Betula</i> spp. (birch)	
	<i>Hypodermella abietis</i> , <i>H. mirabilis</i> , <i>H. nervata</i>	Needle cast		<i>Fomes fomentarius</i>	White mottled rot
17	<i>Lophodermium abietis</i> , <i>L. uncinatum</i>	Needle cast	65	<i>F. igniarius</i>	Heartrot, canker
18	<i>Melampsora abieticapræarum</i>	Needle rot	66	<i>Melampsorium betulinum</i>	Leaf rust
19	<i>Melampsorella cerastii</i>	Witches'-broom	67	<i>Nectria galligena</i>	Nectria canker
20	<i>Milesia fructuosa</i> , <i>M. marginalis</i>	Needle rust	68	<i>Pholiota squarrosa</i>	Brown mottled rot
21	<i>M. pycnograndis</i>	Needle-witches'-broom rust	69	<i>Poria obliqua</i>	Canker
22	<i>Peridermium holwayi</i> , <i>P. ornamentale</i> , <i>P. rugosum</i>	Needle rust	70	<i>Stereum murrayi</i>	Rot, canker
23	<i>Phacidium infestans</i>	Needle blight	71	<i>Taphrina boycei</i> , <i>T. flava</i>	Yellow leaf blister
24	<i>Phomopsis boycei</i>	Phomopsis canker		<i>Cupressus arizonica</i> (Arizona cypress)	
25	<i>Polyporus abietinus</i>	White, pitted sap rot	72	<i>Gymnosporangium cupressi</i>	Fusiform gall rust of juniper
26	<i>P. balsameus</i>	Brown butt rot		<i>Fagus grandifolia</i> (American beech)	
27	<i>P. dryadeus</i>	White root rot	73	<i>Fomes fomentarius</i>	White mottled rot
28	<i>P. schweinitzii</i>	Root and butt rot	74	<i>F. igniarius</i>	Heartrot
29	<i>Poria subacida</i>	Butt rot	75	<i>Nectria coccinea</i>	Beech bark disease
30	<i>Rehmiellopsis balsamiae</i>	Tip blight	76	<i>Phytophthora cactorum</i>	Phytophthora blight
31	<i>Scleroderris abieticola</i>	Scleroderris canker	77	<i>Polyporus glomeratus</i>	Light-brown spongy heartrot
32	<i>Spicaria anomala</i>	Brown stain of fir	78	<i>Poria obliqua</i>	Canker
33	<i>Stereum chailleti</i>	Patchy rot		<i>Fraxinus</i> spp. (ash)	
34	<i>S. sanguinolentum</i>	Red heartrot	79	<i>Armillaria mellea</i>	Root rot
35	<i>Uredinopsis atkinosinii</i> , <i>U. ceratophora</i> , <i>U. longimucronata</i> , <i>U. macrosperma</i> , <i>U. mirabilis</i>	Needle rust	80	<i>Cytispora annularis</i>	Cytispora canker
36	<i>Acer</i> spp. (maple)		81	<i>Fomes fraxinophilus</i>	White heartrot
	<i>Cristulariella depraehens</i>	Leaf spot and wilt	82	<i>Marssonina fraxini</i>	Leaf spot
37	<i>Daedalea unicolor</i>	Rot, canker	83	<i>Mycosphaerella fraxinicola</i>	Leaf spot
38	<i>Daldinia concentrica</i>	White rot	84	<i>Phymatotrichum omnivorum</i>	Phymatotrichum root rot
39	<i>Endoconidiophora virescens</i>	Sapstreak	85	<i>Polyporus hispidus</i>	Spongy white rot
40	<i>Eutypella parasitica</i>	Eutypella canker	86	<i>Puccinia peridermiospora</i>	Leaf rust
41	<i>Fomes connatus</i>	Butt rot		<i>Juglans nigra</i> (black walnut)	
42	<i>F. igniarius</i>	Heartrot	87	<i>Armillaria mellea</i>	Shoestring root rot
43	<i>Hydnum septentrionale</i>	Soft, spongy white rot	88	<i>Fomes everhartii</i>	White heartrot
44	<i>Hymenochaete agglutinans</i>	Hymenochaete canker	89	<i>Phymatotrichum omnivorum</i>	Phymatotrichum root rot
45	<i>Hypoxylon blakei</i>	Hypoxylon canker	90	<i>Phytophthora cinnamomi</i>	Root rot
46	<i>Nectria cinnabarina</i>	Nectria dieback		<i>Juniperus</i> spp. (juniper)	
47	<i>N. galligena</i>	Nectria canker	91	<i>Coccodotthis sphaeroidea</i>	Leaf blight
48	<i>Phleospora aceris</i>	Leaf spot	92	<i>Fomes juniperinus</i>	Juniper pocket rot
49	<i>Phytophthora cactorum</i>	Bleeding canker	93	<i>F. subroseus</i>	Brown pocket rot
50	<i>Polyporus glomeratus</i>	Rot, canker	94	<i>Gymnosporangium aurantiacum</i>	Mountain ash and mountain juniper rot
51	<i>Rhytisma acerinum</i>	Tar spot	95	<i>G. bermudianum</i>	Juniper rust
52	<i>Schizoxylon microsporium</i>	Schizoxylon canker	96	<i>G. betheli</i>	Elongate gall rust of juniper
53	<i>Stereum murrayi</i>	Rot, canker	97	<i>G. clavariaeforme</i> , <i>G. clavipes</i>	Fusiform gall rust of juniper
54	<i>Uncinula circinatis</i>	Powdery mildew	98	<i>G. corniculans</i>	Serviceberry-juniper rust
55	<i>Venturia acerina</i>	Red-brown spot	99	<i>G. davisii</i>	Chokeberry-mountain juniper rust
56	<i>Verticillium albo-atrum</i>	Verticillium wilt	100	<i>G. effusum</i>	Fusiform gall rust of juniper
57	<i>Xylaria digitata</i>	Xylaria root rot	101	<i>G. exiguum</i>	Hawthorne-alligator juniper rust
	<i>Alnus</i> spp. (alder)		102	<i>G. externum</i>	Porteranthus, fusiform gall rust of juniper
58	<i>Cytispora pulcherrima</i>	Cytispora canker	103	<i>G. floriforme</i>	Hawthorne-juniper rust
59	<i>Didymosphaeria oregonensis</i>	Didymosphaeria canker			
60	<i>Erysiphe aggregata</i>	Powdery mildew			
61	<i>Melampsorium alni</i>	Leaf rust			
62	<i>Taphrina amentorum</i> , <i>T. occidentalis</i> , <i>T. robinsoniana</i> , <i>T. rugosa</i>	Leaf blister			
63	<i>T. macrophylla</i>	Yellow leaf blister			

continued



## 137. FUNGAL PARASITES: PLANTS

### Part II. FOREST TREES

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
104	<i>Juniperus</i> spp. (juniper)		143	<i>Pinus strobus</i> (eastern white pine) <sup>1,2</sup>	
	<i>Gymnosporangium globosum</i>	Hawthorne-cedar rust		<i>Fomes laricis</i>	Brown cubical rot
105	<i>G. harnessianum</i>	Serviceberry-western juniper rust	144	<i>F. pini</i>	Red ring rot
106	<i>G. inconspicuum</i>	Serviceberry-Utah twig rust	145	<i>Hypodermma desmazierii</i>	Needle cast
107	<i>G. juniperi-virginiana</i>	Apple-cedar rust	146	<i>Lentinus lepideus</i>	Brown cubical rot
108	<i>G. kernianum</i>	Witches'-broom	147	<i>Lophodermium pinastri</i>	Needle cast
109	<i>G. multiporum</i>	Utah juniper rust	148	<i>Neopeckia coulteris</i>	Brown felt blight
110	<i>G. nelsoni</i>	Hawthorne-western juniper rust	149	<i>Phacidium planum</i>	Needle blight
111	<i>G. trachysorum</i>	Fusiform gall rust of juniper	150	<i>Polyporus circinatus</i>	Red root and butt rot
112	<i>G. tuberculatum</i>	Globoid gall rust of mountain juniper	151	<i>Sparassis radicata</i>	Sparassis root rot
113	<i>G. vauqueliniae</i>	Witches'-broom of vauquelinia		<i>Populus</i> spp. (poplar)	
114	<i>Phomopsis juniperovora</i>	Juniper blight	152	<i>Armillaria mellea</i>	Shoestring root rot
	<i>Larix occidentalis</i> (western larch)		153	<i>Ciborinia whetzelii</i>	Ink spot
115	<i>Fomes laricis</i>	Brown trunk rot	154	<i>Cytispora chrysosperma</i>	Cytispora canker
116	<i>Hypodermella laricis</i>	Needle cast	155	<i>Dothichiza populea</i>	Dothichiza canker
117	<i>Melampsora bigelowii</i> , <i>M. medusae</i>	Needle rust	156	<i>Fomes applanatus</i>	White butt rot
118	<i>Sparassis radicata</i>	Sparassis root rot	157	<i>F. fomentarius</i>	White mottled rot
	<i>Picea glauca</i> (white spruce)		158	<i>F. ignarius</i>	White heartrot
119	<i>Chrysomyxa cassandrae</i> , <i>C. chiogenis</i> , <i>C. empetri</i> , <i>C. ledi</i> , <i>C. ledicola</i>	Needle rust	159	<i>Fusicladium radiosum</i>	Shoot blight
120	<i>C. pyrolae</i>	Spruce cone rust	160	<i>Hypoxylon pruinautum</i>	Hypoxylon canker
121	<i>Peridermium coloradense</i>	Witches'-broom	161	<i>Linospora tetraspora</i>	Leaf blight
122	<i>Pucciniastrum americanum</i> , <i>P. arctium</i>	Needle rust	162	<i>Marssonina populi</i>	Leaf spot, shoot blight
123	<i>Rhizina inflata</i>	Rhizina root rot	163	<i>Melampsora bigelowii</i> , <i>M. medusae</i> , <i>M. occidentalis</i>	Leaf spot
	<i>Pinus ponderosa</i> (western yellow pine) <sup>1</sup>		164	<i>Nectria galligena</i>	Nectria canker
124	<i>Atropellis arizonica</i> , <i>A. piniphila</i>	Atropellis canker	165	<i>Neofabraea populi</i>	Neofabraea canker
125	<i>Coleosporium solidaginis</i>	Needle rust	166	<i>Sclerotinia bifrons</i>	Ink spot
126	<i>Cronartium comandrae</i>	Comandra blister rust	167	<i>Septoria musiva</i>	Septoria canker
127	<i>C. comptoniae</i>	Sweet-fern blister rust	168	<i>S. populicola</i>	Leaf spot
128	<i>C. filamentosum</i>	Paintbrush blister rust	169	<i>Taphrina aurea</i>	Yellow leaf blister
129	<i>C. harknessii</i>	Western gall rust	170	<i>T. johansonii</i>	Catkin blister
130	<i>C. quercuum</i>	Gall rust	171	<i>Trametes suaveolens</i>	Soft white rot
131	<i>Dasyscypha ellisiana</i>	Dasyscypha canker	172	<i>Uncinula salicis</i>	Powdery mildew
132	<i>Elytroderma deformans</i>	Witches'-broom, needle cast	173	<i>Valsa nivea</i> , <i>V. sordida</i>	Valsa canker
133	<i>Fomes laricis</i>	Brown cubical rot	174	<i>Xylaria digitata</i>	Xylaria root rot
134	<i>F. pini</i>	Red ring rot		<i>Quercus</i> spp. (oak)	
135	<i>Hypodermella medusa</i>	Needle cast	175	<i>Aleurodiscus oakii</i>	Smooth patch
136	<i>Lentinus lepideus</i>	Brown cubical rot	176	<i>Armillaria mellea</i>	Shoestring root rot
	<i>Pinus strobus</i> (eastern white pine) <sup>1,2</sup>		177	<i>Cronartium cerebrum</i>	Globose gall rust
137	<i>Armillaria mellea</i>	Shoestring root rot	178	<i>C. fusiforme</i>	Southern fusiform rust
138	<i>Atropellis pinicola</i>	Atropellis canker	179	<i>C. strobilinum</i>	Pinecone rust
139	<i>Bifusella linearis</i>	Needle cast	180	<i>Daedalea quercina</i>	Brown cubical rot
140	<i>Calciopsis pinea</i>	Calciopsis canker	181	<i>Endoconidiophora fagacearum</i>	Oak wilt
141	<i>Cronartium ribicola</i>	White pine blister rust	182	<i>Fistulina hepatica</i>	Brown cubical rot
142	<i>Fomes amosus</i>	Root and butt rot	183	<i>Fomes everhartii</i>	White heartrot
			184	<i>Gnomonia veneta</i>	Anthraxnose
			185	<i>Hydnum erinaceus</i>	White rot
			186	<i>Morenoella quercina</i>	Leaf spot
			187	<i>Polyporus berkeleyi</i>	White butt rot
			188	<i>P. croceus</i>	White pocket rot
			189	<i>P. dryophilus</i>	Piped rot
			190	<i>P. frondosus</i>	Butt rot
			191	<i>P. hispidus</i>	Heartrot, canker
			192	<i>P. spraguei</i>	White rot
			193	<i>P. sulphureus</i>	Brown cubical rot
			194	<i>Sphaeropsis quercina</i>	Sphaeropsis canker
			195	<i>Sphaerotheca lanestris</i>	Witches'-broom
			196	<i>Stereum gausapatum</i>	White mottled rot

<sup>1/1</sup> Hardwood. <sup>1/2</sup> Softwood. <sup>1/3</sup> Data also apply to other softwood pines: *P. lambertiana* (sugar pine), *P. monticola* (western white pine).

continued



## 137. FUNGAL PARASITES: PLANTS

### Part II. FOREST TREES

Host Plant and Pathogen		Disease	Host Plant and Pathogen		Disease
(A)		(B)	(A)		(B)
197	<i>Quercus</i> spp. (oak)	White pocket rot	215	<i>Tsuga canadensis</i> (eastern hemlock)	White rot
198	<i>Stereum subpileatum</i>	Strumella canker	216	<i>Ganoderma lucidum</i>	Cedar leaf blight
199	<i>Strumella coryneoides</i>	Oak leaf blister	217	<i>Keithia tsugae</i>	Needle rust
200	<i>Taphrina caerulescens</i>		218	<i>Melanopsora abietis</i> , <i>M. farlowii</i>	White mottled rot
201	<i>Salix</i> spp. (willow)	Botryosphaeria canker	219	<i>Polyporus borealis</i>	Needle rust
202	<i>Botryosphaeria ribis</i>	Cytispora canker	220	<i>Pucciniastrum hydrangeae</i> , <i>P. myrtilli</i>	Brown felt blight
203	<i>Cytispora pulcherrima</i>	Heartrot	221	<i>Ulmus</i> spp. (elm)	Dutch-elm disease
204	<i>Fomes igniarius</i>	Willow scab	222	<i>Ceratostomella ulmi</i>	Chinese-elm root rot
205	<i>Fusicladium saliciper-dum</i>	Leaf rust	223	<i>Chalaropsis thielavioides</i>	Cytispora canker
206	<i>Melanopsora abieticapraearum</i>	Black canker	224	<i>Cytispora ambiens</i>	Dieback
207	<i>Physalospora miyabeana</i>	White rot	225	<i>Dothiorella ulmi</i>	Leaf spot
208	<i>Polyporus squamosus</i>	Soft white rot	226	<i>Gloeosporium ulmicolum</i>	Leaf spot
209	<i>Trametes suaveolens</i>	Powdery mildew	227	<i>Gnomonia ulnea</i>	Elm leaf spot
210	<i>Uncinula salicis</i>	Valsa canker	228	<i>Phleospora ulmi</i>	Pit canker
211	<i>Valsa nivea</i> , <i>V. sordida</i>		229	<i>Phytophthora inflata</i>	Brown rot
212	<i>Sequoia sempervirens</i> (redwood)	White ring rot	230	<i>Pleurotus ulmaris</i>	Canker
213	<i>Poria albipellucida</i>	Brown pocket rot	231	<i>Sphaeropsis ulmicola</i>	Elm canker
214	<i>P. sequoia</i>		232	<i>Tubercularia</i> sp.	Powdery mildew
	<i>Thuja occidentalis</i> (northern white cedar)	Brown cubical rot	233	<i>Uncinula macrocarpa</i>	Verticillium wilt
	<i>Coniophora puteana</i>	Cedar leaf blight	234	<i>Verticillium</i> spp.	Verticillium root disease
	<i>Keithia thujina</i>			<i>V. rhizophagum</i>	
	<i>Tsuga canadensis</i> (eastern hemlock)				
	<i>Fomes robustus-tsugina</i>	White heartrot			

Contributor: Baxter, Dow V.

References: [1] Baxter, D. V. 1933. Univ. Mich. School Forestry Conserv. Bull. 2. [2] Baxter, D. V. 1937. Univ. Mich. School Forestry Conserv. Circ. 1. [3] Baxter, D. V. 1947. Papers Mich. Acad. Sci. 31:931. [4] Baxter, D. V. 1952. Pathology in forest practice. Ed. 2. J. Wiley, New York. [5] Boyce, J. S. 1961. Forest pathology. Ed. 3. McGraw-Hill, New York. [6] Clapper, R. B. 1943. Am. Forests 49:331. [7] Hepting, G. H. 1961. Forest Farmer 21(1):11. [8] Marshall, R. P., and A. M. Waterman. 1948. U.S. Dept. Agr. Farmers' Bull. 1987. [9] Offord, H. R. 1962. World Forestry Congr., 5th, Seattle, 2:882.

## 138. MISTLETOE PARASITES: FOREST TREES

In forest trees, witches'-broom is the chief disease caused by mistletoe.

Host (Common Name)		Parasite (Common Name)	Host (Common Name)		Parasite (Common Name)
(A)		(B)	(A)		(B)
1	<i>Abies</i> spp. (fir)	<i>Phoradendron pauciflorum</i> (western fir mistletoe) <sup>1</sup>	3	<i>Acer</i> spp. (maple)	<i>Phoradendron flavescens</i> (American mistletoe)
2	<i>Abies anabilis</i> (Cascade fir), <i>A. concolor</i> (white fir), <i>A. grandis</i> (grand fir), <i>A. lasiocarpa</i> (alpine fir), <i>A. lasiocarpa arizonica</i> (cork-bark fir)	<i>Arceuthobium campylo-podium abietinum</i> (western dwarf mistletoe)	4	<i>Alnus</i> spp. (alder)	<i>Phoradendron macrophyllum</i> (big-leaf mistletoe)
			5	<i>Cupressus</i> spp. (cypress)	<i>Phoradendron densum</i> (cypress mistletoe)
			6	<i>Fraxinus</i> spp. (ash)	<i>Phoradendron longispicum</i> (long-spiked mistletoe)

<sup>1/1</sup> Leader injury.

continued

138. MISTLETOE PARASITES: FOREST TREES

Host (Common Name)		Parasite (Common Name)	
(A)	(B)	(A)	(B)
7	<i>Fraxinus</i> spp. (ash)	19	<i>Pinus cembroides</i> (Mexican piñon pine) and other piñons
8	<i>Juglans</i> spp. (walnut)	20	<i>Pinus contorta latifolia</i> (lodgepole pine)
9		21	<i>Pinus lambertiana</i> (sugar pine), <i>P. monticola</i> (western white pine), <i>P. reflexa</i> (Mexican white pine)
10	<i>Juniperus</i> spp. (juniper)	22	<i>Pinus latifolia</i> (Apache pine), <i>P. leiophylla chihuahuana</i> (chihuahua pine), <i>P. ponderosa arizonica</i> (Arizona ponderosa pine)
11		23	<i>Pinus ponderosa scopulorum</i> (Rocky Mountain ponderosa pine)
12		24	<i>Populus</i> spp. (poplar)
13	<i>Larix laricina</i> (eastern larch)	25	
14	<i>Larix lyalli</i> (alpine larch), <i>L. occidentalis</i> (western larch)	26	<i>Quercus</i> spp. (oak)
15	<i>Picea breweriana</i> (Brewer spruce), <i>P. engelmanni</i> (Engelmann spruce), <i>P. pungens</i> (Colorado spruce)	27	
16	<i>Picea glauca</i> (white spruce) <sup>2</sup> , <i>P. mariana</i> (black spruce), <i>P. rubens</i> (red spruce) <sup>2</sup>	28	
17	<i>Pinus albicaulis</i> (whitebark pine), <i>P. aristata</i> (bristlecone pine), <i>P. balfouriana</i> (foxtail pine), <i>P. flexilis</i> (limber pine)	29	
18	<i>Pinus attenuata</i> (knobcone pine) <sup>2</sup> , <i>P. contorta latifolia</i> (lodgepole pine) <sup>2</sup> , <i>P. coulteri</i> (Coulter pine), <i>P. jeffreyi</i> (Jeffrey pine), <i>P. ponderosa</i> (western yellow pine), <i>P. radiata</i> (Monterey pine), <i>P. sabinana</i> (digger pine)	30	
		31	
		32	<i>Tsuga heterophylla</i> (western hemlock), <i>T. mertensiana</i> (mountain hemlock)
		33	<i>Ulmus</i> spp. (elm)
		34	

<sup>1/2</sup> Rarely serves as host.

Contributor: Baxter, Dow V.

References: [1] Baxter, D. V. 1952. Pathology in forest practice. Ed. 2. J. Wiley, New York. [2] Boyce, J. S. 1938. Forest pathology. McGraw-Hill, New York. [3] Gill, L. S. 1935. Trans. Conn. Acad. Arts Sci. 32:111. [4] Hawksworth, F. G. 1954. Phytopathology 44:552. [5] Kimmey, J. W., and D. P. Graham. 1960. U.S. Dept. Agr. Forest Serv. Forest Range Expt. Sta. Res. Paper 60. [6] Trelease, W. 1916. The genus *Phoradendron*. Univ. Illinois Press, Urbana.

	Species (A)	Disease (B)	Natural Occurrence in Animals Other than Man (C)	Microscopic Appearance in Man
				Skin (D)
1	<i>Cladosporium werneckii</i>	Tinea nigra	None	Pigmented (light brown to dark green), branching, septate hyphae; may develop closely septate swollen cells and chlamydo-spores
2	<i>Epidermophyton floccosum</i>	Tinea pedis, T. cruris, T. unguium	None	Abundant-branching, septate hyphae; may segment into chains of arthrospores
3	<i>Malassezia furfur</i>	Tinea versicolor	None	Clusters of spherical, thick-walled, budding cells, 3-8 $\mu$ , and short irregular hyphae
4	<i>Microsporum audouinii</i>	Tinea capitis, T. corporis	Dog (rare), monkey (rare)	Branching, septate hyphae; may segment into chains of arthrospores
5	<i>M. canis</i>	Tinea capitis, T. corporis, T. barbae, T. unguium	Cat <sup>1</sup> , chinchilla, dog <sup>1</sup> , horse, monkey	Branching, septate hyphae; may segment into chains of arthrospores
6	<i>M. distortum</i>	Tinea capitis, T. corporis	Dog, monkey <sup>1</sup>	Branching, septate hyphae; may segment into chains of arthrospores
7	<i>M. gypseum</i> <sup>2</sup>	Tinea capitis, T. corporis	Cat, dog <sup>1</sup> , horse <sup>1</sup> , monkey, mouse, rat	Branching, septate hyphae; may segment into chains of arthrospores
8	<i>M. nanum</i>	Tinea capitis, T. corporis	Swine	Branching, septate hyphae; may segment into chains of arthrospores
9	<i>M. vambreuseghemii</i>	Tinea capitis	Dog, Malabar squirrel	Branching, septate hyphae; may segment into chains of arthrospores
10	<i>Piedraia hortai</i>	Black piedra	Primates	None
11	<i>Trichophyton concentricum</i>	Tinea imbricata	None	Abundant-branching, septate hyphae; may segment into chains of arthrospores
12	<i>T. ferrugineum</i>	Tinea capitis, T. corporis	None	Branching, septate hyphae; may segment into chains of arthrospores
13	<i>T. gallinae</i>	Tinea capitis, T. corporis	Dog, poultry <sup>1</sup> , wild birds	Branching, septate hyphae (rare)
14	<i>T. megninii</i>	Tinea barbae, T. capitis, T. unguium, T. corporis	Cattle?	Branching, septate hyphae
15	<i>T. mentagrophytes</i>	Tinea pedis, T. cruris, T. corporis, T. capitis, T. unguium, T. barbae	Many domestic and wild animals	Branching, septate hyphae; may segment into chains of arthrospores
16	<i>T. rubrum</i>	Tinea pedis, T. unguium, T. cruris, T. corporis, T. barbae, T. capitis	Cattle (rare), dog (rare)	Branching, septate hyphae; may segment into chains of arthrospores
17	<i>T. schoenleimii</i>	Favus, Tinea capitis, T. corporis, T. unguium	Dog (rare)	Abundant hyphae; may segment into chains of arthrospores throughout cellular debris of scutulum
18	<i>T. tonsurans</i>	Tinea capitis, T. corporis, T. unguium	None	Branching, septate hyphae; may segment into chains of arthrospores
19	<i>T. verrucosum</i>	Tinea corporis, T. capitis, T. barbae	Cattle <sup>1</sup> , dog (rare), donkey, goat, horse, sheep	Branching, septate hyphae; may segment into chains of arthrospores

<sup>1/1</sup> The more common hosts. <sup>1/2</sup> A common saprophyte in soil.



**PARASITES: MAN**

**MYCOSES**

Microscopic Appearance in Man		Microscopic Appearance of Culture on Sabouraud's Agar	
Nail (E)	Hair (F)		
None	None	Pigmented hyphae produce blastopores laterally, and 1-3 septate conidia in clusters or in short chains from apiculi or short conidiophores	1
Branching, septate hyphae; may segment into chains of arthrospores	None	Macroconidia abundant, clavate, 2-6 cells, blunt-tipped, smooth thin walls; occur in clusters of 2 or 3; no microconidia; abundant chlamydo-spores	2
None	None	No culture method available	3
None	Ectothrix; sheath of small spores, 2-3 $\mu$	Hyphae with chlamydo-spores; microconidia infrequent, clavate, 2.5-4 x 3-6 $\mu$ ; macroconidia rare, rudimentary, ill-formed	4
Rare	Ectothrix; sheath of small spores, 2-3 $\mu$	Macroconidia numerous, 8-15 cells, spindle-shaped, thick rough walls, 8-15 x 40-150 $\mu$ ; microconidia few, clavate, 2-4 x 3-6 $\mu$	5
None	Ectothrix; sheath of small arthrospores, 2-3 $\mu$	Macroconidia numerous, rough thick walls, distorted in shape, 4-14 x 3-40 $\mu$ ; microconidia numerous, pyriform, 2-4 x 3-6 $\mu$	6
None	May be sheath of small arthrospores, 2-3 $\mu$ ; more commonly large-spored ectothrix, 5-8 $\mu$ ; invasion frequently limited to hyphae inside hair	Macroconidia numerous, 4-6 cells, ellipsoid; thin rough walls, 8-12 x 30-50 $\mu$ ; microconidia few, clavate	7
None	Septate and nonseptate hyphae, air bubbles inside hair	Macroconidia numerous, 2-3 cells, pyriform-to-elliptical, thin walls, finely echinulate or smooth, 5-7 x 12-18 $\mu$ ; microconidia few	8
None	Closely septate hyphae inside hair	Macroconidia numerous, 7-10 cells, cylindro-fusiform, thick walls, densely echinulate, 10 x 60 $\mu$ ; microconidia few to abundant, pyriform-to-obovate, 4 x 9 $\mu$	9
None	Nodule on hair shaft consists of brown, dichotomously branched, closely septate hyphae, 4-8 $\mu$ diameter, and asci containing 2-8 ascospores	Dark, thick-walled, closely septate hyphae, chlamydo-spores	10
None	None	Branching, septate, irregular hyphae, with chlamydo-spores, pectinate hyphae, favic chandeliers	11
None	Ectothrix; sheath of small arthrospores, 2-3 $\mu$	Mycelium with occasional hyphal swellings; arthrospores, chlamydo-spores	12
None	Ectothrix; chains of large spores, 4-8 $\mu$	Macroconidia usually numerous, 2-10 cells, clavate, smooth and slightly thickened walls; microconidia few, small, pyriform-to-elongate	13
Branching, septate hyphae; may segment into chains of arthrospores	Ectothrix; chains of large spherical arthrospores, 6-8 $\mu$	Microconidia numerous, small pyriform to elongate; macroconidia rare, 2-8 cells, slightly clavate	14
Branching, septate hyphae; may segment into chains of arthrospores	Ectothrix; chains of small arthrospores, 3-5 $\mu$	Microconidia numerous, subspherical-to-pyriform, in terminal clusters or singly along hyphae; macroconidia clavate, 2-5 cells, thick walls, 4-6 x 10-50 $\mu$	15
Branching, septate hyphae; may segment into chains of arthrospores	Ectothrix; chains of large arthrospores, approximately 5 $\mu$	Microconidia numerous, singly along hyphae and in clusters; macroconidia infrequent, pencil-shaped thin walls, 4-6 x 10-50 $\mu$	16
Branching, septate hyphae; may segment into chains of arthrospores	Endothrix; hyphae, occasional arthrospore, and numerous air bubbles inside hair	Irregular hyphae, chlamydo-spores, hyphal swellings, pectinate hyphae, favic chandeliers	17
Rare	Endothrix; large spores in chains, 4-7.5 $\mu$	Microconidia numerous, clavate along sides of hyphae or on short conidiophores; spore-bearing hyphae stain poorly with Lacto-Phenol Cotton Blue; numerous chlamydo-spores; thin, smooth-walled macroconidia rare	18
None	Ectothrix; chains of large arthrospores, 5-10 $\mu$	Irregular hyphae, abundant chlamydo-spores; best growth at 37°C	19

*continued*



Species	Disease	Natural Occurrence in Animals Other than Man	Microscopic Appearance in Man Skin
(A)	(B)	(C)	(D)
20 <i>Trichophyton violaceum</i>	Tinea capitis, T. corporis, T. barbae, T. unguium	Cattle (rare)	Branching, septate hyphae; may segment into chains of arthrospores
21 <i>Trichosporon beigelii</i>	White piedra	Monkey	None

Contributors: (a) Halde, Carlyn, (b) Georg, Lucille K., (c) Friedman, Lorraine

References: [1] Ainsworth, G. C. 1952. Medical mycology. Pitman, New York. [2] Ainsworth, G. C., and P. K. C. Public Health Serv. Publ. 994. [4] Conant, N. F., et al. 1954. Manual of clinical mycology. W. B. Saunders, 1963. Medical mycology. Lea and Febiger, Philadelphia. [7] Kaplan, W. 1959. J. Am. Vet. Med. Assoc. 134(3): et al. 1958. An introduction to medical mycology. Yearbook, Chicago. [10] Wilson, J. W. 1957. Clinical and

Part II. DEEP

Species	Disease Produced (Synonym)	Geographical Distribution	Occurrence in Nature		Organs and Tissues of Man Frequently Attacked
			Animal Host	Saprophytic Occurrence	
(A)	(B)	(C)	(D)	(E)	(F)
1 <i>Absidia corymbifera</i> , <i>A. ramosa</i> , <i>Basidiobolus ranarum</i> , <i>Rhizopus arrhizus</i> , <i>R. oryzae</i>	Phycomycosis	Worldwide	Birds, cattle, dog, horse, swine	Soil	Lung, brain, eye, intestinal tract, sinus, cutaneous and subcutaneous tissues
2 <i>Actinomyces israelii</i>	Actinomycosis	Worldwide	Rarely cattle	Man	Cervicofacial region, lung, bone, cecum, appendix, liver
3 <i>Allescheria boydii</i> , <i>Cephalosporium</i> spp., <i>Leptosphaeria senegalensis</i> , <i>Madurella</i> spp., <i>Phialophora jeanselmei</i> , <i>Pyrenochaeta romeroi</i>	Mycetoma (maduro-mycosis, Madura foot)	Worldwide, more frequent in tropics	None	Soil ( <i>A. boydii</i> , <i>P. jeanselmei</i> )	Feet, hands, cutaneous and subcutaneous tissue, bone
4 <i>Aspergillus</i> spp.	Aspergillosis	Worldwide	Birds, cattle	Grain, soil	Ear, sinus, orbit, vagina, lung, brain
5 <i>Blastomyces dermatitidis</i>	North American blastomycosis	United States, Canada, Mexico	Dog, horse	Soil	Lung, skin, bone
6 <i>Candida albicans</i>	Candidiasis (moniliasis, thrush, mycotic vulvovaginitis)	Worldwide	Cattle, young swine, poultry	Man (often), soil (rare)	Mucous membranes, nail, skin, bronchus, lung, vagina
7 <i>Coccidioides immitis</i>	Coccidioidomycosis (coccidioidal granuloma, valley fever, Posada-Wernicke's disease)	Arid southwestern United States, Mexico, Cen. America, Chaco region and arid areas of northern S. America	Cattle, dog, horse, monkey, rodents, sheep	Soil	Lung, skin, bone, meninges

/1/ Animals are often difficult or even impossible to infect because of the great variation in susceptibility. /2/ Such

**PARASITES: MAN**

**MYCOSES**

Microscopic Appearance in Man		Microscopic Appearance of Culture on Sabouraud's Agar	
Nail	Hair		
(E)	(F)	(G)	
Branching, septate hyphae; may segment into chains of arthrospores	Endothrix; chains of large arthrospores, 4-7.5 $\mu$	Irregular hyphae, abundant chlamydospores and hyphal swellings; microconidia rare	20
None	Nodule on hair shaft consists of hyphae which segment into spherical-to-rectangular cells, 2-8 $\mu$ ; budding cells present	Hyphae segment into rectangular-to-spherical arthrospores; budding cells present	21

Austwich. 1958. Commonwealth Agr. Bur. (Gt. Brit.) Animal Health Rev. Ser. 6. [3] Ajello, L., et al. 1963. U.S. Philadelphia. [5] Dodge, C. W. 1935. Medical mycology. C. V. Mosby, St. Louis. [6] Emmons, C. W., et al. 113-117. [8] Langeron, M., and R. Vanbruseghem. 1952. Précis de mycologie. G. Masson, Paris. [9] Lewis, G., immunologic aspects of fungus disease. C. C. Thomas, Springfield.

**MYCOSES**

Susceptible Laboratory Animals <sup>1</sup>	Microscopic Appearance			
	In Human Tissue	Of Culture at 25°C	Of Culture at 37°C	
(G)	(H)	(I)	(J)	
Diabetic rabbit and rat	Nonseptate, coenocytic hyphae, 6-15 $\mu$ in width	Broad, coenocytic mycelium with sporangiophores	Similar to growth at 25°C	1
Hamster, mouse	Granules of filamentous, branching, gram-positive hyphae, 1 $\mu$ or less in width; club-shaped accretions on tips of hyphae may be present	Grows slowly	Microaerophilic-to-anaerobic, filamentous, branching, gram-positive hyphae, 1 $\mu$ or less in width; organism grows only on enriched media <sup>2</sup>	2
Mouse ( <i>A. boydii</i> )	Oval, irregular-shaped granules, 0.5-2 mm, made up of segmented, branched, hyaline or brown hyphae, 2-5 $\mu$ diameter, and chlamydospores	<i>A. boydii</i> : mycelium with oval-to-pyriiform conidia, 5-7 x 8-10 $\mu$ , borne singly at ends of long conidiophores; dark brown, thin-walled perithecia, 50-200 $\mu$ diameter, containing evanescent asci and elliptical ascospores; coremia occasionally present	Similar to cultures at 25°C	3
Birds, guinea pig, rabbit	Branching, septate hyphae	Conidiophore forms vesicle at tip; surface covered with sterigmata bearing long chains of conidia	Similar to growth at 25°C	4
Guinea pig, mouse	Single-budding, thick-walled cells, 8-15 $\mu$	Mycelium with oval-to-pyriiform conidia, 3-5 $\mu$ , on conidiophores or attached directly to hyphae	Similar to forms observed in tissue	5
Guinea pig, mouse, rabbit, rat	Oval-to-spherical budding cells, 2-4 $\mu$ ; frequently hyphae which may show clusters of blastospores attached at septations	Oval-to-spherical, single-budding cells, 2-4 $\mu$ ; pseudohyphae and hyphae; clusters of budding cells often at septations; thick-walled chlamydospores, 6-9 $\mu$ , on special medium	Similar to growth at 25°C	6
Guinea pig, hamster, mouse, other rodents	Thick-walled spherules, 20-60 $\mu$ , containing endospores, 2-5 $\mu$	Mycelium with arthrospores, 2.5-3 x 3-4 $\mu$ , alternating with empty cells	Similar to growth at 25°C; under special conditions with special media, tissue spherules obtained in vitro	7

as beef heart infusion agar at pH 6.8-7.5.

continued

	Species	Disease Produced (Synonym)	Geographical Distribution	Occurrence in Nature		Organs and Tissues of Man Frequently Attacked
				Animal Host	Saprophytic Occurrence	
	(A)	(B)	(C)	(D)	(E)	(F)
8	<i>Cryptococcus neoformans</i>	Cryptococcosis (torulosis, European blastomycosis, Busse-Buschke disease)	Worldwide	Cat, cattle, dog, horse, monkey	Soil, bird droppings	Central nervous system, lung, skin, bone
9	<i>Geotrichum candidum</i>	Geotrichosis	Worldwide	Rodents	Soil, milk products, fruit	Mouth, intestinal tract, bronchus, lung
10	<i>Histoplasma capsulatum</i>	Histoplasmosis (reticuloendothelial cytomycosis, Darling's disease)	At least 30 countries	Cat, cattle, dog, horse, mouse, rat, skunk	Soil, especially from avian and bat habitats	Lung, liver, spleen, lymph nodes, mucous membranes, adrenal, kidney
11	<i>H. duboisii</i>	African histoplasmosis	Africa	Baboon, monkey	None	Lung, skin, bone, lymph nodes, spleen, liver
12	<i>Nocardia asteroides</i> , <i>N. brasiliensis</i>	Nocardiosis	Worldwide	Cat, cattle, dog	Soil	Lung, brain, kidney, heart, spleen, liver
13	<i>N. asteroides</i> , <i>N. brasiliensis</i> , <i>Streptomyces madurae</i> , <i>S. pelletieri</i> , <i>S. somaliensis</i>	Actinomycotic mycetoma	Worldwide, more frequent in tropics	None	None	Skin, subcutaneous tissue, bone, usually on lower extremities
14	<i>Paracoccidioides brasiliensis</i>	Paracoccidioidomycosis (paracoccidioidal granuloma, Lutz-Splendore-Almeida disease)	S. & Cen. America, Mexico	None	None	Mouth, lung, lymph nodes, gastrointestinal tract
15	<i>Phialophora compactum</i> , <i>P. dermatitidis</i> , <i>P. pedrosoi</i> , <i>P. verrucosa</i> , <i>Cladosporium carrionii</i>	Chromoblastomycosis (chromomycosis, verrucous dermatitis)	Worldwide, more frequent in tropics	None	Soil, wood ( <i>P. dermatitidis</i> , <i>P. verrucosa</i> , <i>C. carrionii</i> )	Usually on lower extremities, cutaneous and subcutaneous tissue, lymphatics
16	<i>Rhinosporidium seeberi</i>	Rhinosporidiosis	Worldwide, most frequent in India and Ceylon	Cattle, horse, mule	None	Mucous membranes, nose, eye, vagina, penis, skin
17	<i>Sporotrichum schenckii</i>	Sporotrichosis	Worldwide	Cat, cattle, dog, horse, mouse, mule, rat	Mine timbers, soil, plants	Hands, feet, cutaneous and subcutaneous tissue, lymphatics

/1/ Animals are often difficult or even impossible to infect because of the great variation in susceptibility. /s/ Not resembling many of the common yeasts.

Contributors: (a) Halde, Carlyn, (b) Georg, Lucille K., (c) Friedman, Lorraine

References: [1] Ainsworth, G. C. 1952. Medical mycology. Pitman, New York. [2] Ainsworth, G. C., and P. K. C. Public Health Serv. Publ. 994. [4] Conant, N. F., et al. 1954. Manual of clinical mycology. W. B. Saunders, 1963. Medical mycology. Lea and Febiger, Philadelphia. [7] Kaplan, W. 1959. J. Am. Vet. Med. Assoc. 134(3): et al. 1958. An introduction to medical mycology. Yearbook, Chicago. [10] Wilson, J. W. 1957. Clinical and

**PARASITES: MAN**

**MYCOSES**

Susceptible Laboratory Animals <sup>1</sup>	Microscopic Appearance			
	In Human Tissue	Of Culture at 25°C	Of Culture at 37°C	
(G)	(H)	(I)	(J)	
Mouse, rat	Spherical, single-budding, thick-walled cells, 5-20 μ, surrounded by wide gelatinous capsule	Similar to cells seen in tissue <sup>2</sup> ; abortive hyphae may be seen on primary isolation	Similar to cells seen in tissue	8
None	Oblong-to-rectangular cells with somewhat rounded ends, 4-8 μ	Mycelium segments into rectangular arthrospores; germ tube forms at corners	Similar to growth at 25°C	9
Guinea pig, hamster, mouse	Intracellular, oval, budding cells, 1-5 μ	Delicate mycelium with thin-walled, subspherical-to-pyriform conidia, 2-5 μ, and thick-walled, tuberculated conidia, 8-20 μ	Budding cells, 1-5 μ; must be grown on enriched medium	10
Guinea pig, hamster, mouse	Intracellular, ovoid, budding cells, 7 x 15 μ (occasionally up to 80 μ); walls 1-2 μ thick; fat droplets	Identical to above	Budding cells, 5-15 μ	11
Guinea pig, mouse	Delicate, branched, gram-positive hyphae, 0.5-1 μ diameter; partially acid-fast	Branching hyphae, 0.5-1 μ diameter, break up readily into bacillary or coccoid forms	Similar to cultures at 25°C	12
Guinea pig, mouse	Granules of gram-positive, branching hyphae, 0.5-1 μ diameter	Branching hyphae, 0.5-1 μ diameter; spherical conidia, 0.5-1 μ, sparse to absent	Similar to cultures at 25°C	13
Guinea pig, hamster, mouse	Multiple-budding, thick-walled cells, 10-60 μ	Mycelium with rare oval conidia, 3-5 μ	Similar to forms observed in tissue	14
Mouse, rat	Single or clustered spherical, thick-walled, dark brown cells, 6-12 μ, multiply by splitting, not budding	Three types of sporulation: <i>Phialophora</i> --conidia borne within a terminal cuplike structure on a flask-shaped conidiophore; <i>Cladosporium</i> --conidia in branching chains arising terminally from conidiophore; <i>Acrotheca</i> --conidia borne acropleurogenously on swollen, clublike conidiophore	Similar to cultures at 25°C	15
None	Thick-walled spherule, 50-350 μ, with pore, containing up to 16,000 endospores, 7-9 μ	No culture method available	No culture method available	16
Hamster, mouse, rat	Rarely seen without special stains; gram-positive, cigar-shaped or spherical-to-oval, usually intracellular, budding cells, 3-5 μ; asteroid forms rare	Delicate hyphae, 2 μ in width, pyriform-to-spherical conidia, 2-4 x 2-6 μ, borne in clusters on lateral branches or laterally along hyphae	Cigar-shaped, spherical or oval budding cells; must be grown on enriched medium	17

true of some strains which may be weakly encapsulated in vitro, giving culture a different gross appearance

Austwich. 1958. Commonwealth Agr. Bur. (Gt. Brit.) Animal Health Rev. Ser. 6. [3] Ajello, L., et al. 1963. U.S. Philadelphia. [5] Dodge, C. W. 1935. Medical mycology. C. V. Mosby, St. Louis. [6] Emmons, C. W., et al. 113-117. [8] Langeron, M., and R. Vanbruseghem. 1952. Précis de mycologie. G. Masson, Paris. [9] Lewis, G., immunologic aspects of fungus disease. C. C. Thomas, Springfield.



# *Contrails*

### XIII. MATERIALS AND METHODS

#### 140. CULTURE MEDIA: PROTOZOA

##### Part I. PARASITIC AMOEBAE

Medium	Species Showing Growth
(A)	(B)
<b>Bacteria<sup>1</sup></b>	
<b>Diphasic</b>	
1 <b>Slant:</b> Coagulated whole egg. <b>Overlay:</b> Locke's, Ringer's or saline solution, alone or with one or more of the following: serum, egg white, rice (starch, flour, or powder).	<i>Dientamoeba fragilis</i> [24] <sup>2</sup> , [14] <sup>3</sup> ; <i>Endolimax nana</i> [11,24] <sup>2</sup> ; <i>Entamoeba aulostomi</i> [13] <sup>2</sup> ; <i>E. coli</i> [17] <sup>2</sup> , [23] <sup>2</sup> ; <i>E. gingivalis</i> [12] <sup>2</sup> ; <i>E. histolytica</i> [5, 11] <sup>2</sup> , [6] <sup>2</sup> ; <i>E. invadens</i> [15] <sup>2</sup> ; <i>Iodamoeba buetschlii</i> [24] <sup>2</sup>
2 <b>Slant:</b> Coagulated serum. <b>Overlay:</b> Ringer's or saline solution with serum, egg white and rice.	<i>Endolimax nana</i> , <i>Entamoeba coli</i> , <i>E. gingivalis</i> , <i>E. histolytica</i> [11] <sup>2</sup> ; <i>E. invadens</i> [16] <sup>2</sup> ; <i>E. muris</i> [21] <sup>2</sup>
3 <b>Slant:</b> Liver infusion agar. <b>Overlay:</b> Saline with serum and rice.	<i>Entamoeba histolytica</i> [7] <sup>2</sup> , [8] <sup>2</sup> ; <i>E. invadens</i> [22] <sup>2</sup>
<b>Liquid</b>	
4 Locke's, Ringer's, or saline solution with serum and rice.	<i>Entamoeba thomsoni</i> [27] <sup>2</sup> ; <i>Entamoeba barreti</i> [3] <sup>2</sup> ; <i>E. invadens</i> [22] <sup>2</sup> ; <i>E. ranarum</i> [4] <sup>2</sup>
5 Egg infusion, rice starch, with or without liver extract.	<i>Dientamoeba fragilis</i> [2] <sup>2</sup> ; <i>Entamoeba coli</i> [2] <sup>2</sup> ; <i>E. histolytica</i> [1] <sup>2</sup> ; <i>E. invadens</i> [15, 18] <sup>2</sup> ; <i>E. terrapini</i> [18] <sup>2</sup>
6 Fluid thioglycollate broth with serum <sup>5</sup> .	<i>Entamoeba coli</i> , <i>E. histolytica</i> [25] <sup>3</sup>
<b>Protozoa<sup>1</sup></b>	
7 Trypticase-dextrose broth with serum <sup>6</sup> .	<i>Entamoeba histolytica</i> [20] <sup>2</sup> ; <i>E. invadens</i> , <i>E. terrapini</i> [9] <sup>2</sup>
<b>Tissue<sup>1</sup></b>	
8 Saline, tissue slice.	<i>Entamoeba invadens</i> [19]
9 Hank's salt solution with serum and chick embryo (minced or sliced).	<i>Entamoeba histolytica</i> [26]
<b>Axenic Culture<sup>7</sup></b>	
<b>Diphasic</b>	
10 <b>Slant:</b> Tryptose, trypticase, yeast extract agar with serum. <b>Overlay:</b> Tryptose, trypticase, yeast extract broth with cell-free chick embryo extracts and vitamins.	<i>Entamoeba histolytica</i> [10]
<b>Liquid</b>	
11 Trypticase, yeast extract, maltose broth with serum.	<i>Entamoeba invadens</i> , <i>E. terrapini</i> [9]

/1/ Growth occurs in presence of one or more types of metabolizing cell: bacteria, protozoa, or metazoa.  
 /2/ Xenic growth (unknown number of associates present in culture). /3/ Monoxenic growth (one associate present).  
 /4/ Dixenic growth (two associates present). /5/ Preconditioned with a streptobacillus. /6/ Preconditioned with *Trypanosoma cruzi*. /7/ Growth occurs in absence of any other metabolizing cell.

**Contributors:** Diamond, Louis S., and Bartgis, I. Louise

**References:** [1] Balamuth, W. 1946. Am. J. Clin. Pathol. 16:380. [2] Balamuth, W. 1953. Am. J. Trop. Med. Hyg. 2:191. [3] Barret, H. P., and N. M. Smith. 1924. Am. J. Hyg. 4:155. [4] Barret, H. P., and N. M. Smith. 1926. Ann. Trop. Med. Parasitol. 20:85. [5] Boeck, W. C., and J. Drbohlav. 1925. Am. J. Hyg. 5:371. [6] Chinn, B. D., et al. 1942. Am. J. Trop. Med. 22:137. [7] Cleveland, L. R., and J. Collier. 1930. Am. J. Hyg. 12:606. [8] Cleveland, L. R., and E. P. Sanders. 1930. Science 72:149. [9] Diamond, L. S. 1960. J. Parasitol. 46:484. [10] Diamond, L. S. 1961. Science 134:336. [11] Dobell, C., and P. P. Laidlaw. 1926. Parasitology 18:283. [12] Drbohlav, J. 1925. Ann. Parasitol. Humaine Comparee 3:361. [13] Drbohlav, J. 1925. Ibid. 3:367. [14] Jacobs, L. 1953. Ann. N. Y. Acad. Sci. 56:1057. [15] McConnachie, E. W. 1955. Parasitology 45:452. [16] McConnachie, E. W. 1956. Ibid. 46:117. [17] Mayfield, M. F. 1944. Proc. Soc. Exptl. Biol. Med. 55:20. [18] Meerovitch, E. 1958. Can. J. Zool. 36:513. [19] Miller, M. J. 1953. Nature 172:1192. [20] Phillips, B. P. 1951. Am. J. Trop. Med. 31:290. [21] Pruss, J. 1959. Z. Tropenmed. Parasitol. 10:30. [22] Ratcliffe, H. L.,

*continued*

# Contrails

## 140. CULTURE MEDIA: PROTOZOA

### Part I. PARASITIC AMOEBAE

and Q. M. Geiman. 1934. Science 79:324. [23] Reardon, L. V., E. Verder, and C. W. Rees. 1952. Am. J. Trop. Med. Hyg. 1:155. [24] St. John, J. H. 1926. Am. J. Trop. Med. 6:319. [25] Shaffer, J. G., F. W. Ryden, and W. W. Frye. 1949. Am. J. Hyg. 49:127. [26] Shaffer, J. G., H. S. Sienkiewicz, and J. E. Washington. 1953. Ibid. 57:336. [27] Smith, N. M., and H. P. Barret. 1928. J. Parasitol. 14:272.

### Part II. TRICHOMONADIDAE

Medium (A)	Species Showing Growth <sup>1</sup> (B)
Agnotobiotic Culture <sup>2</sup>	
1 Egg-yolk infusion: Medium, with or without liver extract, with rice starch, 1-10% serum enrichment. For preparation, consult reference 3.	Most species can be isolated and maintained in the presence of balanced contaminants. <sup>3</sup>
Axenic Culture <sup>4</sup>	
2 Diamond's [6]: 2% trypticase, 1% yeast extract, 0.5% maltose, 0.1% L-cysteine HCl, 0.02% L-ascorbic acid, 0.08% KH <sub>2</sub> PO <sub>4</sub> , 0.08% K <sub>2</sub> HPO <sub>4</sub> , 0.05-0.10% agar, 10% inactivated sterile horse or sheep serum. Adjust to pH 6-8 with KOH, NaOH, or HCl (base or acid depends on species). Use with antibiotics for axenic isolation or retardation of bacterial growth. Autoclave 9 ml of medium 15 minutes at 15-lb pressure; add serum and antibiotics aseptically after medium has cooled. Use as soon as possible. Shelf-life 30 days at 4°C.	<i>Hypotrichomonas acosta</i> , <i>Monocercomonas</i> sp. (NS-1:PRR), <i>M. colubrorum</i> , <i>Trichomitis batrachorum</i> , <i>Tritrichomonas augusta</i> [24]; <i>Pentatrichomonas hominis</i> , <i>Tetratrichomonas gallinarum</i> , <i>Trichomonas gallinae</i> , <i>T. vaginalis</i> , <i>Tritrichomonas eberthi</i> [6]; <i>Tetratrichomonas buttrei</i> [6, 9, 15]; <i>Trichomonas rotunda</i> [15]; <i>Tritrichomonas enteris</i> [1]; <i>T. foetus</i> [9]; <i>T. suis</i> [6, 15]
3 CPLM [21]: 3.0% bacto peptone, 0.1% agar, 0.2% cysteine HCl, 0.16% maltose, 20 ml/100 liver infusion, 65 ml/100 Ringer's solution, 10% sterile serum. Prepare as above. 0.002% methylene blue may be added as an indicator. 0.1% Wilson's gastric mucin 1701X stimulatory for <i>Trichomitis batrachorum</i> and <i>Tetratrichomonas prowazeki</i> .	<i>Hypotrichomonas acosta</i> , <i>Tritrichomonas augusta</i> [23]; <i>Pentatrichomonas hominis</i> [4, 41]; <i>Tetratrichomonas buttrei</i> , <i>Trichomonas rotunda</i> , <i>Tritrichomonas suis</i> [15]; <i>Tetratrichomonas prowazeki</i> [17]; <i>Trichomitis batrachorum</i> [17]; <i>Trichomonas gallinae</i> [18]; <i>T. vaginalis</i> [21, 47]; <i>Tritrichomonas enteris</i> [1]; <i>T. foetus</i> [10]
4 B.B.L. <sup>5</sup> fluid thioglycollate <sup>6</sup> : With 5-10% sterile serum.	<i>Hypotrichomonas acosta</i> [23]; <i>Pentatrichomonas hominis</i> , <i>Trichomonas vaginalis</i> [5]; <i>Tetratrichomonas prowazeki</i> [17]; <i>Trichomonas gallinae</i> [18]
5 STS [22]: 2% trypticase, 0.15% cysteine HCl, 0.1% maltose, 0.1% agar, 5% sterile serum.	<i>Trichomonas vaginalis</i> [22]; <i>Tritrichomonas augusta</i> [34]
6 BMH [24]: 0.5% glucose, 1.0% trypticase, 0.25% yeast extract, 0.01% KH <sub>2</sub> PO <sub>4</sub> , 0.25% Na <sub>2</sub> glycerophosphate-5 H <sub>2</sub> O, 0.004% Ca pantothenate, 0.005% cholesterol, 0.0001% TEM 4T, 0.1% agar, 0.04% ascorbic acid, 0.05% thiomalic acid, 0.004% trace-metals mixture #50, 1.0 ml/100 vitamin mixture #12. Grow newly inoculated cultures for 2 days at 25°C, then place at 15°C for 1-2 months.	<i>Hypotrichomonas acosta</i> , <i>Monocercomonas</i> sp. (NS-1:PRR), <i>M. colubrorum</i> , <i>Trichomitis batrachorum</i> , <i>Tritrichomonas augusta</i> [24]

/1/ Species isolated from homeotherms: *Pentatrichomonas hominis*, *Tetratrichomonas buttrei*, *T. gallinarum*, *Trichomonas gallinae*, *T. tenax*, *T. vaginalis*, *Tritrichomonas eberthi*, *T. enteris*, *T. foetus*, *T. suis*. Species isolated from poikilotherms: *Hypotrichomonas acosta*, *Monocercomonas* sp. (NS-1:PRR), *M. colubrorum*, *Tetratrichomonas prowazeki*, *Trichomitis batrachorum*, *Tritrichomonas augusta*. /2/ Many media for parasitic amoeba also support agnotobiotic trichomonad cultures [44, 45] (see Part I). /3/ The following have been cultured agnotobiotically (in the presence of unknown other organisms): *Metatrichomonas termopsidis*, *Monocercomonas verrens*, *Tetratrichomonas limacis*, *T. microti*, *T. ovis*, *Trichomitis marmolae*. /4/ Because these media support high bacterial populations, antibiotics are necessary to retard bacterial overgrowth or for axenic isolation. The following antibiotic combinations have been successful for axenic isolation: 10,000 units/ml Na or K penicillin, 1,000 µg/ml streptomycin [6]; 2,000 µg/ml dehydrostreptomycin, 250 µg/ml chloroamphenicol, 60 µg/ml polymyxin B [23, 24]; for molds and yeasts, 300 µg/ml nystatin [16]; see also references 17, 38. /5/ Baltimore Biological Laboratory, Baltimore 18, Md. /6/ Or without indicator, Brewer-modified.

continued

## 140. CULTURE MEDIA: PROTOZOA

### Part II. TRICHOMONADIDAE

Medium (A)	Species Showing Growth <sup>1</sup> (B)
Special Purpose	
7 Complex chemically better defined media [39]: Mixtures of salts, amino acids, nucleotides, lipids, trace metals, vitamins, and one or more poorly defined complex natural organic substances.	<i>Hypotrichomonas acosta</i> [23]; <i>Monocercomonas</i> sp. (NS-1:PRR), <i>M. colubrurum</i> , <i>Tritrichomonas augusta</i> [24]; <i>Trichomonas gallinae</i> [40]; <i>T. vaginalis</i> [42]
8 Complex medium for axenic <i>Trichomonas tenax</i> .	<i>Trichomonas tenax</i> [7]
9 Media and techniques for freezing cultures.	<i>Pentatrichomonas hominis</i> [8, 31]; <i>Trichomonas gallinae</i> [8, 19, 31]; <i>T. vaginalis</i> [8, 19, 32]; <i>Tritrichomonas foetus</i> [13, 14, 25-29, 31, 33]
10 Solid media for cloning and drug testing.	<i>Pentatrichomonas hominis</i> , <i>Tetratrichomonas gallinarum</i> , <i>Tritrichomonas augusta</i> , <i>T. suis</i> [35]; <i>Trichomonas gallinae</i> [2, 35]; <i>T. vaginalis</i> [12, 20, 35, 37]; <i>Tritrichomonas foetus</i> [35, 46]
11 Tissue culture.	<i>Trichomonas gallinae</i> , <i>T. vaginalis</i> [18]
12 Bulk growth or continuous flow culture.	<i>Pentatrichomonas hominis</i> , <i>Trichomonas vaginalis</i> [11, 30, 36]; <i>Tritrichomonas augusta</i> [36, 43]

<sup>1/1</sup> Species isolated from homeotherms: *Pentatrichomonas hominis*, *Tetratrichomonas buttreysi*, *T. gallinarum*, *Trichomonas gallinae*, *T. tenax*, *T. vaginalis*, *Tritrichomonas eberthi*, *T. enteris*, *T. foetus*, *T. suis*. Species isolated from poikilotherms: *Hypotrichomonas acosta*, *Monocercomonas* sp. (NS-1:PRR), *M. colubrurum*, *Tetratrichomonas prowazeki*, *Trichomitis batrachorum*, *Tritrichomonas augusta*.

**Contributor:** Lee, John J.

**References:** [1] Anderson, F. L., and N. D. Levine. 1962. *J. Protozool.* 9(Suppl.):18. [2] Asami, K., Y. Nodake, and T. Ueno. 1955. *Exptl. Parasitol.* 4:34. [3] Balamuth, W., and J. G. Sandza. 1944. *Proc. Soc. Exptl. Biol. Med.* 57:161. [4] De Carneri, I. 1955. *Nature* 176:605. [5] De Carneri, I. 1956. *Riv. Parassitol.* 17:247. [6] Diamond, L. S. 1957. *J. Parasitol.* 43:488. [7] Diamond, L. S. 1960. *Ibid.* 46:43. [8] Diamond, L. S. 1962. *J. Protozool.* 9:442. [9] Doran, D. J. 1957. *Ibid.* 4:182. [10] Doran, D. J. 1958. *Ibid.* 5:89. [11] Feinberg, F. G. 1953. *Nature* 171:1165. [12] Filadoro, F., and N. Orsi. 1958. *Antibiot. Chemotherapy* 8:561. [13] Fitzgerald, P. R., and N. D. Levine. 1957. *J. Protozool.* 4(Suppl.):5. [14] Fitzgerald, P. R., and N. D. Levine. 1961. *Ibid.* 8:21. [15] Hibler, C. P., et al. 1960. *J. Protozool.* 7:159. [16] Honigberg, B. M. 1957. *J. Parasitol.* 43:43. [17] Honigberg, B. M. 1958. *J. Protozool.* 5(Suppl.):15. [18] Honigberg, B. M. 1961. *Abstr. 1st Intern. Conf. Protozool., Prague*, p. 62. [19] Honigberg, B. M., and V. M. King. 1962. *J. Protozool.* 9(Suppl.):18. [20] Ivey, M. H. 1961. *J. Parasitol.* 47:539. [21] Johnson, G., and M. Trussell. 1943. *Proc. Soc. Exptl. Biol. Med.* 54:245. [22] Kupferberg, A. B., et al. 1953. *Ann. N. Y. Acad. Sci.* 56:1006. [23] Lee, J. J., and S. Pierce. 1960. *J. Protozool.* 7:402. [24] Lee, J. J., et al. 1962. *Ibid.* 9:445. [25] Levine, N. D., W. E. McCaul, and M. Mizell. 1957. *Ibid.* 4(Suppl.):5. [26] Levine, N. D., and W. C. Marquart. 1954. *Ibid.* 1(Suppl.):4. [27] Levine, N. D., and W. C. Marquart. 1955. *Ibid.* 2:100. [28] Levine, N. D., M. Mizell, and D. A. Houlahan. 1958. *Exptl. Parasitol.* 7:236. [29] Levine, N. D., et al. 1962. *J. Protozool.* 9:347. [30] McEntegart, M. G. 1952. *J. Clin. Pathol.* 5:275. [31] McEntegart, M. G. 1954. *J. Hyg.* 52:545. [32] McEntegart, M. G. 1959. *Nature* 183:270. [33] McWade, D. M., and J. A. Williams. 1954. *Mich. State Univ. Agr. Expt. Sta. Quart. Bull.* 37:248. [34] Samuels, R. 1958. *J. Protozool.* 5(Suppl.):9. [35] Samuels, R. 1962. *Ibid.* 9:103. [36] Samuels, R., and E. A. Beil. 1963. *Ibid.* 9(Suppl.):19. [37] Samuels, R., and D. J. Stouder. 1960. *Ibid.* 7:5. [38] Seneca, H., and D. Ides. 1953. *Am. J. Trop. Med. Hyg.* 6:1045. [39] Shorb, M. In press, 1964. In S. H. Hutner, ed. *Nutrition and biochemistry of protozoa*. Academic Press, New York. v. 3. [40] Shorb, M., and P. G. Lund. 1959. *J. Protozool.* 6:122. [41] Solomon, J. M. 1957. *J. Parasitol.* 43:39. [42] Sprince, H., and A. B. Kupferberg. 1947. *J. Bacteriol.* 53:435. [43] Twohy, D. W., and P. A. Tucker. 1961. *J. Protozool.* 8(Suppl.):5. [44] Wenrich, D. H. 1945. *J. Parasitol.* 31:375. [45] Wenrich, D. H. 1946. *Ibid.* 32:40. [46] West, R. A., et al. 1962. *J. Protozool.* 9:65. [47] Wirtschafter, S. K. 1954. *J. Parasitol.* 40:100.

continued



## 140. CULTURE MEDIA: PROTOZOA

### Part III. TRYPANOSOMATIDAE

Medium (A)	Species Showing Growth <sup>2</sup> (B)
Blood Agar <sup>2-5</sup>	
1 <b>Solid phase:</b> 14 g agar, 6 g NaCl, 450 ml defibrinated rabbit blood, 900 ml distilled H <sub>2</sub> O. For variation, consult reference 20.	<i>Leishmania enrietti</i> [4]; <i>L. donovani</i> , <i>L. tropica</i> [27]; <i>L. tarentolae</i> [42]; <i>Trypanosoma ambystomae</i> [16]; <i>T. brucei</i> , <i>T. lewisi</i> [20]; <i>T. cruzi</i> , <i>T. duttoni</i> , <i>T. melophagium</i> , <i>T. rotatorium</i> , <i>T. theileri</i> , 2 species of bird trypanosomes [30]; <i>T. striati</i> [32]
2 <b>Solid phase:</b> 10-15 g agar, 10 g glucose, 1,000 ml horse meat broth, 1,000 ml defibrinated horse blood.	<i>Crithidia melophagia</i> , <i>Leptomonas ctenocephali</i> , <i>L. fasciculata</i> , <i>Trypanosoma cruzi</i> , <i>T. rotatorium</i> , <i>T. syrniai</i> , <i>T. theileri</i> [28]; <i>Leishmania braziliensis</i> [33]; <i>L. donovani</i> [28, 33]; <i>L. tropica</i> [22, 33]; <i>Strigomonas oncopelti</i> [22]; <i>Trypanosoma conorhini</i> [7]
3 <b>Solid phase:</b> 50 g bacto beef, 20 g neopeptone, 5 g NaCl, 20 g bacto agar, 100-150 ml defibrinated human or rabbit blood. Variation: With Locke's solution overlay [14, 39] <sup>5</sup> .	<i>Leishmania braziliensis</i> , <i>L. donovani</i> , <i>L. tropica</i> , <i>Trypanosoma conorhini</i> , <i>T. lewisi</i> , <i>T. pipistrelli</i> [39]; <i>T. cruzi</i> [36, 39]; <i>T. rangeli</i> [38]
4 <b>Solid phase:</b> 31 g nutrient agar, 5 g plain agar, 167 ml inactivated human plasma, 167 ml washed human red cells.	<i>Trypanosoma gambiense</i> , <i>T. rhodesiense</i> [46]
5 <b>Solid phase:</b> 20 g agar, 5 g dextrose, 7 g NaCl, 20 g proteose peptone, 100 ml defibrinated rabbit blood. <b>Liquid phase:</b> Infusion broth containing 2% proteose peptone and 0.5% dextrose. For variation, consult reference 9.	<i>Trypanosoma cruzi</i> [6, 9]; <i>T. lewisi</i> , bird trypanosomes, frog trypanosomes [9]
6 <b>Solid phase:</b> 3 g bacto beef, 5 g bacto peptone, 8 g NaCl, 15 g bacto agar, 333 ml citrated human or rabbit blood. <b>Liquid phase:</b> Locke's solution.	<i>Trypanosoma congolense</i> [37]; <i>T. gambiense</i> , <i>T. rhodesiense</i> [41]
Semi-Solid <sup>3,4,7</sup>	
7 1 part 3% agar, 8 parts Locke's solution with 0.2% glucose, 1 part rabbit serum.	<i>Herpetomonas muscidarum</i> , <i>Leishmania agamiae</i> , <i>L. ceramodactyli</i> , <i>L. donovani</i> , <i>L. tarentolae</i> , <i>L. tropica</i> , <i>Strigomonas oncopelti</i> , <i>Trypanosoma cruzi</i> , <i>T. ptyodactyli</i> , <i>T. rabinowitschi</i> [1]
8 3 g agar, 150 ml defibrinated rabbit blood, 1,000 ml normal saline.	<i>Leishmania donovani</i> , <i>L. tropica</i> [19]
9 Mixture of 21 amino acids, 3 salts, 10 vitamins, glucose, guanosine, adenine SO <sub>4</sub> , uracil, uric acid, urea, creatine, creatinine, nucleic acid, 0.2% agar, heat-coagulated red blood cells.	<i>Trypanosoma cruzi</i> [17]
Liquid <sup>3,4,7</sup>	
10 0.5 ml human or monkey blood, 0.5 ml 2% sodium citrate in 0.85% NaCl solution, 1 ml Ringer's solution (with 0.6% NaCl). For variation, consult reference 3.	<i>Trypanosoma brucei</i> [3]; <i>T. congolense</i> , <i>T. gambiense</i> [3, 34]; <i>T. cruzi</i> [34]
11 Overlay from item 3 for trypanosomes of the <i>lewisi</i> group and from item 6 for African trypanosomes.	<i>Trypanosoma congolense</i> [45]; <i>T. cruzi</i> [44]; <i>T. gambiense</i> , <i>T. rhodesiense</i> [41]
12 10 ml 5% lactoalbumin hydrolysate in Earle's saline, 5 ml filtered and unheated calf serum, 5 ml red cell lysate, 100 ml 0.1% glucose in Earle's saline.	<i>Trypanosoma gambiense</i> [26]
Dialysate <sup>3,7</sup>	
13 Cellophane loop filled with Locke's solution suspended in tubes of diphasic blood agar. Variation: Loop suspended in blood-coagulum-peptone medium [10].	<i>Trypanosoma cruzi</i> [10, 40]
Defined and Partially Defined <sup>3,4,7</sup>	
14 Mixture of amino acids, salts (including trace metals), glucose, purines, pyrimidines, vitamins, growth factors, and hemin. Variation: Only methionine as amino acid and no hemin [25].	<i>Herpetomonas culicidarum</i> [5]; <i>Leishmania tarentolae</i> [42]; <i>Strigomonas oncopelti</i> [25]

/1/ Use of the generic names *Crithidia*, *Herpetomonas*, *Leptomonas*, and *Strigomonas* is not yet uniform. /2/ Test-tube cultures usually contain 5 ml base; flask or plate cultures contain varying amounts of base, depending on size of container. /3/ Cultures usually maintained at 22°-25°C. /4/ Ingredients are given in amounts to be added to one liter of distilled water, unless otherwise specified. /5/ Diphasic test-tube cultures receive 2-3 ml overlay; flask cultures approximately 15 ml for each 25 ml base. /6/ 10% blood for *Trypanosoma cruzi* and *Leishmania* spp.; 30% for other species and all isolations. /7/ Varying amounts of media are used, depending on size of container.

continued

## 140. CULTURE MEDIA: PROTOZOA

### Part III. TRYPANOSOMATIDAE

	Medium (A)	Species Showing Growth (B)
	Defined and Partially Defined <sup>3,4,7</sup>	
15	Item 9 without agar.	<i>Trypanosoma cruzi</i> [17]
16	15 g bacto tryptose, 2 g glucose, 1 mg thiamine, 3 mg folic acid, 20 mg hemin, 25 mg sodium stearate, 4 g NaCl, 5 g Na <sub>3</sub> PO <sub>4</sub> ·12H <sub>2</sub> O, 0.4 g KCl, 1,000 ml twice-distilled H <sub>2</sub> O.	<i>Trypanosoma cruzi</i> [2]
	Avian Embryo <sup>8</sup>	
17	Chorioallantoic membrane.	<i>Leishmania donovani</i> , <i>Trypanosoma gambiense</i> [35]; <i>Leishmania tropica</i> [29]; <i>Trypanosoma brucei</i> , <i>T. evansi</i> [18, 35]; <i>T. cruzi</i> [11, 31]; <i>T. equiperdum</i> , <i>T. rhodesiense</i> [18]
18	Intra-yolk sac.	<i>Leishmania braziliensis</i> , <i>L. donovani</i> , <i>L. tropica</i> [15]; <i>Trypanosoma brucei</i> , <i>T. equiperdum</i> , <i>T. evansi</i> [23]; <i>T. cruzi</i> [21]
	Tissue Culture <sup>9</sup>	
19	Mammalian tissues in nutrient fluid.	<i>Leishmania donovani</i> [13]; <i>Trypanosoma conorhini</i> [7]; <i>T. cruzi</i> [12, 31]; <i>T. gambiense</i> , <i>T. rhodesiense</i> [8]
20	Avian tissues in nutrient fluid.	<i>Trypanosoma cruzi</i> [24]
21	Insect tissues in nutrient fluid.	<i>Trypanosoma brucei</i> , <i>T. congolense</i> , <i>T. vivax</i> [43]

/3/ Cultures usually maintained at 22°-25°C. /4/ Ingredients are given in amounts to be added to one liter of distilled water, unless otherwise specified. /7/ Varying amounts of media are used, depending on size of container. /8/ Cultures usually maintained at 25°-35°C. /9/ Cultures usually maintained at 25°-38°C.

**Contributors:** Tobie, Eleanor J., and von Brand, Theodor

**References:** [1] Adler, S. 1934. Trans. Roy. Soc. Trop. Med. Hyg. 28:201. [2] Boné, G. J., and G. Parent. 1963. J. Gen. Microbiol. 31:261. [3] Brutsaert, P., and C. Henrard. 1938. Compt. Rend. Soc. Biol. 127:1469. [4] Coutinho, J. O. 1955. Folia Clin. Biol. (Sao Paulo) 23:91. [5] Cowperthwaite, J., et al. 1953. Ann. N.Y. Acad. Sci. 56:972. [6] Davis, D. J. 1943. Public Health Rept. (U.S.) 58:775. [7] Deane, M. P., and L. M. Deane. 1961. Rev. Inst. Med. Trop. Sao Paulo 3:149. [8] Demarchi, J., and J. Nicoli. 1960. Ann. Inst. Pasteur 99:120. [9] Diamond, L. S., and C. M. Herman. 1954. J. Parasitol. 40:195. [10] Eife, E. H., and J. F. Kent. 1960. Am. J. Trop. Med. Hyg. 9:512. [11] Ganapati, P. N. 1948. Nature 162:963. [12] Hawking, F. 1946. Trans. Roy. Soc. Trop. Med. Hyg. 40:345. [13] Hawking, F. 1948. Ibid. 41:545. [14] Johnson, E. M. 1947. J. Parasitol. 33:85. [15] Jones, H., G. Rake, and D. Hamre. 1944. Am. J. Trop. Med. 24:381. [16] Lehmann, D. L. 1955. J. Protozool. 2:28. [17] Little, P. A., and J. J. Oleson. 1951. J. Bacteriol. 61:709. [18] Longley, J., N. M. Clausen, and A. L. Tatum. 1939. Proc. Soc. Exptl. Biol. Med. 41:365. [19] Lourie, E. M. 1946. Trans. Roy. Soc. Trop. Med. Hyg. 40:4. [20] MacNeal, W. J. 1904. J. Infect. Diseases 1:517. [21] Manso Soto, A. E., G. A. Loretto, and J. A. Rispoli. 1950. Mision Estud. Patol. Reg. Arg. 21:23. [22] Mayer, M., and B. Malamos. 1936. Zentr. Bakteriolog. Parasitenk., I, 136:412. [23] Merchant, D. J. 1947. Proc. Soc. Exptl. Biol. Med. 64:391. [24] Meyer, H., and M. Xavier de Oliveira. 1948. Parasitology 39:91. [25] Newton, B. A. 1956. Nature 177:279. [26] Nicoli, J. 1961. Bull. Soc. Pathol. Exotique 54:77. [27] Nicolle, C. 1908. Compt. Rend. 146:842. [28] Nöller, W. 1917. Arch. Schiffs- Tropen-Hyg. 21:53. [29] Oberling, C., and N. Ansari. 1951. Bull. Soc. Pathol. Exotique 44:542. [30] Packchanian, A. 1934. Science 80:407. [31] Pipkin, A. C. 1960. Exptl. Parasitol. 9:167. [32] Qadri, S. S. 1962. Parasitology 52:229. [33] Ray, J. C. 1932. Indian J. Med. Res. 20:355. [34] Reichenow, E. 1934. Arch. Schiffs- Tropen-Hyg. 38:292. [35] Rodhain, J., and L. van den Berghe. 1943. Ann. Soc. Belge Med. Trop. 23:141. [36] Senekjic, H. A. 1943. Am. J. Trop. Med. 23:523. [37] Tobie, E. J. 1958. J. Parasitol. 44:241. [38] Tobie, E. J. 1961. Exptl. Parasitol. 11:1. [39] Tobie, E. J. Unpublished. Natl. Institutes of Health,

continued

## 140. CULTURE MEDIA: PROTOZOA

### Part III. TRYPANOSOMATIDAE

Bethesda, Md., 1963. [40] Tobie, E. J., and C. W. Rees. 1948. J. Parasitol. 34:162. [41] Tobie, E. J., T. von Brand, and B. Mehlman. 1950. Ibid. 36:48. [42] Trager, W. 1957. J. Protozool. 4:269. [43] Trager, W. 1959. Nature 184:30. [44] von Brand, T., E. M. Johnson, and C. W. Rees. 1946. J. Gen. Physiol. 30:163. [45] von Brand, T., and E. J. Tobie. 1959. J. Parasitol. 45:204. [46] Weinman, D. 1960. Trans. Roy. Soc. Trop. Med. Hyg. 54:180.

### Part IV. PHYTOMASTIGINA

Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>
(A)	(B)		(A)	(B)		(A)	(B)	
<b>Marine Flagellates<sup>2</sup> [8]</b>			33 Ca	50	68 L-Glutamic acid	3,000		
1 Ca, as Cl	400	34 Co	0.5	69 Glycine	2,500			
2 MgCl <sub>2</sub> ·6H <sub>2</sub> O	4,000	35 Cu	2	70 Cyanocobalamin	0.0002			
3 MgSO <sub>4</sub> ·7H <sub>2</sub> O	7,000	36 Fe	10	71 Thiamine HCl	0.6			
4 KCl	700	37 MgSO <sub>4</sub> ·7H <sub>2</sub> O	500	72	pH = 3-6			
5 K <sub>3</sub> PO <sub>4</sub>	10	38 Mn	4	<b>Ochromonas spp.<sup>3</sup> [1]</b>				
6 NaCl	28,000	39 Mo	4	73 NH <sub>4</sub> Cl	500 <sup>10</sup> ; 400 <sup>11</sup>			
7 NaNO <sub>3</sub>	100	40 K <sub>2</sub> HPO <sub>4</sub>	250	74 B, as H <sub>3</sub> BO <sub>3</sub>	0.1			
8 PII metals <sup>3</sup>	10 ml	41 Zn	20	75 CaCO <sub>3</sub>	50 <sup>10</sup> ; 150 <sup>11</sup>			
9 SII metals <sup>4</sup>	10 ml	42 Ammonium acetate	1,000	76 Co, as SO <sub>4</sub>	0.1			
10 Sodium glycerophosphate	10	43 EDTA <sup>5</sup>	200	77 Cu, as SO <sub>4</sub>	0.08			
11 Sodium metasilicate·9H <sub>2</sub> O	150	44 Glycine	2,000	78 Fe, as SO <sub>4</sub>	2			
12 Nitritotriacetic acid	100	45	pH = 7.5	79 MgCO <sub>3</sub> (basic)	400 <sup>10</sup> ; 500 <sup>11</sup>			
13 Tris <sup>6</sup>	1,000	<b>Chlorogonium spp.<sup>7</sup> [6]</b>		80 MgSO <sub>4</sub> ·7H <sub>2</sub> O	1,000 <sup>11</sup>			
14 Biotin	1 μg	46 NH <sub>4</sub> NO <sub>3</sub>	500	81 Mn, as SO <sub>4</sub>	0.5			
15 Cyanocobalamin	0.2 μg	47 FeCl <sub>3</sub> ·6H <sub>2</sub> O	2.5	82 Mo, as (NH <sub>4</sub> ) <sub>6</sub> Mo <sub>7</sub> -O <sub>24</sub> ·H <sub>2</sub> O	0.35			
16 Thiamine HCl	100 μg	48 MgSO <sub>4</sub> ·7H <sub>2</sub> O	100	83 KH <sub>2</sub> PO <sub>4</sub>	300			
17	pH = 7.8-8.0	49 MnCl <sub>2</sub> ·4H <sub>2</sub> O	0.1	84 V, as Na <sub>3</sub> VO <sub>4</sub> ·16H <sub>2</sub> O	0.01			
<b>Chilomonas paramecium [3]</b>		50 KH <sub>2</sub> PO <sub>4</sub>	500	85 Zn, as SO <sub>4</sub>	1			
18 NH <sub>4</sub> Cl	200	51 NaCl	100	86 Ammonium citrate	1,200 <sup>10</sup>			
19 H <sub>3</sub> BO <sub>3</sub>	115	52	pH = 7.0	87 Nitritotriacetic acid	200 <sup>10</sup> ; 300 <sup>11</sup>			
20 CaCl <sub>2</sub>	55	<b>Euglena gracilis [4]</b>		88 Glucose	10,000			
21 CoSO <sub>4</sub> ·7H <sub>2</sub> O	19	53 B, as H <sub>3</sub> BO <sub>3</sub>	0.1	89 L-Arginine HCl	400 <sup>10</sup> ; 500 <sup>11</sup>			
22 CuSO <sub>4</sub> ·5H <sub>2</sub> O	15.7	54 CaCO <sub>3</sub>	80	90 L-Glutamic acid	10,000 <sup>10</sup> ; 3,000 <sup>11</sup>			
23 FeSO <sub>4</sub> ·7H <sub>2</sub> O	40	55 Co, as SO <sub>4</sub>	0.1	91 Glycine	100 <sup>10</sup>			
24 MgSO <sub>4</sub> ·7H <sub>2</sub> O	800	56 Cu, as SO <sub>4</sub>	0.08	92 L-Histidine HCl	400 <sup>10</sup> ; 500 <sup>11</sup>			
25 MnSO <sub>4</sub> ·4H <sub>2</sub> O	81	57 Fe, as SO <sub>4</sub>	2	93 DL-Methionine	600 <sup>10</sup>			
26 K <sub>2</sub> HPO <sub>4</sub>	200	58 MgSO <sub>4</sub> ·7H <sub>2</sub> O	400	94 Biotin	10 μg <sup>10</sup> ; 4 μg <sup>11</sup>			
27 Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	15	59 Mn, as SO <sub>4</sub>	0.5	95 Cyanocobalamin	1 μg <sup>10</sup>			
28 ZnSO <sub>4</sub> ·7H <sub>2</sub> O	220	60 Mo, as (NH <sub>4</sub> ) <sub>6</sub> Mo <sub>7</sub> -O <sub>24</sub> ·4H <sub>2</sub> O	0.35	96 Thiamine HCl	1 <sup>10</sup> ; 2 <sup>11</sup>			
29 EDTA <sup>5</sup>	500	61 KH <sub>2</sub> PO <sub>4</sub>	300	97	pH = 5.0			
30 Thiamine HCl	10 μg	62 V, as Na <sub>3</sub> VO <sub>4</sub> ·16H <sub>2</sub> O	0.01	<b>Polytoma uvella<sup>11</sup> [2]</b>				
31	pH = 3.5-7.5	63 Zn, as SO <sub>4</sub>	1	98 NH <sub>4</sub> Cl	500			
<b>Chlamydomonas moewusii [5]</b>		64 Ammonium succinate	600	99 H <sub>3</sub> BO <sub>3</sub>	120			
32 B	20	65 Sucrose	15,000	100 CaCl <sub>2</sub>	60			
		66 DL-Malic acid	1,000					
		67 DL-Aspartic acid	2,000					

[1] Unless otherwise specified. [2] Chrysoomonads, cryptomonads, dinoflagellates, and also diatoms. [3] 1 ml of PII metals contains 1 mg ethylenediamine tetra-acetic acid, 0.01 mg Fe (as Cl), 0.2 mg B (as H<sub>3</sub>BO<sub>3</sub>), 0.04 mg Mn (as Cl), 0.005 mg Zn (as Cl), 0.001 mg Co (as Cl). [4] 1 ml of SII metals contains 1.0 mg Br (as Na), 0.2 mg Sr (as Cl), 0.02 mg Rb (as Cl), 0.02 mg Li (as Cl), 0.001 mg I (as K), 0.05 mg Mo (as Na). [5] Tris(hydroxymethyl)-aminomethane. [6] Ethylenediamine tetra-acetic acid. [7] *C. elongatum* and *C. euchlorum*. [8] *O. danica* and *O. malhamensis*. [9] *O. danica*. [10] *O. malhamensis*. [11] Other *Polytoma* species might grow in the same medium if thiamine HCl were added at 100 μg/L.

continued



**140. CULTURE MEDIA: PROTOZOA**

**Part IV. PHYTOMASTIGINA**

Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>
(A)	(B)		(A)	(B)	(A)	(B)		
<i>Polytoma uella</i> <sup>11</sup> [2]			111 Tris <sup>12</sup>	1,210	120 Ca, as Cl		4	
101 CoSO <sub>4</sub> ·7H <sub>2</sub> O	20		112	pH = 8.0	121 Fe, as Cl		0.5	
102 CuSO <sub>4</sub> ·5H <sub>2</sub> O	13	<i>Polytomella caeca</i> <sup>12</sup> [7]			122 Mg, as Cl		0.5	
103 FeSO <sub>4</sub> ·7H <sub>2</sub> O	40	113 MgSO <sub>4</sub> ·7H <sub>2</sub> O	100	123 Mn, as Cl		0.01		
104 MgSO <sub>4</sub> ·7H <sub>2</sub> O	160	114 KH <sub>2</sub> PO <sub>4</sub>	500	124 K, as Cl		2		
105 MnSO <sub>4</sub> ·4H <sub>2</sub> O	50	115 NaCl	100	125 Sodium citrate·H <sub>2</sub> O		20		
106 K <sub>2</sub> HPO <sub>4</sub>	40	116 Ammonium acetate <sup>13</sup>	2,000	126 Sodium glycerophosphate·5H <sub>2</sub> O		50		
107 Na <sub>2</sub> MoO <sub>4</sub> ·7H <sub>2</sub> O	15	117 Thiamine	0.3-1.0	127 Sodium metasilicate·9H <sub>2</sub> O		30		
108 ZnSO <sub>4</sub> ·7H <sub>2</sub> O	220	118	pH = 6.5	128 L-Histidine, free base		200		
109 Sodium acetate·3H <sub>2</sub> O	2,720	<i>Synura</i> spp. <sup>14</sup> [9]		129 Cyanocobalamin		0.4 µg		
110 EDTA <sup>14</sup>	100	119 (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	60	130		pH = 6.0		

<sup>11/</sup> Unless otherwise specified. <sup>12/</sup> Tris(hydroxymethyl)aminomethane. <sup>13/</sup> Ethylenediamine tetra-acetic acid. <sup>14/</sup> Other *Polytoma* species might grow in the same medium if thiamine HCl is added at 100 µg/L. <sup>12/</sup> Sterilize medium, then add CaCl<sub>2</sub> and FeC<sub>6</sub>H<sub>5</sub>O<sub>7</sub>·3H<sub>2</sub>O sufficient to give a final concentration of 10 mg of each per liter. <sup>13/</sup> May be substituted with NH<sub>4</sub>Cl and n-butanol (1 ml/L). <sup>14/</sup> *S. caroliniana* and *S. petersenii*.

**Contributor:** Provasoli, Luigi

**References:** [1] Aaronson, S., and H. Baker. 1959. J. Protozool. 6:282. [2] Cirillo, V. P. 1955. Proc. Soc. Exptl. Biol. Med. 88:352. [3] Holz, G. G. 1954. J. Protozool. 1:114. [4] Hutner, S. H., and M. K. Bach. 1955. Ibid. 3:101. [5] Hutner, S. H., et al. 1950. Proc. Am. Phil. Soc. 94:152. [6] Loefer, J. B. 1934. Biol. Bull. 66:1. [7] Lwoff, A. 1941. Ann. Inst. Pasteur 66:407. [8] Provasoli, L. 1961. In H. Iwasaki. Biol. Bull. 121:176. [9] Provasoli, L., and I. J. Pintner. 1960. Pymatuning Symp. Ecol. Publ. 2.

**141. CULTURE MEDIA: ANIMAL TISSUES**

**Part I. BALANCED SALT SOLUTIONS**

In general, these diluents are used only in combination with naturally occurring body substances (e.g., blood serum, tissue extracts), and/or with more complex, chemically defined, feeding solutions. pH of the final medium must be regulated. Inclusion of the names of commercial suppliers in no way implies endorsement by the Federation of American Societies for Experimental Biology.

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)	(B)		(A)	(B)	(A)	(B)		
Ringer (mammalian) [9]			Locke <sup>2</sup> [6]		15 NaHCO <sub>3</sub>		200	
1 CaCl <sub>2</sub>	250	8 CaCl <sub>2</sub>	240	16 NaCl		9,500		
2 KCl	420	9 KCl	420	17 Glucose		1,000		
3 NaCl	9,000	10 NaHCO <sub>3</sub>	300	Tyrode <sup>2,3</sup> [10]				
Ringer (amphibian) [8]			11 NaCl	9,000	18 CaCl <sub>2</sub>		200	
4 CaCl <sub>2</sub>	120	12 Glucose	1,000	19 MgCl <sub>2</sub> ·6H <sub>2</sub> O		100 <sup>4</sup>		
5 KCl	140	Locke <sup>1</sup> [6]		20 KCl		200		
6 NaHCO <sub>3</sub>	200	13 CaCl <sub>2</sub>	200	21 NaHCO <sub>3</sub>		1,000		
7 NaCl	6,500	14 KCl	200	22 NaCl		8,000		
				23 NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O		50		
				24 Glucose		1,000		

<sup>1/</sup> One of several solutions described by Locke. <sup>2/</sup> Available commercially from Colorado Serum Co., 4950 York Street, Denver 16, Colo. <sup>3/</sup> Available commercially from Difco Laboratories, 920 Henry Street, Detroit 1, Mich. <sup>4/</sup> Or may be 214 mg/L; degree of hydration not reported.

*continued*



## 141. CULTURE MEDIA: ANIMAL TISSUES

### Part I. BALANCED SALT SOLUTIONS

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
Gey (for tubes) <sup>2,3,5-9</sup> [3]			46	NaHCO <sub>3</sub>	2,200	67	MgSO <sub>4</sub> ·7H <sub>2</sub> O	154
25	CaCl <sub>2</sub>	170	47	NaCl	6,800	68	KCl	285
26	MgCl <sub>2</sub> ·6H <sub>2</sub> O	210	48	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140	69	KH <sub>2</sub> PO <sub>4</sub>	83
27	MgSO <sub>4</sub> ·7H <sub>2</sub> O	70	49	Glucose	1,000	70	NaHCO <sub>3</sub>	1,200
28	KCl	370	Hanks <sup>2,3,5-10</sup> [4]			71	NaCl	7,400
29	KH <sub>2</sub> PO <sub>4</sub>	30	50	CaCl <sub>2</sub>	200	72	Na <sub>2</sub> HPO <sub>4</sub> ·7H <sub>2</sub> O	290
30	NaHCO <sub>3</sub>	2,270	51	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	73	Glucose	1,100
31	NaCl	7,000	52	KCl	400	Puck (Saline G) <sup>3</sup> [7]		
32	Na <sub>2</sub> HPO <sub>4</sub> ·2H <sub>2</sub> O	150	53	KH <sub>2</sub> PO <sub>4</sub>	100	74	CaCl <sub>2</sub> ·2H <sub>2</sub> O	16
33	Glucose	1,000	54	NaHCO <sub>3</sub>	1,273	75	MgSO <sub>4</sub> ·7H <sub>2</sub> O	154
Gey (for slides) [3]			55	NaCl	8,000	76	KCl	400
34	CaCl <sub>2</sub>	170	56	Na <sub>2</sub> HPO <sub>4</sub> ·2H <sub>2</sub> O	100	77	KH <sub>2</sub> PO <sub>4</sub>	150
35	MgCl <sub>2</sub> ·6H <sub>2</sub> O	210	57	Glucose	2,000	78	NaHCO <sub>3</sub>	0
36	MgSO <sub>4</sub> ·7H <sub>2</sub> O	70	Hanks <sup>2,3,5-10</sup> [5]			79	NaCl	8,000
37	KCl	370	58	CaCl <sub>2</sub>	140	80	Na <sub>2</sub> HPO <sub>4</sub> ·7H <sub>2</sub> O	290
38	KH <sub>2</sub> PO <sub>4</sub>	30	59	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	81	Glucose	1,100
39	NaHCO <sub>3</sub>	227	60	KCl	400	Dulbecco <sup>2,3</sup> [1]		
40	NaCl	8,000	61	KH <sub>2</sub> PO <sub>4</sub>	60	82	CaCl <sub>2</sub>	100
41	Na <sub>2</sub> HPO <sub>4</sub> ·2H <sub>2</sub> O	150	62	NaHCO <sub>3</sub>	350	83	MgCl <sub>2</sub> ·6H <sub>2</sub> O	100
42	Glucose	1,000	63	NaCl	8,000	84	KCl	200
Earle <sup>2,3,5-9</sup> [2]			64	Na <sub>2</sub> HPO <sub>4</sub> ·2H <sub>2</sub> O	60	85	KH <sub>2</sub> PO <sub>4</sub>	200
43	CaCl <sub>2</sub>	200	65	Glucose	1,000	86	NaCl	8,000
44	MgSO <sub>4</sub>	100	Puck (Saline F) <sup>2,3,9</sup> [7]			87	Na <sub>2</sub> HPO <sub>4</sub>	1,150
45	KCl	400	66	CaCl <sub>2</sub> ·2H <sub>2</sub> O	16			

<sup>2</sup>/ Available commercially from Colorado Serum Co., 4950 York Street, Denver 16, Colo. <sup>3</sup>/ Available commercially from Difco Laboratories, 920 Henry Street, Detroit 1, Mich. <sup>6</sup>/ Available commercially from Baltimore Biological Laboratory, Division of B-D Laboratories, Inc., 2201 Aisquith Street, Baltimore 18, Md. <sup>8</sup>/ Available commercially from Flow Laboratories, Inc., 1710 Chapman Avenue, Rockville, Md. <sup>7</sup>/ Available commercially from Grand Island Biological Co., 959 East River Road, Grand Island, N. Y. <sup>9</sup>/ Available commercially from Hyland Laboratories, 4501 Colorado Boulevard, Los Angeles 39, Calif. <sup>10</sup>/ Available commercially from Microbiological Associates, Inc., 4846 Bethesda Avenue, Bethesda 14, Md. <sup>11</sup>/ One of two solutions described by Hanks.

**Contributors:** (a) Waymouth, Charity, (b) Ambrose, Charles Tesch

**References:** [1] Dulbecco, R., and M. Vogt. 1954. J. Exptl. Med. 99:167. [2] Earle, W. R. 1943. J. Natl. Cancer Inst. 4:165. [3] Gey, G. O., and M. K. Gey. 1936. Am. J. Cancer 27:55. [4] Hanks, J. H. 1948. J. Cellular Comp. Physiol. 31:235. [5] Hanks, J. H., and R. E. Wallace. 1949. Proc. Soc. Exptl. Biol. Med. 71:196. [6] Locke, F. S. 1901. Centr. Physiol. 14:670. [7] Puck, T. T., S. J. Cieciura, and A. Robinson. 1958. J. Exptl. Med. 108:945. [8] Ringer, S. 1883. J. Physiol. (London) 4:222. [9] Ringer, S. 1886. Ibid. 7:291. [10] Tyrode, M. V. 1910. Arch. Intern. Pharmacodyn. 20:205.

### Part II. TISSUE CULTURE MEDIA

Inclusion of names of commercial suppliers in no way implies endorsement by the Federation of American Societies for Experimental Biology.

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
Medium 199 <sup>1-7</sup> [7]			3	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	7	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140
1	CaCl <sub>2</sub>	200	4	KCl	400	8	Sodium acetate	50.0
2	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O	0.1	5	NaHCO <sub>3</sub>	2,200	9	Glucose	1,000
			6	NaCl	6,800	10	2-Deoxy-D-ribose	0.5

*continued*

## 141. CULTURE MEDIA: ANIMAL TISSUES

### Part II. TISSUE CULTURE MEDIA

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
Medium 199 <sup>1-7</sup> [7]			63	KCl	400	116	Pyridoxine HCl	0.025
11	D-Ribose	3.5	64	NaHCO <sub>3</sub>	2,200	117	Riboflavin	0.01
12	DL-Alanine	50.0	65	NaCl	6,800	118	Thiamine HCl	0.01
13	L-Arginine HCl	70.0	66	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140	Eagle's Basal <sup>1-7</sup> [1]		
14	DL-Aspartic acid	60.0	67	Sodium acetate	83.0 <sup>a</sup>	119	CaCl <sub>2</sub>	111
15	L-Cysteine HCl	0.1	68	Sodium glucuronate	4.2	120	MgCl <sub>2</sub>	102
16	L-Cystine	20.0	69	Glucose	1,000	121	KCl	373
17	DL-Glutamic acid	150.0	70	L-Alanine	25.0	122	NaHCO <sub>3</sub>	1,680
18	L-Glutamine	100.0	71	L-Arginine HCl	70.0	123	NaCl	5,845
19	Glycine	50.0	72	L-Aspartic acid	30.0	124	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	138
20	L-Histidine HCl	20.0	73	L-Cysteine HCl	260.0	125	Glucose	900
21	Hydroxy-L-proline	10.0	74	L-Cystine	20.0	126	L-Arginine HCl	17.5
22	DL-Isoleucine	40.0	75	L-Glutamic acid	75.0	127	L-Cystine	12.0
23	DL-Leucine	120.0	76	L-Glutamine	100.0	128	L-Glutamine	292.0
24	L-Lysine HCl	70.0	77	Glycine	50.0	129	L-Histidine HCl	7.75
25	DL-Methionine	30.0	78	L-Histidine HCl	20.0	130	L-Isoleucine	26.0
26	DL-Phenylalanine	50.0	79	Hydroxy-L-proline	10.0	131	L-Leucine	26.0
27	L-Proline	40.0	80	L-Isoleucine	20.0	132	L-Lysine HCl	29.0
28	L-Serine	50.0	81	L-Leucine	60.0	133	L-Methionine	7.5
29	DL-Threonine	60.0	82	L-Lysine HCl	70.0	134	L-Phenylalanine	16.0
30	DL-Tryptophan	20.0	83	L-Methionine	15.0	135	L-Threonine	24.0
31	L-Tyrosine	40.0	84	L-Phenylalanine	25.0	136	L-Tryptophan	4.0
32	DL-Valine	50.0	85	L-Proline	40.0	137	L-Tyrosine	18.0
33	Adenosine triphosphate	10.0	86	L-Serine	25.0	138	L-Valine	23.0
34	Adenine	10.0	87	L-Threonine	30.0	139	Biotin	0.25
35	Guanine HCl	0.3	88	L-Tryptophan	10.0	140	Choline HCl	0.14
36	Hypoxanthine	0.3	89	L-Tyrosine	40.0	141	Folic acid	0.44
37	Thymine	0.3	90	L-Valine	25.0	142	Nicotinamide	0.12
38	Uracil	0.3	91	Deoxyadenosine	10.0	143	Ca pantothenate	0.48
39	Xanthine	0.3	92	Deoxycytidine HCl	10.0	144	Pyridoxal HCl	0.20
40	Adenylic acid	0.2	93	Deoxyguanosine	10.0	145	Riboflavin	0.04
41	Cholesterol	0.2	94	5-Methyl deoxycytidine	0.1	146	Thiamine HCl	0.34
42	Tween 80	20.0	95	Thymidine	10.0	147	Penicillin	50.0
43	Glutathione	0.05	96	Diphosphopyridine nucleotide	7.0	148	Streptomycin	50.0
44	Vitamin A	0.1	97	Flavin adenine dinucleotide	1.0	149	Phenol red	5.0
45	p-Aminobenzoic acid	0.05	98	Triphosphopyridine nucleotide	1.0	Eagle's Minimum Essential <sup>1,4-7</sup> [2]		
46	Ascorbic acid	0.05	99	Uridine triphosphate	1.0	150	CaCl <sub>2</sub>	200
47	Biotin	0.01	100	Coccarboxylase	1.0	151	MgCl <sub>2</sub> ·6H <sub>2</sub> O	200
48	Calciferol	0.1	101	Coenzyme A	2.5	152	KCl	400
49	Choline HCl	0.50	102	Cholesterol	0.2	153	NaHCO <sub>3</sub>	2,000
50	Folic acid	0.01	103	Tween 80	5.0	154	NaCl	6,800
51	m-Inositol	0.05	104	Ethanol	16.0	155	NaH <sub>2</sub> PO <sub>4</sub> ·2H <sub>2</sub> O	150
52	Menadione	0.01	105	Glutathione	10.0	156	Glucose	1,000
53	Nicotinic acid	0.025	106	p-Aminobenzoic acid	0.05	157	L-Arginine HCl	105.0
54	Nicotinamide	0.025	107	Ascorbic acid	50.0	158	L-Cystine	24.0
55	Ca pantothenate	0.01	108	Biotin	0.01	159	L-Glutamine	292.0
56	Pyridoxal HCl	0.025	109	Choline HCl	0.50	160	L-Histidine HCl	31.0
57	Pyridoxine HCl	0.025	110	Folic acid	0.01	161	L-Isoleucine	52.0
58	Riboflavin	0.01	111	m-Inositol	0.05	162	L-Leucine	52.0
59	Thiamine HCl	0.01	112	Nicotinic acid	0.025	163	L-Lysine	58.0
60	α-Tocopherol phosphate	0.01	113	Nicotinamide	0.025	164	L-Methionine	15.0
CMRL 1066 <sup>1,3,7,8</sup> [8]			114	Ca pantothenate	0.01	165	L-Phenylalanine	32.0
61	CaCl <sub>2</sub>	200	115	Pyridoxal HCl	0.025	166	L-Threonine	48.0
62	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200						

/1/ Available commercially from Baltimore Biological Laboratory, Division of B-D Laboratories, Inc., 2201 Aisquith Street, Baltimore 18, Md. /2/ Available commercially from Colorado Serum Co., 4950 York Street, Denver 16, Colo. /3/ Available commercially from Difco Laboratories, 920 Henry Street, Detroit 1, Mich. /4/ Available commercially from Flow Laboratories, Inc., 1710 Chapman Ave., Rockville, Md. /5/ Available commercially from Grand Island Biological Co., 959 East River Road, Grand Island, N. Y. /6/ Available commercially from Hyland Laboratories, 4501 Colorado Boulevard, Los Angeles 39, Calif. /7/ Available commercially from Microbiological Associates, Inc., 4846 Bethesda Avenue, Bethesda 14, Md. /8/ Available commercially from Connaught Medical Research Laboratories, Toronto 4, Canada. /9/ For sodium acetate + 3H<sub>2</sub>O.

*continued*

## 141. CULTURE MEDIA: ANIMAL TISSUES

### Part II. TISSUE CULTURE MEDIA

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)	(B)		(A)	(B)	(A)	(B)		
Eagle's Minimum Essential <sup>1,4-7</sup> [2]			223	Coccarboxylase	1.0	285	Deoxyadenosine	10.0
NCTC 107 <sup>2,3</sup> [4]			224	Coenzyme A	2.5	286	Deoxycytidine	10.0
167	L-Tryptophan	10.0	225	Cholesterol	2.0	287	Deoxyguanosine	10.0
168	L-Tyrosine	36.0	226	Tween 80	22.5	288	5-Methyl cytosine	0.1
169	L-Valine	46.0	227	Methyl linoleate	1.0	289	Thymidine	10.0
170	Choline HCl	1.0	228	Methyl linolenate	1.0	290	Diphosphopyridine nucleotide	7.0
171	Folic acid	1.0	229	Methyl arachidonate	1.0	291	Flavin adenine dinucleotide	1.0
172	m-Inositol	2.0	230	Glutathione	10.1	292	Triphosphopyridine nucleotide	1.0
173	Nicotinamide	1.0	231	Vitamin A	0.25	293	Uridine triphosphate	1.0
174	Ca pantothenate	1.0	232	p-Aminobenzoic acid	0.125	294	Coccarboxylase	1.0
175	Pyridoxal HCl	1.0	233	Ascorbic acid	49.9	295	Coenzyme A	2.5
176	Riboflavin	0.1	234	Biotin	0.025	296	Tween 80	12.5
177	Thiamine HCl	1.0	235	Calciferol	0.25	297	Glutathione	10.1
178	CaCl <sub>2</sub>	200	236	Choline HCl	1.25	298	Vitamin A	0.25
179	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	237	Folic acid	0.025	299	p-Aminobenzoic acid	0.125
180	KCl	400	238	m-Inositol	0.125	300	Ascorbic acid	49.9
181	NaHCO <sub>3</sub>	2,200	239	Menedione	0.025	301	Biotin	0.025
182	NaCl	6,800	240	Nicotinic acid	0.0625	302	Calciferol	0.25
183	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140	241	Nicotinamide	0.0625	303	Choline HCl	1.25
184	Sodium acetate	50.0	242	Ca pantothenate	0.025	304	Cyanocobalamin	1.0
185	Sodium glucuronate	1.8	243	Pyridoxal HCl	0.0625	305	Folic acid	0.025
186	Glucose	1,000	244	Pyridoxine HCl	0.0625	306	m-Inositol	0.125
187	Glucuronolactone	1.8	245	Riboflavin	0.025	307	Menadione	0.025
188	D-Glucosamine HCl	3.2	246	Thiamine HCl	0.025	308	Nicotinic acid	0.0625
189	L-Alanine	31.48	247	a-Tocopherol phosphate	0.025	309	Nicotinamide	0.0625
190	L-a-Aminobutyric acid	5.51	248	Phenol red	20.0	310	Ca pantothenate	0.025
191	L-Arginine HCl	25.76	NCTC 109 <sup>3,4,5,7</sup> [6]			311	Pyridoxal HCl	0.0625
192	L-Asparagine	8.09	249	CaCl <sub>2</sub>	200	312	Pyridoxine HCl	0.0625
193	L-Aspartic acid	9.91	250	MgSO <sub>4</sub> ·7H <sub>2</sub> O	100	313	Riboflavin	0.025
194	L-Cysteine HCl	260.0	251	KCl	400	314	Thiamine HCl	0.025
195	L-Cystine	10.49	252	NaHCO <sub>3</sub>	2,200	315	a-Tocopherol phosphate	0.025
196	L-Glutamic acid	8.26	253	NaCl	6,800	316	Phenol red	20.0
197	L-Glutamine	135.73	254	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140	NCTC 117 [3]		
198	Glycine	13.51	255	Sodium acetate	50.0	317	CaCl <sub>2</sub>	200
199	L-Histidine HCl	19.73	256	Sodium glucuronate	1.8	318	MgSO <sub>4</sub> ·7H <sub>2</sub> O	100
200	Hydroxy-L-proline	4.09	257	Glucose	1,000	319	KCl	400
201	L-Isoleucine	18.04	258	Glucuronolactone	1.8	320	NaHCO <sub>3</sub>	2,200
202	L-Leucine	20.44	259	D-Glucosamine HCl	3.2	321	NaCl	6,800
203	L-Lysine HCl	30.75	260	L-Alanine	31.48	322	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	140
204	L-Methionine	4.44	261	L-a-Aminobutyric acid	5.51	323	Sodium glucuronate	1.8
205	L-Ornithine HCl	7.38	262	L-Arginine HCl	25.76	324	Glucose	1,000
206	L-Phenylalanine	16.53	263	L-Asparagine	8.09	325	Glucuronolactone	1.8
207	L-Proline	6.13	264	L-Aspartic acid	9.91	326	D-Glucosamine HCl	3.2
208	L-Serine	10.75	265	L-Cysteine HCl	260.0	327	L-Alanine	31.48
209	L-Taurine	4.18	266	L-Cystine	10.49	328	L-a-Aminobutyric acid	5.51
210	L-Threonine	18.93	267	L-Glutamic acid	8.26	329	L-Arginine HCl	25.76
211	L-Tryptophan	17.50	268	L-Glutamine	135.73	330	L-Asparagine	8.09
212	L-Tyrosine	16.44	269	Glycine	13.51	331	L-Aspartic acid	9.91
213	L-Valine	25.0	270	L-Histidine HCl	19.73	332	L-Cysteine HCl	260.0
214	Deoxyadenosine	10.0	271	Hydroxy-L-proline	4.09	333	L-Cystine	10.49
215	Deoxycytidine HCl	10.0	272	L-Isoleucine	18.04	334	L-Glutamic acid	8.26
216	Deoxyguanosine	10.0	273	L-Leucine	20.44	335	L-Glutamine	135.73
217	5-Methyl cytosine	0.1	274	L-Lysine HCl	30.75	336	Glycine	13.51
218	Thymidine	10.0	275	L-Methionine	4.44	337	L-Histidine HCl	19.73
219	Diphosphopyridine nucleotide	7.0	276	L-Ornithine HCl	7.38	338	Hydroxy-L-proline	4.09
220	Flavin adenine dinucleotide	1.0	277	L-Phenylalanine	16.53	339	L-Isoleucine	18.04
221	Triphosphopyridine nucleotide	1.0	278	L-Proline	6.13	340	L-Leucine	20.44
222	Uridine triphosphate	1.0	279	L-Serine	10.75	341	L-Lysine HCl	30.75
			280	L-Taurine	4.18	342	L-Methionine	4.44
			281	L-Threonine	18.93	343	L-Ornithine HCl	7.38
			282	L-Tryptophan	17.50			
			283	L-Tyrosine	16.44			
			284	L-Valine	25.0			

continued



*Continails*  
141. CULTURE MEDIA: ANIMAL TISSUES

Part II. TISSUE CULTURE MEDIA

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)	(B)		(A)	(B)	(A)	(B)		
NCTC 117 [3]			396	L-Threonine	37.5	448	L-Glutamic acid	150.0
344	L-Phenylalanine	16.53	397	L-Tryptophan	20.0	449	L-Glutamine	350.0
345	L-Proline	6.13	398	L-Tyrosine	40.0	450	Glycine	50.0
346	L-Serine	10.75	399	DL-Valine	50.0	451	L-Histidine HCl	150.0
347	L-Taurine	4.18	400	Hypoxanthine	25	452	L-Isoleucine	25.0
348	L-Threonine	18.93	401	Biotin	0.1	453	L-Leucine	50.0
349	L-Tryptophan	17.50	402	Choline HCl	3.0	454	L-Lysine HCl	240.0
350	L-Tyrosine	16.44	403	Folic acid	0.1	455	L-Methionine	50.0
351	L-Valine	25.0	404	m-Inositol	1.0	456	L-Phenylalanine	50.0
352	Deoxycytidine HCl	10.0	405	Nicotinamide	3.0	457	L-Proline	50.0
353	Thymidine	10.0	406	Ca pantothenate	3.0	458	L-Threonine	75.0
354	Tween 80	12.5	407	Pyridoxine HCl	0.5	459	L-Tryptophan	40.0
355	Glutathione	10.0	408	Riboflavin	0.5	460	L-Tyrosine	40.0
356	Vitamin A	0.25	409	Thiamine HCl	5.0	461	L-Valine	65.0
357	$\beta$ -Aminobutyric acid	0.125	410	Phenol red	1.2	462	Hypoxanthine	25
358	Ascorbic acid	50.0	Trowell's T8 <sup>e</sup> [10]			463	Glutathione	15.0
359	Biotin	0.025	411	CaCl <sub>2</sub>	220	464	Ascorbic acid	17.5
360	Calciferol	0.25	412	MgSO <sub>4</sub> ·7H <sub>2</sub> O	250	465	Biotin	0.02
361	Choline HCl	1.25	413	KCl	450	466	Choline HCl	250.0
362	Cyanocobalamin	1.0	414	NaHCO <sub>3</sub>	2,820	467	Cyanocobalamin	0.2
363	Folic acid	0.025	415	NaCl	6,100	468	Folic acid	0.4
364	m-Inositol	0.125	416	NaH <sub>2</sub> PO <sub>4</sub> ·2H <sub>2</sub> O	450	469	m-Inositol	1.0
365	Menadione	0.025	417	Glucose	4,000	470	Nicotinamide	1.0
366	Nicotinic acid	0.0625	418	L-Arginine HCl	21.0	471	Ca pantothenate	1.0
367	Nicotinamide	0.0625	419	L-Cysteine HCl	47.0	472	Pyridoxine HCl	1.0
368	Ca pantothenate	0.025	420	L-Histidine HCl	10.0	473	Riboflavin	1.0
369	Pyridoxal HCl	0.0625	421	L-Isoleucine	26.0	474	Thiamine HCl	10.0
370	Pyridoxine HCl	0.0625	422	L-Leucine	26.0	475	Phenol red	10.0
371	Riboflavin	0.025	423	L-Lysine HCl	36.0	MD 705/1 [5]		
372	Thiamine HCl	0.025	424	DL-Methionine	15.0	476	CaCl <sub>2</sub> ·2H <sub>2</sub> O	120
373	$\alpha$ -Tocopherol phosphate	0.025	425	L-Phenylalanine	33.0	477	CoCl <sub>2</sub> ·6H <sub>2</sub> O	0.11
374	Phenol red	20.0	426	L-Threonine	48.0	478	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.25
Puck's N16 <sup>2,3,5,6-7</sup> [9]			427	L-Tryptophan	4.0	479	FeSO <sub>4</sub>	0.26
375	CaCl <sub>2</sub> ·2H <sub>2</sub> O	16	428	L-Tyrosine	18.0	480	MgCl <sub>2</sub> ·6H <sub>2</sub> O	240
376	MgSO <sub>4</sub> ·7H <sub>2</sub> O	154	429	L-Valine	23.0	481	MgSO <sub>4</sub> ·7H <sub>2</sub> O	100
377	KCl	285	430	$\beta$ -Aminobenzoic acid	35.0	482	MnSO <sub>4</sub> ·H <sub>2</sub> O	0.08
378	KH <sub>2</sub> PO <sub>4</sub>	83	431	Thiamine HCl	17.0	483	KCl	150
379	NaHCO <sub>3</sub>	1,200	432	Insulin	50.0	484	KH <sub>2</sub> PO <sub>4</sub>	80
380	NaCl	7,400	433	Chloramphenicol	30.0	485	NaHCO <sub>3</sub>	2,240
381	Na <sub>2</sub> HPO <sub>4</sub> ·7H <sub>2</sub> O	290	434	Phenol red	10.0	486	NaCl	6,000
382	Glucose	1,100	MB 752/1 <sup>1,3,6-7</sup> [11]			487	Na <sub>2</sub> HPO <sub>4</sub>	300
383	L-Arginine HCl	37.5	435	CaCl <sub>2</sub> ·2H <sub>2</sub> O	120	488	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.15
384	L-Aspartic acid	30.0	436	MgCl <sub>2</sub> ·6H <sub>2</sub> O	240	489	Ammonium paramolybdate	0.12
385	L-Cystine	7.5	437	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	490	Glucose	5,000
386	L-Glutamic acid	75.0	438	KCl	150	491	L-Arginine HCl	75.0
387	L-Glutamine	200.0	439	KH <sub>2</sub> PO <sub>4</sub>	80	492	L-Aspartic acid	60.0
388	Glycine	100.0	440	NaHCO <sub>3</sub>	2,240	493	L-Cysteine HCl	90.0
389	L-Histidine HCl	37.5	441	NaCl	6,000	494	L-Cystine	15.0
390	DL-Isoleucine	25.0	442	Na <sub>2</sub> HPO <sub>4</sub>	300	495	L-Glutamic acid	150.0
391	L-Leucine	25.0	443	Glucose	5,000	496	L-Glutamine	350.0
392	L-Lysine HCl	80.0	444	L-Arginine HCl	75.0	497	Glycine	50.0
393	L-Methionine	25.0	445	L-Aspartic acid	60.0	498	L-Histidine HCl	150.0
394	L-Phenylalanine	25.0	446	L-Cysteine HCl	90.0	499	L-Isoleucine	25.0
395	L-Proline	25.0	447	L-Cystine	15.0	500	L-Leucine	50.0

/1/ Available commercially from Baltimore Biological Laboratory, Division of B-D Laboratories, Inc., 2201 Aisquith Street, Baltimore 18, Md. /2/ Available commercially from Colorado Serum Co., 4950 York Street, Denver 16, Colo. /3/ Available commercially from Difco Laboratories, 920 Henry Street, Detroit 1, Mich. /4/ Available commercially from Flow Laboratories, Inc., 1710 Chapman Ave., Rockville, Md. /5/ Available commercially from Grand Island Biological Co., 959 East River Road, Grand Island, New York. /6/ Available commercially from Hyland Laboratories, 4501 Colorado Boulevard, Los Angeles 39, Calif. /7/ Available commercially from Microbiological Associates, Inc., 4846 Bethesda Avenue, Bethesda 14, Md.

*continued*



141. CULTURE MEDIA: ANIMAL TISSUES

Part II. TISSUE CULTURE MEDIA

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
MD 705/1 [5]			507	L-Tyrosine	40.0	515	Folic acid	0.5
501	L-Lysine HCl	240.0	508	L-Valine	65.0	516	m-Inositol	1.0
502	L-Methionine	50.0	509	Hypoxanthine	25	517	Nicotinamide	1.0
503	L-Phenylalanine	50.0	510	Glutathione	15.0	518	Ca pantothenate	1.0
504	L-Proline	50.0	511	Ascorbic acid	17.5	519	Pyridoxine HCl	1.0
505	L-Threonine	75.0	512	Biotin	0.02	520	Riboflavin	1.0
506	L-Tryptophan	40.0	513	Choline HCl	250.0	521	Thiamine HCl	10.0
			514	Cyanocobalamin	0.2	522	Phenol red	10.0

Contributors: (a) Waymouth, Charity, (b) Parker, Raymond C.

References: [1] Eagle, H. 1955. Science 122:501. [2] Eagle, H. 1959. Ibid. 130:432. [3] Evans, V. J. 1961. Pathol. Biol. (Paris) 9:578. [4] Evans, V. J., et al. 1956. Cancer Res. 16:77. [5] Kitos, P. A., R. Sinclair, and C. Waymouth. 1962. Exptl. Cell Res. 27:307. [6] McQuilkin, W. T., V. J. Evans, and W. R. Earle. 1957. J. Natl. Cancer Inst. 19:885. [7] Morgan, J. F., H. J. Morton, and R. C. Parker. 1950. Proc. Soc. Exptl. Biol. Med. 73:1. [8] Parker, R. C. 1961. Methods of tissue culture. Ed. 3. P. B. Hoeber, New York. p. 77. [9] Puck, T. T., S. J. Cieciura, and A. Robinson. 1958. J. Exptl. Med. 108:945. [10] Trowell, O. A. 1959. Exptl. Cell Res. 16:118. [11] Waymouth, C. 1959. J. Natl. Cancer Inst. 22:1003.

142. CULTURE MEDIA: PLANTS

Part I. BACTERIA

Amino acids given as DL-isomers.

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
Heterotrophic Bacteria <sup>1,2</sup>			14	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.02	34	MgSO <sub>4</sub> ·7H <sub>2</sub> O	614
1	Peptone	5,000	15	FeSO <sub>4</sub> ·7H <sub>2</sub> O	20	35	MnSO <sub>4</sub>	15
2	Yeast extract	3,000	16	MgSO <sub>4</sub> ·7H <sub>2</sub> O	250	36	KCl	400
Heterotrophic Bacteria <sup>2,5</sup>			17	MnCl <sub>2</sub> ·4H <sub>2</sub> O	0.05	37	NaCl	300
3	Peptone	5,000	18	K <sub>2</sub> HPO <sub>4</sub>	500	38	Na <sub>2</sub> SO <sub>4</sub>	4,000
4	Yeast extract	3,000	19	Na <sub>2</sub> CO <sub>3</sub>	2,000 <sup>5</sup>	39	ZnCl <sub>2</sub>	10
5	Agar	15,000	20	NaCl	3,000 <sup>5</sup>	40	Citric acid	2,000
Saprophytic Actinomyces and Streptomyces* [7]			21	Na <sub>2</sub> S	1,000 <sup>6</sup>	41	Sucrose	100,000
6	Glycerol	10,000	22	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.5	42	Asparagine	2,000
7	Asparagine	1,000	23	Potassium acetate	1,000 <sup>7</sup>	43	Glutamic acid	2,000
8	Dipotassium phosphate	1,000	24	Sodium succinate	4,000	44	Adenine sulfate	40
9	Agar	15,000	25	Malic acid	3,000 <sup>7</sup>	45	Guanine HCl	40
Photosynthetic Bacteria [4, 5]			26	Glycerol	2,000 <sup>7</sup>	46	Uracil	40
10	NH <sub>4</sub> Cl	1,000	27	Glutamic acid	2,000 <sup>7</sup>	47	Xanthine	40
11	H <sub>3</sub> BO <sub>3</sub>	2.8	28	D-Biotin	0.004 <sup>7</sup>	48	p-Aminobenzoic acid	0.02
12	CaCl <sub>2</sub>	100	29	Niacin	1.0 <sup>7</sup>	49	D-Biotin	0.02
13	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	0.05	30	Thiamine HCl	1.0 <sup>7</sup>	50	Choline Cl	10
			31	pH = 7-9		51	Folacin	0.02
				<i>Bacillus subtilis</i> [1]		52	L-Inositol	10
			32	(NH <sub>4</sub> ) <sub>2</sub> HPO <sub>4</sub>	8,000	53	Niacin	2
			33	FeCl <sub>3</sub> ·6H <sub>2</sub> O	33	54	DL-Ca pantothenate	4
						55	Pyridoxine HCl	2

<sup>1/</sup> Nonsynthetic, nutrient broth prepared by adding specified ingredients to one liter of distilled water. <sup>2/</sup> Sugar broth or agar may be prepared by adding 5,000 mg/L of desired sugar. <sup>3/</sup> Nonsynthetic, nutrient agar prepared by adding specified ingredients to one liter of distilled water. <sup>4/</sup> Nonsynthetic medium prepared by adding specified ingredients to one liter of distilled water. <sup>5/</sup> For purple and green sulfur bacteria. <sup>6/</sup> For marine forms. <sup>7/</sup> For purple nonsulfur bacteria.

continued

## 142. CULTURE MEDIA: PLANTS

### Part I. BACTERIA

	Concentration mg/L		Concentration mg/L		Concentration mg/L			
(A)	(B)	(A)	(B)	(A)	(B)			
<i>Bacillus subtilis</i> [1]		<i>Lactobacillus leichmannii</i> [6]		144	Pyridoxal phosphate	1		
56	Pyridoxamine HCl	2	98	NH <sub>4</sub> Cl	280	145	Pyridoxine HCl	2
57	Riboflavin	4	99	FeSO <sub>4</sub> ·7H <sub>2</sub> O	10	146	Pyridoxamine HCl	0.4
58	Thiamine HCl	4	100	MgSO <sub>4</sub> ·7H <sub>2</sub> O	1,400	147	Riboflavin	1
<i>Haemophilus parainfluenzae</i> [2]		101	MnSO <sub>4</sub>	203	148	Thiamine HCl	1	
59	CaCl <sub>2</sub>	3	102	K <sub>2</sub> HPO <sub>4</sub>	2,000	pH = 5.5		
60	FeSO <sub>4</sub> ·7H <sub>2</sub> O	12.8	103	KH <sub>2</sub> PO <sub>4</sub>	2,000	<i>Streptococcus faecalis</i> [3]		
61	MgSO <sub>4</sub> ·7H <sub>2</sub> O	82	104	Sodium acetate	3,600	150	NH <sub>4</sub> Cl	2,500
62	KH <sub>2</sub> PO <sub>4</sub>	3,120	105	Sodium citrate	5,000	151	FeSO <sub>4</sub> ·7H <sub>2</sub> O	27
63	Sodium acetate	6,000	106	Sodium ethyl oxalacetate	100	152	MgSO <sub>4</sub> ·7H <sub>2</sub> O	512
64	Glucose	1,000	107	Glucose	20,000	153	MnSO <sub>4</sub>	30
65	Alanine	1,000	108	Alanine	200	154	K <sub>2</sub> HPO <sub>4</sub>	5,000
66	Arginine HCl	400	109	Arginine HCl	200	155	NaCl	15
67	Aspartic acid	1,000	110	Asparagine	200	156	Sodium acetate	5,000
68	Cystine	200	111	Aspartic acid	200	157	Sodium citrate	5,000
69	Glutamic acid	2,000	112	Cysteine	800	158	Glucose	20,000
70	Glycine	100	113	Cystine	400	159	Alanine	500
71	Histidine HCl	200	114	Glutamic acid	400	160	Arginine HCl	400
72	Isoleucine	200	115	Glutamine	100	161	Asparagine	500
73	Leucine	200	116	Glycine	300	162	Aspartic acid	500
74	Lysine HCl	400	117	Histidine HCl	200	163	Cystine	200
75	Methionine	200	118	Hydroxyproline	50	164	Glutamic acid	1,000
76	Phenylalanine	200	119	Isoleucine	300	165	Glycine	100
77	Proline	200	120	Leucine	100	166	Histidine HCl	200
78	Serine	200	121	Lysine HCl	600	167	Isoleucine	200
79	Threonine	200	122	Methionine	100	168	Leucine	200
80	Tryptophan	200	123	Norleucine	200	169	Lysine HCl	400
81	Tyrosine	200	124	Phenylalanine	500	170	Methionine	200
82	Valine	200	125	Proline	400	171	Phenylalanine	200
83	Adenine sulfate	10	126	Serine	100	172	Proline	400
84	Guanine HCl	10	127	Threonine	100	173	Serine	500
85	Uracil	10	128	Tryptophan	50	174	Threonine	200
86	Nicotinamide adenine dinucleotide	0.1	129	Tyrosine	400	175	Tryptophan	200
87	Putrescine	500	130	Valine	200	176	Tyrosine	200
88	p-Aminobenzoic acid	0.001	131	Adenine sulfate	5	177	Valine	200
89	D-Biotin	0.001	132	Cytidylic acid	10	178	Adenine sulfate	10
90	Choline Cl	5	133	Guanine HCl	5	179	Uridine	0.2
91	Folacin	0.01	134	Uracil	5	180	Glutathione	20
92	L-Inositol	20	135	Xanthine	8	181	Tween 80	10
93	Niacin	0.5	136	Tween 80	1	182	p-Aminobenzoic acid	0.2
94	D,L-Ca pantothenate	1	137	p-Aminobenzoic acid	0.04	183	D-Biotin	0.01
95	Pyridoxine HCl	2	138	D-Biotin	0.005	184	Folacin	0.02
96	Riboflavin	0.1	139	Cyanocobalamin	0.01	185	Niacin	1
97	Thiamine HCl	1	140	Folacin	0.06	186	D,L-Ca pantothenate	0.5
			141	Niacin	1	187	Pyridoxine HCl	0.5
			142	D,L-Ca pantothenate	1	188	Pyridoxamine HCl	0.5
			143	Pyridoxal HCl	2	189	Riboflavin	0.5
						190	Thiamine HCl	0.5

Contributors: (a) Pavcek, Paul L., (b) Clark, F. M., (c) Allen, Mary Belle

References: [1] Feeney, R. E., J. A. Garibaldi, and E. M. Humphreys. 1948. Arch. Biochem. 17:435. [2] Herbst, E. J., and E. E. Snell. 1949. J. Biol. Chem. 181:47. [3] Hoffmann, H. A., and P. L. Pavcek. 1952. J. Am. Chem. Soc. 74:344. [4] Hutner, S. H. 1950. J. Gen. Microbiol. 4:286. [5] Larsen, H. 1952. J. Bacteriol. 64:187. [6] Shorb, M. S. 1952. Proc. Soc. Exptl. Biol. Med. 79:611. [7] Skinner, C. E., C. W. Emmons, and H. M. Tsuchiya, ed. 1947. Henrici's Molds, yeasts, and actinomycetes. Ed. 2. J. Wiley, New York. p. 59.

continued

## 142. CULTURE MEDIA: PLANTS

### Part II. FUNGI

Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>
(A)	(B)		(A)	(B)	(A)	(B)		
Molds and Yeasts <sup>2</sup> [4]			13	NaNO <sub>3</sub>	3,000	29	n-Biotin	0.005
1	Potato extract <sup>3</sup>	1,000 ml	14	Sucrose	30,000	30	pH = 5.6	
2	Glucose	10,000	pH = 6.8-6.9			Basidiomycetes <sup>5</sup> [2]		
3	Agar	15,000	<i>Neurospora</i> [1]			31	H <sub>3</sub> BO <sub>3</sub>	0.57
Molds and Yeasts <sup>2,4</sup> [5]			16	NH <sub>4</sub> NO <sub>3</sub>	1,000	32	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.04
4	Peptone	5,000	17	H <sub>3</sub> BO <sub>3</sub>	0.06	33	FeSO <sub>4</sub> ·7H <sub>2</sub> O	0.15
5	Yeast extract	3,000	18	CaCl <sub>2</sub>	100	34	MgSO <sub>4</sub> ·7H <sub>2</sub> O	500
6	Malt extract	3,000	19	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.40	35	MnCl <sub>2</sub> ·4H <sub>2</sub> O	0.04
7	Dextrose	10,000	20	FeSO <sub>4</sub> ·7H <sub>2</sub> O	0.72	36	KH <sub>2</sub> PO <sub>4</sub>	1,500
8	Agar	20,000	21	MgSO <sub>4</sub> ·7H <sub>2</sub> O	500	37	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.31
Aspergilli and Penicillia [3]			22	MnCl <sub>2</sub> ·4H <sub>2</sub> O	0.07	38	Ammonium para-molybdate	0.02
9	FeSO <sub>4</sub> ·7H <sub>2</sub> O	10	23	KH <sub>2</sub> PO <sub>4</sub>	1,000	39	Glucose	10,000
10	MgSO <sub>4</sub> ·7H <sub>2</sub> O	500	24	NaCl	100	40	L-Glutamic acid	1,200± <sup>5</sup>
11	KCl	500	25	Na <sub>2</sub> MoO <sub>4</sub>	0.04	41	Thiamine HCl	1
12	K <sub>2</sub> HPO <sub>4</sub>	1,000	26	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	8.8	42	pH = 5.0-5.5	
			27	Ammonium tartrate	5,000			
			28	Sucrose	15,000			

/1/ Unless stated otherwise. /2/ Nonsynthetic medium. /3/ Boil 300 grams sliced potatoes for 20 minutes and strain through cotton. /4/ Prepared by adding specified ingredients to one liter of distilled water. /5/ Wood-rotting types. Biotin and/or riboflavin may be required by some species. /6/ Or DL-glutamic acid, 2,400± mg/L.

Contributors: (a) Clark, F. M., (b) Wolf, Frederick T., (c) Jennison, Marshall W.

References: [1] Beadle, G. W., and E. L. Tatum. 1945. Am. J. Botany 32:678. [2] Jennison, M. W., et al. 1955. Mycologia 47:275. [3] Raper, K. B., and C. Thom. 1949. A manual of the penicillia. Williams and Wilkins, Baltimore. [4] Skinner, C. E., C. W. Emmons, and H. M. Tsuchiya, ed. 1947. Henrici's Molds, yeasts, and actinomyces. Ed. 2. J. Wiley, New York. p. 53. [5] Wickerham, L. J. Unpublished. Northern Regional Research Laboratory, Peoria, Ill., 1963.

### Part III. ALGAE

Variations of Pringsheim's soil-water medium are for nonsterile cultures, especially for isolation purposes and for growing algae to secure "normal" growth forms. Success with soil-water media depends on the selection of a suitable garden soil. This soil should be of medium humus content and should not have been recently fertilized with commercial fertilizers. Soils with a high clay content are usually not the most suitable for most organisms. A variety of soil-water media can be made using a basic formula to which are added additional materials. The basic soil-water medium is made by placing one-quarter to one-half inch of garden soil in the bottom of a test tube, then adding pyrex-distilled water until the tube is three-quarters full. The tube is then plugged with cotton and steamed (not autoclaved) for one hour on two consecutive days. A few algae such as *Spirogyra* grow well in this basic medium. For most presumably phototrophic algae which thrive in an alkaline medium, a small pinch of powdered CaCO<sub>3</sub> is placed in the bottom of the test tube before the soil and water are added. Some algae (*Astasia*, *Euglena*, *Polytoma*, *Polytomella*, *Pyrobotrys*, and others) require additional complex nitrogenous or carbon compounds not present in the basic formula. In the case of *Euglena* and *Pyrobotrys*, the best results have been obtained by adding one-quarter of a garden pea cotyledon to the basic medium (including CaCO<sub>3</sub>) before steaming. For the colorless forms, the addition of a barley grain before steaming supplies the necessary carbon source. A few strains, such as *Botryococcus*, grow best when a pinch of sterile ammonium magnesium phosphate is added after the steaming of the basic medium (including CaCO<sub>3</sub>). [3]

Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>
(A)	(B)		(A)	(B)	(A)	(B)		
Marine Seaweeds [4]			2	Ca, as Cl	150	5	Fe, as Cl	2
1	B, as H <sub>3</sub> BO <sub>3</sub>	2	3	Co, as Cl	10 µg	6	MgSO <sub>4</sub> ·7H <sub>2</sub> O	8,000
			4	Cu, as Cl	20 µg	7	Mn, as Cl	1

/1/ Unless stated otherwise.

continued

## 142. CULTURE MEDIA: PLANTS

### Part III. ALGAE

Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>	Constituent		Concentration mg/L <sup>1</sup>
(A)	(B)		(A)	(B)	(A)	(B)		
Marine Seaweeds [4]			24	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.08	43	Mo	0.1
8	Mo, as Na salt	0.5	25	FeSO <sub>4</sub> ·7H <sub>2</sub> O	20	44	K <sub>2</sub> HPO <sub>4</sub>	200
9	KCl	700	26	MgSO <sub>4</sub> ·7H <sub>2</sub> O	250	45	Si, as orthosilicic acid	35
10	NaCl	24,000	27	MnCl <sub>2</sub> ·4H <sub>2</sub> O	1.8	46	Zn	0.3
11	NaNO <sub>3</sub>	300	28	KNO <sub>3</sub>	2,000 <sup>b</sup>	pH = 7.0-7.5		
12	Na <sub>2</sub> SiO <sub>3</sub> ·9H <sub>2</sub> O	70	29	K <sub>2</sub> HPO <sub>4</sub>	2,580	<i>Chlorella pyrenoidosa</i> [5]		
13	Zn, as Cl	0.5	30	NaCl	40	48	B	0.5
14	Potassium glycerophosphate	100	31	Na <sub>2</sub> CO <sub>3</sub>	1,500 <sup>c</sup>	49	Ca	0.5
15	Sodium versenol	30	32	Na <sub>2</sub> MoO <sub>4</sub>	0.2	50	Co <sup>7</sup>	0.01
16	Tris <sup>2</sup>	1,000	33	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.02	51	Cu <sup>7</sup>	0.04
17	Vitamin mix #8 <sup>3</sup>	1 ml	34	Sodium citrate	200	52	Fe <sup>7</sup>	0.2
18	Cyanocobalamin	0.5 µg	35	pH = 7-9		53	MgSO <sub>4</sub> ·7H <sub>2</sub> O	500
19	pH = 7.6		<i>Navicula pelliculosa</i> [2]			54	Mn <sup>7</sup>	0.5
Cyanophyta [1]			36	B	0.1	55	Mo	0.02
20	NH <sub>4</sub> VO <sub>3</sub>	0.02 <sup>d</sup>	37	Ca(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	1,000	56	KH <sub>2</sub> PO <sub>4</sub>	1,310
21	H <sub>3</sub> BO <sub>3</sub>	2.86	38	Co	0.1	57	V	0.01
22	CaCl <sub>2</sub>	55	39	Cu	0.1	58	Zn <sup>7</sup>	0.5
23	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	0.05 <sup>d</sup>	40	Fe	0.5	59	Urea <sup>e</sup>	440
			41	MgSO <sub>4</sub> ·7H <sub>2</sub> O	200	pH = 6.0		
			42	Mn	0.1	60		

/: Unless stated otherwise. /<sup>2</sup>/ Tris(hydroxymethyl)aminomethane. /<sup>3</sup>/ 1 ml of vitamin mix #8 contains 0.2 mg thiamine HCl, 0.1 mg nicotinic acid, 0.04 mg putrescine·2HCl, 0.1 mg Ca pantothenate, 5.0 µg riboflavin, 0.04 mg pyridoxine·2HCl, 0.02 mg pyridoxamine·2HCl, 0.01 mg *p*-aminobenzoic acid, 0.5 µg biotin, 0.5 mg choline·H<sub>2</sub> citrate, 1.0 mg inositol, 0.8 mg thymine, 0.26 mg orotic acid, 0.05 µg cyanocobalamin, 0.2 µg folic acid, 2.5 µg folic acid. /<sup>4</sup>/ Not yet shown to be generally required. /<sup>5</sup>/ May be omitted for nitrogen-fixing forms. /<sup>6</sup>/ For those strains which grow only at an alkaline pH. /<sup>7</sup>/ These metals were used as compounds chelated by ethylenediamine tetra-acetic acid. /<sup>8</sup>/ Or KNO<sub>3</sub>, 1,440 mg/L.

**Contributors:** (a) Provasoli, Luigi, (b) Allen, Mary Belle, (c) Starr, Richard C.

**References:** [1] Allen, M. B., and D. I. Arnon. 1955. *Plant Physiol.* 30:366. [2] Lewin, J. C. 1955. *Ibid.* 30:129. [3] Pringsheim, E. G. 1950. In J. Brunel, ed. *The culturing of algae*. C. F. Kettering Foundation, Dayton. p. 19. [4] Provasoli, L., J. J. A. McLaughlin, and M. R. Droop. 1957. *Arch. Mikrobiol.* 25:408. [5] Sorokin, C., and R. W. Krauss. 1962. *Plant Physiol.* 37:37.

### Part IV. HIGHER PLANTS

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)	(B)		(A)	(B)	(A)	(B)		
1	H <sub>3</sub> BO <sub>3</sub>	0.57	4	FeSO <sub>4</sub> ·7H <sub>2</sub> O	2.5	7	H <sub>2</sub> MoO <sub>4</sub>	0.02
2	Ca(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	1,180	5	MgSO <sub>4</sub> ·7H <sub>2</sub> O	493	8	KH <sub>2</sub> PO <sub>4</sub>	136
3	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.04	6	MnCl <sub>2</sub> ·4H <sub>2</sub> O	0.90	9	K <sub>2</sub> SO <sub>4</sub>	349
						10	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.22

**Contributor:** Robbins, W. Rei

**Reference:** Robbins, W. R. Unpublished. Rutgers Univ., New Brunswick, N. J., 1963.



## 143. CULTURE MEDIA: PLANT TISSUES

### Part I. BALANCED SALT SOLUTIONS

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L	
(A)		(B)	(A)		(B)	(A)		(B)	
White [3, 4]			8	KNO <sub>2</sub>	80	15	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.03	
	1	H <sub>3</sub> BO <sub>3</sub>	1.5	9	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	16.5	16	FeCl <sub>3</sub> ·6H <sub>2</sub> O	1
2	Ca(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	300	10	NaSO <sub>4</sub>	200	17	MgSO <sub>4</sub> ·7H <sub>2</sub> O	250	
3	Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	2.5	11	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	3	18	MnSO <sub>4</sub> ·4H <sub>2</sub> O	0.1	
4	MgSO <sub>4</sub> ·7H <sub>2</sub> O	720	Heller [1, 2, 4]			19	NiO <sub>2</sub> ·6H <sub>2</sub> O	0.03	
5	MnSO <sub>4</sub> ·4H <sub>2</sub> O	7	12	AlCl <sub>3</sub>	0.03	20	KCl	750	
6	KCl	65	13	H <sub>3</sub> BO <sub>3</sub>	1	21	KI	0.01	
7	KI	0.75	14	CaCl <sub>2</sub> ·2H <sub>2</sub> O	75	22	NaNO <sub>3</sub>	600	
						23	NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	125	
						24	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	1	

*Contributor:* White, Philip R.

*References:* [1] Gautheret, R. J. 1959. La culture des tissus végétaux. G. Masson, Paris. [2] Heller, R. 1953. Ann. Sci. Nat. Bot. Biol. Vegetale, Ser. 11, 14:1. [3] White, P. R. 1943. A handbook of plant tissue culture. J. Cattell Press, Lancaster, Penna. [4] White, P. R. 1963. The cultivation of animal and plant cells. Ed. 2. Ronald Press, New York.

### Part II. TISSUE CULTURE MEDIA

Constituent		Concentration mg/L	Constituent		Concentration mg/L	Constituent		Concentration mg/L
(A)		(B)	(A)		(B)	(A)		(B)
Stem Tips <sup>1</sup> [1]			6	Niacin	0.5	13	Pyridoxine	0.1
1	Glucose	75,000	7	Pyridoxine	0.1	14	Thiamine	0.1
2	Agar	10,000	8	Thiamine	0.1	15	2, 4-D	0.1
3	Gibberellin	1	Callus <sup>1</sup> [2, 4]			Tumor <sup>2</sup> [2, 4]		
Root Tips <sup>2</sup> [3, 4]			9	Sucrose	50,000	16	Sucrose	50,000
4	Sucrose	20,000	10	Agar	5,000	17	Agar	5,000
5	Glycine	3	11	Glycine	3	18	Niacin	0.5
			12	Niacin	0.5	19	Pyridoxine	0.1
						20	Thiamine	0.1

<sup>1/1</sup> Add specified ingredients to either White's or Heller's balanced salt solution (*see* Part I). <sup>1/2</sup> Add specified ingredients to White's balanced salt solution only (*see* Part I).

*Contributor:* White, Philip R.

*References:* [1] Ball, E. 1960. Growth 24:91. [2] Gautheret, R. J. 1959. La culture des tissus végétaux. G. Masson, Paris. [3] White, P. R. 1943. A handbook of plant tissue culture. J. Cattell Press, Lancaster, Penna. [4] White, P. R. 1963. The cultivation of animal and plant cells. Ed. 2. Ronald Press, New York.

## 144. NATURAL SEA WATER

### Part I. GENERAL CHARACTERISTICS, SALINITY, AND CONSTITUENTS

Values are per kilogram of sea water, unless otherwise specified.

Specification	Value	Specification	Value
(A)	(B)	(A)	(B)
General Characteristics			
1 Density	1.02-1.03	38 Calcium	0.40 g
2 Temperature	-1.5 to +30°C	39 Carbon	28 mg
3 pH, surface water	8.1-8.3	40 Carbon dioxide <sup>a</sup>	64-107 mg
4 pH, at depth	7.5-8.1	41 Cerium	0.4 µg
5 Freezing point <sup>1</sup>	-2°C	42 Cesium	2 µg
6 Specific heat <sup>2</sup>	0.955 cal/g	43 Chlorine	18.98 g
7 Velocity of sound	1,450-1,550 m/sec	44 Chromium	Present
8 Transparency, maximum <sup>3</sup>	66 m	45 Cobalt	0.1 µg
9 Hydrostatic pressure <sup>4</sup>	1 atm/10 m	46 Copper	1-10 µg
Salinity		47 Fluorine	1.4 mg
10 All oceans, average	33-37 g	48 Gallium	0.5 µg
11 Below 1,000 m (-0.5 to +5°C)	34.6-35.0 g	49 Gold	0.006 µg
12 At equator	35 g	50 Helium and neon <sup>5</sup>	0.03 µg
13 20th-40th parallel, N. latitude	35.5 g	51 Iodine	50 µg
14 10th-30th parallel, S. latitude	35.5 g	52 Iron	2-20 µg
15 Average, 60° N. and S. latitudes to Poles	35 g	53 Lanthanum	0.3 µg
16 North Pacific	34.5 g	54 Lead	4 µg
17 North Sea, off Denmark	34 g	55 Lithium	0.1 mg
18 Indian Ocean, near Australia	35.5 g	56 Magnesium	1.27 g
19 South Pacific, off Peru	35.5 g	57 Manganese	1-10 µg
20 Arabian Sea	36-37 g	58 Mercury	0.03 µg
21 Sargasso Sea, N. Atlantic	36.5-37.0 g	59 Molybdenum	0.5 µg
22 South Atlantic, off Brazil	36-37 g	60 Nickel	0.1 µg
23 Red Sea (surface)	38-41 g	61 Nitrogen <sup>6</sup>	10-18 mg
24 Mediterranean Sea (surface)	37-39 g	62 Nitrogen <sup>7</sup>	0.006-0.700 mg
25 Gulf of Mexico (surface)	36-37 g	63 Oxygen <sup>6</sup>	0-12 mg
26 Antarctic Ocean (surface)	34.0-34.6 g	64 Phosphorus	1-100 µg
27 Arctic Ocean (surface)	32-33 g	65 Potassium	0.38 g
Constituents <sup>8</sup>		66 Radium	0.2-3.0 × 10 <sup>-7</sup> µg
28 Aluminum	0.5 mg	67 Rubidium	0.2 mg
29 Argon <sup>9</sup>	0.4-0.7 mg	68 Scandium	0.04 µg
30 Arsenic	10-20 µg	69 Selenium	4 µg
31 Barium	54 µg	70 Silicon	0.02-4.00 mg
32 Bicarbonate	0.14 g	71 Silver	0.3 µg
33 Bismuth	0.2 µg	72 Sodium	10.56 g
34 Boric acid	26 mg	73 Strontium	13 mg
35 Boron	4.6 mg	74 Sulfate	2.65 g
36 Bromine	65 mg	75 Sulfur	0.88 g
37 Cadmium	Present	76 Thorium	0.4 µg
		77 Tin	3 µg
		78 Uranium	1.5 µg
		79 Vanadium	0.3 µg
		80 Yttrium	0.3 µg
		81 Zinc	5 µg

<sup>1</sup>/ For water with salinity of slightly more than 35 g/kg. <sup>2</sup>/ For sea water with a salinity of 35 g/kg at 20°C and atmospheric pressure (760 mm Hg). <sup>3</sup>/ Depth at which a 30-cm Secchi disk disappears from sight in the Sargasso Sea. <sup>4</sup>/ Hydrostatic pressure increases approximately one atmosphere (760 mm Hg) for each ten meters of depth, the exact value being affected by salinity, temperature, and latitude. <sup>5</sup>/ Based on total salinity of 34.325 g/kg, or standard chlorinity of 19 g/kg. <sup>6</sup>/ As dissolved gas. <sup>7</sup>/ In combined form.

**Contributors:** (a) Bowman, H. H. M., (b) Olson, F. C. W., (c) Redfield, Alfred C.

**References:** [1] Bowman, H. H. M. 1956. Ohio J. Sci. 56(2):101. [2] Bruns, E. 1962. Ozeanologie. Deutscher Verlag der Wissenschaften, Berlin. Bd. 2. [3] Marmer, H. A. 1930. The sea. D. Appleton, New York.

[4] Olson, F. C. W. Unpublished. Radio Corp. of America, Princeton, N. J., 1963. [5] Sverdrup, H. U., M. W. Johnson, and R. H. Fleming. 1942. The oceans. Prentice-Hall, New York.

*continued*

# Contrails

## 144. NATURAL SEA WATER

### Part II. SURFACE TEMPERATURE OF THE OCEANS

Values are degrees centigrade.

Latitude		Atlantic Ocean	Indian Ocean	Pacific Ocean	Latitude		Atlantic Ocean	Indian Ocean	Pacific Ocean
(A)		(B)	(C)	(D)	(A)		(B)	(C)	(D)
North					South				
1	70-60	5.60	.....	.....	8	70-60	-1.30	-1.50	-1.30
2	60-50	8.66	.....	5.74	9	60-50	1.76	1.63	5.00
3	50-40	13.16	.....	9.99	10	50-40	8.68	8.67	11.16
4	40-30	20.40	.....	18.62	11	40-30	16.90	17.00	16.98
5	30-20	24.16	26.14	23.38	12	30-20	21.20	22.53	21.53
6	20-10	25.81	27.23	26.42	13	20-10	23.16	25.85	25.11
7	10-0	26.66	27.88	27.20	14	10-0	25.18	27.41	26.01

*Contributor:* Sverdrup, H. U.

*Reference:* Sverdrup, H. U., M. W. Johnson, and R.H. Fleming. 1942. The oceans. Prentice-Hall, New York.

### Part III. RELATION OF CHLORINITY AND SALINITY TO DENSITY

Chlorinity is defined as the amount, in grams, of precipitated halides (chlorine, bromine, and iodine), as determined by precipitation with a silver salt, in one kilogram of salt water. Salinity is defined as the total weight, in grams, of dissolved solids in one kilogram of sea water, when all carbonate has been converted to oxide, the bromine and iodine replaced by chlorine, and all organic matter completely oxidized. The standard chlorinity of sea water is 19, and equals a salinity of 34.325 (salinity = 0.03 + 1.805 x chlorinity).

Chlorinity g/kg		Salinity g/kg	Density at 15°C g/cm <sup>3</sup>	Chlorinity g/kg		Salinity g/kg	Density at 15°C g/cm <sup>3</sup>
(A)		(B)	(C)	(A)		(B)	(C)
1	1	1.84	1.000578	12	12	21.69	1.015789
2	2	3.64	1.001967	13	13	23.50	1.017169
3	3	5.45	1.003354	14	14	25.30	1.018550
4	4	7.25	1.004739	15	15	27.11	1.019932
5	5	9.06	1.006123	16	16	28.91	1.021314
6	6	10.86	1.007506	17	17	30.72	1.022698
7	7	12.67	1.008888	18	18	32.52	1.024084
8	8	14.47	1.010268	19	19	34.33	1.025471
9	9	16.28	1.011649	20	20	36.13	1.026860
10	10	18.08	1.013029	21	21	37.94	1.028251
11	11	19.89	1.014409	22	22	39.74	1.029645
				23	23	41.54	1.031041

*Contributor:* Olson, F. C. W.

*Reference:* Japan Meteorological Agency. 1955. Oceanographical tables. Tokyo.

### Part IV. OXYGEN SATURATION FROM NORMAL DRY ATMOSPHERE

Oxygen saturation values may be 3% too high.

Chlorinity g/kg	Salinity g/kg	Oxygen Saturation in ml/L <sup>1</sup> at								
		-2°C	0°C	5°C	10°C	15°C	20°C	25°C	30°C	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	
1	15	27.11	9.01	8.55	7.56	6.77	6.14	5.63	5.17	4.74
2	16	28.91	8.89	8.43	7.46	6.69	6.07	5.56	5.12	4.68
3	17	30.72	8.76	8.32	7.36	6.60	6.00	5.50	5.06	4.63
4	18	32.52	8.64	8.20	7.26	6.52	5.93	5.44	5.00	4.58

<sup>1</sup>/ Milligram-atoms of oxygen per liter = 0.08931 x ml/L.

*continued*

## 144. NATURAL SEA WATER

### Part IV. OXYGEN SATURATION FROM NORMAL DRY ATMOSPHERE

	Chlorinity g/kg	Salinity g/kg	Oxygen Saturation in ml/L <sup>1</sup> at							
			-2°C	0°C	5°C	10°C	15°C	20°C	25°C	30°C
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
5	19	34.33	8.52	8.08	7.16	6.44	5.86	5.38	4.95	4.52
6	20	36.11	8.39	7.97	7.07	6.35	5.79	5.31	4.86	4.46

<sup>1</sup>/ Milligram-atoms of oxygen per liter = 0.08931 x ml/L.

*Contributor:* Sverdrup, H. U.

*Reference:* Sverdrup, H. U., M. W. Johnson, and R. H. Fleming. 1942. The oceans. Prentice-Hall, New York.

### Part V. PRESSURE - DEPTH GRADIENT

Hydrostatic pressure increases approximately one atmosphere (760 mm Hg) for each ten meters of depth, the exact value being affected by salinity, temperature, and latitude.

Depth meters	Salinity g/kg	Temp. °C	Latitude		Depth meters	Salinity g/kg	Temp. °C	Latitude			
			30° atm/meter	60° atm/meter				30° atm/meter	60° atm/meter		
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)		
1	0	32	0	0.099141	0.099403	5	5000	35	0	0.101757	0.102026
2	0	32	20	0.098831	0.099092	6	5000	35	5	0.101660	0.101929
3	0	35	0	0.099375	0.099638	7	10,000	35	0	0.103952	0.104225
4	0	35	20	0.099052	0.099314						

*Contributor:* ZoBell, Claude E.

*Reference:* Bjerknes, V. F. K., and J. W. Sandström. 1910. Carnegie Inst. Wash. Publ. 88.

## 145. ARTIFICIAL SEA WATER

Salt		Concentration mg/L		Salt		Concentration mg/L		Salt		Concentration mg/L	
(A)	(B)	(A)	(B)	(A)	(B)	(A)	(B)	(A)	(B)	(A)	(B)
Brujewicz [4]				Allen and Nelson [2]				Robertson and Webb [15]			
1	CaCl <sub>2</sub>	1,141		18	CaCl <sub>2</sub>	510		36	CaCl <sub>2</sub>	1,117	
2	MgCl <sub>2</sub>	2,447		19	MgCl <sub>2</sub>	3,420		37	MgCl <sub>2</sub>	4,978	
3	MgSO <sub>4</sub>	3,305		20	MgSO <sub>4</sub>	2,100		38	KCl	725	
4	KCl	725		21	KCl	750		39	NaCl	23,465	
5	NaHCO <sub>3</sub>	202		22	NaCl	26,750		40	Na <sub>2</sub> SO <sub>4</sub>	3,928	
6	NaBr	83			TOTAL	33,530			TOTAL	34,213	
7	NaCl	26,518		ZoBell [17]				Allen [1]			
	TOTAL	34,421		23	NH <sub>4</sub> NO <sub>3</sub>	2		41	CaCl <sub>2</sub>	1,200	
Lyman and Fleming [11]				24	H <sub>3</sub> BO <sub>3</sub>	27		42	MgCl <sub>2</sub>	2,550	
8	H <sub>3</sub> BO <sub>3</sub>	26		25	CaCl <sub>2</sub>	1,140		43	MgSO <sub>4</sub>	3,500	
9	CaCl <sub>2</sub>	1,102		26	FePO <sub>4</sub>	1		44	KCl	770	
10	MgCl <sub>2</sub>	4,981		27	MgCl <sub>2</sub>	5,143		45	NaHCO <sub>3</sub>	250	
11	KBr	96		28	KBr	100		46	NaCl	28,130	
12	KCl	664		29	KCl	690			TOTAL	36,400	
13	NaHCO <sub>3</sub>	194		30	NaHCO <sub>3</sub>	200		Levring [9]			
14	NaCl	23,277		31	NaCl	24,320		47	CaCl <sub>2</sub>	1,180	
15	NaF	3		32	NaF	3		48	MgCl <sub>2</sub>	2,540	
16	Na <sub>2</sub> SO <sub>4</sub>	3,917		33	Na <sub>2</sub> SiO <sub>3</sub>	2		49	MgSO <sub>4</sub>	3,482	
17	SrCl <sub>2</sub>	24		34	Na <sub>2</sub> SO <sub>4</sub>	4,060		50	KCl	770	
	TOTAL	34,284		35	SrCl <sub>2</sub>	26		51	NaHCO <sub>3</sub>	200	
					TOTAL	35,714					

*continued*



## 145. ARTIFICIAL SEA WATER

Salt		Concentration mg/L	Salt		Concentration mg/L	Salt		Concentration mg/L
(A)	(B)		(A)	(B)		(A)	(B)	
Levring [9]			85	NaCl	26,726	124	CoSO <sub>4</sub>	0.2
52	NaCl	28,320	86	Na <sub>2</sub> SiO <sub>3</sub>	2.4	125	FeCl <sub>3</sub>	2
53	NaNO <sub>3</sub>	100	87	Na <sub>2</sub> Si <sub>4</sub> O <sub>9</sub>	1.5	126	MgCl <sub>2</sub>	1,000
54	Na <sub>2</sub> HPO <sub>4</sub>	10		TOTAL	34,442.4	127	MgSO <sub>4</sub>	2,439
	TOTAL	36,602	Chu [6]			128	MnSO <sub>4</sub>	6
Fowler and Allen [7]			88	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	3	129	KH <sub>2</sub> PO <sub>4</sub>	100
55	CaCl <sub>2</sub>	100	89	As <sub>2</sub> O <sub>3</sub>	0.03	130	NaCl	25,000
56	CaSO <sub>4</sub>	1,300	90	BaCl <sub>2</sub>	0.077	131	Na <sub>2</sub> MoO <sub>4</sub>	0.2
57	MgBr <sub>2</sub>	100	91	H <sub>3</sub> BO <sub>3</sub>	26	132	ZnSO <sub>4</sub>	5
58	MgCl <sub>2</sub>	3,800	92	CaCl <sub>2</sub>	1,102	133	EDTA <sup>a</sup>	500
59	MgSO <sub>4</sub>	1,600	93	CuSO <sub>4</sub>	0.036		TOTAL	29,453.6
60	K <sub>2</sub> SO <sub>4</sub>	900	94	FeC <sub>6</sub> H <sub>5</sub> O <sub>7</sub>	0.54	McElroy and Farghaly <sup>a</sup> [13]		
61	NaCl	27,200	95	LiNO <sub>3</sub>	1.1	134	NH <sub>4</sub> NO <sub>3</sub>	1,000
	TOTAL	35,000	96	MgCl <sub>2</sub>	4,981	135	FeCl <sub>3</sub>	0.0288
Lewin <sup>a</sup> [10]			97	MnCl <sub>2</sub>	0.2	136	MgSO <sub>4</sub>	100
62	Ca(NO <sub>3</sub> ) <sub>2</sub>	68	98	KBr	96	137	MnSO <sub>4</sub>	0.0012
63	MgCl <sub>2</sub>	5,000	99	KCl	664	138	KH <sub>2</sub> PO <sub>4</sub>	700
64	K <sub>2</sub> HPO <sub>4</sub>	20	100	KI	0.06	139	NaCl	30,000
65	NaHCO <sub>3</sub>	200	101	RbHCO <sub>3</sub>	0.34	140	Na <sub>2</sub> HPO <sub>4</sub>	1,784
66	NaCl	23,000	102	NaHCO <sub>3</sub>	192		TOTAL	33,584.03
67	Na <sub>2</sub> SO <sub>4</sub>	4,000	103	NaCl	23,476	Bernhard [3]		
	TOTAL	32,288	104	NaF	3	141	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	0.0247
Harvey [8]			105	NaNO <sub>3</sub>	50	142	As <sub>2</sub> O <sub>3</sub>	0.03
68	Ca(NO <sub>3</sub> ) <sub>2</sub>	678	106	Na <sub>2</sub> HPO <sub>4</sub>	5	143	H <sub>3</sub> BO <sub>3</sub>	15
69	FeCl <sub>3</sub>	25	107	Na <sub>2</sub> SiO <sub>3</sub>	4.3	144	CaCl <sub>2</sub>	1,180
70	MgSO <sub>4</sub>	2,439	108	Na <sub>2</sub> SO <sub>4</sub>	3,917	145	CoSO <sub>4</sub>	0.0048
71	KCl	600	109	SrCl <sub>2</sub>	24	146	CuSO <sub>4</sub>	0.0004
72	K <sub>2</sub> HPO <sub>4</sub>	20	110	ZnSO <sub>4</sub>	0.005	147	FeCl <sub>3</sub>	6.2929
73	NaCl	30,000		TOTAL	34,545.69	148	FeC <sub>6</sub> H <sub>5</sub> O <sub>7</sub>	5
	TOTAL	33,762	Rice [14]			149	LiNO <sub>3</sub>	0.9935
McClendon-Gault-Mulholland [12]			111	MgCl <sub>2</sub>	1,171	150	MgCl <sub>2</sub>	2,540
74	Al <sub>2</sub> Cl <sub>6</sub>	13	112	MgSO <sub>4</sub>	1,612	151	MgSO <sub>4</sub>	3,482
75	NH <sub>3</sub>	2	113	MnCl <sub>2</sub>	2	152	MnCl <sub>2</sub>	0.0036
76	H <sub>3</sub> BO <sub>3</sub>	58	114	KCl	700	153	NiCl <sub>2</sub>	0.0041
77	CaCl <sub>2</sub>	1,153	115	KNO <sub>3</sub>	100	154	KBr	9.679
78	LiNO <sub>3</sub>	1.3	116	KH <sub>2</sub> PO <sub>4</sub>	7.5	155	KCl	770
79	MgCl <sub>2</sub>	2,260	117	NaHCO <sub>3</sub>	200	156	KI	0.0785
80	MgSO <sub>4</sub>	3,248	118	NaCl	26,500	157	RbHCO <sub>3</sub>	0.0343
81	H <sub>3</sub> PO <sub>4</sub>	0.2	119	Na <sub>2</sub> SiO <sub>3</sub>	10	158	NaHCO <sub>3</sub>	200
82	KCl	721	120	ZnSO <sub>4</sub>	0.005	159	NaCl	28,320
83	NaHCO <sub>3</sub>	198		TOTAL	30,302.51	160	NaF	3
84	NaBr	58	Vishniac <sup>a</sup> [16]			161	NaNO <sub>3</sub>	100
			121	(NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub>	200	162	Na <sub>2</sub> HPO <sub>4</sub>	10
			122	H <sub>3</sub> BO <sub>3</sub>	1.2	163	SrCl <sub>2</sub>	1.826
			123	CaCl <sub>2</sub>	200	164	ZnSO <sub>4</sub>	0.022
							TOTAL	36,643.99

/1/ Enriched with trace elements [5]. /a/ Iron (0.2 mg) supplied by the following stock culture: 24.9 mg/ml FeSO<sub>4</sub>·7H<sub>2</sub>O, 30 mg/ml EDTA, and 20 mg/ml (NH<sub>4</sub>)<sub>2</sub>CO<sub>3</sub>. /s/ Ethylenediamine tetra-acetic acid. /4/ Trace elements included by preparing one liter of the following solution, then adding 0.05 ml to one liter of culture medium: 2.7 g CaCl<sub>2</sub>, 0.96 g FeCl<sub>3</sub>·6H<sub>2</sub>O, 36 mg MnSO<sub>4</sub>·4H<sub>2</sub>O, and 39 mg CuSO<sub>4</sub>·5H<sub>2</sub>O. Ten mg L-histidine and 10 mg DL-threonine added to improve luminescence. Ten ml glycerol also added.

**Contributor:** Jones, Galen E.

**References:** [1] Allen, E. J. 1914. J. Marine Biol. Assoc. U. K. 10:417. [2] Allen, E. J., and E. W. Nelson. 1907. Ibid. 8:42. [3] Bernhard, M. 1957. Pubbl. Staz. Zool. Napoli 29:80. [4] Brujewicz, S. W. 1931. In N. N. Subow, et al., ed. Oceanographical tables. Oceanographical Institute, Hydro-Meteorological Committee of USSR, Moscow. p. 146. [5] Burkholder, P. R., and L. G. Nickell. 1949. Botan. Gaz. 110:426. [6] Chu, S. P. 1949. Sci. Technol. China 2(3):38. [7] Fowler, G. H., and E. J. Allen. 1928. Science of the sea. Clarendon Press, Oxford. p. 68. [8] Harvey, G. W. Unpublished. Scripps Institution of Oceanography, La Jolla, Calif.

*continued*

## 145. ARTIFICIAL SEA WATER

[9] Levring, T. 1946. *Fysiograf. Saellskap.* 16(7):1. [10] Lewin, R. A. 1955. *Can. J. Botany* 33:5. [11] Lyman, J., and R. H. Fleming. 1940. *J. Marine Res. (Sears Found. Marine Res.)* 3:134. [12] McClendon, J. F., C. C. Gault, and S. Mulholland. 1917. *Carnegie Inst. Wash. Publ.* 251. [13] McElroy, W. D., and A.H. Farghaly. 1948. *Arch. Biochem.* 17(3):379. [14] Rice, T. R. 1956. *Limnol. Oceanog.* 1(2):123. [15] Robertson, J. D., and D. A. Webb. 1939. *J. Exptl. Biol.* 16:155. [16] Vishniac, H. S. 1955. *Trans. N. Y. Acad. Sci.* 17:352. [17] ZoBell, C. E. 1946. *Marine microbiology. Chronica Botanica*, Waltham, Mass. p. 21.

## 146. NORMAL SOLUTIONS

To prepare a 0.1 N solution, add the specified number of grams of the compound to 1,000 milliliters of solvent.

	Compound	Formula	Grams		Compound	Formula	Grams
	(A)	(B)	(C)		(A)	(B)	(C)
1	Ammonium molybdate	(NH <sub>4</sub> ) <sub>2</sub> MoO <sub>4</sub>	9.79	11	Potassium thiocyanate	KSCN	9.71
2	Ammonium oxalate	(NH <sub>4</sub> ) <sub>2</sub> C <sub>2</sub> O <sub>4</sub>	10.6	12	Silver nitrate	AgNO <sub>3</sub>	16.9
3	Barium chloride	BaCl <sub>2</sub>	12.2	13	Sodium chloride	NaCl	5.9
4	Copper sulfate	CuSO <sub>4</sub>	12.5	14	Sodium hydroxide	NaOH	4.0
5	Ferric chloride	FeCl <sub>3</sub>	5.4	15	Sodium sulfide	Na <sub>2</sub> S	3.90
6	Hydrobromic acid	HBr	8.1	16	Sodium thiosulfate	Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub>	24.80
7	Hydrochloric acid <sup>1</sup>	HCl	3.65	17	Strontium chloride	SrCl <sub>2</sub>	7.93
8	Potassium dichromate	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	4.90	18	Sulfuric acid <sup>1</sup>	H <sub>2</sub> SO <sub>4</sub>	9.80
9	Potassium hydroxide	KOH	5.6	19	Zinc nitrate	Zn(NO <sub>3</sub> ) <sub>2</sub>	9.47
10	Potassium permanganate	KMnO <sub>4</sub>	3.16				

<sup>1/1</sup> Concentrated.

*Contributor:* Carleton, Ralph K.

*References:* [1] Benedetti-Pichler, A. A. 1942. *Introduction to the microtechnique of inorganic analysis.* J. Wiley, New York. [2] Charlot, G., and R. C. Murray. 1954. *Qualitative inorganic analysis.* Ed. 4. J. Wiley, New York. [3] Feigl, F., and R. E. Oesper. 1954. *Spot tests in organic analysis.* Ed. 6. Elsevier, Amsterdam. [4] Fritz, J. S., and G. S. Hammond. 1957. *Quantitative organic analysis.* J. Wiley, New York. [5] Heisig, G. B. 1950. *Theory and practice of semimicro qualitative analysis.* Ed. 2. W. B. Saunders, Philadelphia. [6] Milton, R. F., and W. A. Waters. 1955. *Methods of quantitative micro-analysis.* Ed. 2. St. Martins, New York. [7] Reedy, J. H. 1938. *Elementary qualitative analysis for college students.* McGraw-Hill, New York. [8] Rieman, W., J. D. Neuss, and B. Naiman. 1942. *Quantitative analysis.* Ed. 2. McGraw-Hill, New York. [9] Sandell, E. B. 1959. *Colorimetric determination of traces of metals.* Ed. 3. Interscience, New York. [10] Scott, W. W., ed. 1938. *Standard methods of chemical analysis.* Ed. 5. Van Nostrand, New York. [11] Snell, F. D., and C. T. Snell. 1959. *Colorimetric methods of analysis.* Ed. 3. Van Nostrand, New York. v. 2A. [12] Vogel, A. I. 1961. *A textbook of quantitative inorganic analysis.* Ed. 3. Longmans, Green; London. [13] Vosburgh, W. C. 1938. *Introductory qualitative analysis.* Macmillan, New York. [14] Welcher, F. J. 1947-48. *Organic analytical reagents.* Van Nostrand, New York. [15] Young, R. S. 1953. *Industrial inorganic analysis.* J. Wiley, New York.

## 147. BUFFER SOLUTIONS: pH RANGES

	Acidic Component	Alkaline Component	pH Range
	(A)	(B)	(C)
1	Hydrochloric acid	Glycine	1.0-3.7
2	Hydrochloric acid	Potassium hydrogen phthalate	2.2-4.0
3	Citric acid	Disodium hydrogen phosphate	2.2-8.0
4	Acetic acid	Sodium acetate	3.7-5.6

*continued*

## 147. BUFFER SOLUTIONS: pH RANGES

	Acidic Component	Alkaline Component	pH Range
	(A)	(B)	(C)
5	Potassium hydrogen phthalate	Sodium hydroxide	4.0-6.2
6	Potassium dihydrogen phosphate	Sodium hydroxide	5.8-8.0
7	Hydrochloric acid	Tris(hydroxymethyl)aminomethane	7.0-9.0
8	Diethylbarbituric acid	Sodium diethylbarbiturate	7.0-9.2
9	Hydrochloric acid or boric acid	Borax	8.0-9.2
10	Glycine	Sodium hydroxide	8.2-10.1
11	Borax	Sodium hydroxide	9.2-11.0
12	Sodium bicarbonate	Sodium hydroxide	9.6-11.0
13	Disodium hydrogen phosphate	Sodium hydroxide	11.0-12.0

*Contributor:* Bates, Roger G.

*References:* [1] Bates, R. G., and V. E. Bower. 1956. *Anal. Chem.* 28:1322. [2] Bower, V. E., and R. G. Bates. 1955. *J. Res. Natl. Bur. Std.* 55:197. [3] Clark, W. M. 1928. *The determination of hydrogen ions*. Ed. 3. Williams and Wilkins, Baltimore. p. 192. [4] Clark, W. M., and H. A. Lubs. 1916. *J. Biol. Chem.* 25:479. [5] Cohn, E. J., F. F. Heyroth, and M. F. Menkin. 1928. *J. Am. Chem. Soc.* 50:696. [6] McIlvaine, T. C. 1921. *J. Biol. Chem.* 49:183. [7] Sørensen, S. P. L. 1912. *Ergeb. Physiol.* 12:393.

## 148. WEAK ACIDS AND BASES: pK VALUES

pK values (dissociation constants) were determined at 25°C. pK values quoted for bases are for the *acidic* dissociation process. Thus the pK value 9.245, given for ammonia, corresponds to the dissociation  $\text{NH}_4^+ \rightarrow \text{H}^+ + \text{NH}_3$ . The *basic* dissociation constant, corresponding to the dissociation  $\text{NH}_3 + \text{H}_2\text{O} \rightarrow \text{NH}_4^+ + \text{OH}^-$ , is 4.752 (13.997 minus 9.245), where 13.997 is the dissociation constant of water at 25°C.

	Acid or Base	pK		Acid or Base	pK
	(A)	(B)		(A)	(B)
1	Acetic acid	4.756	26	Isonicotinic acid	pK <sub>1</sub> 1.84; pK <sub>2</sub> 4.86
2	Alanine	pK <sub>1</sub> 2.348; pK <sub>2</sub> 9.866	27	Lactic acid	3.860
3	<i>p</i> -Aminobenzoic acid	pK <sub>1</sub> 2.413; pK <sub>2</sub> 4.853	28	Maleic acid	pK <sub>1</sub> 1.921; pK <sub>2</sub> 6.225
4	Ammonia	9.245	29	Malonic acid	pK <sub>1</sub> 2.847; pK <sub>2</sub> 5.696
5	Aniline	4.603	30	Methylamine	10.624
6	Ascorbic acid	4.25	31	Nicotinic acid	pK <sub>1</sub> 2.07; pK <sub>2</sub> 4.81
7	Benzoic acid	4.201	32	<i>p</i> -Nitrophenol	7.156
8	Boric acid	9.234	33	Oxalic acid	pK <sub>1</sub> 1.271; pK <sub>2</sub> 4.266
9	Carbonic acid	pK <sub>1</sub> 6.352; pK <sub>2</sub> 10.329	34	Oxaloacetic acid	pK <sub>1</sub> 2.555; pK <sub>2</sub> 4.370
10	Chloroacetic acid	2.865	35	Phenol	9.998
11	Citric acid	pK <sub>1</sub> 3.128; pK <sub>2</sub> 4.761; pK <sub>3</sub> 6.396	36	Phenylacetic acid	4.307
12	Diethylbarbituric acid	7.980	37	Phosphoric acid	pK <sub>1</sub> 2.148; pK <sub>2</sub> 7.198; pK <sub>3</sub> 12.375
13	Dihydroxytartaric acid	pK <sub>1</sub> 1.947; pK <sub>2</sub> 4.004	38	<i>o</i> -Phthalic acid	pK <sub>1</sub> 2.950; pK <sub>2</sub> 5.408
14	Ethylenediamine	pK <sub>1</sub> 6.838; pK <sub>2</sub> 9.960	39	Proline	pK <sub>1</sub> 1.952; pK <sub>2</sub> 10.640
15	Formic acid	3.752	40	Propionic acid	4.874
16	Fumaric acid	pK <sub>1</sub> 3.019; pK <sub>2</sub> 4.384	41	Pyridine	5.22
17	Glucose-1-phosphoric acid	pK <sub>2</sub> 6.503	42	Pyrrolidine	11.305
18	Glycine	pK <sub>1</sub> 2.350; pK <sub>2</sub> 9.780	43	Pyruvic acid	2.490
19	Glycolic acid	3.831	44	Salicylic acid	pK <sub>1</sub> 2.996; pK <sub>2</sub> 13.59
20	Histidine	pK <sub>1</sub> 1.82; pK <sub>2</sub> 6.00; pK <sub>3</sub> 9.17	45	Succinic acid	pK <sub>1</sub> 4.207; pK <sub>2</sub> 5.638
21	Hydrocyanic acid	9.22	46	Sulfuric acid	pK <sub>2</sub> 1.983
22	Hydrofluoric acid	3.17	47	Sulfurous acid	pK <sub>1</sub> 1.76; pK <sub>2</sub> 7.20
23	Hydrogen sulfide	7.06	48	Tartaric acid	pK <sub>1</sub> 3.033; pK <sub>2</sub> 4.366
24	Hydroxylamine	5.96	49	Tartronic acid	pK <sub>1</sub> 2.366; pK <sub>2</sub> 4.735
25	Iodic acid	0.775	50	Tris(hydroxymethyl)-aminomethane	8.075

*Contributor:* Robinson, R. A.

*Reference:* Robinson, R. A., and R. H. Stokes. 1959. *Electrolyte solutions*. Ed. 2. Butterworth, London.

## 149. ACID - BASE INDICATORS: pH RANGES

A solution containing 0.1% indicator is generally satisfactory. Although some of the indicators are water-soluble, 70-90% ethanol is recommended as the solvent for most.

Indicator	pH Range	Acid Color	Basic Color	Indicator	pH Range	Acid Color	Basic Color
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
1 Methyl violet 6B	0.1-1.5	Yellow	Blue	13 Neutral red	6.8-8.0	Red	Yellow
2 Thymol blue (acid range)	1.2-2.8	Red	Yellow	14 Cresol red	7.2-8.8	Yellow	Red
3 Tropaeolin 00	1.4-3.2	Red	Yellow	15 Thymol blue (alk. range)	8.0-9.6	Yellow	Blue
4 Dimethyl yellow	2.8-4.0	Red	Yellow	16 Phenolphthalein	8.2-10.0	Colorless	Pink
5 Methyl orange	3.1-4.4	Red	Yellow	17 Thymolphthalein	9.4-10.6	Colorless	Blue
6 Bromphenol blue	3.2-4.6	Yellow	Blue	18 Alizarin yellow R	10.2-12.0	Yellow	Red
7 Bromcresol green	3.8-5.4	Yellow	Blue	19 Nitramine	10.8-12.8	Colorless	Orange-brown
8 Methyl red	4.2-6.2	Red	Yellow	20 Trapaeolin 0	11.2-12.8	Yellow	Orange-brown
9 Chlorphenol red	5.0-6.8	Yellow	Red	21 1, 3, 5-Trinitrobenzene	12.0-14.0	Colorless	Orange
10 Bromcresol purple	5.2-6.8	Yellow	Purple				
11 Bromthymol blue	6.0-7.6	Yellow	Blue				
12 Phenol red	6.6-8.2	Yellow	Red				

Contributor: ZoBell, Claude E.

References: [1] Britton, H. T. S. 1956. Hydrogen ions. Ed. 4. Chapman, London. [2] Clark, W. M. 1928. Determination of hydrogen ions. Williams and Wilkins, Baltimore. [3] Clark, W. M., and H. A. Lubs. 1917. J. Bacteriol. 2(1):191. [4] Conn, H. J. 1961. Biological stains. Williams and Wilkins, Baltimore. [5] Gold, V. 1956. pH measurements. Methuen, London. [6] Kolthoff, I. M., and C. Rosenblum. 1937. Acid-base indicators. Macmillan, New York.

## 150. OXIDATION - REDUCTION INDICATORS

Redox potential ( $E'_0$ ) values are for varying pH at 30°C. Asterisk (\*) denotes that indicator is unstable at designated pH.

Indicator	$E'_0$ (volts) at				
	pH 5.0	pH 6.0	pH 7.0	pH 8.0	pH 9.0
(A)	(B)	(C)	(D)	(E)	(F)
1 <i>m</i> -Bromophenol indophenol	+0.374	+0.311	+0.248	+0.179	+0.102
2 <i>o</i> -Chlorophenol indophenol	*	+0.301	+0.233	+0.155	+0.082
3 Phenol indophenol	*	+0.286	+0.227	+0.155	+0.083
4 Phenol blue	+0.365	+0.290	+0.224	+0.163	+0.091
5 2, 6-Dichlorophenol indophenol	+0.366	+0.295	+0.217	+0.150	+0.089
6 <i>m</i> -Cresol indophenol	*	+0.272	+0.208	+0.148	+0.076
7 <i>o</i> -Cresol indophenol	*	+0.256	+0.191	+0.130	+0.057
8 Thymol indophenol	*	+0.233	+0.174	+0.110	+0.041
9 <i>m</i> -Toluylenediamine indophenol	+0.195	+0.157	+0.125	+0.088	+0.037
10 Toluylene blue	+0.221	+0.162	+0.115	+0.082	+0.051
11 Thionine (Lauth's violet)	+0.138	+0.094	+0.062	+0.030	-0.001
12 Cresyl blue	+0.149	+0.089	+0.047	+0.015	-0.016
13 Galloxyanine	*	+0.080	+0.021	-0.037	-0.095
14 Methylene blue	+0.101	+0.047	+0.011	-0.020	-0.050
15 Toluidine blue	+0.087	+0.042	-0.005	-0.047	-0.099
16 Janus green (blue-red)	+0.050	+0.002	-0.035	-0.080	-0.115
17 Indigo tetrasulphonate	+0.065	+0.006	-0.046	-0.083	-0.114
18 Methyl capri blue	+0.038	-0.021	-0.060	-0.093	-0.123
19 Indigo trisulphonate	+0.032	-0.028	-0.081	-0.121	-0.152
20 Indigo disulphonate	+0.010	-0.069	-0.125	-0.167	-0.199
21 Gallophenine	-0.003	-0.077	-0.142	-0.202	-0.263
22 Brilliant alizarine blue	-0.040	-0.112	-0.173	-0.226	-0.279
23 Phenosafranine	-0.159	-0.219	-0.252	-0.283	-0.313
24 Tetramethyl phenosafranine	-0.156	-0.225	-0.273	-0.305	-0.336
25 Safranin T	-0.198	-0.250	-0.289	-0.318	-0.348
26 Neutral red	-0.204	-0.275	-0.325	-0.380	-0.435
27 Rosindone sulphonate No. 6	-0.287	-0.338	-0.385	-0.441	-0.508
28 Hydrogen electrode (theoretical)	-0.300	-0.361	-0.421	-0.481	-0.541

continued



## 150. OXIDATION - REDUCTION INDICATORS

Contributor: ZoBell, Claude E.

- References: [1] Clark, W. M. 1925. Chem. Rev. 2:127. [2] Clark, W. M., et al. 1928. U. S. Hyg. Lab. Bull. 151. [3] Cohen, B., and M. Phillips. 1929. Public Health Rept. (U.S.), Suppl. 74. [4] Cohen, B., and P. W. Preisler. 1931. Ibid., Suppl. 92. [5] Hewitt, L. F. 1950. Oxidation-reduction potentials in bacteriology and biochemistry. E. and S. Livingstone, Edinburgh. [6] Michaelis, L., and L. B. Flexner. 1930. Oxidation-reduction potentials. J. B. Lippincott, Philadelphia. [7] Whitehead, T. H., and C. C. Wills. 1941. Chem. Rev. 29:69. [8] ZoBell, C. E. 1946. Bull. Am. Assoc. Petrol. Geol. 30(4):477.

## 151. RADIONUCLIDES USED IN BIOLOGICAL RESEARCH

**Type** (column D):  $\alpha$  = alpha particle, a positively charged particle emitted by radioactive atomic nuclei at high speed, and having a mass number 4 and atomic number 2;  $\beta^-$  = beta particle, an electron, negative ( $\beta^-$  = negatron) or positive ( $\beta^+$  = positron), emitted from the nucleus during radioactive disintegration;  $\gamma$  = gamma ray, electromagnetic radiation of short wavelength and correspondingly high frequency, emitted by the nucleus of an atom in the course of radioactive decay;  $\epsilon$  = orbital electron capture, radioactive transformation occurring when a bound electron merges with the nucleus, converting a proton to a neutron, with liberation of energy in the form of a monoenergetic neutrino plus a photon of X ray characteristic of the new substance (it is a type of beta decay).

Radionuclide					Radiation Emitted					Radionuclide					Radiation Emitted				
Z <sup>1</sup>	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>3</sup> (% Disintegration) <sup>4</sup>	Z:	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>3</sup> (% Disintegration) <sup>4</sup>	(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)
1	1	H <sup>3</sup>	12.26 yr	$\beta^-$	0.02(100)	33	27	Co <sup>60</sup>	5.2 yr	$\beta^-$	0.31(100)				34				1.33(100), 1.17(100)
2	6	C <sup>11</sup>	20.4 min	$\beta^+$	0.97(100)	35	29	Cu <sup>64</sup>	12.8 hr	$\beta^-$	0.57(38)				36				0.66(19)
3	6	C <sup>14</sup>	5,770 yr	$\beta^-$	0.16(100)	37				$\beta^+$	0.0075(42)				38				1.34(1)
4	11	Na <sup>22</sup>	2.6 yr	$\beta^+$	0.54(90)	39	30	Zn <sup>65</sup>	244 da	$\beta^+$	0.32(2)				40				0.008(98)
5				$\epsilon$	0.001(10)	41				$\epsilon$	0.008(98)				42	31	Ga <sup>72</sup>	14.3 hr	$\beta^-$
6				$\gamma$	1.28(100)	43				$\gamma$	1.11(49)				44	33	As <sup>76</sup>	26.5 hr	$\beta^-$
7	11	Na <sup>24</sup>	15 hr	$\beta^-$	1.39(100)	46	35	Br <sup>82</sup>	35.7 hr	$\beta^-$	0.44(100)				48	37	Rb <sup>86</sup>	18.7 da	$\beta^-$
8				$\gamma$	2.75(100), 1.37(100)	49				$\gamma$	1.08(10)				50	38	Sr <sup>89</sup>	54 da	$\beta^-$
9	15	P <sup>32</sup>	14.2 da	$\beta^-$	1.71(100)	51	38	Sr <sup>90</sup>	28 yr	$\beta^-$	0.54(100)				52	39	Y <sup>90</sup>	64 hr	$\beta^-$
10	16	S <sup>35</sup>	87.1 da	$\beta^-$	0.17(100)	53	47	Ag <sup>111</sup>	7.5 da	$\beta^-$	1.06(93), 0.81(1), 0.73(6)				54				0.34(6), 0.25(1)
11	17	Cl <sup>36</sup>	3 x 10 <sup>5</sup> yr	$\beta^-$	0.71(98)	55	49	In <sup>113m</sup>	1.7 hr	$\gamma$	0.39(65)								
12				$\epsilon$	0.002(2)														
13	17	Cl <sup>38</sup>	37.3 min	$\beta^-$	4.81(53), 2.77(16), 1.11(31)														
14				$\gamma$	2.15(57), 1.60(43)														
15	19	K <sup>42</sup>	12.4 hr	$\beta^-$	3.54(81), 1.98(18)														
16				$\gamma$	1.53(18)														
17	20	Ca <sup>45</sup>	164 da	$\beta^-$	0.25(100)														
18	24	Cr <sup>51</sup>	27.8 da	$\epsilon$	0.0049(100)														
19				$\gamma$	0.32(10)														
20	25	Mn <sup>52</sup>	5.7 da	$\beta^+$	0.58(33)														
21				$\epsilon$	0.0054(65)														
22				$\gamma$	1.45(100), 0.94(100), 0.73(100)														
23	25	Mn <sup>54</sup>	320 da	$\epsilon$	0.0054(100)														
24				$\gamma$	0.84(100)														
25	26	Fe <sup>55</sup>	2.7 yr	$\epsilon$	0.0059(100)														
26	26	Fe <sup>59</sup>	45 da	$\beta^-$	0.46(53), 0.27(46), 0.13(1)														
27				$\gamma$	1.29(43), 1.10(57), 0.19(3)														
28	27	Co <sup>57</sup>	270 da	$\epsilon$	0.0064(100)														
29				$\gamma$	0.14(6), 0.12(92), 0.01(9)														
30	27	Co <sup>58</sup>	71 da	$\beta^+$	0.47(15)														
31				$\epsilon$	0.0064(85)														
32				$\gamma$	0.81(100)														

<sup>1</sup>/ Z = atomic number. <sup>2</sup>/ Time required for the amount of a radioactive nuclide to decay to half its initial value. <sup>3</sup>/ Mev = million electron volts. <sup>4</sup>/ Values in boldface indicate percent of nuclear transmutations caused by electron capture.

continued

## 151. RADIONUCLIDES USED IN BIOLOGICAL RESEARCH

Radionuclide					Radiation Emitted					Radionuclide					Radiation Emitted					
Z <sup>1</sup>	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>3</sup> (% Disintegration) <sup>4</sup>	Z <sup>1</sup>	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>5</sup> (% Disintegration) <sup>4</sup>	Z <sup>1</sup>	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>5</sup> (% Disintegration) <sup>4</sup>	Z <sup>1</sup>	Symbol and Mass No.	Half-Life <sup>2</sup>	Type	Energies, Mev <sup>5</sup> (% Disintegration) <sup>4</sup>	
(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	
56	50 Sn <sup>113<sup>s</sup></sup>	118 da	ε	0.024(100)	68	53 I <sup>132</sup>	2.33 hr	β <sup>-</sup>	2.12(18), 1.53(24), 1.16(23), 1.0(20), 0.7(15)	69			γ	2.2(2), 1.9(4), 1.40(13), 1.16(10), 0.96(23), 0.78(94), 0.67(100), 0.62(6), 0.53(28)	70	55 Cs <sup>134</sup>	2.2 yr	β <sup>-</sup>	0.89(1), 0.65(75), 0.28(3), 0.09(20)	
57			γ	0.26(2)	71			γ	1.37(3), 1.31(1), 1.17(3), 0.80(8), 0.79(85), 0.60(97), 0.57(10), 0.56(12), 0.47(2), 0.20(10)	72	55 Cs <sup>137<sup>e</sup></sup>	30 yr	β <sup>-</sup>	1.8(8), 0.51(92)	73	56 Ba <sup>137<sup>m</sup></sup>	2.6 min	γ	0.66(81)	
58	51 Sb <sup>124</sup>	60 da	β <sup>-</sup>	2.31(23), 1.66(2), 1.59(7), 0.95(6), 0.61(51), 0.22(11)	74	79 Au <sup>198</sup>	2.69 da	β <sup>-</sup>	0.96(99), 0.28(1)	75			γ	0.68(1), 0.41(95)	76	80 Hg	2.7 da	ε	0.068(100)	
59			γ	2.11(7), 1.61(46), 1.45(2), 1.37(11), 1.33(2), 1.05(2), 0.97(3), 0.72(14), 0.64(7), 0.60(98)	77			γ	0.08(19)	78	80 Hg <sup>203</sup>	47 da	β <sup>-</sup>	0.21(100)	79			γ	0.28(81)	
60	52 Te <sup>121</sup>	17 da	ε	0.026(100)	80	88 Ra <sup>226<sup>7</sup></sup>	1,620 yr	α, β <sup>-</sup> , γ	Many	61			γ	0.58(87), 0.51(13), 0.07(2)						
62	53 I <sup>125</sup>	60 da	ε	0.027(100)					63			γ	0.04(7)							
63			γ	0.04(7)					64	53 I <sup>130</sup>	12.6 hr	β <sup>-</sup>	1.02(46), 0.60(54)							
64			γ	1.15(31), 0.74(69), 0.66(100), 0.53(100), 0.41(23)					65			γ	0.81(1), 0.61(87), 0.34(9), 0.25(3)							
66	53 I <sup>131</sup>	8.05 da	β <sup>-</sup>	0.81(1), 0.61(87), 0.34(9), 0.25(3)					66			β <sup>-</sup>	0.72(3), 0.64(9), 0.51(1), 0.36(80), 0.28(5), 0.08(2)							
67			γ	0.72(3), 0.64(9), 0.51(1), 0.36(80), 0.28(5), 0.08(2)																

/1/ Z = atomic number. /2/ Time required for the amount of a radioactive nuclide to decay to half its initial value. /3/ Mev = million electron volts. /4/ Values in boldface indicate percent of nuclear transmutations caused by electron capture. /5/ Decays to In<sup>113m</sup>. /6/ Decays to Ba<sup>137m</sup>. /7/ In equilibrium with its gamma-emitting decay products.

**Contributors:** (a) Kahn, Bernd, (b) Siri, William E., (c) Way, Katharine

**References:** [1] Slack, L., and K. Way. 1959. Radiations from radioactive atoms in frequent use. U.S. Atomic Energy Commission, Washington, D. C. [2] Strominger, D., J. M. Hollander, and G. T. Seaborg. 1958. Rev. Mod. Phys. 30:585. [3] U.S. National Bureau of Standards. 1961. Handbook 80. U.S. Govt. Printing Office, Washington, D. C.

## 152. ANESTHETICS

**Dose and Route** (column C): iv = intravenous; rec = rectal; po = oral; ip = intraperitoneal; im = intramuscular.

Drug	Animal	Dose and Route	Time to Act min	Duration	Remarks	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
<b>Mammalia<sup>1</sup></b>							
1	Anavenol-K	Sheep and other ungulates	10 ml/50 kg (about 20 mg/kg); iv <sup>2</sup>	....	20 min	Inject at 1 ml/sec <sup>3</sup>	16
2	Avertin	Cat	300 mg/kg as a 2.5% solution; rec	10	24 hr	Long duration due to lengthy recovery	17
3		Dog	500 mg/kg as a 2.5% solution; rec	....	.....	.....	17
4		Chipmunk	6.02 g/lb; po	1	3.5 hr	Lowers respiratory rate	13
5		Opossum	0.11 g/lb; po	6	10+ hr	.....	13

/1/ For information on additional species, consult reference 12. /2/ Injection of additional half-dose ten minutes later prolongs action. /3/ Spinal block occurs; visual and auditory function persists.

*continued*

## 152. ANESTHETICS

Drug	Animal	Dose and Route	Time to Act min	Duration	Remarks	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
<b>Mammalia<sup>1</sup></b>							
6	Nembutal	Small mammals	1/8-1/2 g/lb; iv <sup>4</sup>	....	15 min	Death due to shock is a general hazard; recovery long and violent	17
7		Ferret	1 gr in 1 ml at 0.45 gr/500 g body wt; ip	....	30 min	0.6-1.0 gr is dangerous dose	17
8		Fox, rabbit, raccoon, squirrel	1 gr/lb; ip	....	.....	No ill effects	14
9		Mouse	1/25-1/50 gr for 15-20 g mouse; ip	6-15	1-5 hr	1 g in 10 ml water	8
10		Rat	3-5 mg/100 g in 6% solution; ip	....	15 min	.....	17
11		Sheep	1/5 gr/lb; iv	....	30 min	Administer over 2 minutes	1
<b>Aves</b>							
12	Avertin	Blackbird	0.15 g/lb; po (corn bait)	7	1.5 hr	Drug deteriorates rapidly (4 hr)	13
13		Starling	0.12 g/lb; po (corn bait)	1	3+ hr	Drug deteriorates rapidly (4 hr)	13
14		Wild turkey	0.06-0.09 g/lb; po (corn bait)	5-17	1-3 hr	Drug deteriorates rapidly (4 hr)	13
15		Pheasant	0.3 ml/kg in 3% solution; rec	....	.....	.....	3
16	Chloral hydrate	Domestic fowl	0.2-0.4 g/ml normal saline; iv	1-2	20-60 min	0.4-0.6 g is dangerous dose	10
17	Equithesin	Many species	2.0-2.5 ml/kg; im	1-2	1+ hr	2.5+ ml/kg is dangerous dose	7
18	Nembutal	Many species	60 mg/kg; im (pectoral)	....	.....	.....	6
19		Domestic fowl	1 ml/5 lb; iv	....	1-2 hr	.....	15
20		Domestic pigeon	1.5 ml; im	....	.....	Repeat injections: 1/2 of original	2
21		Robin	0.1 ml; im	....	.....	.....	2
22		Song sparrow	0.5 ml; im	....	.....	.....	2
<b>Reptilia and Amphibia<sup>5</sup></b>							
23	Avertin	Coachwhip snake	0.8 g/lb; po	3-5	6+ hr	Drug deteriorates quickly in solution	13
24		Copperhead	0.6 g/lb; po	3	6+ hr	Drug deteriorates quickly in solution	13
25		Snapping turtle	0.8 g/lb; po	5	48 hr	Prolonged drowsiness	13
26	Ether	Many species of snakes and lizards	On demand; inhalation	....	.....	Artificial respiration often necessary	5
27	MS-222 <sup>6</sup>	Axolotl, leopard frog	1:3,000 concentration; immersion	....	<30 min	No adverse effect	4
28		Common newt	1:3,000-1:1,000 concentration; immersion	....	30-60 min	No adverse effect	4, 11
29		Congo eel, mud puppy	1:250 concentration; immersion	....	30-60 min	No adverse effect	4
30		Frog, salamander (embryos & larvae)	1:10,000 concentration; immersion	....	Few min-2 da	No adverse effect	4
31		Leopard frog (larvae)	1:1,000 concentration; immersion	....	15-30 min	No adverse effect	4
32		Tiger salamander	1:2,000 concentration; immersion	....	15-30 min	No adverse effect	4
<b>Pisces and Chondrichthyes<sup>7</sup></b>							
33	MS-222 <sup>6</sup>	Bluegill, bullhead, goldfish	1:3,500 concentration; immersion	....	4-10 min	No adverse effect; temperature, 20°C±1.5°C	4
34		Brook trout, large-mouth bass	1:31,000 concentration; immersion	....	20 min	Direct relation between amount of MS-222 and size of fish	4

/1/ For information on additional species, consult reference 12. /4/ One-quarter of dose in 30-40 seconds, remainder administered slowly. /5/ For information on additional species of Amphibia and Pisces, consult reference 4. /6/ Tricaine methanesulphonate.

*continued*

## 152. ANESTHETICS

Drug	Animal	Dose and Route	Time to Act min	Duration	Remarks	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Pisces and Chondrichthyes <sup>5</sup>							
35	MS-222 <sup>6</sup>	Brown trout	1:5,530 concentration; immersion	....	18-20 min	No adverse effect	4
36		Ray, shark	1 g/L of sea water (1:1,000 concentration); po and sprayed over gill exits of pharynx	<1	.....	100-1,000 ml of solution, depending on size of animal; recovery within 5-30 minutes after return to water	9
Turbellaria							
37	MS-222 <sup>6</sup>	Planarian	1:1,500 concentration; immersion	15	.....	Recovery approximately 30 minutes after return to well water	11

<sup>5/</sup> For information on additional species of Amphibia and Pisces, consult reference 4. <sup>6/</sup> Tricaine methanesulphonate.

*Contributors:* (a) Cowan, Ian McTaggart, (b) Achor, Leonard B.

*References:* [1] Awad, F. I. 1954. Vet. Record 66:226. [2] Bailey, R. E. 1953. Auk 70:497. [3] Balfour-Jones, S. E. 1936. Proc. Roy. Soc. Med. 29:709. [4] Bové, F. J. 1962. Sandoz News 3. [5] Brazenor, C. W., and K. Geoffrey. 1953. Copeia (3):165. [6] Durant, A. J. 1953. J. Am. Vet. Med. Assoc. 122:14. [7] Gandal, C. P. 1956. Ibid. 128:332. [8] Gates, W. H. 1932. Science 76:349. [9] Gilbert, P. W., and F. G. Wood. 1957. Science 126:212. [10] Hole, N. 1933. J. Comp. Pathol. Therap. 46:47. [11] Manner, H. W. 1957. Turtox News 35:134. [12] Mosby, H. S., ed. 1960. Manual of game investigational techniques. Edwards Brothers, Ann Arbor. [13] Mosby, H. S., and D. E. Cantner. 1956. Southwestern Vet. 9:132. [14] Rausch, R. 1947. J. Wildlife Management 11:189. [15] Warren, D. C., and H. M. Scott. 1953. Poultry Sci. 14:195. [16] Watson, D. F. 1953. North Am. Vet. 34:334. [17] Wright, J. G. 1952. Veterinary anesthesia. Ed. 3. Williams and Wilkins, Baltimore.

## 153. FIXATIVES AND CLEARING AGENTS

### Part I. FIXATIVES

For additional information, consult references 4 and 6. Formol or formalin = 36-40% aqueous solution of formaldehyde, U.S.P. Alcohol may be ethanol, methanol, or isopropanol.

Fixative	Recommended Uses	Ingredients	Duration of Application	Subsequent Treatment	Reference	
(A)	(B)	(C)	(D)	(E)	(F)	
1	Allen's B-15	Mammalian cytology	75 ml saturated aqueous solution of picric acid <sup>1</sup> , 25 ml formol, 5 ml glacial acetic acid, 2 g urea. Just before use, add 1.5 g chromic acid.	12-24 hr	Brief tap-water rinse. Alcohol series to 70% overnight (small pieces of material).	5
2	Bouin's (picro-formol-acetic)	General fixative	75 ml saturated aqueous solution of picric acid <sup>2</sup> , 25 ml formol, 5 ml glacial acetic acid.	24+ hr	Brief rinse in 30-50% alcohol. Store in 70% alcohol.	5
3	Carnoy's	Eggs of various invertebrates, glycogen, Nissl substance	10 ml glacial acetic acid, 30 ml chloroform, 60 ml 100% ethanol.	2-12 hr, according to size	Wash in 95% alcohol. Store in 70% alcohol.	5

<sup>1/1</sup> Remove picric acid from spread sections with a few drops of saturated aqueous solution of lithium carbonate in any dilution of alcohol.

*continued*



## 153. FIXATIVES AND CLEARING AGENTS

### Part I. FIXATIVES

Fixative	Recommended Uses	Ingredients	Duration of Application	Subsequent Treatment	Reference
(A)	(B)	(C)	(D)	(E)	(F)
4 Champy's	Small animals; for good detail	0.4 g chromic anhydride CrO <sub>3</sub> , 1.2 g potassium bichromate. Dissolve in 60 ml distilled water; add 40 ml 1% aqueous solution of osmic acid.	6-24 hr	Wash in tap water 12 <sup>+</sup> hr. Store in 70% alcohol.	1
5 Craff	Plants, general fixative	<b>Solution A:</b> 1 g chromic anhydride, 7 g glacial acetic acid, 92 ml distilled water. <b>Solution B:</b> 30 ml formalin, 70 ml distilled water. Just before use, mix equal parts of <b>A</b> and <b>B</b> .	12-24 hr	Wash in tap water 12-24 hr. Store in 70% alcohol.	1
6 Flemming's weak	Minute and delicate objects; for good cytoplasmic detail	0.25 g chromic acid, 0.10 g osmic acid, 0.10 ml glacial acetic acid, 100 ml water.	2-24 hr, according to size; best results at 0.5°C	Wash in tap water 2-24 hr. Store in 70% alcohol.	1
7 Formalin solution	Animals, general fixative	10 ml formalin, 90 ml distilled water.	24 <sup>+</sup> hr	Wash in tap water. Store in 10% formalin or 70% alcohol.	1
8 Formalin-acetic	Plant and animal tissues, general fixative	10 ml formalin, 5 ml glacial acetic acid, 85 ml 70% ethanol.	24 <sup>+</sup> hr	Wash in tap water. Store in 70% alcohol.	1
9 Gilson's	Small invertebrates, amphibian eggs, general purpose	15 ml 70-80% nitric acid, 4 ml glacial acetic acid, 100 ml 60% ethanol, 880 ml distilled water, 20 g mercuric chloride <sup>2</sup> .	6 <sup>+</sup> hr	Wash in tap water. Store in 70% alcohol.	5
10 Helly's	Hematopoietic tissue	2.5 g potassium dichromate, 5 g mercuric chloride <sup>2</sup> , 100 ml distilled water. Just before use, add 5 ml formalin.	6-24 hr	Wash 12 <sup>+</sup> hr in tap water, or in weak formol in dark.	1,3
11 Petrunkevitch's	General purpose	<b>Solution A:</b> 100 ml distilled water, 12 ml nitric acid, 8 g cupric nitrate. <b>Solution B:</b> 100 ml 80% ethanol, 4 g phenol, 6 ml ethyl ether. Just before use, mix 1 part <b>A</b> with 3 parts <b>B</b> .	24-72 hr	Wash in tap water. Store in 70% ethanol.	3
12 Susa-Heidenhain's	General purpose	4.5 g mercuric chloride <sup>2</sup> , 80 ml distilled water. Mix, then add 0.5 g sodium chloride, 2 g trichloroacetic acid, 4 ml glacial acetic acid, 20 ml formalin.	3-24 hr, according to size	Wash in tap water 12-24 hr, or in 70% alcohol.	1
13 Worcester's	Eggs of fish and amphibia; embryos; general fixative	90 ml 10% formalin saturated with mercuric chloride <sup>2</sup> , 10 ml glacial acetic acid.	12-24 hr, according to size	Wash in tap water. Store in 70% alcohol.	5
14 Zenker's	Animal tissues, small pieces	25 g potassium dichromate, 5 g mercuric chloride <sup>2</sup> , 5 ml glacial acetic acid, 100 ml water.	4-24 hr	Wash in tap water 12-24 hr. Store in 70% alcohol.	1

<sup>1/2</sup> Remove mercuric chloride from spread sections with Gram's solution (formula: 5 ml water, 1 g iodine, 2 g potassium iodide; add water to make 300-ml solution). [2]

*Contributor:* Jones, Ruth McClung

*References:* [1] Davenport, H. A. 1960. Histological and histochemical technics. W. B. Saunders, Philadelphia. [2] Gray, P. 1954. The microtome's formulary and guide. Blakiston, New York. [3] Gray, P. 1963. Handbook of basic microtechnique. Ed. 3. McGraw-Hill, New York. [4] Humason, G. L. 1962. Animal tissue techniques. W. H. Freeman, San Francisco. [5] Jones, R. M., ed. 1950. McClung's Handbook of microscopical technique. P. B. Hoeber, New York. [6] Lillie, R. D. 1954. Histopathologic technic and practical histochemistry. Blakiston, New York.

*continued*

## 153. FIXATIVES AND CLEARING AGENTS

### Part II. CLEARING AGENTS

Recommended for use in cytology and histology, unless otherwise indicated. Alcohol may be ethanol, methanol, or isopropanol.

Clearing Agent	Preceding Dehydrator	Method of Application	Duration of Application	No. of Changes	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 Anilin oil	30% alcohol	Immersion in mixtures 50:50 of anilin oil, and 30%, 50%, 70% alcohol; then pure anilin oil	1-2 hr each mixture; 2-4 hr pure anilin	1 for mixture; 2 for pure oil	1
2 Benzene	95 or 100% alcohol, or triethyl phosphate, or cellosolve	Immersion	Until translucent (1-3 hr)	3-5	1
3 Cedar wood oil <sup>1</sup>	80-100% alcohol	Layer alcohol over oil in vial; add specimen	Until specimen sinks to bottom	Fresh oil once after sinking	1
4 Petrolatum ether	95 or 100% alcohol, or triethyl phosphate, or cellosolve	Immersion	Until translucent (1-3 hr)	3-5	2
5 Toluene	95 or 100% alcohol, or triethyl phosphate, or cellosolve	Immersion	Until translucent (1-3 hr)	3-5	1
6 Xylene	95 or 100% alcohol, or triethyl phosphate, or cellosolve	Immersion	Until translucent (approximately 3 hr)	3-5	1

<sup>1</sup>/ For use in cytology only.

*Contributor:* Jones, Ruth McClung

*References:* [1] Jones, R. M., ed. 1950. McClung's Handbook of microscopical technique. P. B. Hoeber, New York. [2] Lillie, R. D. 1954. Histopathologic technic and practical histochemistry. Blakiston, New York

## 154. STAINING METHODS

### Part I. LIVING MATERIALS

Material to be Stained	Animal	Stain	Application	Dose		Subsequent Treatment	Reference
				Unit	No.		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Bone matrix	Young mammals	1-2% aqueous solution of alizarin red; pH 8.0	Intraperitoneal	50-80 mg/kg	1 to several at desired intervals	Fix in 10% formalin (sections not decalcified).	2
2 Eggs	Amphibians, fishes	1% Nile blue sulfate adsorbed on small, dry slips of agar	Apply agar slips to egg surface	15-30 min	1	At desired stage of development, fix in Zenker's fluid; wash briefly. Transfer to 1% aqueous solution of phosphomolybdic acid for 1 hr. Dehydrate in alcohol + 1% phosphomolybdic acid.	2
3	Amphibians, lamprey	1% bismarck brown, adsorbed on small, dry slips of agar	Apply agar slips to egg surface	15-30 min	1	Fix in 8.5 ml 100% alcohol, 15 ml glacial acetic acid. Transfer directly to cedar oil for storage.	2
4 Epithelial cells of convoluted tubules in kidney	Mouse	0.5% trypan blue in sterile water	Intraperitoneal	0.1-0.2 ml	4 at 48 hr intervals	Fix in 10% formalin; embed in paraffin.	2
5 Fats, fat cells	Cat	20% Sudan III, or Sudan IV, in olive oil	Pass by tube to stomach	5-10 ml	1	After 3-7 da, freeze sections of fat deposit areas.	1

*continued*

# Contrails

## 154. STAINING METHODS

### Part I. LIVING MATERIALS

Material to be Stained	Animal	Stain	Application	Dose		Subsequent Treatment	Reference
				Unit	No.		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
6 Mitochondria	Mammals	Janus green-B, 1:10,000 in saline	Ventricle or aorta by gravity pressure, with temporary occlusion of inferior or superior vena cava	About 10 min, or until pancreas shows staining		Examine in saline.	2
7 Blood monocytes, tissue phagocytes	Rabbits	50% Higgins ink in sterile distilled water	Injection into marginal ear vein	5 ml	Daily for 3 da; twice thereafter at 3-da intervals	For blood smears, use Wright's stain. Fix tissues in 10% formalin; embed in paraffin.	2
8 Blood monocytes	Mouse	0.5% trypan blue in sterile water	Intraperitoneal	0.2-0.5 ml	4 at 48 hr intervals	Withdraw blood from tail vein and examine fresh or in dry smear.	1
9 Peripheral nerve endings	Mammals (usually cat)	0.2% aqueous methylene blue at 37°C	Intravenous injection after saline perfusion	Same amount as saline	1	After 30 min, immerse desired part in 0.1% methylene blue, at 37°C, until bluish, then in 8% ammonium molybdate for 30 min. Dehydrate in alcohol.	3
10 Phagocytes, loose connective tissue	Mouse	0.5% trypan blue in sterile water	Subcutaneous	0.2-0.5 ml	4 at 48 hr intervals	Fix in 10% formalin; embed in paraffin.	1
11	Common laboratory mammals	50% Higgins ink in sterile water	Subcutaneous	0.1-1.0 ml according to size		Fix in any good fixative; embed in paraffin.	2
12 Reticulo-endothelial phagocytes of spleen, liver, bone marrow, adrenals, lymph nodes	Mouse	0.5% trypan blue in sterile water	Intraperitoneal	0.2-0.5 ml	4 at 48 hr intervals	Fix in 10% formalin; embed in paraffin.	1
13 Teeth (dentin and enamel)	Young dog, before formation of permanent teeth	1% solution trypan blue in sterile water	Intraperitoneal	2-5 ml	5 at 1, 2, 3, 4, and 5 da intervals	Place head in 10% formalin at 35°C. Dehydrate jaws in alcohol, benzene; impregnate with benzodamar for grinding.	2
14 Teeth	Young dog	1% aqueous solution of alizarin red	Intravenous and/or intraperitoneal	50-80 mg/kg	Several	Fix in formalin; grind.	2

*Contributor:* Jones, Ruth McClung

*References:* [1] Cowdry, E. V. 1952. Laboratory technique. Ed. 3. Williams and Wilkins, Baltimore. [2] Jones, R. M., ed. 1950. McClung's Handbook of microscopical technique. P. B. Hoeber, New York. [3] Lillie, R. D. 1954. Histopathologic technic and practical histochemistry. Blakiston, New York.

*continued*

## 154. STAINING METHODS

### Part II. FIXED MATERIALS

**Procedure** (columns C-G): all solutions are aqueous, unless otherwise stated; H<sub>2</sub>O refers to distilled water; ethanol, without qualification, refers to commercial "190 proof neutral grain spirits," and absolute (abs.) ethanol refers to "pure ethyl alcohol, U.S.P., 200 proof." **Abbreviations:** sat. = saturated; sol. = solution(s).

	Dye or Technique	Recommended Material	Procedure					Reference
			1st step	2nd step	3rd step	4th step	5th step	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Heidenhain's iron hematoxylin	Nuclei, particularly mitotic figures, in sections	Mordant in 2.5% ferric alum overnight.	Rinse in H <sub>2</sub> O. Stain 12-24 hr in 0.5% hematoxylin at least 1 mo old.	Rinse in H <sub>2</sub> O. Differentiate in 2.5% ferric alum.	Place in alkaline tap water until blue. Steps 3 and 4 may be repeated.	Wash, dehydrate, clear, and mount in neutral balsam or synthetic substitute.	23
2	Ehrlich's hematoxylin	Nuclei in sections subsequently to be counterstained	Dissolve 0.7 g hematoxylin in 30 ml ethanol and 3 ml acetic acid. Mix 30 ml sat. sol. ammonia alum with 30 ml glycerol and add to dye sol. Ripen 3 mo.	Take sections to ethanol, then stain 0.5-2 min.	Apply ethanol from drop bottle until nuclei are differentiated.	Place in alkaline tap water until blue.	Wash, dehydrate, clear, and mount in neutral balsam or synthetic substitute.	6
3	Delafield's hematoxylin	Whole mounts of plants or animals	Dissolve 0.6 g hematoxylin in 70 ml sat. sol. ammonia alum; add 15 ml glycerol, 15 ml methanol, and 5 ml ethanol. Ripen 3 mo.	Take object to H <sub>2</sub> O. Stain from 2 min (in full-strength stain for minute objects) to 2 da (in 1:10 dilution for a liver fluke).	Wash in H <sub>2</sub> O until no color comes away.	Differentiate, if necessary, in 0.1% HCl in 70% ethanol. Place in alkaline tap water until blue throughout.	Wash, dehydrate, clear, and mount in neutral balsam or synthetic substitute.	1
4	Gray's celestin blue-B	Nuclei and mitotic figures in sections	Mix 1 g celestin blue-B with 0.5 ml H <sub>2</sub> SO <sub>4</sub> . Break up mass and add slowly, with constant stirring, 100 g 2.5% ferric alum with 14 ml glycerol heated to 50°C. Cool and adjust to pH 0.8 with H <sub>2</sub> SO <sub>4</sub> .	Take sections to H <sub>2</sub> O. Stain 1 min or more (overstaining is impossible).	Rinse in tap water until no color comes away.	Dehydrate, clear, and mount in balsam.		2
5	Johansen's safranin	Nuclei in sections of animal tissue; nuclei and xylem in sections of plant tissues	Dissolve 0.1 g safranin-O in 50 ml methyl cellosolve; add 25 ml ethanol. Dissolve 1 g sodium acetate in 25 ml H <sub>2</sub> O with 2 ml 40% formaldehyde; add to dye sol.	Take sections to H <sub>2</sub> O. Stain 1-3 da.	Differentiate in 0.5% picric acid in 65% ethanol.	Wash in running tap water until yellow color is removed.	Dehydrate, clear, and mount in balsam.	11
6	Bismarck brown	Whole mounts of small plants or animals	Take objects to H <sub>2</sub> O. Stain 1-5 min in 1% bismarck brown.	Wash until no color comes away.	Dehydrate, clear, and mount in balsam.			11
7	Grenacher's alcoholic borax carmine	Whole mounts of plants or animals; nuclei in sections of plant or animal tissues	Boil 1.5 g carmine with 2 g borax in 50 ml H <sub>2</sub> O for 30 min. Cool; add 50 ml 70% ethanol. Leave 2 da and filter.	Take objects to H <sub>2</sub> O. Stain overnight.	Differentiate objects in 0.1% HCl in 70% ethanol until clear pink (sections rarely require differentiation).	Dehydrate, clear, and mount in balsam.		9

continued



## 154. STAINING METHODS

### Part II. FIXED MATERIALS

Dye or Technique	Recommended Material	Procedure					Reference
		1st step	2nd step	3rd step	4th step	5th step	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
8 Mayer's paracarmine	Whole mounts of small invertebrates, particularly marine forms and their larvae	Dissolve 1 g carmine acid in 70% ethanol; add 4 g strontium chloride and 0.5 g aluminum chloride.	Take objects to 50% ethanol. Stain overnight.	Differentiate in 0.1% strontium chloride in 70% ethanol.	Dehydrate, clear, and mount in balsam.		18
9 La Cour's aceto-orcein	Chromosomes in squashes	Dissolve 1 g orcein in equal parts acetic acid and water.	Squash material in stain under cover. Examine at intervals until maximum intensity of stain is reached.	Place slide on "dry ice" until cover slip is well frosted.	Pry off cover slip with safety razor blade. Place slide in 70% ethanol.	Dehydrate, clear, and mount in balsam.	13
10 Eosin-Y; ethyl eosin	Contrast stain for black or blue nuclei, particularly in sections of animal material	Dissolve 0.5 g eosin-Y in H <sub>2</sub> O, or 0.5 g ethyl eosin in ethanol.	Take sections with pre-stained nuclei to H <sub>2</sub> O for eosin-Y, or to ethanol for ethyl eosin.	Place sections in stain until preferred intensity is reached.	Dehydrate, clear, and mount in balsam.		13
11 Light green; fast green FCF	Contrast stain for red nuclei, particularly in sections of plant tissues	Dissolve 0.5 g of either dye in ethanol.	Take sections with pre-stained nuclei to ethanol.	Place sections in stain until preferred intensity is reached.	Dehydrate, clear, and mount in balsam.		13
12 Smith's picrospect blue	Double contrast stain for red nuclei, particularly in sections of embryos with heavy yolk	Dissolve 1 g picric acid in 100 ml abs. ethanol. Saturate with spirit blue (about 1.2 g).	Bulk stain amphibian embryos with Grenacher's alcoholic borax carmine. Cut 10- $\mu$ sections. Take to abs. ethanol.	Place in stain 2 min.	Differentiate in abs. ethanol.	Clear in xylene. Steps 3, 4, and 5 may be repeated. Mount in balsam.	21
13 Shumway's picro-indigo carmine	Double contrast stain for red nuclei, particularly in sections of amniote embryos	Mix equal parts of sat. sol. of picric acid and indigo carmine.	Take sections with pre-stained nuclei to water. Stain 5 min.	Differentiate in 70% ethanol.	Transfer to abs. ethanol for not more than 15 sec.	Clear in xylene, mount in balsam.	20
14 Patay's ponceau 2R-light green	Double contrast stain for blue or black nuclei in sections	Take sections with pre-stained nuclei to H <sub>2</sub> O.	Stain 2 min in 1% ponceau 2R.	Rinse briefly in H <sub>2</sub> O. Differentiate 2 min in 1% phosphomolybdic acid.	Rinse briefly in H <sub>2</sub> O. Stain 30 sec in 0.5% light green in 90% ethanol.	Dehydrate, clear, and mount in balsam.	19
15 Gray's ponceau 2R-orange 2	Double contrast stain for blue or black nuclei in sections	Take sections with pre-stained nuclei to H <sub>2</sub> O.	Stain 2 min in 100 ml H <sub>2</sub> O, 0.4 g ponceau 2R, 0.6 g orange 2.	Blot slides, then dip up and down in abs. ethanol until differentiated.	Clear in xylene, mount in balsam.		8
16 Masson's triple contrast	Triple contrast stain for blue or black nuclei in sections	Take sections with nuclei, pre-stained preferably with Gray's celestin blue-B, to H <sub>2</sub> O.	Stain 5 min in 100 ml H <sub>2</sub> O, 1 ml acetic acid, 0.35 g acid fuchsin, 0.65 g ponceau 2R.	Rinse briefly in H <sub>2</sub> O. Differentiate 5 min in 1% phosphomolybdic acid.	Rinse briefly in H <sub>2</sub> O; drain. Flood slide with sat. sol. anilin blue in 2.5% acetic acid; leave 2-5 min. Differentiate in 1% acetic acid.	Pass directly to 0.1% acetic acid in abs. ethanol. Dehydrate, clear in xylene containing 1% salicylic acid; mount in balsam similarly acidified.	17

*continued*

## 154. STAINING METHODS

### Part II. FIXED MATERIALS

	Dye or Technique	Recommended Material	Procedure					Reference
			1st step	2nd step	3rd step	4th step	5th step	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
17	Mallory's triple stain	General stain for sections of animal tissues	Stain 2 min in 1% acid fuchsin.	Rinse in H <sub>2</sub> O. Differentiate 2 min in 1% phosphotungstic acid.	Rinse in H <sub>2</sub> O. Stain 10 min in 100 ml H <sub>2</sub> O, 0.5 g methyl blue, 1 g orange G2, 1 g oxalic acid.	Wash in H <sub>2</sub> O until no color comes away. Dip up and down in abs. ethanol until differentiated.	Clear in xylene and mount in balsam preferably containing 1% salicylic acid.	15
18	Johansen's quadruple stain	General stain for sections of plant tissues	Stain 24 hr in Johansen's safranin, but do not differentiate. Rinse in H <sub>2</sub> O. Stain 10-15 min in 1% methyl violet.	Differentiate 15 sec in mixture of equal parts ethanol, methyl cellosolve, tertiary butanol.	Stain 10-15 min in sat. sol. of fast green FCF in 13 ml equal parts ethanol and methyl cellosolve, 36 ml ethanol, 36 ml tertiary butanol, 12 ml acetic acid.	Rinse in 0.5% acetic acid in equal parts ethanol and tertiary butanol. Stain 3 min in sat. sol. of orange G in equal parts ethanol, tertiary butanol, and methyl cellosolve.	Rinse in equal parts ethanol, methyl cellosolve, and clove oil. Rinse in equal parts ethanol, clove oil, and xylene. Clear in xylene and mount in balsam.	10
19	Wright's methylene blue-eosin	Blood smears	Prepare blood smear; allow to dry.	Place 3 drops of Wright's stain on smear.	Exactly 1 min later, add 6 drops of phosphate pH 6.4 buffer.	2 min later, wash thoroughly with H <sub>2</sub> O.	Allow slide to dry.	24
20	Anderson's "Weigert-Pal" stain	Demonstration of nerve tracts by staining myelin sheaths in 30- $\mu$ frozen sections	Boil together 100 ml H <sub>2</sub> O, 5 g potassium dichromate, 2.5 g chromic fluoride; cool, filter. To this stock sol., add immediately before use, 10% of 2% calcium hypochlorite. Stain sections 2-3 da.	Transfer sections to stock mordant sol. without hypochlorite for 15-60 min.	Dissolve 0.5 g hematoxylin in 10 ml abs. ethanol; add 3 ml 2% calcium hypochlorite. Shake well; dilute with H <sub>2</sub> O to 100 ml; add 3 ml acetic acid. Transfer sections from mordant, without rinsing, and heat to 50°C for 1 hr.	Transfer sections, without washing, to 100 ml H <sub>2</sub> O, 2.5 g boron dichromate, 1 g sodium sulfate. Wash thoroughly. Transfer to fresh 0.25% potassium permanganate until brown.	Transfer, without washing, to 100 ml H <sub>2</sub> O, 0.5 g potassium sulfite, 0.5 g oxalic acid until bleached. Repeat until differentiation is complete. Wash thoroughly, dehydrate, clear, and mount in balsam.	3
21	Davenport-Windle-Rhine protein silver stain	Demonstration of nerve tracts by staining axis cylinders in paraffin sections	Cut 10- $\mu$ paraffin sections of material fixed in 50 ml H <sub>2</sub> O, 50 ml ethanol, 5 ml <i>p</i> -nitrophenol, 10 ml formamide.	Deparaffinize mounted sections and bring to H <sub>2</sub> O. Place in 5% silver nitrate at 60°C for 1 hr.	Wash in 3 successive, 30-sec changes of H <sub>2</sub> O. Transfer to 0.2% protargol for 1 hr.	Rinse quickly in H <sub>2</sub> O. Transfer to 100 ml H <sub>2</sub> O, 5 g sodium sulfite (desiccated), 1 g hydroquinone, 0.5 g potassium metaborate for 1 min. Wash thoroughly in running water.	Transfer to 0.2% gold chloride until gray. Rinse quickly and place in 0.4% oxalic acid until section just starts to darken. Rinse. Fix in 5% sodium thiosulfate. Wash thoroughly, dehydrate, clear, and mount in balsam.	4

*continued*

## 154. STAINING METHODS

### Part II. FIXED MATERIALS

Dye or Technique	Recommended Material	Procedure					Reference	
		1st step	2nd step	3rd step	4th step	5th step		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
22	Kull's method for mitochondria	Mitochondria in 5- $\mu$ paraffin sections	Fix small pieces for 25 hr in 100 ml H <sub>2</sub> O, 0.5 g osmic acid, 0.5 g chromic acid, 1.25 g potassium dichromate.	Wash 30 min in H <sub>2</sub> O. Transfer to 85 ml H <sub>2</sub> O, 15 ml pyroligneous acid, 1 g chromic acid for 24 hr.	Wash 30 min in H <sub>2</sub> O. Transfer to 3% potassium dichromate for 24 hr. Wash 24 hr in running water. Cut 5- $\mu$ paraffin sections; take sections to H <sub>2</sub> O.	Saturate hot anilin with acid fuchsin. Shake thoroughly with equal quantity of H <sub>2</sub> O. Separate and retain aqueous fraction. Flood sol. on slide and heat to steaming for 1 min.	Rinse quickly in H <sub>2</sub> O. Place in 0.5% toluidine blue for 1-2 min. Rinse quickly in H <sub>2</sub> O. Differentiate 0.5% aurantia in 70% ethanol. Dehydrate in acetone, clear in xylene, and mount in balsam.	12
23	Ludford's method for Golgi apparatus	Golgi in 5- $\mu$ paraffin sections	Fix small pieces (earthworm ovary is excellent) for 1 hr in 100 ml H <sub>2</sub> O, 0.5 g osmic acid, 4 g mercuric chloride, 1 g sodium chloride.	Place specimen on bottom of 30-ml stoppered bottle; add just enough 2% osmic acid to cover specimen. Keep 2 wk at room temperature.	Fill bottle with H <sub>2</sub> O and keep 2 da at 38°C. Wash in running water.	Cut 5- $\mu$ paraffin sections. Mount and deparaffinize with xylene.	Examine sections of Golgi apparatus; if not clear, place in turpentine until differentiated. Wash in xylene, and mount in balsam.	14
24	Hucker's crystal violet	Bacterial smears	Dissolve 2 g crystal violet in 20 ml ethanol. Dissolve 0.8 g ammonium oxalate in 80 ml H <sub>2</sub> O; add to dye sol.	Spread thin film of bacteria and allow to dry. Pass through flame.	Flood dye sol. on smear. Leave 30 sec.	Wash off dye sol. with jet H <sub>2</sub> O from wash bottle.	Allow film to dry.	5
25	Gram's stain	Demonstration of gram-positive bacteria	Follow steps 1, 2, and 3 for Hucker's crystal violet.	Rinse quickly in beaker of H <sub>2</sub> O; add few drops of 0.5% iodine in 1% potassium iodide. Leave 1 min.	Rinse quickly in H <sub>2</sub> O. Place in abs. ethanol until no color comes away.	Stain 5-10 sec in 1% safranin-O. Wash and dry.		5
26	Margolena's stain	Demonstration of parasitic fungi in sections of plant tissues	Dissolve 0.1 g thionine and 5 g phenol in 95 ml H <sub>2</sub> O. Stain sections 1 hr.	Rinse in H <sub>2</sub> O. Place in 0.5% light green in ethanol for 1 min.	Wash in H <sub>2</sub> O until no color comes away. Dehydrate in abs. ethanol.	Mix 30 ml sat. sol. orange G in ethanol with 60 ml sat. sol. erythrosin in clove oil. Stain 1-2 min.	Rinse in clove oil, clear in xylene, and mount in balsam.	16
27	Alizarin red	Demonstration of bone, or calcified areas, in whole mounts of small animals	Fix or preserve specimens in ethanol, or formaldehyde made alkaline with borax.	Bring specimens to 70% ethanol. Stain 1-12 hr in mixture of 1 ml sat. ethanol sol. alizarin red S with 100 ml sat. sol. borax in 70% ethanol.	Transfer to sat. sol. borax in 90% ethanol until no color comes away.	Dehydrate, clear, and mount in balsam.		7

*continued*

## 154. STAINING METHODS

### Part II. FIXED MATERIALS

	Dye or Technique	Recommended Material	Procedure					Reference
			1st step	2nd step	3rd step	4th step	5th step	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
28	Van Wijhe's stain	Demonstration of cartilage in whole mounts of embryos	Preserve specimens in any preservative not containing picric acid.	Wash in running water overnight. Take to 70% ethanol.	Stain 24 hr in 70% ethanol containing 0.1% each of HCl and toluidine blue.	Wash in 70% ethanol containing 0.1% HCl until no color comes away.	Dehydrate, clear, and mount in balsam.	22

*Contributor:* Gray, Peter

*References:* [1] Aly, W., and C. J. Eberth. 1885. Z. Wiss. Mikroskopie 2:282. [2] Amenta, P. S. 1961. Stain Technol. 36:15. [3] Anderson, J. 1922. Lab. J. (London) 5:65. [4] Conn, H. J., and M. A. Darrow, ed. 1946. Staining procedures used by the biological stain commission. Biotechnical Publications, Geneva, N. Y. [5] Conn, H. J., F. B. Mallory, and F. Parker, Jr. 1929. In C. E. McClung, ed. Handbook of microscopical technique. P. B. Hoeber, New York. p. 88. [6] Ehrlich, P. 1886. Z. Wiss. Mikroskopie 3:150. [7] Gray, P. 1929. Museums J. (London) 28:341. [8] Gray, P. 1963. Handbook of basic microtechnique. Ed. 3. McGraw-Hill, New York. [9] Grenacher, H. 1879. Arch. Mikroskop. Anat. Entwicklungsmech. 16:465. [10] Johansen, D. A. 1939. Stain Technol. 14:125. [11] Johansen, D. A. 1940. Plant microtechnique. McGraw-Hill, New York. [12] Kull, H. 1913-14. Anat. Anz. 45:153. [13] LaCour, L. 1941. Stain Technol. 16:169. [14] Ludford, R. J. 1925. J. Roy. Microscop. Soc., p. 31. [15] Mallory, F. B. 1900-01. J. Exptl. Med. 5:15. [16] Margolena, L. 1932. Stain Technol. 7:25. [17] Masson, P. 1912. Bull. Mem. Soc. Anat. Paris 87:290. [18] Mayer, P. 1892. Mitt. Zool. Stat. Neapel 10:491. [19] Patay, R. 1934. Bull. Histol. Appl. Physiol. Pathol. Tech. Microscop. 11:408. [20] Shumway, W. 1926. Stain Technol. 1:37. [21] Smith, B. G. 1912. J. Morphol. 23:94. [22] Van Wijhe, J. W. 1902. Koninkl. Ned. Akad. Wetenschap. Proc., Sect. Sci., 5:47. [23] Van Kölliker, A. 1892. Festschrift. Univ. Würzburg, Leipzig. [24] Wright, J. H. 1902. J. Med. Res. 7:138.

## 155. HISTOCHEMICAL TESTS

*Abbreviations:* conc. = concentrated; sat. = saturated; sol. = solution.

	Substance	Preparation of Tissue	Preparation of Reagents	Test Method	Result	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
1	Lipids	Fix in 4% formaldehyde containing 1% calcium chloride. Cut frozen sections, if necessary, after embedding in gelatin.	Digest 1 g Sudan black B in 100 ml 60% triethyl phosphate at 100°C for 5 min with constant agitation. Cool, then filter.	Stain section 2-5 min. Wash in 60% triethyl phosphate. Wash in H <sub>2</sub> O and stain carmine-light green. Mount in aqueous medium.	Lipid granules, black; nuclei, red; cytoplasm, green.	4
2	Cholesterol	Cut frozen sections of fresh or formaldehyde-fixed material.		Strand section on slide. Drain well. Cover with 2 drops conc. H <sub>2</sub> SO <sub>4</sub> for 10 sec. Add 2 drops acetic anhydride, wait 10 sec, then wash thoroughly with acetic anhydride. Place cover slip on section.	Cholesterol shows green, or blue-green. Preparation cannot be preserved.	10

*continued*



## 155. HISTOCHEMICAL TESTS

Substance	Preparation of Tissue	Preparation of Reagents	Test Method	Result	Reference
(A)	(B)	(C)	(D)	(E)	(F)
3 Glycogen	Fix in ethanol at 0°C. Cut 10- $\mu$ paraffin sections and mount on slide; deparaffinize in xylene. Rinse in equal parts ethanol and ether. Dip in collodion U.S.P.	Boil 2 g carmine, 1 g potassium carbonate, and 5 g potassium chloride in 60 ml H <sub>2</sub> O for 5 min. Cool, then add 20 ml ammonium hydroxide. For use, dilute 10 ml of this stock sol. with 15 ml ammonium hydroxide and 15 ml ethanol.	Stain collodionized sections in celestin blue B. Wash thoroughly in water. Stain in carmine 15 min. Rinse thoroughly in methanol, dehydrate in acetone, clear in xylene, and mount in balsam.	Nuclei, black; glycogen granules, scarlet.	1
4 Starch	Fix in any dichromate--chromic acid--formaldehyde fixative. Cut paraffin sections.	Saturate hot anilin with acid fuchsin. Shake well, separate, and retain water fraction.	Pour acid fuchsin stain on sections; heat to steaming for 1 min. Rinse in H <sub>2</sub> O and place in 5% aurantia in ethanol until no color comes away. Rinse in 70% ethanol and transfer to 2% tannic acid for 15 min. Transfer directly to 1% methyl green for 10 min. Differentiate in ethanol until starch grains are sharply distinct.	Plastids, proplastids, and mitochondria, red; starch, green. The standard iodine test for starch does not yield permanent preparations.	6
5 Mucin	Cut paraffin sections of material fixed in any mercuric chloride or dichromate fixative.		Stain 10-40 sec in 1% Alcian Blue. Rinse quickly in H <sub>2</sub> O and transfer for 2 hr to 0.5% borax in 80% ethanol. Dehydrate and mount in balsam.	Mucin, bright blue. Stained sections may be counterstained in hematoxylin-eosin if further histological detail is desired.	12
6 Celluloses	Sections of plant tissues, or teased fibers.	Dissolve 2 g iodine and 5 g potassium iodide in a small amount of H <sub>2</sub> O. Dilute to 100 ml. Add 10 ml iodine sol. and 0.25 ml glycerol to 90 ml H <sub>2</sub> O.	Cover specimen with iodine sol. for 15 sec. Blot dry. Add 1 drop of sat. aqueous sol. lithium carbonate. Apply cover slip.	Pure cellulose, blue; impure celluloses, various shades of green, yellow, and brown.	9
7 Lignin	Sections of plant tissues, or teased fibers.		Place in 1% phloroglucinol for 2 min. Blot and add 1 drop of HCl.	Lignin, red.	9
8 Chitin	Sections of tissues.	Dissolve 10 g anilin hydrochloride in 100 ml 1% HCl. Stain sections 5 min.	Transfer to 7.5% potassium dichromate for 1 min. Rinse in H <sub>2</sub> O and place in alkaline tap water until color changes from green to blue.	Chitin, blue.	2
9 DNA	Sections or smears of either animal or plant material.	Boil 1 g magenta ("basic" fuchsin) in 100 ml H <sub>2</sub> O. Add 20 ml N HCl. Cool, filter, and add 5 ml 10% sodium bisulfite. Leave in dark 24 hr.	Hydrolyze material 20 min in N HCl. Stain 2 hr in dark. Bleach cytoplasm 1-2 min in freshly made 100 ml H <sub>2</sub> O, 5 ml 10% sodium bisulfite, 5 ml N HCl. Counterstain in light green if desired. Dehydrate, clear, and mount in balsam.		3
10 RNA	10- $\mu$ paraffin sections of tissues.	Shake 0.5 g methyl green with successive batches of chloroform until all chloroform-soluble color is removed. Add 13 ml of purified dye sol. to 50 ml pH 4.8 acetate buffer and 37 ml 0.5% pyroam G.	Take sections to H <sub>2</sub> O. Blot. Stain 30 min. Blot. Pass to acetone 1 min, and 50:50 acetone-xylene 1 min. Clear in xylene and mount in balsam.	RNA, blue to blue-green; DNA, red.	5

*continued*

## 155. HISTOCHEMICAL TESTS

Substance	Preparation of Tissue	Preparation of Reagents	Test Method	Result	Reference
(A)	(B)	(C)	(D)	(E)	(F)
11 Proteins	10- $\mu$ sections of neutral formaldehyde-fixed material.	Mix 95 ml ethanol with 0.5 ml 0.2 N sodium hydroxide. Add 0.5 g 2,4-dinitro-fluorobenzene.	Take sections to H <sub>2</sub> O. Stain 24 hr. Rinse thoroughly in ethanol, then H <sub>2</sub> O. Bleach in 5% sodium thiosulfate 40 min at 37°C. Rinse in H <sub>2</sub> O. Add 5 ml ice-cold 4N H <sub>2</sub> SO <sub>4</sub> to 100 ml ice-cold 5% sodium nitrate. Soak bleached sections 4-5 min. Rinse in H <sub>2</sub> O. Transfer to 2% H-acid in barbitone-acetate pH 9.2 buffer for 15 min. Rinse in H <sub>2</sub> O, dehydrate, clear, and mount in balsam.	Protein, purple-red.	11
12 Iron	10- $\mu$ , or thicker, sections of tissues fixed in iron-free, neutral formaldehyde.		Take sections to H <sub>2</sub> O. Place in 2% potassium ferrocyanide with equal volume of 0.2 N HCl and stain 20 min. Dehydrate, clear, mount in balsam.	Reactive iron, blue. Non-reactive iron (e.g., in hemoglobin) may be rendered reactive by treating sections for 30 min, before staining, in alkaline H <sub>2</sub> O <sub>2</sub> .	7
13 Hemoglobin	10- $\mu$ , or thicker, sections of tissues fixed in neutral formaldehyde.	Dissolve 1-2 g benzidine in 100 ml methanol with 1.2 ml acetic acid. Add 0.12 g sodium nitroprusside.	Deparaffinize sections in xylene. Remove xylene completely in several changes of methanol. Stain 10 min. Wash in 50 ml methanol, 25 ml ether, 25 ml 3% H <sub>2</sub> O <sub>2</sub> . Dehydrate, clear, and mount in balsam.	Hemoglobin, bright blue.	8
14 Carotene	Immerse plant tissues in 20 ml sat. aqueous sol. potassium hydroxide, 15 ml ethanol, 85 ml H <sub>2</sub> O in dark until all green removed.		Wash pieces thoroughly in H <sub>2</sub> O. Place fragment on slide, blot, and cover with H <sub>2</sub> SO <sub>4</sub> .	Areas of dark blue crystals indicate carotene locations.	13

*Contributor:* Gray, Peter

*References:* [1] Best, F. 1906. Z. Wiss. Mikroskopie 23:319. [2] Bethe, A. 1895. Zool. Jahrb. Abt. Anat. Ontog. Tiere 8:544. [3] Feulgen, R., and H. Rossenbeck. 1924. Z. Physiol. Chem. 135:203. [4] Gomori, G. 1952. Microscopic histochemistry; principles and practice. Univ. Chicago Press, Chicago. [5] Jordan, B. M., and J. R. Baker. 1955. Quart. J. Microscop. Sci. 96:177. [6] Milovidov, P. F. 1928. Arch. Anat. Microscop. Morphol. Exptl. 24:9. [7] Perls, M. 1867. Arch. Pathol. Anat. Physiol. 39:42. [8] Pickworth, F. A. 1934. J. Anat. 69:62. [9] Post, E. E., and J. D. Lauder milk. 1942. Stain Technol. 17:21. [10] Romieu, M. 1925. Compt. Rend. Assoc. Anat. 20:345. [11] Sanger, F. 1945. Biochem. J. 39:507. [12] Steedman, H. F. 1950. Quart. J. Microscop. Sci. 91:477. [13] Steiger, A. 1941. Mikrokosmos 34:121

# *Contrails*

## APPENDIXES

### Appendix I. ESTIMATED NUMBER OF SPECIES: ANIMAL AND PLANT KINGDOMS

Phylum and Class		No. of Species		Phylum and Class		No. of Species	
(A)		(B)		(A)		(B)	
Animal Kingdom <sup>1</sup> [3]							
1	Chordata	44,794		42	Cnidaria	9,600	
2	Mammalia	4,500		43	Porifera	4,200	
3	Aves	8,590		44	Mesozoa	50	
4	Reptilia	5,000		45	Protozoa	30,000	
5	Amphibia	2,000			<b>TOTAL</b>	999,309	
6	Pisces, Chondrichthyes, Agnatha	23,000		Plant Kingdom <sup>2</sup> [1]			
7	Cephalochordata <sup>2</sup>	13		46	Vira	65	
8	Urochordata <sup>2</sup>	1,600		47	Bacteriophyta	1,478	
9	Hemichordata <sup>2</sup>	91		48	Myxophyta	465	
10	Echinodermata	5,700		49	Myxomyceteae	421	
11	Pogonophora	43		50	Acrasieae	21	
12	Chaetognatha	50		51	Plasmodiophoreae	23	
13	Arthropoda	765,257		52	Fungi	30,021	
14	Tardigrada	280		53	Phycomycetes	1,060	
15	Pentastomida	60		54	Ascomycetes	7,297	
16	Pycnogonida	440		55	Basidiomycetes	15,950	
17	Arachnida	30,000		56	Fungi Imperfecti	5,714	
18	Merostomata	4		57	Lichenes	16,128	
19	Crustacea	25,000		58	Phycolichenes	1	
20	Insecta	700,000		59	Ascolichenes	16,119	
21	Symphyla, Chilopoda, Diplopoda, Pauropoda	9,400		60	Basidiolichenes	8	
22	Onychophora	73		61	Algae	16,951	
23	Annelida	7,000		62	Cyanophyta <sup>4</sup>	1,227	
24	Echiuroidea	80		63	Euglenophyta <sup>4</sup>	481	
25	Sipunculoidea	275		64	Pyrrophyta <sup>4</sup>	1,155	
26	Mollusca	100,000		65	Chrysophyta <sup>4</sup>	5,453	
27	Brachiopoda	260		66	Chlorophyta <sup>4</sup>	4,913	
28	Phoronida	15		67	Charophyta <sup>4</sup>	207	
29	Polyzoa	4,000		68	Phaeophyta <sup>4</sup>	963	
30	Entoprocta	60		69	Rhodophyta <sup>4</sup>	2,552	
31	Acanthocephala	300		70	Bryophyta	22,789	
32	Aschelminthes	11,995		71	Musci	14,252	
33	Nematoda	10,000		72	Hepaticae	8,241	
34	Nematomorpha	250		73	Anthocerotae	296	
35	Priapulida	5		74	Pteridophyta	9,640	
36	Echinoderida	100		75	Spermatophyta	188,991	
37	Gastrotricha	140		76	Gymnospermae	696	
38	Rotifera	1,500		77	Angiospermae	188,295	
39	Nemertina	550		78	Monocotyledoneae <sup>5</sup>	41,714	
40	Platyhelminthes	15,000		79	Dicotyledoneae <sup>5</sup>	146,581	
41	Ctenophora	80			<b>TOTAL</b>	286,528	

/1/ Number of hybrids reported: Mammalia, 300; Aves, 1,599; Reptilia, 40; Amphibia, 271; Pisces, 212; Protochordata, 1; Echinodermata, 36; Arthropoda, 289; Mollusca, 12. [2] /a/ Subphylum. /3/ Number of hybrids reported: Schizomycophyta, 2; Eumycophyta, 67; Chlorophyta, 19; Bryophyta, 43; Pteridophyta, 158; Spermatophyta, 15,000 (exclusive of orchids). [2] /4/ Division. /5/ Subclass.

References: [1] Gould, S. W. 1962. Family names of the plant kingdom. International Plant Index, New Haven, Conn. [2] Knobloch, I. W. Unpublished. Michigan State Univ., East Lansing, 1961. [3] Rothschild, Lord. 1961. A classification of living animals. J. Wiley, New York.



## Appendix II. TAXONOMIC CLASSIFICATION: LIVING ANIMALS

<p>Phylum: <b>CHORDATA</b></p> <p>Subphylum: VERTEBRATA</p> <p>Class: <b>Mammalia</b> [6]</p> <p>Subclass: Theria</p> <p>  Infraclass: Eutheria</p> <p>    Order: Artiodactyla</p> <p>      Suborder: Ruminantia</p> <p>      Suborder: Tylopoda</p> <p>      Suborder: Suiformes</p> <p>    Order: Perissodactyla</p> <p>      Suborder: Ceratomorpha</p> <p>      Suborder: Hippomorpha</p> <p>    Order: Sirenia</p> <p>    Order: Hyracoidea</p> <p>    Order: Proboscidea</p> <p>    Order: Tubulidentata</p> <p>    Order: Carnivora</p> <p>      Suborder: Pinnipedia</p> <p>      Suborder: Fissipeda</p> <p>    Order: Cetacea</p> <p>      Suborder: Mysticeti</p> <p>      Suborder: Odontoceti</p> <p>    Order: Rodentia</p> <p>      Suborder: Hystricomorpha</p> <p>      Suborder: Myomorpha</p> <p>      Suborder: Sciuromorpha</p> <p>    Order: Lagomorpha</p> <p>    Order: Pholidota</p> <p>    Order: Edentata</p> <p>    Order: Primates</p> <p>      Suborder: Anthroipoidea</p> <p>      Suborder: Prosimii</p> <p>    Order: Chiroptera</p> <p>      Suborder: Microchiroptera</p> <p>      Suborder: Megachiroptera</p> <p>    Order: Dermoptera</p> <p>    Order: Insectivora</p> <p>  Infraclass: Metatheria</p> <p>    Order: Marsupialia</p> <p>  Subclass: Prototheria</p> <p>    Order: Monotremata</p> <p>Class: <b>Aves</b> [7]</p> <p>  Order: Passeriformes</p> <p>    Suborder: Passeres</p> <p>    Suborder: Menurae</p> <p>    Suborder: Tyranni</p> <p>    Suborder: Eurylaimi</p> <p>  Order: Piciformes</p> <p>    Suborder: Pici</p> <p>    Suborder: Galbulae</p> <p>  Order: Coraciiformes</p> <p>    Suborder: Bucerotidae</p> <p>    Suborder: Coracii</p> <p>    Suborder: Meropes</p> <p>    Suborder: Alcedines</p> <p>  Order: Trogoniformes</p> <p>  Order: Coliiformes</p> <p>  Order: Apodiformes</p> <p>    Suborder: Trochili</p> <p>    Suborder: Apodi</p> <p>  Order: Caprimulgiformes</p> <p>    Suborder: Caprimulgi</p> <p>    Suborder: Steatornithes</p> <p>  Order: Strigiformes</p> <p>  Order: Cuculiformes</p> <p>    Suborder: Cuculi</p>	<p>Suborder: Musophagi</p> <p>Order: Psittaciformes</p> <p>Order: Columbiformes</p> <p>  Suborder: Columbae</p> <p>  Suborder: Pterocletes</p> <p>Order: Charadriiformes</p> <p>  Suborder: Alcae</p> <p>  Suborder: Lari</p> <p>  Suborder: Charadrii</p> <p>Order: Gruiformes</p> <p>  Suborder: Otides</p> <p>  Suborder: Cariamae</p> <p>  Suborder: Eurypygae</p> <p>  Suborder: Rhynocheti</p> <p>  Suborder: Heliornithes</p> <p>  Suborder: Grues</p> <p>  Suborder: Turnices</p> <p>  Suborder: Mesitornithides</p> <p>Order: Galliformes</p> <p>  Suborder: Opisthocomi</p> <p>  Suborder: Galli</p> <p>Order: Falconiformes</p> <p>  Suborder: Falcones</p> <p>  Suborder: Cathartae</p> <p>Order: Anseriformes</p> <p>  Suborder: Anseres</p> <p>  Suborder: Anhimae</p> <p>Order: Ciconiiformes</p> <p>  Suborder: Phoenicopterii</p> <p>  Suborder: Ciconiae</p> <p>  Suborder: Balaenicipites</p> <p>  Suborder: Ardeae</p> <p>Order: Pelecaniformes</p> <p>  Suborder: Fregatae</p> <p>  Suborder: Pelecani</p> <p>  Suborder: Phaethontes</p> <p>Order: Procellariiformes</p> <p>Order: Podicipediformes</p> <p>Order: Gaviiformes</p> <p>Order: Tinamiformes</p> <p>Order: Apterygiformes</p> <p>Order: Casuariiformes</p> <p>Order: Rheiformes</p> <p>Order: Struthioniformes</p> <p>Order: Sphenisciformes</p> <p>Class: <b>Reptilia</b> [4,5]</p> <p>  Order: Serpentes</p> <p>  Order: Sauria</p> <p>  Order: Crocodylia</p> <p>  Order: Chelonia</p> <p>    Suborder: Pleurodira</p> <p>    Suborder: Cryptodira</p> <p>  Order: Rhynchocephalia</p> <p>Class: <b>Amphibia</b> [2,4]</p> <p>  Order: Gymnophiona</p> <p>  Order: Salientia</p> <p>    Suborder: Diplasiocoela</p> <p>    Suborder: Procoela</p> <p>    Suborder: Anomocoela</p> <p>    Suborder: Opisthocoela</p> <p>    Suborder: Amphicoela</p> <p>  Order: Caudata</p> <p>    Suborder: Meantes</p> <p>    Suborder: Proteida</p> <p>    Suborder: Salamandroidea</p> <p>    Suborder: Ambystomoidea</p> <p>    Suborder: Cryptobranchioidea</p>	<p>Class: <b>Pisces</b> [3,4]</p> <p>Subclass: Crossopterygii</p> <p>  Order: Dipnoi</p> <p>  Order: Actinistia</p> <p>Subclass: Neopterygii</p> <p>  Order: Synbranchii</p> <p>    Suborder: Synbranchioidea</p> <p>    Suborder: Alabetoidea</p> <p>  Order: Opisthomi</p> <p>  Order: Pediculati</p> <p>    Suborder: Ceratioidea</p> <p>    Suborder: Antennarioidea</p> <p>    Suborder: Lophioidea</p> <p>  Order: Haplodoci</p> <p>  Order: Xenopterygii</p> <p>  Order: Malacichthyes</p> <p>  Order: Plectognathi</p> <p>    Suborder: Tetradontoidea</p> <p>    Suborder: Balistoidea</p> <p>  Order: Discocephali</p> <p>  Order: Heterosomata</p> <p>  Order: Hypostomides</p> <p>  Order: Thoracostei</p> <p>  Order: Scleroparei</p> <p>    Suborder: Cephalacanthoidea</p> <p>    Suborder: Scorpaenoidea</p> <p>  Order: Percomorphi</p> <p>    Suborder: Polynemoidea</p> <p>    Suborder: Mugiloidea</p> <p>    Suborder: Channoidea</p> <p>    Suborder: Anabantoidea</p> <p>    Suborder: Stromateoidea</p> <p>    Suborder: Ophidioidea</p> <p>    Suborder: Blennioidea</p> <p>    Suborder: Callionymoidea</p> <p>    Suborder: Gobioidae</p> <p>    Suborder: Scombroidea</p> <p>    Suborder: Trichiuroidea</p> <p>    Suborder: Siganoidae</p> <p>    Suborder: Acanthuroidea</p> <p>    Suborder: Percoidea</p> <p>  Order: Zeomorphi</p> <p>    Order: Berycomorphi</p> <p>      Suborder: Xenoberyces</p> <p>      Suborder: Berycoidea</p> <p>    Order: Allotriognathi</p> <p>    Order: Anacanthini</p> <p>    Order: Solenichthyes</p> <p>      Suborder: Aulostomi</p> <p>      Suborder: Lophobranchi</p> <p>    Order: Salmopercae</p> <p>    Order: Microcyprini</p> <p>    Order: Syntognathi</p> <p>      Suborder: Exocoetoidea</p> <p>      Suborder: Scombresocoidea</p> <p>  Order: Heteromi</p> <p>  Order: Apodes</p> <p>  Order: Ostariophysii</p> <p>    Suborder: Siluroidea</p> <p>    Suborder: Cyprinoidea</p> <p>  Order: Lyomeri</p> <p>  Order: Giganturoidea</p> <p>  Order: Iniomii</p> <p>    Suborder: Alepisauroidae</p> <p>    Suborder: Myctophoidea</p> <p>  Order: Haplomi</p>
---	--	---

*continued*

Appendix II. TAXONOMIC CLASSIFICATION: LIVING ANIMALS

<p>Order: Isospondyli  Suborder: Gonorrhynchoidea  Suborder: Mormyroidea  Suborder: Notopteroidea  Suborder: Osteoglossoidea  Suborder: Salmonoidea  Suborder: Stomiatoida  Suborder: Clupeoidea  Order: Ginglymodi  Order: Protospondyli  Subclass: Palaeopterygii  Order: Chondrostei  Order: Cladistia  Class: <b>Chondrichthyes</b> [3,4]  Subclass: Holocephali  Subclass: Elasmobranchii  Order: Batoidel  Order: Selachii  Suborder: Squaloidea  Suborder: Galeoidea  Suborder: Notidanoidea  Class: <b>Agnatha</b> [3,4]  Order: Petromyzones  Order: Myxini  Subphylum: CEPHALOCHORDATA [4]  Subphylum: UROCHORDATA [4]  Class: <b>Larvacea</b>  Order: Copelata  Class: <b>Thaliacea</b>  Order: Salpida  Order: Doliolida  Order: Pyrosomida  Class: <b>Asciacea</b>  Order: Pleurogona  Suborder: Aspiculata  Suborder: Stolidobranchiata  Order: Enterogona  Suborder: Phlebobranchiata  Suborder: Apiousobranchiata  Subphylum: HEMICHORDATA [4]  Class: <b>Planctosphaeroidea</b>  Class: <b>Pterobranchia</b>  Order: Cephalodiscida  Order: Rhabdopleurida  Class: <b>Entero pneusta</b>  Phylum: <b>ECHINODERMATA</b> [4]  Subphylum: ELEUTHEROZOA  Class: <b>Ophiuroidea</b>  Order: Euryalae  Order: Ophiurae  Class: <b>Asteroidea</b>  Order: Forcipulata  Order: Spinulosa  Order: Phanerozona  Class: <b>Echinoidea</b>  Subclass: Euechinoidea  Superorder: Atelostomata  Order: Spatangoida  Order: Holasteroidea  Order: Cassiduloida  Order: Nucleolitoida  Superorder: Gnathostomata  Order: Clypeasteroidea  Suborder: Rotulina  Suborder: Scutellina  Suborder: Laganina  Suborder: Clypeasterina</p>	<p>Order: Holoctypoida  Suborder: Echinoneina  Superorder: Echinacea  Order: Echinoida  Order: Temnopleuroidea  Order: Arbacioida  Order: Phymosomatoida  Order: Hemicidaroida  Superorder: Diadematacea  Order: Echinothurioida  Order: Diadematoidea  Subclass: Perischoechinoidea  Order: Cidaroida  Class: <b>Holothuroidea</b>  Order: Apoda  Order: Molpadonia  Order: Dendrochirota  Order: Elaspoda  Order: Aspidochirota  Subphylum: PELMATOZOA  Class: <b>Crinoidea</b>  Order: Articulata  Phylum: <b>POGONOPHORA</b> [4]  Order: Thecanephria  Order: Athecanephria  Phylum: <b>CHAETOGNATHA</b> [4]  Phylum: <b>ARTHROPODA</b>  Class: <b>Tardigrada</b> [4]  Order: Eutardigrada  Order: Heterotardigrada  Class: <b>Pentastomida</b> [4]  Order: Porocephalida  Order: Cephalobaenida  Class: <b>Pycnogonida</b> [4]  Order: Pycnogonomorpha  Order: Ascorhynchomorpha  Order: Nymphonomorpha  Order: Colossendeomorpha  Class: <b>Arachnida</b> [4]  Order: Acari  Order: Araneae  Order: Opiliones  Order: Solifugae  Order: Ricinulei  Order: Palpigradi  Order: Amblypygi  Order: Uropygi  Order: Pseudoscorpiones  Order: Scorpiones  Class: <b>Merostomata</b> [4]  Order: Xiphosura  Class: <b>Crustacea</b> [4]  Subclass: Malacostraca  Superorder: Eucarida  Order: Decapoda  Suborder: Reptantia  Suborder: Natantia  Order: Euphausiacea  Superorder: Pancarida  Order: Thermosbaenacea  Superorder: Hoplocarida  Order: Stomatopoda  Superorder: Peracarida  Order: Amphipoda  Order: Spelaeogriphacea</p>	<p>Order: Isopoda  Order: Gnathiidea  Order: Tanaidacea  Order: Cumacea  Order: Mysidacea  Superorder: Syncarida  Order: Bathynellacea  Order: Anaspidacea  Superorder: Leptostraca  Order: Nebaliacea  Subclass: Cirripedia  Order: Ascothoracica  Order: Rhizocephala  Order: Acrothoracica  Order: Thoracica  Subclass: Branchiura  Subclass: Mystacocarida  Order: Derocheilocarida  Subclass: Copepoda  Order: Lernaepodoida  Order: Caligoida  Order: Notodelphyoida  Order: Harpacticoida  Order: Cyclopoidea  Order: Monstrilloidea  Order: Calanoida  Subclass: Ostracoda  Order: Platycopoda  Order: Podocopa  Order: Cladocopa  Order: Myodocopa  Subclass: Branchiopoda  Order: Cephalocarida  Order: Cladocera  Order: Conchostraca  Order: Notostraca  Order: Anostraca  Class: <b>Insecta</b> [1]  Subclass: Pterygota  Order: Hymenoptera  Suborder: Apocrita  Suborder: Symphyta  Order: Siphonaptera  Order: Diptera  Suborder: Brachycera  Suborder: Nematocera  Order: Lepidoptera  Suborder: Jugatae  Suborder: Frenatae  Order: Trichoptera  Order: Mecoptera  Order: Strepsiptera  Order: Coleoptera  Suborder: Polyphaga  Suborder: Adephaga  Suborder: Archostemata  Order: Neuroptera  Order: Homoptera  Suborder: Sternorrhyncha  Suborder: Auchenorrhyncha  Order: Hemiptera  Suborder: Gymnocerata  Suborder: Cryptocerata  Order: Thysanoptera  Suborder: Tubulifera  Suborder: Terebrantia  Order: Anoplura  Order: Mallophaga</p>
--	--	---

*continued*

## Appendix II. TAXONOMIC CLASSIFICATION: LIVING ANIMALS

Order: Zoraptera Order: Psocoptera Suborder: Eupsocida Suborder: Troctomorpha Suborder: Trogiomorpha Order: Embioptera Order: Dermaptera Order: Plecoptera Order: Isoptera Order: Orthoptera Suborder: Grylloblattodea Suborder: Blattodea Suborder: Mantodea Suborder: Phasmatodea Suborder: Ensifera Suborder: Caelifera Order: Odonata Suborder: Zygoptera Suborder: Anisoptera Order: Ephemeroptera Subclass: Apterygota Order: Collembola Suborder: Symphleona Suborder: Arthropleona Order: Thysanura Suborder: Entotrophi Suborder: Ectognatha Order: Protura Class: <b>Symphyla</b> [4] Class: <b>Chilopoda</b> [4] Subclass: Anamorpha Order: Scutigermorpha Order: Heterostigmata Suborder: Craterostigmomorpha Suborder: Lithobiomorpha Subclass: Epimorpha Order: Scolopendromorpha Order: Geophilomorpha Class: <b>Diplopoda</b> [4] Subclass: Chilognatha Superorder: Colobognatha Superorder: Helminthomorpha Order: Cambalida Order: Spirostreptida Order: Spirobolida Order: Julida Order: Polydesmida Order: Stemmiulida Order: Nematophora Superorder: Pentazonia Order: Glomeridesmida Order: Glomerida Subclass: Pselaphognatha Order: Polyxenida Class: <b>Pauropoda</b> [4] Class: <b>Onychophora</b> [4]	Order: Xenopneusta Order: Echiuroinea Phylum: <b>SIPUNCULOIDEA</b> [4] Phylum: <b>MOLLUSCA</b> [4] Class: <b>Cephalopoda</b> Order: Dibranchia Suborder: Octopoda Suborder: Vampyromorpha Suborder: Decapoda Order: Tetrabranchia Class: <b>Bivalvia</b> Order: Septibranchia Order: Eulamellibranchia Order: Filibranchia Order: Protobranchia Class: <b>Scaphopoda</b> Class: <b>Gastropoda</b> Subclass: Pulmonata Order: Stylommatophora Order: Basommatophora Subclass: Opisthobranchia Order: Acoela Suborder: Nudibranchia Suborder: Notaspidea Order: Sacoglossa Order: Pteropoda Order: Pleurocoela Subclass: Prosobranchia Order: Stenoglossa Order: Mesogastropoda Order: Archaeogastropoda Class: <b>Monoplacophora</b> Order: Tryblidiacea Class: <b>Aplacophora</b> Order: Chaetodermomorpha Order: Neomeniomorpha Class: <b>Polyplacophora</b> Order: Chitonida Order: Lepidopleurida	Phylum: <b>ASCHELMINTHES</b> [4] Class: <b>Nematoda</b> Subclass: Aphasmidia Order: Enoplida Suborder: Diotophymatina Suborder: Dorylaimina Suborder: Enoplina Order: Chromadorida Subclass: Phasmidia Order: Spirurida Order: Tylenchida Order: Rhabditida Suborder: Ascaridina Suborder: Strongylina Suborder: Rhabditina Class: <b>Nematomorpha</b> Order: Gordioidea Order: Nectonematoidea Class: <b>Priapulida</b> Class: <b>Echinoderida</b> Class: <b>Gastrotricha</b> Order: Chaetonoidea Order: Macrodasyoidea Class: <b>Rotifera</b> Order: Monogononta Suborder: Collothecacea Suborder: Flosculariacea Suborder: Ploima Order: Bdelloidea Order: Seisonidea Phylum: <b>NEMERTINA</b> [4] Class: <b>Enopla</b> Order: Bdeionemertina Order: Hoplonemertina Suborder: Polystylifera Suborder: Monostylifera Class: <b>Anopla</b> Order: Heteronemertina Order: Palaeonemertina
Phylum: <b>ANNELIDA</b> [4] Class: <b>Archannelida</b> Class: <b>Hirudinea</b> Order: Gnathobdellida Order: Rhyrachobdellida Order: Acanthobdellida Class: <b>Oligochaeta</b> Class: <b>Myzostomaria</b> Class: <b>Polychaeta</b>	Phylum: <b>BRACHIOPODA</b> [4] Class: <b>Articulata</b> Suborder: Terebratelloidea Suborder: Terebratuloidea Suborder: Rhynchonelloidea Suborder: Thecideoidea Class: <b>Inarticulata</b> Order: Neotremata Order: Atremata Phylum: <b>PHORONIDA</b> [4] Phylum: <b>POLYZOA</b> [4] Class: <b>Gymnolaemata</b> Order: Ctenostomata Order: Cheilostomata Order: Cyclostomata Class: <b>Phylactolaemata</b>	Phylum: <b>PLATYHELMINTHES</b> [4] Class: <b>Cestoda</b> Subclass: Eucestoda Order: Pseudophyllidea Order: Nippotaeniidea Order: Caryophyllidea Order: Cyclophyllidea Order: Trypanorhyncha Order: Diphyllidea Order: Disculicepitidea Order: Lecanicephala Order: Tetraphyllidea Order: Proteocephala Subclass: Cestodaria Order: Gyrocotylidea Order: Amphilinidea Class: <b>Trematoda</b> Order: Digenea Suborder: Prosostomata Suborder: Gasterostomata Order: Aspidogastrea Order: Monogenea Suborder: Polyopisthocotylea Suborder: Monopisthocotylea Class: <b>Turbellaria</b> Order: Polycladida Suborder: Cotylea
Phylum: <b>ECHIUROIDEA</b> [4] Order: Heteromyota	Phylum: <b>ACANTHOCEPHALA</b> [4] Order: Eoacanthocephala Order: Palaeacanthocephala Order: Archiacanthocephala	

continued



**Appendix II. TAXONOMIC CLASSIFICATION: LIVING ANIMALS**

Suborder: Acotylea Order: Tricladida Suborder: Terricola Suborder: Paludicola Suborder: Maricola Order: Alloeocoela Order: Rhabdocoela Order: Acoela	Phylum: <b>PORIFERA</b> [4] Class: <b>Demospongiae</b> Subclass: Keratosa Subclass: Monaxonida Order: Epipolasisida Order: Haplosclerina Order: Poecilosclerina Order: Halichondrina Order: Hadromerina Subclass: Tetractinellida Order: Choristida Order: Carnosa Order: Myxospongia Class: <b>Hexactinellida</b> Order: Amphidiscophora Order: Hexasterophora Class: <b>Calcarea</b> Order: Syconosa Order: Asconosa	Order: Chonotrichida Order: Suctorida Order: Gymnostomatida Suborder: Cyrtophorina Suborder: Rhabdophorina Class: <b>Cnidosporidia</b> Order: Haplosporidia Order: Actinomyxidia Order: Microsporidia Order: Myxosporidia Class: <b>Sporozoa</b> Subclass: Coccidiomorpha Order: Eucoccidia Suborder: Haemosporidia Suborder: Eimeriidea Suborder: Adeleidea Order: Prococcidia Subclass: Gregarinomorpha Order: Schizogregarina Order: Eugregarina Order: Archigregarina Class: <b>Actinopoda</b> Order: Heliozoa Order: Radiolaria Class: <b>Rhizopoda</b> Order: Foraminifera Order: Testacea Order: Amoebina Order: Rhizomastigina Class: <b>Mastigophora</b> Subclass: Zoomastigina Order: Opalinina Order: Distomatina Order: Metamonadina Order: Protomonadina Subclass: Phytomastigina Order: Chrysomonadina Order: Coccolithophorida Order: Silicoflagellata Order: Ebrideae Order: Dinoflagellata Order: Cryptomonadina Order: Euglenoidina Order: Chloromonadina Order: Xanthomonadina Order: Phytomonadina
Phylum: <b>CTENOPHORA</b> [4] Class: <b>Nuda</b> Order: Beroida Class: <b>Tentaculata</b> Order: Platyctenea Order: Cestida Order: Lobata Order: Cydippida	Phylum: <b>MESOOZOA</b> [4] Order: Orthonectida Order: Dicyemida	
Phylum: <b>CNIDARIA</b> [4] Class: <b>Anthozoa</b> Subclass: Zoantharia Order: Scleractinia Order: Ptychodactiaria Order: Actiniaria Order: Corallimorpharia Order: Zoanthinaria Subclass: Octocorallia Order: Pennatulacea Order: Gorgonacea Order: Alcyonacea Subclass: Ceriantipatharia Order: Ceriantharia Order: Antipatharia Class: <b>Scyphozoa</b> Order: Rhizostomae Order: Semaestomae Order: Coronatae Order: Cubomedusae Order: Stauromedusae Class: <b>Hydrozoa</b> Order: Siphonophora Order: Narcomedusae Order: Trachymedusae Order: Limnomedusae Order: Thecata Order: Athecata	Phylum: <b>PROTOZOA</b> [4] Class: <b>Ciliata</b> Subclass: Spirotricha Order: Hypotrichida Order: Ctenostomatida Order: Entodiniomorpha Order: Tintinnida Order: Oligotrichida Order: Heterotrichida Suborder: Licnophorina Suborder: Heterotrichina Subclass: Holotricha Order: Peritrichida Order: Thigmotrichida Order: Apostomatida Order: Astomatida Order: Hymenostomatida Suborder: Pleuronematina Suborder: Peniculina Suborder: Tetrahymenina Order: Trichostomatida	

*References:* [1] Borror, D. J., and D. M. DeLong. 1954. An introduction to the study of insects. Rinehart, New York. [2] Cochran, D. M. 1961. Living amphibians of the world. Doubleday, Garden City, N. Y. [3] Herald, E. S. 1961. Living fishes of the world. Doubleday, Garden City, N. Y. [4] Rothschild, Lord. 1961. A classification of living animals. J. Wiley, New York. [5] Schmidt, K. P., and R. F. Inger. 1957. Living reptiles of the world. Hanover House, Garden City, N. Y. [6] Simpson, G. G. 1945. Bull. Am. Museum Nat. Hist. 85. [7] Wetmore, A. 1960. Smithsonian Inst. Misc. Collections 139(11).



Appendix III. TAXONOMIC CLASSIFICATION: LIVING PLANTS

Part I. NONVASCULAR PLANTS

Phylum: <b>VIRA</b> [2]	Order: Ustilaginales	Order: Prorocentrales
Order: Virales	Order: Auriculariales	Order: Dinophysalidales
Order: Rickettsiales	Order: Dacrymycetales	Order: Desmocapsales
Phylum: <b>BACTERIOPHYTA</b> [1,2]	Order: Tremellales	Class: <b>Dinophyceae</b>
Class: <b>Schizomycetes</b>	Order: Tulasnellales	Subclass: Dinoflagellatae
Order: Pseudomonadales	Order: Polyporales	Order: Gymnodiniales
Order: Chlamydobacteriales	Order: Agaricales	Order: Blastodiniales
Order: Hyphomicrobiales	Order: Protogastrales	Order: Peridiniales
Order: Eubacteriales	Order: Hymenogastrales	Subclass: Phytodiniiformes
Order: Actinomycetales	Order: Phallales	Order: Rhizodiniales
Order: Caryophanales	Order: Sclerodermatales	Order: Dinocapsales
Order: Beggiatoales	Order: Nidulariales	Order: Dinococcales
Order: Myxobacterales	Order: Sphaerobolales	Order: Lycoperdals
Order: Spirochaetales	Class: <b>Fungi Imperfecti</b>	Division: <b>CHRYSOPHYTA</b>
Order: Mycoplasmatales	Order: Sphaeropsidales	Class: <b>Heterokontae</b>
<i>Incertis sedis</i>	Order: Melanconiales	Order: Heterochloridales
Phylum: <b>MYXOPHYTA</b> [2]	Order: Moniliales	Order: Rhizochloridales
Class: <b>Myxomyceteae</b>	<i>Incertis sedis</i>	Order: Heterocapsales
Order: Exosporales	Mycelia sterilia	Order: Heterotrichales
Order: Stemonitales	Phylum: <b>LICHENES</b> [2,3]	Order: Heterosiphonales
Order: Liceales	Class: <b>Phycolichenes</b>	Class: <b>Chrysophyceae</b>
Order: Trichiales	Order: Geosiphonales	Order: Chrysoomonadales
Order: Physarales	Class: <b>Ascolichenes</b>	Suborder: Chromulininales
Order: Hydromyxales	Order: Verrucariales	Suborder: Isochrysidinales
Class: <b>Acrasieae</b>	Order: Pyrenulales	Suborder: Ochromonadinales
Order: Acrasiales	Order: Pyrenidiales	Order: Silicoflagellatae
Order: Labyrinthulales	Order: Dermatinales	Suborder: Siphonotestinales
Class: <b>Plasmodiophoreae</b>	Order: Caliciales	Suborder: Stereotestinales
Order: Plasmodiophorales	Order: Graphidiales	Order: Rhizochrysidales
Phylum: <b>FUNGI</b> [2]	Order: Roccellales	Order: Chrysocapsales
Class: <b>Phycomycetes</b>	Order: Thelotrematales	Order: Chrysochaetales
Order: Chytridiales	Order: Cyanophilales	Class: <b>Bacillariophyceae</b>
Order: Hyphochytriales	Order: Lecidiales	Subclass: Centricae
Order: Blastocladales	Order: Lecanorales	Order: Disciales
Order: Monoblepharidales	Order: Caloplacales	Order: Soleniales
Order: Saprolegniales	Class: <b>Basidiolichenes</b>	Order: Biddulphiales
Order: Lagenidiales	Order: Corales	Subclass: Pennatae
Order: Peronosporales	Class: <b>Lichenes Imperfecti</b>	Order: Araphidiales
Order: Mucorales	Phylum: <b>ALGAE</b> [3]	Order: Raphidioidales
Order: Entomophthorales	Division: <b>CYANOPHYTA</b>	Order: Monoraphidales
Order: Zoopagales	Class: <b>Cyanophyceae</b>	Order: Biraphidales
Order: Eccrinales	Order: Chroococcales	Division: <b>CHLOROPHYTA</b>
Class: <b>Ascomycetes</b>	Order: Pleurocapsales	Class: <b>Chlorophyceae</b>
Order: Endomycetales	Order: Chamaesiphonales	Order: Chlorochytridiales
Order: Taphrinales	Order: Hormogonales	Order: Volvocales
Order: Plectascales	Suborder: Nostocinales	Suborder: Chlamydomonadinales
Order: Myriangiales	Suborder: Stigonematinales	Suborder: Tetrasporinales
Order: Erysiphales	Division: <b>EUGLENOPHYTA</b>	Suborder: Chlorodendrinales
Order: Dothideales	Class: <b>Euglenophyceae</b>	Order: Chlorococcales
Order: Pseudosphaeriales	Order: Euglenales	Order: Ulotrichales
Order: Hemisphaeriales	Suborder: Eugleninales	Suborder: Ulotrichinales
Order: Sphaeriales	Suborder: Colaciinales	Suborder: Ulvinales
Order: Hypocreales	Division: <b>PYRROPHYTA</b>	Suborder: Schizogoniinales
Order: Clavicipitales	Class: <b>Cryptophyceae</b>	Suborder: Sphaeropleinales
Order: Pezizales	Order: Monomastigales	Order: Chaetophorales
Order: Helotiales	Order: Cryptomonadales	Order: Cladophorales
Order: Hysteriales	Order: Phaeocapsales	Order: Oedogoniales
Order: Tuberales	Order: Cryptococcales	Order: Conjugatae
Order: Laboulbeniales	Class: <b>Chloromonadophyceae</b>	Suborder: Euconjugatae
Class: <b>Basidiomycetes</b>	Order: Chloromonadales	Suborder: Desmidiinales
Order: Uredinales	Class: <b>Desmokontae</b>	Order: Siphonales
	Order: Desmomonadales	Division: <b>CHAROPHYTA</b>
		Class: <b>Charophyceae</b>
		Order: Charales

continued

## Appendix III. TAXONOMIC CLASSIFICATION: LIVING PLANTS

### Part I. NONVASCULAR PLANTS

Division: PHAEOPHYTA Class: <b>Isogeneratae</b> Order: Ectocarpales Order: Sphacelariales Order: Cutleriales Order: Tilotpteridales Order: Dictyotales Class: <b>Heterogeneratae</b> Subclass: Haplostichidae Order: Chordariales Order: Sporochnales Order: Desmarestiales Subclass: Polystichidae Order: Dictyosiphonales Order: Laminariales Class: <b>Cyclosporeae</b> Order: Fucales Division: RHODOPHYTA Class: <b>Bangiophyceae</b> Order: Porphyridiales	Order: Goniotrichales Order: Bangiales Order: Compsopogonales Order: Rhodochaetales Class: <b>Florideae</b> Order: Nemalionales Order: Gelidiales Order: Cryptonemiales Order: Gigartinales Order: Rhodymeniales Order: Ceramiales <hr/> Phylum: <b>BRYOPHYTA</b> [2] Class: <b>Musci</b> Order: Sphagnales Order: Andreaeales Order: Archidiales Order: Dicranales Order: Fissidentales Order: Pottiales	Order: Grimmiales Order: Funariales Order: Schistostegales Order: Tetrarhiales Order: Eubryales Order: Isobryales Order: Hookeriales Order: Hypnobryales Order: Buxbaumiales Order: Polytrichales Order: Dawsoniales Class: <b>Hepaticae</b> Order: Jungermaniales Order: Calobryales Order: Acrogynales Order: Sphaerocarpaceae Order: Marchantiales Class: <b>Anthocerotae</b> Order: Anthocerotales
---	---	--

*References:* [1] Breed, R. S., E. G. D. Murray, and N. R. Smith, ed. 1957. *Bergey's Manual of determinative bacteriology*. Ed. 7. Williams and Wilkins, Baltimore. [2] Gould, S. W. 1962. Family names of the plant kingdom. *International Plant Index*, New Haven, Conn. [3] Melchior, H., and E. Werdermann, ed. 1954. *Engler's Syllabus der Pflanzenfamilien*. Bd. 1. Gebrüder Borntraeger, Berlin-Nikolassee.

### Part II. VASCULAR PLANTS

Phylum: <b>PTERIDOPHYTA</b> [1,2] Class: <b>Psilopsida</b> Order: Psilotales Family: Psilotaceae Class: <b>Lycopsidea</b> Order: Lycopodiales Family: Lycopodiaceae Order: Selaginellales Family: Selaginellaceae Order: Isoetales Family: Isoetaceae Class: <b>Sphenopsida</b> Order: Equisetales Family: Equisetaceae Class: <b>Pteropsida</b> Subclass: Eusporangiatae Order: Ophioglossales Family: Ophioglossaceae Order: Marattiales Family: Marattiaceae Subclass: Osmundidae Order: Osmundales Family: Osmundaceae Subclass: Leptosporangiatae Order: Filicales Family: Aspidiaceae Family: Aspleniaceae Family: Blechnaceae Family: Cyatheaceae Family: Davalliaceae Family: Dicksoniaceae	Family: Gleicheniaceae Family: Hymenophyllaceae Family: Hymenophyllopsidaceae Family: Loxsomaceae Family: Matoniaceae Family: Parkeriaceae Family: Plagiogyriaceae Family: Polypodiaceae Family: Pteridaceae Family: Schizaeaceae Family: Vittariaceae Order: Marsileales Family: Marsileaceae Order: Salviniiales Family: Salviniaceae <hr/> Phylum: <b>SPERMATOPHYTA</b> [1] Class: <b>Gymnospermae</b> Order: Pinales Family: Araucariaceae Family: Cephalotaxaceae Family: Cupressaceae Family: Pinaceae Family: Podocarpaceae Family: Taxodiaceae Order: Taxales Family: Taxaceae Order: Ginkgoales Family: Ginkgoaceae Order: Gnetales Family: Ephedraceae Family: Gnetaceae	Family: Welwitschiaceae Order: Cycadales Family: Cycadaceae Class: <b>Angiospermae</b> Subclass: Monocotyledoneae Order: Pandanales Family: Pandanaceae Family: Sparganiaceae Family: Typhaceae Order: Najadales Family: Alismataceae Family: Aponogetonaceae Family: Butomaceae Family: Hydrocharitaceae Family: Liliaceae Family: Najadaceae Family: Potamogetonaceae Family: Scheuchzeriaceae Order: Triuridales Family: Triuraceae Order: Graminales Family: Gramineae Order: Cyperales Family: Cyperaceae Order: Palmales Family: Palmae Order: Cyclanthales Family: Cyclanthaceae Order: Arales Family: Araceae Family: Lemnaceae
---	--	---

*continued*

Appendix III. TAXONOMIC CLASSIFICATION: LIVING PLANTS

Part II. VASCULAR PLANTS

Order: Farinosae	Order: Podostemales	Order: Sarraceniales
Family: Bromeliaceae	Family: Podostemaceae	Family: Droseraceae
Family: Centrolepidaceae	Order: Proteales	Family: Nepenthaceae
Family: Commelinaceae	Family: Proteaceae	Family: Sarraceniaceae
Family: Cyanastraceae	Order: Santalales	Order: Rosales
Family: Eriocaulaceae	Family: Grubbiaceae	Family: Brunelliaceae
Family: Flagellariaceae	Family: Loranthaceae	Family: Bruniaceae
Family: Mayacaceae	Family: Misodendraceae	Family: Byblidaceae
Family: Philydraceae	Family: Octoknemaceae	Family: Cephalotaceae
Family: Pontederiaceae	Family: Olacaceae	Family: Connaraceae
Family: Rapateaceae	Family: Opiliaceae	Family: Crassulaceae
Family: Restionaceae	Family: Santalaceae	Family: Crossosomataceae
Family: Thurniaceae	Order: Aristolochiales	Family: Cunoniaceae
Family: Xyridaceae	Family: Aristolochiaceae	Family: Eucommiaceae
Order: Liliales	Family: Hydnoraceae	Family: Hamamelidaceae
Family: Amaryllidaceae	Family: Rafflesiaceae	Family: Leguminosae
Family: Dioscoreaceae	Order: Balanophorales	Family: Myrothamnaceae
Family: Haemodoraceae	Family: Balanophoraceae	Family: Pittosporaceae
Family: Iridaceae	Order: Polygonales	Family: Platanaceae
Family: Juncaceae	Family: Polygonaceae	Family: Rosaceae
Family: Liliaceae	Order: Caryophyllales	Family: Saxifragaceae
Family: Stemonaceae	Family: Achatocarpaceae	Order: Pandales
Family: Taccaceae	Family: Aizoaceae	Family: Pandaceae
Family: Velloziaceae	Family: Amaranthaceae	Order: Geraniales
Order: Zingiberales	Family: Basellaceae	Family: Burseraceae
Family: Cannaceae	Family: Caryophyllaceae	Family: Callitrichaceae
Family: Marantaceae	Family: Chenopodiaceae	Family: Cneoraceae
Family: Musaceae	Family: Gyrostemonaceae	Family: Daphniphyllaceae
Family: Zingiberaceae	Family: Nyctaginaceae	Family: Dichapetalaceae
Order: Orchidales	Family: Phytolaccaceae	Family: Erythroxylaceae
Family: Burmanniaceae	Family: Portulacaceae	Family: Euphorbiaceae
Family: Orchidaceae	Order: Ranales	Family: Geraniaceae
Subclass: Dicotyledoneae	Family: Annonaceae	Family: Linaceae
Order: Casuarinales	Family: Berberidaceae	Family: Malpighiaceae
Family: Casuarinaceae	Family: Calycanthaceae	Family: Meliaceae
Order: Piperales	Family: Ceratophyllaceae	Family: Oxalidaceae
Family: Chloranthaceae	Family: Cercidiphyllaceae	Family: Polygalaceae
Family: Piperaceae	Family: Degeneriaceae	Family: Rutaceae
Family: Saururaceae	Family: Eupomatiaceae	Family: Simaroubaceae
Order: Hydrostachyales	Family: Eupteleaceae	Family: Tremandraceae
Family: Hydrostachyaceae	Family: Gomortegaceae	Family: Trigoniaceae
Order: Salicales	Family: Hernandiaceae	Family: Tropaeolaceae
Family: Salicaceae	Family: Himantandraceae	Family: Vochysiaceae
Order: Garryales	Family: Illiciaceae	Family: Zygophyllaceae
Family: Garryaceae	Family: Lactoridaceae	Order: Sapindales
Order: Myricales	Family: Lardizabalaceae	Family: Aceraceae
Family: Myricaceae	Family: Lauraceae	Family: Aextoxicaceae
Order: Balanopsidales	Family: Magnoliaceae	Family: Anacardiaceae
Family: Balanopaceae	Family: Menispermaceae	Family: Aquifoliaceae
Order: Leitneriales	Family: Monimiaceae	Family: Balsaminaceae
Family: Leitneriaceae	Family: Myristicaceae	Family: Buxaceae
Order: Juglandales	Family: Nymphaeaceae	Family: Celastraceae
Family: Juglandaceae	Family: Ranunculaceae	Family: Coriariaceae
Order: Julianales	Family: Schisandraceae	Family: Corynocarpaceae
Family: Julianiaceae	Family: Tetracentraceae	Family: Cyrillaceae
Order: Batidales	Family: Trochodendraceae	Family: Didiereaceae
Family: Bataceae	Family: Winteraceae	Family: Empetraceae
Order: Fagales	Order: Rhoeadales	Family: Hippocastanaceae
Family: Betulaceae	Family: Bretschneideraceae	Family: Icacinaceae
Family: Fagaceae	Family: Capparaceae	Family: Limnanthaceae
Order: Urticales	Family: Cruciferae	Family: Melianthaceae
Family: Moraceae	Family: Fumariaceae	Family: Pentaphragaceae
Family: Rhoipteleaceae	Family: Moringaceae	Family: Sabiaceae
Family: Ulmaceae	Family: Papaveraceae	Family: Salvadoraceae
Family: Urticaceae	Family: Resedaceae	Family: Sapindaceae
	Family: Tovariaceae	

*continued*



## Appendix III. TAXONOMIC CLASSIFICATION: LIVING PLANTS

### Part II. VASCULAR PLANTS

Family: Stackhouseiaceae Family: Staphyleaceae Order: Rhamnales Family: Rhamnaceae Family: Vitaceae Order: Malvales Family: Bombacaceae Family: Elaeocarpaceae Family: Malvaceae Family: Sarcocaulaceae Family: Scytopetalaceae Family: Sterculiaceae Family: Tiliaceae Order: Parietales Family: Achariaceae Family: Actinidiaceae Family: Ancistrocladaceae Family: Begoniaceae Family: Bixaceae Family: Canellaceae Family: Caricaceae Family: Caryocaraceae Family: Cistaceae Family: Cochlospermaceae Family: Datisceae Family: Dilleniaceae Family: Dipterocarpaceae Family: Elatinaceae Family: Eucryphiaceae Family: Flacourtiaceae Family: Frankeniaceae Family: Guttiferae Family: Hypericaceae Family: Loasaceae Family: Malesherbiaceae Family: Marcgraviaceae Family: Medusagynaceae Family: Ochnaceae Family: Passifloraceae Family: Quinaceae Family: Stachyuraceae Family: Strasburgeriaceae Family: Tamaricaceae Family: Theaceae Family: Turneraceae Family: Violaceae Order: Opuntiales Family: Cactaceae	Order: Myrtales Family: Alangiaceae Family: Combretaceae Family: Crypteroniaceae Family: Cynomoriaceae Family: Elaeagnaceae Family: Geissolomataceae Family: Haloragaceae Family: Heteropyxidaceae Family: Hippuridaceae Family: Lecythidaceae Family: Lythraceae Family: Melastomataceae Family: Myrtaceae Family: Nyssaceae Family: Oliaceae Family: Onagraceae Family: Penaeaceae Family: Punicaceae Family: Rhizophoraceae Family: Sonneratiaceae Family: Theligonaceae Family: Thymelaeaceae Family: Trapaceae Order: Umbellales Family: Araliaceae Family: Cornaceae Family: Umbelliferae Order: Diapensiales Family: Diapensiaceae Order: Ericales Family: Clethraceae Family: Epacridaceae Family: Ericaceae Family: Pyrolaceae Order: Primulales Family: Myrsinaceae Family: Primulaceae Family: Theophrastaceae Order: Plumbaginales Family: Plumbaginaceae Order: Ebenales Family: Dielidanthraceae Family: Ebenaceae Family: Hoplostigmataceae Family: Lissocarpaceae Family: Sapotaceae Family: Styracaceae	Family: Symplocaceae Order: Gentianales Family: Apocynaceae Family: Asclepiadaceae Family: Desfontainiaceae Family: Gentianaceae Family: Loganiaceae Family: Oleaceae Order: Tubiflorae Family: Acanthaceae Family: Bignoniaceae Family: Boraginaceae Family: Columelliaceae Family: Convolvulaceae Family: Fouquieriaceae Family: Gesneriaceae Family: Globulariaceae Family: Hydrophyllaceae Family: Labiatae Family: Lennoaceae Family: Lentibulariaceae Family: Martyniaceae Family: Myoporaceae Family: Nolanaceae Family: Orobanchaceae Family: Pedaliaceae Family: Phrymaceae Family: Polemoniaceae Family: Scrophulariaceae Family: Solanaceae Family: Verbenaceae Order: Plantaginales Family: Plantaginaceae Order: Rubiales Family: Adoxaceae Family: Caprifoliaceae Family: Dipsacaceae Family: Rubiaceae Family: Valerianaceae Order: Cucurbitales Family: Cucurbitaceae Order: Campanulatae Family: Brunoniaceae Family: Calyceraceae Family: Campanulaceae Family: Compositae Family: Goodeniaceae Family: Stylidiaceae
---	---	---

*References:* [1] Gould, S. W. 1962. Family names of the plant kingdom. International Plant Index, New Haven, Conn. [2] Melchior, H., and E. Werdermann, ed. 1954. Engler's Syllabus der Pflanzenfamilien. Bd. 1. Gebrüder Borntraeger, Berlin-Nikolassee.



# Contrails

## Appendix IV. GEOLOGIC DISTRIBUTION: ANIMALS AND PLANTS

Era (Duration, yr)	Period (Duration, yr) [Years Ago]	Epoch	Advances in Life	Dominant Life
(A)	(B)	(C)	(D)	(E)
1 Cenozoic (60 million)	Quaternary (2 million) [2 million]	Recent	Rise of civilization	Man and herbaceous plants
		Pleistocene	Periodic glaciation; extinction of great mammals and many trees; rise of modern herbs; elevation of continents	Mammals, birds, and flowering plants
	Tertiary (58 million) [60 million]	Pliocene	Continued cooling of climate; elevation of Andes; increasing restriction of plant distribution and forests; appearance of man	
		Miocene	Greatly changing climate, becoming cool and semiarid; elevation of Alps; restriction of distribution of plants; beginning of forest reduction; culmination of many mammals	
		Oligocene	Warm, humid climate; elevation of Pyrenees; culmination of Eocene floras; worldwide distribution of tropical forests; disappearance of primitive mammals; rise of higher mammals and birds	
		Eocene	Cool climate, semiarid, then warm and humid; modernization of flowering plants; development of extensive forests, widespread in polar regions; appearance of modern birds and marine mammals	
7 Mesozoic (125 million)	Cretaceous (65 million) [125 million]	Upper Cretaceous	Fluctuating climate; angiosperms dominant in floras, dicotyledons, and monocotyledons of numerous existing genera well-developed; disappearance of Bennettitales; rise of primitive mammals	Reptiles and higher gymnosperms
		Middle Cretaceous	Fluctuating climate; rapid development of angiosperms, appearance of many existing genera; specialization and extinction of great reptiles	
		Lower Cretaceous	Very warm climate; rise of angiosperms (?); conifers and cycads still dominant; earliest known pines (Pinaceae)	
	Jurassic (32 million) [157 million]	Jurassic	Warm climate; beginning of Sierra Nevada mountains; first definitely known angiosperms; conifers and cycads dominant; Ginkgoales and conifers worldwide; disappearance of cordaites; primitive birds and flying reptiles; rise of higher insects	
	Triassic (28 million) [185 million]	Triassic	Warm, semiarid climate; nonluxuriant floras; increase in gymnosperms, spread of conifers, rise of cycads and Bennettitales; disappearance of seed ferns; diversification of modern fern families well under way; first mammals; rise of giant reptiles (dinosaurs)	
12 Paleozoic (368 million)	Carboniferous	Permian (38 million) [223 million]	Dry climate; periodic glaciation, severe in southern hemisphere; elevation of Appalachians; dwindling of ancient groups, extinction of many; rise of ferns and conifers and of land vertebrates; expansion of reptiles; Gondwana flora of southern hemisphere distinct from those in northern hemisphere	Amphibians, lycopods, and seed ferns
		Pennsylvanian (48 million) [271 million]	Lycopods, seed ferns, and horsetails dominant; sphenophylls, Coniferales and calamites present; extensive coal formation	
		Mississippian (38 million) [309 million]	Lycopods, horsetails, and seed ferns dominant; cordaites, sphenophylls, and calamites present; rise of primitive reptiles and insects	
	Devonian (45 million) [354 million]	Devonian	Early land floras: Psilophytales ( <i>Rhynia</i> , <i>Horneophyton</i> ); lycopods; primitive horsetails, including sphenophylls; primitive gymnosperms (earliest cordaites and seed ferns, their seeds not yet known); first forests; rise of amphibians and fishes	Fishes and early land plants
	Silurian (27 million) [381 million]	Silurian	Algae dominant (e.g., <i>Nematophyton hicksi</i> ); rise of lungfishes and scorpions (first air-breathing animals)	Higher invertebrates and algae
	Ordovician (67 million) [448 million]	Ordovician	Rise of corals, armored fishes (and land plants ?); marine algae dominant; first known freshwater fishes	
	Cambrian (105 million) [553 million]	Cambrian	Warm climate, uniform over earth; first abundance of fossils; many groups of marine invertebrates; dominant trilobites; marine plants, few algae determinable; evidence from microfossils may indicate land plants	

continued

## Appendix IV. GEOLOGIC DISTRIBUTION: ANIMALS AND PLANTS

Era (Duration, yr)	Period (Duration, yr) [Years Ago]	Epoch (Duration, yr) [Years Ago]	Advances in Life	Dominant Life	
(A)	(B)	(C)	(D)	(E)	
19	Proterozoic (900 million)	Precambrian [1.5 billion]	Precambrian	Rocks chiefly sedimentary, and of enormous thickness; glaciation; first fossils: worms, crustaceans, brachiopods; evidence of algal and bacterial life	Primitive marine invertebrates (fossils rare)
20	Archeozoic (550+ million)	[3.5-5.0 billion]		Igneous rocks: lavas and metamorphosed rocks; few sedimentary; no direct evidence of life; graphites of possible organic origin in sedimentary rocks of Grenville series	Unicellular life? (fossils unknown)

*Contributors:* (a) Wetmore, Ralph H., (b) ZoBell, Claude E., (c) Eames, A. J., and Banks, Harlan P.

*References:* [1] Andrews, H. N., Jr. 1961. Studies in paleobotany. J. Wiley, New York. [2] Augusta, J., and Z. Burian. 1957. Prehistoric animals. Spring Books, London. [3] Clark, W. E. le G. 1957. History of primates. Univ. Chicago Press, Chicago. [4] Dunbar, C. O. 1960. Historical geology. J. Wiley, New York. [5] Dunbar, C. O., and J. Rodgers. 1957. Principles of stratigraphy. J. Wiley, New York. [6] Holmes, A. 1959. Trans. Edinburgh Geol. Soc. 17:183. [7] Jaeger, H. 1962. Palaeontol. Z. 36:7. [8] Knopf, A. 1957. Sci. Monthly 85:225. [9] Kulp, J. L. 1961. Science 133:1105. [10] Kummel, B. 1961. The history of the earth. W. H. Freeman, San Francisco. [11] Life Magazine Editorial Staff, and L. Barnett. 1955. The world we live in. Time, New York. [12] Marble, J. P., et al. 1952. Trans. Am. Geophys. Union 33:149. [13] Schuchert, C. 1955. Atlas of paleogeographic maps of North America. J. Wiley, New York. [14] Seward, A. C. 1931. Plant life through the ages. Cambridge Univ. Press, London. [15] Simpson, G. G. 1953. An introduction to paleontology. Yale Univ. Press, New Haven. [16] Weller, J. M. 1960. Stratigraphic principles and practice. Harper, New York. [17] Wilmarth, M. G. 1925. U.S. Geol. Surv. Bull. 769. [18] Zeuner, F. E. 1952. Dating the past. Methuen, London.

## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part I. CONVERSION FORMULAS

<p>mEq/L ↔ mg/100 ml</p> $\text{mEq/L} = \frac{\text{mg/100 ml} \times \text{valence} \times 10}{\text{atomic weight}}$ $\text{mg/100 ml} = \frac{\text{mEq/L} \times \text{atomic weight}}{\text{valence} \times 10}$	<p>mM/L ↔ mg/100 g</p> $\text{mM/L} = \frac{\text{mg/100 g} \times 10}{\text{atomic weight}}$ $\text{mg/100 g} = \frac{\text{mM/L} \times \text{atomic weight}}{10}$
<p>ml (milliliters) ↔ g (grams)</p> $\text{ml} = \frac{\text{g}}{\text{specific gravity}}$ $\text{g} = \text{ml} \times \text{specific gravity}$	<p>Wet basis ↔ dry basis</p> $\text{Wet basis} \rightarrow \text{dry basis} = \frac{100 \times a}{b}$ $\text{Dry basis} \rightarrow \text{wet basis} = \frac{a \times b}{100}$ <p>a = % of material determined (wet or dry)                      b = % of dry sample (100 - c = b)                      c = % moisture in sample</p>
<p>Temperature: °C ↔ °F</p> $^{\circ}\text{C} = (^{\circ}\text{F} - 32) \times 5/9$ $^{\circ}\text{F} = (^{\circ}\text{C} \times 9/5) + 32$	

*continued*

## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Multiply the value for the unit of measurement in column A by the factor in column B to convert to the unit of measurement in column C.

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Abamperes		10,0000 2.99796 x 10 <sup>10</sup>		amperes statamperes	BTU		777.9 3.929 x 10 <sup>-4</sup> 1,055		ft-lb horsepower-hr joules
Abcoulombs		10,0000 2.99796 x 10 <sup>10</sup>		coulombs (abs.) statcoulombs	Bushels (U.S., dry)		0.304785 35,239 1.2444 2,150.42 0.035239 0.35238329 35.238329 4		barrels cu cm cu ft cu inches cu m hectoliters liters pecks
Abcoulombs/kg		30,577		statcoulombs/dyne	Bushels (U.S., dry)/acre		0.870754		Hectoliters/hectare
Abcoulombs/lb		6.7411 x 10 <sup>4</sup> 166.2		statcoulombs/dyne	Calorie (15°C) sec		6.3854 x 10 <sup>33</sup>		quanta
Abfarads		1.0000 x 10 <sup>9</sup> 1 x 10 <sup>15</sup> 8.98776 x 10 <sup>20</sup>		farads (abs.) microfarads staffarads	Calorie (15°C)		1.0535 x 10 <sup>10</sup>		quanta
Abhenries		1.0000 x 10 <sup>-9</sup> 1 x 10 <sup>-6</sup> 1.11263 x 10 <sup>-21</sup>		henries (abs.) millihenries stathenries	Calories (15°C)/sec/No <sup>1</sup>		0.011625		joules/abcoulomb
Abmhos/cu cm		1,000 1.00052 x 10 <sup>9</sup> 166.2		megmhos/cu cm mhos (Int.)/cu cm mhos/million ft	Calories (15°C)/amp-hr		41.850		joules/abcoulomb
Abohms		1 x 10 <sup>-15</sup> 0.001 1.0000 x 10 <sup>-9</sup> 1.11263 x 10 <sup>-21</sup>		megohms microhms ohms (abs.) statohms	Calories (15°C)/coulomb		41.850		joules/abcoulomb
Abvolts		0.010000 3.33560 x 10 <sup>-1</sup> 1.0000 x 10 <sup>-8</sup>		microvolts statvolts volts (abs.)	Calories, gram		3.968 x 10 <sup>-3</sup> 1.5591 x 10 <sup>-6</sup>		BTU horsepower-hr
Abvolts/°F		0.018000		microvolts/°C	Calories, gram (mean)		0.001469 99.334 3.0874 0.42685 4.1311 x 10 <sup>-2</sup> 0.0011628		cu ft-atm ft-poundals ft-lb kg-m L-atm watt-hr
Abvolts/cm		1.0000 x 10 <sup>-8</sup>		volts (abs.)/cm	Calories, gram (mean)/g		1.8		BTU (mean)/lb
Abvolts/inch		3.9370 x 10 <sup>-9</sup>		volts (abs.)/cm	Calories, gram (mean)/g/°C		4.186		BTU (60°F)/lb/°F joules/g/°C
Acres (U.S.)		40.46873 0.4046873 43,560 4,046.873 0.0015625 160 4,840		ares hectares sq ft sq m sq mi sq rods sq yd	Calories, gram (15°C)		4.185		joules (abs.)
Ampere-hours (absolute)		3,600.0		coulombs (abs.)	Calories, gram (15°C)/°C		0.0022046 4.185		BTU (60°F)/°F joules/°C
Amperes (absolute)		0.1 1.00007 1.0363 x 10 <sup>-5</sup> 2.99796 x 10 <sup>9</sup>		abamperes amperes (Int.) faradays/sec statamperes	Calories, kilogram (mean)		3.9685 1,000 4.186 x 10 <sup>10</sup> 3,087.4 4.2686 x 10 <sup>7</sup> 0.0015593 4,186 426.85 0.0011628		BTU (mean) gram calories (mean) ergs ft-lb g-cm horsepower-hr joules kg-m kw-hr
Angstrom units		3.937 x 10 <sup>-9</sup> 1 x 10 <sup>-10</sup> 100 1 x 10 <sup>-4</sup> 0.1		inches meters micromicrons microns millimicrons	Calories, kilogram (mean)/min		51.457 0.093558 69.769		ft-lb/sec horsepower watts
Atmospheres		1.0133 1.01325 x 10 <sup>6</sup> 1,033.3 10,333 2,116.32 14.696 76 760 760,000 29.921 33.899		bars dynes/sq cm g/sq cm kg/sq m lb/sq ft lb/sq inch cm Hg at 0°C mm Hg microns Hg inches Hg at 32°F ft water at 39.1°F	Calories, kilogram (mean)/sec		4.186		kilowatts
Barrels (U.S., dry)		3.281 7,056 0.11562 105.0		bushels cu inches cu m quarts (dry)	Candlepower (spherical)		12.566		lumens
Barrels (U.S., liquid)		0.11924 31.5		cu m gallons	Candles (International)		1.0000		lumens (Int.)/steradian
BTU		252 25,030		gram calories ft-poundals	Candles/sq cm		3.1416		lamberts
					Candles/sq inch		0.48695		lamberts
					Centigrade thermal units (15°C)		1,898.3		joules (abs.)
					Centimeters		1 x 10 <sup>8</sup> 0.032808 0.393700 10,000 393.70 0.01093611		angstrom units feet (U.S. or British) inches (U.S.) microns mils yards (U.S.)

1/2 Avogadro's number.

continued



## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Centimeters/sec	1.9685	ft/min			Cubic inches (U.S.)	$1.63871 \times 10^{-5}$	cu m		
	0.03600	km/hr				$2.143347 \times 10^{-5}$	cu yards		
	0.6000	m/min				4.43322	drams (fluid)		
	0.02237	mi/hr				0.0043290	gallons (U.S.)		
	$3.728 \times 10^{-4}$	mi/min				0.0163868	liters		
Centimeters/sec/	0.036	km/hr/sec				0.5541	ounces (U.S., fluid)		
sec	0.02237	mi/hr/sec				0.00186010	pecks (U.S.)		
Centimeters mer-	0.013158	atmospheres				0.0297616	pints (U.S., dry)		
cury at 0°C	$1.33322 \times 10^4$	dynes/sq cm				0.0148808	quarts (U.S., dry)		
	135.95	kg/sq m				0.017316	quarts (U.S., liquid)		
	27.845	lb/sq ft				$4.65025 \times 10^{-4}$	bushels (U.S.)		
	0.19337	lb/sq inch							
	0.44604	ft water at 39.1°F			Cubic inches	16.3870253	cu cm		
Circles	6.28319	radians			(British)	$5.7870 \times 10^{-4}$	cu ft (British)		
Circular inches	5.0671	sq cm			Cubic kilometers	$1 \times 10^9$	cu m		
	0.78540	sq inches							
Circular milli-	0.78540	sq mm			Cubic meters	28.3776	bushels (U.S.)		
meters						$1 \times 10^6$	cu cm		
Circumferences	400	grades				35.314445	cu ft (U.S.)		
Coulombs (abs.)	0.1000	abcoulombs or				61,023	cu inches		
		electromagnetic				1.3079428	cu yd (U.S.)		
		cgs units				999.973	liters		
	1.00010	coulombs (Int.)				2,113.4	pints (U.S., liquid)		
	$6.281 \times 10^{18}$	electronic charges				1,056.7	quarts (U.S., liquid)		
	$2.99796 \times 10^9$	electrostatic cgs			Cubic millimeters	$6.1023 \times 10^{-5}$	cu inches		
		units or statcou-				$1 \times 10^{-9}$	cu m		
		lombs				0.016231	minims (U.S.)		
Coulombs (inter-	0.99990	coulombs (abs.)			Cubic yards	764,559.45	cu cm		
national)						764.54	liters		
Coulombs/kg	3,057.7	statcoulombs/dyne			Cubic yards (U.S.)	27	cu ft		
Cubic centimeter-	0.101325	joules (abs.)				46,656	cu inches		
atmospheres						0.76455945	cu m		
(normal)						202.0	gallons (U.S.)		
Cubic centimeters	$3.531445 \times 10^{-5}$	cu ft (U.S.)				1,616	pints (U.S., liquid)		
	0.061023	cu inches				807.9	quarts (U.S., liquid)		
	$1.3079 \times 10^{-6}$	cu yd			Days (mean solar)	24	hours (mean solar)		
	0.27053	drams (U.S., fluid)				1,440	minutes (mean solar)		
	$2.6417 \times 10^{-4}$	gallons (U.S.)				86,400	seconds (mean solar)		
	$9.9997 \times 10^{-4}$	liters			Days (siderial)	86,164	seconds (mean solar)		
	16.231	minims (U.S.)			Degrees	0.00277778	circumferences		
	0.033814	ounces (U.S., fluid)				60	minutes		
	0.0021134	pints (U.S., liquid)				1/90	quadrants		
	$9.0808 \times 10^{-4}$	quarts (U.S., dry)				0.0174533	radians		
	0.0010567	quarts (U.S., liquid)				0.00277778	revolutions		
Cubic centimeters/	0.0021186	cu ft/min				3,600	seconds		
sec					Degrees/sec	0.1667	revolutions/min		
Cubic feet (U.S.)	28,317	cu cm				0.002778	revolutions/sec		
	1,728	cu inches			Drams (apothecary or troy)	2,194286	drams (avoir.)		
	0.02831701	cu m				60	grains		
	0.037037	cu yd				3.8879351	grams		
	7.481	gallons (U.S.)				0.1371429	ounces (avoir.)		
	28.316	liters				0.125	ounces (troy)		
	25.714	quarts (U.S., dry)				2.5	pennyweights		
	29.922	quarts (U.S., liquid)				0.008571429	pounds (avoir.)		
Cubic feet/min	472.0	cu cm/sec				1/96	pounds (troy)		
	0.1247	gal/sec			Drams (avoirdupois)	0.4557292	drams (apoth. or troy)		
	0.4720	L/sec				27.34375	grains		
Cubic feet/sec	2.22222	cu yd/min				1.771845	grams		
Cubic feet water at	62.37	pounds				0.0625	ounces (avoir.)		
60°F						0.056966146	ounces (troy)		
Cubic foot-atmo-	680.74	gram calories (mean)				1/256	pounds (avoir.)		
spheres	2,116.3	ft-lb				0.0047471788	pounds (troy)		
	2,869.4	joules (abs.)			Drams (U.S., fluid or apothecary)	3.6967	cu cm		
	292.59	kg-m				0.225570	cu inches		
	28.316	L-atm				3.6966	milliliters		
Cubic inches (U.S.)	16.387162	cu cm				60	minims (U.S., fluid)		
	$5.78704 \times 10^{-4}$	cu ft (U.S.)				0.125	ounces (fluid)		
					Dyne-centimeters	$1.0197 \times 10^{-8}$	kg-m		
					(torque)	$7.3757 \times 10^{-8}$	lb-ft		
						$8.8511 \times 10^{-7}$	lb-inches		

*continued*



## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Dynes		0.015368 0.00101972 $2.2481 \times 10^{-6}$		grains grams pounds	Feet (U.S.)		$1.6447 \times 10^{-4}$ $1.893939 \times 10^{-4}$ 473,404		miles (nautical) miles (statute) wavelengths of cadmium red line
Dynes/cm		1 0.01 2,5901 0,10197		ergs/sq cm ergs/sq mm mg/inch mg/mm	Feet/min		0.508001 0.01829 0,00508001 0.011364		cm/sec km/hr mi/sec m/hr
Dynes/sq cm		$9.8692 \times 10^{-7}$ 0.0101971 0.0020886 $1.4504 \times 10^{-5}$ $7.5006 \times 10^{-4}$ $2.9530 \times 10^{-5}$ $4.0148 \times 10^{-4}$		atmospheres kg/sq m lb/sq ft lb/sq inch mm Hg inches Hg at 32°F inches water at 39.2°F (4°C)	Feet/sec		1.0973 0.5921 18,29 0.6818 0,011364		km/hr knots/hr m/min mi/hr mi/min
Electromagnetic cgs units of field strength		1,0000		gauss (abs.)	Feet/sec/sec		1.0973		km/hr/sec
Electromagnetic cgs units of magnetic permeability		$8.9916 \times 10^{20}$		electrostatic cgs units of magnetic permeability	Feet water at 39.2°F (4°C)		0.029499 304.79 0.43352 62.427 0.88265		atmospheres kg/sq m lb/sq inch lb/sq ft inches Hg at 32°F (0°C)
Electromagnetic cgs units of mass resistance		$9.9948 \times 10^{-6}$		ohm (Int.)-meter-gram	Foot-pounds		0.32389 $4.7253 \times 10^{-4}$ $1.35582 \times 10^7$ $1.3825 \times 10^4$ $5.0505 \times 10^{-7}$ 1.35582 0.138255 $3.7662 \times 10^{-7}$ 0.013381		gram calories (mean) cu ft-atm ergs g-cm horsepower-hr joules (abs.) kg-m kw-hr L-atm
Electromagnetic cgs units of reluctance		1,0000		oersteds (abs.)	Foot-pounds/min		$2.2597 \times 10^{-5}$		kilowatts
Electronic charges		$1.5921 \times 10^{-20}$ $1.5921 \times 10^{-19}$ $4.774 \times 10^{-10}$		abcoulombs coulombs (abs.) statcoulombs	Foot-pounds/sec		0.0018182 0.00135582 1.35582		horsepower kilowatts watts (abs.)
Electrons/kg		$4.868 \times 10^{-16}$		statcoulombs/dyne	Footcandles		10.764		lumens/sq m
Electrostatic cgs units of field strength		$3.33560 \times 10^{-11}$		gauss (abs.)	Gallons (U.S.)		3,785.4 0.13368 231 0.0037854 0.004951 3.78533 61,440 128 8 4 8.3378		cu cm cu ft cu inches cu m cu yd liters minims ounces (U.S., fluid) pints (U.S., liquid) quarts (U.S., liquid) pounds (avoir.) water at 62°F (16.7°C)
Ergs		$9.4805 \times 10^{-11}$ $2.3889 \times 10^{-8}$ $2.3889 \times 10^{-11}$ $7.3756 \times 10^{-8}$ 0.00101972 $1 \times 10^{-7}$ $1.0197 \times 10^{-8}$		BTU (mean) gram calories (mean) kg calories (mean) ft-lb g-cm joules kg-m	Gallons (U.S.) water/min		6.0086		tons water/24 hr
Ergs/sec		$5.6883 \times 10^{-9}$ $1.4333 \times 10^{-9}$  $4.4254 \times 10^{-6}$ $7.3756 \times 10^{-8}$ $1.3410 \times 10^{-10}$ $1 \times 10^{-10}$ $1 \times 10^{-7}$		BTU (mean)/min kg calories (mean)/min ft-lb/min ft-lb/sec horsepower kilowatts watts	Grains		63.5453 0.064798918 64.798918 0.0022857 0.0020833 1/7000 1/5760		dynes grams milligrams ounces (avoir.) ounces (troy) pounds (avoir.) pounds (troy)
Ergs/sq cm		1 0.01 100 100		dynes/cm ergs/sq mm dynes/cm ergs/sq cm	Gram-centimeters		$2.3427 \times 10^{-5}$ 980.665 $7.233 \times 10^{-5}$ $9.80665 \times 10^{-5}$ $1 \times 10^{-5}$		gram calories (mean) ergs ft-lb joules (abs.) kg-m
Faradays		$9.6500 \times 10^4$ $9.6507 \times 10^4$		coulombs (abs.) coulombs (Int.)	Gram-centimeters/sec		$9.80665 \times 10^{-5}$		watts (abs.)
Faradays/kg		$2.9507 \times 10^8$		statcoulombs/dyne	Gram-square centimeters (moment of inertia)		$2.37305 \times 10^{-6}$ $3.4172 \times 10^{-4}$		lb/sq ft lb/sq inch
Faradays/sec		96,500		amperes (abs.)					
Farads (abs.)		$1 \times 10^{-9}$ 1.00052 $1 \times 10^6$ $8.98776 \times 10^{11}$		abfarads farads (Int.) microfarads statfarads					
Farads (International)		0.99948		farads (abs.)					
Fathoms		6		feet					
Feet (U.S.)		30.48006096 0.3048006096		centimeters meters					

continued

## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C	
Grams		980.665		dynes	Joules (absolute)		$3.485 \times 10^{-4}$		cu ft-atm	
		15.4324		grains			$1 \times 10^7$		ergs	
		$1 \times 10^6$		micrograms			0.73756		ft-lb	
		1,000		milligrams			$1.0197 \times 10^4$		g-cm	
		0.0352739		ounces (avoir.)			$3.72508 \times 10^{-7}$		horsepower-hr	
		0.0321507		ounces (troy)			0.999680		joules (Int.)	
		0.00220462		pounds (avoir.)			0.101972		kg-m	
		0.00267923		pounds (troy)			$2.77778 \times 10^{-7}$		kw-hr	
		$1 \times 10^{-6}$		tons (metric)			0.0098689		L-atm	
	Grams/cu cm		62.43			lb/cu ft	Joules/°C		$5.2679 \times 10^{-4}$	
		0.03613		lb/cu inch		0.23889			gram calories/°C	
		8.3454		lb/gal (U.S.)	Joules/electron			$6.2811 \times 10^{19}$		joules/aboulomb
		0.35757		grains/cu ft		Joules/electron/°C			$6.2811 \times 10^{18}$	
Grams/L		58.417		grains/gal (U.S.)	Joules/faraday		$1.0363 \times 10^{-4}$		joules/aboulomb	
		1,000		parts per million		Joules/faraday/°C		$1.0363 \times 10^{-5}$		joules/coulomb/°C
		0.062427		lb/cu ft			Joules/gram/°C		0.23889	
		8.345		lb/1000 gal (U.S.)		Kilogram-meters			0.00929667	
Grams/ml		0.999973		g/cu cm			2.3427		gram calories (mean)	
Grams/sq cm		$9.6784 \times 10^{-4}$		atmospheres			0.0034177		cu ft-atm	
		980.665		dynes/sq cm			$9.80665 \times 10^7$		ergs	
		10		kg/sq m			232.71		ft-poundals	
		2.04817		lb/sq ft			7.2330		ft-lb	
		0.0142234		lb/sq inch			$3.6529 \times 10^{-6}$		horsepower-hr	
		0.73556		mm Hg at 0°C			$2.72407 \times 10^{-6}$		kw-hr	
Gravity		980.665		cm/sec/sec	Kilograms			$9.80665 \times 10^5$		dynes
		32.174		ft/sec/sec				2.204622341		pounds (avoir.)
Hectares		2.471044		acres (U.S.)		2.6792285		pounds (troy)		
		10,000		sq m	Kilograms/m		0.67197		lb/ft	
		11,959.85		sq yd (U.S.)		Kilograms/sq cm		14.223		lb/sq inch
Horsepower		42.418		BTU (mean)/min			73.556		cm Hg at 0°C	
		10.688		kg calories (mean)/min	Kilograms/sq m			$9.6777 \times 10^{-5}$		atmospheres
		33,000		ft-lb/min				98.0665		dynes/sq cm
		550		ft-lb/sec				0.1		g/sq cm
		1.0139		horsepower (metric)				0.204817		lb/sq ft
		0.74570		kilowatts				0.00142234		lb/sq inch
		745.70		watts				0.0032809		ft water
	32.549		ft-lb/min				0.0028959		inches Hg	
Horsepower (metric)		0.98632		horsepower (U.S.)			0.073556		mm Hg	
		75		kg-m/sec	Kilograms/sq mm		$9.80665 \times 10^7$		dynes/sq cm	
Horsepower-hour		641.304		kg calories (mean)		Kiloliters		35.317		cu ft
		$1.9800 \times 10^6$		ft-lb	Kilometers			3,280.83		feet
		$2.6845 \times 10^6$		joules (abs.)				0.539593		miles (nautical)
		$2.7374 \times 10^5$		kg-m				0.6213699495		miles (U.S.)
	0.7457		kw-hr			0.1		myriameters		
Hours (mean solar)		0.041667		days (mean solar)		1,093.6		yards		
		0.0059524		weeks	Kilometers/hr		27.7778		cm/sec	
Inches (U.S.)		$2.5400 \times 10^8$		angstrom units			54.68		ft/min	
		2.540005		centimeters			0.9113		ft/sec	
		$1.57828 \times 10^{-5}$		miles			0.5396		knots/hr	
		39,450.45		wavelengths of cadmium red line			16.667		m/min	
Inches mercury at 32°F		0.033421		atmospheres		Kilometers/hr/sec		27.778		cm/sec/sec
		$3.38639 \times 10^4$		dynes/sq cm			0.9113		ft/sec/sec	
		345.31		kg/sq m			0.27778		m/sec/sec	
		70.727		lb/sq ft	Kilometers/min			1,666.7		cm/sec
Inches water at 39.2°F (4°C)		0.0024583		atmospheres			37.2822		mi/hr	
		2,490.82		dynes/sq cm	Kilowatt-hours		3,413.0		BTU (mean)	
		25.399		kg/sq m			$8.6001 \times 10^5$		gram calories (mean)	
		5.2022		lb/sq ft			$2.6552 \times 10^6$		ft-lb	
			quanta			1.3410		horsepower-hr		
Joule-seconds		$1.5258 \times 10^{33}$		quanta		$3.6000 \times 10^6$		joules (abs.)		
Joule-seconds/ $N_0$		$2.5173 \times 10^9$		quanta		$3.6709 \times 10^5$		kg-m		
Joules (absolute)		$9.480 \times 10^{-4}$		BTU (mean)	Kilowatts		56.884		BTU (mean)/min	
		0.23895		gram calories (mean)			14.3334		kg calories (mean)/min	
		0.23918		gram calories (20°C)			$2.6552 \times 10^6$		ft-lb/hr	
		$2.3889 \times 10^{-4}$		kg calories (mean)			1.3410		horsepower	

/1/ Avogadro's number.

continued

## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Knots/hr	51.479	cm/sec			Miles (nautical)	6,080.20	feet		
	1.15155	mi/hr				1.85325	kilometers		
Lamberts	0.3183	candles/sq cm			Miles (U.S., statute)	63,360	inches		
	2.054	candles/sq inch				5,280	feet		
	1	lumens emitted/sq cm (perfectly diffusing surface)				1.609347219	kilometers		
Liter-atmospheres (normal)	24.206	gram calories (mean)			Miles/hr	44.7041	cm/sec		
	0.035316	cu ft-atm				88	ft/min		
Liters	0.028378	bushels (U.S.)				1.4667	ft/sec		
	0.035316	cu ft				0.8684	knots/hr		
	61.025	cu inches				26.82	m/min		
	0.001000027	cu m			Miles/hr/min	0.74507	cm/sec/sec		
	0.0013080	cu yd			Miles/hr/sec	44.704	cm/sec/sec		
	270.5179	drams (U.S., fluid)				1.4667	ft/sec/sec		
	0.26417762	gallons (U.S.)				0.44704	m/sec/sec		
	33.8143	ounces (U.S., fluid)			Miles/min	2,682.2	cm/sec		
	1.816192	pints (U.S., dry)				88	ft/sec		
	2.11336	pints (U.S., liquid)			Milligrams	0.01543236	grains		
	0.908096	quarts (U.S., dry)				$5.64383 \times 10^{-4}$	drams (avoir.)		
	1.056681869	quarts (U.S., liquid)				$2.57206 \times 10^{-4}$	drams (troy)		
Liters/cm/da	0.011574	sq cm/sec				$3.52739 \times 10^{-5}$	ounces (avoir.)		
Liters/min	$5.886 \times 10^{-4}$	cu ft/sec				$3.215074 \times 10^{-5}$	ounces (troy)		
	0.004403	gal/sec				$2.2046 \times 10^{-6}$	pounds (avoir.)		
Liters/sec	2.11896	cu ft/min				$2.67923 \times 10^{-6}$	pounds (troy)		
	15.8507	gallons (U.S.)/min			Milligrams/inch	0.38609	dynes/cm		
Lumens	0.001496	watts			Milligrams/L	1.0	parts per million		
Lumens/sq ft	1	footcandles			Milligrams/mm	9.80665	dynes/cm		
	10.764	lumens/sq m			Millilamberts	0.929	lumens/sq ft (with perfect diffusion)		
Lumens/sq m	0.092902	footcandles/sq ft			Milliliters	1.000027	cu cm		
	$1 \times 10^{-4}$	phots				0.061025	cu inches		
Megmhos	0.001	abmhos				0.2705179	drams (U.S., fluid)		
	2.540	megmhos/cu inch				16.2311	minims (U.S.)		
Megmhos/cu inch	0.39370	megmhos/cu cm				0.0338147	ounces (U.S., fluid)		
Mercury at 0°C	13.5951	g/cu cm			Millimeters	0.0393700	inches (U.S.)		
Meter-candles	1.000	lumens/sq m			Millimeters mercury at 0°C	0.00131579	atmospheres		
Meters	$1 \times 10^{10}$	angstrom units				1,333.22	dynes/sq cm		
	3.280833333	feet (U.S.)				1.3595	g/sq cm		
	39.3700	inches (U.S.)				13.595	kg/sq m		
	$5.39593 \times 10^{-4}$	miles (nautical)				2.7845	lb/sq ft		
	$6.2137 \times 10^{-4}$	miles (statute)				0.019337	lb/sq inch		
	$1 \times 10^9$	millimicrons				1,000	microns Hg at 0°C		
	$1 \times 10^{12}$	millionth microns			Millimicrons	10	angstrom units		
	$1.55316412 \times 10^6$	wavelengths of cadmium red line				$1 \times 10^{-7}$	centimeters		
	1.093611	yards (U.S.)			Minims (U.S.)	61.612	cu cm		
Meters/min	1.6667	cm/sec				1/60	drams (U.S., fluid)		
	0.05468	ft/sec				0.0616102	milliliters		
	0.06	km/hr				1/480	ounces (U.S., fluid)		
	0.03728	mi/hr			Minims (U.S., fluid)	0.061612	cu cm		
Meters/sec	196.8	ft/min			Minutes (time)	$6.94444 \times 10^{-4}$	days		
	3.6000	km/hr				$9.9206 \times 10^{-5}$	weeks		
	0.060000	km/min			Months (mean calendar)	30.4202	days		
	2.2369	mi/hr				730.085	hours		
	0.03728	mi/min				43,805.1	minutes		
Meters/sec/sec	3.6	km/hr/sec				$2.6283 \times 10^6$	seconds		
	2.2369	mi/hr/sec			Oersteds (abs.)	1.00052	oersteds (Int.)		
Mhos (International)/cu cm	0.99948	mhos (abs.)/cu cm			Ohm-mile-pounds	$1.7513 \times 10^{-4}$	ohm-meter-grams		
Microfarads	$1 \times 10^{-15}$	abfarads			Ohms (absolute)	0.99948	ohms (Int.)		
Microns	$1 \times 10^4$	angstrom units			Ounces (avoirdupois)	16	drams (avoir.)		
	$1 \times 10^{-4}$	centimeters				7.29166	drams (troy or apoth.)		
	$3.937 \times 10^{-5}$	inches				437.5	grains		
	0.001	millimeters				28.349527	grams		
Microvolts/°F	1.8000	microvolts/°C				0.9114583	ounces (troy or apoth.)		
						1/16	pounds (avoir.)		
						0.075954861	pounds (troy)		

continued



## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Ounces (troy or apothecary)		17.55428		drams (avoir.)	Pounds/inch		178.6		g/cm
		8		drams (troy or apoth.)	Pounds/sq ft		$4.7252 \times 10^{-4}$		atmospheres
		480		grains			478.78		dynes/sq cm
		31.103481		grams			0.48824		g/sq cm
		1.09714		ounces (avoir.)			4.8824		kg/sq m
		0.06857143		pounds (avoir.)			0.0069445		lb/sq inch
Ounces (U.S., fluid)		1/12		pounds (troy)			0.016018		ft water at 39.1°F
		29.5737		cu cm			0.35913		mm Hg at 0°C
		1.80469		cu inches	Pounds/sq ft (moment of inertia)		$4.2140 \times 10^5$		g/sq cm
		8		drams (fluid)			421.40		kg/sq cm
		1/128		gallons (U.S.)			144		lb/sq inch
		0.0295729		liters	Pounds/sq inch		0.068046		atmospheres
Pints (U.S., dry)		480		minims (U.S.)			68.946		dynes/sq cm
		1/16		pints (U.S., liquid)			70.307		g/sq cm
		0.015625		bushels (U.S.)			0.070307		kg/sq cm
		550.61		cu cm			703.07		kg/sq m
		33.6003		cu inches			144		lb/sq ft
		0.550599		liters			27.673		inches water at 39.2°F (4°C)
Pints (U.S., liquid)		0.5		quarts (U.S., dry)			51.715		mm Hg
		473.179		cu cm	Pounds/sq inch (moment of inertia)		2.9264		g/sq cm
		0.016711		cu ft			2.9264		kg/sq cm
		28.875		cu inches			0.00694444		lb/sq ft
		$6.1881 \times 10^{-4}$		cu yd	Pounds water		27.68		cu inches
		128		drams (fluid)			0.1198		gallons (U.S.)
		0.125		gallons (U.S.)	Pounds water (62°F)		0.016033		cu ft
		0.473168		liters	Pounds water/min		0.016021		cu ft/min
		7,680		minims (U.S.)			$2.670 \times 10^{-4}$		cu ft/sec
		16		ounces (U.S., fluid)	Quarts (U.S., dry)		0.03125		bushels (U.S.)
Planck's quanta		$6.554 \times 10^{-27}$		quarts (U.S., liquid)		1,101.23		cu cm	
Poises		1.000		erg-seconds		0.038889		cu ft	
Pound-feet (torque)		$1.3558 \times 10^7$		g/cm/sec		67.2006		cu inches	
Pound-inches (torque)		$1.1298 \times 10^6$		dyne-cm		1.10120		liters	
Pounds		32.174		dyne-cm		0.125		pecks (U.S.)	
Pounds (avoirdupois)		256		poundals		2		pints (dry)	
		116.6667		drams (avoir.)	Quarts (U.S., liquid)		946.358		cu cm
		$4.44852 \times 10^5$		drams (troy)			57.749		cu inches
		7,000		dynes			0.033420		cu ft
		453.5924277		grains			256.00		drams (fluid)
		0.4535924277		grams			0.25		gallons (U.S.)
		16		kilograms			0.946333		liters
		14.5833		ounces (avoir.)			32		ounces (fluid, U.S.)
		1.2152778		ounces (troy)			1.96841		quintals (metric)
		$4.464286 \times 10^{-4}$		pounds (troy)	Square centimeters		0.0010764		sq ft
Pounds (troy)		$4.5359243 \times 10^{-4}$		tons (long)		0.15500		sq inches	
		$5 \times 10^{-4}$		tons (metric)		0.00247107		sq links (Gunter's)	
		210.6514		tons (short)		$1 \times 10^{-4}$		sq m	
		96		drams (avoir.)		100		sq mm	
		5,760		drams (troy)		$1.1960 \times 10^{-4}$		sq yd	
		373.2418		grains	Square centimeters/day		$1.1574 \times 10^{-5}$		sq cm/sec
		13.165714		grams	Square feet		929.0341		sq cm
		12		ounces (avoir.)	Square feet (U.S.)		$2.29568 \times 10^{-5}$		acres
		0.822857		pounds (avoir.)			144		sq inches
		$3.6735 \times 10^{-4}$		pounds (troy)			0.09290341		sq m
Pounds/cu ft		$3.7324 \times 10^{-4}$		tons (long)		$3.58701 \times 10^{-8}$		sq mi	
		$4.1143 \times 10^{-4}$		tons (metric)		1/9		sq yd (U.S.)	
		0.016018		tons (short)	Square inches (U.S.)		6.4516258		sq cm
		16.018		g/cu cm			1/144		sq ft (U.S.)
		$5.787 \times 10^{-4}$		kg/cu m			$6.4516258 \times 10^{-4}$		sq m
		27.68		lb/cu inch			1/1296		sq yd (U.S.)
		$2.768 \times 10^4$		g/cu cm	Square kilometers		247.1044		acres (U.S.)
		1.48816		kg/cu m			$1.0764 \times 10^7$		sq ft
		0.119826		kg/m			$1 \times 10^6$		sq m
	Pounds/gal (U.S.)		0.099776		g/cu cm		0.3861006		sq mi (U.S.)
Pounds/gal (British)		0.099776		g/cu cm		$1.1960 \times 10^6$		sq yd	

continued



## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part II. CONVERSION FACTORS

Column A	x	Column B	=	Column C	Column A	x	Column B	=	Column C
Square meters		$2.471044 \times 10^{-4}$		acres (U.S.)	Tons (metric)		1,000		kilograms
		$1 \times 10^4$		sq cm			2,204.62		pounds (avoir.)
		10.76387		sq ft (U.S.)	Tons (short)		$8.8964 \times 10^8$		dynes
		1,550.0		sq inches			907.1846		kilograms
		$1 \times 10^{-6}$		sq km			2,000		pounds (avoir.)
		$1 \times 10^6$		sq mm			2,430.56		pounds (troy)
		$3.8610 \times 10^{-7}$		sq mi			0.892857		tons (long)
Square miles		1.195985		sq yd (U.S.)			0.907185		tons (metric)
		640		acres	Tons (short)/sq ft		0.94509		atmospheres
		$2.78784 \times 10^7$		sq ft	Volts (absolute)		$1 \times 10^8$		abvolts
		2,589,998		sq km			0.0033356		statvolts
		2,589,998		sq m			0.99955		volts (Int.)
Square millimeters		$3.0976 \times 10^6$		sq yd	Volts/°C		1.0000		joules/coulomb/°C
		0.01		sq cm	Watt-hours		3.4130		BTU (mean)
		0.0015500		sq inches			2,655.3		ft-pounds
Square mils		$1 \times 10^{-6}$		sq m			860.01		gram calories (mean)
		$6.4516 \times 10^{-6}$		sq cm			0.0013410		horsepower-hr
		$1 \times 10^{-6}$		sq inches			3,600		joules
Square yards (U.S.)		$6.4516 \times 10^{-4}$		sq mm			0.86001		kg calories (mean)
		$2.06612 \times 10^{-4}$		acres (U.S.)			367.09		kg-m
		8,361.31		sq cm	Watts (absolute)		$1 \times 10^7$		ergs/sec
		9		sq ft			44.254		ft-lb/min
		1,296		sq inches			0.73756		ft-lb/sec
		0.83613		sq m			0.0013410		horsepower
Statamperes		$3.22831 \times 10^{-7}$		sq mi			0.0013596		horsepower (metric)
Statcoulombs		$3.33560 \times 10^{-10}$		amperes (abs.)			1		joules/sec
Statcoulombs/kg		$3.33560 \times 10^{-9}$		coulombs (abs.)			0.014333		kg calories (mean)/min
Statcoulombs/lb		$2.0947 \times 10^{-9}$		electronic charges	Watts (International)		1.00032		watts (abs.)
Statfarads		$1.0197 \times 10^{-6}$		statcoulombs/dyne			8.1913		BTU/sq ft/min
Stathenries		$2.2481 \times 10^{-6}$		statcoulombs/dyne	Watts/sq inches		6,372.6		ft-lb/sq ft/min
Stathmos		$1.11263 \times 10^{-12}$		farads (abs.)			0.19310		horsepower
Statohms		$8.98776 \times 10^{11}$		henries (abs.)	Weeks		168		hours
Statvolts		$1.11263 \times 10^{-12}$		mhos (Int.)/cu cm			10,080		minutes
Tons (long)		$8.98776 \times 10^{11}$		ohms (abs.)			604,800		seconds
		299.796		volts (abs.)	Yards (U.S.)		91,440.183		centimeters
		1,016.0470		kilograms			$5.68182 \times 10^{-4}$		miles
		2,240		pounds (avoir.)	Years (sidereal)		365,256		days (mean solar)
		2,722.22		pounds (troy)			8,766.144		hours (mean solar)

### Part III. NUMERICAL CONSTANTS AND BINOMIAL COEFFICIENTS

Constant	Value	Reciprocal	Logarithm	n	Binomial Coefficients									
					Value of $n_x$									
					4	5	6	7	8	9	10	11	12	
$1/4 \pi$	0.7853981634	1.2732395447	1.8950898814	0	1	1	1	1	1	1	1	1	1	
$1/2 \pi$	1.5707963268	0.6366197724	0.1961198770	1	4	5	6	7	8	9	10	11	12	
$\pi$	3.1415926536	0.3183098862	0.4971498727	2	6	10	15	21	28	36	45	55	66	
$2 \pi$	6.2831853072	0.1591549431	0.7981798634	3	4	10	20	35	56	84	120	165	220	
$\sqrt{\pi}$	1.7724538509	0.5641895835	0.2485749363	4	1	5	15	35	70	126	210	330	495	
$\sqrt{2 \pi}$	2.5066282746	0.3989422804	0.3990899342	5		1	6	21	56	126	252	462	792	
$\sqrt{1/2 \pi}$	1.2533141373	0.7978845608	0.0980599385	6			1	7	28	84	210	462	924	
$1/2 \sqrt{\pi}$	0.8862269255	1.1283791671	1.9475449407	7				1	8	36	120	330	792	
e	2.7182818285	0.3678794412	0.4342944819	8					1	9	45	165	495	
e <sup>2</sup>	7.3890560989	0.1353352832	0.8685889638	9						1	10	55	220	
$\sqrt{2}$	1.4142135624	0.7071067812	0.1505149978	10							1	11	66	
$\sqrt{3}$	1.7320508076	0.5773502692	0.2385606274	11								1	12	
$\sqrt{10}$	3.1622776602	0.3162277660	0.5000000000	12									1	
Log <sub>10</sub> e	0.4342944819	2.3025850930	1.6377843113											
Radian	$57.2957795131^\circ$	0.0174532925	1.7581226324											

continued

## Appendix V. FORMULAS, FACTORS, AND CONSTANTS

### Part IV. PHYSICAL CONSTANTS

Speed of light in a vacuum (also, ratio of emu to esu of electric charge ( $c$ )) . . .	$(2.99776 \pm 0.00004) \times 10^8$ m/sec
Charge of an electron ( $e$ ) . . . . .	$(1.6020 \pm 0.0002) \times 10^{-19}$ coulomb
Faraday's constant ( $F$ ) <sup>1</sup> . . . . .	$(96,522 \pm 7)$ coulombs/mole
Avogadro's number ( $N_0$ ) <sup>2</sup> . . . . .	$(6.0251 \pm 0.0004) \times 10^{23}$
Standard atmospheric pressure ( $P_0$ ) . . . . .	$(101,324.6 \pm 0.4)$ nt/sq m
Freezing point of water on the absolute (Kelvin) scale ( $T_0$ ) . . . . .	$(273.16 \pm 0.02)$ °K
Density of mercury at STP . . . . .	$(13,595.04 \pm 0.06)$ kg/cu m
Atomic weight of oxygen, physical scale <sup>3</sup> . . . . .	$(16.00436 \pm 0.00009)$
Volume of a mole of perfect gas at STP ( $V_0$ ) . . . . .	$(22,420.7 \pm 0.6)$ cu cm
Universal gas constant ( $R_0$ ) . . . . .	$(8,316.6 \pm 0.4)$ joules/kg.°K
Boltzmann's constant ( $k$ ), the gas constant per molecule . . . . .	$(1.3803 \pm 0.0001) \times 10^{-23}$ joule/°K
Mass of atom of unit atomic weight ( $m_1$ ), physical scale <sup>3</sup> . . . . .	$(1.6589 \pm 0.0014) \times 10^{-24}$ g
Mass of electron ( $m_e$ ) . . . . .	$(9.103 \pm 0.008) \times 10^{-28}$ g
Mechanical equivalent of heat . . . . .	$(4,185.5 \pm 0.4)$ joules/kcal
Gravitation constant ( $G$ ) . . . . .	$(6.670 \pm 0.005) \times 10^{-11}$ nt·sq m/sq kg
Planck's (quantum) constant ( $h$ ) . . . . .	$(6.623 \pm 0.001) \times 10^{-34}$ joule·sec

<sup>1/1</sup> The charge transported by a gram atom of a univalent element. <sup>1/2</sup> The number of molecules in a gram molecule, or of atoms in a gram atom. <sup>1/3</sup> An atomic weight of 16 for oxygen (as determined by chemical analysis) is the basis for the chemical scale of atomic weights. In the physical scale the value of 16 is assigned to the most abundant isotope of oxygen. Physical scale atomic weights are larger than those in the chemical scale, by a ratio of 1.00027 to 1.

## Appendix VI. ATOMIC WEIGHTS

Values in parentheses are mass numbers of the most stable known isotopes.

Element	Symbol	Atomic		Element	Symbol	Atomic		Element	Symbol	Atomic	
		No.	Wt.			No.	Wt.			No.	Wt.
(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)	(A)	(B)	(C)	(D)
1 Actinium	Ac	89	(227)	35 Gold	Au	79	197.0	69 Praseodymium	Pr	59	140.92
2 Aluminum	Al	13	26.98	36 Hafnium	Hf	72	178.50	70 Promethium	Pm	61	(147)
3 Americium	Am	95	(243)	37 Helium	He	2	4.003	71 Protactinium	Pa	91	(231)
4 Antimony	Sb	51	121.76	38 Holmium	Ho	67	164.94	72 Radium	Ra	88	(226)
5 Argon	Ar	18	39.944	39 Hydrogen	H	1	1.0080	73 Radon	Rn	86	(222)
6 Arsenic	As	33	74.91	40 Indium	In	49	114.82	74 Rhenium	Re	75	186.22
7 Astatine	At	85	(210)	41 Iodine	I	53	126.91	75 Rhodium	Rh	45	102.91
8 Barium	Ba	56	137.36	42 Iridium	Ir	77	192.2	76 Rubidium	Rb	37	85.48
9 Berkelium	Bk	97	(249)	43 Iron	Fe	26	55.85	77 Ruthenium	Ru	44	101.1
10 Beryllium	Be	4	9.013	44 Krypton	Kr	36	83.80	78 Samarium	Sm	62	150.35
11 Bismuth	Bi	83	209.00	45 Lanthanum	La	57	138.92	79 Scandium	Sc	21	44.96
12 Boron	B	5	10.82	46 Lead	Pb	82	207.21	80 Selenium	Se	34	78.96
13 Bromine	Br	35	79.916	47 Lithium	Li	3	6.940	81 Silicon	Si	14	28.09
14 Cadmium	Cd	48	112.41	48 Lutetium	Lu	71	174.99	82 Silver	Ag	47	107.880
15 Calcium	Ca	20	40.08	49 Magnesium	Mg	12	24.32	83 Sodium	Na	11	22.991
16 Californium	Cf	98	(251)	50 Manganese	Mn	25	54.94	84 Strontium	Sr	38	87.63
17 Carbon	C	6	12.011	51 Mendeleevium	Md	101	(256)	85 Sulfur	S	16	32.066 <sup>1</sup>
18 Cerium	Ce	58	140.13	52 Mercury	Hg	80	200.61	86 Tantalum	Ta	73	180.95
19 Cesium	Cs	55	132.91	53 Molybdenum	Mo	42	95.95	87 Technetium	Tc	43	(99)
20 Chlorine	Cl	17	35.457	54 Neodymium	Nd	60	144.27	88 Tellurium	Te	52	127.61
21 Chromium	Cr	24	52.01	55 Neon	Ne	10	20.183	89 Terbium	Tb	65	158.93
22 Cobalt	Co	27	58.94	56 Neptunium	Np	93	(237)	90 Thallium	Tl	81	204.39
23 Copper	Cu	29	63.54	57 Nickel	Ni	28	58.71	91 Thorium	Th	90	232.05
24 Curium	Cm	96	(247)	58 Niobium	Nb	41	92.91	92 Thulium	Tm	69	168.94
25 Dysprosium	Dy	66	162.51	59 Nitrogen	N	7	14.008	93 Tin	Sn	50	118.70
26 Einsteinium	Es	99	(254)	60 Nobelium	No	102	(254)	94 Titanium	Ti	22	47.90
27 Erbium	Er	68	167.27	61 Osmium	Os	76	190.2	95 Tungsten	W	74	183.86
28 Europium	Eu	63	152.0	62 Oxygen	O	8	16	96 Uranium	U	92	238.07
29 Fermium	Fm	100	(255)	63 Palladium	Pd	46	106.4	97 Vanadium	V	23	50.95
30 Fluorine	F	9	19.00	64 Phosphorus	P	15	30.975	98 Xenon	Xe	54	131.30
31 Francium	Fr	87	(223)	65 Platinum	Pt	78	195.09	99 Ytterbium	Yb	70	173.04
32 Gadolinium	Gd	64	157.26	66 Plutonium	Pu	94	(242)	100 Yttrium	Y	39	88.92
33 Gallium	Ga	31	69.72	67 Polonium	Po	84	(210)	101 Zinc	Zn	30	65.38
34 Germanium	Ge	32	72.60	68 Potassium	K	19	39.100	102 Zirconium	Zr	40	91.22

<sup>1/1</sup> The Atomic Weights Commission recommends a range of  $\pm 0.003$ .

# Contrails

## Appendix VII. LOGARITHMS AND ANTILOGARITHMS

### Part I. FOUR-PLACE LOGARITHMS

No.											Proportional Parts								
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
10	0000	0043	0086	0128	0170	0212	0253	0294	0334	0374	4	8	12	17	21	25	29	33	37
11	0414	0453	0492	0531	0569	0607	0645	0682	0719	0755	4	8	11	15	19	23	26	30	34
12	0792	0828	0864	0899	0934	0969	1004	1038	1072	1106	3	7	10	14	17	21	24	28	31
13	1139	1173	1206	1239	1271	1303	1335	1367	1399	1430	3	6	10	13	16	19	23	26	29
14	1461	1492	1523	1553	1584	1614	1644	1673	1703	1732	3	6	9	12	15	18	21	24	27
15	1761	1790	1818	1847	1875	1903	1931	1959	1987	2014	3	6	8	11	14	17	20	22	25
16	2041	2068	2095	2122	2148	2175	2201	2227	2253	2279	3	5	8	11	13	16	18	21	24
17	2304	2330	2355	2380	2405	2430	2455	2480	2504	2529	2	5	7	10	12	15	17	20	22
18	2553	2577	2601	2625	2648	2672	2695	2718	2742	2765	2	5	7	9	12	14	16	19	21
19	2788	2810	2833	2856	2878	2900	2923	2945	2967	2989	2	4	7	9	11	13	16	18	20
20	3010	3032	3054	3075	3096	3118	3139	3160	3181	3201	2	4	6	8	11	13	15	17	19
21	3222	3243	3263	3284	3304	3324	3345	3365	3385	3404	2	4	6	8	10	12	14	16	18
22	3424	3444	3464	3483	3502	3522	3541	3560	3579	3598	2	4	6	8	10	12	14	15	17
23	3617	3636	3655	3674	3692	3711	3729	3747	3766	3784	2	4	6	7	9	11	13	15	17
24	3802	3820	3838	3856	3874	3892	3909	3927	3945	3962	2	4	5	7	9	11	12	14	16
25	3979	3997	4014	4031	4048	4065	4082	4099	4116	4133	2	3	5	7	9	10	12	14	15
26	4150	4166	4183	4200	4216	4232	4249	4265	4281	4298	2	3	5	7	8	10	11	13	15
27	4314	4330	4346	4362	4378	4393	4409	4425	4440	4456	2	3	5	6	8	9	11	13	14
28	4472	4487	4502	4518	4533	4548	4564	4579	4594	4609	2	3	5	6	8	9	11	12	14
29	4624	4639	4654	4669	4683	4698	4713	4728	4742	4757	1	3	4	6	7	9	10	12	13
30	4771	4786	4800	4814	4829	4843	4857	4871	4886	4900	1	3	4	6	7	9	10	11	13
31	4914	4928	4942	4955	4969	4983	4997	5011	5024	5038	1	3	4	6	7	8	10	11	12
32	5051	5065	5079	5092	5105	5119	5132	5145	5159	5172	1	3	4	5	7	8	9	11	12
33	5185	5198	5211	5224	5237	5250	5263	5276	5289	5302	1	3	4	5	6	8	9	10	12
34	5315	5328	5340	5353	5366	5378	5391	5403	5416	5428	1	3	4	5	6	8	9	10	11
35	5441	5453	5465	5478	5490	5502	5514	5527	5539	5551	1	2	4	5	6	7	9	10	11
36	5563	5575	5587	5599	5611	5623	5635	5647	5658	5670	1	2	4	5	6	7	8	10	11
37	5682	5694	5705	5717	5729	5740	5752	5763	5775	5786	1	2	3	5	6	7	8	9	10
38	5798	5809	5821	5832	5843	5855	5866	5877	5888	5899	1	2	3	5	6	7	8	9	10
39	5911	5922	5933	5944	5955	5966	5977	5988	5999	6010	1	2	3	4	5	7	8	9	10
40	6021	6031	6042	6053	6064	6075	6085	6096	6107	6117	1	2	3	4	5	6	8	9	10
41	6128	6138	6149	6160	6170	6180	6191	6201	6212	6222	1	2	3	4	5	6	7	8	9
42	6232	6243	6253	6263	6274	6284	6294	6304	6314	6325	1	2	3	4	5	6	7	8	9
43	6335	6345	6355	6365	6375	6385	6395	6405	6415	6425	1	2	3	4	5	6	7	8	9
44	6435	6444	6454	6464	6474	6484	6493	6503	6513	6522	1	2	3	4	5	6	7	8	9
45	6532	6542	6551	6561	6571	6580	6590	6599	6609	6618	1	2	3	4	5	6	7	8	9
46	6628	6637	6646	6656	6665	6675	6684	6693	6702	6712	1	2	3	4	5	6	7	7	8
47	6721	6730	6739	6749	6758	6767	6776	6785	6794	6803	1	2	3	4	5	5	6	7	8
48	6812	6821	6830	6839	6848	6857	6866	6875	6884	6893	1	2	3	4	4	5	6	7	8
49	6902	6911	6920	6928	6937	6946	6955	6964	6972	6981	1	2	3	4	4	5	6	7	8
50	6990	6998	7007	7016	7024	7033	7042	7050	7059	7067	1	2	3	3	4	5	6	7	8
51	7076	7084	7093	7101	7110	7118	7126	7135	7143	7152	1	2	3	3	4	5	6	7	8
52	7160	7168	7177	7185	7193	7202	7210	7218	7226	7235	1	2	2	3	4	5	6	7	7
53	7243	7251	7259	7267	7275	7284	7292	7300	7308	7316	1	2	2	3	4	5	6	6	7
54	7324	7332	7340	7348	7356	7364	7372	7380	7388	7396	1	2	2	3	4	5	6	6	7
No.	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9

*continued*



# Contrails

## Appendix VII. LOGARITHMS AND ANTILOGARITHMS

### Part I. FOUR-PLACE LOGARITHMS

No.	0	1	2	3	4	5	6	7	8	9	Proportional Parts								
											1	2	3	4	5	6	7	8	9
55	7404	7412	7419	7427	7435	7443	7451	7459	7466	7474	1	2	2	3	4	5	5	6	7
56	7482	7490	7497	7505	7513	7520	7528	7536	7543	7551	1	2	2	3	4	5	5	6	7
57	7559	7566	7574	7582	7589	7597	7604	7612	7619	7627	1	2	2	3	4	5	5	6	7
58	7634	7642	7649	7657	7664	7672	7679	7686	7694	7701	1	1	2	3	4	4	5	6	7
59	7709	7716	7723	7731	7738	7745	7752	7760	7767	7774	1	1	2	3	4	4	5	6	7
60	7782	7789	7796	7803	7810	7818	7825	7832	7839	7846	1	1	2	3	4	4	5	6	6
61	7853	7860	7868	7875	7882	7889	7896	7903	7910	7917	1	1	2	3	4	4	5	6	6
62	7924	7931	7938	7945	7952	7959	7966	7973	7980	7987	1	1	2	3	3	4	5	6	6
63	7993	8000	8007	8014	8021	8028	8035	8041	8048	8055	1	1	2	3	3	4	5	5	6
64	8062	8069	8075	8082	8089	8096	8102	8109	8116	8122	1	1	2	3	3	4	5	5	6
65	8129	8136	8142	8149	8156	8162	8169	8176	8182	8189	1	1	2	3	3	4	5	5	6
66	8195	8202	8209	8215	8222	8228	8235	8241	8248	8254	1	1	2	3	3	4	5	5	6
67	8261	8267	8274	8280	8287	8293	8299	8306	8312	8319	1	1	2	3	3	4	5	5	6
68	8325	8331	8338	8344	8351	8357	8363	8370	8376	8382	1	1	2	3	3	4	4	5	6
69	8388	8395	8401	8407	8414	8420	8426	8432	8439	8445	1	1	2	2	3	4	4	5	6
70	8451	8457	8463	8470	8476	8482	8488	8494	8500	8506	1	1	2	2	3	4	4	5	6
71	8513	8519	8525	8531	8537	8543	8549	8555	8561	8567	1	1	2	2	3	4	4	5	5
72	8573	8579	8585	8591	8597	8603	8609	8615	8621	8627	1	1	2	2	3	4	4	5	5
73	8633	8639	8645	8651	8657	8663	8669	8675	8681	8686	1	1	2	2	3	4	4	5	5
74	8692	8698	8704	8710	8716	8722	8727	8733	8739	8745	1	1	2	2	3	4	4	5	5
75	8751	8756	8762	8768	8774	8779	8785	8791	8797	8802	1	1	2	2	3	3	4	5	5
76	8808	8814	8820	8825	8831	8837	8842	8848	8854	8859	1	1	2	2	3	3	4	5	5
77	8865	8871	8876	8882	8887	8893	8899	8904	8910	8915	1	1	2	2	3	3	4	4	5
78	8921	8927	8932	8938	8943	8949	8954	8960	8965	8971	1	1	2	2	3	3	4	4	5
79	8976	8982	8987	8993	8998	9004	9009	9015	9020	9025	1	1	2	2	3	3	4	4	5
80	9031	9036	9042	9047	9053	9058	9063	9069	9074	9079	1	1	2	2	3	3	4	4	5
81	9085	9090	9096	9101	9106	9112	9117	9122	9128	9133	1	1	2	2	3	3	4	4	5
82	9138	9143	9149	9154	9159	9165	9170	9175	9180	9186	1	1	2	2	3	3	4	4	5
83	9191	9196	9201	9206	9212	9217	9222	9227	9232	9238	1	1	2	2	3	3	4	4	5
84	9243	9248	9253	9258	9263	9269	9274	9279	9284	9289	1	1	2	2	3	3	4	4	5
85	9294	9299	9304	9309	9315	9320	9325	9330	9335	9340	1	1	2	2	3	3	4	4	5
86	9345	9350	9355	9360	9365	9370	9375	9380	9385	9390	1	1	2	2	3	3	4	4	5
87	9395	9400	9405	9410	9415	9420	9425	9430	9435	9440	0	1	1	2	2	3	3	4	4
88	9445	9450	9455	9460	9465	9469	9474	9479	9484	9489	0	1	1	2	2	3	3	4	4
89	9494	9499	9504	9509	9513	9518	9523	9528	9533	9538	0	1	1	2	2	3	3	4	4
90	9542	9547	9552	9557	9562	9566	9571	9576	9581	9586	0	1	1	2	2	3	3	4	4
91	9590	9595	9600	9605	9609	9614	9619	9624	9628	9633	0	1	1	2	2	3	3	4	4
92	9638	9643	9647	9652	9657	9661	9666	9671	9675	9680	0	1	1	2	2	3	3	4	4
93	9685	9689	9694	9699	9703	9708	9713	9717	9722	9727	0	1	1	2	2	3	3	4	4
94	9731	9736	9741	9745	9750	9754	9759	9763	9768	9773	0	1	1	2	2	3	3	4	4
95	9777	9782	9786	9791	9795	9800	9805	9809	9814	9818	0	1	1	2	2	3	3	4	4
96	9823	9827	9832	9836	9841	9845	9850	9854	9859	9863	0	1	1	2	2	3	3	4	4
97	9868	9872	9877	9881	9886	9890	9894	9899	9903	9908	0	1	1	2	2	3	3	4	4
98	9912	9917	9921	9926	9930	9934	9939	9943	9948	9952	0	1	1	2	2	3	3	4	4
99	9956	9961	9965	9969	9974	9978	9983	9987	9991	9996	0	1	1	2	2	3	3	4	4

*continued*



# Contrails

## Appendix VII. LOGARITHMS AND ANTILOGARITHMS

### Part II. FOUR-PLACE ANTILOGARITHMS

Log 10											Proportional Parts								
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
.00	1000	1002	1005	1007	1009	1012	1014	1016	1019	1021	0	0	1	1	1	1	2	2	2
.01	1023	1026	1028	1030	1033	1035	1038	1040	1042	1045	0	0	1	1	1	1	2	2	2
.02	1047	1050	1052	1054	1057	1059	1062	1064	1067	1069	0	0	1	1	1	1	2	2	2
.03	1072	1074	1076	1079	1081	1084	1086	1089	1091	1094	0	0	1	1	1	1	2	2	2
.04	1096	1099	1102	1104	1107	1109	1112	1114	1117	1119	0	1	1	1	1	2	2	2	2
.05	1122	1125	1127	1130	1132	1135	1138	1140	1143	1146	0	1	1	1	1	2	2	2	2
.06	1148	1151	1153	1156	1159	1161	1164	1167	1169	1172	0	1	1	1	1	2	2	2	2
.07	1175	1178	1180	1183	1186	1189	1191	1194	1197	1199	0	1	1	1	1	2	2	2	2
.08	1202	1205	1208	1211	1213	1216	1219	1222	1225	1227	0	1	1	1	1	2	2	2	3
.09	1230	1233	1236	1239	1242	1245	1247	1250	1253	1256	0	1	1	1	1	2	2	2	3
.10	1259	1262	1265	1268	1271	1274	1276	1279	1282	1285	0	1	1	1	1	2	2	2	3
.11	1288	1291	1294	1297	1300	1303	1306	1309	1312	1315	0	1	1	1	1	2	2	2	3
.12	1318	1321	1324	1327	1330	1334	1337	1340	1343	1346	0	1	1	1	1	2	2	2	3
.13	1349	1352	1355	1358	1361	1365	1368	1371	1374	1377	0	1	1	1	1	2	2	2	3
.14	1380	1384	1387	1390	1393	1396	1400	1403	1406	1409	0	1	1	1	1	2	2	2	3
.15	1413	1416	1419	1422	1426	1429	1432	1435	1439	1442	0	1	1	1	1	2	2	2	3
.16	1445	1449	1452	1455	1459	1462	1466	1469	1472	1476	0	1	1	1	1	2	2	2	3
.17	1479	1483	1486	1489	1493	1496	1500	1503	1507	1510	0	1	1	1	1	2	2	2	3
.18	1514	1517	1521	1524	1528	1531	1535	1538	1542	1545	0	1	1	1	1	2	2	2	3
.19	1549	1552	1556	1560	1563	1567	1570	1574	1578	1581	0	1	1	1	1	2	2	2	3
.20	1585	1589	1592	1596	1600	1603	1607	1611	1614	1618	0	1	1	1	1	2	2	2	3
.21	1622	1626	1629	1633	1637	1641	1644	1648	1652	1656	0	1	1	1	1	2	2	2	3
.22	1660	1663	1667	1671	1675	1679	1683	1687	1690	1694	0	1	1	1	1	2	2	2	3
.23	1698	1702	1706	1710	1714	1718	1722	1726	1730	1734	0	1	1	1	1	2	2	2	3
.24	1738	1742	1746	1750	1754	1758	1762	1766	1770	1774	0	1	1	1	1	2	2	2	3
.25	1778	1782	1786	1791	1795	1799	1803	1807	1811	1816	0	1	1	1	1	2	2	2	3
.26	1820	1824	1828	1832	1837	1841	1845	1849	1854	1858	0	1	1	1	1	2	2	2	3
.27	1862	1866	1871	1875	1879	1884	1888	1892	1897	1901	0	1	1	1	1	2	2	2	3
.28	1905	1910	1914	1919	1923	1928	1932	1936	1941	1945	0	1	1	1	1	2	2	2	3
.29	1950	1954	1959	1963	1968	1972	1977	1982	1986	1991	0	1	1	1	1	2	2	2	3
.30	1995	2000	2004	2009	2014	2018	2023	2028	2032	2037	0	1	1	1	1	2	2	2	3
.31	2042	2046	2051	2056	2061	2065	2070	2075	2080	2084	0	1	1	1	1	2	2	2	3
.32	2089	2094	2099	2104	2109	2113	2118	2123	2128	2133	0	1	1	1	1	2	2	2	3
.33	2138	2143	2148	2153	2158	2163	2168	2173	2178	2183	0	1	1	1	1	2	2	2	3
.34	2188	2193	2198	2203	2208	2213	2218	2223	2228	2234	1	1	1	1	1	2	2	2	3
.35	2239	2244	2249	2254	2259	2265	2270	2275	2280	2286	1	1	1	1	1	2	2	2	3
.36	2291	2296	2301	2307	2312	2317	2323	2328	2333	2339	1	1	1	1	1	2	2	2	3
.37	2344	2350	2355	2360	2366	2371	2377	2382	2388	2393	1	1	1	1	1	2	2	2	3
.38	2399	2404	2410	2415	2421	2427	2432	2438	2443	2449	1	1	1	1	1	2	2	2	3
.39	2455	2460	2466	2472	2477	2483	2489	2495	2500	2506	1	1	1	1	1	2	2	2	3
.40	2512	2518	2523	2529	2535	2541	2547	2553	2559	2564	1	1	1	1	1	2	2	2	3
.41	2570	2576	2582	2588	2594	2600	2606	2612	2618	2624	1	1	1	1	1	2	2	2	3
.42	2630	2636	2642	2649	2655	2661	2667	2673	2679	2685	1	1	1	1	1	2	2	2	3
.43	2692	2698	2704	2710	2716	2723	2729	2735	2742	2748	1	1	1	1	1	2	2	2	3
.44	2754	2761	2767	2773	2780	2786	2793	2799	2805	2812	1	1	1	1	1	2	2	2	3
.45	2818	2825	2831	2838	2844	2851	2858	2864	2871	2877	1	1	1	1	1	2	2	2	3
.46	2884	2891	2897	2904	2911	2917	2924	2931	2938	2944	1	1	1	1	1	2	2	2	3
.47	2951	2958	2965	2972	2979	2985	2992	2999	3006	3013	1	1	1	1	1	2	2	2	3
.48	3020	3027	3034	3041	3048	3055	3062	3069	3076	3083	1	1	1	1	1	2	2	2	3
.49	3090	3097	3105	3112	3119	3126	3133	3141	3148	3155	1	1	1	1	1	2	2	2	3
Log 10	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9

*continued*

# Contrails

## Appendix VII. LOGARITHMS AND ANTILOGARITHMS

### Part II. FOUR-PLACE ANTILOGARITHMS

Log 10											Proportional Parts								
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
.50	3162	3170	3177	3184	3192	3199	3206	3214	3221	3228	1	1	2	3	4	4	5	6	7
.51	3236	3243	3251	3258	3266	3273	3281	3289	3296	3304	1	2	2	3	4	5	5	6	7
.52	3311	3319	3327	3334	3342	3350	3357	3365	3373	3381	1	2	2	3	4	5	5	6	7
.53	3388	3396	3404	3412	3420	3428	3436	3443	3451	3459	1	2	2	3	4	5	6	6	7
.54	3467	3475	3483	3491	3499	3508	3516	3524	3532	3540	1	2	2	3	4	5	6	6	7
.55	3548	3556	3565	3573	3581	3589	3597	3606	3614	3622	1	2	2	3	4	5	6	7	7
.56	3631	3639	3648	3656	3664	3673	3681	3690	3698	3707	1	2	3	3	4	5	6	7	8
.57	3715	3724	3733	3741	3750	3758	3767	3776	3784	3793	1	2	3	3	4	5	6	7	8
.58	3802	3811	3819	3828	3837	3846	3855	3864	3873	3882	1	2	3	4	4	5	6	7	8
.59	3890	3899	3908	3917	3926	3936	3945	3954	3963	3972	1	2	3	4	5	5	6	7	8
.60	3981	3990	3999	4009	4018	4027	4036	4046	4055	4064	1	2	3	4	5	6	6	7	8
.61	4074	4083	4093	4102	4111	4121	4130	4140	4150	4159	1	2	3	4	5	6	7	8	9
.62	4169	4178	4188	4198	4207	4217	4227	4236	4246	4256	1	2	3	4	5	6	7	8	9
.63	4266	4276	4285	4295	4305	4315	4325	4335	4345	4355	1	2	3	4	5	6	7	8	9
.64	4365	4375	4385	4395	4406	4416	4426	4436	4446	4457	1	2	3	4	5	6	7	8	9
.65	4467	4477	4487	4498	4508	4519	4529	4539	4550	4560	1	2	3	4	5	6	7	8	9
.66	4571	4581	4592	4603	4613	4624	4634	4645	4656	4667	1	2	3	4	5	6	7	9	10
.67	4677	4688	4699	4710	4721	4732	4742	4753	4764	4775	1	2	3	4	5	7	8	9	10
.68	4786	4797	4808	4819	4831	4842	4853	4864	4875	4887	1	2	3	4	6	7	8	9	10
.69	4898	4909	4920	4932	4943	4955	4966	4977	4989	5000	1	2	3	5	6	7	8	9	10
.70	5012	5023	5035	5047	5058	5070	5082	5093	5105	5117	1	2	4	5	6	7	8	9	11
.71	5129	5140	5152	5164	5176	5188	5200	5212	5224	5236	1	2	4	5	6	7	8	10	11
.72	5248	5260	5272	5284	5297	5309	5321	5333	5346	5358	1	2	4	5	6	7	9	10	11
.73	5370	5383	5395	5408	5420	5433	5445	5458	5470	5483	1	3	4	5	6	8	9	10	11
.74	5495	5508	5521	5534	5546	5559	5572	5585	5598	5610	1	3	4	5	6	8	9	10	12
.75	5623	5636	5649	5662	5675	5689	5702	5715	5728	5741	1	3	4	5	7	8	9	10	12
.76	5754	5768	5781	5794	5808	5821	5834	5848	5861	5875	1	3	4	5	7	8	9	11	12
.77	5888	5902	5916	5929	5943	5957	5970	5984	5998	6012	1	3	4	5	7	8	10	11	12
.78	6026	6039	6053	6067	6081	6095	6109	6124	6138	6152	1	3	4	6	7	8	10	11	13
.79	6166	6180	6194	6209	6223	6237	6252	6266	6281	6295	1	3	4	6	7	9	10	11	13
.80	6310	6324	6339	6353	6368	6383	6397	6412	6427	6442	1	3	4	6	7	9	10	12	13
.81	6457	6471	6486	6501	6516	6531	6546	6561	6577	6592	2	3	5	6	8	9	11	12	14
.82	6607	6622	6637	6653	6668	6683	6699	6714	6730	6745	2	3	5	6	8	9	11	12	14
.83	6761	6776	6792	6808	6823	6839	6855	6871	6887	6902	2	3	5	6	8	9	11	13	14
.84	6918	6934	6950	6966	6982	6998	7015	7031	7047	7063	2	3	5	6	8	10	11	13	15
.85	7079	7096	7112	7129	7145	7161	7178	7194	7211	7228	2	3	5	7	8	10	12	13	15
.86	7244	7261	7278	7295	7311	7328	7345	7362	7379	7396	2	3	5	7	8	10	12	13	15
.87	7413	7430	7447	7464	7482	7499	7516	7534	7551	7568	2	3	5	7	9	10	12	14	16
.88	7586	7603	7621	7638	7656	7674	7691	7709	7727	7745	2	4	5	7	9	11	12	14	16
.89	7762	7780	7798	7816	7834	7852	7870	7889	7907	7925	2	4	6	7	9	11	13	14	16
.90	7943	7962	7980	7998	8017	8035	8054	8072	8091	8110	2	4	6	7	9	11	13	15	17
.91	8128	8147	8166	8185	8204	8222	8241	8260	8279	8299	2	4	6	8	9	11	13	15	17
.92	8318	8337	8356	8375	8395	8414	8433	8453	8472	8492	2	4	6	8	10	12	14	15	17
.93	8511	8531	8551	8570	8590	8610	8630	8650	8670	8690	2	4	6	8	10	12	14	16	18
.94	8710	8730	8750	8770	8790	8810	8831	8851	8872	8892	2	4	6	8	10	12	14	16	18
.95	8913	8933	8954	8974	8995	9016	9036	9057	9078	9099	2	4	6	8	10	12	15	17	19
.96	9120	9141	9162	9183	9204	9226	9247	9268	9290	9311	2	4	6	8	11	13	15	17	19
.97	9333	9354	9376	9397	9419	9441	9462	9484	9506	9528	2	4	7	9	11	13	15	17	20
.98	9550	9572	9594	9616	9638	9661	9683	9705	9727	9750	2	4	7	9	11	13	16	18	20
.99	9772	9795	9817	9840	9863	9886	9908	9931	9954	9977	2	5	7	9	11	14	16	18	20
Log 10	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9

# *Contrails*

# Contrails

## INDEX

It is suggested that the index be used in conjunction with the table of contents: the index to locate data for a specific organism, and the table of contents to determine the scope of the data for a particular topic. To facilitate identification, the index includes the taxonomic order for animals, and the family for plants, unless otherwise specified. As a further aid, the index lists the animals and plants as they are presented in the tables. Entries for a particular organism may therefore be found under the common name, under the scientific name, or under both. Where information is available under both, cross-references make the data easily accessible.

\* indicates diagram, drawing, or graph  
Fn. indicates footnote material

- Abbreviations and symbols, page xviii
- Abies* (fir), Pinaceae (*see also* Fir)  
parasites, 511, 512, 514  
propagation methods, 73  
soil pH, 442
- A. alba* (silver fir), 428
- A. anabilis* (Cascade fir), 514
- A. concolor* (white fir)  
chromosome number, 8  
first flowering, 110  
life span, 110  
measurements, 110  
parasites, 514  
seed germination, 76  
shade tolerance, 118
- A. grandis* (grand fir), 514
- A. lasiocarpa* (alpine fir), 514
- A. lasiocarpa arizonica* (cork-bark fir), 514
- A. procera* (noble fir), 111, 113
- A-B-O blood group system  
agglutinins, 245  
antiserums, 245  
distribution in various populations, 250, 251  
heredity, 249  
phenotypes and genotypes, 245
- Abomasum, parasites, 490, 491
- Abrus precatorius* (rosary pea), Leguminosae, 344
- Absidia*, Mucoraceae, 518
- Absorption maxima of  
cytochromes, 206, 207  
provitamins, 397  
vitamins, 395, 397
- Acanthophis antarcticus* (death adder), SERPENTES, 330
- Acanthoscelides obtectus* (bean weevil), COLEOPTERA, 481
- Acanthurus* (surgeonfish), PERCOMORPHI, 336, 338
- Acer* (maple), Aceraceae  
breeding system, 72  
parasites, 506, 512, 514  
pollen life span, 114  
propagation methods, 73  
soil pH, 442
- A. platanoides* (Norway maple), 213
- A. pseudoplatanus* (plane-tree maple), 214, 230
- A. rubrum* (red maple), 229, 453
- A. saccharinum* (silver maple)  
chromosome number, 8  
measurements, 111  
seed germination, 77  
seed life span, 111  
shade tolerance, 443
- A. saccharum* (sugar maple), 228, 406, 412 (*see also* Maple, sugar)
- A. tschonoski* (Tschonoski maple), 213
- Aceratogallia sanguinolenta* (leafhopper), HOMOPTERA, 502
- Acetabularia*, Dasycladaceae, 7
- Acid(s) (*see also* Amino acids, Carbohydrates, Lipids, Purines, Pyrimidines, Vitamins)  
antimetabolites, 309-311  
effect on cell elongation, 307, 308  
in feces, 190  
as growth stimulators, 175, 176  
pK values, 544  
in urine, 186, 187
- Acid-base, blood (*see* table of contents, page vii)
- Acid-base indicators, 545
- Acidosis, 262, 263
- Acipenser* (sturgeon), CHONDROSTEI, 145, 153
- A. fulvescens* (lake sturgeon), 64, 104, 107
- A. ruthenus* (sterlet), 107
- Acmaea dorsuosa* (limpet), ARCHAEOGASTROPODA, 109
- Aconitum napellus* (aconite monkshood), Ranunculaceae, 344
- Acorn worm (*see* *Saccoglossus*)
- Actinia* (sea anemone), ACTINIARIA, 110, 341
- Actinobacillus*, Brucellaceae, 504
- Actinomyces*, Actinomycetaceae  
antibiotic activity against, 322  
culture medium for, 534  
parasitism, 504, 518  
temperature for growth, 438  
thermal death time, 439
- Adamsia* (cloak anemone), ACTINIARIA, 341
- Adder  
death (*see* *Acanthophis*)  
puff (*see* *Bitis*)
- Adelopus*, Venturiaceae, 511
- Adenohypophysis  
comparative anatomy, 152  
hormones, 290-293
- Adenovirus, 431, 498
- Adiantum pedatum* (American maidenhair), Polypodiaceae, 8, 443
- Adrenal(s)  
cells, staining methods for, 552  
comparative anatomy, 154  
parasite, 520  
tissue growth, 46
- Adrenal cortex  
effect of hormones on, 291  
hormones of, 296-299
- Adrenal medulla, hormones, 298
- Aedes* (mosquito), DIPTERA  
chromosome number, 4  
diapause, 419  
dispersion, 420  
life span, 109  
parasite vector, 478, 486, 490, 498
- A. aegypti* (yellow-fever mosquito)  
dispersion, 420  
hemolymph volume, 266



- Aedes aegypti* (concluded)  
 metamorphosis, 67  
 nutrition, 175Fn.  
 oxygen consumption, 223  
 propagation, 67
- A. leucocelaenus* (forest mosquito), 420
- Aeonium haworthi* (Haworth aeonium), Crassulaceae, 451
- Aerobacter*, Enterobacteriaceae  
 antibiotic activity against, 322  
 generation time, 51  
 nutrition, 165Fn.  
 respiration rate, 225  
 temperature for growth, 438  
 thermal death time, 439
- Aetobatus narinari* (spotted duck-billed ray), BATOIDEI, 338
- Afferents, thalamic nuclei, 125, 126
- Agalliopsis novella* (leafhopper), HOMOPTERA, 502
- Agammaglobulinemia, 11
- Agaricus*, Agaricaceae, 6, 226
- Age and  
 arterial blood pressure, 239  
 body composition, 119\*  
 body height, 93, 94  
 body length, 102-105  
 body weight, 93, 102, 104, 105  
 developmental stages, 82-92  
 erythrocyte values, 268  
 heart rate, 234  
 hemoglobin values, 268  
 leukocyte counts, 272, 273  
 platelet count, 271  
 seed-bearing, 76, 77
- Agglutinin, 245
- Agglutininogen, 245, 248
- Aglycone fractions, glycosides, 369, 371
- Agriotes* (click beetle), COLEOPTERA, 420, 481
- Agrobacterium*, Rhizobiaceae, 438, 507, 508
- Agropyron* (wheatgrass), Gramineae, 428
- Agrostemma githago* (corn cockle), Caryophyllaceae, 344
- Agrostis* (bent grass), Gramineae, 494
- A. nebulosa* (cloud bent grass), 444
- Air layering, 73, 74
- Alabama argillacea* (cotton leafworm), LEPIDOPTERA, 481
- Albinism, ocular, 11
- Albula vulpes* (ladyfish), ISOSPONDYLI, 336, 338
- Alca* (auk), ALCIDAE, 148
- Aldaric acids, properties, 359
- Alder (see *Alnus*)
- Alditols, properties, 356
- Aldonic acids, properties, 358
- Aldosamines, properties, 355
- Aldoses, properties, 351, 352
- Aldrich syndrome, 11
- Alectoria*, Usneaceae, 227
- Aleurites fordii* (tung oil tree), Euphorbiaceae, 380
- Aleurodiscus*, Theleporaceae, 511, 513
- Alfalfa (see also *Medicago*)  
 parasites  
 arthropod, 482, 483, 485  
 nematode, 494, 496  
 viral, 432, 433, 501  
 sterol source, 386
- Alfalfa bug (see *Spissistilus*)
- Algae (see specific genus)
- Alimentary canal (see also Gastrointestinal tract)  
 tissue growth, 46  
 water content, 401, 402
- Alkalosis, 263
- Allelic genes (see specific blood group system)
- Allescheria*, Eurotiaceae, 518
- Alligator mississippiensis* (American alligator), CROCODYLIA  
 acid-base, blood, 260  
 blood volumes, 265, 269  
 brachial vein, 148  
 chromosome number, 1  
 erythrocyte values, 269  
 heart rate, 235  
 hemoglobin values, 269  
 life span, 107  
 oxygen consumption, 222  
 propagation, 62
- Allium* (onion), Liliaceae, 508 (see also Onion)
- A. cepa* (garden onion)  
 breeding system, 72  
 cell division, 53, 54  
 chromosome number, 8  
 light and temperature for flowering, 446  
 mineral content, 405, 411  
 parasites, 500, 506  
 pollen life span, 114  
 respiration rate, 228, 230  
 seed germination, 75  
 seed life span, 111, 113  
 soil pH, 442
- A. victorialis* (long-root onion), 213
- Almond, 368, 388 (see also *Prunus*)
- Alnus* (alder), Betulaceae  
 breeding system, 72  
 parasites, 512, 514  
 propagation methods, 73  
 soil pH, 442
- A. glutinosa* (European alder), 114, 406
- A. rubra* (red alder)  
 chromosome number, 9  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 seed germination, 77
- Aloe* (aloe), Liliaceae, 368
- Alouatta balzabul* (howler monkey), PRIMATES, 158, 160, 162
- Alternaria*, Dematiaceae  
 parasitism, 509, 510  
 respiration rate, 226  
 temperature for growth, 440  
 thermal death time, 440
- Althea* (see *Hibiscus*)
- Altica* (flea beetle), COLEOPTERA, 481
- Altitude, respiratory gases at, 219
- Alutera scripta* (longtail filefish), PLECTOGNATHI, 336, 338
- Amanita phalloides* (death cup), Agaricaceae, 344
- Amaranthus retroflexus* (redroot amaranth), Amaranthaceae, 453
- Ambergris, 385
- Amblyomma americanum* (lone star tick), ACARI, 477
- Ambrosia trifida* (giant ragweed), Compositae, 454
- Ambystoma maculatum* (spotted salamander), CAUDATA, 63, 107
- A. tigrinum* (tiger salamander) (see also Salamander, tiger)  
 blood volumes, 269  
 chromosome number, 2  
 erythrocyte and hemoglobin values, 269  
 life span, 107  
 propagation, 63
- Amelogenesis imperfecta, 11
- Amia* (bowfin), PROTOSPONDYLI, 64, 107, 153
- Amino acid(s)  
 antimetabolites, 309-311

- in biosynthesis, 208\*-210\*  
 digestion, 183\*  
 in enzymes, 289, 291  
 in feces, 190  
 metabolism, 199, 200, 203\*  
 in nitrogen cycle, 218\*  
 in nucleoprotein catabolism, 201\*  
 nutritional requirement for, 168, 169, 177, 179, 181  
 properties, 392, 393  
 RNA codewords for, 43  
 in urine, 186
- Amino sugars, properties, 355
- Amoeba*, AMOEBINA  
 cell division, 53  
 chromosome number, 5  
 oxygen consumption, 224  
 propagation, 71
- Amorpha canescens* (leadplant), Leguminosae, 456  
*Amorphophallus rivieri* (devil's-tongue), Araceae, 115  
*Amphibolips confluenta* (oak gall wasp), HYMENOPTERA, 481
- Amphioxus (*see Branchiostoma*)
- Amphiuma means* (two-toed amphiuma), CAUDATA  
 blood volumes, 269  
 chromosome number, 2  
 erythrocyte and hemoglobin values, 269  
 life span, 107
- A. tridactylum* (three-toed amphiuma), 63  
*Anabaena*, Nostocaceae, 218\*, 227, 441  
*Anabrus simplex* (Mormon cricket), ORTHOPTERA, 481  
*Anacystis*, Chroococcaceae, 227  
*Ananas comosus* (pineapple), Bromeliaceae, 451 (*see also* Pineapple)
- Anaphase, effect of compounds on, 53  
*Anaplasma*, Anaplasmataceae, 503  
*Anas* (duck), ANSERIFORMES (*see also* Duck)  
 leukocyte counts, 274  
 lung ventilation, 220  
 oxygen consumption, 221  
 propagation, 59
- A. platyrhynchos* (mallard duck)  
 arterial blood pressure, 240  
 blood volumes, 269  
 chromosome number, 1  
 clutch size, 60  
 erythrocyte values, 269  
 hatching success, 60  
 heart rate, 235  
 hemoglobin values, 269  
 life span, 106
- A. platyrhynchos domesticus* (Pekin duck), 101, 264  
*Anasa tristis* (squash bug), HEMIPTERA, 481  
*Anastatus bifasciatus* (gypsy moth egg parasite), HYMENOPTERA, 420  
*Anax imperator* (darnier), ODONATA, 419  
 Anchovy (*see Engraulis*)  
*Ancistrodon acutus* (Mexican copperhead), SERPENTES, 1  
*A. contortrix* (southern U.S. copperhead), 328  
*A. contortrix mokeson* (northern U.S. copperhead), 62, 102, 107  
*A. piscivorus* (eastern cottonmouth), 328  
*Ancylostoma* (hookworm), RHABDITIDA, 486  
*A. caninum* (dog hookworm), 490  
*Andropogon scoparius* (little bluestem), Gramineae, 454, 456
- Anemia, hypochromic, 11  
*Anemone* (sea anemone), ACTINIARIA, 223, 341  
 Anesthetics, animal dosages, 547-549  
*Angelica* (angelica), Umbelliferae, 379  
 Angiokeratoma, diffuse, 11  
*Anguilla* (freshwater eel), APODES, 236, 240  
*A. anguilla* (European freshwater eel), 2, 107, 222  
*A. rostrata* (American freshwater eel), 64, 107, 270  
*Anguina* (wheat gall eelworm), TYLENCHIDA, 494  
*Anguis fragilis* (slowworm), SAURIA  
 chromosome number, 1  
 heart rate, 235  
 life span, 107  
 oxygen consumption, 222  
 propagation, 62
- Animals, number of species, 561
- Anions, metabolism, 192\*, 194, 195  
*Anisotarsus* (beetle), COLEOPTERA, 490  
*Anodonta* (mussel), EULAMELLIBRANCHIA, 70  
*Anolis carolinensis* (American "chameleon"), SAURIA  
 acid-base, blood, 260  
 body length, 102  
 chromosome number, 1  
 propagation, 62
- A. equestris* (giant Cuban "chameleon"), 107  
*Anopheles* (mosquito), DIPTERA  
 dispersion, 420, 423  
 parasite vector, 478, 488, 490
- A. gambiae* (African malaria mosquito), 420  
*A. quadrimaculatus* (malaria mosquito), 237, 420, 424  
*Anser* (goose), ANSERIFORMES (*see also* Goose)  
 arterial blood pressure, 240  
 heart rate, 235  
 lung ventilation, 220  
 lymph hearts, 148  
 oxygen consumption, 221  
 propagation, 59
- A. albifrons* (white-fronted goose), ANSERIFORMES, 1  
*A. anser* (graylag goose), 101  
*A. domesticus* (common goose), 106, 269
- Ant  
 cornfield (*see Lasius*)  
 red, 371 (*see also Formica*)
- Anteater (*see Tachyglossus*)
- Antedon* (feather star), ARTICULATA, 69
- Antelope, 491
- Anthoceros*, Anthocerataceae, 7  
*Anthonomus grandis* (boll weevil), COLEOPTERA, 420, 482  
*Anthophora retusa* (digger bee), HYMENOPTERA, 237  
*Anthrenus verbasci* (varied carpet beetle), COLEOPTERA, 419
- Antiaris toxicaria* (upas tree), Moraceae, 344
- Antibiotics  
 biological activity, 319-324  
 effect on cell division, 54  
 properties, 312-317
- Anticoagulants, 257\*, 325-327  
 Antilogarithms, 582, 583  
 Antimetabolites, 309-311  
*Antirrhinum* (snapdragon), Scrophulariaceae, 75  
*A. majus* (snapdragon)  
 chromosome number, 9  
 light exposures for development, 444  
 light and temperature for flowering, 446  
 parasite, 506  
 pollen life span, 114  
 respiration rates, 230, 231  
 soil pH, 442
- Antiserums, phenotypic reactions with, 245-248
- Aorta  
 blood pressure, 241  
 comparative anatomy, 144, 145  
 nerve connections, 133, 136, 138  
 sterol source, 385
- Apanteles glomeratus* (little braconid), HYMENOPTERA, 419  
*A. militaris* (parasitic wasp), 67
- Ape, 486, 491, 503

- Aphanomyces*, Saprolegniaceae, 509, 510  
*Aphelenchoides besseyi* (summer crimp nematode of strawberry), TYLENCHIDA, 494  
*A. cocophilus* (coconut palm nematode), 494  
*A. fragariae* (spring crimp nematode of strawberry), 494  
*A. ritzema-bosi* (chrysanthemum nematode), 494  
Aphids, 426 (see also *Aphis*, *Macrosiphum*)  
    green peach (see *Myzus*)  
    strawberry (see *Capitophorus*)  
    woolly apple (see *Eriosoma*)  
*Aphis* (aphid), HOMOPTERA, 500-502  
*A. pomi* (apple aphid), 482  
*Aphodius* (beetle), COLEOPTERA, 490  
Apical meristem, cell description, 44  
*Apis* (honeybee), HYMENOPTERA, 69 (see also Honeybee)  
*A. mellifera* (honeybee)  
    chromosome number, 4  
    dispersion, 420, 421  
    hemolymph volume, 266  
    life span, 109  
    metamorphosis, 67  
    oxygen consumption, 223  
    propagation, 67  
*Apistus carinatus* (scorpion fish), SCLEROPAREI, 338  
*Aplysia* (sea hare), PLEUROCOELA  
    arterial blood pressure, 241  
    chromosome number, 4  
    heart rate, 237  
    life span, 109  
    oxygen consumption, 223  
*Apocynum* (dogbane), Apocynaceae, 368  
Apple, 426, 481-483, 485 (see also *Malus*)  
    crab (see *Malus*)  
    of Peru (see *Nicandra*)  
Apricot, 482  
*Aprion virescens* (snapper), PERCOMORPHI, 336, 338  
*Aptenodytes* (penguin), SPHENISCIFORMES, 59  
*A. patagonica* (king penguin), 106  
*Apus apus* (swift), APODIFORMES, 418  
*Arachis hypogaea* (peanut), Leguminosae, 216, 380 (see also Peanut)  
*Arbacia* (sea urchin), ARBACIOIDA  
    cell division, 53, 54  
    chromosome number, 3  
    propagation, 69  
Arborvitae (see *Thuja*)  
*Arceuthobium* (mistletoe), LORANTHACEAE, 514, 515  
*Archilochus* (hummingbird), APODIFORMES, 59  
*A. colubris* (ruby-throated hummingbird), 235  
*Arctium minus* (smaller burdock), COMPOSITAE, 453  
*Arenicola* (lugworm), POLYCHAETA, 223, 237  
*Argas persicus* (fowl tick), ACARI, 477  
Argus (see *Cephalopholis*)  
Armadillo, 493 (see also *Dasybus*, *Euphractus*)  
*Armigeres* (mosquito), DIPTERA, 486  
*Armillaria*, Agaricaceae, 509-513  
Armyworm (see *Prodenia*, *Pseudaletia*)  
*Arothron* (puffer), PLECTOGNATHI, 338, 339  
*A. meleagris* (white-spotted puffer), 338, 339  
*A. nigropunctatus* (black-spotted puffer), 338, 339  
*Artemia salina* (brine shrimp), ANOSTRACA, 4  
Arterial blood, acid-base balance, 259-262\*  
Arterial blood pressure, 238-242  
Artery(ies)  
    blood pressure, 241, 242  
    comparative anatomy, 144-147  
    nerve connections, 131, 132, 136, 138  
    parasite, 491  
Artificial sea water, 541, 542  
*Ascaridia galli* (large roundworm of chicken), RHABDITIDA, 490  
*Ascaris* (large roundworm), RHABDITIDA, 70  
*A. lumbricoides* (large roundworm of man), 4, 223, 486  
*A. lumbricoides suum* (large roundworm of swine), 490  
*Ascidia* (sea squirt), ENTEROGONA, 223  
*Asclepias syriaca* (milkweed), ASCLEPIADACEAE, 373  
*Ascochyta*, Sphaerioidaceae, 509, 510  
Ash, 483 (see also *Fraxinus*)  
*Asparagus albus* (white asparagus), LILIACEAE, 230  
*A. officinalis* (garden asparagus)  
    breeding system, 72  
    chromosome number, 8  
    mineral content, 405, 411  
    parasites, 508  
    respiration rate, 229  
    seed germination, 75  
    soil pH, 442  
Aspen, 483 (see also *Populus*)  
Aspergilli, culture medium for, 536  
*Aspergillus*, Aspergillaceae, Moniliaceae  
    antibiotic source, 314  
    chromosome number, 6  
    nutrition, 174Fn., 178Fn., 179Fn., 181Fn.  
    parasitism, 518  
    respiration rate, 226  
    sterol source, 385  
    temperature for growth, 440  
    thermal death time, 440  
*Aspidiotus perniciosus* (San Jose scale), HOMOPTERA, 482  
*Asplanchna* (rotifer), MONOGONONTA, 4, 110  
Ass, 479  
Assassin bug (see *Rhodinus*)  
*Astacus* (crayfish), DECAPODA  
    arterial blood pressure, 241  
    chromosome number, 4  
    heart rate, 237  
    life span, 109  
    oxygen consumption, 223  
*Astasia* (flagellate), EUGLENOIDINA, 71, 175Fn., 177Fn., 179Fn.  
*Asterias* (starfish), FORCIPULATA  
    chromosome number, 3  
    life span, 109  
    oxygen consumption, 223  
    propagation, 66, 69  
*Asthenosoma* (sea urchin), ECHINOTHURIOIDA, 340  
*Astragalus* (locoweed), Leguminosae, 344  
*A. crassicarpus* (ground plum milk vetch), 456  
Atmosphere, respiratory gases of the, 219  
Atomic weights, 579  
Atrium(a)  
    blood pressure, 241  
    water content, 401, 402  
*Atropa belladonna* (belladonna), SOLANACEAE, 345  
*Atropellis*, Helotiaceae, 513  
Atrophy, peroneal, 11  
*Attalea funifera* (piassava attalea), PALMAE, 380  
Auk (see *Alca*)  
*Aurelia* (scyphomedusa), SEMAEOSTOMAE, 5, 70, 223  
Autonomic nervous system (see table of contents, page vi)  
*Avena* (oat), Gramineae (see also Oat)  
    cell elongation, 307, 308  
    photosynthesis, 213  
    protoplasmic streaming, 448-450  
*A. fatua* (wild oat), 112  
*A. sativa* (common oat)  
    breeding system, 72  
    cell sap composition, 404  
    chromosome number, 8  
    light and temperature for flowering, 446  
    mineral content, 405, 411  
    parasites, 506, 509

† Class



- respiration rate, 228
- seed germination, 75
- seed life span, 111
- soil pH, 442
- transpiration rate, 453
- Avocado, 496 (see also *Persea*)
- Axolotl, anesthetic for, 548
- Azalea, 427, 496 (see also *Rhododendron*)
- Azotobacter*, Azotobacteraceae
  - generation time, 51
  - in nitrogen cycle, 218\*
  - nitrogen fixation, 217
  - nutrition, 165Fn.
  - respiration rate, 225
  - temperature for growth, 438
- Babesia*, EUCOCCIDIA, 492
- Baboon, 520
- Bacillus*, Bacillaceae
  - antibiotic activity against, 319, 320, 322, 323
  - antibiotic source, 312, 314
  - chemical constituents, 400
  - culture medium for, 534, 535
  - generation time, 51
  - in nitrogen cycle, 218\*
  - nutrition, 171Fn.
  - parasitism, 504, 508
  - respiration rate, 225
  - temperature for growth, 438
  - thermal death time, 439
- Bacteria (see also specific genus)
  - culture media for, 534
  - staining methods for, 556
- Badger, 491
- Bagre marina* (sea catfish), OSTARIOPHYSI, 336
- Bagworm (see *Thyridopteryx*)
- Balaena mysticetus* (Greenland whale), CETACEA, 380
- Balaenoptera physalus* (finback whale), CETACEA
  - body surface area constants, 120
  - body weight, 120
  - life span, 106
  - propagation, 57
  - skeletal system, 159, 161, 163
- Balantidium*, TRICHOSTOMATIDA, 488, 492
- Balistoides niger* (triggerfish), PLECTOGNATHI, 336, 338
- Balloonfish (see *Diodon*)
- Bamboo, 382
- Bamboo worm (see *Clymenella*)
- Banana, 496
- Band cell counts, bone marrow, 275, 276
- Baptisia leucophaea* (Plains wild indigo), Leguminosae, 456
- Barbados nut (see *Jatropha*)
- Bark
  - diseases affecting, 507, 512
  - glycoside source, 368, 370
  - mineral content, 406-409, 412, 413
  - sterol source, 386
  - toxins, 346, 348, 349
- Barley (see also *Hordeum*)
  - enzyme source, 388
  - parasites, 427, 482, 484
  - protein source, 388
- Barnacle (see *Lepas*)
- Barracuda (see *Sphyraena*)
- Bartonella*, Bartonellaceae, 438, 503
- Bases, pK values, 544
- Basidiobolus*, Entomophthoraceae, 518
- Basidiomycetes (see also specific genus)
  - culture medium for, 536
- Basophil, facing page 276\*
- Basophil count, 272-274
- Bass
  - largemouth (see also *Micropterus*)
    - anesthetic for, 548
    - sea (see *Mycteroperca*, *Plectropomus*)
- Basswood, 483
- Bat, 493
  - brown (see *Eptesicus*, *Myotis*, *Pipistrellus*)
  - horseshoe (see *Rhinolophus*)
  - long-eared (see *Plecotus*)
  - noctule (see *Nyctalus*)
  - parti-colored (see *Vespertilio*)
- Batrachus cirrhosus* (toadfish), HAPLODOCI, 339
- B. didactylus* (paddefisk), 339
- B. grunniens* (toadfish), 339
- Bdellonyssus sylviarum* (northern fowl mite), ACARI, 477
- Bean
  - leaf expansion, 308, 309
  - parasites
    - arthropod, 481, 483, 485
    - nematode, 494
    - viral, 433, 501
  - broad (see *Vicia*)
  - horse- (see *Vicia*)
  - kidney (see *Phaseolus*)
  - soy- (see *Glycine*)
  - velvet (see *Stizolobium*)
- Bear, 486, 491
- Bedbug (see *Cimex*)
- Bedstraw (see *Galium*)
- Bee
  - digger (see *Anthophora*)
  - honey- (see *Apis*)
  - leaf-cutting (see *Megachile*)
- Beech (see *Fagus*)
- Beet (see also *Beta*)
  - parasites
    - nematode, 494, 496
    - fungal, 426
    - viral, 426, 500
  - sugar, 426, 435, 500 (see also *Beta*)
- Beetle
  - bean (see *Epilachna*)
  - blister (see *Epicauta*)
  - carpet (see *Anthrenus*)
  - click (see *Agriotes*)
  - cucumber (see *Diabrotica*)
  - diving (see *Dytiscus*)
  - dung, 492
  - elm bark, 427 (see also *Hylurgopinus*, *Scolytus*)
  - flea (see *Altica*, *Epitrix*, *Phyllotreta*)
  - flour (see *Tribolium*)
  - grain, 486 (see also *Laemophloeus*, *Oryzaephilus*)
  - ground, 492
  - Japanese (see *Popillia*)
  - June (see *Melolontha*, *Phyllophaga*)
  - lady (see *Hippodamia*)
  - ladybird (see *Epilachna*)
  - mountain pine (see *Dendroctonus*)
  - potato (see *Leptinotarsa*)
  - turpentine (see *Dendroctonus*)
- Beggiatoa*, Beggiatoaceae, 181Fn.
- Belladonna (see *Atropa*)
- Bellis perennis* (English daisy), Compositae, 451
- Belonolaimus gracilis* (sting nematode), TYLENCHIDA, 494
- Bemisia gossypiperda* (whitefly), HOMOPTERA, 501
- Benzoic acids
  - nutritional requirement for, 175
  - as plant growth regulators, 307-309
- Beroe (comb jelly), BEROIDA, 70



- Beta* (beet), Chenopodiaceae, 428 (see also Beet)
- B. saccharifera* (sugar beet), 404, 446
- B. vulgaris* (common beet)
- breeding system, 72
  - chromosome number, 9
  - light and temperature for flowering, 446
  - mineral content, 406, 412
  - osmotic potential, 453
  - parasites, 435, 500, 506, 509
  - photosynthesis, 213
  - pollen dispersion, 430
  - respiration rates, 228, 230
  - seed germination, 75
  - seed life span, 112
  - shade tolerance, 443
  - soil pH, 442
- Betula* (birch), Betulaceae, 72, 73, 512 (see also Birch)
- B. lenta* (sweet birch)
- chromosome number, 9
  - first flowering, 111
  - life span, 111
  - measurements, 111
  - seed germination, 77
  - shade tolerance, 443
  - soil pH, 442
- B. lutea* (yellow birch), 114, 453
- B. nana* (dwarf arctic birch), 230, 232
- B. pendula* (European white birch), 213
- B. populifolia* (gray birch), 406
- Bichir (see *Polypterus*)
- Bifusella*, Phacidiaceae, 511, 513
- Bile, minerals excreted in, 192\*, 193-195
- Bile ducts, parasites, 489, 493
- Billbug (see *Calendra*)
- Biosynthesis (see table of contents, page vii)
- Birch, 483 (see also *Betula*)
- Bitis arietans* (puff adder), SERPENTES, 330
- Black snake (see *Coluber*, *Pseudochis*)
- Blackbird (see also *Turdus*)
- anesthetic for, 548
- Bladder
- nerve connections, 130\*, 132, 133, 135
  - parasite, 489
  - radiation effect on, 469
- Blastodermic vesicle, 79\*-81\*
- Blastomyces*, Moniliaceae, 319, 321, 518
- Blastula, developmental stages, 82, 84, 86, 89
- Blatta orientalis* (Oriental cockroach), ORTHOPTERA, 223
- Blattella germanica* (German cockroach), ORTHOPTERA, 168Fn., 176Fn.
- Blissus leucopterus* (chinch bug), HEMIPTERA, 482
- Blood (see also table of contents, pages xiii, xiv)
- parasites, 488-493, 498-500
  - radiation effect on, 468-475
- Blood agar media for protozoa, 526
- Blood factors, 253\*-258\*
- Blood group systems (see table of contents, page vii)
- Blood loss, erythrocyte recovery after, 47
- Blood pressures, 239-243
- Blood proteins, properties, 388, 389
- Blood smears, staining method for, 555
- Blood vessels
- comparative anatomy, 144-149
  - nerve connections, 130\*, 131-136
- Blood volumes, 263-265
- Blueberry, 485
- Bluegill (see also *Lepomis*)
- anesthetic for, 548
- Bluetongue virus, 498
- BMH medium for axenic culture, 524
- Boa* (boa), SERPENTES, 146
- Body composition, with increasing weight and age, 119\*
- Body fat
- with increasing weight and age, 119\*
  - radiation effect on, 472
- Body height
- and body surface area, 120
  - various races and nationalities, 93, 94
- Body length, 102-105
- Body organs (see also specific organs)
- permissible radiation exposure for, 457
  - permissible radionuclide concentration in, 458-467
- Body surface area formula, constants for, 120, 121
- Body temperature
- brain involvement, 127-129
  - during hibernation, 417, 418
- Body weight, 94-102, 104, 105
- at birth, 82
  - and body composition, 119\*
  - and body surface area, 120, 121
  - at hatching, 82
  - various races and nationalities, 93, 94
- Boiling point, fatty acids, 370-378
- Bollworm (see *Pectinophora*)
- Bombyx mori* (silkworm), LEPIDOPTERA
- chromosome linkage groups, 31, 32
  - chromosome number, 4, 31
  - diapause, 419
  - heart rate, 237
  - hemolymph volume, 266
  - metamorphosis, 67
  - mutations, 31, 32
  - parasite, 499
  - propagation, 67
- Bone(s)
- effect of hormones on, 293, 297, 301
  - with increasing weight and age, 119\*
  - parasites, 518, 520
  - permissible radiation exposure for, 457
  - permissible radionuclide concentration in, 459-463, 465, 466
  - radiation effect on, 468, 469, 471-474
  - water content, 401-403
- Bone marrow
- cell chemical constituents, 398, 399
  - cells, staining method for, 552
  - differential cell counts, 275, 276
  - effect of hormones on, 297, 299
  - radiation effect on, 469, 470, 473, 474
- Bone matrix, staining method for, 551
- Boomslang (see *Dispholidus*)
- Boophilus* (tick), ACARI, 492
- B. annulatus* (cattle tick), 477
- Bordetella*, Brucellaceae, 323, 504
- Borer
- apple tree (see *Chrysobothris*, *Saperda*)
  - corn (see *Pyrausta*)
  - currant (see *Ramosia*)
  - peach tree (see *Sammnoidea*)
  - strawberry (see *Tyloclerma*)
- Borrelia*, Treponemataceae, 320, 322, 504
- Bos taurus* (cattle), ARTIODACTYLA (see also Cattle)
- acid-base, blood, 259
  - arterial blood pressure, 239
  - blood volumes, 264, 268
  - body surface area constants, 121
  - body weight, 97, 121
  - breeds, body weight, 97
  - chromosome number, 1
  - digestive enzymes, 139
  - erythrocyte values, 268
  - fatty acid source, 380

- heart, anatomy, 142
- heart rate, 234
- hemoglobin values, 268
- leukocyte counts, 273
- life span, 106
- lung ventilation, 220
- oxygen consumption, 221
- platelet count, 271
- propagation, 57
- skeletal system, 158, 160, 162
- Bothrops atrox* (fer-de-lance), SERPENTES, 328
- B. jararaca* (jararaca), 328
- Botryosphaeria*, Botryosphaeriaceae, 510, 514
- Botrytis*, Moniliaceae, 508-510
- Bouteloua gracilis* (blue grama), Gramineae, 453
- Bovicola bovis* (cattle-biting louse), MALLOPHAGA, 478
- Bovista*, Lycoperdaceae, 426
- Bowfin (*see Amia*)
- Boxfish (*see Chilomycteris*)
- Brain, 122\*
  - cell chemical constituents, 398
  - fatty acid source, 375
  - nerve connections, 130\*
  - parasites, 488, 489, 518, 520
  - regions and functions, 123, 124\*, 125\*
  - thalamic nuclei, 125, 126
  - tissue growth, 47
  - tracts, 127-129
  - water content, 401-403
- Branchiostoma* (amphioxus), CEPHALOCHORDATA††
  - chromosome number, 3
  - life span, 109
  - oxygen consumption, 223
  - propagation, 69
- Brassica campestris* (bird rape), Cruciferae, 115, 380
- B. hirta* (white mustard), 380
- B. napus* (winter rape), 375
- B. rapa* (turnip), 428
- Brazil nut, 388
- Breeding seasons
  - amphibians, 63
  - angiosperms, 72
  - aquatic invertebrates, 66
  - mammals, 57
  - reptiles, 62
- Bremia*, Peronosporaceae, 509
- Brevoortia tyrannus* (Atlantic menhaden), ISOSPONDYLI, 380
- Brine shrimp (*see Artemia*)
- Bristle worm (*see Eurythroë*)
- Brittle star (*see Ophioderma, Ophiopholis*)
- Broccoli, 494
- Brodmann, areas of, 122\*, 124
- Bromus* (bromegrass), Gramineae, 428
- Bronchus(i), parasites, 491, 518, 520
- Brood size
  - amphibians, 63
  - reptiles, 62
- Brown snake (*see Demansia*)
- Brucella*, Brucellaceae
  - antibiotic activity against, 321, 323
  - parasitism, 504
  - temperature for growth, 438
  - thermal death time, 439
- Bruchus pisorum* (pea weevil), COLFOPTERA, 421
- Brugia malayi* (Malayan filarial worm), SPIRURIDA, 486
- Bryonia alba* (white bryony), Cucurbitaceae, 368
- Bryophytes (*see specific genus*)
- Buccinum* (whelk), STENOGLOSSA, 70
- Buckwheat, 363 (*see also Fagopyrum*)
- Budworm (*see Choristoneura*)
- Buffer base, blood, 259, 262\*, 263
- Buffer solutions, pH, 543, 544
- Bufo* (toad), SALIENTIA, 235
  - B. alvarius* (Colorado River toad), 334
  - B. americanus* (American toad)
    - chromosome number, 2
    - life span, 107
    - propagation, 63
    - toxins, 334
  - B. arenarum* (sand toad), 2, 107, 334
  - B. bufo* (European toad), 334
  - B. formosus* (Japanese toad), 334
  - B. gargarizans* (Cantor's toad), 334
  - B. marinus* (marine toad), 334
  - B. quercicus* (oak toad), 334
  - B. regularis* (leopard toad), 336
  - B. terrestris* (southern toad), 240
  - B. valliceps* (Mexican toad), 103, 336
  - B. viridis* (green toad), 336
  - B. woodhousii fowleri* (Fowler's toad), 334
- Bug (*see specific genus*)
- Bugula*, CHEILOSTOMATA, 70
- Bulbs
  - disease affecting, 500
  - mineral content, 405, 411
  - parasites, 481, 483, 495
- Bull snake (*see Pitnophis*)
- Bullhead (*see also Ictalurus*)
  - anesthetic for, 548
- Bungarus candidus caeruleus* (Indian krait), SERPENTES, 330
- Burdock (*see Arctium*)
- Bushmaster (*see Lachesis*)
- Busycon canaliculatum* (whelk), STENOGLOSSA, 66
- Cabbage, parasites
  - arthropod, 483-485
  - nematode, 494-496
  - viral, 426
- Cabbage bug (*see Murgantia*)
- Cabbageworm (*see Pieris*)
- Cacao (*see Theobroma*)
- Cadelle (*see Tenebroides*)
- Caeoma*, Melampsoraceae, 511
- Calendra maidis* (maize billbug), COLEOPTERA, 421
- Caliciopsis*, Coryneliaceae, 513
- Callinectes sapidus* (blue crab), DECAPODA, 66, 109, 237
- Calliphora* (blowfly), DIPTERA, 237
- C. erythrocephala* (bluebottle fly), 4
- Callus, tissue culture
  - growth rate, 116
  - medium for, 538
- Calothrix*, Rivulariaceae, 217
- Calotropis procera* (faftan calotrope), Asclepiadaceae, 379
- Calypte anna* (Anna's hummingbird), APODIFORMES, 418
- Calyptospora*, Melampsoraceae, 511
- Camas, death (*see Ziganenus*)
- Cambarus* (crayfish), DECAPODA, 69
- Cambium
  - cell description, 44
  - cell division, 53
  - mineral content, 409
  - osmotic potential, 455
  - tissue culture, 116
- Camel, 490, 491
- Camelus bactrianus* (Bactrian camel), ARTIODACTYLA
  - chromosome number, 1
  - heart rate, 234
  - life span, 106
  - propagation, 57

†† Subphylum

- Camelus dromedarius* (Arabian camel), 264
- Camnula pellucida* (clear-winged grasshopper), ORTHOPTERA, 421
- Campion (*see* *Lychnis*)
- Canary (*see* *Serinus*)
- Cancer irroratus* (edible crab), DECAPODA, 241
- Candida*, Cryptococcaceae  
antibiotic activity against, 319-321  
parasitism, 518, 519  
respiration rate, 226  
temperature for growth, 440  
thermal death time, 440
- Canis familiaris* (dog), CARNIVORA (*see also* Dog)  
acid-base, blood, 260  
arterial blood pressure, 239  
blood volumes, 264, 268  
body surface area constants, 121  
body weight, 97, 98, 121  
breeds, body weight, 97, 98  
chromosome number, 1  
clotting time, 326, 327  
digestive enzymes, 140  
erythrocyte values, 268  
heart, anatomy, 142  
heart rate, 234  
hemoglobin values, 268  
life span, 106  
leukocyte counts, 274  
lung ventilation, 220  
oxygen consumption, 221  
platelet count, 271  
propagation, 57  
skeletal system, 159, 161, 163
- Cankerworm (*see* *Paleacrita*)
- Canna, 496
- Cannabis sativa* (hemp), Moraceae, 345
- Canthigaster* (sharp-nosed puffer), PLECTOGNATHI, 339
- Cape berry, 382
- Capillary(ies), effect of hormones on, 294-299
- Capillary blood pressure, 241-243
- Capitophorus fragae-folii* (strawberry aphid), HOMOPTERA, 501
- Capra hircus* (goat), ARTIODACTYLA (*see also* Goat)  
arterial blood pressure, 239  
blood volumes, 264, 268  
body surface area constants, 121  
body weight, 98, 99, 121  
breeds, body weight, 98, 99  
chromosome number, 1  
digestive enzymes, 139  
erythrocyte and hemoglobin values, 268  
heart rate, 234  
leukocyte counts, 274  
life span, 106  
lung ventilation, 220  
platelet count, 271  
propagation, 57
- Capsicum* (pepper), Solanaceae, 75
- C. frutescens* (bush red pepper)  
breeding system, 72  
chromosome number, 9  
light and temperature for flowering, 446  
mineral content, 412  
parasites, 506, 509  
respiration rate, 231  
seed life span, 111, 113  
soil pH, 442  
transpiration rates, 451
- Caranx hippos* (jack), PERCOMORPHI, 336, 338
- Carassius auratus* (goldfish), OSTARIOPHYSI (*see also* Goldfish)  
body weight, 104  
chromosome number, 2  
heart rate, 236  
life span, 107  
oxygen consumption, 222  
propagation, 64
- Carbohydrates (*see also* table of contents, page viii)  
aerobic oxidation, 204\*  
digestion, 184\*  
metabolic interrelationships, 203\*  
metabolism, 197\*, 198\*  
nutritional requirement for, 177  
properties, 351-366  
in urine, 187
- Carbon  
in enzymes, 288, 290  
nutritional requirement for, 165, 177  
production, plants: various regions, 216
- Carbon dioxide  
in amino acid metabolism, 199, 200  
in biosynthesis, 208\*, 209\*  
in carbohydrate metabolism, 198\*, 203\*  
and growth stimulation, 175  
in Krebs cycle, 205Fn.  
in lipid metabolism, 197\*, 203\*  
in nucleoprotein catabolism, 201\*  
nutritional requirement for, 177  
in protein metabolism, 201Fn., 203\*  
in purine and pyrimidine catabolism, 201Fn., 202\*
- Carbon dioxide content  
blood, 259-261  
respiratory media, 219
- Carbon dioxide fixation, in photosynthesis, 212, 214
- Carbon dioxide pressure, blood, 259-263
- Carbon dioxide production  
during hibernation, 417, 418  
by plants, 225-232
- Carbon dioxide reduction, in photosynthesis, 211\*
- Carbon monoxide, nutritional requirement for, 177
- Carboxydomonas*, Methanomonadaceae, 177
- Carcharhinus* (requiem shark), SELACHII, 157
- Carcinus maenas* (shore crab), DECAPODA, 223
- Cardiac reflexes, brain involvement, 123
- Cardiac sphincter, nerve connections, 132
- Cardium edule* (cockle), EULAMELLIBRANCHIA, 340
- Caretta* (loggerhead turtle), CHELONIA, 235
- C. caretta* (loggerhead turtle), 1, 62, 107
- Carotene, histochemical test for, 559
- Carp, 403 (*see also* *Cyprinus*)
- Carpocapsa pomonella* (codling moth), LEPIDOPTERA, 421, 482
- Carrot, 484, 494 (*see also* *Daucus*)
- Carthamus tinctorius* (safflower), Compositae, 380
- Carya*, Juglandaceae, 72, 73
- C. illinoensis* (pecan) (*see also* Pecan)  
first flowering, 111  
life span, 111  
measurements, 111  
mineral content, 406, 412  
parasites, 509  
pollen life span, 114  
propagation method, 73  
seed germination, 77  
shade tolerance, 443
- C. ovata* (shagbark hickory), 442
- C. tomentosa* (mockernut hickory), 9
- Carybdea alta* (sea wasp), HYMENOPTERA, 341
- Cascabel (*see* *Crotalus*)
- Cassava (*see* *Manihot*)
- Cassowary (*see* *Casuarius*)
- Castanea sativa* (European chestnut), Fagaceae, 455



- Castor bean (*see Ricinus*)
- Casuarium* (cassowary), CASUARIIFORMES, 148
- Cat (*see also Felis*)
- alimentary canal, tissue growth, 46
  - anesthetic for, 547
  - cells, staining method for, 551
  - granulocytes, growth, 47
  - heart, effect of toad toxins on, 334-337
  - lymphocytes, entry into circulation, 47
  - nerve fiber regeneration, 47
  - parasites
    - arthropod, 478, 479
    - bacterial, 505
    - fungal, 516, 520
    - helminthic, 488, 491, 493
    - protozoan, 493
    - rickettsial, 503
  - parathyroid, 154
  - platelet life span, 47
  - water content, tissues and organs, 401, 402
- Catabolism
- nucleoprotein, 201\*
  - purine and pyrimidine, 202\*
- Catalpa* (catalpa), BIGNONIACEAE, 73, 442
- C. bignonioides* (southern catalpa), 213, 230, 443
- C. ovata* (Chinese catalpa), 375
- C. speciosa* (northern catalpa)
- breeding system, 72
  - chromosome number, 9
  - first flowering, 111
  - life span, 111
  - measurements, 111
  - mineral content, 406, 412
  - seed germination, 77
- Caterpillar, forest tent (*see Malacosoma*)
- Catfish
- freshwater (*see Ictalurus*)
  - sea (*see Bagre, Clarias, Galeichthys, Plotosus*)
- Catfish sting, 336
- Cations, metabolism, 192\*, 193, 194
- Catocala* (moth), LEPIDOPTERA, 421
- Cattail (*see Typha*)
- Cattle (*see also Bos*)
- cell chemical constituents, 398
  - effect of plant toxins on, 344-349
  - enzyme source, 388, 389
  - fatty acid source, 373, 375, 377, 380
  - hormone source, 389
  - parasites
    - arthropod, 477-480
    - bacterial, 504-506
    - fungal, 516, 518, 520
    - helminthic, 486, 490, 491
    - protozoan, 493
    - rickettsial, 503
    - viral, 498-500
  - protein source, 388, 389
  - sterol source, 385
  - water content, tissues and organs, 402
- Cauliflower, 494, 496
- Cavia* (guinea pig), RODENTIA (*see also* Guinea pig)
- acid-base, blood, 260
  - blood volumes, 264
  - body surface area constants, 121
  - body weight, 121
  - leukocyte counts, 274
  - oxygen consumption, 221
- C. porcellus* (guinea pig)
- arterial blood pressure, 239
  - blood volumes, 268
  - body weight, 94, 95
  - capillary blood pressure, 242
  - chromosome linkage groups, 13
  - chromosome number, 1, 13
  - digestive enzymes, 140
  - erythrocyte values, 268
  - heart rate, 234
  - hemoglobin values, 268
  - life span, 106
  - lung ventilation, 220
  - mutations, 13
  - platelet count, 271
  - propagation, 57
  - strains, body weight, 94, 95
- C. tschudi pallidor* (guinea pig), 159, 161, 163
- Cecal worm (*see Heterakis*)
- Cecum
- and colon: enzymes, 139-141
  - parasites, 487, 490-493, 518
  - water content, 402
- Cedar, 426, 485 (*see also Cedrus*)
- red (*see Juniperus*)
  - white (*see Thuja*)
- Cedrus atlantica* (atlas cedar), PINACEAE, 428
- C. libani* (cedar of Lebanon), 428
- Celery
- glycoside source, 368
  - parasites
    - arthropod, 482, 484
    - nematode, 494, 496
    - viral, 426
  - wild (*see Vallisneria*)
- Cell(s)
- blood, *facing page* 276\*
  - chemical constituents, 398-400
  - elongation, effect of acids on, 307, 308
  - osmotic potential, 454
  - sap composition, 404
  - staining methods for, 551, 552
  - types, seed plants, 44-46
- Cell division, 51, 53, 54
- Celluloses, histochemical test for, 558
- Centipede (*see Lithobius*)
- Central nervous system
- parasites, 489, 498-500, 520
  - radiation effect on, 469, 471
- Centropogon australis* (waspfish), SCLEROPAREI, 338
- Centruroides* (scorpion), SCORPIONES, 69
- Cephalopholis argus* (blue-spotted argus), PERCOMORPHI, 336, 338
- Cephalosporium*, MONILIACEAE
- antibiotic source, 314
  - parasitism, 509, 511, 518
- Cephus cinctus* (wheat stem sawfly), HYMENOPTERA, 419
- C. pygmaeus* (European wheat stem sawfly), 482
- Ceratitis capitata* (Mediterranean fruit fly), DIPTERA, 482
- Ceratophyllus fasciatus* (European rat flea), SIPHONAPTERA, 419
- Ceratostomella*, OPHIOSTOMATACEAE, 427, 514
- Cercospora*, DEMATIACEAE, 509, 510
- Cereals, arthropod pests, 481, 483-485
- Cerebellum, regions, 123
- Cerebral cortex, areas and functions, 122\*, 123, 124\*, 125\*
- Cerebratulus* (ribbon worm), HETERONEMERTINA, 70
- Cerebrosides
- fatty acid source, 373, 377
  - properties, 384
- Cerebrospinal fluid, parasites, 499
- Ceroplastes rubens* (red wax scale), HEMIPTERA, 373
- Chalaropsis*, DEMATIACEAE, 514
- Chalcid (*see Melittobia*)



- Chalcides ocellatus* (sand skink), SAURIA, 107  
*Chalcodermus aeneus* (cowpea curculio), COLEOPTERA, 421  
 "Chameleon" (see *Anolis*)  
*Chara foetida* (stonewort), Characeae, 448  
 Chaulmoogra tree (see *Hydnocarpus*)  
*Chelonia mydas* (green sea turtle), CHELONIA, 339  
*Chelydra serpentina* (snapping turtle), CHELONIA, 62, 107 (see also Turtle)  
 Chemical elements (see also Minerals)  
   atomic weights, 579  
   in enzymes, 288, 290  
*Chenopodium album* (lamb's-quarters), Chenopodiaceae, 453  
*C. quinoa* (quinoa), 436  
 Cherry, 482 (see also *Prunus*)  
 Chestnut, 426 (see also *Castanea*)  
   Chinese water, 494  
 Chick(en) (see also *Gallus*)  
   cell growth, 47  
   developmental stages, 82, 88, 89  
   incubation time, 82  
   nutrition, 168Fn., 173Fn.  
   parasites  
     arthropod, 477-479  
     bacterial, 504, 505  
     helminthic, 491, 493  
     protozoan, 493  
     viral, 498, 499  
   protein source, 388, 389  
   water content, tissues and organs, 403  
 Chicken pox virus, 498  
 Chicory (see *Cichorium*)  
 Chigger (see *Eutrombicula*)  
*Chilomonas*, CRYPTOMONADINA  
   cell division frequency, 51  
   culture medium for, 528  
   nutrition, 177Fn.  
   oxygen consumption, 224  
*Chilomycterus spinosus* (spiny boxfish), PLECTOGNATHI, 339  
*Chilopsis linearis* (desert willow), BIGNONIACEAE, 375  
*Chimaera* (ratfish), HOLOCEPHALI†, 157  
 Chimpanzee, 486 (see also *Pan*)  
 Chinaberry (see *Melia*)  
 Chinaman fish (see *Paradicichthys*)  
 Chinch bug (see *Blissus*)  
 Chinchilla, 516  
*Chionaspis furfura* (scurfy scale), HEMIPTERA, 421  
 Chipmunk (see also *Eutamias*, *Tamias*)  
   anesthetic for, 547  
*Chiropsalmus quadrigatus* (sea wasp), CUBOMEDUSAE, 341  
 Chitin, histochemical test for, 558  
 Chiton (see *Ischnochiton*)  
*Chlamydia*, Chlamydiaceae, 503  
*Chlamydomonas*, PHYTOMONADINA or Chlamydomonadaceae  
   chromosome linkage groups, 37-39  
   chromosome number, 7, 37  
   culture medium for, 528  
   mutations, 37-39  
   temperature tolerances, 441  
*Chlorella*, Chlorellaceae  
   culture medium for, 537  
   nutrition, 168Fn., 178Fn., 179Fn., 181Fn.  
   photosynthesis, 214, 215  
   respiration rates, 227  
   temperature tolerances, 441  
 Chlorinity, sea water, 540  
*Chlorobacterium*, Chlorobacteriaceae, 217  
*Chlorogonium*, PHYTOMONADINA, culture medium for, 528  
 Chlorophyll biosynthesis, 210\*  
 Cholesterol, histochemical test for, 557  
*Choridactylus multibarbis* (stonefish), SCLEROPAREI, 338  
*Choristoneura fumiferana* (spruce budworm), LEPIDOPTERA, 482  
 Choroideremia, 11  
*Chromatium*, Thiorhodaceae, 217  
 Chromosome linkage groups (see table of contents, page xi)  
 Chromosome numbers (see table of contents, page v)  
 Chromosomes, squash: staining method for, 554  
*Chroococcus*, Chroococcaceae, 441  
*Chrysanthemum* (chrysanthemum), Compositae  
   light exposures for development, 444  
   parasites, 494, 506  
   propagation methods, 73  
   shade tolerance, 443  
*C. alpinum* (alpine chrysanthemum), 213  
*C. frutescens* (marguerite chrysanthemum), 116  
*C. leucanthemum* (oxeye daisy), 112  
*C. maximum* (Pyrenees chrysanthemum), 9, 75, 446  
*C. morifolium* (florist's chrysanthemum), 228, 442  
*C. segetum* (corn chrysanthemum), 406  
*Chrysemys* (painted turtle), CHELONIA, 154  
*C. marginata* (painted turtle), 1  
*Chrysobothris femorata* (flatheaded apple tree borer), COLEOPTERA, 482  
*Chrysomya chloropyga* (Old World screwworm), DIPTERA, 480Fn.  
*Chrysomyxa*, Melampsoraceae, 513  
*Chrysops* (mango fly), DIPTERA, 486  
*C. discalis* (deerfly), 478  
*Ciborinia*, Helotiaceae, 513  
 Cicada (see *Magicicada*)  
*Cicadulina* (leafhopper), HOMOPTERA, 502  
*Cicer arietinum* (garbanzo), Leguminosae, 216  
*Cichorium intybus* (chicory), Compositae, 116  
*Cicuta* (water hemlock), Umbelliferae, 345  
 Ciguatera poisoning, 336-338  
 Ciliate (see specific genus)  
*Cimex lectularius* (bedbug), HEMIPTERA, 4, 109, 478  
*Cinchona* (cinchona), Rubiaceae, 72, 386  
*C. ledgeriana* (ledger-bark cinchona), 9, 114, 406  
*Ciona* (sea squirt), ENTEROGONA  
   chromosome number, 3  
   heart rate, 237  
   life span, 109  
   propagation, 69  
 Circulation (see table of contents, page vii)  
 Circulatory system, comparative anatomy, 142-151  
*Circulifer tenellus* (beet leafhopper), HOMOPTERA, 421, 424, 500  
*Citellus citellus* (suslik), RODENTIA, 417  
*C. tridecemlineatus* (thirteen-lined ground squirrel), 417  
*C. undulatus* (Arctic ground squirrel), 417  
*C. undulatus parryi* (Parry's Arctic ground squirrel), 221  
*Citrullus vulgaris* (watermelon), Cucurbitaceae, 428  
*Citrus*, Rutaceae  
   breeding system, 72  
   glycoside source, 368  
   parasites  
   arthropod, 482, 484  
   bacterial, 506  
   nematode, 494, 496  
   viral, 426  
   pollen life span, 114  
   propagation method, 73  
*C. limon* (lemon)  
   chromosome number, 9  
   mineral content, 406, 412  
   osmotic potential, 453  
   photosynthesis, 213, 215

† Subclass

- respiration rates, 230, 231  
seed life span, 113  
soil pH, 442  
transpiration rates, 451
- C. paradisi* (grapefruit), 451
- C. sinensis* (sweet orange)  
chromosome number, 9  
mineral content, 406, 412  
photosynthesis, 215  
respiration rates, 230, 232  
soil pH, 442
- Cladius isomerus* (rose slug), HYMENOPTERA, 482
- Cladonia*, Cladoniaceae, 6, 227
- Cladophora*, Cladophoraceae  
chromosome number, 7  
photosynthesis, 215  
respiration rate, 227  
temperature tolerances, 441
- Cladosporium*, Dematiaceae  
antibiotic activity against, 319  
parasitism, 509, 516  
respiration rate, 226  
temperature for growth, 440  
thermal death time, 440
- Clam  
bar (*see Mactra*)  
butter (*see Saxidomus*)  
gaper (*see Schizothaerus*)  
razor (*see Ensis*)  
soft shell (*see Mya*)
- Clam worm (*see Nereis*)
- Clarias batrachus* (catfish), OSTARIOPHYSI, 336
- Classification, taxonomic (*see* Taxonomic classification)
- Claviceps purpurea* (ergot claviceps), Clavicipitaceae, 345
- Clearing agents, 551
- Clitocybe*, Agaricaceae, 509, 510
- Clitoris, nerve connections, 130\*, 133, 136
- Cloak anemone (*see Adamsia*)
- Clonorchis sinensis* (Chinese liver fluke), DIGENEA, 488
- Clostridium*, Bacillaceae  
antibiotic activity against, 320, 322  
generation time, 51  
in nitrogen cycle, 218\*  
nitrogen fixation, 217  
nutrition, 171Fn., 178Fn., 179Fn.  
parasitism, 504  
temperature for growth, 438  
thermal death time, 439  
toxin source, 388
- Clotting time, effect of anticoagulants on, 325-327
- Clove, 496
- Clover, 168Fn., 494, 502 (*see also* *Medicago*, *Melilotus*, *Trifolium*)
- Clupanodon thrissa* (gizzard shad), ISOSPONDYLI, 336, 338
- Clupea harengus* (Atlantic herring), ISOSPONDYLI, 64, 107, 380
- C. pallasii* (Pacific herring), 104
- Clutch sizes, 60, 62
- Clymenella torquata* (bamboo worm), POLYCHAETA†, 54
- CMRL 1066 culture medium for animal tissue, 531
- Coachwhip snake, anesthetic for, 548
- Coagulation, blood (*see also* table of contents, page vii)  
clotting time, 325-327  
effect of reptile toxins on 328-332
- Cobra, 388 (*see also* *Naja*, *Ophiophagus*)
- Coccidioides*, Mucoraceae, 319, 321, 518
- Coccodopsis*, Dothideaceae, 512
- Coccus hesperidum* (brown scale), HOMOPTERA, 482
- Cochliobolus*, Pseudosphaeriaceae, 510, 511
- Cochliomyia hominivorax* (screwworm), DIPTERA, 67, 421, 478
- C. macellaria* (secondary screwworm), 421, 478Fn.
- Cockle (*see* *Cardium*)
- Cocklebur (*see* *Xanthium*)
- Cockroach (*see* *Blatta*, *Blattella*, *Periplaneta*)
- Coconut, 494
- Cocos nucifera* (coconut), Palmae, 380
- Cod, 400 (*see also* *Gadus*)
- Coefficients (*see* specific coefficient)
- Coelacanth (*see* *Latimeria*)
- Colchicum autumnale* (autumn crocus), Liliaceae, 345
- Cold virus, 431
- Coleosporium*, Melampsoraceae, 513
- Colesiota*, Chlamydiaceae, 503
- Colinus* (quail), GALLIFORMES, 59 (*see also* Quail)
- C. virginianus* (bobwhite quail), 60, 101
- Colladonus citellarius* (saddled leafhopper), HOMOPTERA, 502
- Collenchyma, cell description, 45
- Colletotrichum*, Melanconiaceae, 509, 510
- Collettsia*, Chlamydiaceae, 503
- Colomesus psittacus* (freshwater puffer), PLECTOGNATHI, 339
- Colon  
and cecum: enzymes, 139-141  
nerve connections, 130\*, 132, 135  
parasites, 487  
water content, 402
- Color blindness, 11
- Coluber constrictor* (American black snake), SERPENTES, 62, 107
- Columba* (pigeon), COLUMBIFORMES, 59, 222, 235 (*see also* Pigeon)
- C. leucocephala* (white-crowned pigeon), 106
- C. livia* (street pigeon)  
arterial blood pressure, 240  
blood volumes, 265, 269  
chromosome number, 1  
erythrocyte and hemoglobin values, 269  
life span, 106  
lung ventilation, 220
- Comatricha*, Stemonitaceae, 6
- Comb jelly (*see* *Beroe*)
- Cone (*see* *Conus*)
- Cone-nose bug (*see* *Triatoma*)
- Cone sting, 340
- Congo eel, anesthetic for, 548
- Coniophora*, Thelephoraceae, 514
- Conium maculatum* (poison hemlock), Umbelliferae, 345
- Conotrachelus nemphar* (plum curculio), COLEOPTERA, 421, 482
- Constants (*see* specific constant)
- Conus* (cone), STENOGLOSSA, 340
- Conversion factors, 572-578
- Conversion formulas, 571
- Convolvulus sepium* (hedge glorybind), Convolvulaceae, 444
- Copperhead (*see also* *Ancistrodon*, *Denisonia*)  
anesthetic for, 548
- Coral, stinging (*see* *Millepora*)
- Coral plant (*see* *Jatropha*)
- Coral snake (*see* *Micrurus*)
- Coregonus albula* (European lake whitefish), ISOSPONDYLI, 2
- C. clupeaformis* (North American lake whitefish), 64, 104, 107
- Corn (*see also* *Zea*)  
parasites  
arthropod, 482-484  
fungal, 427  
nematode, 494, 496  
photosynthesis, 216  
protein source, 389
- Corn cockle (*see* *Agrostemma*)
- Cornus* (dogwood), Cornaceae, 73

† Class

- Cornus florida* (flowering dogwood)  
 chromosome number, 9  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 mineral content, 406, 412  
 osmotic potential, 453  
 photosynthesis, 215  
 shade tolerance, 443  
 soil pH, 442
- C. mas* (cornelian cherry dogwood), 114, 406
- Corpus luteum, effect of hormones on, 293, 301
- Corpus striatum, regions and functions, 123
- Cortex, plant: osmotic potential, 455
- Corticium*, Thelephoraceae, 510, 511
- Corvus*, PASSERIFORMES, 59
- C. brachyrhynchos* (American crow), 106
- C. corax* (raven), 106, 222
- C. cornix* (hooded crow), 235, 240
- Corynebacterium*, Corynebacteriaceae  
 antibiotic activity against, 322  
 generation time, 51  
 nutrition, 176Fn.  
 parasitism, 504-508  
 respiration rate, 225  
 temperature for growth, 438  
 thermal death time, 439  
 toxin source, 388
- Corynespora*, Dematiaceae, 509
- Coryneum*, Melanconiaceae, 510
- Corythuca arcuata* (oak lace bug), HEMIPTERA, 482
- Cotinis* (beetle), COLEOPTERA, 490
- Cotton (*see also Gossypium*)  
 parasites  
   arthropod, 481-485  
   nematode, 494, 496  
   viral, 501  
   sterol source, 386  
   wax source, 382
- Cotton leaf curl virus, parasitism, 501
- Cowdria*, Rickettsiaceae, 503
- Cowpea, parasites  
 arthropod, 481, 483  
 nematode, 494, 496
- Cowpox virus, 498
- Coxiella*, Rickettsiaceae, 438, 439, 503
- Coxsackie virus, 431, 498
- Coyote, 491, 493, 503
- CPLM medium for axenic culture, 524
- Crab, 488  
 blue (*see Callinectes*)  
 edible (*see Cancer*)  
 king (*see Limulus, Tachypleus*)  
 river (*see Potamon*)  
 shore (*see Carcinus*)
- Cranberry, 368, 382
- Crappie (*see Pomoxis*)
- Crassostrea virginica* (eastern oyster), EULAMELLI-BRANCHIA, 66
- Crayfish, 488 (*see also Astacus, Cambarus, Orconectes*)
- Cricket (*see Anabrus, Gryllus*)
- Criconemoides* (ring nematode), TYLENCHIDA, 494
- Cristulariella*, Moniliaceae, 512
- Crithidia*, PROTOMONADINA, culture medium for, 526
- Crocus, 368 (*see also Colchicum*)
- Cronartium*, Melampsoraceae, 427, 513
- Crotalus adamanteus* (eastern diamondback rattlesnake), SERPENTES, 328
- C. atrox* (western diamondback rattlesnake), 222, 328
- C. durissus terrificus* (cascabel), 328
- C. viridis* (prairie rattlesnake), 62, 107, 328
- C. viridis lutosus* (Great Basin rattlesnake), 102
- Croton* (croton), Euphorbiaceae, 373
- C. tiglium* (purging croton), 346, 379
- Crow (*see Corvus*)
- Cryptobranchus alleganiensis* (hellbender), CAUDATA  
 blood volumes, 269  
 chromosome number, 2  
 erythrocyte and hemoglobin values, 269  
 life span, 107  
 propagation, 63  
 segmental arteries, 147
- Cryptococcus*, Cryptococcaceae, 319, 321, 520
- Cryptosporella*, Hyalosporae, 511
- Ctenocephalides* (flea), SIPHONAPTERA, 490
- C. canis* (dog flea), 4, 479 (*see also* Flea, cat and dog)
- C. felis* (cat flea), 67, 479 (*see also* Flea, cat and dog)
- Cuclotogaster heterographus* (chicken head louse), MALLOPHAGA, 479
- Cucumaria* (sea cucumber), DENDROCHIROTA, 69, 109
- Cucumber (*see also Cucumis*)  
 parasites  
   nematode, 496  
   viral, 426, 433, 501
- Cucumis melo* (muskmelon), Cucurbitaceae, 114
- C. sativus* (cucumber)  
 chromosome number, 9  
 light for flowering, 446  
 mineral content, 406, 412  
 parasites  
   bacterial, 507  
   viral, 432, 433, 501  
 photosynthesis, 213  
 respiration rates, 231, 232  
 seed germination, 75  
 soil pH, 442
- Cucurbita* (gourd), Cucurbitaceae, 75
- C. moschata* (cushaw), 114
- C. pepo* (pumpkin)  
 chromosome number, 9  
 mineral content, 406, 412  
 parasite, 507  
 photosynthesis, 213, 215  
 respiration rate, 228  
 soil pH, 442
- Culex* (mosquito), DIPTERA  
 dispersion, 421  
 parasite vector, 486, 490, 498
- C. quinquefasciatus* (southern house mosquito), 421, 479
- Culicoides* (punkie), DIPTERA, 421, 498
- Culture media (*see table of contents, page ix*)
- Culture medium 199 for animal tissue, 530, 531
- Cupressus* (cypress), Cupressaceae, 73, 514
- C. arizonica* (Arizona cypress), 76, 110, 512
- C. sempervirens* (Italian cypress), 8
- Curculio (*see Chalcodermus, Conotrachelus*)
- Currant, 484 (*see also Ribes*)
- Curry fish (*see Stichopus*)
- Cushaw (*see Cucurbita*)
- Cutaneous blood, acid-base, 259, 262
- Cuttings, plant propagation with, 73
- Cuttlefish (*see Sepia*)
- Cutworm (*see Peridroma*)
- Cyanea capillata* (giant jellyfish), SEMAEOSTOMAE, 341
- Cyanophyta, culture medium for, 537
- Cyclops* (cyclops), CYCLOPOIDA  
 chromosome number, 4  
 life span, 109  
 parasites, 486, 490  
 propagation, 66
- Cygnus* (swan), ANSERIFORMES, 59
- C. buccinator* (trumpeter swan), 106
- C. cygnus* (whooper swan), 1



- C. olor* (mute swan), 235  
*Cylas formicarius elegantulus* (sweet potato weevil), COLEOPTERA, 421  
*Cymadothea*, Dothideaceae, 511  
 Cypress (see *Cupressus*, *Taxodium*)  
*Cyprinus carpio* (carp), OSTARIOPHYSI (see also Carp)  
   acid-base, blood, 261  
   arterial blood pressure, 240  
   blood volumes, 270  
   body weight and length, 104  
   chromosome number, 2  
   erythrocyte values, 270  
   heart rate, 236  
   hemoglobin values, 270  
   life span, 107  
   oxygen consumption, 222  
   propagation, 64  
*Cytispora*, Phyllostictaceae, 511-514  
 Cytochrome(s)  
   in chlorophyll biosynthesis, 210\*  
   and oxidation, 204\*, 205\*  
   properties, 206, 207  
 Cytology (see table of contents, page v)
- Dactylis* (orchard grass), Gramineae, 429  
*Dactylometra quinquecirrha* (pink-fringed jellyfish), SEMAEOSTOMAE, 341  
*Daedalea*, Polyporaceae, 512, 513  
 Daisy (see *Bellis*, *Chrysanthemum*)  
*Daldinia*, Xylariaceae, 512  
 Damsel fly (see *Lestes*)  
 Dandelion (see *Taraxacum*)  
*Daphne* (daphne), Thymelaceae, 368  
*Daphnia* (water flea), CLADOCERA  
   chromosome number, 4  
   dispersion, 421  
   heart rate, 237  
   life span, 109  
   propagation, 66  
 Darner (see *Anax*)  
*Dasyatis* (stingray), BATOIDEI, 157  
*D. americana* (southern stingray), 64  
*D. centroura* (rough tail stingray), 270  
*D. dipterurus* (diamond stingray), 338  
*D. pastinaca* (stingray), 108, 338  
*Dasyfus* (armadillo), EDENTATA, 221 (see also Armadillo)  
*D. novemcinctus* (nine-banded armadillo)  
   chromosome number, 1  
   heart rate, 234  
   propagation, 57  
   skeletal system, 159, 161, 163  
*Dasyscypha*, Helotiaceae, 511, 513  
 Date palm, 494 (see also *Phoenix*)  
*Datura metel* (Hindu datura), Solanaceae, 346, 434  
*D. stramonium* (jimsonweed datura), 346  
*Daucus carota* (carrot), Umbelliferae (see also Carrot)  
   breeding system, 72  
   cell sap composition, 404  
   chromosome number, 9  
   light and temperature for flowering, 446  
   mineral content, 407, 412  
   parasites, 507, 509  
   respiration rates, 228  
   seed germination, 75  
   seed life span, 112, 113  
   soil pH, 442  
   tissue culture, 116  
 Deaf-mutism, 11  
 Deer, 491, 504  
*Delphinapterus leucas* (beluga whale), CETACEA, 234  
*Delphinium* (larkspur), Ranunculaceae, 346  
*Deltocephalus striatus* (leafhopper), HOMOPTERA, 502  
*Demansia textilis* (brown snake), SERPENTES, 330  
*Demodex canis* (dog follicle mite), ACARI, 477  
*Dendroaspis angusticeps* (eastern green mamba), SERPENTES, 330  
*Dendroctonus monticolae* (mountain pine beetle), COLEOPTERA, 421, 482  
*D. valens* (red turpentine beetle), 421  
*Dendrophoma*, Sphaerioidaceae, 509  
 Dengue virus, 431, 498  
*Denisonia superba* (Australian copperhead), SERPENTES, 330  
 Density of  
   fats, 380  
   oils, 380  
   sea water, 539, 540  
   waxes, 382  
*Dermacentor andersoni* (Rocky Mountain wood tick), ACARI, 109, 477  
*D. variabilis* (American dog tick), 477  
*Dermanyssus gallinae* (chicken mite), ACARI, 477  
*Dermatobia hominis* (human botfly), DIPTERA, 479  
*Dermatocarpon*, Dermatocarpaceae, 6  
*Dermodochelys coriacea* (leatherback sea turtle), CHELONIA, 339  
 Desert plants, carbon production, 216  
*Desulfovibrio*, Spirillaceae  
   in nitrogen cycle, 218\*  
   nitrogen fixation, 217  
   nutrition, 180Fn., 181Fn.  
 Devil's-tongue (see *Amorphophallus*)  
 Diabetes insipidus, nephrogenic, 11  
*Diabrotica duodecimpunctata* (spotted cucumber beetle), COLEOPTERA, 421  
*D. undecimpunctata* (spotted cucumber beetle), 482  
*D. vittata* (striped cucumber beetle), 421  
*Diadema setosum* (reef urchin), DIADEMATOIDA, 340  
 Diamond's medium for axenic culture, 524  
 Diapause, 419  
*Diaportha*, Diaporthaceae, 509  
*Diaptomus*, CALANOIDA, 486, 490  
 Diastolic blood pressures, 239-242  
*Dibothriocephalus latus* (fish tapeworm), PSEUDOPHYLLIDEA, 490, 491 (see also *Diphyllobothrium*)  
*Dibotryon*, Dothideaceae, 510  
*Dictyocaulus filaria* (thread lungworm of sheep), RHABDITIDA, 490  
*D. viviparus* (lungworm of cattle), 490  
*Dictyostelium*, Dictyosteliaceae, 6  
*Dicyema*, DICYEMIDA, 71  
*Didelphis* (opossum), MARSUPIALIA (see also Opossum)  
   arterial blood pressure, 239  
   blood volumes, 264  
   body surface area constants, 121  
   body weight, 121  
   subintestinal vein, 146  
*D. marsupialis virginiana* (Virginia opossum)  
   chromosome number, 1  
   life span, 106  
   heart rate, 234  
   propagation, 57  
   skeletal system, 159, 161, 163  
*Didinium* (carnivorous ciliate), GYMNOSTOMATIDA, 5, 51, 110  
*Didymosphaeria*, Phaeodidymae, 512  
 Diencephalon, regions and functions, 123  
*Dientamoeba*, RHIZOMASTIGINA, culture media for, 523  
 Diffusion coefficients for  
   enzymes, 282, 284, 286  
   gases in respiratory media, 219  
   hormones, 292



- Digestion (see table of contents, page vi)
- Digestive enzymes, 139-141
- Digitalis lanata* (Grecian foxglove), Scrophulariaceae, 346, 369
- D. purpurea* (common foxglove)
- breeding system, 72
  - chromosome number, 9
  - glycoside source, 368
  - light and temperature for flowering, 446
  - pollen life span, 114
  - shade tolerance, 443
  - toxin, 346
- Diodon holacanthus* (balloonfish), PLECTOGNATHI, 339
- Dioscorea hispida* (wild yam), Dioscoreaceae, 346
- Diphyllobothrium latum* (fish tapeworm), PSEUDOPHYLLIDEA, 110, 223, 486 (see also *Dibothriocephalus*)
- Diplocarpon*, Microthyriaceae, 509
- Diplococcus*, Lactobacillaceae
- antibiotic activity against, 320, 322, 323
  - generation time, 51
  - temperature for growth, 438
  - thermal death time, 439
- Diplopida*, Sphaeropsidaceae, 509-511
- Diprion hercyniae* (European spruce sawfly), HYMENOPTERA, 482
- Dipylidium caninum* (double-pored dog tapeworm), CYCLOPHYLLIDEA, 486, 490
- Dirofilaria immitis* (dog heartworm), SPIRURIDA, 490
- Disaccharides, digestion, 184\*
- Dispholidus typus* (boomslang), SERPENTES, 332
- Dissociation constants (see pK values)
- Dissosteira longipennis* (high plains grasshopper), ORTHOPTERA, 421
- Distemper virus, canine, 431, 498
- Ditylenchus destructor* (potato rot nematode), TYLENCHIDA, 494
- D. dipsaci* (bulb and stem nematode), 494
- Diurnal variation, effect on
- osmotic potential, 454
  - transpiration rates, 452
- DNA, histochemical test for, 558
- DNA content, cells, 398-400
- Dog (see also *Canis*)
- anesthetic for, 547
  - cell chemical constituents, 398
  - differential cell count, rib marrow, 275
  - lymphocytes, growth, 47
  - parasites
    - arthropod, 477-480
    - bacterial, 505
    - fungus, 516, 518, 520
    - helminthic, 486, 488, 490-493
    - protozoan, 488, 493
    - rickettsial, 503
    - viral, 498
  - parathyroid, 154
  - platelet replacement, 47
  - radiation effect on, 470
  - teeth, staining method for, 552
  - thyroid tissue growth, 49
  - tissue regeneration, 46, 48
  - water content, tissues and organs, 402
- Dogbane (see *Apocynum*)
- Dogfish (see *Musculus*, *Squalus*)
- Dogwood (see *Cornus*)
- Dolichodoris heterocephalus* (awl nematode), TYLENCHIDA, 494
- Dolphin, 379
- Donax serra* (white mussel), EULAMELLIBRANCHIA, 340, 341
- Doris* (sea lemon), ACOELA, 4, 109
- Dormouse (see *Glis*, *Muscardinus*)
- Dothichiza*, Sphaerioidaceae, 513
- Dothiorella*, Sphaerioidaceae, 509, 514
- Dove, 493 (see also *Zenaidura*)
- Dracunculus medinensis* (guinea worm), SPIRURIDA, 486
- Drosophila* (fruit fly), DIPTERA
- cell division, 54
  - chromosome linkage groups, 20-29
  - chromosome number, 4, 20
  - diapause, 419
  - dispersion, 421, 422, 424
  - heart rate, 237
  - life span, 109
  - metamorphosis, 67
  - mutations, 20-29
  - nutrition, 168Fn., 175Fn.
  - oxygen consumption, 223
  - parasitism, 482
  - propagation, 67
- Dryopteris spinulosa* (wood fern), Polypodiaceae, 213
- Duck, 499, 504 (see also *Anas*)
- Dugesia* (planarian), TRICLADIDA, 70
- Dulbecco's solution for animal tissue culture, 530
- Duodenum
- cell life span, 46
  - nerve connections, 138
  - parasites, 486-489
  - water content, 402
- Dysplasia, 11
- Dystrophy, 11
- Dytiscus* (diving beetle), COLEOPTERA
- chromosome number, 4
  - diapause, 419
  - heart rate, 237
  - hemolymph volume, 266
- Eagle's culture media for animal tissue, 531, 532
- Ear(s)
- parasites, 478, 518
  - radiation effect on, 468, 471
- Earle's solution for animal tissue culture, 530
- Earthworm (see *Eisenia*, *Lumbricus*)
- Earwig (see *Forficula*)
- Earworm (see *Heliothis*)
- Echinophaga gallinacea* (sticktight flea), SIPHONAPTERA, 479
- Echinarachnius* (sand dollar), CLYPEASTEROIDA, 3, 53
- Echinococcus granulosus* (hydatid tapeworm), CYCLOPHYLLIDEA, 486, 490
- Echis carinatus* (saw-scaled viper), SERPENTES, 330
- Ectocarpus*, Ectocarpaceae, 7, 227
- Ectoderm, 81\*
- Eel, 403 (see also *Anguilla*, *Electrophorus*, *Gymnothorax*)
- Eelworm (see *Anguina*)
- Efferents, thalamic nuclei, 125, 126
- Egg(s)
- cell division, 53, 54
  - fertilized: developmental stages, 82-92
  - fixatives for, 549, 550
  - protein source, 388, 389
  - staining methods for, 551
- Egg-yolk infusion for agnotobiotic culture, 524
- Eggplant, 483 (see also *Solanum*)
- Ehrlichia*, Rickettsiaceae, 503
- Eimeria*, EUCOCCIDIA, 71, 492
- Eisenia* (earthworm), OLIGOCHAETA†, 490
- Elaeis guineensis* (African oil palm), Palmae, 380
- Elder, 484 (see also *Sambucus*)
- Electrolytes, 190 (see also Minerals)
- Electrophorus electricus* (electric eel), OSTARIOPHYSI, 107
- Elements (see Chemical elements, Minerals)
- Elephant, 493
- Elephas maximus* (Asiatic elephant), PROBOSCIDEA
- heart rate, 235
  - life span, 106

† Class

- oxygen consumption, 221  
 propagation, 57  
 skeletal system, 158, 160, 162
- Elk, 504**
- Elm, 427 (see also *Ulmus*)**
- Elodea canadensis* (Canada waterweed), Hydrocharitaceae**  
 chromosome number, 8  
 mineral content, 405  
 protoplasmic streaming, 448  
 respiration rates, 229, 230
- Elsimoe*, Myriangiaceae, 509, 511**
- Elytoderma*, Phacidiaceae, 513**
- Embryo**  
 developmental stages, 82-92  
 early development, 79\*  
 germ layers, 80\*, 81\*
- Empoasca fabae* (potato leafhopper), HOMOPTERA, 422, 482**
- Emys* (turtle), CHELONIA, 148**
- E. orbicularis* (European pond turtle), 1, 235**
- Encephalomyelitis virus, equine**  
 arthropod vectors, 477-479  
 effect of temperature on, 431  
 parasitism, 498  
 size, 498
- Enchytraeus humiculator* (white worm), OLIGOCHAETA†, 4**
- Entamoeba*, AMOEBINA (see also *Entamoeba*)**  
 culture media for, 523  
 propagation, 71
- Endoconidiophora*, Hyalosporae, 509, 512, 513**
- Endocrine organs**  
 invertebrate, 304-306  
 vertebrate, 290-303  
 comparative anatomy, 152-157
- Endoderm, 80\***
- Endolimax*, AMOEBINA, culture media for, 523**
- Endoplasmic reticulum, cytochromes, 206, 207**
- Endosperm, tissue culture, 117**
- Endothia*, Diaporthaceae, 426**
- Energy (see also Cytochromes, Krebs cycle)**  
 utilization in photosynthesis, 216
- Engraulis japonica* (anchovy), ISOSPONDYLLI, 336, 338**
- Enhydrina schistosa* (beaked sea snake), SERPENTES, 330**
- Ensis directus* (razor clam), EULAMELLIBRANCHIA, 340, 341**
- Entamoeba*, AMOEBINA (see also *Entamoeba*)**  
 antibiotic activity against, 320, 323  
 chromosome number, 5  
 culture media for, 523  
 parasitism, 488
- Enderobius vermicularis* (pinworm), RHABDITIDA, 109, 486**
- Enzyme(s)**  
 amino acid content, 289, 291  
 digestive, 139-141  
 mineral content, 288, 290  
 occurrence, 277-281  
 properties, 282-287, 388  
 in urine, 188
- Enzyme activity**  
 in biosynthesis, 208\*, 209\*  
 in carbohydrate digestion, 184\*  
 in carbohydrate metabolism, 198\*  
 catalytic, 227-281  
 in cytochrome system, 205\*  
 effect of reptile toxins on, 328, 330, 332  
 in Krebs cycle, 204\*  
 in lipid digestion, 185\*  
 in photosynthesis, 211\*  
 in protein digestion, 183\*
- in purine and pyrimidine catabolism, 201\*, 202\*  
 in sucrose synthesis, 212\*
- Eosinophil(s), facing page 276\***  
 life span, 47
- Eosinophil count, 273, 274**
- Epeyithroozoon*, Bartonellaceae, 503**
- Ephemera* (mayfly), EPHEMEROPTERA, 69**
- Ephestia eludella* (tobacco moth), LEPIDOPTERA, 419**
- E. kuehniella* (Mediterranean flour moth)**  
 chromosome number, 4  
 heart rate, 237  
 metamorphosis, 68  
 parasitism, 482  
 propagation, 68
- Epibolus insidiator* (Indo-Pacific long-jawed wrasse), PERCOMORPHI, 336, 338**
- Epicauda vittata* (blister beetle), COLEOPTERA, 482**
- Epidermis, plant**  
 cell description, 44  
 osmotic potential, 454
- Epidermophyton*, Moniliaceae, 321, 516**
- Epilachna corrupta* (ladybird beetle), COLEOPTERA, 419**
- E. varivestis* (Mexican bean beetle), 483**
- Epinephelus fuscoguttatus* (mottled grouper), PERCOMORPHI, 336, 338**
- Epithalamus, functions, 123**
- Epitrix cucumeris* (potato flea beetle), COLEOPTERA, 422**
- E. hirtipennis* (tobacco flea beetle), COLEOPTERA, 424, 483**
- Eptesicus fuscus* (big brown bat), CHIROPTERA**  
 hibernation, 417  
 skeletal system, 159, 161, 163
- Equine encephalomyelitis virus (see Encephalomyelitis)**
- Equisetum arvense* (field horsetail), Equisetaceae, 8**
- E. hyemale* (scouring rush), 443**
- E. telmateia* (giant horsetail), 229**
- Equus caballus* (horse), PERISSODACTYLIA (see also Horse)**  
 acid-base, blood, 260  
 arterial blood pressure, 239, 240  
 blood volumes, 264, 268  
 body surface area constants, 121  
 body weight, 99, 121  
 chromosome number, 1  
 digestive enzymes, 139, 140  
 erythrocyte values, 268  
 heart rate, 235  
 hemoglobin values, 268  
 leukocyte counts, 274  
 life span, 106  
 lung ventilation, 220  
 oxygen consumption, 221  
 platelet count, 271  
 propagation, 57  
 skeletal system, 158, 160, 162
- Eretmochelys imbricata* (hawksbill sea turtle), CHELONIA, 339**
- Eriaceus europaeus* (European hedgehog), INSECTIVORA**  
 body surface area constants, 121  
 body weight, 121  
 chromosome number, 1  
 heart rate, 235  
 hibernation, 417  
 life span, 106  
 propagation, 57
- Eriophyes pyri* (pear leaf blister mite), ACARI, 481**
- Eriosoma lanigerum* (woolly apple aphid), HOMOPTERA, 483**
- Erwinia*, Enterobacteriaceae, 52, 438, 506-508**

† Class

- Erysiphe*, Erysiphaceae, 427, 509-512
- Erythroblast, facing page 276\*
- Erythrocyte(s), facing page 276\*
- chemical constituents, 400
  - dimensions, 267-270
  - growth, 47
  - hemoglobin content, 259-261, 267-270
  - life span, 47
  - parasites, 489, 493
  - pH, 262\*
  - radiation effect on, 470, 473-475
  - total ion content, 262\*
- Erythrocyte count, 267-270
- Erythrocyte volume, 259-261, 263-265, 267-270
- Erythroxylon coca* (Huanuco cocaine tree), Erythroxylaceae, 346
- Escherichia*, Enterobacteriaceae
- antibiotic activity against, 320-322
  - chemical constituents, 400
  - generation time, 52
  - parasitism, 505
  - respiration rate, 225
  - temperature for growth, 438
  - thermal death time, 439
- Esophagus
- enzymes, 139-141
  - nerve connections, 132, 134
  - parasites, 493
- Esox* (pike), HAPLOMI, 157
- E. lucius* (northern pike)
- arterial blood pressure, 240
  - body weight and length, 104
  - chromosome number, 2
  - heart rate, 236
  - life span, 107
  - oxygen consumption, 222
  - propagation, 64
- Estrus cycle, 57
- Euapta lappa* (sea cucumber), APODA, 340
- Eubranchipus* (fairy shrimp), ANOSTRACA, 69
- Euglena*, EUGLENOIDINA or Euglenaceae
- cell division frequency, 51
  - chromosome number, 5
  - culture medium for, 528
  - nutrition, 171 Fn., 175Fn.
- Eumeces elegans* (elegant skink), SAURIA, 1
- E. fasciatus* (five-lined skink), 62, 102, 103
- Euphorbia* (euphorbia), Euphorbiaceae, 346
- E. capitellata* (head euphorbia), 451
- Euphractus villosus* (six-banded armadillo), EDENTATA, 106
- Eurydema ornatum* (ornate vegetable bug), HEMIPTERA, 419
- Eurygaster integriceps* (soun pest), HEMIPTERA, 419
- Eurythoe complanata* (bristle worm), POLYCHAETA†, 340
- Eutamias minimus* (least chipmunk), RODENTIA, 235
- Eutrombicula alfreddugesi* (chigger), ACARI, 477
- Eutypella*, Diatrypaceae, 512
- Excretion products
- feces, 190
  - minerals, 192\*, 193-195
  - in nitrogen cycle, 218\*
  - urine, 186-188
- Exosporina*, Tuberculariaceae, 509
- Extracellular fluids, mineral retention in, 192\*, 193-195
- Extremities, bones comprising, 160-163
- Eye(s)
- nerve connections, 130\*, 131, 134
  - parasites
    - fungal, 518, 520
    - helminthic, 487
    - viral, 498
  - permissible radiation exposure for, 457
  - radiation effect on, 468-475
  - water content, 401-403
- Face, nerve connections, 130\*, 136
- Factors, blood, 253\*-258\*
- Factors, conversion, 572-578
- Fagopyrum esculentum* (buckwheat), Polygonaceae (see also Buckwheat)
- breeding system, 72
  - cell sap composition, 404
  - chromosome number, 9
  - light for flowering, 446
  - mineral content, 407, 412
  - photosynthesis, 213
  - respiration rate, 228
  - seed germination, 75
  - soil pH, 442
- Fagus* (beech), Fagaceae, 72, 73, 455
- F. grandifolia* (American beech)
- first flowering, 111
  - life span, 111
  - measurements, 111
  - parasites, 512
  - seed germination, 77
  - shade tolerance, 443
- F. sylvatica* (European beech)
- chromosome number, 9
  - mineral content, 407
  - photosynthesis, 213
  - pollen life span, 114
  - respiration rate, 230
  - soil pH, 442
- Fairy shrimp (see *Eubranchipus*)
- Fairyfly (see *Trichogramma*)
- Fascioli of the brain, 127-129
- Fasciola* (liver fluke), DIGENEA, 70
- F. hepatica* (liver fluke)
- chromosome number, 4
  - oxygen consumption, 223
  - parasitism, 488, 492
- Fasciolopsis buski* (intestinal fluke), DIGENEA, 488
- Fat(s)
- aerobic oxidation, 204\*
  - fatty acid source, 373, 379, 380
  - in feces, 190
  - metabolic interrelationships, 203\*
  - nutritional requirements for, 177
  - properties, 380, 381
- Fat, body (see Body fat)
- Fat cells, staining method for, 551
- Fatty acid(s)
- digestion, 185\*
  - metabolism, 197\*
  - nutritional requirement for, 167
  - properties, 370-379
  - sources, 370-380
- Fatty acid content
- cells, 398
  - feces, 190
- Feather star (see *Antedon*)
- Feathers, water content, 403
- Feces
- constituents, 190
  - minerals excreted in, 192\*, 193-195
  - parasites, 498
  - sterol source, 385
- Felis catus* (cat), CARNIVORA (see also Cat)
- acid-base, blood, 260
  - arterial blood pressure, 240
  - blood volumes, 264, 268
  - body surface area constants, 121

† Class



- body weight, 99, 121
- chromosome number, 1
- digestive enzymes, 140
- erythrocyte values, 268
- heart rate, 235
- hemoglobin values, 268
- leukocyte counts, 274
- life span, 106
- lung ventilation, 220
- oxygen consumption, 221
- platelet count, 271
- propagation, 57
- skeletal system, 159, 161, 163
- Fer-de-lance (*see Bothriops*)
- Fern (*see Dryopteris, Osmunda*)
- Ferret
  - anesthetic for, 548
  - ovum, fertility loss, 48
  - parasites, 478, 498
- Fetus (*see also Embryo*)
  - developmental stages, 83-87
- Fiber, plant: cell description, 45
- Fibroblasts, cell division, 53, 54
- Fibulia nolitangere* (brown sponge), HAPLOSCLERINA, 341
- Fig, parasites, 496
- Filarial worm (*see Setaria*)
  - African (*see Loa*)
  - Bancroft's (*see Wuchereria*)
  - convoluted (*see Onchocerca*)
- Filefish (*see Alutera*)
- Fir, 482 (*see also Abies*)
  - Douglas, 382 (*see also Pseudotsuga*)
- Firebrat (*see Thermobia*)
- Fish (*see specific genus*)
- Fistulina*, *Fistulinaceae*, 513
- Fixatives, 549, 550
- Flagellates, marine (*see also specific genus*)
  - culture medium for, 528
- Flannel mullein (*see Verbascum*)
- Flatworm (*see specific genus*)
- Flax, 382 (*see also Linum*)
- Flea, 503
  - cat and dog, 486 (*see also Ctenocephalides*)
  - human, 486 (*see also Pulex*)
  - rodent, 486 (*see also Ceratophyllus*)
  - sticktight (*see Echinophaga*)
- Fledging success, 61
- Flounder (*see Limanda, Pseudopleuronectes*)
- Flower(s)
  - diseases affecting, 507, 510
  - mineral content, 406, 407, 409, 413
  - respiration rates, 231
  - toxins, 344, 345, 348, 349
- Flowering
  - light and temperature for, 446, 447
  - tree age at first, 110, 111
- Fluke
  - blood (*see Schistosoma*)
  - intestinal (*see Fasciolopsis*)
  - liver (*see Clonorchis, Fasciola*)
  - lung (*see Paragonimus*)
  - "salmon poisoning" (*see Nanophyetus*)
- Fly
  - black (*see Simulium*)
  - blow- (*see Calliphora, Phaenicia, Phormia*)
  - bluebottle (*see Calliphora*)
  - bot- (*see Dermatobia, Gasterophilus, Oestrus*)
  - bulb (*see Lampetia, Merodon*)
  - deer- (*see Chrysops*)
  - flesh (*see Wohlfahrtia*)
  - frit (*see Oscinella*)
  - fruit (*see Ceratitis, Drosophila*)
  - greenbottle (*see Phaenicia*)
  - Hessian (*see Phytophaga*)
  - horn (*see Siphona*)
  - horse- (*see Tabanus*)
  - house- (*see Musca*)
  - mango (*see Chrysops*)
  - rust (*see Psila*)
  - sand (*see Phlebotomus*)
  - stable (*see Stomoxys*)
  - tachinid (*see Lydella*)
  - tsetse (*see Glossina*)
- Follicle (*see specific follicle*)
- Fomes, Polyporaceae, 511-514
- Foot-and-mouth disease virus, 431, 498
- Forest plants, carbon production, 216
- Forficula auricularia* (European earwig), DERMAPTERA, 483
- Formica* (ant), HYMENOPTERA, 223
- F. fusca* (black ant), 109
- F. sanguinea* (red ant), 4, 109 (*see also Ant, red*)
- Formulas, conversion, 571
- Fossaria (freshwater snail), BASOMMATOPHORA, 492
- Fowl (*see also specific genus*)
  - anesthetic for, 548
  - cell chemical constituents, 400
  - parasites
    - bacterial, 504, 505
    - rickettsial, 503
    - viral, 431, 498, 499
- Fox
  - anesthetic for, 548
  - parasites, 490-493, 503
- Foxglove (*see Digitalis*)
- Fragaria* (strawberry), Rosaceae (*see also Strawberry*)
  - mineral content, 412
  - parasite, 501
  - pollen life span, 114
  - propagation method, 73
  - respiration rates, 230-232
  - soil pH, 442
- F. chiloensis* (chiloe strawberry), 446, 509
- F. virginiana* (Virginia strawberry), 9, 72, 507
- Frankliniella*, THYSANOPTERA, 501
- Fraxinus* (ash), Oleaceae (*see also Ash*)
  - breeding system, 72
  - parasites, 507, 512, 514, 515
  - pollen dispersion, 429
  - propagation method, 73
- F. americana* (white ash)
  - chromosome number, 9
  - first flowering, 111
  - life span, 111
  - measurements, 111
  - mineral content, 412
  - osmotic potential, 453
  - seed germination, 77
  - seed life span, 112
  - shade tolerance, 443
- F. excelsior* (European ash), 213, 230, 407
- F. nigra* (black ash), 229
- F. pennsylvanica* (green ash), 112, 113
- Freezing point, sea water, 539
- Freshwater, respiratory medium, 219
- Frog (*see also Rana*)
  - cardiac arrest, effect of toad toxins on, 335, 337
  - embryonic development, 89, 90
  - water content, tissues and organs, 403
- clawed (*see Xenopus*)
- leopard, 548 (*see also Rana*)
- tree (*see Hyla*)



- Fruit(s)  
 diseases affecting, 501, 502, 506, 507, 510  
 mineral content, 405-409, 411-413  
 parasites, 482-485  
 respiration rates, 231, 232  
 toxins, 345, 347, 348
- Fucus serratus* (rockweed), Fucaceae, 213  
*F. vesiculosus* (rockweed), 7, 227, 441  
*Fugu* (puffer), PLECTOGNATHI, 339  
*Fundulus heteroclitus* (mummichog), MICROCYPRINI, 2, 64
- Fungi (see also specific genus)  
 tissue culture media for, 536
- Fusarium*, Tuberculariaceae  
 nutrition, 180Fn.  
 parasitism, 508-510  
 respiration rate, 226  
 temperature for growth, 440  
 thermal death time, 440
- Fusicladium*, Pseudosphaeriaceae, 513, 514  
*Fusobacterium*, Bacteroidaceae, 505
- Gadus callarias* (rock cod), ANACANTHINI, 7, 270  
*G. morhua* (Atlantic cod)  
 body weight and length, 104  
 fatty acid source, 380  
 heart rate, 236  
 life span, 107  
 propagation, 64
- Galba* (freshwater snail), BASOMMATOPHORA, 492  
*Galeichthys felis* (sea catfish), OSTARIOPHYSI, 336  
*Galium aparine* (catchweed bedstraw), Rubiaceae, 453  
 Gall, plant: tissue culture, 116, 117  
 Gallbladder  
 effect of hormones on, 303  
 nerve connections, 130\*, 132, 135
- Galleria mellonella* (greater wax moth), LEPIDOPTERA, 266
- Gallus domesticus* (chicken), GALLIFORMES (see also Chicken)  
 acid-base, blood, 260  
 arterial blood pressure, 240  
 blood volumes, 265, 269  
 body weight, 101  
 breeds, body weight, 101  
 cell division, 53, 54  
 chromosome linkage groups, 19, 20  
 chromosome number, 1, 19  
 clotting time, 327  
 digestive enzymes, 141  
 erythrocyte values, 269  
 heart rate, 235  
 hemoglobin values, 269  
 leukocyte counts, 274  
 life span, 106  
 lung ventilation, 220  
 mutations, 19, 20  
 oxygen consumption, 222
- Galumna* (grass mite), ACARI, 490  
 Ganglionic connections, 130\*, 131-138  
*Ganoderma*, Polyporaceae, 514  
 Gar (see *Lepisosteus*)  
 Garbanzo (see *Cicer*)  
 Gardenia, 368  
 Garlic, 483  
 Garter snake (see *Thamnophis*)  
 Gases in respiratory media, 219  
*Gasterophilus* (botfly), DIPTERA, 479  
*Gasterosteus* (stickleback), THORACOSTEI, 157  
 Gastrointestinal tract (see also Alimentary tract)  
 carbohydrate digestion, 184\*  
 hormones, 302, 303  
 lipid digestion, 185\*  
 parasite, 520  
 permissible radionuclide concentration in, 458-467
- Gastrula, developmental stages, 82, 84, 86, 89, 91  
 Gene(s) (see specific blood group system)  
 Gene linkage (see table of contents, page v)  
 Generation time, bacteria and viruses, 51  
 Genetic code, 43  
 Genitals (see also specific organs)  
 nerve connections, 130\*  
 parasites, 487, 493
- Genotype (see specific blood group system)  
 Geologic time divisions, 570, 571  
*Geotrichum*, Moniliaceae, 520  
 Gerbil, 488, 493 (see also *Rhombomys*)  
 Germ layers, mammals, 79\*-81\*  
 German measles virus, 499  
 Germination, seeds, 75-77  
 Gestation period, 57, 62, 82  
 Gey's solution for animal tissue culture, 530  
*Giardia*, DISTOMATINA, 488  
*Gibberella*, Nectriaceae, 509, 511  
*Gigartina harveyana* (seaweed), Gigartinaceae, 214  
 Gila monster (see *Heloderma*)  
*Gilia*, Polemoniaceae, 429  
*Ginkgo biloba* (ginkgo), Ginkgoaceae  
 chromosome number, 8  
 pollen life span, 114  
 propagation methods, 73  
 shade tolerance, 443  
 soil pH, 442
- Gladiolus* (*gladiolus*), Iridaceae  
 chromosome number, 8  
 parasites, 507  
 respiration rate, 229  
 seed life span, 113  
 soil pH, 442
- G. gandavensis* (breeder's *gladiolus*), 230, 231  
*G. hortulanus* (horticultural *gladiolus*), 73  
*G. hybrida* (hybrid *gladiolus*), 114  
*Glaucoma*, HYMENOSTOMATIDA, 168Fn.  
*Glis glis* (fat dormouse), RODENTIA, 417  
*Gloeodes*, Leptostromataceae, 510  
*Gloeosporium*, Melanconiaceae, 509, 514  
*Glomerella*, Gnomoniaceae, 509, 510  
 Glomerulus, diameter, 47  
 Glorybind (see *Convolvulus*)  
*Glossina* (tsetse fly), DIPTERA  
 dispersion, 422  
 parasite vector, 479, 488, 492
- Glycera dibranchiata* (bloodworm), POLYCHAETA†, 340  
*Glycine soja* (soybean), Leguminosae (see also Soybean)  
 breeding system, 72  
 cell sap composition, 404  
 chromosome number, 9  
 fatty acid source, 380, 381  
 light exposures for development, 443-445  
 light and temperature for flowering, 446  
 mineral content, 407, 412  
 nitrogen fixation, 217  
 parasites, 433, 507, 509  
 photosynthesis, 213  
 respiration rate, 228  
 seed germination, 76  
 seed life span, 112
- Glycogen, histochemical test for, 558  
 Glycolysis, 198\*  
 Glycosides  
 nutritional requirements for, 177  
 properties, 368-371  
 sources, 368, 370  
 uses, 369, 371

† Class

- Gnathodentex aureolineatus* (snapper), PERCOMORPHI, 337, 338
- Gnomonia*, Gnomoniaceae, 509, 513, 514
- Goat (see also *Capra*)  
parasites  
arthropod, 477, 480  
bacterial, 504-506  
fungal, 516  
helminthic, 490, 491, 493  
protozoan, 493  
ricketsial, 503  
viral, 498
- Goldenrod (see *Solidago*)
- Goldfish (see also *Carassius*)  
anesthetic for, 548
- Golgi apparatus, staining method for, 556
- Gonads (see also specific organ)  
comparative anatomy, 156, 157  
parasite, 499  
permissible radiation exposure for, 457
- Goniobasis* (North American river snail), MESOGASTROPODA, 492
- Gonionemus* (hydromedusa), LIMNOMEDUSAE, 5
- Gonyaulax* (dinoflagellate), DINOFLAGELLATA, 341, 342
- Goose, 491 (see also *Anser*)
- Gooseberry, 484
- Goosefish (see *Lophius*)
- Gordius* (horsehair worm), GORDIOIDEA, 70
- Gossypium* (cotton), Malvaceae (see also Cotton)  
breeding system, 72  
mineral content, 412  
osmotic potential, 453  
parasites, 507, 509  
seed germination, 76  
seed life span, 112, 113
- G. arboreum* (Asiatic tree cotton), 429
- G. barbadense* (Pima cotton), 114
- G. herbaceum* (Levant cotton), 228-230, 232, 451
- G. hirsutum* (upland cotton)  
chromosome number, 9  
fatty acid source, 380  
light and temperature for flowering, 446  
mineral content, 407  
parasite, 501  
photosynthesis, 213  
pollen dispersion, 429  
respiration rate, 228  
soil pH, 442  
transpiration rates, 453
- Gourd (see *Cucurbita*, *Trichosanthes*)
- Grafting, plant propagation by, 72, 73
- Grahamella*, Bartonellaceae, 503
- Grain  
arthropod pests, 481-485  
mineral content, 405, 411
- Grantia* (marine sponge), SYCONOSA, 110
- Granulocytes, growth, 47
- Grape, 484 (see also *Vitis*)
- Grapefruit (see *Citrus*)
- Grapholitha molesta* (oriental fruit moth), LEPIDOPTERA, 422
- Grass(es), 482-484, 494  
Bahia (see *Paspalum*)  
bent (see *Agrostis*)  
blue- (see *Poa*)  
blue gramma (see *Bouteloua*)  
bluestem (see *Andropogon*)  
brome- (see *Bromus*)  
June (see *Koeleria*)  
orchard (see *Dactylis*)  
rye- (see *Lolium*)  
Sudan (see *Sorghum*)  
switch (see *Panicum*)  
wheat- (see *Agropyron*)
- Grasshopper  
clear-winged (see *Cammula*)  
high plains (see *Dissosteira*)  
spur-throated (see *Melanoplus*, *Romalea*)
- Grassland, plant carbon production, 216
- Gratiola* (hedge hyssop), Scrophulariaceae, 368
- Grebe (see *Podiceps*)
- Groundsel (see *Senecio*)
- Grouper (see *Epinephelus*)
- Grouse, ruffed, 493
- Growth (see table of contents, pages v, vi)
- Grub, cattle (see *Hypoderma*)
- Gryllus* (field cricket), ORTHOPTERA, 483
- Guard cells  
description, 44  
osmotic potential, 454
- Guignardia*, Sphaeriaceae, 511
- Guinea fowl, 479
- Guinea pig (see also *Cavia*)  
cell chemical constituents, 398, 399  
effect of toad toxins on, 334-337  
ovum, fertility loss, 48  
parasites, 480, 505  
radiation effect on, 470  
thyroid, tissue growth, 49  
tissue regeneration, 46, 48  
water content, tissues and organs, 402
- Guinea worm (see *Dracunculus*)
- Guizotia abyssinica* (Ethiopian niger seed), Compositae, 380
- Gull (see *Larus*)
- Gum, sweet (see *Liquidambar*)
- Gymnodinium* (dinoflagellate), DINOFLAGELLATA, 341, 342
- Gymnosporangium*, Pucciniaceae  
parasitism, 510, 512, 513  
spore dispersion, 426
- Gymnothorax* (moray eel), APODES, 338
- Gynmura marmorata* (butterfly stingray), BATOIDEI, 338
- Gyps (vulture), FALCONIFORMES, 59
- G. fulvus* (griffon vulture), 106, 235
- Habrobracon juglandis* (parasitic wasp), HYMENOPTERA, 4, 29, 30
- Habu, Taiwan (see *Trimeresurus*)
- Haddock (see *Melanogrammus*)
- Haemagogus spegazzinii* (forest mosquito), DIPTERA, 420
- Haematopinus eurysternus* (short-nosed cattle louse), ANOPLURA, 479
- Haemobartonella*, Bartonellaceae, 503
- Haemonchus contortus* (twisted stomach worm), RHABDITIDA, 490
- Haemophilus*, Brucellaceae  
antibiotic activity against, 320-323  
culture medium for, 535  
nutrition, 171Fn.  
parasitism, 505
- Hagfish (see *Myxine*)
- Hair  
growth rate, 47  
parasites, 517, 519  
radiation effect on, 468-471, 473, 474  
water content, 401-403
- Hair follicles  
effect of hormones on, 301  
nerve connections, 130\*  
parasite, 477

- Hairworm (see *Trichostrongylus*)  
 Half-life, radionuclides, 546, 547  
 Halibut (see *Hippoglossus*)  
 Hamster, 82 (see also *Mesocricetus*)  
 Hanks' solution for animal tissue culture, 530  
 Hare, 492, 493 (see also *Lepus*)  
*Harmolita grandis* (wheat strawworm), HYMENOPTERA, 422  
*H. tritici* (wheat jointworm), 422, 483  
 Hatching success, 60, 61  
 Hawk, 82, 493  
 Hawkbit (see *Leontodon*)  
 Hawkweed (see *Hieracium*)  
 Hearing, brain involvement, 123, 124, 127  
 Heart  
   cell chemical constituents, 398, 399  
   comparative anatomy, 142, 143  
   effect of hormones on, 299  
   effect of toad toxins on, 334-337  
   nerve connections, 130\*-132, 134, 136, 138  
   parasites, 490, 491  
   radiation effect on, 469, 470, 472  
   tissue growth, 47  
   water content, 401, 402  
 Heart rate, 234-238  
   during hibernation, 417, 418  
 Heartworm (see *Dirofilaria*)  
 Hedgehog (see *Erinaceus*)  
 Height, body (see Body height)  
 Height, trees, 110, 111  
*Helianthus* (sunflower), Compositae, 429  
*H. annuus* (common sunflower)  
   breeding system, 72  
   chromosome number, 9  
   fatty acid source, 380  
   light for flowering, 446  
   mineral content, 407, 412  
   osmotic potential, 453  
   parasite, 507  
   photosynthesis, 213-215  
   respiration rates, 228-232  
   seed germination, 76  
   seed life span, 112  
   soil pH, 442  
   tissue culture, 116  
   transpiration rates, 451  
*H. rigidus* (stiff sunflower), 456  
*H. tuberosus* (Jerusalem artichoke sunflower), 116  
*Helicotylenchus* (spiral nematode), TYLENCHIDA, 494  
*Heliothis armigera* (corn earworm), LEPIDOPTERA, 68, 422  
*H. zea* (corn earworm), 483  
*Helix* (land snail), STYLOMMATOPHORA, 70  
*H. pomatia* (land snail)  
   chromosome number, 4  
   heart rate, 238  
   life span, 109  
   oxygen consumption, 223  
   propagation, 66  
   protein source, 389  
 Hellbender (see *Cryptobranchus*)  
 Hellebore, false (see *Veratrum*)  
*Helleborus* (hellebore), Ranunculaceae, 455  
*H. niger* (black Christmas rose), 346  
 Heller's solution for tissue culture, 538  
*Helminthosporium*, Dematiaceae  
   parasitism, 509, 511  
   respiration rate, 226  
   spore dispersion, 427  
   temperature for growth, 440  
*Heloderma suspectum* (Gila monster), SAURIA  
   body length, 332  
   chromosome number, 1  
   life span, 107  
   propagation, 62  
   venom, 332  
*Hemachatus haemachates* (ringhals), SERPENTES, 330  
 Hematocrit, 267-270  
 Hemeralopia, 11  
 Hemizygote, sex-linked mutations, 11, 12  
 Hemlock, 482 (see also *Tsuga*)  
   poison (see *Conium*)  
   water (see *Cicuta*)  
 Hemocytoblast, facing page 276\*  
 Hemoglobin, histochemical test for, 559  
 Hemoglobin concentration, 267-270  
 Hemoglobin volume, 259-261  
 Hemolymph volume, 266, 267  
 Hemophilia, 11  
 Hemp, 375, 382, 388 (see also *Cannabis*)  
   Canadian, 368  
 Henbane (see *Hyoscyamus*)  
 Hepatitis virus, infectious, 431, 498  
 Heredity of blood groups and types, 249, 250  
 Herpes simplex virus, 431, 498  
*Herpetomonas*, PROTOMONADINA  
   culture media for, 526  
   nutrition, 168Fn., 171Fn.  
*Herpobdella bistrata* (leech), GNATHOBDELLIDA, 4  
 Herring, 375, 380, 400 (see also *Clupea*)  
*Heterakis* (cecal worm), RHABDITIDA, 109, 490  
*Heterodera glycine* (soybean cyst nematode), TYLENCHIDA, 494  
*H. rostochiensis* (golden nematode of potato), 424, 494  
*H. schachtii* (sugar beet nematode), 494  
*Heteropneustes fossilis* (catfish), OSTARIOPHYSI, 336  
 Heterozygote, sex-linked mutations, 11, 12  
 Hibernation, 417, 418  
*Hibiscus syriacus* (shrub althea), Malvaceae, 445  
 Hickory (see *Carya*)  
*Hieracium pilosella* (mouse-ear hawkweed), Compositae, 451  
*Hippocampus hudsonius* (Atlantic sea horse), SOLENICHTHYES, 64, 107  
*Hippodamia convergens* (convergent lady beetle), COLEOPTERA, 422  
*Hippoglossus hippoglossus* (Atlantic halibut), HETEROSOMATA, 64, 107  
*Hippomane mancinella* (manchineel), Euphorbiaceae, 347  
*Hippospongia* (commercial sponge), KERATOSA†, 110  
*Hirudo* (leech), GNATHOBDELLIDA, 70, 237  
*H. medicinalis* (medicinal leech), 109  
 Histochemical tests, 557-559  
*Histomonas*, RHIZOMASTIGINA, 492  
*Histoplasma*, Moniliaceae, 319, 321, 520  
 Hog cholera virus, 431, 498  
 Hogfish (see *Scorpaena*)  
 Holly (see *Ilex*)  
 Hollyhock, 427  
*Holothuria* (sea cucumber), ASPIDOCIROTA, 223, 340  
*Homalodisca triquetra* (leafhopper), HOMOPTERA, 502  
*Homarus* (lobster), DECAPODA, 4  
*H. americanus* (American lobster), 66, 223, 241  
*H. gammarus* (European lobster), 109, 237  
*Homo sapiens* (man), PRIMATES (see also Man)  
   acid-base, blood, 259  
   blood volumes, 263, 264, 267, 268  
   capillary blood pressure, 242  
   chromosome number, 1  
   clotting time, 325-327  
   digestive enzymes, 139  
   erythrocyte values, 267, 268  
   fatty acid source, 380  
   heart, anatomy, 142

† Subclass



- hemoglobin values, 267, 268
- life span, 106
- lung ventilation, 220
- oxygen consumption, 221
- platelet count, 271
- propagation, 57
- skeletal system, 158, 160, 162
- Homozygote, sex-linked mutations, 11, 12
- Honeybee, 385 (*see also* *Apis*)
- Hookworm (*see* *Ancylostoma*, *Necator*)
- Hoplolaimus tylenchiformis* (lance nematode), TYLENCHIDA, 494
- Hordeum* (barley), Gramineae, 213 (*see also* Barley)
- H. vulgare* (barley)
  - breeding system, 72
  - cell sap composition, 404
  - chromosome number, 8
  - light exposures for development, 444
  - light and temperature for flowering, 446
  - mineral content, 405, 411
  - parasites, 507, 509
  - photosynthesis, 215
  - pollen life span, 114
  - respiration rates, 228, 230
  - seed germination, 75
  - seed life span, 112
  - soil pH, 442
  - transpiration rates, 453
- Horistonotus* (wireworm), COLEOPTERA, 481
- Hormidium flaccidum* (water net), Ulotrichaceae, 214
- Hormone(s), invertebrate, 304-306
- Hormone(s), vertebrate
  - metabolites, 290, 292, 294, 296, 298, 300, 302
  - properties, 290-303, 388, 389
  - in urine, 187, 188
- Hornworm (*see* *Protoparce*)
- Horse (*see also* *Equus*)
  - antitoxin source, 388
  - effect of plant toxins on, 344-349
  - parasites
    - arthropod, 477-479
    - bacterial, 504-506
    - fungal, 516, 518, 520
    - helminthic, 490, 491, 493
    - viral, 498
  - sperm survival time, 49
  - sterol source, 385
  - water content, tissues and organs, 402
- Horsehair worm (*see* *Gordius*)
- Horsetail (*see* *Equisetum*)
- Humidity, effect on transpiration rates, 452
- Hummingbird (*see* *Archilochus*, *Calypte*, *Selasphorus*)
- Hurler syndrome, 11
- Hyacinth, 494
- Hyalomma* (tick), ACARI, 492
- Hyalophora cecropia* (cecropia moth), LEPIDOPTERA, 266
- Hyalopsora*, Melampsoraceae, 511
- Hydnocarpus* (chaulmoogra tree), Flacourtiaceae, 377
- Hydnum*, Hydnaceae, 512, 513
- Hydra (*see* specific genus)
- Hydra* (freshwater hydra), ATHECATA, 5, 110
- Hydrocephalus, congenital, 11
- Hydroid sting, 341
- Hyla arborea* (tree frog), SALIENTIA, 2, 107
- H. regilla* (Pacific tree frog), 63, 103
- Hylemya antiqua* (onion maggot), DIPTERA, 483
- Hylocomium* (hylocomium), Hylocomiaceae, 7, 215, 227
- Hylurgopinus rufipes* (elm bark beetle), COLEOPTERA, 422
- Hymenochaete*, Thelephoraceae, 512
- Hymenolepis carioca* (thread tapeworm), CYCLOPHYL-LIDEA, 490
- H. diminuta* (rat tapeworm), 486
- H. nana* (dwarf tapeworm), 486
- Hyoscyamus niger* (black henbane), Solanaceae, 347, 444
- Hypercapnia, effect on blood acid-base balance, 263
- Hyphantria cunea* (fall webworm), LEPIDOPTERA, 483
- Hypnum*, Hypnaceae, 7, 227
- Hypocapnia, effect on blood acid-base balance, 263
- Hypoderma* (cattle grub), DIPTERA, 479
- Hypoderma*, Phacidiaceae, 511, 513
- Hypodermella*, Phacidiaceae, 512, 513
- Hypogastric plexus, nerve connections, 130\*, 133, 135-138
- Hypoparathyroidism, 11
- Hypophosphatemia, 11
- Hypophysis, 122\*
  - comparative anatomy, 152, 153
  - hormone source, 388, 389
  - sterol source, 385
  - tissue growth, 47
- Hypothalamus, regions and functions, 123
- Hypotrachomonas*, METAMONADINA, culture media for, 524, 525
- Hypoxylon*, Xylariaceae, 512, 513
- Hyssop, hedge (*see* *Gratiola*)
- Ichthyosis simplex, 11
- Ictalurus* (bullhead), OSTARIOPHYSI, 236 (*see also* Bullhead)
  - I. catus* (white catfish), 107, 270
  - I. natalis* (yellow bullhead), 265
  - I. punctatus* (channel catfish)
    - arterial blood pressure, 240
    - body weight and length, 105
    - life span, 107
    - propagation, 64
- Idiocy, 11
- Iguana tuberculata* (tuberculate iguana), SAURIA, 222
- Ileum
  - cell life span, 46
  - effect of hormones on, 303
  - water content, 402
- Ilex* (holly), Aquifoliaceae, 72-74
- I. aquifolium* (English holly)
  - chromosome number, 9
  - light for flowering, 446
  - mineral content, 407
  - respiration rate, 230
  - soil pH, 442
- I. opaca* (American holly), 111, 412, 443
- Impatiens* (snapweed), Balsaminaceae, 54, 375
- I. biflora* (spotted snapweed), 453
- Incubation period, 59, 62, 82
- Indigo, (*see* *Baptisia*, *Indigofera*)
- Indigofera* (indigo), Leguminosae, 368
- Indole acids
  - as growth regulators, 307, 308
  - nutritional requirement for, 175
- Influenza viruses, 51, 431, 498
- Inimicus didactylus* (lumpfish), SCLEROPAREI, 338
- I. japonicus* (stonefish), 338
- Inositols, properties, 356, 357
- Interphase, effect of compounds on, 53
- Interstitial fluid, with increasing weight and age, 119\*
- Intestinal tract (*see also* Gastrointestinal tract)
  - parasites
    - fungal, 518, 520
    - viral, 498, 499
- Intestine(s)
  - cell division, compounds affecting, 53
  - enzymes, 139-141
  - nerve connections, 130\*, 135



- Intestine(s) (*concluded*)  
 parasites, 486-493  
 permissible radionuclide concentration in, 458-467  
 protein digestion, 183\*  
 radiation effect on, 468, 470, 474  
 water content, 401, 402
- Intracellular fluid and solids, with increasing weight and age, 119\*
- Iodamoeba*, AMOEBINA  
 culture medium for, 523  
 parasitism, 492
- Iodine value  
 fats, 380  
 fatty acids, 371, 373, 375, 377, 379  
 oils, 380  
 waxes, 382
- Ipomoea batatas* (sweet potato), Convolvulaceae (*see also* Sweet potato)  
 breeding system, 72  
 chromosome number, 9  
 light for flowering, 446  
 mineral content, 407, 412  
 parasites, 507, 509  
 pollen life span, 114  
 respiration rates, 228, 229, 232  
 soil pH, 442
- I. grandiflora* (large moonflower), 230  
*I. purpurea* (common morning glory), 377
- Iris* (iris), Iridaceae  
 breeding system, 72  
 glycoside source, 368  
 parasites, 494, 507  
 soil pH, 442
- I. cristata* (crested iris), 443  
*I. germanica* (German iris), 73, 230, 453  
*I. graminea* (grass iris), 114  
*I. versicolor* (blue-flag iris), 8
- Iron in tissues, histochemical test for, 559
- Ironweed (*see Vernonia*)
- Ischnochiton* (chiton), CHITONIDA  
 heart rate, 238  
 life span, 109  
 propagation, 66, 70
- Isoelectric point  
 amino acids, 392, 393  
 enzymes, 282, 284, 286, 388, 389  
 hormones, 290, 292, 294, 300, 302, 388, 389  
 proteins, 388, 389  
 toxins, 388
- Isospora, EUCOCCIDIA, 492
- Ivy, poison (*see Rhus*)
- Ixodes* (tick), ACARI, 69
- I. ricinus* (sheep tick), 3  
*I. ricinus scapularis* (black-legged tick), 478
- Jacaranda* (jacaranda), Bignoniaceae, 375
- Jack bean, 388, 389
- Jatropha curcas* (Barbados nut), Euphorbiaceae, 347
- J. multifida* (coral plant), 347
- Jejunum  
 effect of hormones on, 303  
 parasites, 489
- Jellyfish sting, 341
- Jobba, wax source, 382
- Juglans* (walnut), Juglandaceae (*see also* Walnut)  
 breeding system, 72  
 mineral content, 412  
 parasites, 507, 515  
 propagation methods, 74  
 soil pH, 442
- J. nigra* (black walnut)  
 chromosome number, 9  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 mineral content, 407, 412  
 osmotic potential, 453  
 parasites, 512  
 seed germination, 77  
 shade tolerance, 443
- J. regia* (Persian walnut)  
 mineral content, 407  
 parasites, 509  
 pollen dispersion, 429  
 respiration rate, 228
- J. sieboldiana* (Siebold walnut), 114
- Julus* (millipede), JULIDA, 70, 485
- Juniperus* (juniper), Cupressaceae, 73, 377, 515
- J. scopularum* (western red cedar), 429
- J. virginiana* (eastern red cedar)  
 chromosome number, 8  
 first flowering, 110  
 life span, 110  
 measurements, 110  
 respiration rate, 228  
 seed germination, 77  
 shade tolerance, 443  
 soil pH, 442
- Kabatiella*, Tuberculariaceae, 511
- Kale, 494
- Kalmia angustifolia* (lambkill kalmia), Ericaceae, 347
- K. latifolia* (mountain laurel), 347
- Kamala, 379
- Kangaroo, 493
- Katsuwonus pelamis* (skipjack), PERCOMORPHI, 337, 338
- Katydid (*see Microcentrum*)
- Ked, sheep (*see Melophagus*)
- Keithia*, Stictidiaceae, 514
- Keratosis follicularis, 11
- Ketosamines, properties, 355
- Ketoses, properties, 352, 353
- Kidney  
 cell chemical constituents, 398, 399  
 cells, staining method for, 551  
 effect of hormones on, 295, 297, 299  
 nerve connections, 130\*, 132, 135, 136, 138  
 parasites, 520  
 permissible radionuclide concentration in, 460-466  
 radiation effect on, 469, 470, 472, 474  
 tissue growth, 47  
 tissue regeneration, 47  
 tubules, diameter, 47  
 water content, 401, 402
- Kissing bug (*see Triatoma*)
- Klebsiella*, Enterobacteriaceae  
 antibiotic activity against, 320, 321, 323  
 parasitism, 505
- Knemidokoptes mutans* (scaly-leg mite), ACARI, 478
- Koeleria cristata* (prairie June grass), Gramineae, 456
- Krait (*see Bungarus*)
- Krebs cycle, 204\*  
 in metabolic processes, 197\*, 198\*, 203\*
- Kudzu (*see Pueraria*)
- Labia minora, nerve connections, 133, 136
- Labyrinthula*, Labyrinthulaceae, 167Fn.

- Lace bug (*see Corythuca*)
- Lachesis muta* (bushmaster), SERPENTES, 328
- Lacrimal gland, nerve connections, 130\*, 131, 134
- Lactarius*, Agaricaceae, 377
- Lactobacillus*, Lactobacillaceae  
 culture medium for, 535  
 fatty acid source, 377  
 generation time, 52  
 nutrition, 171Fn., 173Fn.  
 respiration rate, 225  
 temperature for growth, 438  
 thermal death time, 439
- Lactophrys trigonus* (trunkfish), PLECTOGNATHI, 337, 338
- Lactoria cornutus* (trunkfish), PLECTOGNATHI, 337, 338
- Lactuca sativa* (lettuce), Compositae (*see also* Lettuce)  
 breeding system, 72  
 cell sap composition, 404  
 chromosome number, 9  
 light exposures for development, 444  
 light and temperature for flowering, 446  
 mineral content, 407, 412  
 parasites  
   bacterial, 507  
   fungal, 509  
   viral, 435, 501  
 respiration rates, 230  
 seed germination, 76  
 seed life span, 112, 113  
 shade tolerance, 443  
 soil pH, 442
- Ladyfish (*see Albula*)
- Laemophloeus minutum* (flat grain beetle), COLEOPTERA, 422
- Lagocephalus* (smooth puffer), PLECTOGNATHI, 339
- Lamb's-quarters (*see Chenopodium*)
- Laminaria* (kelp), Laminariaceae  
 chromosome number, 7  
 photosynthesis, 213  
 respiration rate, 227  
 temperature tolerances, 441
- Lampetia equestris* (narcissus bulb fly), DIPTERA, 483
- Lampetra fluviatilis* (river lamprey), PETROMYZONES, 108
- L. lamottei* (American brook lamprey), 65
- Lamprey (*see Lampetra, Petromyzon*)
- Land plants, carbon production, 216
- Larix* (larch), Pinaceae  
 parasite, 482  
 propagation methods, 73  
 shade tolerance, 443  
 soil pH, 442
- L. laricina* (eastern larch), 515
- L. lyalli* (alpine larch), 515
- L. occidentalis* (western larch)  
 chromosome number, 8  
 first flowering, 110  
 life span, 110  
 measurements, 110  
 parasites, 513, 515  
 seed germination, 77
- Larkspur (*see Delphinium*)
- Larus* (gull), CHARADRIIFORMES, 59
- L. argentatus* (herring gull), 60, 106
- L. canus* (mew gull), 235, 240
- L. crassirostris* (black-tailed gull), 1
- L. hyperboreus* (glaucous gull), 222
- Lasallia papulosa* (rock tripe), Umbilicariaceae, 212
- Lasius alienus americanus* (cornfield ant), HYMENOPTERA, 483
- Lathyrus cicera* (flatpod pea), Leguminosae, 347
- L. odoratus* (sweet pea), 347
- L. sativus* (grass peavine), 347
- Laticifer, cell description, 46
- Latimeria* (coelacanth), ACTINISTIA, 153
- Laurel (*see Kalmia*)
- Laurel oak, parasites, 496
- Lead tree (*see Leucaena*)
- Leadplant (*see Amorpha*)
- Leaf(ves)  
 cell sap composition, 404  
 diseases affecting, 500-502, 506-514  
 fatty acid source, 373, 375  
 glycoside source, 368  
 mineral content, 405-409, 411-413  
 osmotic potential, 453-455  
 parasites, 481-485, 494, 495  
 respiration rates, 230-232  
 toxins, 344-349
- Leaf miner (*see Liriomyza*)
- Leafhopper(s), 426 (*see also Aceratogallia, Cicadulina, Deltocephalus, Homalodisca, Macropsis, Nephrotettix*)  
 beet (*see Circulifer*)  
 cotton (*see Psallus*)  
 potato (*see Empoasca*)  
 saddled (*see Colladonus*)  
 six-spotted (*see Macrosteles*)
- Leafworm (*see Alabama*)
- Lecanora*, Lecanoraceae, 6
- Lecidea*, Lecideaceae, 6
- Leech (*see Herpobdella*)
- Leishmania*, PROTOMONADINA  
 culture media for, 526, 527  
 parasitism, 488, 492
- Lemon (*see Citrus*)
- Lemur macaco* (lemur), PRIMATES, 158, 160, 162
- Lens culinaris* (lentil), Leguminosae, 217
- Lentil (*see Lens*)
- Lentinus*, Agaricaceae, 513
- Leontodon* (hawkbit), Compositae, 429
- Lepas anatifera* (goose barnacle), THORACICA, 4
- Lepidosiren paradoxa* (South American lungfish), DIPNOI, 2, 107, 222
- Lepisma saccharina* (silverfish), THYSANURA, 109, 483
- Lepisosteus* (gar), GINGLYMODI, 153
- L. osseus* (longnose gar), 64, 105, 107
- Lepomis cyanellus* (green sunfish), PERCOMORPHI, 107
- L. macrochirus* (bluegill), 64, 105
- Leptinotarsa decemlineata* (Colorado potato beetle), COLEOPTERA, 68, 419, 483
- L. signatolicis* (potato beetle), 4
- Leptomonas*, PROTOMONADINA, culture medium for, 526
- Leptosphaeria*, Pseudosphaeriaceae, 509-511, 518
- Leptospira*, Treponemataceae, 322, 505
- Leptosynapta* (sea cucumber), APODA, 340
- Lepus americanus virginianus* (varying hare), LAGOMORPHA, 159, 161, 163
- Lespedeza* (lespedeza), Leguminosae, 217, 494
- Lestes sponsa* (damselfly), ODONATA, 419
- Lethrinus miniatus* (snapper-like fish), PERCOMORPHI, 337, 338
- Lettuce, 501 (*see also Lactuca*)
- Leucaena glauca* (white popinac lead tree), Leguminosae, 347
- Leuconostoc*, Lactobacillaceae, 171Fn., 173Fn.
- Leukocyte(s), *facing page* 276\*  
 chemical constituents, 398  
 parasite, 493  
 radiation effect on, 468-475
- Leukocyte counts  
 blood, 272-274  
 bone marrow, 275, 276
- Licania rigida* (licania), Rosaceae, 380

- Lichens (*see specific genus*)
- Life spans (*see also* table of contents, page vi)  
 radiation effect on, 470-472, 474  
 tissues, 46-49
- Light  
 effect on plant development, 443-445  
 effect on protoplasmic streaming, 449, 450  
 exposures for flowering, 446, 447  
 speed of, 579
- Lignin, histochemical test for, 558
- Lilium* (lily), Liliaceae  
 chromosome number, 8  
 mineral content, 411  
 propagation methods, 73  
 shade tolerance, 443  
 sterol source, 385
- L. bulbiferum* (bulbil lily), 231  
*L. longiflorum* (Easter lily), 442  
*L. regale* (regal lily), 72, 112-114  
 Lily of the valley, 368
- Limanda ferruginea* (yellowtail flounder), HETEROSOMATA, 270
- Limoni* (wireworm), COLEOPTERA, 481
- Limpet (*see Acmaea*)
- Limulus* (king crab), XIPHOSURA, 66, 69, 237
- Linden (*see Tilia*)
- Lindera* (spicebush), Lauraceae, 373
- Linguatula serrata* (tongue worm), POROCEPHALIDA, 477
- Lingula* (brachiopod), ATREMATA, 70
- Linkage groups (*see* table of contents, page v)
- Limnaea borealis* (twinflower), Caprifoliaceae, 455
- Linognathus vituli* (long-nosed cattle louse), ANOPLURA, 479Fn.
- Linospora*, Diaporthaceae, 513
- Linum usitatissimum* (flax), Linaceae, 380 (*see also* Flax)
- Lipid(s) (*see also* Cerebrosides, Fats, Fatty acids, Waxes)  
 digestion, 185\*  
 fatty acid source, 375, 377, 379  
 histochemical tests for, 557  
 metabolism, 197\*  
 nutritional requirement for, 167
- Lipid content  
 cells, 398-400  
 feces, 190  
 urine, 187
- Liposcelis divinatorius* (book louse), PSOCOPTERA, 483
- Liquidambar styraciflua* (American sweet gum), Hamamelidaceae, 453
- Liriodendron tulipifera* (yellow poplar), Magnoliaceae, 453
- Liriomyza pusilla* (serpentine leaf miner), DIPTERA, 422
- Listeria*, Corynebacteriaceae, 505
- Lithobius* (centipede), HETEROSTIGMATA, 70
- Litter size, 57
- Littorina* (periwinkle), MESOGASTROPODA, 66, 109, 422
- Liver  
 and blood coagulation, 253\*  
 cell chemical constituents, 398-400  
 cells, staining method for, 552  
 effect of hormones on, 295, 297, 299, 303  
 fatty acid source, 373, 375, 380  
 nerve connections, 130\*, 132, 135, 138  
 parasites  
   fungal, 518, 520  
   helminthic, 488, 489  
   protozoan, 492, 493  
   viral, 499, 500  
 permissible radionuclide concentration in, 459-463, 465, 466  
 radiation effect on, 472-475  
 sterol source, 385
- tissue growth, 48  
 tissue regeneration, 48  
 water content, 401-403
- Lizard (*see specific genus*)
- Loa loa* (African filarial worm), SPIRURIDA, 109, 486
- Lobaria pulmonaria* (lungwort), Stictaceae, 212
- Lobster (*see Homarus*)
- Locke's solution for animal tissue culture, 529
- Locoweed (*see Astragalus*)
- Locust (*see Locusta, Schistocerca*)
- Locust, black (*see Robinia*)
- Locusta migratoria* (migratory locust), ORTHOPTERA, 4, 237, 266
- L. migratoria gallica* (migratory locust), 419
- Logarithms, 578, 580, 581
- Loligo* (squid), DIBRANCHIA, 70, 237
- L. pealeii* (squid), 109
- Lolium* (ryegrass), Gramineae, 429
- Lomatium foeniculaceum* (lomatium), Umbelliferae, 456
- Lophius* (goosefish), PEDICULATI, 157
- Lophodermium*, Phasidiaceae, 512, 513
- Lotus (*see Nelumbium*)
- Louping ill virus, 431, 478, 498
- Louse, 503 (*see also Pediculus*)  
 book (*see Liposcelis*)  
 cattle (*see Bovicola, Haematopinus, Linognathus*)  
 chicken (*see Cuculotogaster, Menacanthus*)  
 dog (*see Trichodectes*)  
 shaft (*see Menopon*)
- Lowe's syndrome, 11
- Lucerne flea (*see Sminthurus*)
- Lugworm (*see Arenicola*)
- Lumbriconereis heteropoda* (marine worm), POLYCHAETA†, 340
- Lumbricus* (earthworm), OLIGOCHAETA†, 70, 490
- L. terrestris* (earthworm)  
 chromosome number, 4  
 heart rate, 237  
 life span, 109  
 oxygen consumption, 223
- Lumpfish (*see Inimicus*)
- Lung  
 cell chemical constituents, 399  
 nerve connections, 130\*, 131, 134, 136-138  
 parasites  
   fungal, 518, 520  
   helminthic, 488, 489  
   viral, 499  
 permissible radionuclide concentration in, 458-467  
 radiation effect on, 469  
 water content, 401, 402
- Lung ventilation, 220
- Lungfish (*see Lepidosiren, Protopterus*)
- Lungworm (*see Dictyocaulus, Metastrongylus*)
- Lupinus* (lupine), Leguminosae, 217, 347
- Lutjanus bohar* (twinspot snapper), PERCOMORPHI, 337, 338
- L. gibbus* (red snapper), 337, 338
- L. monostigma* (one-spot snapper), 337, 338
- L. vaigiensis* (red snapper), 337, 338
- Lychnis dioica* (red campion), Caryophyllaceae, 337
- Lycoperdon*, Lycoperdaceae, 6
- Lycopersicon esculentum* (tomato), Solanaceae (*see also* Tomato)  
 breeding system, 72  
 cell sap composition, 404  
 chromosome linkage groups, 41-43  
 chromosome number, 9, 41  
 light exposures for development, 444  
 light and temperature for flowering, 446  
 mineral content, 408, 412  
 mutations, 41-43

† Class



- parasites
  - bacterial, 507
  - fungal, 509, 510
  - viral, 433, 434, 436, 501
- photosynthesis, 213, 215
- pollen dispersion, 429
- pollen life span, 114
- respiration rates, 229, 230, 232
- seed germination, 76
- seed life span, 112, 113
- shade tolerance, 443
- soil pH, 442
- Lycopodium clavatum* (club moss), Lycopodiaceae, 8
- L. lucidulum* (shining club moss), 443
- Lydella stabulans grisescens* (tachinid fly), DIPTERA, 422
- Lygus lineolaris* (tarnished plant bug), HEMIPTERA, 483
- Lymnaea* (freshwater snail), BASOMMATOPHORA
  - chromosome number, 4
  - heart rate, 238
  - life span, 109
  - oxygen consumption, 223
  - parasites, 492
  - propagation, 66
- Lymph nodes (*see also* Lymphatic system)
  - cells, staining method for, 552
  - parasites, 498, 500, 520
  - radiation effect on, 470, 474
- Lymphatic system
  - comparative anatomy, 148-151
  - effect of hormones on, 297, 299
  - parasites, 487, 520
- Lymphocyte(s), *facing page* 276\*
  - life span, 47
  - parasite, 493
  - radiation effect on, 470, 473-475
- Lymphocyte counts
  - blood, 273, 274
  - bone marrow, 275, 276
- Lymphogranuloma venereum virus, 431, 498
- Lynx, 493
- Lytechinus variegatus* (sea urchin), TEMNOPLEUROIDA, 54
  
- Macaca irus* (crab-eating macaque), PRIMATES, 235
- M. mulatta* (rhesus monkey) (*see also* Monkey, rhesus)
  - arterial blood pressure, 240
  - blood volumes, 264, 268
  - body surface area constants, 121
  - body weight, 99, 121
  - chromosome number, 1
  - erythrocyte and hemoglobin values, 268
  - life span, 106
  - lung ventilation, 220
  - platelet count, 271
  - propagation, 57
  - skeletal system, 158, 160, 162
- Mackerel (*see* *Scomber*)
- Macracanthorhynchus* (thorny-headed worm), ARCHIA-CANTHOCEPHALA, 70, 490
- Macrophomina*, Sphaeropsidaceae, 509, 510
- Macropsis trimaculata* (leafhopper), HOMOPTERA, 502
- Macrosiphum* (aphid), HOMOPTERA, 501
- Macrosteles divisus* (six-spotted leafhopper), HOMOPTERA, 422
- Mactra* (bar clam), EULAMELLIBRANCHIA, 4
- Madurella*, Dematiaceae, 518
- Maggot (*see* *Hylemya*, *Rhagoletis*)
- Magiccada septendecim* (periodical cicada), HOMOPTERA
  - chromosome number, 4
  - life span, 109
  - metamorphosis, 68
- parasitism, 483
- propagation, 68
- Magnolia* (magnolia), Magnoliaceae, 9, 74
- M. grandiflora* (southern magnolia), 111, 442
- M. macrophylla* (big-leaf magnolia), 408, 412
- Magpie (*see* *Pica*)
- Maidenhair (*see* *Adiantum*)
- Maize streak virus, 502
- Malaclemys terrapin* (diamondback terrapin), CHELONIA, 62
- M. terrapin centrata* (southern diamondback terrapin), 107, 220, 222
- M. terrapin pileata* (Mississippi diamondback terrapin), 103
- Malacosoma disstria* (forest tent caterpillar), LEPIDOPTERA, 419, 483
- Malassezia*, Cryptococcaceae, 516
- Malus coronaria* (sweet crab apple), Rosaceae, 443
- M. pumila* (common apple)
  - breeding system, 72
  - chromosome number, 9
  - parasites, 507, 510
  - photosynthesis, 213
  - pollen dispersion, 429
  - pollen life span, 114
  - propagation method, 74
  - respiration rate, 228-230, 232
  - soil pH, 442
  - transpiration rates, 451
- M. sylvestris* (apple), 215, 408, 412
- Mamba (*see* *Dendroaspis*)
- Mammary glands
  - effect of hormones on, 293, 299, 301
  - radiation effect on, 471, 473, 474
- Man (*see also* *Homo sapiens*)
  - acid-base, blood, 259, 262, 263
  - autonomic nervous system, 130\*, 131-138
  - blood group systems, 245-252
  - blood pressures, 238, 239, 241, 242
  - body composition, with increasing weight and age, 119\*
  - body surface area, 120
  - body weight and height, 93, 94
  - brain, regions and functions, 122\*, 123-129
  - carbohydrate digestion, 184\*
  - cell chemical constituents, 398
  - cells, blood and bone marrow, *facing page* 276\*
  - differential cell count, sternal marrow, 275, 276
  - effect of
    - animal toxins on, 329, 331, 333, 336-342
    - hormones on, 290-303
    - plant toxins on, 344-349
  - enzyme source, 388
  - erythrocyte growth, 47
  - feces, 190
  - gestation period, 82
  - glomerulus diameter, 47
  - granulocyte growth, 47
  - hair growth, 47
  - heart fibers, diameter, 47
  - heart rate, 234
  - hormone source, 388
  - leukocyte counts, 272, 273
  - lipid digestion, 185\*
  - mutations, 11, 12
  - nerve connections, 130\*, 131-138
  - parasites
    - arthropod, 477-480
    - bacterial, 504-506
    - fungal, 516-521
    - helminthic, 486-493



- Man (concluded)  
 parasites (concluded)  
   protozoan, 488, 489, 492, 493  
   ricketsial, 503  
   viral, 498-500  
 parathyroid, 154  
 permissible radiation exposure for, 457-467  
 prenatal development, 79\*-81\*, 82, 83  
 protein digestion, 183\*  
 protein source, 388, 389  
 races and nationalities, body weight and height,  
   93, 94  
 radiation effect on, 468-470  
 sex linkage, 11, 12  
 sperm survival time, 49  
 thymus gland weight, 49  
 tissue growth, 46-49  
 tissue regeneration, 46-49  
 urine, 186-188  
 water content, tissues and organs, 401
- Man-o'-war (see *Physalia*)  
 Manatee (see *Trichechus*)  
 Manchineel (see *Hippomane*)  
 Mangel-wurzel, 494  
*Manihot esculenta* (cassava), Euphorbiaceae, 348  
*Mansonia* (mosquito), DIPTERA, 486  
 Mantis (see *Mantis*, *Stagmomantis*)  
*Mantis religiosa* (praying mantis), ORTHOPTERA, 4, 109  
 Maple (see *Acer*)  
   sugar, 483
- Marchantia*, Marchantiaceae, 7, 227  
 Margosa (see *Melia*)  
 Marigold (see *Tagetes*)  
 Marine worm (see *Lumbriconereis*)  
*Marmota marmota* (Eurasian marmot), RODENTIA, 417  
*M. monax* (woodchuck), 417  
 Marrow cells, facing page 276\*  
*Marssonina*, Melanconiaceae, 513  
 Masked hunter (see *Reduvius*)  
 Massasauga (see *Sistrurus*)  
 Maternity, putative: phenotypes refuting, 249, 250  
 Mayfly (see *Ephemera*)  
 MB 752/1 culture medium for animal tissue, 533  
 MD 705/1 culture medium for animal tissue, 533, 534  
 Mealworm (see *Tenebrio*)  
 Mealybug (see *Pseudococcus*)  
 Measles virus, 431, 499  
 Media, culture (see table of contents, page ix)  
*Medicago hispida denticulata* (toothed bur clover), Legu-  
   minosae, 217
- M. sativa* (alfalfa) (see also Alfalfa)  
   breeding system, 72  
   chromosome number, 9  
   light and temperature for flowering, 447  
   mineral content, 408, 413  
   nitrogen fixation, 217  
   parasites, 433, 501, 507, 510  
   photosynthesis, 213, 215  
   pollen life span, 114  
   respiration rates, 228  
   seed germination, 76  
   seed life span, 112  
   soil pH, 442  
   transpiration rates, 453
- Medulla oblongata, 122\*  
   nerve connections, 130\*  
   regions and functions, 123
- Megachile latimanus* (leaf-cutting bee), HYMENOPTERA,  
 483  
 Megakaryocyte, facing page 276\*  
 Megakaryocyte count, bone marrow, 275, 276  
 Megaloblast, facing page 276\*  
 Megalocornea, 11
- Melampsora*, Melampsoraceae, 512-514  
*Melampsorella*, Melampsoraceae, 512  
*Melampsoridium*, Melampsoraceae, 512  
 Melanocytes, effect of hormones on, 295  
*Melanogrammus* (haddock), ANACANTHINI, 236  
*M. aeglefinus* (haddock), 64, 105, 107  
*Melanoplus* (grasshopper), ORTHOPTERA, 422  
*M. bivittatus* (two-striped grasshopper), 419  
*M. differentialis* (differential grasshopper)  
   chromosome number, 4  
   diapause, 419  
   heart rate, 237  
   life span, 109  
*M. femur-rubrum* (red-legged grasshopper), 483  
*M. mexicanus* (migratory grasshopper), 68  
*Melanotus* (wireworm), COLEOPTERA, 481  
*Meleagris* (turkey), GALLIFORMES, 59 (see also  
 Turkey)  
*M. gallopavo* (turkey)  
   arterial blood pressure, 240  
   blood volumes, 269  
   body weight, 101, 102  
   breeds, body weight, 101, 102  
   chromosome number, 1  
   erythrocyte values, 269  
   heart rate, 235  
   hemoglobin values, 269  
   leukocyte counts, 274  
   life span, 106  
   lung ventilation, 220
- Melia azadirachta* (margosa), Meliaceae, 380  
*M. azedarach* (chinaberry), 348  
*Melilotus* (clover), Leguminosae, 217  
*M. alba* (white sweet clover), 217  
*M. indica* (annual yellow sweet clover), 217  
*Melittobia chalybii* (chalcid), HYMENOPTERA, 419  
*Meloidogyne* (root-knot nematode), TYLENCHIDA, 496  
*Melolontha* (June beetle), COLEOPTERA, 69, 223  
*M. vulgaris* (June beetle), 109  
*Melophagus ovinus* (sheep ked), DIPTERA, 479  
*Melospiza melodia* (song sparrow), PASSERIFORMES,  
 59-61 (see also Sparrow, song)  
 Melting point  
   aglycones, 369, 371  
   aldaric acids, 359  
   alditols, 356  
   aldonic acids, 358  
   aldoses, 351, 352  
   amino acids, 392, 393  
   amino sugars, 355  
   antibiotics, 313, 315, 317  
   carbohydrates, 351-359, 364-366  
   fats, 380  
   fatty acids, 370, 372, 374, 376, 378  
   glycosides, 368, 370  
   inositols, 356, 357  
   ketoses, 352, 353  
   monosaccharides, 351-356  
   oils, 380  
   oligosaccharides, 364-366  
   phosphatides, 383, 384  
   provitamins, 396  
   sphingolipids, 384  
   sterols, 385, 386  
   uronic acids, 359  
   vitamins, 394, 396  
   waxes, 382
- Memory, brain involvement, 123-125  
*Menacanthus stramineus* (chicken body louse), MAL-  
 LOPHAGA, 479  
 Menhaden (see *Brevoortia*)  
*Menopon gallinae* (shaft louse), MALLOPHAGA, 479

- Mercenaria* (quahog), EULAMELLIBRANCHIA, 66, 70, 109
- Merodon equestris* (narcissus bulb fly), DIPTERA, 422
- Mesencephalon  
     nerve connections, 130\*  
     regions and functions, 123
- Mesenteric ganglia, connections, 130\*, 137
- Mesenteric plexus, connections, 138
- Mesocricetus auratus* (golden hamster), RODENTIA  
     acid-base, blood, 260  
     arterial blood pressure, 240  
     blood volumes, 268  
     body weight, 95  
     chromosome number, 1  
     clotting time, 326  
     erythrocyte values, 268  
     heart rate, 235  
     hemoglobin values, 268  
     hibernation, 260, 417  
     life span, 106  
     lung ventilation, 220  
     oxygen consumption, 221  
     platelet count, 271  
     propagation, 57
- Mesoderm, eutherian mammals, 80\*
- Metabolism (see table of contents, pages xii, xiii)
- Metabolites of hormones, 290, 292, 294, 296, 298, 300, 302
- Metamorphosis  
     amphibians, 63, 90  
     insects, 67, 68
- Metamyelocyte, facing page 276\*
- Metamyelocyte count, bone marrow, 275, 276
- Metaphase, effect of compounds on, 53
- Metastrongylus elongatus* (swine lungworm), RHABDITIDA, 490
- Metatetranychus ulmi* (European red mite), ACARI, 419
- Metathalamus, regions and functions, 123
- Metencephalon, regions and functions, 123
- Metopium toxiferum* (poisonwood), ANACARDIACEAE, 348
- Metridium* (sea anemone), ACTINIARIA, 70
- Michaelis constants for enzymes, 283, 285, 287
- Microcentrum rhombifolium* (broad-winged katydid), ORTHOPTERA, 483
- Micrococcus*, Micrococcaceae, 218\*, 225, 322
- Microphthalmia, 11
- Micropterus salmoides* (largemouth black bass), PERCOMORPHI (see also Bass, largemouth)  
     arterial blood pressure, 240  
     body weight and length, 105  
     heart rate, 236  
     life span, 107  
     propagation, 64
- Microsome, chemical constituents, 399, 400
- Microsporium*, Moniliaceae, 320, 321, 516
- Micrurus corallinus* (coral snake), SERPENTES, 330
- M. fulvius* (eastern coral snake), 330
- Milesia*, Melampsoraceae, 512
- Milk  
     fatty acid source, 371, 373, 379-381  
     parasites, 498  
     protein source, 388, 389
- Milkweed (see *Asclepias*)
- Millepora alcicornis* (stinging coral), ATHECATA, 341
- Millet, 494 (see also *Pennisetum*, *Setaria*)
- Millipede (see *Julus*)
- Mineral(s) (see also Chemical elements)  
     metabolism, 192\*, 193-195  
     nutritional requirements for, 165
- Mineral content  
     cell sap, 404  
     feces, 190  
     plant tissues, 405-409, 411-413  
     sea water, 539  
     urine, 186, 187
- Mink, 480, 493 (see also *Mustela*)
- Minous monodactylus* (stonefish), SCLEROPAREI, 338
- Minute volume, lungs, 220
- Mistletoe (see *Arceuthobium*, *Phoradendron*)
- Mite, 503  
     blister (see *Eriophes*)  
     bulb (see *Rhizoglyphus*)  
     chicken (see *Dermanyssus*)  
     cyclamen (see *Steneotarsonemus*)  
     ear (see *Otodectes*)  
     follicle (see *Demodex*)  
     fowl (see *Bdellonyssus*)  
     grass (see *Oribatula*)  
     itch (see *Sarcoptes*)  
     red (see *Metastrongylus*)  
     scab (see *Psoroptes*)  
     scaly-leg (see *Knemidokoptes*)  
     spider (see *Tetranychus*)
- Mitochondria  
     chemical constituents, 399, 400  
     cytochromes, 206  
     staining methods for, 552, 556
- Miyagawanella*, Chlamydiaceae, 438, 439, 503
- M-N blood group system, 246  
     distribution in various populations, 251, 252  
     heredity, 249
- Mnium*, Mniaceae, 7, 227
- Modiolus modiolus* (horse mussel), FILIBRANCHIA, 340
- Mola mola* (common ocean sunfish), PLECTOGNATHI, 339
- Molds, culture media for, 536
- Mole, 503
- Molecular activity, enzymes, 283, 285, 287
- Molecular weights  
     amino acids, 392, 393  
     antibiotics, 312, 314, 316  
     cytochromes, 206, 207  
     enzymes, 282, 284, 286, 388, 389  
     fatty acids, 370, 372, 374, 376, 378  
     hormones, 290, 292, 294, 296, 298, 300, 302, 388, 389  
     phosphatides, 383, 384  
     proteins, 388, 389  
     sphingolipids, 384  
     toxins, 388, 389
- Molgula manhattensis* (sea squirt), PLEUROGONA, 237
- Molluscum contagiosum virus, 499
- Moniezia expansa* (sheep tapeworm), CYCLOPHYLLIDEA, 110, 490
- Monilia*, Sclerotiniaceae, 510
- Monilochaetes*, Dematiaceae, 509
- Monkey (see also *Alouatta*, *Macaca*)  
     parasites  
         bacterial, 505  
         fungal, 516, 518, 520  
         helminthic, 486, 488, 490, 491  
         protozoan, 493  
         rickettsial, 503  
         viral, 498-500  
         rhesus, 82 (see also *Macaca*)
- Monkshood (see *Aconitum*)
- Monocercomonas*, METAMONADINA, culture media for, 524, 525
- Monocystis*, EUGREGARINA, 71
- Monocyte(s), facing page 276\*  
     staining method for, 552
- Monocyte count  
     blood, 272-274  
     bone marrow, 275, 276
- Monoglycerides  
     digestion, 185\*  
     metabolism, 197\*

- Monosaccharides  
     digestion, 184\*  
     properties, 351-355  
 Moonflower (see *Ipomoea*)  
 Moose, 490, 504  
*Moraxella*, Brucellaceae, 505  
*Morenoella*, Microthyriaceae, 513  
 Morning glory (see *Ipomoea*)  
 Morphology (see table of contents, page vi)  
 Morula, 79\*-81\*  
 Mosaic viruses, 426, 501, 502  
 Mosquito (see specific genus)  
 Moss (see specific genus)  
 Moth  
     cecropia (see *Hyalophora*)  
     codling (see *Carpocapsa*)  
     cynthia (see *Samia*)  
     diamondback (see *Plutella*)  
     flour (see *Ephestia*)  
     fruit (see *Grapholitha*)  
     gypsy (see *Anastatus*, *Porthetria*)  
     Indian meal (see *Plodia*)  
     meal, 486  
     tobacco (see *Ephestia*)  
     underwing (see *Catocala*)  
 Motor reflexes, brain involvement, 123-129  
 Mouse (see also *Mus*)  
     anesthetic for, 548  
     cell chemical constituents, 399  
     cells, staining method for, 551, 552  
     developmental stages, 84-86  
     effect of animal toxins, 328, 330, 332, 338-341  
     parasites  
         fungal, 516, 519-521  
         helminthic, 486  
         ricketsial, 503  
         viral, 498, 499  
     radiation effect on, 470-472  
     sperm survival time, 49  
     tissue growth, 48, 49  
     tissue regeneration, 49  
 Mucin, histochemical test for, 558  
 Mud puppy (see also *Necturus*)  
     anesthetic for, 548  
 Mule, 479, 504, 520  
 Mummichog (see *Fundulus*)  
 Mumps virus, 431, 499  
*Murgantia histrionica* (harlequin cabbage bug), HEMIP-  
     TERA, 483  
*Mus* (mouse), RODENTIA, 53, 264, 274 (see also Mouse)  
*M. musculus* (house mouse)  
     arterial blood pressure, 240  
     blood volumes, 268  
     body surface area constants, 121  
     body weight, 95, 121  
     chromosome linkage groups, 13-16  
     chromosome number, 1, 13  
     clotting time, 326  
     erythrocyte values, 268  
     heart rate, 235  
     hemoglobin values, 268  
     life span, 106  
     lung ventilation, 220  
     mutations, 13-16  
     oxygen consumption, 221  
     platelet count, 271  
     propagation, 57  
     strains, body weight, 95  
*Musca domestica* (housefly), DIPTERA  
     chromosome number, 4  
     dispersion, 422  
     life span, 109  
     metamorphosis, 68  
     oxygen consumption, 223  
     parasitism, 480  
     propagation, 68  
*Muscardinus avellanarius* (common dormouse), RODEN-  
     TIA, 417  
 Muscle(s)  
     effect of hormones on, 293, 295, 297, 299, 301,  
         303  
     enzyme source, 388, 389  
     mineral retention in, 192\*, 193-195  
     parasites, 490, 491  
     permissible radionuclide concentration in,  
         462Fn.  
     protein source, 388, 389  
     tissue growth, 48  
     tissue regeneration, 48  
     water content, 401-403  
 Muskmelon (see *Cucumis*)  
 Muskrat (see *Ondatra*)  
 Mussel (see *Mytilus*)  
     horse (see *Modiolus*)  
     pearl (see *Unio*)  
     swan (see *Anodonta*)  
     white (see *Donax*)  
 Mustard (see *Brassica*)  
*Mustela vison* (least weasel), CARNIVORA, 221  
*M. vison* (mink)  
     chromosome number, 1  
     heart rate, 235  
     life span, 106  
     propagation, 57  
*Mustelus* (dogfish), SELACHII, 157  
 Mutations (see Linkage groups, table of contents, page  
     xi)  
*Mya arenaria* (soft shell clam), EULAMELLIBRANCHIA,  
     341  
*Mycobacterium*, Mycobacteriaceae  
     antibiotic activity against, 320, 322, 323  
     generation time, 52  
     nutrition, 176Fn.  
     parasitism, 505  
     protein source, 389  
     respiration rate, 225  
     temperature for growth, 438  
     thermal death time, 439  
*Mycosphaerella*, Mycosphaerellaceae, 509, 510, 512  
*Mycteroperca venenosa* (sea bass), PERCOMORPHI,  
     337, 338  
 Myelencephalon, regions and functions, 123  
 Myelin sheath, staining method for, 555  
 Myeloblast count, bone marrow, 275, 276  
 Myelocyte(s), facing page 276\*  
 Myelocyte count, bone marrow, 275, 276  
*Myliobatis californicus* (bat stingray), BATOIDEI, 338  
 Myocardium, parasite, 493  
*Myotis keenii* (long-eared little brown bat), CHIROP-  
     TERA, 417  
*M. lucifugus* (little brown bat)  
     blood volumes, 264  
     body surface area constants, 121  
     body weight, 121  
     heart rate, 235  
     hibernation, 417  
     life span, 106  
     oxygen consumption, 221  
     propagation, 57  
*M. myotis* (common brown bat), 1, 417  
 Myrtle, 382  
*Mytilus* (mussel), FILIBRANCHIA, 223  
*M. californianus* (ocean mussel), 341



- M. edulis* (mussel)  
 heart rate, 237  
 life span, 109  
 propagation, 66  
 toxin, 341
- Myxine glutinosa* (Atlantic hagfish), MYXINI, 2, 65, 270
- Myxophyta (see specific genus)
- Myzus persicae* (green peach aphid), HOMOPTERA, 422, 483, 500-502
- Nail(s)  
 growth, 48  
 parasites, 516-519  
 radiation effect on, 469
- Naja naja* (Indian cobra), SERPENTES, 1, 107, 330
- N. nigricollis* (black-necked cobra), 332
- Nanophyetus salmincola* ("salmon-poisoning" fluke), DIGENEA, 492
- Naphthalene compounds, as growth regulators, 308, 309
- Naphthoxy compounds, as growth regulators, 308, 309
- Narcissus, 483, 494
- Nasturtium (see *Tropaeolum*)
- Natrix erythrogaster* (copper-bellied water snake), SERPENTES, 62
- N. natrix* (European water snake), 222, 236, 240
- N. septemvittata* (queen snake), 103
- N. sipedon* (North American water snake), 107, 269
- N. tigrina* (Japanese water snake), 1
- Navicula*, Naviculaceae, culture medium for, 537
- NCTC culture media for animal tissue, 532
- Necator americanus* (hookworm), RHABDITIDA, 109, 486
- Nectria*, Nectriaceae, 512, 513
- Necturus* (mud puppy), CAUDATA, 154 (see also Mud puppy)
- N. maculosus* (mud puppy)  
 blood volumes, 269  
 chromosome number, 2  
 erythrocyte and hemoglobin values, 269  
 life span, 107  
 propagation, 63
- Neisseria*, Neisseriaceae  
 antibiotic activity against, 320-323  
 nutrition, 175Fn.  
 parasitism, 505  
 temperature for growth, 438  
 thermal death time, 439
- Nehumbium nelumbo* (Hindu lotus), Nymphaeaceae, 112
- Nematode (see specific genus)
- Neofabraea*, Dermaceae, 510, 513
- Neomentia* (solenogaster), NEOMENIOMORPHA, 70
- Neopeckia*, Phaeodidymae, 513
- Neorickettsia*, Rickettsiaceae, 503
- Nephotettix apicalis* (leafhopper), HOMOPTERA, 501
- Nereis* (clam worm), POLYCHAETA†  
 chromosome number, 4  
 heart rate, 237  
 life span, 109  
 oxygen consumption, 223  
 propagation, 70
- Nerium oleander* (oleander), Apocynaceae 348
- Nerve(s)  
 connections, 130\*, 131-138  
 fiber regeneration, 47  
 water content, 401-403
- Nerve endings, staining method for, 552
- Nervous system (see also Autonomic nervous system, table of contents, page vi)  
 hormone, 302, 303
- Nesting, 59
- † Class
- Nettle (see *Urtica*)
- Neurohypophysis  
 comparative anatomy, 152, 153  
 hormones, 292-295
- Neurons, ganglionic, 132-136
- Neurospora crassa*, Melanosporaceae  
 chromosome linkage groups, 33-36  
 chromosome number, 6, 33  
 culture medium for, 536  
 mutations, 33-36  
 nutrition, 168Fn., 171Fn.  
 respiration rate, 225
- Neurula, developmental stages, 83, 85, 87, 89, 91
- Neutrophil(s), facing page 276\*  
 life span, 47  
 radiation effect on, 470
- Neutrophil count, 272-274
- Newcastle disease virus, 432, 499
- Newt, common (see also *Triturus*)  
 anesthetic for, 548
- Nicandra physalodes* (apple of Peru), Solanaceae, 433
- Nicotiana* (tobacco), Solanaceae (see also Tobacco)  
 mineral content, 413  
 nutrition, 178Fn.  
 parasites, 507  
 respiration rates, 229, 230  
 tissue culture, 116
- N. alata* (winged tobacco), 436
- N. glutinosa* (tobacco), 433, 435, 436
- N. repanda*, 434
- N. rustica* (Mahorka tobacco), 434, 435
- N. sylvestris* (tobacco), 114, 435
- N. tabacum* (common tobacco)  
 breeding system, 72  
 cell division, 53, 54  
 chromosome number, 9  
 light exposures for development, 445  
 light and temperature for flowering, 447  
 mineral content, 408  
 parasites, 432-436, 501, 510  
 photosynthesis, 213  
 respiration rate, 232  
 seed germination, 76  
 seed life span, 112  
 soil pH, 442  
 transpiration rates, 451
- Niger seed (see *Guizotia*)
- Nightshade (see *Solanum*)
- Nitella*, Characeae, 448
- Nitrobacter*, Nitrobacteraceae, 218\*
- Nitrogen, nutritional requirement for, 165, 168, 178, 179
- Nitrogen content  
 cells, 398-400  
 enzymes, 288, 290  
 feces, 190  
 respiratory media, 219  
 urine, 187
- Nitrogen cycle, 218\*
- Nitrogen fixation, plants, 216, 217
- Nitrosomonas*, Nitrobacteraceae, 179Fn., 218\*
- Nocardia*, Actinomycetaceae, 314, 520
- Nodular worm (see *Oesophagostomum*)
- Normal solutions, formulas for, 543
- Normoblast(s), facing page 276\*
- Normoblast count, bone marrow, 275, 276
- Nose, parasites, 477, 480, 520
- Nostoc*, Nostocaceae  
 in nitrogen cycle, 218\*  
 nitrogen fixation, 217  
 photosynthesis, 215  
 respiration rate, 227  
 temperature for growth, 441



- Notechis scutatus* (tiger snake), SERPENTES, 332  
*Notesthes robusta* (scorpion fish), SCLEROPAREI, 338  
 Nuclei, staining methods for, 553, 554  
 Nuclei, thalamic (see Thalamic nuclei)  
 Nucleoproteins, catabolism, 201\*  
 Numerical constants, 578  
 Nutrition (see table of contents, page vi)  
*Nyctalus noctula* (noctule bat), CHIROPTERA, 417  
*Nymphon* (sea spider), NYMPHONOMORPHA, 69  
 Nystagmus, 11
- Oak, 370, 481, 483 (see also *Quercus*)  
 Oat (see also *Avena*)  
   parasites, 426, 494, 496  
   sterol source, 386  
*Obelia* (marine hydra), THECATA, 5, 70  
 Ocean  
   plants, carbon production, 216  
   as respiratory medium, 219  
   salinity, 539  
*Ochromonas*, CHRYSOMONADINA, culture medium for, 528  
*Octopus* (octopus), DIBRANCHIA, 340  
 Octopus bite, 340  
 Oculomotor nerve, 130\*, 134  
*Oedogonium*, Oedogoniaceae, 7  
*Oenothera biennis* (common evening primrose), Oenotheraceae  
   chromosome number, 9  
   fatty acid source, 374, 375  
   light and temperature for flowering, 447  
   pollen life span, 114  
   respiration rate, 230  
   seed life span, 113  
   soil pH, 442  
*O. speciosa* (white evening primrose), 445  
*Oesophagostomum columbianum* (sheep nodular worm), RHABDITIDA, 490  
*Oestrus ovis* (sheep botfly), DIPTERA, 480  
 Oil(s)  
   fatty acid source 372-380  
   properties, 380  
 Oil palm, 494 (see also *Elaeis*)  
 Okra, 484, 494, 496  
*Olea europaea sativa* (olive), Oleaceae, 380 (see also Olive)  
 Oleander (see *Nerium*)  
 Olfaction, brain involvement, 123, 124, 128  
 Oligosaccharides, properties, 364-366  
 Olive, 496 (see also *Olea*)  
*Onchocerca volvulus* (convoluted filarial worm), SPIRURIDA, 486  
*Oncorhynchus* (salmon), ISOSPONDYLI, 157, 492  
*Ondatra zibethica* (muskrat), RODENTIA  
   chromosome number, 1  
   heart rate, 235  
   life span, 106  
   propagation, 57  
 Onion (see also *Allium*)  
   parasites  
     arthropod, 481, 483, 485  
     fungal, 427  
     nematode, 494  
     viral, 500  
*Ophiobolus*, Pseudosphaeriaceae, 511  
*Ophioderma longicauda* (brittle star), OPHIURAE, 223  
*Ophiophagus hannah* (king cobra), SERPENTES, 332  
*Ophiopholis* (brittle star), OPHIURAE, 69  
 Ophthalmoplegia, 11  
 Opossum, 82, 493, 547 (see also *Didelphis*)  
*Opsanus tau* (oyster toadfish), HAPLODOCI, 339  
 Orange, 382 (see also *Citrus*)  
*Orconectes immunis* (crayfish), DECAPODA, 66  
*Oribatula* (grass mite), ACARI, 490  
*Ornithorhynchus* (platypus), MONOTREMATA  
   arterial blood pressure, 240  
   interosseal artery, 146  
   life span, 106  
   oxygen consumption, 221  
   skeletal system, 159, 161, 163  
*O. anatinus* (platypus), 1, 57  
*Oryctolagus cuniculus* (European rabbit), LAGOMORPHA (see also Rabbit)  
   acid-base, blood, 260  
   arterial blood pressure, 240  
   blood volumes, 264, 269  
   body surface area constants, 121  
   body weight, 99, 100, 121  
   cell division, 53  
   chromosome linkage groups, 17, 18  
   chromosome number, 1, 17  
   clotting time, 326, 327  
   digestive enzymes, 140, 141  
   erythrocyte values, 269  
   heart rate, 235  
   hemoglobin values, 269  
   leukocyte counts, 274  
   life span, 106  
   lung ventilation, 220  
   mutations, 17, 18  
   oxygen consumption, 221  
   platelet count, 271  
   propagation, 57  
*Oryza sativa* (rice), Gramineae (see also Rice)  
   breeding system, 72  
   chromosome number, 8  
   light and temperature for flowering, 446  
   mineral content, 405, 411  
   parasites, 501, 507, 510  
   photosynthesis, 215  
   pollen dispersion, 429  
   respiration rates, 228, 229  
   seed germination, 75  
   soil pH, 442  
*Oryzaephilus surinamensis* (saw-toothed grain beetle), COLEOPTERA, 485  
*Oscillatoria*, Oscillatoriaceae, 181Fn., 441  
*Oscinella frit* (frit fly), COLEOPTERA, 484  
*Osmerus eperlanus* (European smelt), ISOSPONDYLI, 2  
*O. mordax* (American smelt), 64, 105, 107  
 Osmotic potential, plants, 453-456  
*Osmunda cinnamomea* (cinnamon fern), Osmundaceae, 453  
*Ostertagia circumcincta* (brown stomach worm of sheep), RHABDITIDA, 490  
*O. ostertagi* (brown stomach worm of cattle), 490  
*Ostrea edulis* (oyster), EULAMELLIBRANCHIA, 109, 237  
 Ostrich (see *Struthio*)  
*Otobius megnini* (ear tick), ACARI, 478  
*Otodectes cyanotis* (ear mite), ACARI, 478  
 Ovarian follicle, effect of hormones on, 293  
 Ovary(ies)  
   hormones, 298-301  
   nerve connections, 133, 138  
   permissible radionuclide concentration in, 465Fn.  
   protein source, 389  
   radiation effect on, 468, 469, 471-473, 475  
   tissue growth, 48  
   tissue regeneration, 48  
   water content, 401, 402  
*Ovis aries* (sheep), ARTIODACTYLA (see also Sheep)  
   acid-base, blood, 260

- arterial blood pressure, 240
- blood volumes, 264, 269
- body surface area constants, 121
- body weight, 100, 121
- breeds, body weight, 100
- chromosome number, 1
- digestive enzymes, 139
- erythrocyte values, 269
- fatty acid source, 380
- heart rate, 235
- hemoglobin values, 269
- leukocyte counts, 274
- life span, 106
- oxygen consumption, 221
- propagation, 57
- tallow source, 380
- Ovulinia*, Helotiaceae, 427
- Ovum
  - chemical composition, 400
  - fertility loss, 48
  - fertilized, 80\*, 81\*
- Oxidation (see Cytochromes, Krebs cycle)
- Oxidation-reduction indicators, 545
- Oxidation-reduction potential, cytochromes, 206, 207
- Oxygen
  - in metabolism, 203\*
  - in nitrogen fixation, 217
  - in respiratory media, 219
- Oxygen consumption (see also table of contents, page xiii)
  - during hibernation, 417, 418
- Oxygen release, via cytochrome system, 205\*
- Oxygen saturation
  - plants: effect on protoplasmic streaming, 450
  - sea water, 540, 541
- Oxyuramus scutellatus* (taipan), SERPENTES, 332
- Oyster (see *Crassostrea*, *Ostrea*)
  
- Paddefisk (see *Batrachus*)
- Paddlefish (see *Polyodon*)
- Paeonia tenuifolia* (fernleaf peony), Ranunculaceae, 54
- Pagellus erythrinus* (porgy), PERCOMORPHI, 337, 338
- Pagrus pagrus* (porgy), PERCOMORPHI, 337, 338
- Pain, brain involvement, 127-129
- Palaearcta vernata* (spring cankerworm), LEPIDOPTERA, 484
- Palm, 373, 380, 382 (see also specific genus)
- Pan troglodytes* (chimpanzee), PRIMATES 158, 160, 162 (see also Chimpanzee)
- Pancreas
  - cell chemical constituents, 398
  - comparative anatomy, 156, 157
  - effect of hormones on, 297, 299, 303
  - enzymes, 139-141, 388, 389
  - hormones, 302, 303
  - islets of Langerhans, 156, 157
  - nerve connections, 132, 134, 138
  - parasites, 489, 499
  - permissible radionuclide concentration in, 460
  - tissue growth, 48
  - tissue regeneration, 48
  - water content, 401, 402
- Panicum virgatum* (switch grass), Gramineae, 429
- Panstrongylus*, HFMIPTERA, 492
- Panus*, Agaricaceae, 6
- Papaver somniferum* (opium poppy), Papaveraceae, 348, 380
- Paracaudina chilensis* (sea cucumber), MOLPADONIA, 340
- Paracentrotus lividus* (sea urchin), ECHINOIDA, 400
- Paracoccidioides*, Moniliaceae, 319, 520
- Paradicichthys venenatus* (Chinaman fish), PERCOMORPHI, 337, 338
- Paragonimus westermani* (lung fluke), DIGENEA, 488
- Parainfluenza virus, 499
- Paramecium* (ciliate), HYMENOSTOMATIDA
  - cell division frequency, 51
  - nutrition, 167Fn., 168Fn.
  - oxygen consumption, 224
  - propagation, 71
- Paraplegia, spastic, 12
- Parascaris equorum* (large roundworm), RHABDITIDA, 490
- Parasites (see table of contents, page ix)
- Parasympathetic nervous system, 130\*, 134-137
- Parathyroid
  - comparative anatomy, 154, 155
  - hormones, 296, 297
  - tissue growth, 48
- Paratylenchus* (pin nematode), TYLENCHIDA, 496
- Pardosa* (spider), ARANEAE, 69
- Parenchyma, plant
  - cell description, 44, 45
  - osmotic potential, 455
- Parinarium* (parinarium), Rosaceae, 375
- Parmelia*, Parmeliaceae, 227
- Parotid gland, nerve connections, 130\*, 132, 134
- Parrot fish (see *Scarus*)
- Pars intermedia, hormones, 294, 295
- Parsley, 368, 373
- Parsnip (see *Pastinaca*)
- Parthenium argentatum* (guayule parthenium), Compositae, 429
- Partridge, 479, 493
  - European (see *Perdix*)
- Parupeneus chryserydros* (surmullet), PERCOMORPHI, 337, 338
- Paspalum notatum* (Pensacola Bahia grass), Gramineae, 429
- Passer (sparrow), PASSERIFORMES, 59 (see also Sparrow)
- P. domesticus* (house sparrow)
  - arterial blood pressure, 240
  - chromosome number, 1
  - hatching success, 61
  - heart rate, 235
  - oxygen consumption, 222
- P. italiae* (Italian sparrow), 106
- Pasteurella*, Brucellaceae
  - antibiotic activity against, 320, 321, 323
  - parasitism, 505
- Pastinaca sativa* (parsnip), Umbelliferae
  - breeding system, 72
  - chromosome number, 9
  - mineral content, 408, 413
  - parasite, 507
  - respiration rate, 229
  - seed germination, 76
- Patellina*, FORAMINIFERA, 71
- Paternity, putative: phenotypes refuting, 249, 250
- Patience dock (see *Rumex*)
- Pea (see also *Lathyrus*, *Pisum*)
  - nutrition, 168Fn.
  - parasites, 433, 481
  - stem curvature, 308, 309
- cow- (see also *Vigna*)
  - parasites
    - arthropod, 481 483
    - nematode, 494, 496
  - rosary (see *Abrus*)
  - scurf (see *Psoralea*)
- Peach (see also *Prunus*)
  - parasites
    - arthropod, 482, 485
    - nematode, 494
    - viral, 426, 502

- Peafowl, parasites, 493  
 Peanut, 373, 380, 494 (see also *Arachis*)  
 Pear, 368, 481-485 (see also *Pyrus*)  
 Pecan, 496 (see also *Carya*)  
*Pecten* (scallop), FILIBRANCHIA, 66, 109, 237  
*Pectinatella*, PHYLACTOLAEMATA†, 70  
*Pectinophora gossypiella* (pink bollworm), LEPIDOPTERA, 423, 484  
 Pectoral girdle, bones comprising, 160, 161  
*Pediculus* (louse), ANOPLURA, 68, 237  
*P. capitis* (head louse), 4  
*Pellicularia*, Theleporaceae, 509, 510  
*Peltigera aphthosa*, Peltigeraceae, 227  
 Pelvic girdle, bones comprising, 162, 163  
 Pelvic organs (see also specific organs)  
   nerve connections, 137, 138  
   parasites, 489  
 Penguin (see *Aptenodytes*)  
 Penicillia, culture medium for, 536  
*Penicillium*, Aspergillaceae, Moniliaceae  
   antibiotic source, 312, 314  
   chromosome number, 6  
   nutrition, 181Fn.  
   respiration rates, 226  
   temperature for growth, 441  
 Penis, 520  
*Pennisetum glaucum* (pearl millet), Gramineae, 429  
*Pentatrichomonas*, METAMONADINA  
   culture media for, 524, 525  
 Peony (see *Paeonia*)  
 Pepper (see *Capsicum*)  
 Peptides, nutritional requirement for, 168, 177, 179  
*Peranema*, EUGLENOIDINA, 176Fn.  
*Perca flavescens* (yellow perch), PERCOMORPHI, 105  
*P. fluviatilis* (European perch), 2, 107, 236  
*Perdix perdix* (European partridge), GALLIFORMES, 60, 107  
*Peridermium*, Pucciniaceae, 512, 513  
*Peridroma* (cutworm), LEPIDOPTERA, 484  
*Perilla frutescens* (common perilla), Labiatae, 380  
*Peripatus* (peripatus), ONYCHOPHORA†, 4, 70, 223  
*Periplaneta americana* (American cockroach), ORTHOPTERA  
   chromosome number, 4  
   heart rate, 237  
   hemolymph volume, 266  
   life span, 109  
   metamorphosis, 68  
   propagation, 68  
 Periwinkle (see *Littorina*)  
 Periwinkle, Madagascar (see *Vinca*)  
*Peronospora*, Peronosporaceae, 426, 427, 508-510  
*Persea* (persea), Lauraceae, 413, 429  
*P. americana* (American avocado)  
   breeding system, 72  
   chromosome number, 9  
   parasites, 510  
   pollen dispersion, 429  
   pollen life span, 114  
   propagation methods, 74  
   respiration rate, 232  
 Petiole  
   cell sap composition, 404  
   mineral content, 407-409, 413  
*Petromyzon marinus* (sea lamprey), PETROMYZONES  
   blood volumes, 270  
   body weight and length, 105  
   erythrocyte and hemoglobin values, 270  
   life span, 108  
   propagation, 65  
*Petunia hybrida* (common petunia), Solanaceae, 435, 436  
 Peziza, Pezizaceae, 6
- pH  
 acid-base indicators, 545  
 blood, 259-262\*, 263  
 buffer solutions, 543, 544  
 enzymes, 282-287  
 for enzyme activity, 277-281  
 freshwater, 219  
 and nitrogen fixation, 217  
 ocean water, 219, 539  
 soil, 442  
*Phacidium*, Phacidiaceae, 512  
*Phaenicia* (blowfly), DIPTERA, 423  
*P. sericata* (greenbottle fly)  
   diapause, 419  
   dispersion, 423  
   oxygen consumption, 223  
   parasitism, 480  
 Phagocytes, staining methods for, 552  
*Phaseolus* (bean), Leguminosae, 53, 217  
*P. coccineus* (scarlet runner bean), 445  
*P. limensis* (lima bean), 429  
*P. lunatus* (sieva bean), 429  
*P. vulgaris* (kidney bean)  
   breeding system, 72  
   cell sap composition, 404  
   chromosome number, 9  
   light for flowering, 447  
   mineral content, 408, 413  
   nitrogen fixation, 217  
   parasites  
   bacterial, 507  
   fungal, 510  
   viral, 433, 435, 436, 501  
   photosynthesis, 213, 215  
   pollen dispersion, 429  
   respiration rates, 228-230, 232  
   seed germination, 76  
   seed life span, 112  
   soil pH, 442  
*Phasianus* (pheasant), GALLIFORMES, 59  
*P. colchicus* (ring-necked pheasant)  
   blood volumes, 265  
   chromosome number, 1  
   clutch size, 60  
   hatching success, 60, 61  
   life span, 107  
 Pheasant, 491, 493, 548 (see also *Phasianus*)  
 Phellem, cell description, 44  
 Phellogen, cell description, 44  
 Phenotype (see specific blood group system)  
 Phenoxy acids, as growth regulators, 307  
 Phenoxy compounds, as growth regulators, 308, 309  
 Phenyl acids, as growth regulators, 309  
 Phenyl compounds, as growth regulators, 307  
*Phialophora*, Dematiaceae, 518, 520  
*Philaenus leucophthalmus* (meadow spittlebug), HOMOPTERA, 484  
*Philodina* (rotifer), BDELLOIDEA, 70  
*Phlebotomus* (sand fly), DIPTERA, 480, 488, 492  
*Phleospora*, Sphaerioidaceae, 512, 514  
*Phleum pratense* (timothy), Gramineae (see also Timothy)  
   breeding system, 72  
   chromosome number, 8  
   light for flowering, 446  
   mineral content, 405, 411  
   parasite, 507  
   photosynthesis, 213  
   pollen dispersion, 430  
   respiration rate, 230  
   seed germination, 75

† Class



- seed life span, 112, 113  
soil pH, 442
- Phloem  
diseases affecting, 500  
tissue culture, 116
- Phlox* (phlox), Polemoniaceae, 9, 494
- P. divaricata* (Sweet William phlox), 443
- P. paniculata* (summer phlox), 447
- Phoca vitulina* (harbor seal), CARNIVORA  
arterial blood pressure, 240  
heart rate, 235  
life span, 106  
lung ventilation, 220  
oxygen consumption, 221  
propagation, 57  
skeletal system, 159, 161, 163
- Phocaena dalli* (Dall's porpoise), CETACEA, 1
- P. phocaena* (harbor porpoise)  
heart rate, 235  
oxygen consumption, 221  
propagation, 57  
skeletal system, 159, 161, 163
- Phoenix dactylifera* (date palm), Palmae (see also Date palm)  
breeding system, 72  
chromosome number, 8  
mineral content, 405  
photosynthesis, 213  
pollen life span, 114  
propagation method, 73  
respiration rate, 230
- Pholiota*, Agaricaceae, 512
- Phoma*, Sphaeropsidaceae, 510, 511
- Phomopsis*, Sphaeropsidaceae, 510, 513
- Phoradendron* (mistletoe), Loranthaceae, 514, 515
- Phormia regina* (black blowfly), DIPTERA  
dispersion, 423  
hemolymph volume, 266  
nutrition, 168Fn.  
parasitism, 480
- Phormidium*, Oscillatoriaceae, 441
- Phoronis* (phoronid), PHORONIDA††, 70
- Phosphate(s)  
fatty acid source, 375  
properties, 383, 384
- Phosphate esters, carbohydrate: properties, 360-362
- Phospholipid(s)  
and blood coagulation, 257\*  
digestion, 185\*  
metabolism, 197\*  
nutritional requirement for, 167
- Phospholipid content, cells, 398-400
- Photobacterium*, Pseudomonadaceae, 438
- Photosynthesis  
carbon dioxide reduction, 211\*  
efficiency, 216  
rates, 212, 214, 215
- Phrynosoma cornutum* (horned lizard), SAURIA, 62
- Phycomyces*, Mucoraceae, 6, 178Fn., 225
- Phyllophaga* (June beetle), COLEOPTERA, 484, 490
- Phyllosticta*, Sphaeropsidaceae, 509, 510
- Phyllotreta* (flea beetle), COLEOPTERA, 481
- Phylloxera vitifoliae* (grape phylloxera), HOMOPTERA, 484
- Phymatotrichum*, Moniliaceae, 509, 512
- Physalia physalis* (Portuguese man-o'-war), SIPHONOPHORA, 341
- Physalia sting, 341
- Physalospora*, Pleosporaceae, 511, 514
- Physarum*, Physaraceae, 6, 225, 450
- Physeter macrocephalus* (sperm whale), CETACEA, 380
- Physical constants, 579
- Phytolacca americana* (pokeberry), Phytolaccaceae, 348, 454
- Phytophaga destructor* (Hessian fly), DIPTERA, 423, 484
- Phytophthora*, Pythiaceae  
parasitism, 508-510, 512, 514  
spore dispersion, 427
- Pica* (magpie), PASSERIFORMES, 59
- P. nuttalli* (yellow-billed magpie), 60
- P. pica* (black-billed magpie), 1
- Picea* (spruce), Pinaceae, 73, 430, 442
- P. abies* (Norway spruce), 113, 114, 213
- P. breweriana* (Brewer spruce), 515
- P. engelmanni* (Engelmann spruce), 454, 515
- P. glauca* (white spruce)  
chromosome number, 8  
first flowering, 110  
life span, 110  
measurements, 110  
osmotic potential, 455  
parasites, 513, 515  
seed germination, 77  
shade tolerance, 443
- P. mariana* (black spruce), 430, 515
- P. pungens* (Colorado spruce), 215, 515
- P. rubens* (red spruce), 515
- Piedraia*, Myriangiaceae, 516
- Pieris* (cabbage butterfly), LEPIDOPTERA, 69
- P. brassicae* (European cabbageworm), 4, 237
- P. rapae* (imported cabbageworm), 68, 419, 484
- Pigeon (see also *Columba*)  
anesthetic for, 548  
effect of toad toxin on, 334-337  
parasites, 493  
water content, tissues and organs, 403
- Pigment(s)  
in chlorophyll biosynthesis, 210\*  
in feces, 190  
sources, 206, 207  
in urine, 186, 187
- Pigments, respiratory (see Cytochromes)
- Pike (see *Esox*)
- Pilchard, 375
- Pine, 482, 494 (see also *Pinus*)
- Pineal body, hormones, 294, 295
- Pineapple, 494, 496 (see also *Ananas*)
- Pinna*, FILIBRANCHIA, 165Fn.
- Pinnae, mineral content, 405, 411
- Pinnularia*, Naviculaceae, 181Fn.
- Pinus* (pine), Pinaceae (see also Pine)  
osmotic potential, 454  
parasites, 513, 515  
propagation methods, 73  
seed dispersion, 430  
soil pH, 442
- P. caribaea* (slash pine), 112, 113
- P. densiflora* (Japanese red pine), 231
- P. echinata* (shortleaf pine), 430
- P. palustris* (longleaf pine), 8
- P. pinea* (Italian stone pine), 230
- P. radiata* (Monterey pine), 228, 515
- P. strobus* (eastern white pine)  
first flowering, 111  
life span, 111  
measurements, 111  
parasites, 513  
pollen life span, 114  
seed germination, 77  
shade tolerance, 443
- Pinworm (see *Enterobius*)
- Pipa pipa* (Surinam toad), SALIENTIA, 63, 107
- Pipistrellus pipistrellus* (European brown bat), CHIROPTERA, 417

†† Phylum



- Pisum* (pea), Leguminosae, 54, 217
- P. sativum* (garden pea)
- breeding system, 72
  - cell division, 53
  - chromosome number, 9
  - light exposures for development, 444
  - light for flowering, 447
  - mineral content, 408, 413
  - nitrogen fixation, 217
  - osmotic potential, 454
  - parasites, 433, 507, 510
  - pollen life span, 114
  - respiration rates, 228-230, 232
  - seed germination, 76
  - soil pH, 442
  - tissue culture, 116
- P. sativum arvense* (field pea), 217
- Pith, osmotic potential, 455
- Pituophis sayi* (bull snake), SERPENTES, 274
- pK values, weak acids and bases, 544
- Placenta, hormones, 300, 301
- Plaice (*see Pleuronectes*)
- Planaria torva* (flatworm), TRICLADIDA, 5, 110, 223
- Planarian (*see also Dugesia, Planaria*)
- anesthetic for, 549
- Plants (*see also specific genus*)
- number of species, 561
- Plasma
- acid-base balance, 259-262\*
  - with increasing weight and age, 119\*
  - pH, 262\*
- Plasma cell count, bone marrow, 275, 276
- Plasma volumes, 263-265
- Plasmodium*, EUCOCCIDIA, 71, 171Fn., 175Fn., 488
- Plasmopara*, Peronosporaceae, 511
- Platanus occidentalis* (American sycamore), Platanaceae, 454
- Platelet(s), *facing page 276\**
- and blood coagulation, 253\*, 258\*
  - growth, 47
  - life span, 47
  - radiation, 470, 475
- Platelet count, 271
- Platypus (*see Ornithorhynchus*)
- Plecotus auritus* (long-eared bat), CHIROPTERA, 418
- Plectropomus oligacanthus* (sea bass), PERCOMORPHI, 337, 338
- Plenodomus*, Sphaerioidaceae, 509
- Pleospora*, Pleosporaceae, 509
- Pleuronectes platessa* (European plaice), HETEROSOMATA, 107, 236
- Pleurotus*, Agaricaceae, 514
- Plexus(es), 131-138
- parasite, 489
- Plodia interpunctella* (Indian meal moth), LEPIDOPTERA, 484
- Plotosus lineatus* (sea catfish), OSTARIOPHYSI, 336
- Plum, 482 (*see also Prunus*)
- Plutella maculipennis* (diamondback moth), LEPIDOPTERA, 484
- Poa compressa* (Canada bluegrass), Gramineae, 114
- P. pratensis* (Kentucky bluegrass)
- chromosome number, 8
  - light and temperature for flowering, 446
  - mineral content, 405, 411
  - osmotic potential, 454
  - seed germination, 75
  - seed life span, 113
  - soil pH, 442
- P. trivialis* (roughstalk bluegrass), 443
- Podiceps* (grebe), PODICIPEDIFORMES, 148
- Podophrya* (suctorian), SUCTORIDA, 71
- Podosphaera*, Erysiphaceae, 510
- Poikilocytes, *facing page 276\**
- Poison nut (*see Strychnos*)
- Poisonwood (*see Metopium*)
- Pokeberry (*see Phytolacca*)
- Poliomyelitis virus, 432, 499
- Pollen
- dispersion, 428-430
  - life span, 114, 115
  - mineral content, 406
  - tissue culture, 117
- Polyodon spathula* (paddlefish), CHONDROSTEI, 64, 105, 107
- Polyoma virus, 499
- Polypodium virginianum* (rock polypody), Polypodiaceae, 8, 443
- P. vulgare* (common polypody), 213, 231
- Polypody (*see Polypodium*)
- Polyporus*, Polyporaceae, 174Fn., 512-514
- Polypterus* (bichir), CLADISTIA, 153
- Polysaccharides, digestion, 184\*
- Polysiphonia*, Rhodomelaceae, 7, 227, 441
- Polytoma*, PHYTOMONADINA, culture medium for, 528, 529
- Polytomella*, PHYTOMONADINA, 177Fn.
- culture medium for, 529
- Polytrichum*, Polytrichaceae, 7, 227
- Pomoxis annularis* (white crappie), PERCOMORPHI, 64, 105, 107
- P. nigromaculatus* (black crappie), 107
- Pondweed (*see Potamogeton*)
- Pons, 122\*, 123
- Popillia japonica* (Japanese beetle), COLEOPTERA
- chromosome number, 4
  - diapause, 419
  - dispersion, 423
  - hemolymph volume, 266
  - metamorphosis, 68
  - parasitism, 484
  - propagation, 68
- Poplar, 368 (*see also Liriodendron, Populus*)
- Populus* (poplar), Salicaceae
- breeding system, 72
  - parasites, 507, 513, 515
  - pollen dispersion, 430
  - propagation methods, 74
  - respiration rate, 231
- P. alba* (white poplar), 214, 454
- P. deltoides* (eastern poplar), 430, 443, 454
- P. suaveolens* (Mongolian poplar), 114
- P. tremula* (European aspen), 408
- P. tremuloides* (quaking aspen)
- chromosome number, 9
  - first flowering, 111
  - life span, 111
  - measurements, 111
  - osmotic potential, 455
  - seed germination, 77
  - seed life span, 112
  - soil pH, 442
- Poppy (*see Papaver*)
- Porcellio laevis* (sowbug), ISOPODA, 481
- Porgy (*see Pagellus, Pagrus*)
- Poria*, Polyporaceae, 512, 514
- Porphyra*, Bangiaceae
- chromosome number, 7
  - photosynthesis, 213
  - respiration rate, 227
  - temperature tolerances, 441
- Porpoise (*see Phocaena*)
- Porthetria dispar* (gypsy moth), LEPIDOPTERA, 423, 484

- Posture  
 brain involvement, 123  
 effect on blood pressure, 242  
 effect on heart rate, 234
- Potamogeton crispus* (curly pondweed), Potamogetonaceae, 454
- Potamon dehaanii* (river crab), DECAPODA, 4
- Potamotrygon motoro* (freshwater stingray), BATOIDEI, 338
- Potato (*see also Solanum*)  
 parasites  
 arthropod, 482, 483  
 fungal, 427  
 nematode, 494  
 viral, 426, 433-435, 502
- Potato, sweet, 494
- Poultry, 516, 518 (*see also* specific genus)
- Prenatal development (*see* table of contents, page v)
- Pressure, blood (*see* Blood pressures)
- Pressure, gases in respiratory media, 219
- Pressure, hydrostatic: sea water, 539
- Pressure, standard atmospheric, 579
- Pressure-depth gradient, sea water, 541
- Primitive streak, developmental stages, 82, 84, 86, 88, 91
- Primrose (*see* *Oenothera*)
- Procyon cancrivorus* (crab-eating raccoon), CARNIVORA, 221
- P. lotor* (raccoon), 57, 106
- Prodenia euidania* (southern armyworm), LEPIDOPTERA, 237, 267
- Proerythroblast count, bone marrow, 275, 276
- Promyelocyte count, bone marrow, 275, 276
- Propagation (*see* table of contents, page v)
- Propagation methods, cultivated plants, 73, 74
- Prophase, effect of compounds on, 53
- Prosencephalon, regions and functions, 123
- Prostate  
 nerve connections, 133, 135, 138  
 permissible radionuclide concentration in, 459, 460  
 tissue growth, 48  
 water content, 401
- Protaminobacter*, Pseudomonadaceae, 218\*
- Protein(s)  
 in acid-base, blood, 259-262\*  
 aerobic oxidation, 204\*  
 catabolism, 201\*  
 digestion, 183\*  
 histochemical test for, 559  
 metabolic interrelationships, 203\*  
 metabolism, 197\*  
 in nitrogen cycle, 218\*  
 nutritional requirement for, 168, 177, 179  
 properties, 388, 389
- Protein content  
 cells, 398-400  
 urine, 186
- Proteus*, Enterobacteriaceae  
 antibiotic activity against, 320-322  
 generation time, 52  
 and nitrogen cycle, 218\*  
 parasitism, 505  
 temperature for growth, 438  
 thermal death time, 439
- Prothrombin time, 326, 327
- Protococcus*, Pleurococcaceae, 441
- Protoparce quinque maculata* (tomato hornworm), LEPIDOPTERA, 484
- Protoplasmic streaming, 448, 449
- Protopterus* (lungfish), DIPNOI, 153
- P. aethiopicus* (East African lungfish), 222
- P. annectans* (West African lungfish), 2, 108
- Protozoa (*see also* specific genus)  
 culture media for, 523-529
- Provitamins, properties, 396, 397
- Prunus*, Rosaceae, 348
- P. americana* (American plum), 112
- P. amygdalus* (almond) (*see also* Almond)  
 chromosome number, 9  
 mineral content, 408, 413  
 parasite, 510  
 pollen life span, 114  
 propagation method, 74  
 respiration rates, 228, 231
- P. domestica* (garden plum)  
 breeding system, 72  
 chromosome number, 9  
 mineral content, 408, 413  
 parasites, 507, 510  
 pollen life span, 114  
 propagation methods, 74  
 respiration rates, 228, 232
- P. laurocerasus* (laurel cherry)  
 photosynthesis, 213, 215  
 respiration rates, 229, 231
- P. persica* (peach) (*see also* Peach)  
 breeding system, 72  
 chromosome number, 9  
 mineral content, 409, 413  
 parasites, 502, 507, 510  
 photosynthesis, 215  
 pollen life span, 114  
 propagation method, 74  
 respiration rates, 228, 232  
 soil pH, 442
- P. serotina* (black cherry)  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 seed germination, 77  
 shade tolerance, 443
- Psallus seriatus* (cotton fleahopper), HEMIPTERA, 424
- Pseudaletia unipuncta* (armyworm), LEPIDOPTERA, 484
- Pseudemys floridana peninsularis* (peninsular turtle), CHELONIA, 62
- P. scripta elegans* (red-eared turtle)  
 acid-base, blood, 261  
 arterial blood pressure, 240  
 blood volumes, 265, 269  
 erythrocyte and hemoglobin values, 269  
 life span, 107
- P. terrapen rugosa* (Cuban freshwater turtle), 236
- Pseudochis porphyriacus* (Australian black snake), SERPENTES, 332
- Pseudococcus citri* (mealybug), HOMOPTERA, 484
- Pseudomonas*, Pseudomonadaceae  
 antibiotic activity against, 320-323  
 generation time, 52  
 and nitrogen cycle, 218\*  
 nutrition, 178Fn., 181Fn.  
 parasitism, 505-508  
 protein source, 389  
 respiration rate, 225  
 temperature for growth, 438  
 thermal death time, 439
- Pseudoperonospora*, Peronosporaceae, 426, 509
- Pseudopeziza*, Dermaceae, 510
- Pseudoplea*, Hyalodictyae, 511
- Pseudopleuronectes americanus* (winter flounder), HETEROSOMATA, 64
- Pseudosuccinea* (freshwater snail), BASOMMATOPHORA, 492
- Pseudotsuga taxifolia* (Douglas fir), Pinaceae, 430 (*see also* Fir, Douglas)

- Psila rosae* (carrot rust fly), DIPTERA, 423, 484  
 Psittacosis virus, 432, 499  
*Psoralea tenuiflora* (slim flower scurf pea), Leguminosae, 456  
*Psorophora* (rice-field mosquito), DIPTERA, 423  
*Psoroptes equi ovis* (sheep-scab mite), ACARI, 478  
*Psyllia pyricola* (pear psylla), HOMOPTERA, 484  
*Pterois antennata* (thread-finned zebra fish), SCLEROPAREI, 338  
*P. lunulata* (tiger fish), 338  
*P. volitans* (turkey fish), 338  
 Puberty, age at, 57  
*Puccinia*, Pucciniaceae  
   chromosome number, 6  
   parasitism, 508, 509, 511, 512  
   respiration rate, 226  
   spore dispersion, 426, 427  
   temperature for growth, 441  
*Pucciniastrum*, Melampsoraceae, 513, 514  
 Puck's N16 culture medium for animal tissue, 533  
 Puck's solution for animal tissue culture, 530  
*Pueraria thumbergiana* (kudzu), Leguminosae, 216  
 Puffer, 338, 339  
*Pulex* (flea), SIPHONAPTERA, 490  
*P. irritans* (human flea), 480  
 Pulvini, osmotic potential, 455  
 Pumpkin (*see Cucurbita*)  
 Punkie (*see Culicoides*)  
 Purine(s)  
   biosynthesis, 208\*  
   catabolism, 202\*  
   effect on cell division, 53  
   metabolism, 200  
   in nitrogen cycle, 218\*  
   in nucleoprotein catabolism, 201\*  
   nutritional requirement for, 171, 179  
 Purine content  
   feces, 190  
   urine, 186, 187  
*Pyrausta nubilalis* (European corn borer), LEPIDOPTERA  
   diapause, 419  
   dispersion, 423, 424  
   parasitism, 484  
*Pyrenochaeta*, Sphaerioidaceae, 508, 518  
*Pyrenophora*, Pseudosphaeriaceae, 509  
 Pyrimidines  
   biosynthesis, 209\*  
   catabolism, 202\*  
   metabolism, 199  
   in nucleoprotein catabolism, 201\*  
   nutritional requirement for, 171, 173, 179  
*Pyrodinium* (dinoflagellate), DINOFLAGELLATA, 341, 342  
*Pyrola rotundifolia* (European pyrola), Ericaceae, 455  
*Pyrus communis* (pear), Rosaceae (*see also* Pear)  
   breeding system, 72  
   chromosome number, 9  
   mineral content, 409, 413  
   parasites, 507, 510  
   pollen life span, 114  
   propagation methods, 74  
   respiration rate, 232  
   soil pH, 442  
*Pythium*, Pythiaceae, 508  
 Quahog (*see Mercenaria*)  
 Quail, 491, 493 (*see also* *Colinus*)  
 Queen snake (*see Natrix*)  
 Quercitron, 368  
*Quercus* (oak), Fagaceae, 72, 74, 513-515 (*see also* Oak)  
*Q. alba* (white oak)  
   chromosome number, 9  
   first flowering, 111  
   life span, 111  
   measurements, 111  
   osmotic potential, 454  
   respiration rate, 232  
   seed germination, 77  
   shade tolerance, 443  
   soil pH, 442  
*Q. coccifera* (kermes oak), 229, 231  
*Q. coccinea* (scarlet oak), 114, 454  
*Q. ilex* (holly oak), 213  
*Q. lyrata* (overcup oak), 213  
*Q. robur* (English oak), 409  
*Q. rubra* (eastern red oak), 215  
*Q. velutina* (black oak), 413  
 Quince, 482, 485  
 Rabbit (*see also* *Oryctolagus*)  
   anesthetic for, 548  
   cell chemical constituents, 399, 400  
   cells, staining method for, 552  
   developmental stages, 82  
   effect of toad toxins on, 334, 336  
   enzyme source, 388, 389  
   gestation time, 82  
   heart fibers, diameter, 47  
   parasites  
     arthropod, 480  
     bacterial, 505  
     helminthic, 490, 493  
     rickettsial, 503  
     viral, 499, 500  
   protein source, 388, 389  
   radiation effect on, 474, 475  
   sperm survival time, 49  
   tissue growth, 47-49  
   tissue regeneration, 48, 49  
   water content, tissues and organs, 402  
 Rabies virus, 432, 499  
 Raccoon (*see also* *Procyon*)  
   anesthetic for, 548  
   parasites, 493  
 Races, human: body weight and height, 93, 94  
 Radiation  
   late effects, 468-476  
   permissible occupational exposure, 457-467  
 Radionuclides  
   permissible body concentration, 458-467  
   radiation emitted, 546, 547  
 Radish (*see* *Raphanus*)  
*Radopholus oryzae* (rice-root nematode), TYLENCHIDA, 496  
*R. similis* (burrowing nematode), 496  
 Ragweed (*see* *Ambrosia*)  
*Raillietina cesticillus* (broad-headed tapeworm), CYCLOPHYLLIDEA, 492  
*Raja* (skate), BATOIDEI  
   acid-base, blood, 261  
   endocrine tissue, 155, 157  
   heart rate, 236  
*R. erinacea* (little skate), 64, 270  
*R. maculata* (skate), 108  
*R. meerdervoortii* (skate), 2  
*R. punctulata* (skate), 240  
*R. rhina* (longnose skate), 265  
 Rami, nerve connections, 130\*, 131, 132, 134  
*Ramosia tipuliformis* (currant borer), LEPIDOPTERA, 484



- Rana* (frog), SALIENTIA  
 capillary blood pressure, 242, 243  
 cell division, 53  
 digestive enzymes, 141  
 endocrine system, 152, 154
- R. catesbeiana* (American bullfrog)  
 arterial blood pressure, 240  
 blood volumes, 265, 270  
 body length, 103, 104  
 chromosome number, 2  
 erythrocyte and hemoglobin values, 270  
 life span, 107  
 propagation, 63
- R. esculenta* (edible frog), 222
- R. pipiens* (leopard frog) (see also Frog, leopard)  
 arterial blood pressure, 240  
 body length, 104  
 chromosome number, 2  
 heart rate, 236  
 life span, 107  
 propagation, 63  
 sterol source, 385
- Rape (see *Brassica*)
- Raphanus* (radish), Cruciferae, 430
- R. raphanistrum* (wild radish), 229, 231
- R. sativus* (garden radish)  
 breeding system, 72  
 chromosome number, 9  
 light for flowering, 447  
 mineral content, 409, 413  
 parasites, 507  
 pollen dispersion, 430  
 respiration rates, 228, 229  
 seed germination, 76  
 soil pH, 442
- Raspberry viruses, 426
- Rat (see also *Rattus*)  
 anesthetic for, 548  
 cell chemical constituents, 399  
 cell growth, 47  
 developmental stages, 82, 84-86  
 gestation time, 82  
 nutrition, 168Fn.  
 parasites  
   fungal, 516, 520  
   helminthic, 486, 491, 492  
   protozoan, 488, 493  
 parathyroid, 154  
 radiation effect on, 472-474  
 tissue growth, 46-49  
 tissue regeneration, 49  
 water content, tissues and organs, 402
- Rat, wood, 493
- Ratfish (see *Chimaera*)
- Rattlesnake, 388 (see also *Crotalus*)
- Rattus* (rat), RODENTIA (see also Rat)  
 acid-base, blood, 260  
 capillary blood pressure, 242  
 cell division, 53  
 digestive enzymes, 140  
 leukocyte counts, 274  
 oxygen consumption, 221
- R. norvegicus* (Norway rat)  
 arterial blood pressure, 240  
 blood volumes, 264, 269  
 body surface area constants, 121  
 body weight, 95, 96, 121  
 chromosome linkage groups, 18, 19  
 chromosome number, 1, 18  
 clotting time, 326, 327  
 erythrocyte values, 269  
 heart rate, 235  
 hemoglobin values, 269  
 life span, 106  
 lung ventilation, 220  
 mutations, 18, 19  
 platelet count, 271  
 propagation, 57  
 skeletal system, 159, 161, 163  
 strains, body weight, 95, 96
- Raven (see *Corvus*)
- Ray (see also *Aetobatus*, *Torpedo*)  
 anesthetic for, 549  
 sting- (see Stingray)
- Rectum  
 nerve connections, 132  
 parasite, 489
- Red blood cells, facing page 276\* (see also Erythrocytes)
- Red fish (see *Theleleota*)
- Redox potential, 545
- Reduvius personatus* (masked hunter), HEMIPTERA, 419
- Redwood (see *Sequoia*)
- Refractive index  
 fats, 380  
 fatty acids, 371, 373, 375, 377, 379  
 hormones, 292, 296, 298, 300  
 oils, 380  
 waxes, 382
- Regeneration, tissue, 46-49
- Rehmiellosis*, Hyalodidymae, 512
- Reovirus, 499
- Reproduction (see also table of contents, page v)  
 radiation effect on, 469, 470
- Resin  
 fatty acids source, 377  
 glycoside source, 368
- Respiration rate (see also table of contents, page vii)  
 during hibernation, 417, 418
- Respiratory media, 219
- Respiratory quotient, 225-232
- Respiratory rate, 220
- Respiratory reflexes, brain involvement, 123, 127
- Respiratory tract, 498, 499
- Reticulitermes flavipes* (eastern subterranean termite), ISOPTERA, 484
- Reticulocytes, facing page 276\*  
 chemical composition, 400
- Reticuloendothelial phagocytes, staining methods for, 552
- Reticuloendothelium, 489, 493
- Reticulum cell count, bone marrow, 275, 276
- Retinal detachment, 12
- Retinitis pigmentosa, 12
- Rh-Hr blood group system, 247, 248  
 distribution in various populations, 252  
 heredity, 249, 250
- Rhabditis* (free-living roundworm), RHABDITIDA, 4, 109
- Rhynchospora obscura* (New Guinea sugarcane weevil), COLEOPTERA, 423
- Rheum* (rhubarb), Polygonaceae, 413
- R. officinale* (medicinal rhubarb), 9
- R. palmatum* (sorrel rhubarb), 213, 409
- R. rhaponticum* (garden rhubarb)  
 parasites, 507  
 photosynthesis, 215  
 respiration rate, 231  
 seed germination, 76
- Rhagoletis pomonella* (apple maggot), DIPTERA, 423, 485
- Rhinencephalon, regions and functions, 123
- Rhinolophus ferrum-equinum* (greater horseshoe bat), CHIROPTERA, 418
- R. hipposideros* (lesser horseshoe bat), 418
- Rhinosporidium*, Endomycetales\*, 520
- Rhipicephalus* (tick), ACARI, 492

• Order



- Rhipicephalus sanguineus* (brown dog tick), 478  
*Rhizina*, Rhizinaceae, 513  
*Rhizobium*, Rhizobiaceae, 52, 216-218\*, 438  
*Rhizoctonia*, Mycelia sterilia<sup>•</sup>, 508  
*Rhizoglyphus echinopus* (bulb mite), ACARI, 481  
*Rhizopus*, Mucoraceae  
   nutrition, 173Fn.  
   parasitism, 518  
   respiration rate, 225  
   temperature for growth, 440, 441  
   thermal death time, 440, 441  
*Rhodinus* (assassin bug), HEMIPTERA, 492  
*Rhododendron* (rhododendron), Ericaceae  
   chromosome number, 9  
   light for flowering, 447  
   pollen life span, 114  
   propagation methods, 74  
   shade tolerance, 443  
*R. brachycarpum* (Fujiyama rhododendron), 213  
*R. fargesii* (Père Farges' rhododendron), 231  
*R. obtusum amoenum* (amoena azalea), 442  
*Rhodomicrobium*, Hypomicrobiaceae, 217  
*Rhodopseudomonas*, Athiorhodaceae, 217  
*Rhodospirillum*, Athiorhodaceae, 217  
Rhombencephalon, regions and functions, 123  
*Rhombomys* (great gerbil), RODENTIA, 493Fn.  
Rhubarb (*see Rheum*)  
*Rhus toxicodendron* (poison ivy), Anacardiaceae, 348  
*R. vernix* (poison sumac), 348  
*Rhynchosporium*, Moniliaceae, 509  
*Rhytisma*, Phacidiaceae, 512  
Rib marrow, differential cell counts, 275  
Ribbon worm (*see Cerebratulus*)  
*Ribes* (currant), Saxifragaceae, 9, 72, 413 (*see also*  
 Currant)  
*R. americanum* (American black currant), 443  
*R. aureum* (golden currant), 507  
*R. glutinosum* (nutmeg currant), 114  
*R. nigrum* (European black currant), 409  
*R. rubrum* (northern red currant), 232  
*R. sativum* (common red currant), 74  
Ribosome, chemical constituents, 398, 400  
Ribs, number, 158, 159  
*Riccia*, Ricciaceae, 7, 227  
Rice (*see also Oryza*)  
 parasites, 494, 496, 501  
 wax source, 382  
*Ricinus communis* (castor bean), Euphorbiaceae, 348  
 349, 380  
Rickettsia (*see specific genus*)  
*Rickettsia*, Rickettsiaceae, 438, 439, 503  
*Ricoleisia*, Chlamydiaceae, 503  
Rift Valley fever virus, 432, 499  
Rinderpest virus, 499  
Ringer's solution for animal tissue culture, 529  
Ringhals (*see Hemachatus*)  
RNA, histochemical test for, 558  
RNA codewords for amino acids, 43  
RNA content, cells, 398-400  
Robin (*see Turdus*)  
*Robinia pseudoacacia* (black locust), Leguminosae, 454  
*Rochalimaea*, Rickettsiaceae, 503  
Rockweed (*see Fucus*)  
*Romalea* (grasshopper), ORTHOPTERA, 69  
Root(s)  
 cell division, 53, 54  
 cell sap composition, 404  
 depth: effect on osmotic potential, 456  
 diseases affecting, 500-502, 507-514  
 mineral content, 405-409, 411-413  
 osmotic potential, 454, 455  
 parasites, 481-485, 494-497  
 respiration rates, 228, 229  
 sterol source, 386  
 tissue culture, 115, 116  
 tissue culture medium for, 538  
 toxins, 344-349  
Root graft, 73  
*Rosa* (rose), Rosaceae (*see also* Rose)  
 breeding system, 72  
 chromosome number, 9  
 propagation methods, 74  
 respiration rates, 231, 232  
 soil pH, 442  
 tissue culture, 116  
*R. centifolia* (cabbage rose), 409  
*R. multiflora* (Japanese rose), 507, 508  
Rose, 482, 496 (*see also Rosa*)  
 black Christmas (*see Helleborus*)  
*Rosellinia*, Sphaeriaceae, 514  
Rotifer (*see specific genus*)  
*Rotylenchus* (spiral nematode), TYLENCHIDA, 494  
Roundworm  
 dog (*see Toxascaris*)  
 free-living (*see Rhabditis*)  
 large (*see Ascaris, Parascaris*)  
Rous sarcoma virus, 499  
*Rumex acetosa* (garden sorrel), Polygonaceae, 116, 451  
*R. patientia* (patience dock), 454  
Rush (*see Equisetum*)  
Rutabaga, 494  
Rye (*see also Secale*)  
 parasites  
   arthropod, 482, 484  
   fungal, 427  
   nematode, 494  
   sterol source, 386  
  
*Saccharomyces*, Saccharomycetaceae  
 antibiotic activity against, 320  
 chromosome number, 6  
 nutrition, 168Fn., 171Fn., 173Fn.  
 respiration rate, 225  
*Saccoglossus* (acorn worm), ENTEROPNEUSTA†, 69  
Safflower (*see Carthamus*)  
Saffron, 368  
*Sagartia elegans* (sea anemone), ACTINIARIA, 341  
*Saissetia oleae* (black scale), HOMOPTERA, 423  
Salamander (*see also Ambystoma, Salamandra*)  
 tiger, anesthetic for, 548  
*Salamandra* (salamander), CAUDATA, 147, 154, 236  
Salinity, sea water, 539, 540  
Saliva, 498, 499  
Salivary glands  
 cell division, 54  
 enzymes, 139-141  
 parasite, 498  
 tissue growth, 48  
 tissue regeneration, 48  
*Salix* (willow), Salicaceae (*see also* Willow)  
 breeding system, 72  
 parasites, 508, 514, 515  
 shade tolerance, 443  
 soil pH, 442  
*S. alba* (white willow), 9, 454  
*S. glauca* (gray-leaf willow), 213, 231  
*S. gracistyla* (big catkin willow), 114  
*S. herbacea* (pygmy willow), 229  
*S. humilis* (prairie willow), 445  
*S. nigra* (black willow), 111  
*S. viminalis* (basket willow), 409  
*Salmo*, ISOSPONDYLI  
 adrenals, 155

• Order  
 † Class

- arterial blood pressure, 240
- developmental stages, 91, 92
- parasite, 492
- subclavian vein, 149
- S. gairdneri* (rainbow trout), 157, 261
- S. salar* (Atlantic salmon)
  - body weight and length, 105
  - chromosome number, 2
  - life span, 108
  - propagation, 64
- S. trutta* (brown trout) (*see also* Trout, brown)
  - body weight and length, 105
  - chromosome number, 2
  - heart rate, 236
  - life span, 108
  - oxygen consumption, 222
  - propagation, 64
- Salmon, 389, 400 (*see also* *Oncorhynchus*, *Salmo*)
- Salmonella*, Enterobacteriaceae
  - antibiotic activity against, 320-323
  - chemical composition, 400
  - generation time, 52
  - parasitism, 505, 506
  - temperature for growth, 438
  - thermal death time, 439
- Salpa* (tunicate), SALPIDA, 69
- Salsify (*see* *Tragopogon*)
- Salt solutions
  - for animal tissue culture, 529, 530
  - for plant tissue culture, 538
- Salvelinus* (trout), ISOSPONDYLI, 492
- S. fontinalis* (eastern brook trout) (*see also* Trout, brook)
  - blood volumes, 270
  - body weight and length, 105
  - chromosome number, 2
  - erythrocyte and hemoglobin values, 270
  - propagation, 64
- S. namaycush* (lake trout), 108
- Sambucus nigra* (European elder), Caprifoliaceae, 213, 214
- Samia cynthia* (cynthia moth), LEPIDOPTERA, 4, 419
- Sand dollar (*see* *Echinarachnius*)
- Sanninoidea exitiosa* (peach tree borer), LEPIDOPTERA, 485
- Sap
  - chemical composition, 404
  - fatty acid source, 379
  - toxins, 344, 346-349
- Saperda candida* (roundheaded apple tree borer), COLEOPTERA, 485
- Saprolegnia*, Saprolegniaceae, 6
- Sarcina*, Micrococcaceae, 177Fn.
- Sarcoptes scabiei* (itch mite), ACARI, 478
- Sardines, 375, 380
- Sardinops caerulea* (Pacific American sardine), ISOSPONDYLI, 380
- Sarsaparilla, 368
- Saturated fatty acids
  - in fats and oils, 381
  - properties, 370-373
- Sawfly (*see* *Cephus*, *Diprion*)
- Saxidomus* (butter clam), EULAMELLIBRANCHIA, 341
- Scale
  - black (*see* *Saissetia*)
  - brown (*see* *Coccus*)
  - San Jose (*see* *Aspidiotus*)
  - scurfy (*see* *Chionaspis*)
  - wax (*see* *Ceroplastes*)
- Scallop (*see* *Pecten*)
- Scammony, 368
- Scarus* (parrot fish), PERCOMORPHI, 155, 337, 338
- Sceloporus graciosus* (sagebrush lizard), SAURIA, 62, 107
- S. spinosus* (spiny fence lizard), 2
- Scenedesmus*, Coelastraceae, 181Fn.
- Schistocerca gregaria* (desert locust), ORTHOPTERA, 485
- Schistosoma haematobium* (human blood fluke), DIGENEA, 5, 488
- S. japonicum* (blood fluke), 488
- S. mansoni* (blood fluke), 488
- Schizophyllum*, Tricholomataceae, 181Fn.
- Schizosaccharomyces*, Schizosaccharomycetaceae, 6, 226
- Schizothaerus nuttalli* (gaper clam), EULAMELLIBRANCHIA, 341
- Schizoxylon*, Ostropaceae, 512
- Sciurus carolinensis* (gray squirrel), RODENTIA
  - chromosome number, 1
  - heart rate, 235
  - life span, 106
  - propagation, 57
- Sclereid, description, 45
- Scleroderris*, HELOTIALES, 512
- Sclerosis, diffuse cerebral, 12
- Sclerospora*, Peronosporaceae, 511
- Sclerotinia*, Sclerotiniaceae
  - parasitism, 509, 510, 513
  - spore dispersion, 426
- Sclerotium*, Mycelia sterilia<sup>o</sup>, 510
- Scolytus multistriatus* (smaller European elm bark beetle), COLEOPTERA, 423
- Scomber scombrus* (Atlantic mackerel), PERCOMORPHI
  - blood volumes, 270
  - erythrocyte and hemoglobin values, 270
  - life span, 108
  - oxygen consumption, 222
  - propagation, 64
- Scorpaena guttata* (California scorpion fish), SCLEROPAREL, 338
- S. plumieri* (sculpin), 338
- S. porcus* (hogfish), 338
- Scorpaenopsis diabolus* (scorpion fish), SCLEROPAREL, 338
- Scorpion (*see* *Centrurus*)
- Scorpion fish (*see* *Apistus*, *Notesthes*, *Scorpaena*, *Scorpaenopsis*)
- Scorpion-fish sting, 338
- Scorzonera hispanica* (black salsify serpentroot), Compositae, 116
- Screwworm (*see* *Chrysomya*, *Cochliomyia*)
- Sculpin (*see* *Scorpaena*)
- Scutigerebella immaculata* (garden symphyliid), SYMPHYLA†, 109, 485
- Scyliorhinus* (cat shark), SELACHII, 157
- Scyphu* (marine sponge), SYCONOSA, 5, 70
- Scyphomedusa (*see* *Aurelia*)
- Sea anemone, 386 (*see also* specific genus)
- Sea-anemone sting, 341
- Sea cucumber (*see* specific genus)
- Sea-cucumber poisoning, 340
- Sea hare (*see* *Aplysia*)
- Sea horse (*see* *Hippocampus*)
- Sea lemon (*see* *Doris*)
- Sea lettuce (*see* *Ulva*)
- Sea snake (*see* *Enhydrina*)
- Sea spider (*see* *Nymphon*)
- Sea squirt (*see* *Ascidia*, *Ciona*, *Molgula*)
- Sea urchin (*see* specific genus)
- Sea-urchin sting, 340
- Sea water, artificial, 541, 542
- Sea water, natural, 539-541
- Seal (*see* *Phoca*)
- Seaweed (*see* *Gigartina*)
  - culture medium for, 536, 537
- Sebaceous glands, 477

• Order  
† Class

- Sebastiania fruticosa* (sebastiania), Euphorbiaceae, 373  
*Secale cereale* (rye), Gramineae 430, 453, (see also Rye)  
 Sedimentation coefficient for  
   enzymes, 282-287, 388, 389  
   hormones, 290, 292, 388, 389  
   proteins, 388, 389
- Seed(s)  
   disease affecting, 500  
   dispersion, 428-430  
   fatty acid source, 373, 375, 377, 379-381  
   germination, 75-77  
   glycoside source, 368  
   life span, 111-113  
   mineral content, 406-409, 412, 413  
   parasites, 481  
   propagation by, 73, 74  
   protein source, 388  
   respiration rates, 228  
   sterol source, 385, 386  
   toxins, 344-349
- Seed plants, cell types, 44-46  
 Segmented cells, bone marrow, 275, 276  
*Selaginella selaginoides* (spike moss), Selaginellaceae, 8  
*Selasphorus sasin* (Allen's hummingbird), APODIFORMES, 22, 418  
 Seminal vesicles, nerve connections, 133, 135  
*Senecio* (groundsel), Compositae, 349  
 Sensory reflexes, brain involvement, 123, 124, 127-129  
*Sepia officinalis* (cuttlefish), DIBRANCHIA, 4, 223, 237  
*Septoria*, Sphaeropsidaceae, 509-511, 513  
*Sequoia gigantea* (giant sequoia), Taxodiaceae  
   chromosome number, 8  
   first flowering, 111  
   life span, 111  
   measurements, 111  
   seed germination, 77  
   shade tolerance, 443
- S. sempervirens* (redwood), 514  
*Serinus* (canary), PASSERIFORMES, 59  
*S. canarius* (canary)  
   arterial blood pressure, 240  
   heart rate, 235  
   life span, 107  
   lung ventilation, 220  
   oxygen consumption, 222
- Serpentroot (see *Scorzonera*)  
*Sesamum indicum* (Oriental sesame), Pedaliaceae, 380  
*Setaria equinum* (filarial worm), SPIRURIDA, 223  
*Setaria italica* (foxtail millet), Gramineae, 453  
 Sex-linked mutations, 11, 12, 16, 19  
 Sexual differences in  
   arterial blood pressure, 239  
   body height, 93, 94  
   body length, 102-104  
   body weight, 93-102  
   erythrocyte and hemoglobin values, 268
- Sexual maturity, 62, 63, 66  
 Shad (see *Clupanodon*)  
 Shade tolerance, spermatophytes, 443  
 Shark (see also *Carcharhinus*, *Scyliorhinus*, *Sphyrna*)  
   anesthetic for, 549  
   fatty acid source, 373, 375
- Sheep (see also *Ovis*)  
   anesthetics for, 547, 548  
   effect of plant toxins on, 344, 347  
   fatty acid source, 373, 379-381  
   gestation time, 82  
   hormone source, 388, 389  
   parasites  
     arthropod, 477-480  
     bacterial, 504-506  
     fungal, 516, 518  
     helminthic, 486, 490, 491, 493  
     protozoan, 493  
     ricketsial, 503  
     viral, 498, 499  
   prenatal development, 82  
   protein source, 388, 389  
   sterol source, 385  
   water content, tissues and organs, 402, 403  
   mountain, 504
- Shellfish, 385, 386  
 Shellfish poisoning, 340-342  
*Shigella*, Enterobacteriaceae  
   antibiotic activity against, 320, 321, 323  
   generation time, 52  
   parasitism, 506  
   temperature for growth, 438  
   thermal death time, 439
- Shipworm (see *Teredo*)  
 Shope papilloma virus, 499  
 Shrew (see *Sorex*)  
 Sieve cell, description, 45  
 Sieve-tubes  
   cell description, 45, 46  
   osmotic potential, 455
- Silkworm (see *Bombyx*)  
 Silkworm jaundice virus, 499  
 Silverfish (see *Lepisma*)  
*Simulium* (blackfly), DIPTERA, 423, 480, 486  
*Siphona irritans* (horn fly), DIPTERA, 480  
*Siren* (siren), CAUDATA, 147, 154  
*Sistrurus catenatus* (eastern massasauga), SERPENTES, 328  
*Sitophilus oryza* (rice weevil), COLEOPTERA, 423, 485  
 Skate (see *Raja*)  
 Skeletal system, 158-163  
 Skin  
   growth, 49  
   mineral retention in, 192\*, 193-195  
   parasites  
     arthropod, 477-480  
     fungal, 516, 518, 520  
     helminthic, 487, 489  
     protozoan, 489  
     viral, 498-500  
   permissible radiation exposure for, 457, 464  
   radiation effect on, 468, 469, 471, 472  
   regeneration, 49  
   sterol source, 385  
   water content, 401-403
- Skink (see *Chalcides*, *Eumeces*)  
 Skipjack (see *Katsuwonus*)  
 Skull, bones comprising, 158, 159  
 Skunk, 520  
 Sleep, heart rate during, 234  
 Slowworm (see *Anguis*)  
 Slug (see *Cladius*)  
 Smallpox virus, 432, 500  
 Smelt (see *Osmerus*)  
*Sminthurus viridis* (lucerne flea), COLLEMBOLA, 485  
 Snail, 385, 488  
   freshwater (see *Fossaria*, *Galba*, *Lymnaea*, *Pseudosuccinea*)  
   land (see *Helix*)  
   river (see *Goniobasis*)
- Snake (see specific genus)  
 Snapdragon (see *Antirrhinum*)  
 Snapper (see *Aprion*, *Gnathodentex*, *Lutjanus*)  
 Snapweed (see *Impatiens*)  
 Soil  
   effect on transpiration rates, 452  
   life span of seeds in, 112, 113  
   pH for plant growth, 442  
   water content: effect on osmotic potential, 455



- Solanum*, Solanaceae, 370
- S. lycopersicum* (nightshade), 232
- S. melongena* (eggplant), 76, 113, 443
- S. nigrum* (black nightshade), 113
- S. tuberosum* (potato) (*see also* Potato)
- breeding system, 72
  - cell sap composition, 404
  - chromosome number, 9
  - light for flowering, 447
  - mineral content, 409, 413
  - parasites
    - bacterial, 508
    - fungal, 510
    - viral, 433, 434, 502
  - photosynthesis, 213, 215
  - pollen life span, 114
  - respiration rates, 231
  - seed life span, 112
  - soil pH, 442
  - tissue culture, 116
- Solar energy for photosynthesis, 216
- Solenogaster (*see Neomenia*)
- Solidago* (goldenrod), Compositae, 454, 456
- Sorex araneus* (European shrew), INSECTIVORA, 1, 57
- S. cinereus* (gray shrew)
- body surface area constants, 121
  - body weight, 121
  - heart rate, 235
  - oxygen consumption, 221
  - skeletal system, 159, 161, 163
- S. palustris* (water shrew), 106
- Sorghum vulgare sudanense* (Sudan grass), Gramineae, 453
- Sorrel (*see Rumex*)
- Soun pest (*see Eurygaster*)
- Sowbug (*see Porcellio*)
- Soybean (*see also Glycine*)
- parasites, 483, 494
  - sterol source, 385, 386
- Sparassis*, Thelephoraceae, 513
- Sparrow, 82 (*see also Passer*)
- song (*see also Melospiza*)
  - anesthetic for, 548
- Spawning seasons, 64, 65
- Specific gravity
- fats, 380
  - fatty acids, 370, 372, 374, 376, 378
  - oils, 380
  - waxes, 382
- Specific rotation
- aglycones, 369, 371
  - aldaric acids, 359
  - alditols, 356
  - aldonic acids, 358
  - aldoses, 351, 352
  - amino acids, 392, 393
  - amino sugars, 355
  - carbohydrate phosphate esters, 360-362
  - carbohydrates, 351-366
  - glycosides, 368, 370
  - inositols, 356, 357
  - ketoses, 352, 353
  - monosaccharides, 351-356
  - oligosaccharides, 364-366
  - phosphatides, 383, 384
  - provitamins, 397
  - sphingolipids, 384
  - sterols, 385, 386
  - uronic acids, 359
  - vitamins, 395, 397
- Speech, brain involvement, 124
- Speedwell (*see Veronica*)
- Sperm
- cell division, 53
  - chemical constituents, 398, 400
  - effect of hormone on, 293
  - radiation effect on, 470, 473
  - survival time, 49
- Sphaceloma*, Melanconiaceae, 510
- Sphaeroides amulatus* (bull's-eye puffer), PLECTOGNATHI, 339
- S. maculatus* (northern puffer), 339
- S. spengleri* (bandtail puffer), 339
- Sphaerophorus*, Bacteroidaceae, 506
- Sphaeropsis*, Sphaeropsidaceae, 513, 514
- Sphaerotheca*, Erysiphaceae, 513
- Sphagnum girgensohnii* (sphagnum moss), Sphagnaceae, 7, 215, 227
- Sphaciospongia* (loggerhead sponge), HADROMERINA, 375
- Sphenodon*, RHYNCHOCEPHALIA, 144, 152
- S. punctatus* (tuatara), 2, 107
- Sphincter ani, nerve connections, 132
- Sphingolipids, properties, 384
- Sphyraena barracuda* (great barracuda), PERCOMORPHI, 337, 338
- Sphyrna zygaena* (hammerhead shark), SELACHII, 64, 270
- Spicaria*, Moniliaceae, 512
- Spicebush (*see Lindera*)
- Spider (*see Pardosa, Tegenaria*)
- Spiderwort (*see Tradescantia*)
- Spinach, 386
- Spinal cord
- tissue growth, 47
  - water content, 401, 402
- Spirillum*, Spirillaceae, 506
- Spirogyra*, Zygnemataceae, 7, 227
- Spissistilus festinus* (three-cornered alfalfa bug), HOMOPTERA, 485
- Spisula solidissima* (surf clam), EULAMELLIBRANCHIA, 341
- Spittlebug (*see Philaenus*)
- Spleen
- cell chemical constituents, 398, 399
  - cells, staining methods for, 552
  - nerve connections, 134, 138
  - parasites, 498-500, 520
  - permissible radionuclide concentration in, 459, 461-465
  - radiation effect on, 470, 473, 474
  - sterol source, 385
  - water content, 401-403
- Sponge, 385, 386
- brown (*see Fibulia*)
  - commercial (*see Hippospongia*)
  - fire (*see Tedania*)
  - freshwater (*see Spongilla*)
  - loggerhead (*see Sphaciospongia*)
  - marine (*see Grantia, Scypha*)
- Sponge sting, 341
- Spongilla lacustra* (freshwater sponge), HAPLOSCLE-RINA, 5
- Spore dispersion, 426, 427
- Sporotrichum*, Moniliaceae, 319, 520
- Spruce, 482 (*see also Picea*)
- Squalus acanthias* (Atlantic spiny dogfish), SELACHII
- arterial blood pressure, 240
  - blood volumes, 265, 270
  - endocrine tissue, 157
  - erythrocyte values, 270
  - heart rate, 236
  - hemoglobin values, 270
  - propagation, 64



- Squalus suckleyi* (Pacific spiny dogfish), 2  
 Squaretail (see *Tetragomurus*)  
 Squash, 481, 554  
 Squash bug (see *Anasa*)  
 Squid (see *Loligo*)  
 Squirrel  
   anesthetic for, 548  
   parasites, 492, 503  
   gray (see *Sciurus*)  
   ground (see *Citellus*)  
   Malabar, 516  
*Stagmomantis carolina* (Carolina mantis), ORTHOPTERA, 68  
 Staining methods, 551-557  
 Standard stages in development, 82-92  
*Staphylococcus*, Micrococcaceae  
   antibiotic activity against, 320-323  
   chemical constituents, 400  
   generation time, 52  
   parasitism, 506  
   temperature for growth, 438  
   thermal death time, 439  
 Starch, histochemical test for, 558  
 Starfish, 385, 386 (see also *Asterias*)  
 Starling (see also *Sturnus*)  
   anesthetic for, 548  
 Stem(s)  
   cell sap composition, 404  
   diseases affecting, 502, 507-511  
   mineral content, 405, 407-409, 411-413  
   osmotic potential, 455  
   parasites, 482-485  
   respiration rates, 229, 230  
   tissue culture, 116, 117  
   tissue culture medium for, 538  
   toxins, 345, 348, 349  
*Stemphylium*, Dematiaceae, 510  
*Steneotarsonemus pallidus* (cyclamen mite), ACARI, 481  
*Stentor coeruleus* (heterotrichous ciliate), HETEROTRICHIDA, 5, 51  
*Sterculia foetida* (hazel sterculia), Sterculiaceae, 377  
*Stereum*, Thelephoraceae, 512-514  
 Sterlet (see *Acipenser*)  
 Sternal marrow, differential cell counts, 275, 276  
*Sternotherus odoratus* (musk turtle), CHELONIA  
   body length, 103  
   chromosome number, 2  
   life span, 107  
   propagation, 62  
 Sternum, bones comprising, 158, 159  
 Sterols  
   nutritional requirement for, 167  
   properties, 385, 386  
*Stichopus regalis* (sea cucumber), ASPIDOCHIROTA, 3  
*S. variegatus* (curry fish), 340  
 Stickleback (see *Gasterosteus*)  
 Stigma, mineral content, 411  
*Stillingia* (stillingia), Euphorbiaceae, 373  
 Stingray (see *Dasyatis*)  
   bat (see *Myliobatis*)  
   butterfly (see *Gymnura*)  
   freshwater (see *Potamotrygon*)  
   round (see *Urolophus*)  
 Stingray sting, 338  
*Stizolobium deeringianum* (Florida velvet bean), Leguminosae, 216  
 Stomach  
   effect of hormones on, 303  
   enzymes, 139-141  
   nerve connections, 130\*, 132, 134  
   permissible radionuclide concentration in, 459-462, 466, 467  
   protein digestion, 183\*  
   radiation effect on, 470  
   tissue regeneration, 46  
   water content, 401, 402  
 Stomach worm (see *Haemonchus*, *Ostertagia*, *Trichostrongylus*)  
*Stomoxys* (stable fly), DIPTERA, 490  
*S. calcitrans* (stable fly), 480  
 Stonefish (see specific genus)  
 Stonewort (see *Chara*)  
 Strawberry (see also *Fragaria*)  
   parasites  
     arthropod, 481  
     nematode, 494, 496  
     viral, 501  
 Strawworm (see *Harmolita*)  
*Streptococcus*, Lactobacillaceae  
   antibiotic activity against, 320-323  
   culture medium for, 535  
   generation time, 52  
   nutrition, 171Fn., 181Fn.  
   parasitism, 506  
   respiration rate, 225  
   temperature for growth, 438  
   thermal death time, 439  
*Streptomyces*, Streptomycetaceae  
   antibiotics source, 312, 314, 316  
   culture medium for, 534  
   glycoside source, 370  
   parasitism, 507-510, 520, 521  
   respiration rate, 225  
   temperature for growth, 438  
*Strigomonas*, PROTOMONADINA, culture media for, 526  
*Strongyloides stercoralis* (intestinal threadworm), RHABDITIDA, 486, 490  
*Strongylus vulgaris* (single-toothed strongyle), RHABDITIDA, 490  
*Strophanthus* (strophanthus), Apocynaceae, 349, 368, 377  
*Strumella*, Tuberculariaceae, 514  
*Struthio* (ostrich), STRUTHIONIFORMES, 59, 148  
*S. camelus* (African ostrich), 107, 235  
*Strychnos nuxvomica* (nuxvomica poison nut), Loganiaceae, 349  
 STS medium for axenic culture, 524  
 Sturgeon (see *Acipenser*)  
*Sturnus* (starling), PASSERIFORMES, 59  
*S. vulgaris*  
   arterial blood pressure, 240  
   clutch size, 60  
   hatching success, 61  
   heart rate, 235  
   life span, 107  
 Sublingual gland, nerve connections, 130\*, 132, 134, 137  
 Submandibular gland, nerve connections, 137  
 Submaxillary gland  
   nerve connections, 130\*, 132, 134  
   tissue regeneration, 49  
 Subthalamus, 122\*  
   functions, 123  
 Sucrose (see also Carbohydrates)  
   synthesis, 212\*  
 Suctorian (see *Podophrya*)  
 Sugar (see also Carbohydrates)  
   photosynthetic production, 216  
 Sugar fractions of glycosides, 369, 371  
 Sugarcane, 382, 494, 496  
 Sulfur, nutritional requirement for, 180-182  
 Sulfur content, enzymes, 288, 290  
 Sumac, 370, 484 (see also *Rhus*)  
 Sunfish (see *Lepomis*, *Mola*)  
 Sunflower (see *Helianthus*)

- Suprarenal gland, nerve connections, 130\*, 131, 134, 138  
 Surgeonfish (*see Acanthurus*)  
 Surmullet (*see Parupeneus, Upeneus*)  
*Sus scrofa* (swine), ARTIODACTYLA (*see also* Swine)  
   arterial blood pressure, 240  
   blood volumes, 264, 269  
   body surface area constants, 121  
   body weight, 100, 121  
   chromosome number, 1  
   digestive enzymes, 139  
   erythrocyte values, 269  
   fatty acid source, 380  
   heart rate, 235  
   hemoglobin values, 269  
   leukocyte counts, 274  
   life span, 106  
   lung ventilation, 220  
   oxygen consumption, 221  
   platelet count, 271  
   propagation, 57  
   skeletal system, 158, 160, 162
- Suslik (*see Citellus*)  
 Swan (*see Cygnus*)  
 Sweat glands, nerve connections, 130\*, 131  
 Sweet potato, 388 (*see also Ipomoea*)  
 Swift (*see Apus*)  
 Swine (*see also Sus*)  
   developmental stages, 82, 86, 87  
   effect of plant toxins on, 345-349  
   enzyme source, 388, 389  
   fatty acid source, 373, 375, 380, 381  
   gestation time, 82  
   hormone source, 389  
   parasites  
     arthropod, 479, 480  
     bacterial, 504-506  
     fungal, 516  
     helminthic, 486, 488, 490-493  
     protozoan, 492, 493  
     rickettsial, 503  
     viral, 498  
   protein source, 388, 389  
   sterol source, 385  
   water content, tissues and organs, 403
- Swine influenza virus, generation time, 51  
 Sycamore (*see Platanus*)  
 Symbiosis, plant: in nitrogen fixation, 216, 217  
 Sympathetic nervous system, 130\*-133, 136-138  
   effect of hormones on, 299  
 Symphylid (*see Scutigera*)  
*Synanceja* (poison stonefish), SCLEROPAREI, 338  
*Synechococcus*, Chroococcaceae, 181Fn., 441  
*Synura*, CHIRYSOMONADINA, culture medium for, 529  
 Systolic blood pressure, 239-242
- Tabanus* (horsefly), DIPTERA, 492  
*T. atratus* (black horsefly), 480  
*Tachyglossus* (spiny anteater), MONOTREMATA, 159, 161, 163  
*Tachypleus tridentatus* (king crab), XIPHOSURA, 4  
 Tadpole, developmental stages, 90  
*Taenia* (tapeworm), CYCLOPHYLLIDEA, 70  
*T. pisiformis* (dog tapeworm), 4, 492  
*T. saginata* (beef tapeworm), 110, 486  
*T. solium* (pork tapeworm), 486  
*Tagetes erecta* (Aztec marigold), Compositae, 116  
 Taipan (*see Oxyuramus*)  
 Tallowwood (*see Ximemia*)  
*Tamias striatus* (eastern chipmunk), RODENTIA, 57, 106  
 Tapeworm (*see Hymenolepis, Taenia*)  
   broadheaded (*see Raillietina*)  
   dog (*see Dipylidium*)  
   fish (*see Diphyllobothrium*)  
   hydatid (*see Echinococcus*)  
   sheep (*see Moniezia*)
- Taphrina*, Taphrinaceae, 510, 512-514  
*Taraxacum officinale* (dandelion), Compositae, 454  
 Tarnished bug (*see Lygus*)  
*Taxodium distichum* (bald cypress), Taxodiaceae  
   chromosome number, 8  
   life span, 111  
   measurements, 111  
   seed germination, 77  
   shade tolerance, 443
- Taxonomic classification  
   animals, 563-565  
   plants, 566-569
- Taxus* (yew), Taxaceae, 73, 442  
*T. baccata* (English yew), 8, 229, 231  
*T. brevifolia* (Pacific yew), 111, 117  
*T. canadensis* (Canada yew), 443  
*Tedania* (fire sponge), POECILOSCLERINA, 341
- Teeth  
   radiation effect on, 468, 469, 473, 474  
   staining method for, 552  
   water content, 401, 403
- Tegenaria derhami* (spider), ARANEAE, 109  
*T. domestica* (house spider), 4
- Telencephalon, regions and functions, 123  
 Telophase, effect of compounds on, 54  
 Temperature (*see also* Body temperature)  
   conversion formula, 571  
   effect on  
     osmotic potential, 454  
     protoplasmic streaming, 448, 449  
     seed life span, 113  
     survival: bacteria, rickettsia, 439  
     transpiration rates, 451, 452  
     viruses, 432-436  
   for enzyme activity, 277-281  
   for growth: bacteria, rickettsia, 438, 440, 441  
   and light for flowering, 446, 447  
   respiratory media, 219  
   sea water, 539, 540  
   tolerances: algae, 441
- Tenebrio molitor* (yellow mealworm), COLEOPTERA  
   chromosome number, 4  
   heart rate, 237  
   hemolymph volume, 267  
   metamorphosis, 68  
   nutrition, 175Fn.  
   parasitism, 485  
   propagation, 68
- Tenebroides mauritanicus* (cadelle), COLEOPTERA, 485  
*Teredo navalis* (shipworm), EULAMELLIBRANCHIA, 109  
 Termite (*see Reticulitermes*)  
*Terrapene carolina* (box turtle), CHELONIA  
   blood volumes, 269  
   erythrocyte and hemoglobin values, 269  
   leukocyte counts, 274  
   life span, 107  
   propagation, 62
- T. ornata* (ornate box turtle), 103
- Testis(es)  
   hormones, 300, 301  
   nerve connections, 130\*, 133  
   permissible radionuclide concentration in, 459, 462  
   protein source, 389  
   radiation effect on, 469, 470, 473-475  
   sterol source, 385  
   tissue growth, 49  
   tissue regeneration, 49  
   water content, 401, 402

- Tetragonurus cuvieri* (squaretail), PERCOMORPHI, 337, 338
- Tetrahymena*, HYMENOSTOMATIDA, 168Fn., 171Fn., 175Fn.-177Fn., 181Fn.
- Tetranychus telarius* (two-spotted spider mite), ACARI, 419, 481
- Tetraodon lineatus* (puffer), PLECTOGNATHI, 339
- Tetraodon poisoning, 338
- Tetratrichomonas*, METAMONADINA, culture media for, 524, 525
- Thalamic nuclei, 125, 126
- Thalamus, regions and functions, 123, 125, 126, 129
- Thalassophryne reticulata* (venomous toadfish), HAPLODOCI, 339
- Thamnophis* (garter snake), SERPENTES, 261
- T. butleri* (Butler's garter snake), 2
- T. sirtalis* (common garter snake)
- blood volumes, 269
  - body length, 103
  - erythrocyte and hemoglobin values, 269
  - life span, 107
  - propagation, 62
- Theileria*, EUCOCCIDIA, 492
- Thelenota ananas* (prickly red fish), ASPIDOCYROTA, 340
- Theobroma cacao* (cacao), Sterculiaceae, 380
- Thermal death time: bacteria, fungi, rickettsia, 439-441
- Thermobia domestica* (firebrat), THYSANURA
- chromosome number, 4
  - metamorphosis, 68
  - parasitism, 485
  - propagation, 68
- Thevetia peruviana* (thevetia), Apocynaceae, 349
- Thielaviopsis*, Dematiaceae, 510
- Thiobacillus*, Thiobacteriaceae, 180Fn., 181Fn.
- Thioploca*, Beggiatoaceae, 181Fn.
- Thiothrix*, Beggiatoaceae, 181Fn.
- Thorny-headed worm (see *Macracanthorhynchus*)
- Threadworm (see *Strongyloides*)
- Thrips tabaci* (onion thrips), THYSANOPTERA, 485, 501
- Thuja* (arborvitae), Cupressaceae, 73
- T. occidentalis* (northern white cedar)
- chromosome number, 8
  - first flowering, 111
  - life span, 111
  - measurements, 111
  - parasites, 514
  - seed germination, 77
  - shade tolerance, 443
  - soil pH, 442
- Thunnus thynnus* (bluefin tuna), PERCOMORPHI, 105, 108
- Thymus
- cell chemical constituents, 398, 400
  - protein source, 389
  - radiation effect on, 472
  - tissue growth, 49
- Thyridopteryx ephemeriformis* (bagworm), LEPIDOPTERA, 485
- Thyroid
- comparative anatomy, 154, 155
  - effect of hormones on, 293
  - hormone source, 389
  - hormones, 294-297
  - permissible radiation exposure for, 457
  - permissible radionuclide concentration in, 462, 465
  - protein source, 389
  - tissue growth, 49
- Tick, 498, 503 (see also *Hyalomma*, *Ixodes*, *Rhipicephalus*)
- cattle (see *Boophilus*)
  - dog (see *Dermacentor*)
  - ear (see *Otobius*)
  - fowl (see *Argas*)
  - wood (see *Dermacentor*)
- Tidal volume, lungs, 220
- Tiger fish (see *Pterois*)
- Tiger snake (see *Notechis*)
- Tilia cordata* (little-leaf linden), Tiliaceae, 214
- Tilletia*, Tilletiaceae, 427, 511
- Timothy, 482 (see also *Phleum*)
- Tiphia vernalis* (Japanese beetle parasite), HYMENOPTERA, 423
- Tissue(s), animal (see also specific genus)
- culture media for, 529-534
  - fixatives, 549, 550
  - growth, 46-49
  - mineral retention in, 192\*, 193-195
  - staining methods for, 551-557
  - water content, 401-403
- Tissue(s), plant (see also specific genus)
- culture media for, 538
  - fixatives, 550
  - staining methods for, 553, 555, 556
- Tissue culture
- fungi, 516-521
  - spermatophytes, 115-117
- Tissue culture media, 529-534, 538
- Toad (see *Bufo*, *Pipa*)
- Toadfish (see *Batrachus*, *Opsanus*, *Thalassophryne*)
- Toadfish sting, 339
- Tobacco (see also *Nicotiana*)
- nutrition, 168Fn.
  - parasites
    - arthropod, 483, 484
    - fungi, 427
    - nematode, 496
    - viral, 435, 436, 501  - protein source, 389
- Tomato (see also *Lycopersicon*)
- nutrition, 168Fn.
  - parasites
    - arthropod, 483-485
    - nematode, 494
    - viral, 436, 501  - protein source, 388
- Tongue worm (see *Linguatula*)
- Torpedo* (electric ray), BATOIDEI, 155
- Torquigener hamiltoni* (puffer), PLECTOGNATHI, 339
- Torula*, Saccharomycetaceae, 182Fn.
- Torulopsis*, Cryptococcaceae, 226, 320, 440
- Touch, brain involvement, 123, 127, 129
- Toxascaris leonina* (dog roundworm), RHABDITIDA, 490
- Toxins (see table of contents, page xiv)
- Toxocara canis* (dog roundworm), RHABDITIDA, 490
- Toxoplasma*, EUCOCCIDIA, 324, 488, 492
- Toxopneustes* (sea urchin), TEMNOPLEUROIDA, 340
- Trachea, nerve connections, 130\*, 134, 137
- Tracheid, description, 45
- Trachinus* (weever), PERCOMORPHI, 339
- Trachoma virus, 432, 500
- Tradescantia* (spiderwort), Commelinaceae, 53, 54, 455
- T. virginiana* (Virginia spiderwort)
- chromosome number, 8
  - pollen life span, 114
  - shade tolerance, 443
  - soil pH, 442
- T. viridis* (wandering Jew), 231
- Tragopogon porrifolius* (salsify), Compositae, 377
- Trametes*, Polyporaceae, 513, 514
- Treponema*, Treponemataceae, 319, 322, 506
- Trialeurodes vaporariorum* (greenhouse whitefly), HOMOPTERA, 485



- Triatoma* (kissing bug), HEMIPTERA, 492  
*T. sanguisuga* (cone-nose bug), 480  
*Tribolium* (flour beetle), COLEOPTERA, 109  
*T. confusum* (confused flour beetle), 68, 485  
*Trichechus* (manatee), SIRENIA, 158, 160, 162  
*Trichinella spiralis* (trichina worm), ENOPLIDA, 109, 486, 490  
*Trichodectes canis* (dog-biting louse), MALLOPHAGA, 480, 490  
*Trichodorus* (stubby root nematode), ENOPLIDA, 496  
*Trichogramma cacaeciae* (fairyfly), HYMENOPTERA, 419  
*Trichomitis*, METAMONADINA, culture media for, 524  
*Trichomonas*, METAMONADINA  
 antibiotic activity against, 320  
 culture media for, 524, 525  
 nutrition, 167Fn., 168Fn., 173Fn.  
 parasitism, 488, 492  
*Trichophyton*, Moniliaceae, 320, 321, 516-518  
*Trichosanthes* (snake gourd), Cucurbitaceae, 375  
*Trichosporon*, Moniliaceae, 518  
*Trichostrongylus axei* (minute stomach worm), RHABDITIDA, 490  
*T. colubriformis* (hairworm), 490  
*Trichuris suis* (swine whipworm), ENOPLIDA, 490  
*T. trichiura* (human whipworm), 486  
*T. vulpis* (dog whipworm), 490  
*Trifolium* (clover), Leguminosae  
 nitrogen fixation, 217  
 parasites, 508, 510, 511  
*T. hybridum* (alsike clover)  
 breeding system, 72  
 pollen life span, 115  
 seed dispersion, 430  
 seed germination, 76  
*T. incarnatum* (crimson clover), 217, 502  
*T. pratense* (red clover)  
 chromosome number, 9  
 light for flowering, 447  
 mineral content, 409, 413  
 nitrogen fixation, 217  
 photosynthesis, 213  
 seed dispersion, 430  
 seed life span, 112, 113  
 soil pH, 442  
*T. repens* (white clover), 217, 430  
 Triggerfish (see *Balistoides*)  
 Triglycerides  
 digestion, 185\*  
 metabolism, 197\*  
*Trigonella foenum-graecum* (fenugreek), Leguminosae, 217  
*Trimeresurus mucrosquamatus* (Taiwan habu), SERPENTES, 330  
*Trioza tripunctata* (blackberry psyllid), HOMOPTERA, 423  
 Tripe, rock (see *Lasallia*)  
*Tripneustes esculentus* (sea urchin), TEMNOPLEUROIDA, 54  
*Triticum* (wheat), Gramineae, 53, 72, 75 (see also Wheat)  
*T. aestivum* (wheat)  
 cell sap composition, 404  
 chromosome number, 8  
 fatty acid source, 380, 381  
 light and temperature for flowering, 446  
 mineral content, 405, 411  
 osmotic potential, 455  
 parasites, 502, 508, 511  
 photosynthesis, 213  
 pollen life span, 114  
 respiration rates, 228, 229, 231, 232  
 seed life span, 112  
 soil pH, 442  
 transpiration rates, 451, 453  
*Tritrichomonas*, METAMONADINA  
 antibiotic activity against, 320  
 culture media for, 524, 525  
 parasitism, 492  
*Triturus* (newt), CAUDATA, 222 (see also Newt)  
*T. cristatus* (crested newt), 2, 107  
*T. viridescens* (common newt), 2, 63, 107  
*Troglodytes* (wren), PASSERIFORMES, 59  
*T. aedon* (house wren), 60, 61, 235  
*Tropaeolum majus* (nasturtium), Tropaeolaceae, 436  
 Trout  
 brook (see also *Salvelinus*)  
 anesthetic for, 548  
 brown (see also *Salmo*)  
 anesthetic for, 549  
 Trowell's T8 culture medium for animal tissue, 533  
 Trunk diameter, trees, 110, 111  
 Trunkfish (see *Lactophrys*, *Lactoria*)  
*Trypanosoma*, PROTOMONADINA  
 chromosome number, 5  
 culture media for, 526, 527  
 oxygen consumption, 224  
 parasitism, 488, 492  
 propagation, 71  
*Tsuga* (hemlock), Pinaceae, 73  
*T. canadensis* (eastern hemlock)  
 chromosome number, 8  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 parasites, 514  
 pollen life span, 114  
 seed germination, 77  
 shade tolerance, 443  
 soil pH, 442  
*T. heterophylla* (western hemlock), 430, 515  
*T. mertensiana* (mountain hemlock), 515  
 Tuatara (see *Sphenodon*)  
 Tuber  
 diseases affecting, 502, 510  
 parasites, 494-497  
 tissue culture, 115, 116  
 toxins, 346  
*Tubercularia*, Tuberculariaceae, 514  
*Tubifex* (oligochaete worm), OLIGOCHAETA†, 54  
 Tubules, kidney  
 cells, staining method for, 551  
 effect of hormones on, 295, 297, 301  
 Tubules, seminiferous: effect of hormones on, 293  
 Tulips, 427, 494  
 Tuna (see *Thunnus*)  
 Tung oil tree (see *Aleurites*)  
*Turdus*, PASSERIFORMES, 59  
*T. merula* (blackbird), 1 (see also Blackbird)  
*T. migratorius* (American robin)  
 arterial blood pressure, 240  
 clutch size, 60  
 hatching success, 61  
 heart rate, 235  
 life span, 107  
 Turkey, 478, 491, 493 (see also *Meleagris*)  
 wild, anesthetic for, 548  
 Turkey fish (see *Pterois*)  
 Turnip, 494 (see also *Brassica*)  
 Turtle (see also specific genus)  
 anesthetic for, 548  
 Turtle poisoning, 339  
 Twinflower (see *Linnaea*)  
*Tylenchorhynchus* (stunt nematode), TYLENCHIDA, 496  
*Tylenchulus semipenetrans* (citrus nematode), TYLENCHIDA, 496  
*Tylosiderma fragariae* (strawberry crown borer), COLEOPTERA, 423  
*Typha latifolia* (cattail), Typhaceae, 454

† Class



- Typhula*, Clavariaceae, 509  
 Tyrode's solution for animal tissue culture, 529
- Ulmus* (elm), Ulmaceae  
 breeding system, 72  
 parasites, 508, 514, 515  
 pollen dispersion, 430  
 propagation methods, 74
- U. americana* (American elm)  
 chromosome number, 9  
 first flowering, 111  
 life span, 111  
 measurements, 111  
 mineral content, 409, 413  
 seed germination, 77  
 seed life span, 112, 113  
 shade tolerance, 443  
 soil pH, 442
- U. glabra* (Scotch elm), 231
- Ulothrix*, Ulotrichaceae, 7, 227, 441
- Ultimobranchial bodies, comparative anatomy, 154, 155
- Uva lactuca* (sea lettuce), Ulvaceae, 213, 214
- Umbilicaria*, Umbilicariaceae, 227
- Uncinula*, Erysiphaceae, 511-514
- Unio* (pearl mussel), EULAMELLIBRANCHIA, 4
- Unsaturated fatty acids  
 in fats and oils, 381  
 properties, 372-379
- Upas tree (*see Antiaris*)
- Upeneus arge* (surmullet), PERCOMORPHI, 337, 338
- Uredinopsis*, Melampsoraceae, 512
- Ureter, nerve connections, 132, 135
- Urethra, nerve connections, 133, 135
- Urine  
 composition, 186-188  
 hormone source, 388  
 minerals excreted in, 192\*, 193-195  
 parasites, 498  
 protein source, 388
- Urobacillus*, Bacillaceae, 218\*
- Urocystis*, Tilletiaceae, 508, 511
- Urohypophysis, comparative anatomy, 156, 157
- Urolophus halleri* (round stingray), BATOIDEI, 338
- Uromyces*, Pucciniaceae, 509, 510
- Uronic acids, properties, 358, 359
- Urophlyctis*, Physodermataceae, 510
- Ursus americanus* (black bear), CARNIVORA, 418
- Urtica* (nettle), Urticaceae, 455
- Usnea*, Usneaceae, 227
- Ustilago*, Ustilaginaceae  
 chromosome number, 6  
 parasitism, 509, 511  
 respiration rate, 226  
 spore dispersion, 427  
 temperature for growth, 441  
 thermal death time, 441
- Uterine tube, nerve connections, 133
- Uterus  
 effect of hormones on, 293, 299  
 nerve connections, 133, 136-138  
 parasite, 493  
 tissue growth, 49  
 tissue regeneration, 49  
 water content, 401
- Vaccinia virus, 432, 500
- Vagina  
 nerve connections, 133, 138  
 parasites, 488, 489, 518, 520
- Vallisneria spiralis* (spiral wild celery), Hydrocharitaceae, 448
- Valsa*, Allantosporae, 513, 514
- Vas deferens*, nerve connections, 133, 135
- Vaucheria*, Vaucheriaceae, 6
- Vein(s)  
 blood pressure, 241, 242  
 comparative anatomy, 146-149
- Veneer graft, 73, 74
- Venoms (*see* Toxins)
- Venous blood  
 acid-base balance, 259-261  
 carbohydrate absorption by, 184\*  
 fatty acid absorption, 185\*  
 pressure, 241, 242
- Venous hematocrit, 263-265
- Ventricle(s)  
 blood pressure, 241  
 water content, 401, 402
- Venturia*, Pleosporaceae  
 chromosome number, 6  
 parasitism, 510, 512  
 temperature for growth, 441
- Veratrum viride* (American false hellebore), Liliaceae, 349
- Verbascum thapsus* (flannel mullein), Scrophulariaceae, 454
- Verbena, 494
- Vernonia anthelmintica* (kinka oil ironweed), Compositae, 377
- Veronica beccabunga* (beccabunga speedwell), Scrophulariaceae, 451
- Vertebrae, number, 158, 159
- Verticillium*, Moniliaceae, 509, 510, 512, 514
- Vespertilio murinus* (parti-colored bat), CHIROPTERA, 418
- Vetch, 494 (*see also Astragalus, Vicia*)
- Vibrio*, Spirillaceae  
 antibiotic activity against, 320, 321  
 generation time, 52  
 parasitism, 506  
 temperature for growth, 438  
 thermal death time, 439
- Vicia*, Leguminosae, 217
- V. faba* (broad bean)  
 breeding system, 72  
 cell division, 54  
 chromosome number, 9  
 light and temperature for flowering, 447  
 mineral content, 409, 413  
 osmotic potential, 454  
 parasites, 433, 508  
 pollen life span, 115  
 respiration rates, 228-231  
 seed germination, 76
- V. faba equina* (horsebean), 442
- V. faba minor* (small horsebean), 433
- V. sativa* (vetch), 213
- Vigna sinensis* (cowpea), Leguminosae (*see also* Pea)  
 cell sap composition, 404  
 nitrogen fixation, 216, 217  
 transpiration rate, 453
- Vinca rosea* (Madagascar periwinkle), Apocynaceae, 117
- Violet, 373
- Viper (*see Echis, Vipera*)
- Vipera berus* (European viper), SERPENTES, 330
- V. russelli* (Russell's viper), 330
- Viruses (*see* specific virus)
- Vision, brain involvement, 123, 124, 128
- Vitamin(s)  
 antimetabolites, 309-311  
 in feces, 190

- nutritional requirement for, 172-175, 181  
 properties, 394-397  
 in urine, 187
- Vitis* (grape), Vitaceae (see also Grape)  
 mineral content, 413  
 parasites, 508, 511  
 pollen life span, 115  
 propagation methods, 74  
 soil pH, 442
- V. vinifera* (European grape)  
 breeding system, 72  
 chromosome number, 9  
 mineral content, 409  
 photosynthesis, 213  
 respiration rates, 231, 232  
 tissue culture, 117
- Vole, 506
- Volvox* (pale-green flagellate), PHYTOMONADINA, 5, 72
- Vulture (see *Gyps*)
- Walnut, 482 (see also *Juglans*)
- Walrus, 486
- Wart virus, 500
- Wasp  
   gall (see *Amphibolips*)  
   parasitic (see *Apanteles*, *Habrobracon*)  
   sea (see *Carybdea*, *Chiropsalmus*)
- Waspfish (see *Centropogon*)
- Water  
   in acid-base balance, blood, 259-261  
   in amino acid metabolism, 200  
   in carbohydrate metabolism, 198\*  
   freezing point, 579  
   in Krebs cycle, 205Fn.  
   in metabolism, 197\*, 203\*  
   in nucleoprotein catabolism, 201\*  
   as respiratory medium, 219
- Water content  
   feces, 190  
   organs and tissues, 401-403  
   urine, 186
- Water flea (see *Daphnia*)
- Water net (see *Hormidium*)
- Water snake (see *Natrix*)
- Waterweed (see *Elodea*)
- Waxes  
   fatty acid source, 373, 377  
   properties, 382
- Weasel (see *Mustela*)
- Webworm (see *Hyphantria*)
- Weeds, 482, 484
- Weever (see *Trachinus*)
- Weever-fish sting, 339
- Weevil  
   bean (see *Acanthoscelides*)  
   boll (see *Anthonomus*)  
   pea (see *Bruchus*)  
   rice (see *Sitophilus*)  
   sugarcane (see *Rhagoletis*)  
   sweet potato (see *Cylas*)
- Weight (see Atomic, Body, and Molecular Weights)
- Whale, 373, 375, 380  
   beluga (see *Delphinapterus*)  
   finback (see *Balaenoptera*)  
   Greenland (see *Balaena*)  
   sperm (see *Physeter*)
- Wheat (see also *Triticum*)  
   parasites  
     arthropod, 482-484  
     fungal, 427  
     nematode, 494, 496  
     viral, 426, 502  
   protein source, 388  
   sterol source, 385, 386
- Whelk (see *Buccinum*, *Busycon*)
- Whip graft, 73, 74
- Whipworm (see *Trichuris*)
- White pine, 427
- White worm (see *Enchytraeus*)
- White's solution for tissue culture, 538
- Whitefish (see *Coregonus*)
- Whitefly (see *Bemisia*, *Trialeurodes*)
- Willow, 368 (see also *Chilopsis*, *Salix*)
- Wind, effect on transpiration rates, 452
- Wintergreen, glycoside source, 368
- Wireworm, 481
- Wohlfahrtia vigil* (flesh fly), DIPTERA, 480
- Wolf, 491, 493
- Woodchuck (see *Marmota*)
- Worm bite, 340
- Wormwood, 368
- Wound tumor virus, 502
- Wrasse (see *Epibolus*)
- Wren (see *Troglodytes*)
- Wuchereria bancrofti* (Bancroft's filarial worm), SPIRURIDA, 110, 486
- Xanthium* (cocklebur), Compositae, 443, 454
- Xanthomonas*, Pseudomonadaceae, 52, 438, 506-508
- Xenopus laevis* (clawed frog), SALIENTIA, 2, 63, 107
- Ximenia* (tallowwood), Olacaceae, 373
- Xiphinema* (dagger nematode), ENOPLIDA, 496
- Xylaria*, Xylariaceae, 510, 512, 513
- Xylem, staining method for, 553
- Yam (see *Dioscorea*)
- Yeast(s)  
   culture media for, 536  
   enzyme source, 388, 389  
   sterol source, 385
- Yellow fever virus, 432, 500
- Yew (see *Taxus*)
- Yucca* (yucca), Liliaceae, 8, 72
- Y. gloriosa* (mound lily yucca), 231
- Zea mays* (corn), Gramineae (see also Corn)  
   breeding system, 72  
   cell division, 53  
   cell sap composition, 404  
   chromosome linkage groups, 39-41  
   chromosome number, 8, 39  
   fatty acid source, 380, 381  
   light for flowering, 446  
   mineral content, 405, 406, 411  
   mutations, 39-41  
   osmotic potential, 455  
   parasites, 433, 502, 508, 511  
   photosynthesis, 213, 215  
   pollen dispersion, 430  
   pollen life span, 114  
   respiration rates, 228, 230-232  
   seed dispersion, 430  
   seed germination, 75  
   seed life span, 112  
   soil pH, 442  
   tissue culture, 117  
   transpiration rates, 451, 453
- Zebra fish* (see *Pterois*)
- Zebu, 493
- Zenaidura* (dove), COLUMBIFORMES, 59
- Z. macroura* (mourning dove), 61, 235
- Zigadenus* (death camas), Liliaceae, 349
- Zinnia, 494
- Zygote development, 69-72

# *Contrails*

UNCLASSIFIED  
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
<b>1. ORIGINATING ACTIVITY (Corporate author)</b> Federation of American Societies for Experimental Biology, Washington, D. C.	<b>2a. REPORT SECURITY CLASSIFICATION</b> UNCLASSIFIED	
	<b>2b. GROUP</b> N/A	
<b>3. REPORT TITLE</b>  BIOLOGY DATA BOOK		
<b>4. DESCRIPTIVE NOTES (Type of report and inclusive dates)</b> Supersedes WADC TR 56-273 (AD 110501) Handbook of Biological Data, October 1956		
<b>5. AUTHOR(S) (Last name, first name, initial)</b>  Compiled and edited by: Altman, Philip L. Dittmer, Dorothy S.		
<b>6. REPORT DATE</b>  October 1964	<b>7a. TOTAL NO. OF PAGES</b>  646	<b>7b. NO. OF REFS included after each table</b>
<b>8a. CONTRACT OR GRANT NO.</b> AF 33(657)-1082; NIH GM 06533*; and NSF GN 255** <b>b. PROJECT NO</b> 7164 <b>c. Task No.</b> 716406 <b>d.</b>	<b>9a. ORIGINATOR'S REPORT NUMBER(S)</b>  <b>9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)</b> AMRL-TR-64-100 Library of Congress cat. card no. 64-20958	
<b>10. AVAILABILITY/LIMITATION NOTICES</b> Qualified requesters may obtain copies of this report from DDC. Available, for sale to the public, from Federation of American Societies for Experimental Biology, Washington, D. C.		
<b>11. SUPPLEMENTARY NOTES</b> Prepared under the joint sponsorship of: * National Institutes of Health ** National Science Foundation	<b>12. SPONSORING MILITARY ACTIVITY</b> Aerospace Medical Research Laboratories, Aerospace Medical Division, Air Force Systems Command, Wright-Patterson AFB, Ohio	
<b>13. ABSTRACT</b>  The Biology Data Book has been compiled to present numerical data of biology and medicine in a convenient and accessible form for reference, and to standardize accepted constants as a basis for correlation, establish common standards for statistical studies, and provide normal values for research. The biology data are organized in the form of tables, diagrams, charts, and graphs, arranged under the following headings: Genetics and Cytology, Reproduction, Development and Growth, Morphology, Nutrition and Digestion, Metabolism, Respiration and Circulation, Blood, Biological Regulators and Toxins, Biophysical and Biochemical Characteristics, Environment and Survival, Parasitism, and Materials and Methods. Seven appendices provide information concerning estimated number of species, taxonomic classification for living plants and animals, geologic distribution, atomic weights, as well as logarithms and antilogarithms. A detailed index completes the book. The contents have been authenticated by 470 authorities in the fields of biology and medicine. The review process of the tables was designed to eliminate, insofar as possible, errors of transcription and material of questionable validity.		

DD FORM 1473  
1 JAN 64  
AF-WP-B-AUG 64 400

UNCLASSIFIED  
Security Classification



UNCLASSIFIED  
Security Classification

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Genetics						
	Cytology						
	Reproduction						
	Morphology						
	Nutrition						
	Digestion						
	Metabolism						
	Respiration						
	Circulation						
	Blood						
	Enzymes						
	Hormones						
	Antibiotics						
	Anticoagulants						

**INSTRUCTIONS**

**1. ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

**2a. REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

**2b. GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

**3. REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

**4. DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

**5. AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

**6. REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

**7a. TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

**7b. NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

**8a. CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

**8b, 8c, & 8d. PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

**9a. ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

**9b. OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

**10. AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

**11. SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

**12. SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

**13. ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

**14. KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

UNCLASSIFIED  
Security Classification